

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages^3				
...	All causes	122,026	100.0	305.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,298	23.2	70.9
2	Malignant neoplasms (C00-C97)	24,070	19.7	60.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,418	8.5	26.1
4	Cerebrovascular diseases (I60-I69)	6,658	5.5	16.7
5	Diabetes mellitus (E10-E14)	6,179	5.1	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,382	2.8	8.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,355	2.7	8.4
8	Chronic lower respiratory diseases (J40-J47)	3,174	2.6	8.0
9	Influenza and pneumonia (J10-J18)	2,948	2.4	7.4
10	Certain conditions originating in the perinatal period (P00-P96)	2,628	2.2	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,170	1.8	5.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,007	1.6	5.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,890	1.5	4.7
14	Alzheimer's disease (G30)	1,835	1.5	4.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,787	1.5	4.5
...	All other causes (Residual)	21,227	17.4	53.2
Hispanic, both sexes, 1-4 years				
...	All causes	993	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	32.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	12.3	3.7
3	Malignant neoplasms (C00-C97)	98	9.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	8.3	2.5
5	Influenza and pneumonia (J10-J18)	37	3.7	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.5	1.1
7	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
9	Septicemia (A40-A41)	10	1.0	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
13	Anemias (D50-D64)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	224	22.6	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,173	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	36.7	5.8
2	Malignant neoplasms (C00-C97)	217	18.5	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	5.5	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.5	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.6	0.4
7	Influenza and pneumonia (J10-J18)	22	1.9	0.3
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	Cerebrovascular diseases (I60-I69)	14	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Septicemia (A40-A41)	7	0.6	*
13	Diseases of appendix (K35-K38)	5	0.4	*
14	Anemias (D50-D64)	4	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	241	20.5	3.2
Hispanic, both sexes, 15-24 years				
...	All causes	5,394	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,366	43.9	33.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,286	23.8	18.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	491	9.1	7.0
4	Malignant neoplasms (C00-C97)	288	5.3	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	2.5	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	1.1	0.8
7	Influenza and pneumonia (J10-J18)	36	0.7	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	28	0.5	0.4
9	Legal intervention (Y35,Y89.0)	28	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-099)	24	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.3
13	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
14	Diabetes mellitus (E10-E14)	19	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
...	All other causes (Residual)	545	10.1	7.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,765	100.0	111.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,044	27.4	30.4
2	Malignant neoplasms (C00-C97)	1,877	12.7	14.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,473	10.0	11.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,226	8.3	9.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	971	6.6	7.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	901	6.1	6.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	569	3.9	4.3
8	Cerebrovascular diseases (I60-I69)	356	2.4	2.7
9	Diabetes mellitus (E10-E14)	261	1.8	2.0
10	Influenza and pneumonia (J10-J18)	128	0.9	1.0
11	Septicemia (A40-A41)	121	0.8	0.9
12	Viral hepatitis (B15-B19)	117	0.8	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.7	0.8
15	Chronic lower respiratory diseases (J40-J47)	87	0.6	0.7
...	All other causes (Residual)	2,426	16.4	18.2
Hispanic, both sexes, 45-64 years				
...	All causes	27,429	100.0	464.4
1	Malignant neoplasms (C00-C97)	7,563	27.6	128.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,489	20.0	92.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,893	6.9	32.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,729	6.3	29.3
5	Diabetes mellitus (E10-E14)	1,714	6.2	29.0
6	Cerebrovascular diseases (I60-I69)	1,305	4.8	22.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	810	3.0	13.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	492	1.8	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	420	1.5	7.1
10	Viral hepatitis (B15-B19)	417	1.5	7.1
11	Chronic lower respiratory diseases (J40-J47)	400	1.5	6.8
12	Influenza and pneumonia (J10-J18)	375	1.4	6.3
13	Septicemia (A40-A41)	338	1.2	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	321	1.2	5.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	196	0.7	3.3
...	All other causes (Residual)	3,967	14.5	67.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 65 years and over			
...	All causes	66,944	100.0	3,257.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,301	31.8	1,036.4
2	Malignant neoplasms (C00-C97)	14,013	20.9	681.8
3	Cerebrovascular diseases (I60-I69)	4,930	7.4	239.9
4	Diabetes mellitus (E10-E14)	4,180	6.2	203.4
5	Chronic lower respiratory diseases (J40-J47)	2,634	3.9	128.2
6	Influenza and pneumonia (J10-J18)	2,284	3.4	111.1
7	Alzheimer's disease (G30)	1,795	2.7	87.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,512	2.3	73.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,213	1.8	59.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,081	1.6	52.6
11	Septicemia (A40-A41)	960	1.4	46.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	819	1.2	39.8
13	Parkinson's disease (G20-G21)	504	0.8	24.5
14	Pneumonitis due to solids and liquids (J69)	444	0.7	21.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	384	0.6	18.7
...	All other causes (Residual)	8,890	13.3	432.5
	Hispanic, male, all ages^3			
...	All causes	68,119	100.0	330.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,867	21.8	72.2
2	Malignant neoplasms (C00-C97)	12,671	18.6	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,899	11.6	38.3
4	Cerebrovascular diseases (I60-I69)	3,070	4.5	14.9
5	Diabetes mellitus (E10-E14)	2,976	4.4	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,809	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,387	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,711	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,697	2.5	8.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,465	2.2	7.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,459	2.1	7.1
12	Influenza and pneumonia (J10-J18)	1,429	2.1	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	937	1.4	4.5
15	Septicemia (A40-A41)	721	1.1	3.5
...	All other causes (Residual)	10,952	16.1	53.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 1-4 years			
...	All causes	569	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	35.1	11.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	10.9	3.7
3	Malignant neoplasms (C00-C97)	60	10.5	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.3	2.1
5	Influenza and pneumonia (J10-J18)	24	4.2	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Septicemia (A40-A41)	6	1.1	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Anemias (D50-D64)	2	0.4	*
14	Nutritional deficiencies (E40-E64)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	118	20.7	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	706	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	40.9	7.5
2	Malignant neoplasms (C00-C97)	123	17.4	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.1	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.4	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	2.7	*
7	Influenza and pneumonia (J10-J18)	12	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
11	Septicemia (A40-A41)	4	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
15	Diabetes mellitus (E10-E14)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Diseases of appendix (K35-K38)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
...	All other causes (Residual)	126	17.8	3.3
Hispanic, male, 15-24 years				
...	All causes	4,244	100.0	112.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,905	44.9	50.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,139	26.8	30.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	9.9	11.2
4	Malignant neoplasms (C00-C97)	186	4.4	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.4	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	0.8
7	Legal intervention (Y35,Y89.0)	28	0.7	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	0.5
8	Influenza and pneumonia (J10-J18)	20	0.5	0.5
10	Cerebrovascular diseases (I60-I69)	15	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
14	Diabetes mellitus (E10-E14)	11	0.3	*
15	Septicemia (A40-A41)	8	0.2	*
...	All other causes (Residual)	322	7.6	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 25-44 years			
...	All causes	10,629	100.0	149.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,315	31.2	46.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,253	11.8	17.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	898	8.4	12.6
4	Malignant neoplasms (C00-C97)	840	7.9	11.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	775	7.3	10.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	734	6.9	10.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	459	4.3	6.4
8	Cerebrovascular diseases (I60-I69)	213	2.0	3.0
9	Diabetes mellitus (E10-E14)	154	1.4	2.2
10	Viral hepatitis (B15-B19)	88	0.8	1.2
11	Influenza and pneumonia (J10-J18)	76	0.7	1.1
12	Septicemia (A40-A41)	68	0.6	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.6	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.6	0.9
15	Chronic lower respiratory diseases (J40-J47)	45	0.4	0.6
...	All other causes (Residual)	1,584	14.9	22.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	17,458	100.0	601.8
1	Malignant neoplasms (C00-C97)	3,992	22.9	137.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,854	22.1	132.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,415	8.1	48.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,384	7.9	47.7
5	Diabetes mellitus (E10-E14)	963	5.5	33.2
6	Cerebrovascular diseases (I60-I69)	776	4.4	26.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	652	3.7	22.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	349	2.0	12.0
9	Viral hepatitis (B15-B19)	310	1.8	10.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	1.5	9.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	253	1.4	8.7
12	Chronic lower respiratory diseases (J40-J47)	225	1.3	7.8
13	Influenza and pneumonia (J10-J18)	215	1.2	7.4
14	Septicemia (A40-A41)	189	1.1	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	126	0.7	4.3
...	All other causes (Residual)	2,488	14.3	85.8
Hispanic, male, 65 years and over				
...	All causes	31,534	100.0	3,634.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,929	31.5	1,144.4
2	Malignant neoplasms (C00-C97)	7,466	23.7	860.5
3	Cerebrovascular diseases (I60-I69)	2,040	6.5	235.1
4	Diabetes mellitus (E10-E14)	1,845	5.9	212.6
5	Chronic lower respiratory diseases (J40-J47)	1,392	4.4	160.4
6	Influenza and pneumonia (J10-J18)	1,042	3.3	120.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	703	2.2	81.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	2.2	78.7
9	Alzheimer's disease (G30)	616	2.0	71.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	541	1.7	62.4
11	Septicemia (A40-A41)	416	1.3	47.9
12	Parkinson's disease (G20-G21)	303	1.0	34.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	287	0.9	33.1
14	Aortic aneurysm and dissection (I71)	220	0.7	25.4
15	Pneumonitis due to solids and liquids (J69)	211	0.7	24.3
...	All other causes (Residual)	3,840	12.2	442.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, all ages^3				
...	All causes	53,907	100.0	279.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,431	24.9	69.6
2	Malignant neoplasms (C00-C97)	11,399	21.1	59.1
3	Cerebrovascular diseases (I60-I69)	3,588	6.7	18.6
4	Diabetes mellitus (E10-E14)	3,203	5.9	16.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,519	4.7	13.1
6	Influenza and pneumonia (J10-J18)	1,519	2.8	7.9
7	Chronic lower respiratory diseases (J40-J47)	1,477	2.7	7.7
8	Alzheimer's disease (G30)	1,202	2.2	6.2
9	Certain conditions originating in the perinatal period (P00-P96)	1,169	2.2	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	2.0	5.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	995	1.8	5.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	850	1.6	4.4
13	Septicemia (A40-A41)	784	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	616	1.1	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	1.0	2.8
...	All other causes (Residual)	9,508	17.6	49.3
Hispanic, female, 1-4 years				
...	All causes	424	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	29.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	14.2	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.8	2.9
4	Malignant neoplasms (C00-C97)	38	9.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.8	*
6	Influenza and pneumonia (J10-J18)	13	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Septicemia (A40-A41)	4	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
10	Anemias (D50-D64)	3	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Meningitis (G00,G03)	2	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Cerebrovascular diseases (I60-I69)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	97	22.9	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 5-14 years			
...	All causes	467	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	30.4	3.9
2	Malignant neoplasms (C00-C97)	94	20.1	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.9	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.6	*
7	Influenza and pneumonia (J10-J18)	10	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
11	Diseases of appendix (K35-K38)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
13	Septicemia (A40-A41)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	Anemias (D50-D64)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
...	All other causes (Residual)	106	22.7	2.9
	Hispanic, female, 15-24 years			
...	All causes	1,150	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	461	40.1	14.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	12.8	4.5
3	Malignant neoplasms (C00-C97)	102	8.9	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	6.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.0	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	2.3	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	24	2.1	0.7
8	Influenza and pneumonia (J10-J18)	16	1.4	*
9	Cerebrovascular diseases (I60-I69)	13	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.0	*
11	Septicemia (A40-A41)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	206	17.9	6.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,136	100.0	66.9
1	Malignant neoplasms (C00-C97)	1,037	25.1	16.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	17.6	11.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	328	7.9	5.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	237	5.7	3.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	5.3	3.6
6	Cerebrovascular diseases (I60-I69)	143	3.5	2.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	3.0	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	110	2.7	1.8
9	Diabetes mellitus (E10-E14)	107	2.6	1.7
10	Pregnancy, childbirth and the puerperium (O00-099)	74	1.8	1.2
11	Septicemia (A40-A41)	53	1.3	0.9
12	Influenza and pneumonia (J10-J18)	52	1.3	0.8
13	Chronic lower respiratory diseases (J40-J47)	42	1.0	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.0	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	0.6
...	All other causes (Residual)	797	19.3	12.9
Hispanic, female, 45-64 years				
...	All causes	9,971	100.0	331.8
1	Malignant neoplasms (C00-C97)	3,571	35.8	118.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,635	16.4	54.4
3	Diabetes mellitus (E10-E14)	751	7.5	25.0
4	Cerebrovascular diseases (I60-I69)	529	5.3	17.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	478	4.8	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	345	3.5	11.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	2.3	7.5
8	Chronic lower respiratory diseases (J40-J47)	175	1.8	5.8
9	Influenza and pneumonia (J10-J18)	160	1.6	5.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	158	1.6	5.3
11	Septicemia (A40-A41)	149	1.5	5.0
12	Viral hepatitis (B15-B19)	107	1.1	3.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	0.7	2.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	70	0.7	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	0.7	2.3
...	All other causes (Residual)	1,479	14.8	49.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	35,410	100.0	2,981.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,372	32.1	957.5
2	Malignant neoplasms (C00-C97)	6,547	18.5	551.2
3	Cerebrovascular diseases (I60-I69)	2,890	8.2	243.3
4	Diabetes mellitus (E10-E14)	2,335	6.6	196.6
5	Influenza and pneumonia (J10-J18)	1,242	3.5	104.6
5	Chronic lower respiratory diseases (J40-J47)	1,242	3.5	104.6
7	Alzheimer's disease (G30)	1,179	3.3	99.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	809	2.3	68.1
9	Septicemia (A40-A41)	544	1.5	45.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	540	1.5	45.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	532	1.5	44.8
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	1.5	44.6
13	Pneumonitis due to solids and liquids (J69)	233	0.7	19.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	208	0.6	17.5
15	Parkinson's disease (G20-G21)	201	0.6	16.9
...	All other causes (Residual)	5,006	14.1	421.5
White, non-Hispanic, both sexes, all ages^3				
...	All causes	1,979,465	100.0	993.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	565,808	28.6	284.0
2	Malignant neoplasms (C00-C97)	457,127	23.1	229.5
3	Cerebrovascular diseases (I60-I69)	127,969	6.5	64.2
4	Chronic lower respiratory diseases (J40-J47)	113,541	5.7	57.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82,815	4.2	41.6
6	Alzheimer's disease (G30)	57,268	2.9	28.7
7	Influenza and pneumonia (J10-J18)	54,617	2.8	27.4
8	Diabetes mellitus (E10-E14)	52,894	2.7	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,497	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,434	1.3	13.3
11	Septicemia (A40-A41)	25,664	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,602	1.0	10.3
13	Parkinson's disease (G20-G21)	16,499	0.8	8.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,878	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	14,845	0.7	7.5
...	All other causes (Residual)	316,007	16.0	158.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,539	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	38.8	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	10.7	3.0
3	Malignant neoplasms (C00-C97)	221	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	4.8	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.3	0.9
6	Influenza and pneumonia (J10-J18)	80	3.2	0.9
7	Septicemia (A40-A41)	43	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	31	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
12	Meningitis (G00,G03)	14	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	579	22.8	6.3
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,884	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,535	39.5	6.2
2	Malignant neoplasms (C00-C97)	637	16.4	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	3.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	2.8	0.4
7	Influenza and pneumonia (J10-J18)	86	2.2	0.3
8	Septicemia (A40-A41)	47	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	39	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	37	1.0	0.1
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Anemias (D50-D64)	17	0.4	*
14	Pneumonitis due to solids and liquids (J69)	11	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (Residual)	774	19.9	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,640	100.0	75.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,638	54.2	41.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,832	14.4	10.9
3	Malignant neoplasms (C00-C97)	1,039	5.3	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	907	4.6	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	605	3.1	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	1.5	1.1
7	Influenza and pneumonia (J10-J18)	137	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	124	0.6	0.5
9	Diabetes mellitus (E10-E14)	90	0.5	0.3
10	Septicemia (A40-A41)	86	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	82	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	41	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.2	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	31	0.2	0.1
...	All other causes (Residual)	2,647	13.5	10.2
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	81,652	100.0	148.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,344	24.9	37.0
2	Malignant neoplasms (C00-C97)	13,077	16.0	23.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,682	13.1	19.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,374	11.5	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,254	2.8	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,100	2.6	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,078	2.5	3.8
8	Diabetes mellitus (E10-E14)	1,606	2.0	2.9
9	Cerebrovascular diseases (I60-I69)	1,537	1.9	2.8
10	Influenza and pneumonia (J10-J18)	843	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	757	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	715	0.9	1.3
13	Septicemia (A40-A41)	668	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	468	0.6	0.9
15	Viral hepatitis (B15-B19)	464	0.6	0.8
...	All other causes (Residual)	14,685	18.0	26.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	319,430	100.0	612.1
1	Malignant neoplasms (C00-C97)	111,901	35.0	214.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,073	23.2	142.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,764	5.9	36.0
4	Chronic lower respiratory diseases (J40-J47)	13,190	4.1	25.3
5	Diabetes mellitus (E10-E14)	10,285	3.2	19.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10,016	3.1	19.2
7	Cerebrovascular diseases (I60-I69)	9,587	3.0	18.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,236	2.9	17.7
9	Septicemia (A40-A41)	3,766	1.2	7.2
10	Influenza and pneumonia (J10-J18)	3,634	1.1	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,174	1.0	6.1
12	Viral hepatitis (B15-B19)	2,208	0.7	4.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,845	0.6	3.5
14	Aortic aneurysm and dissection (I71)	1,726	0.5	3.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,503	0.5	2.9
...	All other causes (Residual)	44,522	13.9	85.3
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,538,976	100.0	5,182.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	480,003	31.2	1,616.5
2	Malignant neoplasms (C00-C97)	330,191	21.5	1,112.0
3	Cerebrovascular diseases (I60-I69)	116,614	7.6	392.7
4	Chronic lower respiratory diseases (J40-J47)	99,440	6.5	334.9
5	Alzheimer's disease (G30)	56,707	3.7	191.0
6	Influenza and pneumonia (J10-J18)	49,693	3.2	167.4
7	Diabetes mellitus (E10-E14)	40,885	2.7	137.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,031	2.0	101.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,717	1.8	93.3
10	Septicemia (A40-A41)	20,958	1.4	70.6
11	Parkinson's disease (G20-G21)	16,142	1.0	54.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,260	0.9	48.0
13	Pneumonitis due to solids and liquids (J69)	13,749	0.9	46.3
14	Atherosclerosis (I70)	10,933	0.7	36.8
15	Aortic aneurysm and dissection (I71)	10,709	0.7	36.1
...	All other causes (Residual)	220,944	14.4	744.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages^3				
...	All causes	956,194	100.0	979.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	276,241	28.9	282.9
2	Malignant neoplasms (C00-C97)	236,150	24.7	241.8
3	Chronic lower respiratory diseases (J40-J47)	53,593	5.6	54.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51,888	5.4	53.1
5	Cerebrovascular diseases (I60-I69)	48,549	5.1	49.7
6	Diabetes mellitus (E10-E14)	25,940	2.7	26.6
7	Influenza and pneumonia (J10-J18)	23,541	2.5	24.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,085	2.2	21.6
9	Alzheimer's disease (G30)	16,427	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,323	1.6	15.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,254	1.4	13.6
12	Septicemia (A40-A41)	11,181	1.2	11.4
13	Parkinson's disease (G20-G21)	9,303	1.0	9.5
14	Aortic aneurysm and dissection (I71)	7,731	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,454	0.8	7.6
...	All other causes (Residual)	138,534	14.5	141.9
White, non-Hispanic, male, 1-4 years				
...	All causes	1,423	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	610	42.9	12.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	9.6	2.9
3	Malignant neoplasms (C00-C97)	123	8.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	4.3	1.3
5	Influenza and pneumonia (J10-J18)	44	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.4	0.7
7	Septicemia (A40-A41)	29	2.0	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
10	Cerebrovascular diseases (I60-I69)	8	0.6	*
11	Meningitis (G00,G03)	7	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
13	Meningococcal infection (A39)	6	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	307	21.6	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,319	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	41.1	7.5
2	Malignant neoplasms (C00-C97)	348	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	5.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	5.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	3.8	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	3.1	0.6
7	Influenza and pneumonia (J10-J18)	39	1.7	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.0	0.2
9	Cerebrovascular diseases (I60-I69)	23	1.0	0.2
10	Septicemia (A40-A41)	22	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	21	0.9	0.2
12	Diabetes mellitus (E10-E14)	12	0.5	*
13	Anemias (D50-D64)	9	0.4	*
14	Meningitis (G00,G03)	6	0.3	*
14	Pneumonitis due to solids and liquids (J69)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
...	All other causes (Residual)	444	19.1	3.5
White, non-Hispanic, male, 15-24 years				
...	All causes	14,022	100.0	105.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	54.9	58.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,412	17.2	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	665	4.7	5.0
4	Malignant neoplasms (C00-C97)	607	4.3	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	368	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	1.2	1.3
7	Influenza and pneumonia (J10-J18)	75	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	72	0.5	0.5
9	Diabetes mellitus (E10-E14)	54	0.4	0.4
10	Septicemia (A40-A41)	48	0.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	48	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.2	0.3
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
15	Aortic aneurysm and dissection (I71)	21	0.1	0.2
...	All other causes (Residual)	1,696	12.1	12.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	53,220	100.0	192.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,712	27.6	53.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,641	14.4	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,395	13.9	26.8
4	Malignant neoplasms (C00-C97)	6,021	11.3	21.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,723	3.2	6.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,492	2.8	5.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,395	2.6	5.1
8	Diabetes mellitus (E10-E14)	1,006	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	796	1.5	2.9
10	Influenza and pneumonia (J10-J18)	487	0.9	1.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	0.7	1.4
12	Chronic lower respiratory diseases (J40-J47)	339	0.6	1.2
13	Septicemia (A40-A41)	329	0.6	1.2
14	Viral hepatitis (B15-B19)	322	0.6	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	0.5	1.0
...	All other causes (Residual)	8,908	16.7	32.3
White, non-Hispanic, male, 45-64 years				
...	All causes	195,489	100.0	761.1
1	Malignant neoplasms (C00-C97)	59,627	30.5	232.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,375	27.3	207.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,060	6.7	50.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,179	3.7	28.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,035	3.6	27.4
6	Chronic lower respiratory diseases (J40-J47)	6,753	3.5	26.3
7	Diabetes mellitus (E10-E14)	6,163	3.2	24.0
8	Cerebrovascular diseases (I60-I69)	5,240	2.7	20.4
9	Influenza and pneumonia (J10-J18)	2,063	1.1	8.0
10	Septicemia (A40-A41)	1,978	1.0	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,793	0.9	7.0
12	Viral hepatitis (B15-B19)	1,607	0.8	6.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,600	0.8	6.2
14	Aortic aneurysm and dissection (I71)	1,325	0.7	5.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	948	0.5	3.7
...	All other causes (Residual)	25,743	13.2	100.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	682,034	100.0	5,486.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	214,615	31.5	1,726.4
2	Malignant neoplasms (C00-C97)	169,391	24.8	1,362.6
3	Chronic lower respiratory diseases (J40-J47)	46,408	6.8	373.3
4	Cerebrovascular diseases (I60-I69)	42,387	6.2	341.0
5	Influenza and pneumonia (J10-J18)	20,746	3.0	166.9
6	Diabetes mellitus (E10-E14)	18,699	2.7	150.4
7	Alzheimer's disease (G30)	16,185	2.4	130.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,552	2.1	117.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,174	1.9	106.0
10	Parkinson's disease (G20-G21)	9,089	1.3	73.1
11	Septicemia (A40-A41)	8,729	1.3	70.2
12	Pneumonitis due to solids and liquids (J69)	6,792	1.0	54.6
13	Aortic aneurysm and dissection (I71)	6,188	0.9	49.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,703	0.7	37.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,610	0.7	37.1
...	All other causes (Residual)	85,766	12.6	689.9
White, non-Hispanic, female, all ages^3				
...	All causes	1,023,271	100.0	1,007.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,567	28.3	285.1
2	Malignant neoplasms (C00-C97)	220,977	21.6	217.6
3	Cerebrovascular diseases (I60-I69)	79,420	7.8	78.2
4	Chronic lower respiratory diseases (J40-J47)	59,948	5.9	59.0
5	Alzheimer's disease (G30)	40,841	4.0	40.2
6	Influenza and pneumonia (J10-J18)	31,076	3.0	30.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,927	3.0	30.5
8	Diabetes mellitus (E10-E14)	26,954	2.6	26.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,174	1.6	15.9
10	Septicemia (A40-A41)	14,483	1.4	14.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,205	1.0	10.0
12	Pneumonitis due to solids and liquids (J69)	7,391	0.7	7.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,348	0.7	7.2
14	Atherosclerosis (I70)	7,294	0.7	7.2
15	Parkinson's disease (G20-G21)	7,196	0.7	7.1
...	All other causes (Residual)	173,470	17.0	170.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,116	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	33.6	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	12.2	3.0
3	Malignant neoplasms (C00-C97)	98	8.8	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.4	1.1
6	Influenza and pneumonia (J10-J18)	36	3.2	0.8
7	Cerebrovascular diseases (I60-I69)	15	1.3	*
8	Septicemia (A40-A41)	14	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Meningitis (G00,G03)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Meningococcal infection (A39)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	272	24.4	6.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,565	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	581	37.1	4.8
2	Malignant neoplasms (C00-C97)	289	18.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	6.7	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.7	0.5
5	Influenza and pneumonia (J10-J18)	47	3.0	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	2.4	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	1.9	0.2
8	Septicemia (A40-A41)	25	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
11	Cerebrovascular diseases (I60-I69)	14	0.9	*
12	Anemias (D50-D64)	8	0.5	*
13	Diabetes mellitus (E10-E14)	7	0.4	*
14	Pneumonitis due to solids and liquids (J69)	5	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	318	20.3	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,618	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,940	52.3	23.2
2	Malignant neoplasms (C00-C97)	432	7.7	3.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	420	7.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	242	4.3	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	237	4.2	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	2.0	0.9
7	Influenza and pneumonia (J10-J18)	62	1.1	0.5
8	Cerebrovascular diseases (I60-I69)	52	0.9	0.4
9	Pregnancy, childbirth and the puerperium (O00-099)	41	0.7	0.3
10	Septicemia (A40-A41)	38	0.7	0.3
11	Diabetes mellitus (E10-E14)	36	0.6	0.3
12	Chronic lower respiratory diseases (J40-J47)	34	0.6	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.4	0.2
14	Meningococcal infection (A39)	15	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
...	All other causes (Residual)	921	16.4	7.3
White, non-Hispanic, female, 25-44 years				
...	All causes	28,432	100.0	103.8
1	Malignant neoplasms (C00-C97)	7,056	24.8	25.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,632	19.8	20.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,041	10.7	11.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,979	7.0	7.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	762	2.7	2.8
6	Cerebrovascular diseases (I60-I69)	741	2.6	2.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	705	2.5	2.6
8	Diabetes mellitus (E10-E14)	600	2.1	2.2
9	Chronic lower respiratory diseases (J40-J47)	418	1.5	1.5
10	Influenza and pneumonia (J10-J18)	356	1.3	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	355	1.2	1.3
12	Septicemia (A40-A41)	339	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	330	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	0.7	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	153	0.5	0.6
...	All other causes (Residual)	5,766	20.3	21.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 45-64 years			
...	All causes	123,941	100.0	467.8
1	Malignant neoplasms (C00-C97)	52,274	42.2	197.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,698	16.7	78.1
3	Chronic lower respiratory diseases (J40-J47)	6,437	5.2	24.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,704	4.6	21.5
5	Cerebrovascular diseases (I60-I69)	4,347	3.5	16.4
6	Diabetes mellitus (E10-E14)	4,122	3.3	15.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,837	2.3	10.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,201	1.8	8.3
9	Septicemia (A40-A41)	1,788	1.4	6.7
10	Influenza and pneumonia (J10-J18)	1,571	1.3	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,381	1.1	5.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	678	0.5	2.6
13	Viral hepatitis (B15-B19)	601	0.5	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	565	0.5	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	540	0.4	2.0
...	All other causes (Residual)	18,197	14.7	68.7
	White, non-Hispanic, female, 65 years and over			
...	All causes	856,942	100.0	4,964.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	265,388	31.0	1,537.3
2	Malignant neoplasms (C00-C97)	160,800	18.8	931.5
3	Cerebrovascular diseases (I60-I69)	74,227	8.7	430.0
4	Chronic lower respiratory diseases (J40-J47)	53,032	6.2	307.2
5	Alzheimer's disease (G30)	40,522	4.7	234.7
6	Influenza and pneumonia (J10-J18)	28,947	3.4	167.7
7	Diabetes mellitus (E10-E14)	22,186	2.6	128.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,479	1.8	89.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,543	1.7	84.2
10	Septicemia (A40-A41)	12,229	1.4	70.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,557	1.1	55.4
12	Atherosclerosis (I70)	7,110	0.8	41.2
13	Parkinson's disease (G20-G21)	7,053	0.8	40.9
14	Pneumonitis due to solids and liquids (J69)	6,957	0.8	40.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,944	0.6	28.6
...	All other causes (Residual)	133,968	15.6	776.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, all ages ³			
...	All causes	287,968	100.0	788.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,452	26.5	209.4
2	Malignant neoplasms (C00-C97)	62,054	21.5	170.0
3	Cerebrovascular diseases (I60-I69)	18,649	6.5	51.1
4	Diabetes mellitus (E10-E14)	12,787	4.4	35.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,154	4.2	33.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,276	2.9	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,804	2.7	21.4
8	Chronic lower respiratory diseases (J40-J47)	7,632	2.7	20.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,351	2.6	20.1
10	Septicemia (A40-A41)	6,141	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,798	2.0	15.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,716	1.6	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,445	1.5	12.2
14	Alzheimer's disease (G30)	3,693	1.3	10.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,582	0.9	7.1
...	All other causes (Residual)	47,434	16.5	129.9
	Black, non-Hispanic, both sexes, 1-4 years			
...	All causes	1,169	100.0	48.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	28.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	155	13.3	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	10.1	4.8
4	Malignant neoplasms (C00-C97)	55	4.7	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.7	2.3
6	Certain conditions originating in the perinatal period (P00-P96)	29	2.5	1.2
7	Influenza and pneumonia (J10-J18)	27	2.3	1.1
8	Septicemia (A40-A41)	24	2.1	1.0
8	Chronic lower respiratory diseases (J40-J47)	24	2.1	1.0
10	Cerebrovascular diseases (I60-I69)	14	1.2	*
11	Anemias (D50-D64)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.8	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
15	Meningococcal infection (A39)	6	0.5	*
...	All other causes (Residual)	292	25.0	12.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,530	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	517	33.8	8.0
2	Malignant neoplasms (C00-C97)	173	11.3	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	9.6	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	5.0	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	4.8	1.1
6	Chronic lower respiratory diseases (J40-J47)	62	4.1	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	2.8	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	31	2.0	0.5
9	Influenza and pneumonia (J10-J18)	28	1.8	0.4
10	Anemias (D50-D64)	25	1.6	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.1	*
12	Cerebrovascular diseases (I60-I69)	16	1.0	*
13	Septicemia (A40-A41)	15	1.0	*
14	Diabetes mellitus (E10-E14)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	293	19.2	4.6
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,021	100.0	116.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,955	42.1	49.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,590	22.6	26.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	438	6.2	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	4.9	5.7
5	Malignant neoplasms (C00-C97)	245	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	117	1.7	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.3	1.5
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.3
9	Anemias (D50-D64)	78	1.1	1.3
10	Cerebrovascular diseases (I60-I69)	58	0.8	1.0
11	Diabetes mellitus (E10-E14)	45	0.6	0.7
12	Influenza and pneumonia (J10-J18)	42	0.6	0.7
12	Pregnancy, childbirth and the puerperium (O00-099)	42	0.6	0.7
14	Septicemia (A40-A41)	41	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.5	0.6
...	All other causes (Residual)	817	11.6	13.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	29,356	100.0	274.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,366	14.9	40.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,797	12.9	35.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,778	12.9	35.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	3,724	12.7	34.8
5	Malignant neoplasms (C00-C97)	3,443	11.7	32.2
6	Cerebrovascular diseases (I60-I69)	953	3.2	8.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	907	3.1	8.5
8	Diabetes mellitus (E10-E14)	755	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	473	1.6	4.4
10	Septicemia (A40-A41)	385	1.3	3.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	364	1.2	3.4
12	Influenza and pneumonia (J10-J18)	352	1.2	3.3
13	Chronic lower respiratory diseases (J40-J47)	349	1.2	3.3
14	Anemias (D50-D64)	247	0.8	2.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	225	0.8	2.1
...	All other causes (Residual)	5,238	17.8	48.9
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	78,934	100.0	1,077.7
1	Malignant neoplasms (C00-C97)	21,705	27.5	296.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,289	25.7	277.0
3	Cerebrovascular diseases (I60-I69)	4,398	5.6	60.0
4	Diabetes mellitus (E10-E14)	3,778	4.8	51.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,441	4.4	47.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,184	4.0	43.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,952	2.5	26.7
8	Chronic lower respiratory diseases (J40-J47)	1,706	2.2	23.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,636	2.1	22.3
10	Septicemia (A40-A41)	1,531	1.9	20.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,105	1.4	15.1
12	Influenza and pneumonia (J10-J18)	1,052	1.3	14.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	949	1.2	13.0
14	Viral hepatitis (B15-B19)	573	0.7	7.8
15	Aortic aneurysm and dissection (I71)	398	0.5	5.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	398	0.5	5.4
...	All other causes (Residual)	10,839	13.7	148.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 65 years and over			
...	All causes	161,779	100.0	5,463.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,189	31.6	1,728.7
2	Malignant neoplasms (C00-C97)	36,426	22.5	1,230.2
3	Cerebrovascular diseases (I60-I69)	13,179	8.1	445.1
4	Diabetes mellitus (E10-E14)	8,199	5.1	276.9
5	Chronic lower respiratory diseases (J40-J47)	5,403	3.3	182.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,274	3.3	178.1
7	Influenza and pneumonia (J10-J18)	4,205	2.6	142.0
8	Septicemia (A40-A41)	4,025	2.5	135.9
9	Alzheimer's disease (G30)	3,660	2.3	123.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,103	1.9	104.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,192	1.4	74.0
12	Pneumonitis due to solids and liquids (J69)	1,321	0.8	44.6
13	Atherosclerosis (I70)	861	0.5	29.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	704	0.4	23.8
15	Aortic aneurysm and dissection (I71)	693	0.4	23.4
...	All other causes (Residual)	21,345	13.2	720.9
	Black, non-Hispanic, male, all ages^3			
...	All causes	146,136	100.0	840.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,962	25.3	212.6
2	Malignant neoplasms (C00-C97)	32,131	22.0	184.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,237	5.6	47.4
4	Cerebrovascular diseases (I60-I69)	7,798	5.3	44.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,980	4.8	40.1
6	Diabetes mellitus (E10-E14)	5,380	3.7	30.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,947	3.4	28.5
8	Chronic lower respiratory diseases (J40-J47)	4,202	2.9	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,614	2.5	20.8
10	Influenza and pneumonia (J10-J18)	2,861	2.0	16.5
11	Septicemia (A40-A41)	2,748	1.9	15.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,687	1.8	15.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,875	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,696	1.2	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,566	1.1	9.0
...	All other causes (Residual)	22,452	15.4	129.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	28.9	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	14.1	7.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	9.1	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.6	2.5
5	Malignant neoplasms (C00-C97)	30	4.4	2.4
6	Chronic lower respiratory diseases (J40-J47)	19	2.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.3	*
8	Septicemia (A40-A41)	14	2.1	*
9	Influenza and pneumonia (J10-J18)	13	1.9	*
10	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
12	Anemias (D50-D64)	7	1.0	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.7	*
15	Meningococcal infection (A39)	3	0.4	*
...	All other causes (Residual)	165	24.2	13.4
Black, non-Hispanic, male, 5-14 years				
...	All causes	911	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	36.2	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	10.8	3.0
3	Malignant neoplasms (C00-C97)	96	10.5	2.9
4	Chronic lower respiratory diseases (J40-J47)	43	4.7	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	4.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	4.2	1.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.7	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.9	*
9	Influenza and pneumonia (J10-J18)	15	1.6	*
10	Anemias (D50-D64)	12	1.3	*
11	Cerebrovascular diseases (I60-I69)	9	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
13	Septicemia (A40-A41)	6	0.7	*
14	Diseases of appendix (K35-K38)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
...	All other causes (Residual)	156	17.1	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,343	100.0	176.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,642	49.4	87.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	22.6	39.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	377	7.1	12.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	4.2	7.4
5	Malignant neoplasms (C00-C97)	141	2.6	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.1	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.0	1.8
8	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.6
9	Anemias (D50-D64)	39	0.7	1.3
10	Legal intervention (Y35,Y89.0)	31	0.6	1.0
11	Diabetes mellitus (E10-E14)	26	0.5	0.9
12	Cerebrovascular diseases (I60-I69)	25	0.5	0.8
13	Influenza and pneumonia (J10-J18)	24	0.4	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.7
15	Septicemia (A40-A41)	19	0.4	*
...	All other causes (Residual)	409	7.7	13.5
Black, non-Hispanic, male, 25-44 years				
...	All causes	17,997	100.0	355.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,206	17.8	63.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,736	15.2	54.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,638	14.7	52.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,351	13.1	46.4
5	Malignant neoplasms (C00-C97)	1,375	7.6	27.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	748	4.2	14.8
7	Cerebrovascular diseases (I60-I69)	471	2.6	9.3
8	Diabetes mellitus (E10-E14)	431	2.4	8.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	1.4	5.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	213	1.2	4.2
11	Influenza and pneumonia (J10-J18)	200	1.1	3.9
12	Septicemia (A40-A41)	190	1.1	3.8
13	Chronic lower respiratory diseases (J40-J47)	178	1.0	3.5
14	Anemias (D50-D64)	126	0.7	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	125	0.7	2.5
...	All other causes (Residual)	2,754	15.3	54.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	46,176	100.0	1,383.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,483	27.0	373.9
2	Malignant neoplasms (C00-C97)	11,668	25.3	349.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,449	5.3	73.4
4	Cerebrovascular diseases (I60-I69)	2,394	5.2	71.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,307	5.0	69.1
6	Diabetes mellitus (E10-E14)	1,910	4.1	57.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,156	2.5	34.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,094	2.4	32.8
9	Chronic lower respiratory diseases (J40-J47)	908	2.0	27.2
10	Septicemia (A40-A41)	800	1.7	24.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	754	1.6	22.6
12	Influenza and pneumonia (J10-J18)	641	1.4	19.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	637	1.4	19.1
14	Viral hepatitis (B15-B19)	395	0.9	11.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	0.7	9.0
...	All other causes (Residual)	6,279	13.6	188.1
Black, non-Hispanic, male, 65 years and over				
...	All causes	70,410	100.0	6,218.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,462	30.5	1,895.6
2	Malignant neoplasms (C00-C97)	18,817	26.7	1,662.0
3	Cerebrovascular diseases (I60-I69)	4,871	6.9	430.2
4	Diabetes mellitus (E10-E14)	3,009	4.3	265.8
5	Chronic lower respiratory diseases (J40-J47)	3,004	4.3	265.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,209	3.1	195.1
7	Influenza and pneumonia (J10-J18)	1,915	2.7	169.1
8	Septicemia (A40-A41)	1,648	2.3	145.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,161	1.6	102.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,106	1.6	97.7
11	Alzheimer's disease (G30)	1,060	1.5	93.6
12	Pneumonitis due to solids and liquids (J69)	615	0.9	54.3
13	Parkinson's disease (G20-G21)	324	0.5	28.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	324	0.5	28.6
15	Atherosclerosis (I70)	322	0.5	28.4
...	All other causes (Residual)	8,563	12.2	756.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, all ages^3			
...	All causes	141,832	100.0	741.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,490	27.8	206.5
2	Malignant neoplasms (C00-C97)	29,923	21.1	156.5
3	Cerebrovascular diseases (I60-I69)	10,851	7.7	56.7
4	Diabetes mellitus (E10-E14)	7,407	5.2	38.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,190	3.0	21.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,917	2.8	20.5
7	Chronic lower respiratory diseases (J40-J47)	3,430	2.4	17.9
8	Septicemia (A40-A41)	3,393	2.4	17.7
9	Influenza and pneumonia (J10-J18)	2,937	2.1	15.4
10	Alzheimer's disease (G30)	2,614	1.8	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,570	1.8	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,404	1.7	12.6
13	Certain conditions originating in the perinatal period (P00-P96)	2,029	1.4	10.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,296	0.9	6.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	886	0.6	4.6
...	All other causes (Residual)	24,495	17.3	128.1
	Black, non-Hispanic, female, 1-4 years			
...	All causes	488	100.0	40.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	27.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	12.1	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	11.5	4.7
4	Malignant neoplasms (C00-C97)	25	5.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.9	2.0
6	Influenza and pneumonia (J10-J18)	14	2.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.7	*
8	Septicemia (A40-A41)	10	2.0	*
9	Anemias (D50-D64)	5	1.0	*
9	Cerebrovascular diseases (I60-I69)	5	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Meningococcal infection (A39)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
15	Hernia (K40-K46)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	120	24.6	10.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	619	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	30.2	5.9
2	Malignant neoplasms (C00-C97)	77	12.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.9	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.5	1.1
6	Chronic lower respiratory diseases (J40-J47)	19	3.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.3	*
8	Anemias (D50-D64)	13	2.1	*
8	Influenza and pneumonia (J10-J18)	13	2.1	*
10	Septicemia (A40-A41)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.5	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.5	*
13	Cerebrovascular diseases (I60-I69)	7	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Diabetes mellitus (E10-E14)	4	0.6	*
...	All other causes (Residual)	135	21.8	4.3
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,678	100.0	56.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	22.9	12.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	18.7	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	7.2	4.0
4	Malignant neoplasms (C00-C97)	104	6.2	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	3.6	2.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	2.0
7	Pregnancy, childbirth and the puerperium (O00-099)	42	2.5	1.4
8	Anemias (D50-D64)	39	2.3	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	2.2	1.2
10	Cerebrovascular diseases (I60-I69)	33	2.0	1.1
10	Chronic lower respiratory diseases (J40-J47)	33	2.0	1.1
12	Septicemia (A40-A41)	22	1.3	0.7
13	Diabetes mellitus (E10-E14)	19	1.1	*
14	Influenza and pneumonia (J10-J18)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
...	All other causes (Residual)	377	22.5	12.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-44 years				
...	All causes	11,359	100.0	201.3
1	Malignant neoplasms (C00-C97)	2,068	18.2	36.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,728	15.2	30.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,373	12.1	24.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,061	9.3	18.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	572	5.0	10.1
6	Cerebrovascular diseases (I60-I69)	482	4.2	8.5
7	Diabetes mellitus (E10-E14)	324	2.9	5.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	1.9	3.9
9	Septicemia (A40-A41)	195	1.7	3.5
10	Chronic lower respiratory diseases (J40-J47)	171	1.5	3.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	1.4	2.8
12	Influenza and pneumonia (J10-J18)	152	1.3	2.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	151	1.3	2.7
14	Pregnancy, childbirth and the puerperium (O00-099)	140	1.2	2.5
15	Anemias (D50-D64)	121	1.1	2.1
...	All other causes (Residual)	2,444	21.5	43.3
Black, non-Hispanic, female, 45-64 years				
...	All causes	32,758	100.0	821.9
1	Malignant neoplasms (C00-C97)	10,037	30.6	251.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,806	23.8	195.9
3	Cerebrovascular diseases (I60-I69)	2,004	6.1	50.3
4	Diabetes mellitus (E10-E14)	1,868	5.7	46.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	992	3.0	24.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	877	2.7	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	858	2.6	21.5
8	Chronic lower respiratory diseases (J40-J47)	798	2.4	20.0
9	Septicemia (A40-A41)	731	2.2	18.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	480	1.5	12.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	468	1.4	11.7
12	Influenza and pneumonia (J10-J18)	411	1.3	10.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	0.6	4.9
14	Viral hepatitis (B15-B19)	178	0.5	4.5
15	Aortic aneurysm and dissection (I71)	126	0.4	3.2
...	All other causes (Residual)	4,929	15.0	123.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	91,369	100.0	4,996.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,727	32.5	1,625.4
2	Malignant neoplasms (C00-C97)	17,609	19.3	962.8
3	Cerebrovascular diseases (I60-I69)	8,308	9.1	454.3
4	Diabetes mellitus (E10-E14)	5,190	5.7	283.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,065	3.4	167.6
6	Alzheimer's disease (G30)	2,600	2.8	142.2
7	Chronic lower respiratory diseases (J40-J47)	2,399	2.6	131.2
8	Septicemia (A40-A41)	2,377	2.6	130.0
9	Influenza and pneumonia (J10-J18)	2,290	2.5	125.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,997	2.2	109.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,031	1.1	56.4
12	Pneumonitis due to solids and liquids (J69)	706	0.8	38.6
13	Atherosclerosis (I70)	539	0.6	29.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	399	0.4	21.8
15	Aortic aneurysm and dissection (I71)	382	0.4	20.9
...	All other causes (Residual)	12,750	14.0	697.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable