

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	131,161	100.0	307.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,555	22.5	69.2
2	Malignant neoplasms (C00-C97)	26,156	19.9	61.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,464	8.7	26.9
4	Cerebrovascular diseases (I60-I69)	6,830	5.2	16.0
5	Diabetes mellitus (E10-E14)	6,665	5.1	15.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,555	2.7	8.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,520	2.7	8.2
8	Chronic lower respiratory diseases (J40-J47)	3,457	2.6	8.1
9	Influenza and pneumonia (J10-J18)	3,085	2.4	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,816	2.1	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,329	1.8	5.5
12	Alzheimer's disease (G30)	2,201	1.7	5.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,188	1.7	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,896	1.4	4.4
15	Septicemia (A40-A41)	1,704	1.3	4.0
...	All other causes (Residual)	23,740	18.1	55.6
Hispanic, both sexes, 1-4 years				
...	All causes	1,041	100.0	28.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	36.5	10.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	11.4	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	7.9	2.3
4	Malignant neoplasms (C00-C97)	80	7.7	2.2
5	Influenza and pneumonia (J10-J18)	32	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.2	0.6
7	Cerebrovascular diseases (I60-I69)	15	1.4	*
8	Septicemia (A40-A41)	14	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Meningitis (G00,G03)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	252	24.2	7.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,068	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	370	34.6	4.7
2	Malignant neoplasms (C00-C97)	194	18.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	8.0	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.8	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	3.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.2	0.4
7	Cerebrovascular diseases (I60-I69)	19	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
9	Septicemia (A40-A41)	12	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
11	Influenza and pneumonia (J10-J18)	9	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
13	Anemias (D50-D64)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	194	18.2	2.5
Hispanic, both sexes, 15-24 years				
...	All causes	5,837	100.0	81.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	45.7	37.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,307	22.4	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	553	9.5	7.7
4	Malignant neoplasms (C00-C97)	323	5.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	2.3	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	1.3	1.0
7	Cerebrovascular diseases (I60-I69)	44	0.8	0.6
8	Pregnancy, childbirth and the puerperium (O00-099)	35	0.6	0.5
9	Legal intervention (Y35,Y89.0)	32	0.5	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.5	0.4
11	Septicemia (A40-A41)	18	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
13	Diabetes mellitus (E10-E14)	14	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
15	Influenza and pneumonia (J10-J18)	11	0.2	*
...	All other causes (Residual)	560	9.6	7.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,287	100.0	107.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,222	27.6	29.7
2	Malignant neoplasms (C00-C97)	1,945	12.7	13.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,605	10.5	11.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,311	8.6	9.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,006	6.6	7.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	808	5.3	5.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	545	3.6	3.8
8	Cerebrovascular diseases (I60-I69)	378	2.5	2.7
9	Diabetes mellitus (E10-E14)	285	1.9	2.0
10	Influenza and pneumonia (J10-J18)	133	0.9	0.9
11	Viral hepatitis (B15-B19)	124	0.8	0.9
12	Septicemia (A40-A41)	110	0.7	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	92	0.6	0.6
...	All other causes (Residual)	2,526	16.5	17.8
Hispanic, both sexes, 45-64 years				
...	All causes	29,658	100.0	446.1
1	Malignant neoplasms (C00-C97)	8,160	27.5	122.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,875	19.8	88.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,175	7.3	32.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,880	6.3	28.3
5	Diabetes mellitus (E10-E14)	1,800	6.1	27.1
6	Cerebrovascular diseases (I60-I69)	1,434	4.8	21.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	728	2.5	11.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	516	1.7	7.8
9	Viral hepatitis (B15-B19)	495	1.7	7.4
10	Chronic lower respiratory diseases (J40-J47)	449	1.5	6.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	1.4	6.4
12	Influenza and pneumonia (J10-J18)	419	1.4	6.3
13	Septicemia (A40-A41)	387	1.3	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	1.1	5.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	232	0.8	3.5
...	All other causes (Residual)	4,348	14.7	65.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	72,490	100.0	3,173.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,099	30.5	967.4
2	Malignant neoplasms (C00-C97)	15,434	21.3	675.7
3	Cerebrovascular diseases (I60-I69)	4,916	6.8	215.2
4	Diabetes mellitus (E10-E14)	4,563	6.3	199.8
5	Chronic lower respiratory diseases (J40-J47)	2,891	4.0	126.6
6	Influenza and pneumonia (J10-J18)	2,415	3.3	105.7
7	Alzheimer's disease (G30)	2,160	3.0	94.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,657	2.3	72.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,460	2.0	63.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,120	1.5	49.0
11	Septicemia (A40-A41)	1,097	1.5	48.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,028	1.4	45.0
13	Parkinson's disease (G20-G21)	695	1.0	30.4
14	Pneumonitis due to solids and liquids (J69)	504	0.7	22.1
15	Atherosclerosis (I70)	366	0.5	16.0
...	All other causes (Residual)	10,085	13.9	441.5
Hispanic, male, all ages				
...	All causes	73,788	100.0	334.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,900	21.5	72.1
2	Malignant neoplasms (C00-C97)	13,896	18.8	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,612	11.7	39.0
4	Diabetes mellitus (E10-E14)	3,296	4.5	14.9
5	Cerebrovascular diseases (I60-I69)	3,188	4.3	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,008	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,561	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,841	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,839	2.5	8.3
10	Certain conditions originating in the perinatal period (P00-P96)	1,612	2.2	7.3
11	Influenza and pneumonia (J10-J18)	1,506	2.0	6.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,309	1.8	5.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,186	1.6	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	996	1.3	4.5
15	Septicemia (A40-A41)	838	1.1	3.8
...	All other causes (Residual)	12,200	16.5	55.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	610	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	40.0	13.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	10.7	3.5
3	Malignant neoplasms (C00-C97)	52	8.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.8	*
6	Influenza and pneumonia (J10-J18)	15	2.5	*
7	Cerebrovascular diseases (I60-I69)	8	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
9	Septicemia (A40-A41)	6	1.0	*
10	Meningococcal infection (A39)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	131	21.5	7.1
Hispanic, male, 5-14 years				
...	All causes	612	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	36.3	5.6
2	Malignant neoplasms (C00-C97)	99	16.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	7.7	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.5	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.4	0.5
7	Cerebrovascular diseases (I60-I69)	9	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
9	Influenza and pneumonia (J10-J18)	7	1.1	*
11	Septicemia (A40-A41)	6	1.0	*
12	Anemias (D50-D64)	4	0.7	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	106	17.3	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,604	100.0	120.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,133	46.3	55.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,187	25.8	31.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	462	10.0	12.1
4	Malignant neoplasms (C00-C97)	202	4.4	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	2.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.9	1.1
7	Legal intervention (Y35,Y89.0)	32	0.7	0.8
8	Cerebrovascular diseases (I60-I69)	25	0.5	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.6
10	Septicemia (A40-A41)	10	0.2	*
11	Diabetes mellitus (E10-E14)	9	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
12	Influenza and pneumonia (J10-J18)	8	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.2	*
...	All other causes (Residual)	356	7.7	9.3
Hispanic, male, 25-44 years				
...	All causes	11,105	100.0	144.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,443	31.0	44.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,384	12.5	18.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,005	9.0	13.1
4	Malignant neoplasms (C00-C97)	881	7.9	11.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	860	7.7	11.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	638	5.7	8.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	431	3.9	5.6
8	Cerebrovascular diseases (I60-I69)	227	2.0	3.0
9	Diabetes mellitus (E10-E14)	193	1.7	2.5
10	Viral hepatitis (B15-B19)	92	0.8	1.2
11	Influenza and pneumonia (J10-J18)	85	0.8	1.1
12	Septicemia (A40-A41)	58	0.5	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.5	0.7
15	Chronic lower respiratory diseases (J40-J47)	50	0.5	0.7
...	All other causes (Residual)	1,649	14.8	21.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	18,948	100.0	575.7
1	Malignant neoplasms (C00-C97)	4,328	22.8	131.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,171	22.0	126.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,671	8.8	50.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,478	7.8	44.9
5	Diabetes mellitus (E10-E14)	1,040	5.5	31.6
6	Cerebrovascular diseases (I60-I69)	842	4.4	25.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	591	3.1	18.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	353	1.9	10.7
9	Viral hepatitis (B15-B19)	352	1.9	10.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	1.7	9.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	1.4	8.2
12	Influenza and pneumonia (J10-J18)	257	1.4	7.8
12	Chronic lower respiratory diseases (J40-J47)	257	1.4	7.8
14	Septicemia (A40-A41)	214	1.1	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	115	0.6	3.5
...	All other causes (Residual)	2,695	14.2	81.9
Hispanic, male, 65 years and over				
...	All causes	34,666	100.0	3,573.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,552	30.4	1,087.6
2	Malignant neoplasms (C00-C97)	8,328	24.0	858.4
3	Cerebrovascular diseases (I60-I69)	2,063	6.0	212.6
4	Diabetes mellitus (E10-E14)	2,052	5.9	211.5
5	Chronic lower respiratory diseases (J40-J47)	1,505	4.3	155.1
6	Influenza and pneumonia (J10-J18)	1,097	3.2	113.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	2.3	82.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	790	2.3	81.4
9	Alzheimer's disease (G30)	750	2.2	77.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	643	1.9	66.3
11	Septicemia (A40-A41)	507	1.5	52.3
12	Parkinson's disease (G20-G21)	404	1.2	41.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	396	1.1	40.8
14	Pneumonitis due to solids and liquids (J69)	256	0.7	26.4
15	Aortic aneurysm and dissection (I71)	190	0.5	19.6
...	All other causes (Residual)	4,335	12.5	446.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, all ages			
...	All causes	57,373	100.0	278.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,655	23.8	66.2
2	Malignant neoplasms (C00-C97)	12,260	21.4	59.5
3	Cerebrovascular diseases (I60-I69)	3,642	6.3	17.7
4	Diabetes mellitus (E10-E14)	3,369	5.9	16.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,852	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,618	2.8	7.8
7	Influenza and pneumonia (J10-J18)	1,579	2.8	7.7
8	Alzheimer's disease (G30)	1,437	2.5	7.0
9	Certain conditions originating in the perinatal period (P00-P96)	1,204	2.1	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,143	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	994	1.7	4.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	900	1.6	4.4
13	Septicemia (A40-A41)	866	1.5	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	767	1.3	3.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	512	0.9	2.5
...	All other causes (Residual)	10,575	18.4	51.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	431	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	31.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	12.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	8.4	2.0
4	Malignant neoplasms (C00-C97)	28	6.5	1.6
5	Influenza and pneumonia (J10-J18)	17	3.9	*
6	Septicemia (A40-A41)	8	1.9	*
7	Cerebrovascular diseases (I60-I69)	7	1.6	*
8	Meningitis (G00,G03)	6	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
12	Anemias (D50-D64)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
15	Shigellosis and amebiasis (A03,A06)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	108	25.1	6.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	456	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	32.5	3.9
2	Malignant neoplasms (C00-C97)	95	20.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	8.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.9	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.6	*
7	Cerebrovascular diseases (I60-I69)	10	2.2	*
8	Septicemia (A40-A41)	6	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
13	Meningitis (G00,G03)	2	0.4	*
13	Influenza and pneumonia (J10-J18)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	90	19.7	2.4
Hispanic, female, 15-24 years				
...	All causes	1,233	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	43.6	15.9
2	Malignant neoplasms (C00-C97)	121	9.8	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	9.7	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.4	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	3.7	1.4
6	Pregnancy, childbirth and the puerperium (O00-099)	35	2.8	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.5	0.9
8	Cerebrovascular diseases (I60-I69)	19	1.5	*
9	Septicemia (A40-A41)	8	0.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
12	Diabetes mellitus (E10-E14)	5	0.4	*
14	Tuberculosis (A16-A19)	3	0.2	*
14	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Influenza and pneumonia (J10-J18)	3	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (Residual)	188	15.2	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,182	100.0	63.9
1	Malignant neoplasms (C00-C97)	1,064	25.4	16.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	779	18.6	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	7.3	4.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	221	5.3	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	170	4.1	2.6
6	Cerebrovascular diseases (I60-I69)	151	3.6	2.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	3.5	2.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	114	2.7	1.7
9	Diabetes mellitus (E10-E14)	92	2.2	1.4
9	Pregnancy, childbirth and the puerperium (O00-099)	92	2.2	1.4
11	Septicemia (A40-A41)	52	1.2	0.8
12	Influenza and pneumonia (J10-J18)	48	1.1	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.1	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	1.0	0.7
15	Viral hepatitis (B15-B19)	32	0.8	0.5
...	All other causes (Residual)	827	19.8	12.6
Hispanic, female, 45-64 years				
...	All causes	10,710	100.0	319.1
1	Malignant neoplasms (C00-C97)	3,832	35.8	114.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,704	15.9	50.8
3	Diabetes mellitus (E10-E14)	760	7.1	22.6
4	Cerebrovascular diseases (I60-I69)	592	5.5	17.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	504	4.7	15.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	402	3.8	12.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	1.9	6.0
8	Chronic lower respiratory diseases (J40-J47)	192	1.8	5.7
9	Septicemia (A40-A41)	173	1.6	5.2
10	Influenza and pneumonia (J10-J18)	162	1.5	4.8
11	Viral hepatitis (B15-B19)	143	1.3	4.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	137	1.3	4.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.1	3.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	0.7	2.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	0.6	1.9
...	All other causes (Residual)	1,653	15.4	49.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	37,824	100.0	2,878.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,547	30.5	878.7
2	Malignant neoplasms (C00-C97)	7,106	18.8	540.8
3	Cerebrovascular diseases (I60-I69)	2,853	7.5	217.1
4	Diabetes mellitus (E10-E14)	2,511	6.6	191.1
5	Alzheimer's disease (G30)	1,410	3.7	107.3
6	Chronic lower respiratory diseases (J40-J47)	1,386	3.7	105.5
7	Influenza and pneumonia (J10-J18)	1,318	3.5	100.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	867	2.3	66.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	662	1.8	50.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	632	1.7	48.1
11	Septicemia (A40-A41)	590	1.6	44.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	477	1.3	36.3
13	Parkinson's disease (G20-G21)	291	0.8	22.1
14	Pneumonitis due to solids and liquids (J69)	248	0.7	18.9
15	Atherosclerosis (I70)	207	0.5	15.8
...	All other causes (Residual)	5,719	15.1	435.2
White, non-Hispanic, both sexes, all ages				
...	All causes	967,142	100.0	981.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	535,101	27.2	267.1
2	Malignant neoplasms (C00-C97)	456,133	23.2	227.7
3	Chronic lower respiratory diseases (J40-J47)	117,337	6.0	58.6
4	Cerebrovascular diseases (I60-I69)	115,074	5.8	57.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88,970	4.5	44.4
6	Alzheimer's disease (G30)	63,958	3.3	31.9
7	Diabetes mellitus (E10-E14)	53,159	2.7	26.5
8	Influenza and pneumonia (J10-J18)	52,431	2.7	26.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32,492	1.7	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,305	1.4	13.6
11	Septicemia (A40-A41)	25,496	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,428	1.0	10.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17,949	0.9	9.0
14	Parkinson's disease (G20-G21)	17,767	0.9	8.9
15	Pneumonitis due to solids and liquids (J69)	14,739	0.7	7.4
...	All other causes (Residual)	328,803	16.7	164.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,404	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	37.4	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.2	2.9
3	Malignant neoplasms (C00-C97)	215	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	5.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	2.9	0.8
6	Influenza and pneumonia (J10-J18)	46	1.9	0.5
7	Septicemia (A40-A41)	44	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	30	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.8	0.2
12	Meningitis (G00,G03)	12	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	573	23.8	6.2
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,681	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,423	38.7	5.9
2	Malignant neoplasms (C00-C97)	597	16.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	5.6	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	4.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.5	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	3.0	0.5
7	Influenza and pneumonia (J10-J18)	67	1.8	0.3
8	Septicemia (A40-A41)	47	1.3	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	45	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.1
12	Diabetes mellitus (E10-E14)	22	0.6	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	19	0.5	*
14	Meningitis (G00,G03)	16	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	739	20.1	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,683	100.0	74.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,711	54.4	40.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,992	15.2	11.4
3	Malignant neoplasms (C00-C97)	1,048	5.3	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	895	4.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	604	3.1	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	313	1.6	1.2
7	Influenza and pneumonia (J10-J18)	123	0.6	0.5
8	Diabetes mellitus (E10-E14)	110	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	92	0.5	0.3
10	Septicemia (A40-A41)	77	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	76	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	64	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	31	0.2	0.1
...	All other causes (Residual)	2,439	12.4	9.3
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	78,612	100.0	146.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,599	27.5	40.3
2	Malignant neoplasms (C00-C97)	12,151	15.5	22.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,895	12.6	18.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,142	11.6	17.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,072	2.6	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,970	2.5	3.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,629	2.1	3.0
8	Diabetes mellitus (E10-E14)	1,521	1.9	2.8
9	Cerebrovascular diseases (I60-I69)	1,432	1.8	2.7
10	Influenza and pneumonia (J10-J18)	817	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	722	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	668	0.8	1.2
13	Septicemia (A40-A41)	637	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	456	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	359	0.5	0.7
...	All other causes (Residual)	13,542	17.2	25.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	332,747	100.0	608.2
1	Malignant neoplasms (C00-C97)	114,050	34.3	208.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,879	22.2	135.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,960	6.6	40.1
4	Chronic lower respiratory diseases (J40-J47)	14,112	4.2	25.8
5	Diabetes mellitus (E10-E14)	10,747	3.2	19.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10,581	3.2	19.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,030	3.0	18.3
8	Cerebrovascular diseases (I60-I69)	9,812	2.9	17.9
9	Septicemia (A40-A41)	3,983	1.2	7.3
10	Influenza and pneumonia (J10-J18)	3,906	1.2	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,376	1.0	6.2
12	Viral hepatitis (B15-B19)	2,460	0.7	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,922	0.6	3.5
14	Aortic aneurysm and dissection (I71)	1,621	0.5	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,582	0.5	2.9
...	All other causes (Residual)	48,726	14.6	89.1
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	516,916	100.0	5,041.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	450,345	29.7	1,496.7
2	Malignant neoplasms (C00-C97)	328,024	21.6	1,090.1
3	Cerebrovascular diseases (I60-I69)	103,607	6.8	344.3
4	Chronic lower respiratory diseases (J40-J47)	102,376	6.7	340.2
5	Alzheimer's disease (G30)	63,323	4.2	210.4
6	Influenza and pneumonia (J10-J18)	47,377	3.1	157.5
7	Diabetes mellitus (E10-E14)	40,757	2.7	135.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,794	2.1	105.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28,531	1.9	94.8
10	Septicemia (A40-A41)	20,608	1.4	68.5
11	Parkinson's disease (G20-G21)	17,331	1.1	57.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16,181	1.1	53.8
13	Pneumonitis due to solids and liquids (J69)	13,511	0.9	44.9
14	Atherosclerosis (I70)	9,927	0.7	33.0
15	Aortic aneurysm and dissection (I71)	9,796	0.6	32.6
...	All other causes (Residual)	233,428	15.4	775.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	954,402	100.0	970.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,311	27.6	267.8
2	Malignant neoplasms (C00-C97)	236,659	24.8	240.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,021	5.9	57.0
4	Chronic lower respiratory diseases (J40-J47)	55,015	5.8	56.0
5	Cerebrovascular diseases (I60-I69)	44,038	4.6	44.8
6	Diabetes mellitus (E10-E14)	26,365	2.8	26.8
7	Influenza and pneumonia (J10-J18)	22,903	2.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,605	2.3	22.0
9	Alzheimer's disease (G30)	18,208	1.9	18.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,953	1.7	16.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,088	1.4	13.3
12	Septicemia (A40-A41)	11,337	1.2	11.5
13	Parkinson's disease (G20-G21)	10,232	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,430	0.8	7.6
15	Aortic aneurysm and dissection (I71)	6,877	0.7	7.0
...	All other causes (Residual)	145,360	15.2	147.8
White, non-Hispanic, male, 1-4 years				
...	All causes	1,408	100.0	29.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	39.6	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	9.9	3.0
3	Malignant neoplasms (C00-C97)	118	8.4	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.8
6	Septicemia (A40-A41)	26	1.8	0.6
7	Influenza and pneumonia (J10-J18)	23	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.4	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	18	1.3	*
11	Meningitis (G00,G03)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
...	All other causes (Residual)	335	23.8	7.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,163	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	886	41.0	7.1
2	Malignant neoplasms (C00-C97)	333	15.4	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	6.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	4.7	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.2	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	2.8	0.5
7	Septicemia (A40-A41)	29	1.3	0.2
7	Influenza and pneumonia (J10-J18)	29	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	26	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Diabetes mellitus (E10-E14)	9	0.4	*
12	Meningitis (G00,G03)	9	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	422	19.5	3.4
White, non-Hispanic, male, 15-24 years				
...	All causes	14,267	100.0	105.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,942	55.7	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,487	17.4	18.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	639	4.5	4.7
4	Malignant neoplasms (C00-C97)	628	4.4	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	3.0	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	1.5	1.6
7	Influenza and pneumonia (J10-J18)	82	0.6	0.6
8	Diabetes mellitus (E10-E14)	64	0.4	0.5
9	Septicemia (A40-A41)	46	0.3	0.3
10	Cerebrovascular diseases (I60-I69)	40	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.3	0.3
13	Legal intervention (Y35,Y89.0)	26	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
15	Aortic aneurysm and dissection (I71)	17	0.1	*
...	All other causes (Residual)	1,561	10.9	11.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	51,085	100.0	189.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,551	30.4	57.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,159	14.0	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,126	13.9	26.5
4	Malignant neoplasms (C00-C97)	5,583	10.9	20.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,374	2.7	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,317	2.6	4.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,259	2.5	4.7
8	Diabetes mellitus (E10-E14)	949	1.9	3.5
9	Cerebrovascular diseases (I60-I69)	728	1.4	2.7
10	Influenza and pneumonia (J10-J18)	451	0.9	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	358	0.7	1.3
12	Chronic lower respiratory diseases (J40-J47)	331	0.6	1.2
13	Septicemia (A40-A41)	315	0.6	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	268	0.5	1.0
15	Viral hepatitis (B15-B19)	225	0.4	0.8
...	All other causes (Residual)	8,091	15.8	30.1
White, non-Hispanic, male, 45-64 years				
...	All causes	204,089	100.0	757.3
1	Malignant neoplasms (C00-C97)	60,702	29.7	225.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,196	26.1	197.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,211	7.5	56.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,609	3.7	28.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,537	3.7	28.0
6	Chronic lower respiratory diseases (J40-J47)	7,191	3.5	26.7
7	Diabetes mellitus (E10-E14)	6,563	3.2	24.4
8	Cerebrovascular diseases (I60-I69)	5,402	2.6	20.0
9	Influenza and pneumonia (J10-J18)	2,246	1.1	8.3
10	Septicemia (A40-A41)	2,124	1.0	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,001	1.0	7.4
12	Viral hepatitis (B15-B19)	1,817	0.9	6.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,618	0.8	6.0
14	Aortic aneurysm and dissection (I71)	1,219	0.6	4.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,019	0.5	3.8
...	All other causes (Residual)	28,634	14.0	106.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	673,920	100.0	5,312.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	202,357	30.0	1,595.2
2	Malignant neoplasms (C00-C97)	169,262	25.1	1,334.3
3	Chronic lower respiratory diseases (J40-J47)	47,420	7.0	373.8
4	Cerebrovascular diseases (I60-I69)	37,789	5.6	297.9
5	Influenza and pneumonia (J10-J18)	20,011	3.0	157.7
6	Diabetes mellitus (E10-E14)	18,780	2.8	148.0
7	Alzheimer's disease (G30)	17,925	2.7	141.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,557	2.3	122.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,616	2.0	107.3
10	Parkinson's disease (G20-G21)	9,949	1.5	78.4
11	Septicemia (A40-A41)	8,734	1.3	68.8
12	Pneumonitis due to solids and liquids (J69)	6,719	1.0	53.0
13	Aortic aneurysm and dissection (I71)	5,454	0.8	43.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,344	0.8	42.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,676	0.7	36.9
...	All other causes (Residual)	90,327	13.4	712.0
White, non-Hispanic, female, all ages				
...	All causes	012,740	100.0	992.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	271,790	26.8	266.4
2	Malignant neoplasms (C00-C97)	219,474	21.7	215.1
3	Cerebrovascular diseases (I60-I69)	71,036	7.0	69.6
4	Chronic lower respiratory diseases (J40-J47)	62,322	6.2	61.1
5	Alzheimer's disease (G30)	45,750	4.5	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,949	3.3	32.3
7	Influenza and pneumonia (J10-J18)	29,528	2.9	28.9
8	Diabetes mellitus (E10-E14)	26,794	2.6	26.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,539	1.6	16.2
10	Septicemia (A40-A41)	14,159	1.4	13.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,475	1.1	11.2
12	Parkinson's disease (G20-G21)	7,535	0.7	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,340	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	7,309	0.7	7.2
15	Atherosclerosis (I70)	6,523	0.6	6.4
...	All other causes (Residual)	182,217	18.0	178.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	996	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	34.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	13.2	2.9
3	Malignant neoplasms (C00-C97)	97	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	6.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.2	0.7
6	Influenza and pneumonia (J10-J18)	23	2.3	0.5
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Cerebrovascular diseases (I60-I69)	10	1.0	*
12	Meningococcal infection (A39)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	231	23.2	5.2
White, non-Hispanic, female, 5-14 years				
...	All causes	1,518	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	35.4	4.6
2	Malignant neoplasms (C00-C97)	264	17.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	7.0	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	3.3	0.4
6	Influenza and pneumonia (J10-J18)	38	2.5	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	19	1.3	*
10	Septicemia (A40-A41)	18	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
12	Diabetes mellitus (E10-E14)	13	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
14	Meningitis (G00,G03)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	316	20.8	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,416	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,769	51.1	21.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	505	9.3	3.9
3	Malignant neoplasms (C00-C97)	420	7.8	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	4.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	182	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	76	1.4	0.6
8	Cerebrovascular diseases (I60-I69)	52	1.0	0.4
9	Diabetes mellitus (E10-E14)	46	0.8	0.4
10	Influenza and pneumonia (J10-J18)	41	0.8	0.3
11	Septicemia (A40-A41)	31	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	24	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.2
15	Pneumonitis due to solids and liquids (J69)	15	0.3	*
...	All other causes (Residual)	851	15.7	6.6
White, non-Hispanic, female, 25-44 years				
...	All causes	27,527	100.0	103.3
1	Malignant neoplasms (C00-C97)	6,568	23.9	24.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,048	22.0	22.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,769	10.1	10.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,983	7.2	7.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	711	2.6	2.7
6	Cerebrovascular diseases (I60-I69)	704	2.6	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	698	2.5	2.6
8	Diabetes mellitus (E10-E14)	572	2.1	2.1
9	Chronic lower respiratory diseases (J40-J47)	391	1.4	1.5
10	Influenza and pneumonia (J10-J18)	366	1.3	1.4
11	Septicemia (A40-A41)	322	1.2	1.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	312	1.1	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	310	1.1	1.2
14	Pregnancy, childbirth and the puerperium (O00-099)	215	0.8	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	0.7	0.7
...	All other causes (Residual)	5,370	19.5	20.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	128,658	100.0	463.5
1	Malignant neoplasms (C00-C97)	53,348	41.5	192.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,683	16.1	74.5
3	Chronic lower respiratory diseases (J40-J47)	6,921	5.4	24.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,749	5.2	24.3
5	Cerebrovascular diseases (I60-I69)	4,410	3.4	15.9
6	Diabetes mellitus (E10-E14)	4,184	3.3	15.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,044	2.4	11.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,421	1.9	8.7
9	Septicemia (A40-A41)	1,859	1.4	6.7
10	Influenza and pneumonia (J10-J18)	1,660	1.3	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,375	1.1	5.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	665	0.5	2.4
13	Viral hepatitis (B15-B19)	643	0.5	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	571	0.4	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	561	0.4	2.0
...	All other causes (Residual)	19,564	15.2	70.5
White, non-Hispanic, female, 65 years and over				
...	All causes	842,996	100.0	4,843.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,988	29.4	1,424.9
2	Malignant neoplasms (C00-C97)	158,762	18.8	912.2
3	Cerebrovascular diseases (I60-I69)	65,818	7.8	378.2
4	Chronic lower respiratory diseases (J40-J47)	54,956	6.5	315.8
5	Alzheimer's disease (G30)	45,398	5.4	260.8
6	Influenza and pneumonia (J10-J18)	27,366	3.2	157.2
7	Diabetes mellitus (E10-E14)	21,977	2.6	126.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,237	1.9	93.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,915	1.8	85.7
10	Septicemia (A40-A41)	11,874	1.4	68.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,837	1.3	62.3
12	Parkinson's disease (G20-G21)	7,382	0.9	42.4
13	Pneumonitis due to solids and liquids (J69)	6,792	0.8	39.0
14	Atherosclerosis (I70)	6,343	0.8	36.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,908	0.6	28.2
...	All other causes (Residual)	141,443	16.8	812.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	289,163	100.0	774.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,302	25.3	196.3
2	Malignant neoplasms (C00-C97)	62,510	21.6	167.4
3	Cerebrovascular diseases (I60-I69)	17,378	6.0	46.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,393	4.6	35.9
5	Diabetes mellitus (E10-E14)	12,835	4.4	34.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,522	2.9	22.8
7	Chronic lower respiratory diseases (J40-J47)	8,136	2.8	21.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,017	2.8	21.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,908	2.4	18.5
10	Septicemia (A40-A41)	6,159	2.1	16.5
11	Influenza and pneumonia (J10-J18)	5,716	2.0	15.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,906	1.7	13.1
13	Certain conditions originating in the perinatal period (P00-P96)	4,837	1.7	13.0
14	Alzheimer's disease (G30)	4,579	1.6	12.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,495	0.9	6.7
...	All other causes (Residual)	49,470	17.1	132.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,065	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	29.7	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	13.0	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.8	4.1
4	Malignant neoplasms (C00-C97)	62	5.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	30	2.8	1.2
7	Influenza and pneumonia (J10-J18)	28	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.2	0.9
9	Septicemia (A40-A41)	20	1.9	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	255	23.9	10.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,500	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	33.1	7.9
2	Malignant neoplasms (C00-C97)	165	11.0	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	9.7	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	51	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.7	0.7
8	Cerebrovascular diseases (I60-I69)	23	1.5	0.4
9	Influenza and pneumonia (J10-J18)	22	1.5	0.4
10	Septicemia (A40-A41)	20	1.3	0.3
11	Anemias (D50-D64)	15	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
13	Diabetes mellitus (E10-E14)	12	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.5	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	325	21.7	5.2
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,244	100.0	115.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,030	41.8	48.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,734	23.9	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	5.9	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	4.6	5.3
5	Malignant neoplasms (C00-C97)	259	3.6	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	108	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	1.4	1.6
8	Anemias (D50-D64)	77	1.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	74	1.0	1.2
10	Diabetes mellitus (E10-E14)	73	1.0	1.2
11	Pregnancy, childbirth and the puerperium (O00-099)	69	1.0	1.1
12	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
13	Septicemia (A40-A41)	44	0.6	0.7
14	Influenza and pneumonia (J10-J18)	31	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.4	0.5
...	All other causes (Residual)	805	11.1	12.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	27,780	100.0	258.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,173	15.0	38.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,915	14.1	36.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,889	14.0	36.2
4	Malignant neoplasms (C00-C97)	3,195	11.5	29.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,122	11.2	29.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	889	3.2	8.3
7	Cerebrovascular diseases (I60-I69)	846	3.0	7.9
8	Diabetes mellitus (E10-E14)	752	2.7	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	1.6	4.0
10	Septicemia (A40-A41)	364	1.3	3.4
11	Chronic lower respiratory diseases (J40-J47)	313	1.1	2.9
12	Influenza and pneumonia (J10-J18)	291	1.0	2.7
13	Anemias (D50-D64)	275	1.0	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	267	1.0	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	245	0.9	2.3
...	All other causes (Residual)	4,812	17.3	44.7
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	82,235	100.0	1,045.2
1	Malignant neoplasms (C00-C97)	22,640	27.5	287.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,590	25.0	261.7
3	Cerebrovascular diseases (I60-I69)	4,406	5.4	56.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,088	5.0	52.0
5	Diabetes mellitus (E10-E14)	3,811	4.6	48.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,377	4.1	42.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,085	2.5	26.5
8	Chronic lower respiratory diseases (J40-J47)	1,892	2.3	24.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,635	2.0	20.8
10	Septicemia (A40-A41)	1,588	1.9	20.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,211	1.5	15.4
12	Influenza and pneumonia (J10-J18)	1,097	1.3	13.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,029	1.3	13.1
14	Viral hepatitis (B15-B19)	602	0.7	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	449	0.5	5.7
...	All other causes (Residual)	11,735	14.3	149.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	160,966	100.0	5,244.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,974	29.8	1,562.9
2	Malignant neoplasms (C00-C97)	36,177	22.5	1,178.6
3	Cerebrovascular diseases (I60-I69)	11,995	7.5	390.8
4	Diabetes mellitus (E10-E14)	8,186	5.1	266.7
5	Chronic lower respiratory diseases (J40-J47)	5,764	3.6	187.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,408	3.4	176.2
7	Alzheimer's disease (G30)	4,528	2.8	147.5
8	Influenza and pneumonia (J10-J18)	4,152	2.6	135.3
9	Septicemia (A40-A41)	3,996	2.5	130.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,438	2.1	112.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,532	1.6	82.5
12	Pneumonitis due to solids and liquids (J69)	1,326	0.8	43.2
13	Aortic aneurysm and dissection (I71)	749	0.5	24.4
14	Atherosclerosis (I70)	732	0.5	23.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	709	0.4	23.1
...	All other causes (Residual)	23,300	14.5	759.1
Black, non-Hispanic, male, all ages				
...	All causes	147,010	100.0	825.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,874	24.4	201.5
2	Malignant neoplasms (C00-C97)	32,363	22.0	181.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,130	6.2	51.3
4	Cerebrovascular diseases (I60-I69)	7,448	5.1	41.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,288	5.0	40.9
6	Diabetes mellitus (E10-E14)	5,666	3.9	31.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,600	3.1	25.8
8	Chronic lower respiratory diseases (J40-J47)	4,406	3.0	24.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,612	2.5	20.3
10	Septicemia (A40-A41)	2,755	1.9	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,712	1.8	15.2
12	Influenza and pneumonia (J10-J18)	2,697	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,108	1.4	11.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,634	1.1	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	1.1	8.9
...	All other causes (Residual)	23,125	15.7	129.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	606	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	29.5	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.5	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	7.6	3.6
4	Malignant neoplasms (C00-C97)	36	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	21	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.1	*
7	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.7	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	147	24.3	11.5
Black, non-Hispanic, male, 5-14 years				
...	All causes	885	100.0	27.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	35.3	9.8
2	Malignant neoplasms (C00-C97)	98	11.1	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	10.5	2.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.2	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.0	1.1
6	Chronic lower respiratory diseases (J40-J47)	33	3.7	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	3.1	0.8
8	Cerebrovascular diseases (I60-I69)	14	1.6	*
9	Septicemia (A40-A41)	13	1.5	*
9	Influenza and pneumonia (J10-J18)	13	1.5	*
11	Anemias (D50-D64)	9	1.0	*
12	Meningitis (G00,G03)	6	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Diseases of appendix (K35-K38)	3	0.3	*
...	All other causes (Residual)	180	20.3	5.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,615	100.0	177.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,750	49.0	87.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,301	23.2	41.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	6.7	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	4.1	7.2
5	Malignant neoplasms (C00-C97)	151	2.7	4.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.1	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.0	1.8
8	Chronic lower respiratory diseases (J40-J47)	47	0.8	1.5
9	Anemias (D50-D64)	45	0.8	1.4
10	Diabetes mellitus (E10-E14)	39	0.7	1.2
11	Septicemia (A40-A41)	24	0.4	0.8
11	Cerebrovascular diseases (I60-I69)	24	0.4	0.8
11	Legal intervention (Y35,Y89.0)	24	0.4	0.8
14	Influenza and pneumonia (J10-J18)	23	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	453	8.1	14.3
Black, non-Hispanic, male, 25-44 years				
...	All causes	17,157	100.0	336.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,345	19.5	65.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,853	16.6	55.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,593	15.1	50.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,909	11.1	37.4
5	Malignant neoplasms (C00-C97)	1,273	7.4	24.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	727	4.2	14.2
7	Diabetes mellitus (E10-E14)	434	2.5	8.5
8	Cerebrovascular diseases (I60-I69)	418	2.4	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	224	1.3	4.4
10	Influenza and pneumonia (J10-J18)	171	1.0	3.3
11	Septicemia (A40-A41)	170	1.0	3.3
12	Chronic lower respiratory diseases (J40-J47)	155	0.9	3.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	155	0.9	3.0
14	Anemias (D50-D64)	146	0.9	2.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	131	0.8	2.6
...	All other causes (Residual)	2,453	14.3	48.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	48,342	100.0	1,347.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,901	26.7	359.7
2	Malignant neoplasms (C00-C97)	12,004	24.8	334.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,968	6.1	82.7
4	Cerebrovascular diseases (I60-I69)	2,450	5.1	68.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,417	5.0	67.4
6	Diabetes mellitus (E10-E14)	2,046	4.2	57.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,184	2.4	33.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,127	2.3	31.4
9	Chronic lower respiratory diseases (J40-J47)	1,011	2.1	28.2
10	Septicemia (A40-A41)	829	1.7	23.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	817	1.7	22.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	702	1.5	19.6
13	Influenza and pneumonia (J10-J18)	671	1.4	18.7
14	Viral hepatitis (B15-B19)	405	0.8	11.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	344	0.7	9.6
...	All other causes (Residual)	6,466	13.4	180.3
Black, non-Hispanic, male, 65 years and over				
...	All causes	69,716	100.0	5,919.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,033	28.7	1,701.1
2	Malignant neoplasms (C00-C97)	18,793	27.0	1,595.8
3	Cerebrovascular diseases (I60-I69)	4,506	6.5	382.6
4	Diabetes mellitus (E10-E14)	3,144	4.5	267.0
5	Chronic lower respiratory diseases (J40-J47)	3,133	4.5	266.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,153	3.1	182.8
7	Influenza and pneumonia (J10-J18)	1,758	2.5	149.3
8	Septicemia (A40-A41)	1,639	2.4	139.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,346	1.9	114.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,270	1.8	107.8
11	Alzheimer's disease (G30)	1,262	1.8	107.2
12	Pneumonitis due to solids and liquids (J69)	648	0.9	55.0
13	Parkinson's disease (G20-G21)	390	0.6	33.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	346	0.5	29.4
15	Aortic aneurysm and dissection (I71)	327	0.5	27.8
...	All other causes (Residual)	8,968	12.9	761.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	142,153	100.0	727.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,428	26.3	191.6
2	Malignant neoplasms (C00-C97)	30,147	21.2	154.3
3	Cerebrovascular diseases (I60-I69)	9,930	7.0	50.8
4	Diabetes mellitus (E10-E14)	7,169	5.0	36.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,405	3.1	22.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,263	3.0	21.8
7	Chronic lower respiratory diseases (J40-J47)	3,730	2.6	19.1
8	Septicemia (A40-A41)	3,404	2.4	17.4
9	Alzheimer's disease (G30)	3,296	2.3	16.9
10	Influenza and pneumonia (J10-J18)	3,019	2.1	15.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,798	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,308	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,125	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	0.9	6.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	861	0.6	4.4
...	All other causes (Residual)	26,036	18.3	133.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	459	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	29.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	4.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	10.9	4.1
4	Malignant neoplasms (C00-C97)	26	5.7	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	2.1
6	Influenza and pneumonia (J10-J18)	12	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Meningitis (G00,G03)	3	0.7	*
12	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	106	23.1	8.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	615	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	30.1	6.0
2	Malignant neoplasms (C00-C97)	67	10.9	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	8.5	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	7.0	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.5	1.3
6	Chronic lower respiratory diseases (J40-J47)	18	2.9	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.6	*
9	Diabetes mellitus (E10-E14)	9	1.5	*
9	Cerebrovascular diseases (I60-I69)	9	1.5	*
9	Influenza and pneumonia (J10-J18)	9	1.5	*
12	Septicemia (A40-A41)	7	1.1	*
13	Anemias (D50-D64)	6	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	140	22.8	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,629	100.0	52.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	26.6	14.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	17.2	9.0
3	Malignant neoplasms (C00-C97)	108	6.6	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	6.4	3.4
5	Pregnancy, childbirth and the puerperium (O00-099)	69	4.2	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.3	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	51	3.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.2	1.2
9	Diabetes mellitus (E10-E14)	34	2.1	1.1
10	Anemias (D50-D64)	32	2.0	1.0
11	Cerebrovascular diseases (I60-I69)	27	1.7	0.9
11	Chronic lower respiratory diseases (J40-J47)	27	1.7	0.9
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.1	*
15	Influenza and pneumonia (J10-J18)	8	0.5	*
...	All other causes (Residual)	328	20.1	10.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-44 years				
...	All causes	10,623	100.0	188.1
1	Malignant neoplasms (C00-C97)	1,922	18.1	34.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,580	14.9	28.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,213	11.4	21.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,062	10.0	18.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	5.1	9.6
6	Cerebrovascular diseases (I60-I69)	428	4.0	7.6
7	Diabetes mellitus (E10-E14)	318	3.0	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.0	3.7
9	Septicemia (A40-A41)	194	1.8	3.4
10	Pregnancy, childbirth and the puerperium (O00-099)	180	1.7	3.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	1.5	2.9
12	Chronic lower respiratory diseases (J40-J47)	158	1.5	2.8
13	Anemias (D50-D64)	129	1.2	2.3
14	Influenza and pneumonia (J10-J18)	120	1.1	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	1.1	2.0
...	All other causes (Residual)	2,291	21.6	40.6
Black, non-Hispanic, female, 45-64 years				
...	All causes	33,893	100.0	791.7
1	Malignant neoplasms (C00-C97)	10,636	31.4	248.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,689	22.7	179.6
3	Cerebrovascular diseases (I60-I69)	1,956	5.8	45.7
4	Diabetes mellitus (E10-E14)	1,765	5.2	41.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,120	3.3	26.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	960	2.8	22.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	901	2.7	21.0
8	Chronic lower respiratory diseases (J40-J47)	881	2.6	20.6
9	Septicemia (A40-A41)	759	2.2	17.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	509	1.5	11.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	508	1.5	11.9
12	Influenza and pneumonia (J10-J18)	426	1.3	10.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	212	0.6	5.0
14	Viral hepatitis (B15-B19)	197	0.6	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.4	3.3
...	All other causes (Residual)	5,233	15.4	122.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65 years and over				
...	All causes	91,250	100.0	4,823.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,941	30.6	1,476.9
2	Malignant neoplasms (C00-C97)	17,384	19.1	918.9
3	Cerebrovascular diseases (I60-I69)	7,489	8.2	395.9
4	Diabetes mellitus (E10-E14)	5,042	5.5	266.5
5	Alzheimer's disease (G30)	3,266	3.6	172.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,255	3.6	172.1
7	Chronic lower respiratory diseases (J40-J47)	2,631	2.9	139.1
8	Influenza and pneumonia (J10-J18)	2,394	2.6	126.5
9	Septicemia (A40-A41)	2,357	2.6	124.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,168	2.4	114.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,186	1.3	62.7
12	Pneumonitis due to solids and liquids (J69)	678	0.7	35.8
13	Atherosclerosis (I70)	486	0.5	25.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	425	0.5	22.5
15	Aortic aneurysm and dissection (I71)	422	0.5	22.3
...	All other causes (Residual)	14,126	15.5	746.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable