

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	122,416	100.0	296.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,788	22.7	67.2
2	Malignant neoplasms (C00-C97)	24,522	20.0	59.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,408	8.5	25.2
4	Cerebrovascular diseases (I60-I69)	6,781	5.5	16.4
5	Diabetes mellitus (E10-E14)	5,999	4.9	14.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,383	2.8	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,271	2.7	7.9
8	Chronic lower respiratory diseases (J40-J47)	3,107	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,912	2.4	7.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,681	2.2	6.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,207	1.8	5.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,196	1.8	5.3
13	Alzheimer's disease (G30)	1,897	1.5	4.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,765	1.4	4.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,758	1.4	4.3
...	All other causes (Residual)	21,741	17.8	52.6
Hispanic, both sexes, 1-4 years				
...	All causes	947	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	33.1	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	138	14.6	4.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	8.4	2.3
4	Malignant neoplasms (C00-C97)	70	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.1	1.1
6	Septicemia (A40-A41)	24	2.5	0.7
6	Influenza and pneumonia (J10-J18)	24	2.5	0.7
8	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Meningococcal infection (A39)	5	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
10	Anemias (D50-D64)	5	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Meningitis (G00,G03)	3	0.3	*
15	Cerebrovascular diseases (I60-I69)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	201	21.2	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,105	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	35.4	5.1
2	Malignant neoplasms (C00-C97)	211	19.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	6.1	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	3.5	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.1	0.4
7	Influenza and pneumonia (J10-J18)	13	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	12	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
10	Cerebrovascular diseases (I60-I69)	10	0.9	*
11	Septicemia (A40-A41)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
...	All other causes (Residual)	212	19.2	2.8
Hispanic, both sexes, 15-24 years				
...	All causes	5,428	100.0	76.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	43.7	33.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,235	22.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	571	10.5	8.0
4	Malignant neoplasms (C00-C97)	305	5.6	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	2.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	1.1	0.9
7	Cerebrovascular diseases (I60-I69)	36	0.7	0.5
8	Influenza and pneumonia (J10-J18)	32	0.6	0.5
9	Legal intervention (Y35,Y89.0)	30	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	29	0.5	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	17	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
14	Septicemia (A40-A41)	13	0.2	*
14	Diabetes mellitus (E10-E14)	13	0.2	*
...	All other causes (Residual)	545	10.0	7.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,505	100.0	105.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,909	26.9	28.4
2	Malignant neoplasms (C00-C97)	1,871	12.9	13.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,439	9.9	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,236	8.5	9.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	966	6.7	7.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	887	6.1	6.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	541	3.7	3.9
8	Cerebrovascular diseases (I60-I69)	397	2.7	2.9
9	Diabetes mellitus (E10-E14)	251	1.7	1.8
10	Influenza and pneumonia (J10-J18)	130	0.9	0.9
11	Septicemia (A40-A41)	123	0.8	0.9
12	Viral hepatitis (B15-B19)	119	0.8	0.9
13	Chronic lower respiratory diseases (J40-J47)	96	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.6	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	0.6	0.7
...	All other causes (Residual)	2,353	16.2	17.1
Hispanic, both sexes, 45-64 years				
...	All causes	27,826	100.0	445.2
1	Malignant neoplasms (C00-C97)	7,699	27.7	123.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,570	20.0	89.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,927	6.9	30.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,767	6.4	28.3
5	Diabetes mellitus (E10-E14)	1,608	5.8	25.7
6	Cerebrovascular diseases (I60-I69)	1,366	4.9	21.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	755	2.7	12.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	1.8	8.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	460	1.7	7.4
10	Viral hepatitis (B15-B19)	446	1.6	7.1
11	Chronic lower respiratory diseases (J40-J47)	396	1.4	6.3
12	Septicemia (A40-A41)	378	1.4	6.0
13	Influenza and pneumonia (J10-J18)	367	1.3	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	1.2	5.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	210	0.8	3.4
...	All other causes (Residual)	4,054	14.6	64.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	67,236	100.0	3,105.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,696	30.8	955.9
2	Malignant neoplasms (C00-C97)	14,344	21.3	662.5
3	Cerebrovascular diseases (I60-I69)	4,947	7.4	228.5
4	Diabetes mellitus (E10-E14)	4,125	6.1	190.5
5	Chronic lower respiratory diseases (J40-J47)	2,566	3.8	118.5
6	Influenza and pneumonia (J10-J18)	2,293	3.4	105.9
7	Alzheimer's disease (G30)	1,866	2.8	86.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,538	2.3	71.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,319	2.0	60.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,068	1.6	49.3
11	Septicemia (A40-A41)	1,046	1.6	48.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	870	1.3	40.2
13	Parkinson's disease (G20-G21)	568	0.8	26.2
14	Pneumonitis due to solids and liquids (J69)	458	0.7	21.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	372	0.6	17.2
...	All other causes (Residual)	9,160	13.6	423.1
Hispanic, male, all ages				
...	All causes	68,544	100.0	321.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,983	21.9	70.2
2	Malignant neoplasms (C00-C97)	13,002	19.0	60.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,796	11.4	36.5
4	Cerebrovascular diseases (I60-I69)	3,207	4.7	15.0
5	Diabetes mellitus (E10-E14)	2,855	4.2	13.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,782	4.1	13.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,376	3.5	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,840	2.7	8.6
9	Chronic lower respiratory diseases (J40-J47)	1,631	2.4	7.6
10	Certain conditions originating in the perinatal period (P00-P96)	1,538	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,403	2.0	6.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,371	2.0	6.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	902	1.3	4.2
15	Septicemia (A40-A41)	801	1.2	3.8
...	All other causes (Residual)	10,956	16.0	51.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	535	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	34.8	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	15.3	4.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.4	2.5
4	Malignant neoplasms (C00-C97)	29	5.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.5	1.4
6	Septicemia (A40-A41)	14	2.6	*
7	Influenza and pneumonia (J10-J18)	9	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
10	Anemias (D50-D64)	3	0.6	*
10	Cerebrovascular diseases (I60-I69)	3	0.6	*
12	Meningococcal infection (A39)	2	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.7	6.3
Hispanic, male, 5-14 years				
...	All causes	656	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	36.7	6.2
2	Malignant neoplasms (C00-C97)	118	18.0	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	7.5	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.1	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.3	*
7	Influenza and pneumonia (J10-J18)	10	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
9	Septicemia (A40-A41)	6	0.9	*
10	Cerebrovascular diseases (I60-I69)	5	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Anemias (D50-D64)	2	0.3	*
15	Meningitis (G00,G03)	2	0.3	*
15	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	119	18.1	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,376	100.0	115.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,926	44.0	50.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,125	25.7	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	11.1	12.8
4	Malignant neoplasms (C00-C97)	207	4.7	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.1
7	Legal intervention (Y35,Y89.0)	30	0.7	0.8
8	Influenza and pneumonia (J10-J18)	24	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	20	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.3	*
13	Diabetes mellitus (E10-E14)	8	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
15	Septicemia (A40-A41)	7	0.2	*
15	Aortic aneurysm and dissection (I71)	7	0.2	*
...	All other causes (Residual)	345	7.9	9.1
Hispanic, male, 25-44 years				
...	All causes	10,520	100.0	142.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,187	30.3	43.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,228	11.7	16.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	959	9.1	13.0
4	Malignant neoplasms (C00-C97)	871	8.3	11.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	818	7.8	11.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	691	6.6	9.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	438	4.2	5.9
8	Cerebrovascular diseases (I60-I69)	251	2.4	3.4
9	Diabetes mellitus (E10-E14)	148	1.4	2.0
10	Viral hepatitis (B15-B19)	91	0.9	1.2
11	Influenza and pneumonia (J10-J18)	79	0.8	1.1
12	Septicemia (A40-A41)	69	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	65	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.5	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.4	0.6
...	All other causes (Residual)	1,526	14.5	20.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	17,697	100.0	574.5
1	Malignant neoplasms (C00-C97)	4,094	23.1	132.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,945	22.3	128.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,442	8.1	46.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,379	7.8	44.8
5	Diabetes mellitus (E10-E14)	925	5.2	30.0
6	Cerebrovascular diseases (I60-I69)	808	4.6	26.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	597	3.4	19.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	362	2.0	11.8
9	Viral hepatitis (B15-B19)	316	1.8	10.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	1.6	9.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	1.5	8.5
12	Influenza and pneumonia (J10-J18)	222	1.3	7.2
13	Septicemia (A40-A41)	215	1.2	7.0
14	Chronic lower respiratory diseases (J40-J47)	201	1.1	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	126	0.7	4.1
...	All other causes (Residual)	2,516	14.2	81.7
Hispanic, male, 65 years and over				
...	All causes	31,766	100.0	3,465.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,895	31.1	1,079.6
2	Malignant neoplasms (C00-C97)	7,673	24.2	837.2
3	Cerebrovascular diseases (I60-I69)	2,107	6.6	229.9
4	Diabetes mellitus (E10-E14)	1,772	5.6	193.3
5	Chronic lower respiratory diseases (J40-J47)	1,334	4.2	145.5
6	Influenza and pneumonia (J10-J18)	1,022	3.2	111.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	722	2.3	78.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	710	2.2	77.5
9	Alzheimer's disease (G30)	592	1.9	64.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	555	1.7	60.6
11	Septicemia (A40-A41)	454	1.4	49.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	345	1.1	37.6
13	Parkinson's disease (G20-G21)	330	1.0	36.0
14	Pneumonitis due to solids and liquids (J69)	218	0.7	23.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	174	0.5	19.0
...	All other causes (Residual)	3,863	12.2	421.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	53,872	100.0	269.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,805	23.8	64.1
2	Malignant neoplasms (C00-C97)	11,520	21.4	57.7
3	Cerebrovascular diseases (I60-I69)	3,574	6.6	17.9
4	Diabetes mellitus (E10-E14)	3,144	5.8	15.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,612	4.8	13.1
6	Influenza and pneumonia (J10-J18)	1,509	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,476	2.7	7.4
8	Alzheimer's disease (G30)	1,291	2.4	6.5
9	Certain conditions originating in the perinatal period (P00-P96)	1,143	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,095	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	1.9	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	863	1.6	4.3
13	Septicemia (A40-A41)	847	1.6	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	624	1.2	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	0.9	2.4
...	All other causes (Residual)	9,873	18.3	49.4
Hispanic, female, 1-4 years				
...	All causes	412	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	30.8	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	13.6	3.3
3	Malignant neoplasms (C00-C97)	41	10.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	8.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.6	*
5	Influenza and pneumonia (J10-J18)	15	3.6	*
7	Septicemia (A40-A41)	10	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.0	*
10	Meningococcal infection (A39)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	88	21.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	449	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	33.4	4.0
2	Malignant neoplasms (C00-C97)	93	20.7	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.0	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	4.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.2	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.6	*
8	Cerebrovascular diseases (I60-I69)	5	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
11	Septicemia (A40-A41)	3	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	Influenza and pneumonia (J10-J18)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (Residual)	88	19.6	2.4
Hispanic, female, 15-24 years				
...	All causes	1,052	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	42.4	13.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	10.5	3.3
3	Malignant neoplasms (C00-C97)	98	9.3	3.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	8.0	2.5
5	Pregnancy, childbirth and the puerperium (O00-099)	29	2.8	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	2.7	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	2.0	0.6
8	Cerebrovascular diseases (I60-I69)	16	1.5	*
9	Anemias (D50-D64)	8	0.8	*
9	Influenza and pneumonia (J10-J18)	8	0.8	*
11	Septicemia (A40-A41)	6	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Diabetes mellitus (E10-E14)	5	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	173	16.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	3,985	100.0	62.5
1	Malignant neoplasms (C00-C97)	1,000	25.1	15.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	722	18.1	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	7.0	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	5.3	3.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	196	4.9	3.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	3.7	2.3
7	Cerebrovascular diseases (I60-I69)	146	3.7	2.3
8	Diabetes mellitus (E10-E14)	103	2.6	1.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	103	2.6	1.6
10	Pregnancy, childbirth and the puerperium (O00-099)	75	1.9	1.2
11	Septicemia (A40-A41)	54	1.4	0.8
12	Influenza and pneumonia (J10-J18)	51	1.3	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.2	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.0	0.6
15	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.5
...	All other causes (Residual)	780	19.6	12.2
Hispanic, female, 45-64 years				
...	All causes	10,129	100.0	319.6
1	Malignant neoplasms (C00-C97)	3,605	35.6	113.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,625	16.0	51.3
3	Diabetes mellitus (E10-E14)	683	6.7	21.5
4	Cerebrovascular diseases (I60-I69)	558	5.5	17.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	4.8	15.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	388	3.8	12.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	2.1	6.8
8	Chronic lower respiratory diseases (J40-J47)	195	1.9	6.2
9	Septicemia (A40-A41)	163	1.6	5.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	158	1.6	5.0
11	Influenza and pneumonia (J10-J18)	145	1.4	4.6
12	Viral hepatitis (B15-B19)	130	1.3	4.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	1.0	3.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	84	0.8	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.6	2.0
...	All other causes (Residual)	1,536	15.2	48.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	35,470	100.0	2,841.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,801	30.5	865.1
2	Malignant neoplasms (C00-C97)	6,671	18.8	534.3
3	Cerebrovascular diseases (I60-I69)	2,840	8.0	227.5
4	Diabetes mellitus (E10-E14)	2,353	6.6	188.5
5	Alzheimer's disease (G30)	1,274	3.6	102.0
6	Influenza and pneumonia (J10-J18)	1,271	3.6	101.8
7	Chronic lower respiratory diseases (J40-J47)	1,232	3.5	98.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	816	2.3	65.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	609	1.7	48.8
10	Septicemia (A40-A41)	592	1.7	47.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	525	1.5	42.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	513	1.4	41.1
13	Pneumonitis due to solids and liquids (J69)	240	0.7	19.2
14	Parkinson's disease (G20-G21)	238	0.7	19.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.6	15.9
...	All other causes (Residual)	5,297	14.9	424.3
White, non-Hispanic, both sexes, all ages				
...	All causes	1,933,382	100.0	967.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	537,512	27.8	269.1
2	Malignant neoplasms (C00-C97)	453,534	23.5	227.0
3	Cerebrovascular diseases (I60-I69)	121,065	6.3	60.6
4	Chronic lower respiratory diseases (J40-J47)	109,670	5.7	54.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,466	4.4	42.8
6	Alzheimer's disease (G30)	59,153	3.1	29.6
7	Diabetes mellitus (E10-E14)	52,101	2.7	26.1
8	Influenza and pneumonia (J10-J18)	49,456	2.6	24.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,508	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,001	1.4	13.5
11	Septicemia (A40-A41)	25,041	1.3	12.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,144	1.0	10.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16,569	0.9	8.3
14	Parkinson's disease (G20-G21)	16,428	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	14,411	0.7	7.2
...	All other causes (Residual)	314,323	16.3	157.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,458	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	933	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.0	2.9
3	Malignant neoplasms (C00-C97)	244	9.9	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	5.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	0.9
6	Influenza and pneumonia (J10-J18)	55	2.2	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.6	0.4
8	Septicemia (A40-A41)	36	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	25	1.0	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	1.0	0.3
11	Meningitis (G00,G03)	16	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	10	0.4	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (Residual)	564	22.9	6.1
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,820	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,573	41.2	6.4
2	Malignant neoplasms (C00-C97)	587	15.4	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	3.3	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	3.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.4	0.2
8	Cerebrovascular diseases (I60-I69)	48	1.3	0.2
9	Influenza and pneumonia (J10-J18)	43	1.1	0.2
10	Septicemia (A40-A41)	41	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	41	1.1	0.2
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	18	0.5	*
14	Anemias (D50-D64)	17	0.4	*
14	Meningitis (G00,G03)	17	0.4	*
...	All other causes (Residual)	723	18.9	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,684	100.0	75.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,847	55.1	41.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,045	15.5	11.6
3	Malignant neoplasms (C00-C97)	1,016	5.2	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	917	4.7	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	290	1.5	1.1
7	Cerebrovascular diseases (I60-I69)	117	0.6	0.4
8	Influenza and pneumonia (J10-J18)	108	0.5	0.4
9	Diabetes mellitus (E10-E14)	87	0.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.4	0.3
12	Septicemia (A40-A41)	70	0.4	0.3
13	Pregnancy, childbirth and the puerperium (O00-099)	60	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.2
15	Anemias (D50-D64)	29	0.1	0.1
...	All other causes (Residual)	2,357	12.0	9.0
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	78,940	100.0	145.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,737	26.3	38.2
2	Malignant neoplasms (C00-C97)	12,332	15.6	22.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,156	12.9	18.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,345	11.8	17.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,100	2.7	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,041	2.6	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,825	2.3	3.4
8	Diabetes mellitus (E10-E14)	1,571	2.0	2.9
9	Cerebrovascular diseases (I60-I69)	1,536	1.9	2.8
10	Influenza and pneumonia (J10-J18)	735	0.9	1.4
11	Chronic lower respiratory diseases (J40-J47)	703	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	664	0.8	1.2
13	Septicemia (A40-A41)	646	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	502	0.6	0.9
15	Viral hepatitis (B15-B19)	421	0.5	0.8
...	All other causes (Residual)	13,626	17.3	25.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	321,879	100.0	602.6
1	Malignant neoplasms (C00-C97)	112,416	34.9	210.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,792	22.6	136.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,134	6.3	37.7
4	Chronic lower respiratory diseases (J40-J47)	12,885	4.0	24.1
5	Diabetes mellitus (E10-E14)	10,427	3.2	19.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10,168	3.2	19.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,685	3.0	18.1
8	Cerebrovascular diseases (I60-I69)	9,676	3.0	18.1
9	Septicemia (A40-A41)	3,872	1.2	7.2
10	Influenza and pneumonia (J10-J18)	3,537	1.1	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,243	1.0	6.1
12	Viral hepatitis (B15-B19)	2,283	0.7	4.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,829	0.6	3.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,511	0.5	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,470	0.5	2.8
...	All other causes (Residual)	45,951	14.3	86.0
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,493,442	100.0	5,003.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	453,580	30.4	1,519.5
2	Malignant neoplasms (C00-C97)	326,892	21.9	1,095.1
3	Cerebrovascular diseases (I60-I69)	109,604	7.3	367.2
4	Chronic lower respiratory diseases (J40-J47)	95,939	6.4	321.4
5	Alzheimer's disease (G30)	58,585	3.9	196.3
6	Influenza and pneumonia (J10-J18)	44,866	3.0	150.3
7	Diabetes mellitus (E10-E14)	39,991	2.7	134.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,650	2.1	102.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,647	1.9	92.6
10	Septicemia (A40-A41)	20,263	1.4	67.9
11	Parkinson's disease (G20-G21)	16,070	1.1	53.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,913	1.0	50.0
13	Pneumonitis due to solids and liquids (J69)	13,365	0.9	44.8
14	Atherosclerosis (I70)	9,992	0.7	33.5
15	Aortic aneurysm and dissection (I71)	9,965	0.7	33.4
...	All other causes (Residual)	221,120	14.8	740.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	938,143	100.0	957.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,638	28.1	269.1
2	Malignant neoplasms (C00-C97)	234,392	25.0	239.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,465	5.7	54.6
4	Chronic lower respiratory diseases (J40-J47)	51,909	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	46,087	4.9	47.0
6	Diabetes mellitus (E10-E14)	25,773	2.7	26.3
7	Influenza and pneumonia (J10-J18)	21,905	2.3	22.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,214	2.3	21.6
9	Alzheimer's disease (G30)	16,989	1.8	17.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,247	1.6	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,931	1.4	13.2
12	Septicemia (A40-A41)	11,188	1.2	11.4
13	Parkinson's disease (G20-G21)	9,469	1.0	9.7
14	Pneumonitis due to solids and liquids (J69)	7,367	0.8	7.5
15	Aortic aneurysm and dissection (I71)	6,906	0.7	7.0
...	All other causes (Residual)	139,663	14.9	142.5
White, non-Hispanic, male, 1-4 years				
...	All causes	1,365	100.0	29.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	553	40.5	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	10.6	3.1
3	Malignant neoplasms (C00-C97)	134	9.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.7	0.8
6	Influenza and pneumonia (J10-J18)	23	1.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.5	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
9	Septicemia (A40-A41)	15	1.1	*
9	Cerebrovascular diseases (I60-I69)	15	1.1	*
11	Meningitis (G00,G03)	10	0.7	*
12	Meningococcal infection (A39)	8	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	296	21.7	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,264	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	991	43.8	7.9
2	Malignant neoplasms (C00-C97)	339	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	5.3	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	4.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	3.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.0	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	24	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.2
10	Influenza and pneumonia (J10-J18)	22	1.0	0.2
11	Septicemia (A40-A41)	20	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.5	*
13	Meningitis (G00,G03)	9	0.4	*
14	Meningococcal infection (A39)	7	0.3	*
14	Diabetes mellitus (E10-E14)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	408	18.0	3.2
White, non-Hispanic, male, 15-24 years				
...	All causes	14,017	100.0	104.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,885	56.3	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,535	18.1	19.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	624	4.5	4.7
4	Malignant neoplasms (C00-C97)	593	4.2	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	363	2.6	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	1.1	1.2
7	Influenza and pneumonia (J10-J18)	73	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	63	0.4	0.5
9	Diabetes mellitus (E10-E14)	45	0.3	0.3
9	Chronic lower respiratory diseases (J40-J47)	45	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.3
12	Septicemia (A40-A41)	39	0.3	0.3
13	Legal intervention (Y35,Y89.0)	26	0.2	0.2
14	Aortic aneurysm and dissection (I71)	22	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.1	0.2
...	All other causes (Residual)	1,486	10.6	11.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	51,351	100.0	188.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,870	29.0	54.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,313	14.2	26.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,274	14.2	26.7
4	Malignant neoplasms (C00-C97)	5,611	10.9	20.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,486	2.9	5.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,423	2.8	5.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,361	2.7	5.0
8	Diabetes mellitus (E10-E14)	986	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	790	1.5	2.9
10	Influenza and pneumonia (J10-J18)	415	0.8	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	373	0.7	1.4
12	Chronic lower respiratory diseases (J40-J47)	333	0.6	1.2
13	Septicemia (A40-A41)	328	0.6	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	275	0.5	1.0
15	Viral hepatitis (B15-B19)	268	0.5	1.0
...	All other causes (Residual)	8,245	16.1	30.2
White, non-Hispanic, male, 45-64 years				
...	All causes	196,912	100.0	748.7
1	Malignant neoplasms (C00-C97)	60,024	30.5	228.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,407	26.6	199.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,890	7.1	52.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,219	3.7	27.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,152	3.6	27.2
6	Chronic lower respiratory diseases (J40-J47)	6,524	3.3	24.8
7	Diabetes mellitus (E10-E14)	6,312	3.2	24.0
8	Cerebrovascular diseases (I60-I69)	5,288	2.7	20.1
9	Influenza and pneumonia (J10-J18)	2,069	1.1	7.9
10	Septicemia (A40-A41)	2,023	1.0	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,820	0.9	6.9
12	Viral hepatitis (B15-B19)	1,668	0.8	6.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,578	0.8	6.0
14	Aortic aneurysm and dissection (I71)	1,097	0.6	4.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	992	0.5	3.8
...	All other causes (Residual)	26,849	13.6	102.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent	
			of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	664,751	100.0	5,301.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	203,361	30.6	1,621.7
2	Malignant neoplasms (C00-C97)	167,666	25.2	1,337.1
3	Chronic lower respiratory diseases (J40-J47)	44,968	6.8	358.6
4	Cerebrovascular diseases (I60-I69)	39,873	6.0	318.0
5	Influenza and pneumonia (J10-J18)	19,238	2.9	153.4
6	Diabetes mellitus (E10-E14)	18,421	2.8	146.9
7	Alzheimer's disease (G30)	16,731	2.5	133.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,912	2.2	118.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,086	2.0	104.4
10	Parkinson's disease (G20-G21)	9,233	1.4	73.6
11	Septicemia (A40-A41)	8,695	1.3	69.3
12	Pneumonitis due to solids and liquids (J69)	6,732	1.0	53.7
13	Aortic aneurysm and dissection (I71)	5,586	0.8	44.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,107	0.8	40.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,605	0.7	36.7
...	All other causes (Residual)	86,537	13.0	690.1
White, non-Hispanic, female, all ages				
...	All causes	995,239	100.0	977.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	273,874	27.5	269.1
2	Malignant neoplasms (C00-C97)	219,142	22.0	215.3
3	Cerebrovascular diseases (I60-I69)	74,978	7.5	73.7
4	Chronic lower respiratory diseases (J40-J47)	57,761	5.8	56.7
5	Alzheimer's disease (G30)	42,164	4.2	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,001	3.2	31.4
7	Influenza and pneumonia (J10-J18)	27,551	2.8	27.1
8	Diabetes mellitus (E10-E14)	26,328	2.6	25.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,261	1.6	16.0
10	Septicemia (A40-A41)	13,853	1.4	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,422	1.0	10.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,213	0.7	7.1
13	Pneumonitis due to solids and liquids (J69)	7,044	0.7	6.9
14	Parkinson's disease (G20-G21)	6,959	0.7	6.8
15	Atherosclerosis (I70)	6,534	0.7	6.4
...	All other causes (Residual)	173,154	17.4	170.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,093	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	34.8	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	11.4	2.8
3	Malignant neoplasms (C00-C97)	110	10.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.9	1.0
6	Influenza and pneumonia (J10-J18)	32	2.9	0.7
7	Septicemia (A40-A41)	21	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.6	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.3	*
...	All other causes (Residual)	266	24.3	5.9
White, non-Hispanic, female, 5-14 years				
...	All causes	1,556	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	37.4	4.9
2	Malignant neoplasms (C00-C97)	248	15.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	6.8	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	3.9	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.7	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	3.1	0.4
7	Cerebrovascular diseases (I60-I69)	24	1.5	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.5	0.2
9	Septicemia (A40-A41)	21	1.3	0.2
9	Influenza and pneumonia (J10-J18)	21	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
12	Diabetes mellitus (E10-E14)	12	0.8	*
13	Anemias (D50-D64)	11	0.7	*
14	Meningitis (G00,G03)	8	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	7	0.4	*
...	All other causes (Residual)	307	19.7	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,667	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,962	52.3	23.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	9.0	4.0
3	Malignant neoplasms (C00-C97)	423	7.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	293	5.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	3.4	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	2.4	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	60	1.1	0.5
8	Cerebrovascular diseases (I60-I69)	54	1.0	0.4
9	Diabetes mellitus (E10-E14)	42	0.7	0.3
10	Influenza and pneumonia (J10-J18)	35	0.6	0.3
11	Septicemia (A40-A41)	31	0.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	0.2
13	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.3	*
...	All other causes (Residual)	837	14.8	6.5
White, non-Hispanic, female, 25-44 years				
...	All causes	27,589	100.0	102.1
1	Malignant neoplasms (C00-C97)	6,721	24.4	24.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,867	21.3	21.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,882	10.4	10.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,032	7.4	7.5
5	Cerebrovascular diseases (I60-I69)	746	2.7	2.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	680	2.5	2.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	677	2.5	2.5
8	Diabetes mellitus (E10-E14)	585	2.1	2.2
9	Chronic lower respiratory diseases (J40-J47)	370	1.3	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	339	1.2	1.3
11	Influenza and pneumonia (J10-J18)	320	1.2	1.2
12	Septicemia (A40-A41)	318	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	291	1.1	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	0.8	0.8
15	Pregnancy, childbirth and the puerperium (O00-099)	196	0.7	0.7
...	All other causes (Residual)	5,338	19.3	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	124,967	100.0	460.8
1	Malignant neoplasms (C00-C97)	52,392	41.9	193.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,385	16.3	75.2
3	Chronic lower respiratory diseases (J40-J47)	6,361	5.1	23.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,244	5.0	23.0
5	Cerebrovascular diseases (I60-I69)	4,388	3.5	16.2
6	Diabetes mellitus (E10-E14)	4,115	3.3	15.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,016	2.4	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,466	2.0	9.1
9	Septicemia (A40-A41)	1,849	1.5	6.8
10	Influenza and pneumonia (J10-J18)	1,468	1.2	5.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,423	1.1	5.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	647	0.5	2.4
13	Viral hepatitis (B15-B19)	615	0.5	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	562	0.4	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	521	0.4	1.9
...	All other causes (Residual)	18,515	14.8	68.3
White, non-Hispanic, female, 65 years and over				
...	All causes	828,691	100.0	4,787.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,219	30.2	1,445.5
2	Malignant neoplasms (C00-C97)	159,226	19.2	919.8
3	Cerebrovascular diseases (I60-I69)	69,731	8.4	402.8
4	Chronic lower respiratory diseases (J40-J47)	50,971	6.2	294.5
5	Alzheimer's disease (G30)	41,854	5.1	241.8
6	Influenza and pneumonia (J10-J18)	25,628	3.1	148.1
7	Diabetes mellitus (E10-E14)	21,570	2.6	124.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,738	1.9	90.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,561	1.8	84.1
10	Septicemia (A40-A41)	11,568	1.4	66.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,806	1.2	56.6
12	Parkinson's disease (G20-G21)	6,837	0.8	39.5
13	Pneumonitis due to solids and liquids (J69)	6,633	0.8	38.3
14	Atherosclerosis (I70)	6,344	0.8	36.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,838	0.6	27.9
...	All other causes (Residual)	133,167	16.1	769.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	283,859	100.0	768.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,373	25.8	198.7
2	Malignant neoplasms (C00-C97)	61,842	21.8	167.5
3	Cerebrovascular diseases (I60-I69)	17,980	6.3	48.7
4	Diabetes mellitus (E10-E14)	12,685	4.5	34.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,446	4.4	33.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,000	2.8	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,759	2.7	21.0
8	Chronic lower respiratory diseases (J40-J47)	7,331	2.6	19.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,163	2.5	19.4
10	Septicemia (A40-A41)	5,945	2.1	16.1
11	Influenza and pneumonia (J10-J18)	5,620	2.0	15.2
12	Certain conditions originating in the perinatal period (P00-P96)	4,740	1.7	12.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,709	1.7	12.8
14	Alzheimer's disease (G30)	4,196	1.5	11.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,506	0.9	6.8
...	All other causes (Residual)	47,564	16.8	128.8
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,128	100.0	45.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	28.1	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	13.7	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	11.3	5.1
4	Malignant neoplasms (C00-C97)	62	5.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.4	2.5
6	Influenza and pneumonia (J10-J18)	34	3.0	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	23	2.0	0.9
8	Chronic lower respiratory diseases (J40-J47)	21	1.9	0.8
9	Septicemia (A40-A41)	19	1.7	*
9	Anemias (D50-D64)	19	1.7	*
11	Cerebrovascular diseases (I60-I69)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	257	22.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,551	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	35.9	8.8
2	Malignant neoplasms (C00-C97)	165	10.6	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	8.4	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	64	4.1	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.2	0.8
8	Anemias (D50-D64)	23	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.4	0.3
10	Influenza and pneumonia (J10-J18)	20	1.3	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.1	*
11	Cerebrovascular diseases (I60-I69)	17	1.1	*
13	Diabetes mellitus (E10-E14)	16	1.0	*
14	Septicemia (A40-A41)	12	0.8	*
15	Certain conditions originating in the perinatal period (P00-P96)	9	0.6	*
...	All other causes (Residual)	291	18.8	4.6
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,924	100.0	112.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,771	40.0	45.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,614	23.3	26.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	450	6.5	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	4.5	5.0
5	Malignant neoplasms (C00-C97)	307	4.4	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	141	2.0	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	1.6	1.8
8	Anemias (D50-D64)	90	1.3	1.5
9	Chronic lower respiratory diseases (J40-J47)	79	1.1	1.3
10	Pregnancy, childbirth and the puerperium (O00-099)	73	1.1	1.2
11	Diabetes mellitus (E10-E14)	60	0.9	1.0
12	Cerebrovascular diseases (I60-I69)	48	0.7	0.8
13	Septicemia (A40-A41)	45	0.6	0.7
14	Influenza and pneumonia (J10-J18)	40	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.4	0.5
...	All other causes (Residual)	754	10.9	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	28,005	100.0	261.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,173	14.9	38.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,776	13.5	35.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,681	13.1	34.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	3,458	12.3	32.2
5	Malignant neoplasms (C00-C97)	3,300	11.8	30.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	879	3.1	8.2
7	Cerebrovascular diseases (I60-I69)	871	3.1	8.1
8	Diabetes mellitus (E10-E14)	720	2.6	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	1.5	3.9
10	Septicemia (A40-A41)	357	1.3	3.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	329	1.2	3.1
12	Chronic lower respiratory diseases (J40-J47)	321	1.1	3.0
13	Influenza and pneumonia (J10-J18)	286	1.0	2.7
14	Anemias (D50-D64)	265	0.9	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	232	0.8	2.2
...	All other causes (Residual)	4,938	17.6	46.0
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	79,245	100.0	1,045.6
1	Malignant neoplasms (C00-C97)	21,810	27.5	287.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,113	25.4	265.4
3	Cerebrovascular diseases (I60-I69)	4,325	5.5	57.1
4	Diabetes mellitus (E10-E14)	3,675	4.6	48.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,669	4.6	48.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,264	4.1	43.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,096	2.6	27.7
8	Chronic lower respiratory diseases (J40-J47)	1,722	2.2	22.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,618	2.0	21.3
10	Septicemia (A40-A41)	1,556	2.0	20.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,182	1.5	15.6
12	Influenza and pneumonia (J10-J18)	980	1.2	12.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	979	1.2	12.9
14	Viral hepatitis (B15-B19)	609	0.8	8.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	443	0.6	5.8
...	All other causes (Residual)	11,204	14.1	147.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent	
			of total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	158,751	100.0	5,282.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,529	30.6	1,614.8
2	Malignant neoplasms (C00-C97)	36,190	22.8	1,204.3
3	Cerebrovascular diseases (I60-I69)	12,666	8.0	421.5
4	Diabetes mellitus (E10-E14)	8,211	5.2	273.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,141	3.2	171.1
6	Chronic lower respiratory diseases (J40-J47)	5,113	3.2	170.1
7	Influenza and pneumonia (J10-J18)	4,167	2.6	138.7
8	Alzheimer's disease (G30)	4,154	2.6	138.2
9	Septicemia (A40-A41)	3,865	2.4	128.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,284	2.1	109.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,214	1.4	73.7
12	Pneumonitis due to solids and liquids (J69)	1,214	0.8	40.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	732	0.5	24.4
14	Atherosclerosis (I70)	722	0.5	24.0
15	Aortic aneurysm and dissection (I71)	694	0.4	23.1
...	All other causes (Residual)	21,855	13.8	727.2
Black, non-Hispanic, male, all ages				
...	All causes	144,022	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,749	24.8	203.2
2	Malignant neoplasms (C00-C97)	32,089	22.3	182.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,430	5.9	47.9
4	Cerebrovascular diseases (I60-I69)	7,586	5.3	43.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,730	4.7	38.3
6	Diabetes mellitus (E10-E14)	5,498	3.8	31.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,693	3.3	26.7
8	Chronic lower respiratory diseases (J40-J47)	4,010	2.8	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,505	2.4	19.9
10	Influenza and pneumonia (J10-J18)	2,729	1.9	15.5
11	Septicemia (A40-A41)	2,660	1.8	15.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,653	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,945	1.4	11.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,650	1.1	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,614	1.1	9.2
...	All other causes (Residual)	22,481	15.6	127.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	623	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	29.2	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	12.7	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	11.9	5.9
4	Malignant neoplasms (C00-C97)	35	5.6	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.1	2.5
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.1	*
8	Anemias (D50-D64)	12	1.9	*
8	Influenza and pneumonia (J10-J18)	12	1.9	*
10	Septicemia (A40-A41)	9	1.4	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	141	22.6	11.2
Black, non-Hispanic, male, 5-14 years				
...	All causes	870	100.0	26.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	343	39.4	10.6
2	Malignant neoplasms (C00-C97)	82	9.4	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	9.3	2.5
4	Chronic lower respiratory diseases (J40-J47)	44	5.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	4.4	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.0	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.4	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
8	Anemias (D50-D64)	12	1.4	*
8	Influenza and pneumonia (J10-J18)	12	1.4	*
11	Cerebrovascular diseases (I60-I69)	10	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
15	Septicemia (A40-A41)	4	0.5	*
...	All other causes (Residual)	146	16.8	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,239	100.0	169.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,495	47.6	80.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,191	22.7	38.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	382	7.3	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	3.9	6.5
5	Malignant neoplasms (C00-C97)	174	3.3	5.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	73	1.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.4	2.4
8	Anemias (D50-D64)	50	1.0	1.6
9	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.6
10	Diabetes mellitus (E10-E14)	30	0.6	1.0
10	Cerebrovascular diseases (I60-I69)	30	0.6	1.0
12	Septicemia (A40-A41)	22	0.4	0.7
13	Influenza and pneumonia (J10-J18)	19	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
13	Legal intervention (Y35,Y89.0)	19	0.4	*
...	All other causes (Residual)	412	7.9	13.3
Black, non-Hispanic, male, 25-44 years				
...	All causes	17,116	100.0	336.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,135	18.3	61.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,737	16.0	53.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,524	14.7	49.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,102	12.3	41.4
5	Malignant neoplasms (C00-C97)	1,279	7.5	25.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	713	4.2	14.0
7	Cerebrovascular diseases (I60-I69)	421	2.5	8.3
8	Diabetes mellitus (E10-E14)	415	2.4	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	1.3	4.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	211	1.2	4.2
11	Septicemia (A40-A41)	177	1.0	3.5
12	Chronic lower respiratory diseases (J40-J47)	173	1.0	3.4
13	Influenza and pneumonia (J10-J18)	167	1.0	3.3
14	Anemias (D50-D64)	144	0.8	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	132	0.8	2.6
...	All other causes (Residual)	2,569	15.0	50.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	46,499	100.0	1,346.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,542	27.0	363.2
2	Malignant neoplasms (C00-C97)	11,715	25.2	339.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,615	5.6	75.7
4	Cerebrovascular diseases (I60-I69)	2,393	5.1	69.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,316	5.0	67.1
6	Diabetes mellitus (E10-E14)	1,919	4.1	55.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,114	2.4	32.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,109	2.4	32.1
9	Chronic lower respiratory diseases (J40-J47)	961	2.1	27.8
10	Septicemia (A40-A41)	799	1.7	23.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	764	1.6	22.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	652	1.4	18.9
13	Influenza and pneumonia (J10-J18)	577	1.2	16.7
14	Viral hepatitis (B15-B19)	439	0.9	12.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	354	0.8	10.3
...	All other causes (Residual)	6,230	13.4	180.4
Black, non-Hispanic, male, 65 years and over				
...	All causes	69,037	100.0	5,999.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,350	29.5	1,768.4
2	Malignant neoplasms (C00-C97)	18,801	27.2	1,633.8
3	Cerebrovascular diseases (I60-I69)	4,704	6.8	408.8
4	Diabetes mellitus (E10-E14)	3,124	4.5	271.5
5	Chronic lower respiratory diseases (J40-J47)	2,760	4.0	239.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,110	3.1	183.4
7	Influenza and pneumonia (J10-J18)	1,887	2.7	164.0
8	Septicemia (A40-A41)	1,598	2.3	138.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,202	1.7	104.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,156	1.7	100.5
11	Alzheimer's disease (G30)	1,121	1.6	97.4
12	Pneumonitis due to solids and liquids (J69)	596	0.9	51.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	328	0.5	28.5
14	Parkinson's disease (G20-G21)	321	0.5	27.9
15	Aortic aneurysm and dissection (I71)	318	0.5	27.6
...	All other causes (Residual)	8,661	12.5	752.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,837	100.0	723.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,624	26.9	194.6
2	Malignant neoplasms (C00-C97)	29,753	21.3	153.9
3	Cerebrovascular diseases (I60-I69)	10,394	7.4	53.8
4	Diabetes mellitus (E10-E14)	7,187	5.1	37.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,254	3.0	22.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,016	2.9	20.8
7	Chronic lower respiratory diseases (J40-J47)	3,321	2.4	17.2
8	Septicemia (A40-A41)	3,285	2.3	17.0
9	Alzheimer's disease (G30)	3,058	2.2	15.8
10	Influenza and pneumonia (J10-J18)	2,891	2.1	15.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,764	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,470	1.8	12.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,087	1.5	10.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,270	0.9	6.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	856	0.6	4.4
...	All other causes (Residual)	24,607	17.6	127.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	505	100.0	41.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	26.7	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	14.9	6.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	10.5	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.7	2.4
5	Malignant neoplasms (C00-C97)	27	5.3	2.2
6	Influenza and pneumonia (J10-J18)	22	4.4	1.8
7	Septicemia (A40-A41)	10	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	2.0	*
9	Anemias (D50-D64)	7	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	111	22.0	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	681	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	31.4	6.8
2	Malignant neoplasms (C00-C97)	83	12.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.2	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	6.3	1.4
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	0.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.8	*
8	Anemias (D50-D64)	11	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Septicemia (A40-A41)	8	1.2	*
11	Diabetes mellitus (E10-E14)	8	1.2	*
11	Influenza and pneumonia (J10-J18)	8	1.2	*
14	Cerebrovascular diseases (I60-I69)	7	1.0	*
15	Meningitis (G00,G03)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
...	All other causes (Residual)	141	20.7	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,685	100.0	55.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	423	25.1	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	16.4	9.1
3	Malignant neoplasms (C00-C97)	133	7.9	4.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	6.4	3.5
5	Pregnancy, childbirth and the puerperium (O00-099)	73	4.3	2.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	68	4.0	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	4.0	2.2
8	Anemias (D50-D64)	40	2.4	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	2.3	1.2
10	Chronic lower respiratory diseases (J40-J47)	31	1.8	1.0
11	Diabetes mellitus (E10-E14)	30	1.8	1.0
12	Septicemia (A40-A41)	23	1.4	0.8
13	Influenza and pneumonia (J10-J18)	21	1.2	0.7
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.7	*
...	All other causes (Residual)	323	19.2	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-44 years				
...	All causes	10,889	100.0	192.9
1	Malignant neoplasms (C00-C97)	2,021	18.6	35.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,649	15.1	29.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,356	12.5	24.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,039	9.5	18.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	5.0	9.7
6	Cerebrovascular diseases (I60-I69)	450	4.1	8.0
7	Diabetes mellitus (E10-E14)	305	2.8	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	1.9	3.6
9	Septicemia (A40-A41)	180	1.7	3.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	166	1.5	2.9
11	Pregnancy, childbirth and the puerperium (O00-099)	150	1.4	2.7
12	Chronic lower respiratory diseases (J40-J47)	148	1.4	2.6
13	Anemias (D50-D64)	121	1.1	2.1
14	Influenza and pneumonia (J10-J18)	119	1.1	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	118	1.1	2.1
...	All other causes (Residual)	2,319	21.3	41.1
Black, non-Hispanic, female, 45-64 years				
...	All causes	32,746	100.0	793.8
1	Malignant neoplasms (C00-C97)	10,095	30.8	244.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,571	23.1	183.5
3	Cerebrovascular diseases (I60-I69)	1,932	5.9	46.8
4	Diabetes mellitus (E10-E14)	1,756	5.4	42.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,054	3.2	25.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	982	3.0	23.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	948	2.9	23.0
8	Chronic lower respiratory diseases (J40-J47)	761	2.3	18.4
9	Septicemia (A40-A41)	757	2.3	18.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	530	1.6	12.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	509	1.6	12.3
12	Influenza and pneumonia (J10-J18)	403	1.2	9.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	215	0.7	5.2
14	Viral hepatitis (B15-B19)	170	0.5	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.4	3.2
...	All other causes (Residual)	4,932	15.1	119.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	89,714	100.0	4,837.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,179	31.4	1,519.5
2	Malignant neoplasms (C00-C97)	17,389	19.4	937.7
3	Cerebrovascular diseases (I60-I69)	7,962	8.9	429.3
4	Diabetes mellitus (E10-E14)	5,087	5.7	274.3
5	Alzheimer's disease (G30)	3,033	3.4	163.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,031	3.4	163.4
7	Chronic lower respiratory diseases (J40-J47)	2,353	2.6	126.9
8	Influenza and pneumonia (J10-J18)	2,280	2.5	122.9
9	Septicemia (A40-A41)	2,267	2.5	122.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,128	2.4	114.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	1.1	54.6
12	Pneumonitis due to solids and liquids (J69)	618	0.7	33.3
13	Atherosclerosis (I70)	479	0.5	25.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	421	0.5	22.7
15	Aortic aneurysm and dissection (I71)	376	0.4	20.3
...	All other causes (Residual)	13,099	14.6	706.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable