

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, both sexes				
...	All causes	28,440	100.0	687.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5,552	19.5	134.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,714	16.6	113.9
3	Sudden infant death syndrome (R95)	2,230	7.8	53.9
4	Newborn affected by maternal complications of pregnancy (P01)	1,776	6.2	42.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	1,110	3.9	26.8
6	Accidents (unintentional injuries) (V01-X59)	1,083	3.8	26.2
7	Respiratory distress of newborn (P22)	860	3.0	20.8
8	Bacterial sepsis of newborn (P36)	834	2.9	20.2
9	Neonatal hemorrhage (P50-P52,P54)	665	2.3	16.1
10	Necrotizing enterocolitis of newborn (P77)	546	1.9	13.2
11	Diseases of the circulatory system (I00-I99)	529	1.9	12.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	529	1.9	12.8
13	Atelectasis (P28.0-P28.1)	377	1.3	9.1
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	341	1.2	8.2
15	Assault (homicide) (*U01,X85-Y09)	306	1.1	7.4
...	All other causes (Residual)	6,988	24.6	168.9
Infants, all races, male				
...	All causes	16,018	100.0	755.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,965	18.5	139.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,644	16.5	124.8
3	Sudden infant death syndrome (R95)	1,372	8.6	64.7
4	Newborn affected by maternal complications of pregnancy (P01)	943	5.9	44.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	611	3.8	28.8
6	Accidents (unintentional injuries) (V01-X59)	578	3.6	27.3
7	Respiratory distress of newborn (P22)	518	3.2	24.4
8	Bacterial sepsis of newborn (P36)	478	3.0	22.6
9	Neonatal hemorrhage (P50-P52,P54)	398	2.5	18.8
10	Necrotizing enterocolitis of newborn (P77)	316	2.0	14.9
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	300	1.9	14.2
12	Diseases of the circulatory system (I00-I99)	295	1.8	13.9
13	Atelectasis (P28.0-P28.1)	220	1.4	10.4
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	211	1.3	10.0
15	Assault (homicide) (*U01,X85-Y09)	173	1.1	8.2
...	All other causes (Residual)	3,996	24.9	188.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, female				
...	All causes	12,422	100.0	615.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,587	20.8	128.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,070	16.7	102.5
3	Sudden infant death syndrome (R95)	858	6.9	42.5
4	Newborn affected by maternal complications of pregnancy (P01)	833	6.7	41.3
5	Accidents (unintentional injuries) (V01-X59)	505	4.1	25.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	499	4.0	24.7
7	Bacterial sepsis of newborn (P36)	356	2.9	17.6
8	Respiratory distress of newborn (P22)	342	2.8	16.9
9	Neonatal hemorrhage (P50-P52,P54)	267	2.1	13.2
10	Diseases of the circulatory system (I00-I99)	234	1.9	11.6
11	Necrotizing enterocolitis of newborn (P77)	230	1.9	11.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	229	1.8	11.3
13	Atelectasis (P28.0-P28.1)	157	1.3	7.8
14	Assault (homicide) (*U01,X85-Y09)	133	1.1	6.6
15	Septicemia (A40-A41)	131	1.1	6.5
...	All other causes (Residual)	2,991	24.1	148.1
Infants, white, both sexes				
...	All causes	18,514	100.0	573.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4,187	22.6	129.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,623	14.2	81.2
3	Sudden infant death syndrome (R95)	1,493	8.1	46.2
4	Newborn affected by maternal complications of pregnancy (P01)	1,050	5.7	32.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	760	4.1	23.5
6	Accidents (unintentional injuries) (V01-X59)	734	4.0	22.7
7	Respiratory distress of newborn (P22)	534	2.9	16.5
8	Bacterial sepsis of newborn (P36)	520	2.8	16.1
9	Neonatal hemorrhage (P50-P52,P54)	466	2.5	14.4
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	388	2.1	12.0
11	Diseases of the circulatory system (I00-I99)	339	1.8	10.5
12	Necrotizing enterocolitis of newborn (P77)	313	1.7	9.7
13	Atelectasis (P28.0-P28.1)	263	1.4	8.1
14	Assault (homicide) (*U01,X85-Y09)	195	1.1	6.0
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	191	1.0	5.9
...	All other causes (Residual)	4,458	24.1	138.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, white, male				
...	All causes	10,471	100.0	632.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,239	21.4	135.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,454	13.9	87.8
3	Sudden infant death syndrome (R95)	959	9.2	57.9
4	Newborn affected by maternal complications of pregnancy (P01)	557	5.3	33.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	422	4.0	25.5
6	Accidents (unintentional injuries) (V01-X59)	388	3.7	23.4
7	Respiratory distress of newborn (P22)	331	3.2	20.0
8	Bacterial sepsis of newborn (P36)	306	2.9	18.5
9	Neonatal hemorrhage (P50-P52,P54)	277	2.6	16.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	224	2.1	13.5
11	Diseases of the circulatory system (I00-I99)	188	1.8	11.4
12	Necrotizing enterocolitis of newborn (P77)	184	1.8	11.1
13	Atelectasis (P28.0-P28.1)	155	1.5	9.4
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	115	1.1	6.9
15	Assault (homicide) (*U01,X85-Y09)	109	1.0	6.6
...	All other causes (Residual)	2,563	24.5	154.8
Infants, white, female				
...	All causes	8,043	100.0	511.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,948	24.2	123.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,169	14.5	74.3
3	Sudden infant death syndrome (R95)	534	6.6	33.9
4	Newborn affected by maternal complications of pregnancy (P01)	493	6.1	31.3
5	Accidents (unintentional injuries) (V01-X59)	346	4.3	22.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	338	4.2	21.5
7	Bacterial sepsis of newborn (P36)	214	2.7	13.6
8	Respiratory distress of newborn (P22)	203	2.5	12.9
9	Neonatal hemorrhage (P50-P52,P54)	189	2.3	12.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	164	2.0	10.4
11	Diseases of the circulatory system (I00-I99)	151	1.9	9.6
12	Necrotizing enterocolitis of newborn (P77)	129	1.6	8.2
13	Atelectasis (P28.0-P28.1)	108	1.3	6.9
14	Assault (homicide) (*U01,X85-Y09)	86	1.1	5.5
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	76	0.9	4.8
...	All other causes (Residual)	1,895	23.6	120.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, both sexes				
...	All causes	8,695	100.0	1,373.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,894	21.8	299.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,080	12.4	170.6
3	Sudden infant death syndrome (R95)	663	7.6	104.7
4	Newborn affected by maternal complications of pregnancy (P01)	660	7.6	104.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	320	3.7	50.5
6	Accidents (unintentional injuries) (V01-X59)	314	3.6	49.6
7	Respiratory distress of newborn (P22)	295	3.4	46.6
8	Bacterial sepsis of newborn (P36)	273	3.1	43.1
9	Necrotizing enterocolitis of newborn (P77)	207	2.4	32.7
10	Neonatal hemorrhage (P50-P52,P54)	169	1.9	26.7
11	Diseases of the circulatory system (I00-I99)	157	1.8	24.8
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	138	1.6	21.8
13	Septicemia (A40-A41)	131	1.5	20.7
14	Chronic respiratory disease originating in the perinatal period (P27)	127	1.5	20.1
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	121	1.4	19.1
...	All other causes (Residual)	2,146	24.7	338.9
Infants, black, male				
...	All causes	4,867	100.0	1,515.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,078	22.1	335.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	564	11.6	175.6
3	Sudden infant death syndrome (R95)	368	7.6	114.5
4	Newborn affected by maternal complications of pregnancy (P01)	349	7.2	108.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	174	3.6	54.2
6	Accidents (unintentional injuries) (V01-X59)	172	3.5	53.5
7	Respiratory distress of newborn (P22)	171	3.5	53.2
8	Bacterial sepsis of newborn (P36)	151	3.1	47.0
9	Necrotizing enterocolitis of newborn (P77)	116	2.4	36.1
10	Neonatal hemorrhage (P50-P52,P54)	103	2.1	32.1
11	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	92	1.9	28.6
12	Diseases of the circulatory system (I00-I99)	91	1.9	28.3
13	Chronic respiratory disease originating in the perinatal period (P27)	75	1.5	23.3
14	Septicemia (A40-A41)	69	1.4	21.5
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	63	1.3	19.6
...	All other causes (Residual)	1,231	25.3	383.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, female				
...	All causes	3,828	100.0	1,227.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	816	21.3	261.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	516	13.5	165.5
3	Newborn affected by maternal complications of pregnancy (P01)	311	8.1	99.7
4	Sudden infant death syndrome (R95)	295	7.7	94.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	146	3.8	46.8
6	Accidents (unintentional injuries) (V01-X59)	142	3.7	45.5
7	Respiratory distress of newborn (P22)	124	3.2	39.8
8	Bacterial sepsis of newborn (P36)	122	3.2	39.1
9	Necrotizing enterocolitis of newborn (P77)	91	2.4	29.2
10	Diseases of the circulatory system (I00-I99)	66	1.7	21.2
10	Neonatal hemorrhage (P50-P52,P54)	66	1.7	21.2
12	Septicemia (A40-A41)	62	1.6	19.9
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	58	1.5	18.6
14	Chronic respiratory disease originating in the perinatal period (P27)	52	1.4	16.7
15	Influenza and pneumonia (J10-J18)	51	1.3	16.4
...	All other causes (Residual)	910	23.8	291.8
Infants, American Indian, both sexes				
...	All causes	357	100.0	796.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	21.6	171.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	56	15.7	125.0
3	Sudden infant death syndrome (R95)	37	10.4	82.6
4	Accidents (unintentional injuries) (V01-X59)	19	5.3	*
5	Newborn affected by maternal complications of pregnancy (P01)	12	3.4	*
6	Diseases of the circulatory system (I00-I99)	9	2.5	*
7	Bacterial sepsis of newborn (P36)	8	2.2	*
8	Newborn affected by complications of placenta, cord and membranes (P02)	7	2.0	*
8	Necrotizing enterocolitis of newborn (P77)	7	2.0	*
10	Influenza and pneumonia (J10-J18)	5	1.4	*
10	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	1.4	*
10	Atelectasis (P28.0-P28.1)	5	1.4	*
10	Assault (homicide) (*U01,X85-Y09)	5	1.4	*
14	Respiratory distress of newborn (P22)	4	1.1	*
14	Neonatal hemorrhage (P50-P52,P54)	4	1.1	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	4	1.1	*
...	All other causes (Residual)	93	26.1	207.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, American Indian, male				
...	All causes	196	100.0	864.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	22.4	194.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	36	18.4	158.8
3	Sudden infant death syndrome (R95)	20	10.2	88.2
4	Accidents (unintentional injuries) (V01-X59)	10	5.1	*
5	Newborn affected by maternal complications of pregnancy (P01)	9	4.6	*
6	Necrotizing enterocolitis of newborn (P77)	5	2.6	*
7	Assault (homicide) (*U01,X85-Y09)	4	2.0	*
8	Diseases of the circulatory system (I00-I99)	3	1.5	*
8	Newborn affected by complications of placenta, cord and membranes (P02)	3	1.5	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	3	1.5	*
8	Respiratory distress of newborn (P22)	3	1.5	*
8	Atelectasis (P28.0-P28.1)	3	1.5	*
8	Bacterial sepsis of newborn (P36)	3	1.5	*
14	Whooping cough (A37)	2	1.0	*
14	Meningitis (G00,G03)	2	1.0	*
14	Influenza and pneumonia (J10-J18)	2	1.0	*
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	2	1.0	*
14	Neonatal hemorrhage (P50-P52,P54)	2	1.0	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	2	1.0	*
...	All other causes (Residual)	38	19.4	167.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, American Indian, female				
...	All causes	161	100.0	727.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	20.5	149.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	20	12.4	90.3
3	Sudden infant death syndrome (R95)	17	10.6	*
4	Accidents (unintentional injuries) (V01-X59)	9	5.6	*
5	Diseases of the circulatory system (I00-I99)	6	3.7	*
6	Bacterial sepsis of newborn (P36)	5	3.1	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	4	2.5	*
8	Influenza and pneumonia (J10-J18)	3	1.9	*
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	3	1.9	*
8	Newborn affected by maternal complications of pregnancy (P01)	3	1.9	*
8	Congenital pneumonia (P23)	3	1.9	*
12	Septicemia (A40-A41)	2	1.2	*
12	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	2	1.2	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	2	1.2	*
12	Newborn affected by maternal hypertensive disorders (P00.0)	2	1.2	*
12	Atelectasis (P28.0-P28.1)	2	1.2	*
12	Neonatal hemorrhage (P50-P52,P54)	2	1.2	*
12	Necrotizing enterocolitis of newborn (P77)	2	1.2	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	2	1.2	*
...	All other causes (Residual)	39	24.2	176.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, both sexes				
...	All causes	874	100.0	378.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	23.8	90.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	141	16.1	61.0
3	Newborn affected by maternal complications of pregnancy (P01)	54	6.2	23.4
4	Sudden infant death syndrome (R95)	37	4.2	16.0
5	Bacterial sepsis of newborn (P36)	33	3.8	14.3
6	Respiratory distress of newborn (P22)	27	3.1	11.7
7	Neonatal hemorrhage (P50-P52,P54)	26	3.0	11.3
8	Diseases of the circulatory system (I00-I99)	24	2.7	10.4
9	Newborn affected by complications of placenta, cord and membranes (P02)	23	2.6	10.0
10	Necrotizing enterocolitis of newborn (P77)	19	2.2	*
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	17	1.9	*
12	Accidents (unintentional injuries) (V01-X59)	16	1.8	*
13	Assault (homicide) (*U01,X85-Y09)	11	1.3	*
14	Septicemia (A40-A41)	9	1.0	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	9	1.0	*
...	All other causes (Residual)	220	25.2	95.2
Infants, Asian or Pacific Islander, male				
...	All causes	484	100.0	405.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	24.4	99.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	76	15.7	63.7
3	Newborn affected by maternal complications of pregnancy (P01)	28	5.8	23.5
4	Sudden infant death syndrome (R95)	25	5.2	21.0
5	Bacterial sepsis of newborn (P36)	18	3.7	*
6	Neonatal hemorrhage (P50-P52,P54)	16	3.3	*
7	Diseases of the circulatory system (I00-I99)	13	2.7	*
7	Respiratory distress of newborn (P22)	13	2.7	*
9	Newborn affected by complications of placenta, cord and membranes (P02)	12	2.5	*
10	Necrotizing enterocolitis of newborn (P77)	11	2.3	*
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	10	2.1	*
12	Accidents (unintentional injuries) (V01-X59)	8	1.7	*
13	Atelectasis (P28.0-P28.1)	7	1.4	*
14	Septicemia (A40-A41)	6	1.2	*
14	Pulmonary hemorrhage originating in the perinatal period (P26)	6	1.2	*
14	Assault (homicide) (*U01,X85-Y09)	6	1.2	*
...	All other causes (Residual)	111	22.9	93.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, female				
...	All causes	390	100.0	348.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	23.1	80.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	65	16.7	58.1
3	Newborn affected by maternal complications of pregnancy (P01)			23.2
4	Bacterial sepsis of newborn (P36)	26	6.7	
5	Respiratory distress of newborn (P22)	15	3.8	*
6	Sudden infant death syndrome (R95)	14	3.6	*
7	Diseases of the circulatory system (I00-I99)	12	3.1	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	11	2.8	*
9	Neonatal hemorrhage (P50-P52,P54)	11	2.8	*
10	Necrotizing enterocolitis of newborn (P77)	10	2.6	*
10	Accidents (unintentional injuries) (V01-X59)	8	2.1	*
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	8	2.1	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	1.8	*
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	1.3	*
13	Chronic respiratory disease originating in the perinatal period (P27)	5	1.3	*
13	Hydrops fetalis not due to hemolytic disease (P83.2)	5	1.3	*
13	Assault (homicide) (*U01,X85-Y09)	5	1.3	*
...	All other causes (Residual)	88	22.6	78.7
Neonates, all races, both sexes				
...	All causes	18,770	100.0	453.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,593	24.5	111.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,885	20.7	93.9
3	Newborn affected by maternal complications of pregnancy (P01)	1,755	9.4	42.4
4	Newborn affected by complications of placenta, cord and membranes (P02)			26.3
5	Respiratory distress of newborn (P22)	1,090	5.8	
6	Bacterial sepsis of newborn (P36)	832	4.4	20.1
7	Neonatal hemorrhage (P50-P52,P54)	790	4.2	19.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	652	3.5	15.8
9	Necrotizing enterocolitis of newborn (P77)	501	2.7	12.1
10	Atelectasis (P28.0-P28.1)	468	2.5	11.3
11	Sudden infant death syndrome (R95)	368	2.0	8.9
12	Pulmonary hemorrhage originating in the perinatal period (P26)	199	1.1	4.8
13	Hydrops fetalis not due to hemolytic disease (P83.2)	180	1.0	4.3
14	Newborn affected by other complications of labor and delivery (P03)	154	0.8	3.7
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	132	0.7	3.2
...	All other causes (Residual)	121	0.6	2.9
...	All other causes (Residual)	3,050	16.2	73.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, all races, male				
...	All causes	10,444	100.0	492.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,575	24.7	121.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,096	20.1	98.9
3	Newborn affected by maternal complications of pregnancy (P01)	930	8.9	43.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	597	5.7	28.2
5	Respiratory distress of newborn (P22)	507	4.9	23.9
6	Bacterial sepsis of newborn (P36)	446	4.3	21.0
7	Neonatal hemorrhage (P50-P52,P54)	389	3.7	18.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	285	2.7	13.4
9	Necrotizing enterocolitis of newborn (P77)	265	2.5	12.5
10	Atelectasis (P28.0-P28.1)	217	2.1	10.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	114	1.1	5.4
12	Sudden infant death syndrome (R95)	113	1.1	5.3
13	Hydrops fetalis not due to hemolytic disease (P83.2)	85	0.8	4.0
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	74	0.7	3.5
15	Newborn affected by other complications of labor and delivery (P03)	69	0.7	3.3
...	All other causes (Residual)	1,682	16.1	79.4
Neonates, all races, female				
...	All causes	8,326	100.0	412.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,018	24.2	99.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,789	21.5	88.6
3	Newborn affected by maternal complications of pregnancy (P01)	825	9.9	40.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	493	5.9	24.4
5	Bacterial sepsis of newborn (P36)	344	4.1	17.0
6	Respiratory distress of newborn (P22)	325	3.9	16.1
7	Neonatal hemorrhage (P50-P52,P54)	263	3.2	13.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	216	2.6	10.7
9	Necrotizing enterocolitis of newborn (P77)	203	2.4	10.1
10	Atelectasis (P28.0-P28.1)	151	1.8	7.5
11	Sudden infant death syndrome (R95)	86	1.0	4.3
12	Hydrops fetalis not due to hemolytic disease (P83.2)	69	0.8	3.4
13	Pulmonary hemorrhage originating in the perinatal period (P26)	66	0.8	3.3
13	Accidents (unintentional injuries) (V01-X59)	66	0.8	3.3
15	Newborn affected by other complications of labor and delivery (P03)	63	0.8	3.1
...	All other causes (Residual)	1,349	16.2	66.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, both sexes				
...	All causes	12,239	100.0	379.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,989	24.4	92.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,569	21.0	79.6
3	Newborn affected by maternal complications of pregnancy (P01)	1,036	8.5	32.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	749	6.1	23.2
5	Respiratory distress of newborn (P22)	519	4.2	16.1
6	Bacterial sepsis of newborn (P36)	495	4.0	15.3
7	Neonatal hemorrhage (P50-P52,P54)	455	3.7	14.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	371	3.0	11.5
9	Necrotizing enterocolitis of newborn (P77)	271	2.2	8.4
10	Atelectasis (P28.0-P28.1)	258	2.1	8.0
11	Sudden infant death syndrome (R95)	145	1.2	4.5
12	Hydrops fetalis not due to hemolytic disease (P83.2)	127	1.0	3.9
13	Pulmonary hemorrhage originating in the perinatal period (P26)	109	0.9	3.4
14	Newborn affected by other complications of labor and delivery (P03)	83	0.7	2.6
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	83	0.7	2.6
...	All other causes (Residual)	1,980	16.2	61.3
Neonates, white, male				
...	All causes	6,796	100.0	410.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,605	23.6	96.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,425	21.0	86.1
3	Newborn affected by maternal complications of pregnancy (P01)	549	8.1	33.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	414	6.1	25.0
5	Respiratory distress of newborn (P22)	323	4.8	19.5
6	Bacterial sepsis of newborn (P36)	288	4.2	17.4
7	Neonatal hemorrhage (P50-P52,P54)	270	4.0	16.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	214	3.1	12.9
9	Necrotizing enterocolitis of newborn (P77)	153	2.3	9.2
10	Atelectasis (P28.0-P28.1)	152	2.2	9.2
11	Sudden infant death syndrome (R95)	85	1.3	5.1
12	Pulmonary hemorrhage originating in the perinatal period (P26)	71	1.0	4.3
13	Hydrops fetalis not due to hemolytic disease (P83.2)	69	1.0	4.2
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	49	0.7	3.0
15	Hematological disorders (P60-P61)	46	0.7	2.8
...	All other causes (Residual)	1,083	15.9	65.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, female				
...	All causes	5,443	100.0	345.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,384	25.4	88.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,144	21.0	72.7
3	Newborn affected by maternal complications of pregnancy (P01)	487	8.9	31.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	335	6.2	21.3
5	Bacterial sepsis of newborn (P36)	207	3.8	13.2
6	Respiratory distress of newborn (P22)	196	3.6	12.5
7	Neonatal hemorrhage (P50-P52,P54)	185	3.4	11.8
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	157	2.9	10.0
9	Necrotizing enterocolitis of newborn (P77)	118	2.2	7.5
10	Atelectasis (P28.0-P28.1)	106	1.9	6.7
11	Sudden infant death syndrome (R95)	60	1.1	3.8
12	Hydrops fetalis not due to hemolytic disease (P83.2)	58	1.1	3.7
13	Accidents (unintentional injuries) (V01-X59)	44	0.8	2.8
14	Newborn affected by other complications of labor and delivery (P03)	40	0.7	2.5
15	Pulmonary hemorrhage originating in the perinatal period (P26)	38	0.7	2.4
...	All other causes (Residual)	884	16.2	56.2
Neonates, black, both sexes				
...	All causes	5,740	100.0	906.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,836	32.0	290.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	713	12.4	112.6
3	Newborn affected by maternal complications of pregnancy (P01)	653	11.4	103.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	311	5.4	49.1
5	Respiratory distress of newborn (P22)	282	4.9	44.5
6	Bacterial sepsis of newborn (P36)	254	4.4	40.1
7	Necrotizing enterocolitis of newborn (P77)	174	3.0	27.5
8	Neonatal hemorrhage (P50-P52,P54)	167	2.9	26.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	110	1.9	17.4
10	Atelectasis (P28.0-P28.1)	97	1.7	15.3
11	Pulmonary hemorrhage originating in the perinatal period (P26)	63	1.1	10.0
12	Sudden infant death syndrome (R95)	52	0.9	8.2
13	Newborn affected by other complications of labor and delivery (P03)	44	0.8	6.9
14	Newborn affected by maternal hypertensive disorders (P00.0)	37	0.6	5.8
14	Accidents (unintentional injuries) (V01-X59)	37	0.6	5.8
...	All other causes (Residual)	910	15.9	143.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, male				
...	All causes	3,201	100.0	996.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,045	32.6	325.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	384	12.0	119.5
3	Newborn affected by maternal complications of pregnancy (P01)	344	10.7	107.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	168	5.2	52.3
4	Respiratory distress of newborn (P22)	168	5.2	52.3
6	Bacterial sepsis of newborn (P36)	137	4.3	42.6
7	Neonatal hemorrhage (P50-P52,P54)	101	3.2	31.4
8	Necrotizing enterocolitis of newborn (P77)	97	3.0	30.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	58	1.8	18.1
10	Atelectasis (P28.0-P28.1)	55	1.7	17.1
11	Pulmonary hemorrhage originating in the perinatal period (P26)	37	1.2	11.5
12	Sudden infant death syndrome (R95)	27	0.8	8.4
13	Newborn affected by other complications of labor and delivery (P03)	24	0.7	7.5
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	21	0.7	6.5
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.6	*
...	All other causes (Residual)	516	16.1	160.6
Neonates, black, female				
...	All causes	2,539	100.0	814.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	791	31.2	253.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	329	13.0	105.5
3	Newborn affected by maternal complications of pregnancy (P01)	309	12.2	99.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	143	5.6	45.9
5	Bacterial sepsis of newborn (P36)	117	4.6	37.5
6	Respiratory distress of newborn (P22)	114	4.5	36.6
7	Necrotizing enterocolitis of newborn (P77)	77	3.0	24.7
8	Neonatal hemorrhage (P50-P52,P54)	66	2.6	21.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	52	2.0	16.7
10	Atelectasis (P28.0-P28.1)	42	1.7	13.5
11	Newborn affected by maternal hypertensive disorders (P00.0)	26	1.0	8.3
11	Pulmonary hemorrhage originating in the perinatal period (P26)	26	1.0	8.3
13	Sudden infant death syndrome (R95)	25	1.0	8.0
14	Accidents (unintentional injuries) (V01-X59)	21	0.8	6.7
15	Newborn affected by other complications of labor and delivery (P03)	20	0.8	6.4
...	All other causes (Residual)	381	15.0	122.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, both sexes				
...	All causes	186	100.0	415.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	50	26.9	111.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	26.3	109.3
3	Newborn affected by maternal complications of pregnancy (P01)	12	6.5	*
4	Bacterial sepsis of newborn (P36)	8	4.3	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	7	3.8	*
6	Necrotizing enterocolitis of newborn (P77)	6	3.2	*
7	Atelectasis (P28.0-P28.1)	5	2.7	*
8	Respiratory distress of newborn (P22)	4	2.2	*
8	Neonatal hemorrhage (P50-P52,P54)	4	2.2	*
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	3	1.6	*
10	Congenital pneumonia (P23)	3	1.6	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	3	1.6	*
13	Diseases of the circulatory system (I00-I99)	2	1.1	*
13	Newborn affected by maternal hypertensive disorders (P00.0)	2	1.1	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	0.5	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.5	*
15	Newborn affected by other complications of labor and delivery (P03)	1	0.5	*
15	Birth trauma (P10-P15)	1	0.5	*
15	Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0-P70.2)	1	0.5	*
15	Sudden infant death syndrome (R95)	1	0.5	*
...	All other causes (Residual)	22	11.8	49.1
Neonates, American Indian, male				
...	All causes	107	100.0	471.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	29.0	136.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	27.1	127.9
3	Newborn affected by maternal complications of pregnancy (P01)	9	8.4	*
4	Necrotizing enterocolitis of newborn (P77)	5	4.7	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	3	2.8	*
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	3	2.8	*
5	Respiratory distress of newborn (P22)	3	2.8	*
5	Atelectasis (P28.0-P28.1)	3	2.8	*
5	Bacterial sepsis of newborn (P36)	3	2.8	*
10	Neonatal hemorrhage (P50-P52,P54)	2	1.9	*
11	Diseases of the circulatory system (I00-I99)	1	0.9	*
11	Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0-P70.2)	1	0.9	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.9	*
...	All other causes (Residual)	13	12.1	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, female				
...	All causes	79	100.0	356.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	25.3	90.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	19	24.1	*
3	Bacterial sepsis of newborn (P36)	5	6.3	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	4	5.1	*
5	Newborn affected by maternal complications of pregnancy (P01)	3	3.8	*
5	Congenital pneumonia (P23)	3	3.8	*
7	Newborn affected by maternal hypertensive disorders (P00.0)	2	2.5	*
7	Atelectasis (P28.0-P28.1)	2	2.5	*
7	Neonatal hemorrhage (P50-P52,P54)	2	2.5	*
7	Hydrops fetalis not due to hemolytic disease (P83.2)	2	2.5	*
11	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	1.3	*
11	Diseases of the circulatory system (I00-I99)	1	1.3	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	1.3	*
11	Newborn affected by other complications of labor and delivery (P03)	1	1.3	*
11	Birth trauma (P10-P15)	1	1.3	*
11	Respiratory distress of newborn (P22)	1	1.3	*
11	Necrotizing enterocolitis of newborn (P77)	1	1.3	*
11	Sudden infant death syndrome (R95)	1	1.3	*
...	All other causes (Residual)	9	11.4	*
Neonates, Asian or Pacific Islander, both sexes				
...	All causes	605	100.0	261.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	138	22.8	59.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	22.1	58.0
3	Newborn affected by maternal complications of pregnancy (P01)	54	8.9	23.4
4	Bacterial sepsis of newborn (P36)	33	5.5	14.3
5	Respiratory distress of newborn (P22)	27	4.5	11.7
6	Neonatal hemorrhage (P50-P52,P54)	26	4.3	11.3
7	Newborn affected by complications of placenta, cord and membranes (P02)	23	3.8	10.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	17	2.8	*
8	Necrotizing enterocolitis of newborn (P77)	17	2.8	*
10	Pulmonary hemorrhage originating in the perinatal period (P26)	8	1.3	*
10	Atelectasis (P28.0-P28.1)	8	1.3	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	8	1.3	*
13	Diseases of the circulatory system (I00-I99)	7	1.2	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	5	0.8	*
14	Hematological disorders (P60-P61)	5	0.8	*
...	All other causes (Residual)	95	15.7	41.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Asian or Pacific Islander, male				
...	All causes	340	100.0	285.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	22.9	65.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	74	21.8	62.1
3	Newborn affected by maternal complications of pregnancy (P01)			23.5
4	Bacterial sepsis of newborn (P36)	18	5.3	*
5	Neonatal hemorrhage (P50-P52,P54)	16	4.7	*
6	Respiratory distress of newborn (P22)	13	3.8	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	12	3.5	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	10	2.9	*
8	Necrotizing enterocolitis of newborn (P77)	10	2.9	*
10	Atelectasis (P28.0-P28.1)	7	2.1	*
11	Pulmonary hemorrhage originating in the perinatal period (P26)	6	1.8	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	4	1.2	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	4	1.2	*
14	Diseases of the circulatory system (I00-I99)	3	0.9	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	3	0.9	*
14	Hematological disorders (P60-P61)	3	0.9	*
14	Assault (homicide) (*U01,X85-Y09)	3	0.9	*
...	All other causes (Residual)	48	14.1	40.3
Neonates, Asian or Pacific Islander, female				
...	All causes	265	100.0	236.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	64	24.2	57.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	21.1	50.1
3	Newborn affected by maternal complications of pregnancy (P01)			23.2
4	Bacterial sepsis of newborn (P36)	15	5.7	*
5	Respiratory distress of newborn (P22)	14	5.3	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	11	4.2	*
7	Neonatal hemorrhage (P50-P52,P54)	10	3.8	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	7	2.6	*
8	Necrotizing enterocolitis of newborn (P77)	7	2.6	*
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	4	1.5	*
10	Diseases of the circulatory system (I00-I99)	4	1.5	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	4	1.5	*
13	Slow fetal growth and fetal malnutrition (P05)	3	1.1	*
14	Newborn affected by other complications of labor and delivery (P03)	2	0.8	*
14	Pulmonary hemorrhage originating in the perinatal period (P26)	2	0.8	*
14	Hematological disorders (P60-P61)	2	0.8	*
...	All other causes (Residual)	34	12.8	30.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races, both sexes				
...	All causes	9,670	100.0	233.7
1	Sudden infant death syndrome (R95)	2,031	21.0	49.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,667	17.2	40.3
3	Accidents (unintentional injuries) (V01-X59)	963	10.0	23.3
4	Diseases of the circulatory system (I00-I99)	414	4.3	10.0
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	317	3.3	7.7
6	Septicemia (A40-A41)	302	3.1	7.3
7	Assault (homicide) (*U01,X85-Y09)	270	2.8	6.5
8	Influenza and pneumonia (J10-J18)	263	2.7	6.4
9	Chronic respiratory disease originating in the perinatal period (P27)	240	2.5	5.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	121	1.3	2.9
11	Necrotizing enterocolitis of newborn (P77)	78	0.8	1.9
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	75	0.8	1.8
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	64	0.7	1.5
14	Malignant neoplasms (C00-C97)	59	0.6	1.4
15	Acute bronchitis and acute bronchiolitis (J20-J21)	48	0.5	1.2
...	All other causes (Residual)	2,758	28.5	66.6
Postneonates, all races, male				
...	All causes	5,574	100.0	263.1
1	Sudden infant death syndrome (R95)	1,259	22.6	59.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	869	15.6	41.0
3	Accidents (unintentional injuries) (V01-X59)	524	9.4	24.7
4	Diseases of the circulatory system (I00-I99)	234	4.2	11.0
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	194	3.5	9.2
6	Septicemia (A40-A41)	171	3.1	8.1
7	Assault (homicide) (*U01,X85-Y09)	156	2.8	7.4
8	Influenza and pneumonia (J10-J18)	147	2.6	6.9
9	Chronic respiratory disease originating in the perinatal period (P27)	135	2.4	6.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	69	1.2	3.3
11	Necrotizing enterocolitis of newborn (P77)	51	0.9	2.4
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	43	0.8	2.0
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	41	0.7	1.9
14	Malignant neoplasms (C00-C97)	37	0.7	1.7
15	Bacterial sepsis of newborn (P36)	32	0.6	1.5
...	All other causes (Residual)	1,612	28.9	76.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races, female				
...	All causes	4,096	100.0	202.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	798	19.5	39.5
2	Sudden infant death syndrome (R95)	772	18.8	38.2
3	Accidents (unintentional injuries) (V01-X59)	439	10.7	21.7
4	Diseases of the circulatory system (I00-I99)	180	4.4	8.9
5	Septicemia (A40-A41)	131	3.2	6.5
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	123	3.0	6.1
7	Influenza and pneumonia (J10-J18)	116	2.8	5.7
8	Assault (homicide) (*U01,X85-Y09)	114	2.8	5.6
9	Chronic respiratory disease originating in the perinatal period (P27)	105	2.6	5.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	52	1.3	2.6
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	32	0.8	1.6
12	Necrotizing enterocolitis of newborn (P77)	27	0.7	1.3
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	23	0.6	1.1
14	Malignant neoplasms (C00-C97)	22	0.5	1.1
15	Meningitis (G00,G03)	21	0.5	1.0
...	All other causes (Residual)	1,141	27.9	56.5
Postneonates, white, both sexes				
...	All causes	6,275	100.0	194.3
1	Sudden infant death syndrome (R95)	1,348	21.5	41.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,198	19.1	37.1
3	Accidents (unintentional injuries) (V01-X59)	652	10.4	20.2
4	Diseases of the circulatory system (I00-I99)	264	4.2	8.2
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	177	2.8	5.5
6	Assault (homicide) (*U01,X85-Y09)	170	2.7	5.3
7	Septicemia (A40-A41)	160	2.5	5.0
8	Influenza and pneumonia (J10-J18)	156	2.5	4.8
9	Chronic respiratory disease originating in the perinatal period (P27)	122	1.9	3.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	54	0.9	1.7
11	Malignant neoplasms (C00-C97)	49	0.8	1.5
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	44	0.7	1.4
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	44	0.7	1.4
14	Necrotizing enterocolitis of newborn (P77)	42	0.7	1.3
15	Acute bronchitis and acute bronchiolitis (J20-J21)	28	0.4	0.9
...	All other causes (Residual)	1,767	28.2	54.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, male				
...	All causes	3,675	100.0	221.9
1	Sudden infant death syndrome (R95)	874	23.8	52.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	634	17.3	38.3
3	Accidents (unintentional injuries) (V01-X59)	350	9.5	21.1
4	Diseases of the circulatory system (I00-I99)	149	4.1	9.0
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	107	2.9	6.5
6	Assault (homicide) (*U01,X85-Y09)	98	2.7	5.9
7	Septicemia (A40-A41)	96	2.6	5.8
8	Influenza and pneumonia (J10-J18)	95	2.6	5.7
9	Chronic respiratory disease originating in the perinatal period (P27)	68	1.9	4.1
10	Malignant neoplasms (C00-C97)	31	0.8	1.9
10	Necrotizing enterocolitis of newborn (P77)	31	0.8	1.9
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	29	0.8	1.8
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	26	0.7	1.6
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	24	0.7	1.4
15	Bacterial sepsis of newborn (P36)	18	0.5	*
...	All other causes (Residual)	1,045	28.4	63.1
Postneonates, white, female				
...	All causes	2,600	100.0	165.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	564	21.7	35.8
2	Sudden infant death syndrome (R95)	474	18.2	30.1
3	Accidents (unintentional injuries) (V01-X59)	302	11.6	19.2
4	Diseases of the circulatory system (I00-I99)	115	4.4	7.3
5	Assault (homicide) (*U01,X85-Y09)	72	2.8	4.6
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	70	2.7	4.4
7	Septicemia (A40-A41)	64	2.5	4.1
8	Influenza and pneumonia (J10-J18)	61	2.3	3.9
9	Chronic respiratory disease originating in the perinatal period (P27)	54	2.1	3.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	25	1.0	1.6
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	20	0.8	1.3
12	Malignant neoplasms (C00-C97)	18	0.7	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	0.7	*
14	Meningitis (G00,G03)	13	0.5	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	11	0.4	*
15	Necrotizing enterocolitis of newborn (P77)	11	0.4	*
...	All other causes (Residual)	697	26.8	44.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, both sexes				
...	All causes	2,955	100.0	466.7
1	Sudden infant death syndrome (R95)	611	20.7	96.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	367	12.4	58.0
3	Accidents (unintentional injuries) (V01-X59)	277	9.4	43.8
4	Septicemia (A40-A41)	131	4.4	20.7
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	129	4.4	20.4
6	Diseases of the circulatory system (I00-I99)	126	4.3	19.9
7	Chronic respiratory disease originating in the perinatal period (P27)	111	3.8	17.5
8	Influenza and pneumonia (J10-J18)	98	3.3	15.5
9	Assault (homicide) (*U01,X85-Y09)	88	3.0	13.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	58	2.0	9.2
11	Necrotizing enterocolitis of newborn (P77)	33	1.1	5.2
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	29	1.0	4.6
13	Bacterial sepsis of newborn (P36)	19	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	17	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	17	0.6	*
...	All other causes (Residual)	844	28.6	133.3
Postneonates, black, male				
...	All causes	1,666	100.0	518.6
1	Sudden infant death syndrome (R95)	341	20.5	106.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	10.8	56.0
3	Accidents (unintentional injuries) (V01-X59)	156	9.4	48.6
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	84	5.0	26.1
5	Diseases of the circulatory system (I00-I99)	73	4.4	22.7
6	Septicemia (A40-A41)	69	4.1	21.5
7	Chronic respiratory disease originating in the perinatal period (P27)	65	3.9	20.2
8	Assault (homicide) (*U01,X85-Y09)	51	3.1	15.9
9	Influenza and pneumonia (J10-J18)	47	2.8	14.6
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	33	2.0	10.3
11	Necrotizing enterocolitis of newborn (P77)	19	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	17	1.0	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.9	*
14	Bacterial sepsis of newborn (P36)	14	0.8	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.5	*
...	All other causes (Residual)	493	29.6	153.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, female				
...	All causes	1,289	100.0	413.3
1	Sudden infant death syndrome (R95)	270	20.9	86.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	187	14.5	60.0
3	Accidents (unintentional injuries) (V01-X59)	121	9.4	38.8
4	Septicemia (A40-A41)	62	4.8	19.9
5	Diseases of the circulatory system (I00-I99)	53	4.1	17.0
6	Influenza and pneumonia (J10-J18)	51	4.0	16.4
7	Chronic respiratory disease originating in the perinatal period (P27)	46	3.6	14.7
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	45	3.5	14.4
9	Assault (homicide) (*U01,X85-Y09)	37	2.9	11.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	25	1.9	8.0
11	Necrotizing enterocolitis of newborn (P77)	14	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	12	0.9	*
13	Respiratory distress of newborn (P22)	10	0.8	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.6	*
15	Meningitis (G00,G03)	7	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	7	0.5	*
...	All other causes (Residual)	334	25.9	107.1
Postneonates, American Indian, both sexes				
...	All causes	171	100.0	381.6
1	Sudden infant death syndrome (R95)	36	21.1	80.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	16.4	62.5
3	Accidents (unintentional injuries) (V01-X59)	19	11.1	*
4	Diseases of the circulatory system (I00-I99)	7	4.1	*
5	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	6	3.5	*
6	Influenza and pneumonia (J10-J18)	5	2.9	*
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	2.9	*
6	Assault (homicide) (*U01,X85-Y09)	5	2.9	*
9	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	3	1.8	*
10	Whooping cough (A37)	2	1.2	*
10	Septicemia (A40-A41)	2	1.2	*
10	Meningitis (G00,G03)	2	1.2	*
13	Meningococcal infection (A39)	1	0.6	*
13	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	0.6	*
13	Infantile cerebral palsy (G80)	1	0.6	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.6	*
13	Diseases of the ear and mastoid process (H60-H93)	1	0.6	*
13	Acute upper respiratory infections (J00-J06)	1	0.6	*
13	Bronchitis, chronic and unspecified (J40-J42)	1	0.6	*
13	Pneumonitis due to solids and liquids (J69)	1	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	1	0.6	*
13	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.6	*
...	All other causes (Residual)	41	24.0	91.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, American Indian, male				
...	All causes	89	100.0	392.5
1	Sudden infant death syndrome (R95)	20	22.5	88.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	16.9	*
3	Accidents (unintentional injuries) (V01-X59)	10	11.2	*
4	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	5	5.6	*
5	Assault (homicide) (*U01,X85-Y09)	4	4.5	*
6	Whooping cough (A37)	2	2.2	*
6	Meningitis (G00,G03)	2	2.2	*
6	Diseases of the circulatory system (I00-I99)	2	2.2	*
6	Influenza and pneumonia (J10-J18)	2	2.2	*
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	2	2.2	*
11	Meningococcal infection (A39)	1	1.1	*
11	Infantile cerebral palsy (G80)	1	1.1	*
11	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	1.1	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	1	1.1	*
...	All other causes (Residual)	21	23.6	92.6
Postneonates, American Indian, female				
...	All causes	82	100.0	370.4
1	Sudden infant death syndrome (R95)	16	19.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	15.9	*
3	Accidents (unintentional injuries) (V01-X59)	9	11.0	*
4	Diseases of the circulatory system (I00-I99)	5	6.1	*
5	Influenza and pneumonia (J10-J18)	3	3.7	*
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	3	3.7	*
7	Septicemia (A40-A41)	2	2.4	*
7	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	2	2.4	*
9	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	1.2	*
9	Anoxic brain damage, not elsewhere classified (G93.1)	1	1.2	*
9	Diseases of the ear and mastoid process (H60-H93)	1	1.2	*
9	Acute upper respiratory infections (J00-J06)	1	1.2	*
9	Bronchitis, chronic and unspecified (J40-J42)	1	1.2	*
9	Pneumonitis due to solids and liquids (J69)	1	1.2	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.2	*
9	Necrotizing enterocolitis of newborn (P77)	1	1.2	*
9	Assault (homicide) (*U01,X85-Y09)	1	1.2	*
...	All other causes (Residual)	20	24.4	90.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Asian or Pacific Islander, both sexes				
...	All causes	269	100.0	116.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	27.5	32.0
2	Sudden infant death syndrome (R95)	36	13.4	15.6
3	Diseases of the circulatory system (I00-I99)	17	6.3	*
4	Accidents (unintentional injuries) (V01-X59)	15	5.6	*
5	Septicemia (A40-A41)	9	3.3	*
6	Chronic respiratory disease originating in the perinatal period (P27)	7	2.6	*
6	Assault (homicide) (*U01,X85-Y09)	7	2.6	*
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	6	2.2	*
9	Influenza and pneumonia (J10-J18)	4	1.5	*
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	1.1	*
10	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	3	1.1	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	3	1.1	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
14	Meningitis (G00,G03)	2	0.7	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	0.7	*
14	Necrotizing enterocolitis of newborn (P77)	2	0.7	*
...	All other causes (Residual)	74	27.5	32.0
Postneonates, Asian or Pacific Islander, male				
...	All causes	144	100.0	120.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	27.8	33.5
2	Sudden infant death syndrome (R95)	24	16.7	20.1
3	Diseases of the circulatory system (I00-I99)	10	6.9	*
4	Accidents (unintentional injuries) (V01-X59)	8	5.6	*
5	Septicemia (A40-A41)	6	4.2	*
6	Influenza and pneumonia (J10-J18)	3	2.1	*
6	Acute bronchitis and acute bronchiolitis (J20-J21)	3	2.1	*
6	Assault (homicide) (*U01,X85-Y09)	3	2.1	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	2	1.4	*
9	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	2	1.4	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.4	*
9	Chronic respiratory disease originating in the perinatal period (P27)	2	1.4	*
13	Meningococcal infection (A39)	1	0.7	*
13	Malignant neoplasms (C00-C97)	1	0.7	*
13	Meningitis (G00,G03)	1	0.7	*
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	0.7	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.7	*
13	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	1	0.7	*
13	Slow fetal growth and fetal malnutrition (P05)	1	0.7	*
13	Necrotizing enterocolitis of newborn (P77)	1	0.7	*
...	All other causes (Residual)	31	21.5	26.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Asian or Pacific Islander, female				
...	All causes	125	100.0	111.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	27.2	30.4
2	Sudden infant death syndrome (R95)	12	9.6	*
3	Diseases of the circulatory system (I00-I99)	7	5.6	*
3	Accidents (unintentional injuries) (V01-X59)	7	5.6	*
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	4.0	*
5	Chronic respiratory disease originating in the perinatal period (P27)	5	4.0	*
7	Assault (homicide) (*U01,X85-Y09)	4	3.2	*
8	Septicemia (A40-A41)	3	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.6	*
9	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	1.6	*
11	Candidiasis (B37)	1	0.8	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	0.8	*
11	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	0.8	*
11	Meningitis (G00,G03)	1	0.8	*
11	Infantile spinal muscular atrophy, type I (Werdnig-Hoffman) (G12.0)	1	0.8	*
11	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.8	*
11	Influenza and pneumonia (J10-J18)	1	0.8	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	0.8	*
11	Necrotizing enterocolitis of newborn (P77)	1	0.8	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.8	*
...	All other causes (Residual)	34	27.2	30.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable