

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,598	100.0	563.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,182	25.7	144.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	633	13.8	77.6
3	Sudden infant death syndrome (R95)	302	6.6	37.0
4	Newborn affected by maternal complications of pregnancy (P01)	178	3.9	21.8
5	Respiratory distress of newborn (P22)	166	3.6	20.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	146	3.2	17.9
7	Bacterial sepsis of newborn (P36)	134	2.9	16.4
8	Accidents (unintentional injuries) (V01-X59)	118	2.6	14.5
9	Diseases of the circulatory system (I00-I99)	112	2.4	13.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	99	2.2	12.1
11	Neonatal hemorrhage (P50-P52,P54)	98	2.1	12.0
12	Atelectasis (P28.0-P28.1)	95	2.1	11.6
13	Septicemia (A40-A41)	62	1.3	7.6
14	Influenza and pneumonia (J10-J18)	60	1.3	7.4
15	Necrotizing enterocolitis of newborn (P77)	57	1.2	7.0
...	All other causes (Residual)	1,156	25.1	141.7
Infants, Hispanic, male				
...	All causes	2,514	100.0	603.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	605	24.1	145.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	337	13.4	80.9
3	Sudden infant death syndrome (R95)	183	7.3	43.9
4	Newborn affected by maternal complications of pregnancy (P01)	95	3.8	22.8
5	Respiratory distress of newborn (P22)	91	3.6	21.8
6	Bacterial sepsis of newborn (P36)	83	3.3	19.9
7	Newborn affected by complications of placenta, cord and membranes (P02)	73	2.9	17.5
8	Accidents (unintentional injuries) (V01-X59)	67	2.7	16.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	61	2.4	14.6
10	Neonatal hemorrhage (P50-P52,P54)	60	2.4	14.4
11	Diseases of the circulatory system (I00-I99)	57	2.3	13.7
12	Atelectasis (P28.0-P28.1)	52	2.1	12.5
13	Septicemia (A40-A41)	33	1.3	7.9
14	Necrotizing enterocolitis of newborn (P77)	31	1.2	7.4
15	Influenza and pneumonia (J10-J18)	30	1.2	7.2
...	All other causes (Residual)	656	26.1	157.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,084	100.0	521.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	577	27.7	144.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	296	14.2	74.1
3	Sudden infant death syndrome (R95)	119	5.7	29.8
4	Newborn affected by maternal complications of pregnancy (P01)	83	4.0	20.8
5	Respiratory distress of newborn (P22)	75	3.6	18.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	73	3.5	18.3
7	Diseases of the circulatory system (I00-I99)	55	2.6	13.8
8	Bacterial sepsis of newborn (P36)	51	2.4	12.8
8	Accidents (unintentional injuries) (V01-X59)	51	2.4	12.8
10	Atelectasis (P28.0-P28.1)	43	2.1	10.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	1.8	9.5
11	Neonatal hemorrhage (P50-P52,P54)	38	1.8	9.5
13	Influenza and pneumonia (J10-J18)	30	1.4	7.5
14	Septicemia (A40-A41)	29	1.4	7.3
15	Assault (homicide) (X85-Y09)	28	1.3	7.0
...	All other causes (Residual)	498	23.9	124.7
Infants, white, non-Hispanic, both sexes				
...	All causes	13,513	100.0	571.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,199	23.7	135.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,708	12.6	72.3
3	Sudden infant death syndrome (R95)	1,327	9.8	56.2
4	Newborn affected by maternal complications of pregnancy (P01)	663	4.9	28.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	573	4.2	24.2
6	Accidents (unintentional injuries) (V01-X59)	455	3.4	19.3
7	Respiratory distress of newborn (P22)	451	3.3	19.1
8	Diseases of the circulatory system (I00-I99)	343	2.5	14.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	341	2.5	14.4
10	Bacterial sepsis of newborn (P36)	337	2.5	14.3
11	Neonatal hemorrhage (P50-P52,P54)	289	2.1	12.2
12	Atelectasis (P28.0-P28.1)	252	1.9	10.7
13	Assault (homicide) (X85-Y09)	141	1.0	6.0
14	Necrotizing enterocolitis of newborn (P77)	138	1.0	5.8
15	Chronic respiratory disease originating in the perinatal period (P27)	135	1.0	5.7
...	All other causes (Residual)	3,161	23.4	133.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,660	100.0	632.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,720	22.5	141.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	925	12.1	76.3
3	Sudden infant death syndrome (R95)	798	10.4	65.9
4	Newborn affected by maternal complications of pregnancy (P01)	387	5.1	31.9
5	Respiratory distress of newborn (P22)	322	4.2	26.6
6				
	Newborn affected by complications of placenta, cord and membranes (P02)	308	4.0	25.4
7	Accidents (unintentional injuries) (V01-X59)	266	3.5	22.0
8	Bacterial sepsis of newborn (P36)	200	2.6	16.5
9	Diseases of the circulatory system (I00-I99)	189	2.5	15.6
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	180	2.3	14.9
11	Neonatal hemorrhage (P50-P52,P54)	171	2.2	14.1
12	Atelectasis (P28.0-P28.1)	156	2.0	12.9
13	Assault (homicide) (X85-Y09)	96	1.3	7.9
14	Necrotizing enterocolitis of newborn (P77)	86	1.1	7.1
15	Chronic respiratory disease originating in the perinatal period (P27)	72	0.9	5.9
...	All other causes (Residual)	1,784	23.3	147.2
Infants, white, non-Hispanic, female				
...	All causes	5,853	100.0	508.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,479	25.3	128.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	783	13.4	68.0
3	Sudden infant death syndrome (R95)	529	9.0	46.0
4	Newborn affected by maternal complications of pregnancy (P01)	276	4.7	24.0
5				
	Newborn affected by complications of placenta, cord and membranes (P02)	265	4.5	23.0
6	Accidents (unintentional injuries) (V01-X59)	189	3.2	16.4
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	161	2.8	14.0
8	Diseases of the circulatory system (I00-I99)	154	2.6	13.4
9	Bacterial sepsis of newborn (P36)	137	2.3	11.9
10	Respiratory distress of newborn (P22)	129	2.2	11.2
11	Neonatal hemorrhage (P50-P52,P54)	118	2.0	10.3
12	Atelectasis (P28.0-P28.1)	96	1.6	8.3
13	Chronic respiratory disease originating in the perinatal period (P27)	63	1.1	5.5
14	Hydrops fetalis not due to hemolytic disease (P83.2)	61	1.0	5.3
15	Pulmonary hemorrhage originating in the perinatal period (P26)	57	1.0	5.0
...	All other causes (Residual)	1,356	23.2	117.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,525	100.0	1,410.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,817	21.3	300.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,062	12.5	175.7
3	Sudden infant death syndrome (R95)	809	9.5	133.9
4	Newborn affected by maternal complications of pregnancy (P01)	506	5.9	83.7
5	Respiratory distress of newborn (P22)	341	4.0	56.4
6				
	Newborn affected by complications of placenta, cord and membranes (P02)	295	3.5	48.8
7	Accidents (unintentional injuries) (V01-X59)	268	3.1	44.3
8	Bacterial sepsis of newborn (P36)	264	3.1	43.7
9	Diseases of the circulatory system (I00-I99)	171	2.0	28.3
10	Neonatal hemorrhage (P50-P52,P54)	159	1.9	26.3
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	156	1.8	25.8
12	Necrotizing enterocolitis of newborn (P77)	154	1.8	25.5
13	Assault (homicide) (X85-Y09)	139	1.6	23.0
14	Atelectasis (P28.0-P28.1)	131	1.5	21.7
15	Chronic respiratory disease originating in the perinatal period (P27)	117	1.4	19.4
...	All other causes (Residual)	2,136	25.1	353.4
Infants, black, non-Hispanic, male				
...	All causes	4,757	100.0	1,550.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	997	21.0	324.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	545	11.5	177.6
3	Sudden infant death syndrome (R95)	473	9.9	154.2
4	Newborn affected by maternal complications of pregnancy (P01)	277	5.8	90.3
5	Respiratory distress of newborn (P22)	205	4.3	66.8
6				
	Newborn affected by complications of placenta, cord and membranes (P02)	155	3.3	50.5
7	Bacterial sepsis of newborn (P36)	152	3.2	49.5
8	Accidents (unintentional injuries) (V01-X59)	145	3.0	47.3
9	Neonatal hemorrhage (P50-P52,P54)	98	2.1	31.9
10	Necrotizing enterocolitis of newborn (P77)	95	2.0	31.0
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	91	1.9	29.7
12	Diseases of the circulatory system (I00-I99)	90	1.9	29.3
13	Atelectasis (P28.0-P28.1)	82	1.7	26.7
14	Assault (homicide) (X85-Y09)	73	1.5	23.8
15	Chronic respiratory disease originating in the perinatal period (P27)	68	1.4	22.2
...	All other causes (Residual)	1,211	25.5	394.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,768	100.0	1,266.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	820	21.8	275.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	517	13.7	173.8
3	Sudden infant death syndrome (R95)	336	8.9	112.9
4	Newborn affected by maternal complications of pregnancy (P01)	229	6.1	77.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	140	3.7	47.1
6	Respiratory distress of newborn (P22)	136	3.6	45.7
7	Accidents (unintentional injuries) (V01-X59)	123	3.3	41.3
8	Bacterial sepsis of newborn (P36)	112	3.0	37.6
9	Diseases of the circulatory system (I00-I99)	81	2.1	27.2
10	Assault (homicide) (X85-Y09)	66	1.8	22.2
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	65	1.7	21.8
12	Neonatal hemorrhage (P50-P52,P54)	61	1.6	20.5
13	Necrotizing enterocolitis of newborn (P77)	59	1.6	19.8
14	Chronic respiratory disease originating in the perinatal period (P27)	49	1.3	16.5
14	Atelectasis (P28.0-P28.1)	49	1.3	16.5
...	All other causes (Residual)	925	24.5	310.9
Neonates, Hispanic, both sexes				
...	All causes	3,048	100.0	373.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	826	27.1	101.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	610	20.0	74.8
3	Newborn affected by maternal complications of pregnancy (P01)	177	5.8	21.7
4	Respiratory distress of newborn (P22)	159	5.2	19.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	144	4.7	17.6
6	Bacterial sepsis of newborn (P36)	125	4.1	15.3
7	Neonatal hemorrhage (P50-P52,P54)	96	3.1	11.8
8	Atelectasis (P28.0-P28.1)	93	3.1	11.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	92	3.0	11.3
10	Necrotizing enterocolitis of newborn (P77)	51	1.7	6.3
11	Pulmonary hemorrhage originating in the perinatal period (P26)	45	1.5	5.5
12	Diseases of the circulatory system (I00-I99)	40	1.3	4.9
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	34	1.1	4.2
14	Sudden infant death syndrome (R95)	19	0.6	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	18	0.6	*
...	All other causes (Residual)	519	17.0	63.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,672	100.0	401.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	433	25.9	104.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	328	19.6	78.7
3	Newborn affected by maternal complications of pregnancy (P01)	94	5.6	22.6
4	Respiratory distress of newborn (P22)	87	5.2	20.9
5	Bacterial sepsis of newborn (P36)	79	4.7	19.0
6				
	Newborn affected by complications of placenta, cord and membranes (P02)	73	4.4	17.5
7	Neonatal hemorrhage (P50-P52,P54)	59	3.5	14.2
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	57	3.4	13.7
9	Atelectasis (P28.0-P28.1)	50	3.0	12.0
10	Pulmonary hemorrhage originating in the perinatal period (P26)	28	1.7	6.7
11	Necrotizing enterocolitis of newborn (P77)	27	1.6	6.5
12	Diseases of the circulatory system (I00-I99)	23	1.4	5.5
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	22	1.3	5.3
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	12	0.7	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	11	0.7	*
...	All other causes (Residual)	289	17.3	69.4
Neonates, Hispanic, female				
...	All causes	1,376	100.0	344.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	393	28.6	98.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	282	20.5	70.6
3	Newborn affected by maternal complications of pregnancy (P01)	83	6.0	20.8
4	Respiratory distress of newborn (P22)	72	5.2	18.0
5				
	Newborn affected by complications of placenta, cord and membranes (P02)	71	5.2	17.8
6	Bacterial sepsis of newborn (P36)	46	3.3	11.5
7	Atelectasis (P28.0-P28.1)	43	3.1	10.8
8	Neonatal hemorrhage (P50-P52,P54)	37	2.7	9.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	35	2.5	8.8
10	Necrotizing enterocolitis of newborn (P77)	24	1.7	6.0
11	Diseases of the circulatory system (I00-I99)	17	1.2	*
11	Pulmonary hemorrhage originating in the perinatal period (P26)	17	1.2	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	12	0.9	*
14	Sudden infant death syndrome (R95)	10	0.7	*
15	Newborn affected by other complications of labor and delivery (P03)	7	0.5	*
15	Neonatal aspiration syndromes (P24)	7	0.5	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	7	0.5	*
15	Assault (homicide) (X85-Y09)	7	0.5	*
...	All other causes (Residual)	206	15.0	51.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	9,078	100.0	384.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,399	26.4	101.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,690	18.6	71.5
3	Newborn affected by maternal complications of pregnancy (P01)	658	7.2	27.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	567	6.2	24.0
5	Respiratory distress of newborn (P22)	411	4.5	17.4
6	Bacterial sepsis of newborn (P36)	327	3.6	13.8
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	322	3.5	13.6
8	Neonatal hemorrhage (P50-P52,P54)	283	3.1	12.0
9	Atelectasis (P28.0-P28.1)	244	2.7	10.3
10	Diseases of the circulatory system (I00-I99)	138	1.5	5.8
11	Pulmonary hemorrhage originating in the perinatal period (P26)	123	1.4	5.2
12	Necrotizing enterocolitis of newborn (P77)	122	1.3	5.2
13	Sudden infant death syndrome (R95)	113	1.2	4.8
14	Hydrops fetalis not due to hemolytic disease (P83.2)	110	1.2	4.7
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	105	1.2	4.4
...	All other causes (Residual)	1,466	16.1	62.0
Neonates, white, non-Hispanic, male				
...	All causes	5,106	100.0	421.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,296	25.4	107.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	914	17.9	75.4
3	Newborn affected by maternal complications of pregnancy (P01)	384	7.5	31.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	304	6.0	25.1
5	Respiratory distress of newborn (P22)	294	5.8	24.3
6	Bacterial sepsis of newborn (P36)	195	3.8	16.1
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	171	3.3	14.1
8	Neonatal hemorrhage (P50-P52,P54)	168	3.3	13.9
9	Atelectasis (P28.0-P28.1)	152	3.0	12.5
10	Diseases of the circulatory system (I00-I99)	82	1.6	6.8
11	Necrotizing enterocolitis of newborn (P77)	77	1.5	6.4
12	Pulmonary hemorrhage originating in the perinatal period (P26)	67	1.3	5.5
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	60	1.2	5.0
14	Sudden infant death syndrome (R95)	56	1.1	4.6
15	Hydrops fetalis not due to hemolytic disease (P83.2)	53	1.0	4.4
...	All other causes (Residual)	833	16.3	68.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,972	100.0	345.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,103	27.8	95.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	776	19.5	67.4
3	Newborn affected by maternal complications of pregnancy (P01)	274	6.9	23.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	263	6.6	22.8
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	151	3.8	13.1
6	Bacterial sepsis of newborn (P36)	132	3.3	11.5
7	Respiratory distress of newborn (P22)	117	2.9	10.2
8	Neonatal hemorrhage (P50-P52,P54)	115	2.9	10.0
9	Atelectasis (P28.0-P28.1)	92	2.3	8.0
10	Hydrops fetalis not due to hemolytic disease (P83.2)	57	1.4	5.0
10	Sudden infant death syndrome (R95)	57	1.4	5.0
12	Diseases of the circulatory system (I00-I99)	56	1.4	4.9
12	Pulmonary hemorrhage originating in the perinatal period (P26)	56	1.4	4.9
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	45	1.1	3.9
14	Necrotizing enterocolitis of newborn (P77)	45	1.1	3.9
...	All other causes (Residual)	633	15.9	55.0
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,659	100.0	936.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,784	31.5	295.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	697	12.3	115.3
3	Newborn affected by maternal complications of pregnancy (P01)	502	8.9	83.1
4	Respiratory distress of newborn (P22)	320	5.7	52.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	291	5.1	48.2
6	Bacterial sepsis of newborn (P36)	252	4.5	41.7
7	Neonatal hemorrhage (P50-P52,P54)	154	2.7	25.5
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	143	2.5	23.7
9	Atelectasis (P28.0-P28.1)	128	2.3	21.2
10	Necrotizing enterocolitis of newborn (P77)	125	2.2	20.7
11	Pulmonary hemorrhage originating in the perinatal period (P26)	97	1.7	16.1
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	71	1.3	11.7
13	Sudden infant death syndrome (R95)	66	1.2	10.9
14	Diseases of the circulatory system (I00-I99)	51	0.9	8.4
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	41	0.7	6.8
...	All other causes (Residual)	937	16.6	155.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,177	100.0	1,035.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	984	31.0	320.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	11.8	122.5
3	Newborn affected by maternal complications of pregnancy (P01)	276	8.7	90.0
4	Respiratory distress of newborn (P22)	190	6.0	61.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	155	4.9	50.5
6	Bacterial sepsis of newborn (P36)	146	4.6	47.6
7	Neonatal hemorrhage (P50-P52,P54)	96	3.0	31.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	84	2.6	27.4
9	Atelectasis (P28.0-P28.1)	80	2.5	26.1
10	Necrotizing enterocolitis of newborn (P77)	76	2.4	24.8
11	Pulmonary hemorrhage originating in the perinatal period (P26)	58	1.8	18.9
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	38	1.2	12.4
12	Sudden infant death syndrome (R95)	38	1.2	12.4
14	Diseases of the circulatory system (I00-I99)	25	0.8	8.1
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	25	0.8	8.1
...	All other causes (Residual)	530	16.7	172.7
Neonates, black, non-Hispanic, female				
...	All causes	2,482	100.0	834.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	800	32.2	268.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	321	12.9	107.9
3	Newborn affected by maternal complications of pregnancy (P01)	226	9.1	76.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	136	5.5	45.7
5	Respiratory distress of newborn (P22)	130	5.2	43.7
6	Bacterial sepsis of newborn (P36)	106	4.3	35.6
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	59	2.4	19.8
8	Neonatal hemorrhage (P50-P52,P54)	58	2.3	19.5
9	Necrotizing enterocolitis of newborn (P77)	49	2.0	16.5
10	Atelectasis (P28.0-P28.1)	48	1.9	16.1
11	Pulmonary hemorrhage originating in the perinatal period (P26)	39	1.6	13.1
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	33	1.3	11.1
13	Sudden infant death syndrome (R95)	28	1.1	9.4
14	Diseases of the circulatory system (I00-I99)	26	1.0	8.7
15	Hematological disorders (P60-P61)	17	0.7	*
15	Accidents (unintentional injuries) (V01-X59)	17	0.7	*
...	All other causes (Residual)	389	15.7	130.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,550	100.0	190.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	356	23.0	43.6
2	Sudden infant death syndrome (R95)	283	18.3	34.7
3	Accidents (unintentional injuries) (V01-X59)	107	6.9	13.1
4	Diseases of the circulatory system (I00-I99)	72	4.6	8.8
5	Septicemia (A40-A41)	62	4.0	7.6
6	Influenza and pneumonia (J10-J18)	53	3.4	6.5
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	45	2.9	5.5
7	Assault (homicide) (X85-Y09)	45	2.9	5.5
9	Chronic respiratory disease originating in the perinatal period (P27)	28	1.8	3.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	1.5	2.8
11	Malignant neoplasms (C00-C97)	13	0.8	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	11	0.7	*
13	Meningococcal infection (A39)	9	0.6	*
13	Meningitis (G00,G03)	9	0.6	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.6	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	9	0.6	*
13	Bacterial sepsis of newborn (P36)	9	0.6	*
...	All other causes (Residual)	407	26.3	49.9
Postneonates, Hispanic, male				
...	All causes	842	100.0	202.1
1	Sudden infant death syndrome (R95)	174	20.7	41.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	20.4	41.3
3	Accidents (unintentional injuries) (V01-X59)	61	7.2	14.6
4	Diseases of the circulatory system (I00-I99)	34	4.0	8.2
5	Septicemia (A40-A41)	33	3.9	7.9
6	Influenza and pneumonia (J10-J18)	27	3.2	6.5
7	Assault (homicide) (X85-Y09)	24	2.9	5.8
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	23	2.7	5.5
9	Chronic respiratory disease originating in the perinatal period (P27)	15	1.8	*
10	Malignant neoplasms (C00-C97)	9	1.1	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.1	*
12	Meningitis (G00,G03)	6	0.7	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	6	0.7	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	5	0.6	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	5	0.6	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	5	0.6	*
...	All other causes (Residual)	234	27.8	56.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	708	100.0	177.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	26.0	46.1
2	Sudden infant death syndrome (R95)	109	15.4	27.3
3	Accidents (unintentional injuries) (V01-X59)	46	6.5	11.5
4	Diseases of the circulatory system (I00-I99)	38	5.4	9.5
5	Septicemia (A40-A41)	29	4.1	7.3
6	Influenza and pneumonia (J10-J18)	26	3.7	6.5
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	22	3.1	5.5
8	Assault (homicide) (X85-Y09)	21	3.0	5.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	2.0	*
10	Chronic respiratory disease originating in the perinatal period (P27)	13	1.8	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	6	0.8	*
11	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	6	0.8	*
13	Meningococcal infection (A39)	5	0.7	*
13	Bacterial sepsis of newborn (P36)	5	0.7	*
15	Malignant neoplasms (C00-C97)	4	0.6	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	4	0.6	*
...	All other causes (Residual)	176	24.9	44.1
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,435	100.0	187.7
1	Sudden infant death syndrome (R95)	1,214	27.4	51.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	800	18.0	33.9
3	Accidents (unintentional injuries) (V01-X59)	405	9.1	17.1
4	Diseases of the circulatory system (I00-I99)	205	4.6	8.7
5	Assault (homicide) (X85-Y09)	129	2.9	5.5
6	Septicemia (A40-A41)	108	2.4	4.6
7	Chronic respiratory disease originating in the perinatal period (P27)	106	2.4	4.5
8	Influenza and pneumonia (J10-J18)	103	2.3	4.4
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	65	1.5	2.8
10	Malignant neoplasms (C00-C97)	57	1.3	2.4
11	Respiratory distress of newborn (P22)	40	0.9	1.7
12	Meningitis (G00,G03)	37	0.8	1.6
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	29	0.7	1.2
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	23	0.5	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	0.9
15	Infantile spinal muscular atrophy, type I (Werdnig-Hoffman) (G12.0)	21	0.5	0.9
...	All other causes (Residual)	1,072	24.2	45.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, male				
...	All causes	2,554	100.0	210.8
1	Sudden infant death syndrome (R95)	742	29.1	61.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	424	16.6	35.0
3	Accidents (unintentional injuries) (V01-X59)	237	9.3	19.6
4	Diseases of the circulatory system (I00-I99)	107	4.2	8.8
5	Assault (homicide) (X85-Y09)	88	3.4	7.3
6	Influenza and pneumonia (J10-J18)	63	2.5	5.2
7	Chronic respiratory disease originating in the perinatal period (P27)	58	2.3	4.8
8	Septicemia (A40-A41)	55	2.2	4.5
9	Malignant neoplasms (C00-C97)	34	1.3	2.8
10	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	32	1.3	2.6
11	Respiratory distress of newborn (P22)	28	1.1	2.3
12	Meningitis (G00,G03)	20	0.8	1.7
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	15	0.6	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	15	0.6	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	13	0.5	*
...	All other causes (Residual)	623	24.4	51.4
Postneonates, white, non-Hispanic, female				
...	All causes	1,881	100.0	163.4
1	Sudden infant death syndrome (R95)	472	25.1	41.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	20.0	32.7
3	Accidents (unintentional injuries) (V01-X59)	168	8.9	14.6
4	Diseases of the circulatory system (I00-I99)	98	5.2	8.5
5	Septicemia (A40-A41)	53	2.8	4.6
6	Chronic respiratory disease originating in the perinatal period (P27)	48	2.6	4.2
7	Assault (homicide) (X85-Y09)	41	2.2	3.6
8	Influenza and pneumonia (J10-J18)	40	2.1	3.5
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	33	1.8	2.9
10	Malignant neoplasms (C00-C97)	23	1.2	2.0
11	Meningitis (G00,G03)	17	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	14	0.7	*
13	Respiratory distress of newborn (P22)	12	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.5	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	10	0.5	*
...	All other causes (Residual)	466	24.8	40.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,866	100.0	474.2
1	Sudden infant death syndrome (R95)	743	25.9	122.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	365	12.7	60.4
3	Accidents (unintentional injuries) (V01-X59)	238	8.3	39.4
4	Assault (homicide) (X85-Y09)	124	4.3	20.5
5	Diseases of the circulatory system (I00-I99)	120	4.2	19.9
6	Chronic respiratory disease originating in the perinatal period (P27)	104	3.6	17.2
7	Septicemia (A40-A41)	96	3.3	15.9
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	95	3.3	15.7
9	Influenza and pneumonia (J10-J18)	93	3.2	15.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	33	1.2	5.5
11	Meningitis (G00,G03)	32	1.1	5.3
12	Necrotizing enterocolitis of newborn (P77)	29	1.0	4.8
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	25	0.9	4.1
14	Respiratory distress of newborn (P22)	21	0.7	3.5
15	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.5	*
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	13	0.5	*
...	All other causes (Residual)	722	25.2	119.5
Postneonates, black, non-Hispanic, male				
...	All causes	1,580	100.0	514.9
1	Sudden infant death syndrome (R95)	435	27.5	141.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	10.7	55.1
3	Accidents (unintentional injuries) (V01-X59)	132	8.4	43.0
4	Assault (homicide) (X85-Y09)	69	4.4	22.5
5	Diseases of the circulatory system (I00-I99)	65	4.1	21.2
6	Chronic respiratory disease originating in the perinatal period (P27)	60	3.8	19.6
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	53	3.4	17.3
8	Influenza and pneumonia (J10-J18)	50	3.2	16.3
9	Septicemia (A40-A41)	48	3.0	15.6
10	Necrotizing enterocolitis of newborn (P77)	19	1.2	*
11	Meningitis (G00,G03)	15	0.9	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	15	0.9	*
11	Respiratory distress of newborn (P22)	15	0.9	*
14	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	13	0.8	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	10	0.6	*
...	All other causes (Residual)	412	26.1	134.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2000

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group.

Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
	Postneonates, black, non-Hispanic, female			
...	All causes	1,286	100.0	432.2
1	Sudden infant death syndrome (R95)	308	24.0	103.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	15.2	65.9
3	Accidents (unintentional injuries) (V01-X59)	106	8.2	35.6
4	Diseases of the circulatory system (I00-I99)	55	4.3	18.5
4	Assault (homicide) (X85-Y09)	55	4.3	18.5
6	Septicemia (A40-A41)	48	3.7	16.1
7	Chronic respiratory disease originating in the perinatal period (P27)	44	3.4	14.8
8	Influenza and pneumonia (J10-J18)	43	3.3	14.5
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	42	3.3	14.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	20	1.6	6.7
11	Meningitis (G00,G03)	17	1.3	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.8	*
12	Necrotizing enterocolitis of newborn (P77)	10	0.8	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	6	0.5	*
14	Respiratory distress of newborn (P22)	6	0.5	*
14	Bacterial sepsis of newborn (P36)	6	0.5	*
...	All other causes (Residual)	314	24.4	105.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable