

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,680	100.0	549.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,195	25.5	140.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	672	14.4	78.9
3	Sudden infant death syndrome (R95)	250	5.3	29.3
4	Newborn affected by maternal complications of pregnancy (P01)	185	4.0	21.7
5	Respiratory distress of newborn (P22)	181	3.9	21.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	171	3.7	20.1
7	Accidents (unintentional injuries) (V01-X59)	154	3.3	18.1
8	Bacterial sepsis of newborn (P36)	132	2.8	15.5
9	Diseases of the circulatory system (I00-I99)	103	2.2	12.1
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	93	2.0	10.9
11	Atelectasis (P28.0-P28.1)	88	1.9	10.3
12	Neonatal hemorrhage (P50-P52,P54)	80	1.7	9.4
13	Necrotizing enterocolitis of newborn (P77)	64	1.4	7.5
14	Influenza and pneumonia (J10-J18)	57	1.2	6.7
15	Assault (homicide) (*U01,X85-Y09)	55	1.2	6.5
...	All other causes (Residual)	1,200	25.6	140.9
Infants, Hispanic, male				
...	All causes	2,601	100.0	599.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	639	24.6	147.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	368	14.1	84.8
3	Sudden infant death syndrome (R95)	142	5.5	32.7
4	Newborn affected by maternal complications of pregnancy (P01)	109	4.2	25.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	109	4.2	25.1
6	Respiratory distress of newborn (P22)	98	3.8	22.6
7	Bacterial sepsis of newborn (P36)	78	3.0	18.0
8	Accidents (unintentional injuries) (V01-X59)	75	2.9	17.3
9	Diseases of the circulatory system (I00-I99)	57	2.2	13.1
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	56	2.2	12.9
11	Neonatal hemorrhage (P50-P52,P54)	47	1.8	10.8
12	Atelectasis (P28.0-P28.1)	46	1.8	10.6
13	Necrotizing enterocolitis of newborn (P77)	35	1.3	8.1
14	Assault (homicide) (*U01,X85-Y09)	33	1.3	7.6
15	Influenza and pneumonia (J10-J18)	31	1.2	7.1
...	All other causes (Residual)	678	26.1	156.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,079	100.0	497.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	556	26.7	133.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	304	14.6	72.7
3	Sudden infant death syndrome (R95)	108	5.2	25.8
4	Respiratory distress of newborn (P22)	83	4.0	19.9
5	Accidents (unintentional injuries) (V01-X59)	79	3.8	18.9
6	Newborn affected by maternal complications of pregnancy (P01)	76	3.7	18.2
7	Newborn affected by complications of placenta, cord and membranes (P02)	62	3.0	14.8
8	Bacterial sepsis of newborn (P36)	54	2.6	12.9
9	Diseases of the circulatory system (I00-I99)	46	2.2	11.0
10	Atelectasis (P28.0-P28.1)	42	2.0	10.0
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	37	1.8	8.9
12	Neonatal hemorrhage (P50-P52,P54)	33	1.6	7.9
13	Necrotizing enterocolitis of newborn (P77)	29	1.4	6.9
14	Influenza and pneumonia (J10-J18)	26	1.3	6.2
15	Assault (homicide) (*U01,X85-Y09)	22	1.1	5.3
...	All other causes (Residual)	522	25.1	124.9
Infants, white, non-Hispanic, both sexes				
...	All causes	13,311	100.0	572.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,053	22.9	131.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,773	13.3	76.2
3	Sudden infant death syndrome (R95)	1,173	8.8	50.4
4	Newborn affected by maternal complications of pregnancy (P01)	733	5.5	31.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	519	3.9	22.3
6	Accidents (unintentional injuries) (V01-X59)	500	3.8	21.5
7	Respiratory distress of newborn (P22)	456	3.4	19.6
8	Bacterial sepsis of newborn (P36)	311	2.3	13.4
9	Diseases of the circulatory system (I00-I99)	301	2.3	12.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	293	2.2	12.6
11	Atelectasis (P28.0-P28.1)	254	1.9	10.9
12	Neonatal hemorrhage (P50-P52,P54)	248	1.9	10.7
13	Necrotizing enterocolitis of newborn (P77)	156	1.2	6.7
14	Chronic respiratory disease originating in the perinatal period (P27)	141	1.1	6.1
15	Assault (homicide) (*U01,X85-Y09)	139	1.0	6.0
...	All other causes (Residual)	3,261	24.5	140.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,508	100.0	629.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,664	22.2	139.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	969	12.9	81.3
3	Sudden infant death syndrome (R95)	707	9.4	59.3
4	Newborn affected by maternal complications of pregnancy (P01)	419	5.6	35.1
5	Accidents (unintentional injuries) (V01-X59)	281	3.7	23.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	280	3.7	23.5
7	Respiratory distress of newborn (P22)	252	3.4	21.1
8	Bacterial sepsis of newborn (P36)	193	2.6	16.2
9	Diseases of the circulatory system (I00-I99)	164	2.2	13.8
10	Atelectasis (P28.0-P28.1)	157	2.1	13.2
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	155	2.1	13.0
12	Neonatal hemorrhage (P50-P52,P54)	148	2.0	12.4
13	Necrotizing enterocolitis of newborn (P77)	99	1.3	8.3
14	Chronic respiratory disease originating in the perinatal period (P27)	91	1.2	7.6
15	Assault (homicide) (*U01,X85-Y09)	84	1.1	7.0
...	All other causes (Residual)	1,845	24.6	154.8
Infants, white, non-Hispanic, female				
...	All causes	5,803	100.0	511.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,389	23.9	122.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	804	13.9	70.9
3	Sudden infant death syndrome (R95)	466	8.0	41.1
4	Newborn affected by maternal complications of pregnancy (P01)	314	5.4	27.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	239	4.1	21.1
6	Accidents (unintentional injuries) (V01-X59)	219	3.8	19.3
7	Respiratory distress of newborn (P22)	204	3.5	18.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	138	2.4	12.2
9	Diseases of the circulatory system (I00-I99)	137	2.4	12.1
10	Bacterial sepsis of newborn (P36)	118	2.0	10.4
11	Neonatal hemorrhage (P50-P52,P54)	100	1.7	8.8
12	Atelectasis (P28.0-P28.1)	97	1.7	8.6
13	Hydrops fetalis not due to hemolytic disease (P83.2)	59	1.0	5.2
14	Necrotizing enterocolitis of newborn (P77)	57	1.0	5.0
15	Assault (homicide) (*U01,X85-Y09)	55	0.9	4.8
...	All other causes (Residual)	1,407	24.2	124.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,277	100.0	1,403.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,781	21.5	301.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	999	12.1	169.3
3	Sudden infant death syndrome (R95)	731	8.8	123.9
4	Newborn affected by maternal complications of pregnancy (P01)	524	6.3	88.8
5	Respiratory distress of newborn (P22)	339	4.1	57.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	286	3.5	48.5
7	Accidents (unintentional injuries) (V01-X59)	272	3.3	46.1
8	Bacterial sepsis of newborn (P36)	226	2.7	38.3
9	Diseases of the circulatory system (I00-I99)	189	2.3	32.0
10	Atelectasis (P28.0-P28.1)	137	1.7	23.2
10	Necrotizing enterocolitis of newborn (P77)	137	1.7	23.2
12	Chronic respiratory disease originating in the perinatal period (P27)	128	1.5	21.7
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	125	1.5	21.2
14	Septicemia (A40-A41)	124	1.5	21.0
15	Assault (homicide) (*U01,X85-Y09)	119	1.4	20.2
...	All other causes (Residual)	2,160	26.1	366.2
Infants, black, non-Hispanic, male				
...	All causes	4,651	100.0	1,552.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	988	21.2	329.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	525	11.3	175.3
3	Sudden infant death syndrome (R95)	414	8.9	138.2
4	Newborn affected by maternal complications of pregnancy (P01)	290	6.2	96.8
5	Respiratory distress of newborn (P22)	203	4.4	67.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	163	3.5	54.4
7	Accidents (unintentional injuries) (V01-X59)	158	3.4	52.7
8	Bacterial sepsis of newborn (P36)	121	2.6	40.4
9	Diseases of the circulatory system (I00-I99)	96	2.1	32.0
10	Atelectasis (P28.0-P28.1)	85	1.8	28.4
11	Necrotizing enterocolitis of newborn (P77)	82	1.8	27.4
12	Chronic respiratory disease originating in the perinatal period (P27)	75	1.6	25.0
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	71	1.5	23.7
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	70	1.5	23.4
15	Assault (homicide) (*U01,X85-Y09)	68	1.5	22.7
...	All other causes (Residual)	1,242	26.7	414.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,626	100.0	1,248.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	793	21.9	273.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	474	13.1	163.3
3	Sudden infant death syndrome (R95)	317	8.7	109.2
4	Newborn affected by maternal complications of pregnancy (P01)	234	6.5	80.6
5	Respiratory distress of newborn (P22)	136	3.8	46.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	123	3.4	42.4
7	Accidents (unintentional injuries) (V01-X59)	114	3.1	39.3
8	Bacterial sepsis of newborn (P36)	105	2.9	36.2
9	Diseases of the circulatory system (I00-I99)	93	2.6	32.0
10	Septicemia (A40-A41)	57	1.6	19.6
11	Necrotizing enterocolitis of newborn (P77)	55	1.5	18.9
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	54	1.5	18.6
13	Chronic respiratory disease originating in the perinatal period (P27)	53	1.5	18.3
14	Atelectasis (P28.0-P28.1)	52	1.4	17.9
15	Influenza and pneumonia (J10-J18)	51	1.4	17.6
15	Assault (homicide) (*U01,X85-Y09)	51	1.4	17.6
...	All other causes (Residual)	864	23.8	297.6
Neonates, Hispanic, both sexes				
...	All causes	3,112	100.0	365.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	812	26.1	95.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	659	21.2	77.4
3	Newborn affected by maternal complications of pregnancy (P01)	185	5.9	21.7
4	Respiratory distress of newborn (P22)	169	5.4	19.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	167	5.4	19.6
6	Bacterial sepsis of newborn (P36)	124	4.0	14.6
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	85	2.7	10.0
8	Atelectasis (P28.0-P28.1)	84	2.7	9.9
9	Neonatal hemorrhage (P50-P52,P54)	80	2.6	9.4
10	Necrotizing enterocolitis of newborn (P77)	56	1.8	6.6
11	Pulmonary hemorrhage originating in the perinatal period (P26)	40	1.3	4.7
12	Diseases of the circulatory system (I00-I99)	33	1.1	3.9
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	29	0.9	3.4
14	Birth trauma (P10-P15)	26	0.8	3.1
14	Hematological disorders (P60-P61)	26	0.8	3.1
...	All other causes (Residual)	537	17.3	63.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,770	100.0	408.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	454	25.6	104.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	364	20.6	83.9
3	Newborn affected by maternal complications of pregnancy (P01)	109	6.2	25.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	108	6.1	24.9
5	Respiratory distress of newborn (P22)	89	5.0	20.5
6	Bacterial sepsis of newborn (P36)	74	4.2	17.1
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	54	3.1	12.4
8	Neonatal hemorrhage (P50-P52,P54)	47	2.7	10.8
9	Atelectasis (P28.0-P28.1)	44	2.5	10.1
10	Necrotizing enterocolitis of newborn (P77)	33	1.9	7.6
11	Pulmonary hemorrhage originating in the perinatal period (P26)	27	1.5	6.2
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	16	0.9	*
13	Diseases of the circulatory system (I00-I99)	15	0.8	*
13	Birth trauma (P10-P15)	15	0.8	*
15	Newborn affected by other complications of labor and delivery (P03)	14	0.8	*
...	All other causes (Residual)	307	17.3	70.8
Neonates, Hispanic, female				
...	All causes	1,342	100.0	321.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	358	26.7	85.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	295	22.0	70.6
3	Respiratory distress of newborn (P22)	80	6.0	19.1
4	Newborn affected by maternal complications of pregnancy (P01)	76	5.7	18.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	59	4.4	14.1
6	Bacterial sepsis of newborn (P36)	50	3.7	12.0
7	Atelectasis (P28.0-P28.1)	40	3.0	9.6
8	Neonatal hemorrhage (P50-P52,P54)	33	2.5	7.9
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	2.3	7.4
10	Necrotizing enterocolitis of newborn (P77)	23	1.7	5.5
11	Diseases of the circulatory system (I00-I99)	18	1.3	*
12	Hematological disorders (P60-P61)	14	1.0	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	13	1.0	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	13	1.0	*
15	Birth trauma (P10-P15)	11	0.8	*
...	All other causes (Residual)	228	17.0	54.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	8,888	100.0	382.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,224	25.0	95.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,739	19.6	74.7
3	Newborn affected by maternal complications of pregnancy (P01)	731	8.2	31.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	510	5.7	21.9
5	Respiratory distress of newborn (P22)	429	4.8	18.4
6	Bacterial sepsis of newborn (P36)	296	3.3	12.7
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	285	3.2	12.2
8	Atelectasis (P28.0-P28.1)	250	2.8	10.7
9	Neonatal hemorrhage (P50-P52,P54)	242	2.7	10.4
10	Necrotizing enterocolitis of newborn (P77)	133	1.5	5.7
11	Diseases of the circulatory system (I00-I99)	118	1.3	5.1
11	Sudden infant death syndrome (R95)	118	1.3	5.1
13	Hydrops fetalis not due to hemolytic disease (P83.2)	115	1.3	4.9
14	Pulmonary hemorrhage originating in the perinatal period (P26)	114	1.3	4.9
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	111	1.2	4.8
...	All other causes (Residual)	1,473	16.6	63.3
Neonates, white, non-Hispanic, male				
...	All causes	4,969	100.0	416.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,232	24.8	103.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	947	19.1	79.4
3	Newborn affected by maternal complications of pregnancy (P01)	417	8.4	35.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	274	5.5	23.0
5	Respiratory distress of newborn (P22)	244	4.9	20.5
6	Bacterial sepsis of newborn (P36)	186	3.7	15.6
7	Atelectasis (P28.0-P28.1)	155	3.1	13.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	151	3.0	12.7
9	Neonatal hemorrhage (P50-P52,P54)	145	2.9	12.2
10	Necrotizing enterocolitis of newborn (P77)	79	1.6	6.6
11	Diseases of the circulatory system (I00-I99)	68	1.4	5.7
12	Pulmonary hemorrhage originating in the perinatal period (P26)	65	1.3	5.5
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	63	1.3	5.3
14	Sudden infant death syndrome (R95)	60	1.2	5.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	59	1.2	4.9
...	All other causes (Residual)	824	16.6	69.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,919	100.0	345.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	992	25.3	87.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	792	20.2	69.8
3	Newborn affected by maternal complications of pregnancy (P01)	314	8.0	27.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	236	6.0	20.8
5	Respiratory distress of newborn (P22)	185	4.7	16.3
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	134	3.4	11.8
7	Bacterial sepsis of newborn (P36)	110	2.8	9.7
8	Neonatal hemorrhage (P50-P52,P54)	97	2.5	8.6
9	Atelectasis (P28.0-P28.1)	95	2.4	8.4
10	Sudden infant death syndrome (R95)	58	1.5	5.1
11	Hydrops fetalis not due to hemolytic disease (P83.2)	56	1.4	4.9
12	Necrotizing enterocolitis of newborn (P77)	54	1.4	4.8
13	Diseases of the circulatory system (I00-I99)	50	1.3	4.4
14	Pulmonary hemorrhage originating in the perinatal period (P26)	49	1.3	4.3
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	48	1.2	4.2
...	All other causes (Residual)	649	16.6	57.2
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,427	100.0	920.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,741	32.1	295.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	655	12.1	111.0
3	Newborn affected by maternal complications of pregnancy (P01)	519	9.6	88.0
4	Respiratory distress of newborn (P22)	312	5.7	52.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	280	5.2	47.5
6	Bacterial sepsis of newborn (P36)	207	3.8	35.1
7	Atelectasis (P28.0-P28.1)	136	2.5	23.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	119	2.2	20.2
9	Necrotizing enterocolitis of newborn (P77)	115	2.1	19.5
10	Pulmonary hemorrhage originating in the perinatal period (P26)	104	1.9	17.6
11	Neonatal hemorrhage (P50-P52,P54)	96	1.8	16.3
12	Diseases of the circulatory system (I00-I99)	54	1.0	9.2
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	54	1.0	9.2
14	Sudden infant death syndrome (R95)	45	0.8	7.6
15	Birth trauma (P10-P15)	41	0.8	7.0
...	All other causes (Residual)	949	17.5	160.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,045	100.0	1,016.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	965	31.7	322.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	11.5	116.5
3	Newborn affected by maternal complications of pregnancy (P01)	288	9.5	96.1
4	Respiratory distress of newborn (P22)	189	6.2	63.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	158	5.2	52.7
6	Bacterial sepsis of newborn (P36)	110	3.6	36.7
7	Atelectasis (P28.0-P28.1)	84	2.8	28.0
8	Necrotizing enterocolitis of newborn (P77)	73	2.4	24.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	68	2.2	22.7
10	Pulmonary hemorrhage originating in the perinatal period (P26)	64	2.1	21.4
11	Neonatal hemorrhage (P50-P52,P54)	48	1.6	16.0
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	29	1.0	9.7
13	Hematological disorders (P60-P61)	28	0.9	9.3
14	Birth trauma (P10-P15)	27	0.9	9.0
15	Diseases of the circulatory system (I00-I99)	26	0.9	8.7
...	All other causes (Residual)	539	17.7	179.9
Neonates, black, non-Hispanic, female				
...	All causes	2,382	100.0	820.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	776	32.6	267.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	12.8	105.4
3	Newborn affected by maternal complications of pregnancy (P01)	231	9.7	79.6
4	Respiratory distress of newborn (P22)	123	5.2	42.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	122	5.1	42.0
6	Bacterial sepsis of newborn (P36)	97	4.1	33.4
7	Atelectasis (P28.0-P28.1)	52	2.2	17.9
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	51	2.1	17.6
9	Neonatal hemorrhage (P50-P52,P54)	48	2.0	16.5
10	Necrotizing enterocolitis of newborn (P77)	42	1.8	14.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	40	1.7	13.8
12	Diseases of the circulatory system (I00-I99)	28	1.2	9.6
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	25	1.0	8.6
13	Sudden infant death syndrome (R95)	25	1.0	8.6
15	Congenital pneumonia (P23)	15	0.6	*
15	Neonatal aspiration syndromes (P24)	15	0.6	*
...	All other causes (Residual)	386	16.2	132.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,568	100.0	184.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	383	24.4	45.0
2	Sudden infant death syndrome (R95)	238	15.2	27.9
3	Accidents (unintentional injuries) (V01-X59)	138	8.8	16.2
4	Diseases of the circulatory system (I00-I99)	70	4.5	8.2
5	Influenza and pneumonia (J10-J18)	52	3.3	6.1
6	Septicemia (A40-A41)	50	3.2	5.9
7	Assault (homicide) (*U01,X85-Y09)	46	2.9	5.4
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	34	2.2	4.0
9	Chronic respiratory disease originating in the perinatal period (P27)	23	1.5	2.7
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	19	1.2	*
11	Malignant neoplasms (C00-C97)	14	0.9	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	13	0.8	*
13	Respiratory distress of newborn (P22)	12	0.8	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.6	*
15	Whooping cough (A37)	9	0.6	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	9	0.6	*
...	All other causes (Residual)	448	28.6	52.6
Postneonates, Hispanic, male				
...	All causes	831	100.0	191.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	22.3	42.6
2	Sudden infant death syndrome (R95)	137	16.5	31.6
3	Accidents (unintentional injuries) (V01-X59)	68	8.2	15.7
4	Diseases of the circulatory system (I00-I99)	42	5.1	9.7
5	Septicemia (A40-A41)	29	3.5	6.7
5	Assault (homicide) (*U01,X85-Y09)	29	3.5	6.7
7	Influenza and pneumonia (J10-J18)	26	3.1	6.0
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	19	2.3	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	11	1.3	*
10	Respiratory distress of newborn (P22)	9	1.1	*
10	Chronic respiratory disease originating in the perinatal period (P27)	9	1.1	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	6	0.7	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	6	0.7	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.7	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	5	0.6	*
...	All other causes (Residual)	244	29.4	56.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	737	100.0	176.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	26.9	47.4
2	Sudden infant death syndrome (R95)	101	13.7	24.2
3	Accidents (unintentional injuries) (V01-X59)	70	9.5	16.7
4	Diseases of the circulatory system (I00-I99)	28	3.8	6.7
5	Influenza and pneumonia (J10-J18)	26	3.5	6.2
6	Septicemia (A40-A41)	21	2.8	5.0
7	Assault (homicide) (*U01,X85-Y09)	17	2.3	*
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	15	2.0	*
9	Chronic respiratory disease originating in the perinatal period (P27)	14	1.9	*
10	Malignant neoplasms (C00-C97)	10	1.4	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.2	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	8	1.1	*
13	Whooping cough (A37)	6	0.8	*
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	6	0.8	*
13	Necrotizing enterocolitis of newborn (P77)	6	0.8	*
...	All other causes (Residual)	202	27.4	48.3
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,423	100.0	190.1
1	Sudden infant death syndrome (R95)	1,055	23.9	45.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	829	18.7	35.6
3	Accidents (unintentional injuries) (V01-X59)	447	10.1	19.2
4	Diseases of the circulatory system (I00-I99)	183	4.1	7.9
5	Septicemia (A40-A41)	122	2.8	5.2
6	Chronic respiratory disease originating in the perinatal period (P27)	118	2.7	5.1
6	Assault (homicide) (*U01,X85-Y09)	118	2.7	5.1
8	Influenza and pneumonia (J10-J18)	106	2.4	4.6
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	67	1.5	2.9
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	35	0.8	1.5
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	34	0.8	1.5
12	Malignant neoplasms (C00-C97)	33	0.7	1.4
13	Meningitis (G00,G03)	28	0.6	1.2
14	Respiratory distress of newborn (P22)	27	0.6	1.2
15	Acute bronchitis and acute bronchiolitis (J20-J21)	23	0.5	1.0
15	Necrotizing enterocolitis of newborn (P77)	23	0.5	1.0
...	All other causes (Residual)	1,175	26.6	50.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, male				
...	All causes	2,539	100.0	213.0
1	Sudden infant death syndrome (R95)	647	25.5	54.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	432	17.0	36.2
3	Accidents (unintentional injuries) (V01-X59)	253	10.0	21.2
4	Diseases of the circulatory system (I00-I99)	96	3.8	8.1
5	Chronic respiratory disease originating in the perinatal period (P27)	78	3.1	6.5
6	Assault (homicide) (*U01,X85-Y09)	76	3.0	6.4
7	Septicemia (A40-A41)	71	2.8	6.0
8	Influenza and pneumonia (J10-J18)	64	2.5	5.4
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	33	1.3	2.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	0.9	1.8
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	21	0.8	1.8
12	Necrotizing enterocolitis of newborn (P77)	20	0.8	1.7
13	Malignant neoplasms (C00-C97)	17	0.7	*
14	Meningitis (G00,G03)	13	0.5	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	13	0.5	*
...	All other causes (Residual)	683	26.9	57.3
Postneonates, white, non-Hispanic, female				
...	All causes	1,884	100.0	166.1
1	Sudden infant death syndrome (R95)	408	21.7	36.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	397	21.1	35.0
3	Accidents (unintentional injuries) (V01-X59)	194	10.3	17.1
4	Diseases of the circulatory system (I00-I99)	87	4.6	7.7
5	Septicemia (A40-A41)	51	2.7	4.5
6	Influenza and pneumonia (J10-J18)	42	2.2	3.7
6	Assault (homicide) (*U01,X85-Y09)	42	2.2	3.7
8	Chronic respiratory disease originating in the perinatal period (P27)	40	2.1	3.5
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	34	1.8	3.0
10	Respiratory distress of newborn (P22)	19	1.0	*
11	Malignant neoplasms (C00-C97)	16	0.8	*
12	Meningitis (G00,G03)	15	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	14	0.7	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	13	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.7	*
...	All other causes (Residual)	499	26.5	44.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,850	100.0	483.1
1	Sudden infant death syndrome (R95)	686	24.1	116.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	12.1	58.3
3	Accidents (unintentional injuries) (V01-X59)	242	8.5	41.0
4	Diseases of the circulatory system (I00-I99)	135	4.7	22.9
5	Septicemia (A40-A41)	124	4.4	21.0
6	Chronic respiratory disease originating in the perinatal period (P27)	112	3.9	19.0
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	104	3.6	17.6
8	Assault (homicide) (*U01,X85-Y09)	101	3.5	17.1
9	Influenza and pneumonia (J10-J18)	97	3.4	16.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	1.4	6.8
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	28	1.0	4.7
12	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	27	0.9	4.6
12	Respiratory distress of newborn (P22)	27	0.9	4.6
14	Necrotizing enterocolitis of newborn (P77)	22	0.8	3.7
15	Meningitis (G00,G03)	21	0.7	3.6
...	All other causes (Residual)	740	26.0	125.4
Postneonates, black, non-Hispanic, male				
...	All causes	1,606	100.0	536.1
1	Sudden infant death syndrome (R95)	394	24.5	131.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	11.0	58.8
3	Accidents (unintentional injuries) (V01-X59)	141	8.8	47.1
4	Diseases of the circulatory system (I00-I99)	70	4.4	23.4
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	68	4.2	22.7
6	Septicemia (A40-A41)	67	4.2	22.4
7	Chronic respiratory disease originating in the perinatal period (P27)	66	4.1	22.0
8	Assault (homicide) (*U01,X85-Y09)	55	3.4	18.4
9	Influenza and pneumonia (J10-J18)	48	3.0	16.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	1.4	7.7
11	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	14	0.9	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.9	*
11	Respiratory distress of newborn (P22)	14	0.9	*
14	Bacterial sepsis of newborn (P36)	11	0.7	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	9	0.6	*
15	Meningitis (G00,G03)	9	0.6	*
15	Necrotizing enterocolitis of newborn (P77)	9	0.6	*
...	All other causes (Residual)	418	26.0	139.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2001

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
	Postneonates, black, non-Hispanic, female			
...	All causes	1,244	100.0	428.4
1	Sudden infant death syndrome (R95)	292	23.5	100.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	13.5	57.9
3	Accidents (unintentional injuries) (V01-X59)	101	8.1	34.8
4	Diseases of the circulatory system (I00-I99)	65	5.2	22.4
5	Septicemia (A40-A41)	57	4.6	19.6
6	Influenza and pneumonia (J10-J18)	49	3.9	16.9
7	Chronic respiratory disease originating in the perinatal period (P27)	46	3.7	15.8
7	Assault (homicide) (*U01,X85-Y09)	46	3.7	15.8
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	36	2.9	12.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.4	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	1.1	*
12	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	13	1.0	*
12	Respiratory distress of newborn (P22)	13	1.0	*
12	Necrotizing enterocolitis of newborn (P77)	13	1.0	*
15	Meningitis (G00,G03)	12	1.0	*
...	All other causes (Residual)	302	24.3	104.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable