

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,943	100.0	563.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,257	25.4	143.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	763	15.4	87.0
3	Sudden infant death syndrome (R95)	295	6.0	33.7
4	Newborn affected by maternal complications of pregnancy (P01)	240	4.9	27.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	181	3.7	20.6
6	Respiratory distress of newborn (P22)	157	3.2	17.9
7	Bacterial sepsis of newborn (P36)	127	2.6	14.5
7	Accidents (unintentional injuries) (V01-X59)	127	2.6	14.5
9	Diseases of the circulatory system (I00-I99)	116	2.3	13.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	103	2.1	11.7
11	Neonatal hemorrhage (P50-P52,P54)	77	1.6	8.8
12	Birth trauma (P10-P15)	76	1.5	8.7
13	Atelectasis (P28.0-P28.1)	69	1.4	7.9
14	Necrotizing enterocolitis of newborn (P77)	66	1.3	7.5
15	Septicemia (A40-A41)	65	1.3	7.4
...	All other causes (Residual)	1,224	24.8	139.6
Infants, Hispanic, male				
...	All causes	2,746	100.0	614.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	653	23.8	146.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	422	15.4	94.4
3	Sudden infant death syndrome (R95)	182	6.6	40.7
4	Newborn affected by maternal complications of pregnancy (P01)	139	5.1	31.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	105	3.8	23.5
6	Respiratory distress of newborn (P22)	92	3.4	20.6
7	Accidents (unintentional injuries) (V01-X59)	83	3.0	18.6
8	Bacterial sepsis of newborn (P36)	81	2.9	18.1
9	Diseases of the circulatory system (I00-I99)	57	2.1	12.8
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	55	2.0	12.3
11	Neonatal hemorrhage (P50-P52,P54)	48	1.7	10.7
12	Birth trauma (P10-P15)	47	1.7	10.5
13	Atelectasis (P28.0-P28.1)	38	1.4	8.5
14	Necrotizing enterocolitis of newborn (P77)	36	1.3	8.1
15	Septicemia (A40-A41)	34	1.2	7.6
...	All other causes (Residual)	674	24.5	150.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,197	100.0	511.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	604	27.5	140.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	341	15.5	79.4
3	Sudden infant death syndrome (R95)	113	5.1	26.3
4	Newborn affected by maternal complications of pregnancy (P01)	101	4.6	23.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	76	3.5	17.7
6	Respiratory distress of newborn (P22)	65	3.0	15.1
7	Diseases of the circulatory system (I00-I99)	59	2.7	13.7
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	2.2	11.2
9	Bacterial sepsis of newborn (P36)	46	2.1	10.7
10	Accidents (unintentional injuries) (V01-X59)	44	2.0	10.2
11	Septicemia (A40-A41)	31	1.4	7.2
11	Atelectasis (P28.0-P28.1)	31	1.4	7.2
13	Necrotizing enterocolitis of newborn (P77)	30	1.4	7.0
14	Birth trauma (P10-P15)	29	1.3	6.8
14	Neonatal hemorrhage (P50-P52,P54)	29	1.3	6.8
...	All other causes (Residual)	550	25.0	128.0
Infants, white, non-Hispanic, both sexes				
...	All causes	13,463	100.0	585.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,068	22.8	133.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,814	13.5	78.9
3	Sudden infant death syndrome (R95)	1,218	9.0	53.0
4	Newborn affected by maternal complications of pregnancy (P01)	810	6.0	35.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	495	3.7	21.5
6	Accidents (unintentional injuries) (V01-X59)	462	3.4	20.1
7	Respiratory distress of newborn (P22)	432	3.2	18.8
8	Bacterial sepsis of newborn (P36)	368	2.7	16.0
9	Diseases of the circulatory system (I00-I99)	327	2.4	14.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	308	2.3	13.4
11	Neonatal hemorrhage (P50-P52,P54)	195	1.4	8.5
12	Atelectasis (P28.0-P28.1)	193	1.4	8.4
13	Birth trauma (P10-P15)	171	1.3	7.4
14	Chronic respiratory disease originating in the perinatal period (P27)	149	1.1	6.5
15	Necrotizing enterocolitis of newborn (P77)	139	1.0	6.0
...	All other causes (Residual)	3,314	24.6	144.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,706	100.0	653.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,643	21.3	139.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,010	13.1	85.7
3	Sudden infant death syndrome (R95)	721	9.4	61.1
4	Newborn affected by maternal complications of pregnancy (P01)	449	5.8	38.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	290	3.8	24.6
6	Accidents (unintentional injuries) (V01-X59)	279	3.6	23.7
7	Respiratory distress of newborn (P22)	271	3.5	23.0
8	Bacterial sepsis of newborn (P36)	210	2.7	17.8
9	Diseases of the circulatory system (I00-I99)	185	2.4	15.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	160	2.1	13.6
11	Neonatal hemorrhage (P50-P52,P54)	119	1.5	10.1
12	Birth trauma (P10-P15)	109	1.4	9.2
12	Atelectasis (P28.0-P28.1)	109	1.4	9.2
14	Chronic respiratory disease originating in the perinatal period (P27)	102	1.3	8.7
15	Necrotizing enterocolitis of newborn (P77)	92	1.2	7.8
...	All other causes (Residual)	1,957	25.4	166.0
Infants, white, non-Hispanic, female				
...	All causes	5,757	100.0	514.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,425	24.8	127.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	804	14.0	71.8
3	Sudden infant death syndrome (R95)	497	8.6	44.4
4	Newborn affected by maternal complications of pregnancy (P01)	361	6.3	32.3
5	Newborn affected by complications of placenta, cord and membranes (P02)	205	3.6	18.3
6	Accidents (unintentional injuries) (V01-X59)	183	3.2	16.4
7	Respiratory distress of newborn (P22)	161	2.8	14.4
8	Bacterial sepsis of newborn (P36)	158	2.7	14.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	148	2.6	13.2
10	Diseases of the circulatory system (I00-I99)	142	2.5	12.7
11	Atelectasis (P28.0-P28.1)	84	1.5	7.5
12	Neonatal hemorrhage (P50-P52,P54)	76	1.3	6.8
13	Hydrops fetalis not due to hemolytic disease (P83.2)	64	1.1	5.7
14	Septicemia (A40-A41)	62	1.1	5.5
14	Birth trauma (P10-P15)	62	1.1	5.5
...	All other causes (Residual)	1,325	23.0	118.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,284	100.0	1,432.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,819	22.0	314.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,015	12.3	175.5
3	Sudden infant death syndrome (R95)	689	8.3	119.1
4	Newborn affected by maternal complications of pregnancy (P01)	558	6.7	96.5
5	Respiratory distress of newborn (P22)	326	3.9	56.4
6	Accidents (unintentional injuries) (V01-X59)	324	3.9	56.0
7	Newborn affected by complications of placenta, cord and membranes (P02)	319	3.9	55.2
8	Bacterial sepsis of newborn (P36)	231	2.8	39.9
9	Diseases of the circulatory system (I00-I99)	190	2.3	32.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	145	1.8	25.1
11	Necrotizing enterocolitis of newborn (P77)	132	1.6	22.8
12	Chronic respiratory disease originating in the perinatal period (P27)	122	1.5	21.1
13	Atelectasis (P28.0-P28.1)	120	1.4	20.7
14	Assault (homicide) (*U01,X85-Y09)	114	1.4	19.7
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	101	1.2	17.5
...	All other causes (Residual)	2,079	25.1	359.5
Infants, black, non-Hispanic, male				
...	All causes	4,520	100.0	1,538.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,020	22.6	347.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	506	11.2	172.3
3	Sudden infant death syndrome (R95)	394	8.7	134.1
4	Newborn affected by maternal complications of pregnancy (P01)	320	7.1	108.9
5	Accidents (unintentional injuries) (V01-X59)	172	3.8	58.6
6	Respiratory distress of newborn (P22)	164	3.6	55.8
7	Newborn affected by complications of placenta, cord and membranes (P02)	158	3.5	53.8
8	Bacterial sepsis of newborn (P36)	116	2.6	39.5
9	Diseases of the circulatory system (I00-I99)	98	2.2	33.4
10	Necrotizing enterocolitis of newborn (P77)	74	1.6	25.2
11	Atelectasis (P28.0-P28.1)	73	1.6	24.9
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	71	1.6	24.2
12	Chronic respiratory disease originating in the perinatal period (P27)	71	1.6	24.2
14	Neonatal hemorrhage (P50-P52,P54)	58	1.3	19.7
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	56	1.2	19.1
...	All other causes (Residual)	1,169	25.9	398.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,764	100.0	1,322.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	799	21.2	280.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	509	13.5	178.9
3	Sudden infant death syndrome (R95)	295	7.8	103.7
4	Newborn affected by maternal complications of pregnancy (P01)	238	6.3	83.6
5	Respiratory distress of newborn (P22)	162	4.3	56.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	161	4.3	56.6
7	Accidents (unintentional injuries) (V01-X59)	152	4.0	53.4
8	Bacterial sepsis of newborn (P36)	115	3.1	40.4
9	Diseases of the circulatory system (I00-I99)	92	2.4	32.3
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	74	2.0	26.0
11	Assault (homicide) (*U01,X85-Y09)	60	1.6	21.1
12	Necrotizing enterocolitis of newborn (P77)	58	1.5	20.4
13	Chronic respiratory disease originating in the perinatal period (P27)	51	1.4	17.9
14	Atelectasis (P28.0-P28.1)	47	1.2	16.5
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	45	1.2	15.8
...	All other causes (Residual)	906	24.1	318.4
Neonates, Hispanic, both sexes				
...	All causes	3,331	100.0	380.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	872	26.2	99.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	743	22.3	84.8
3	Newborn affected by maternal complications of pregnancy (P01)	239	7.2	27.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	173	5.2	19.7
5	Respiratory distress of newborn (P22)	149	4.5	17.0
6	Bacterial sepsis of newborn (P36)	119	3.6	13.6
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	98	2.9	11.2
8	Neonatal hemorrhage (P50-P52,P54)	76	2.3	8.7
9	Birth trauma (P10-P15)	74	2.2	8.4
10	Atelectasis (P28.0-P28.1)	68	2.0	7.8
11	Necrotizing enterocolitis of newborn (P77)	59	1.8	6.7
12	Diseases of the circulatory system (I00-I99)	35	1.1	4.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	35	1.1	4.0
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	32	1.0	3.7
15	Pulmonary hemorrhage originating in the perinatal period (P26)	28	0.8	3.2
...	All other causes (Residual)	531	15.9	60.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,847	100.0	413.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	453	24.5	101.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	410	22.2	91.7
3	Newborn affected by maternal complications of pregnancy (P01)	138	7.5	30.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	100	5.4	22.4
5	Respiratory distress of newborn (P22)	90	4.9	20.1
6	Bacterial sepsis of newborn (P36)	79	4.3	17.7
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	51	2.8	11.4
8	Neonatal hemorrhage (P50-P52,P54)	47	2.5	10.5
9	Birth trauma (P10-P15)	46	2.5	10.3
10	Atelectasis (P28.0-P28.1)	38	2.1	8.5
11	Necrotizing enterocolitis of newborn (P77)	32	1.7	7.2
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	19	1.0	*
13	Diseases of the circulatory system (I00-I99)	18	1.0	*
14	Pulmonary hemorrhage originating in the perinatal period (P26)	17	0.9	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	15	0.8	*
...	All other causes (Residual)	294	15.9	65.8
Neonates, Hispanic, female				
...	All causes	1,484	100.0	345.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	419	28.2	97.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	333	22.4	77.5
3	Newborn affected by maternal complications of pregnancy (P01)	101	6.8	23.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	73	4.9	17.0
5	Respiratory distress of newborn (P22)	59	4.0	13.7
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	47	3.2	10.9
7	Bacterial sepsis of newborn (P36)	40	2.7	9.3
8	Atelectasis (P28.0-P28.1)	30	2.0	7.0
9	Neonatal hemorrhage (P50-P52,P54)	29	2.0	6.8
10	Birth trauma (P10-P15)	28	1.9	6.5
11	Necrotizing enterocolitis of newborn (P77)	27	1.8	6.3
12	Hydrops fetalis not due to hemolytic disease (P83.2)	20	1.3	4.7
13	Diseases of the circulatory system (I00-I99)	17	1.1	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	13	0.9	*
15	Pulmonary hemorrhage originating in the perinatal period (P26)	11	0.7	*
...	All other causes (Residual)	237	16.0	55.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	9,014	100.0	392.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,259	25.1	98.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,777	19.7	77.3
3	Newborn affected by maternal complications of pregnancy (P01)	804	8.9	35.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	490	5.4	21.3
5	Respiratory distress of newborn (P22)	404	4.5	17.6
6	Bacterial sepsis of newborn (P36)	346	3.8	15.1
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	288	3.2	12.5
8	Neonatal hemorrhage (P50-P52,P54)	189	2.1	8.2
9	Atelectasis (P28.0-P28.1)	181	2.0	7.9
10	Birth trauma (P10-P15)	170	1.9	7.4
11	Diseases of the circulatory system (I00-I99)	142	1.6	6.2
12	Necrotizing enterocolitis of newborn (P77)	123	1.4	5.4
13	Hydrops fetalis not due to hemolytic disease (P83.2)	116	1.3	5.0
14	Sudden infant death syndrome (R95)	105	1.2	4.6
15	Pulmonary hemorrhage originating in the perinatal period (P26)	88	1.0	3.8
...	All other causes (Residual)	1,532	17.0	66.7
Neonates, white, non-Hispanic, male				
...	All causes	5,090	100.0	431.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,241	24.4	105.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	989	19.4	83.9
3	Newborn affected by maternal complications of pregnancy (P01)	447	8.8	37.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	287	5.6	24.3
5	Respiratory distress of newborn (P22)	254	5.0	21.5
6	Bacterial sepsis of newborn (P36)	199	3.9	16.9
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	152	3.0	12.9
8	Neonatal hemorrhage (P50-P52,P54)	117	2.3	9.9
9	Birth trauma (P10-P15)	108	2.1	9.2
10	Atelectasis (P28.0-P28.1)	99	1.9	8.4
11	Diseases of the circulatory system (I00-I99)	82	1.6	7.0
12	Necrotizing enterocolitis of newborn (P77)	79	1.6	6.7
13	Hydrops fetalis not due to hemolytic disease (P83.2)	55	1.1	4.7
14	Sudden infant death syndrome (R95)	54	1.1	4.6
15	Pulmonary hemorrhage originating in the perinatal period (P26)	44	0.9	3.7
...	All other causes (Residual)	883	17.3	74.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,924	100.0	350.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,018	25.9	91.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	788	20.1	70.4
3	Newborn affected by maternal complications of pregnancy (P01)	357	9.1	31.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	203	5.2	18.1
5	Respiratory distress of newborn (P22)	150	3.8	13.4
6	Bacterial sepsis of newborn (P36)	147	3.7	13.1
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	136	3.5	12.2
8	Atelectasis (P28.0-P28.1)	82	2.1	7.3
9	Neonatal hemorrhage (P50-P52,P54)	72	1.8	6.4
10	Birth trauma (P10-P15)	62	1.6	5.5
11	Hydrops fetalis not due to hemolytic disease (P83.2)	61	1.6	5.5
12	Diseases of the circulatory system (I00-I99)	60	1.5	5.4
13	Sudden infant death syndrome (R95)	51	1.3	4.6
14	Pulmonary hemorrhage originating in the perinatal period (P26)	44	1.1	3.9
14	Necrotizing enterocolitis of newborn (P77)	44	1.1	3.9
...	All other causes (Residual)	649	16.5	58.0
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,470	100.0	945.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,778	32.5	307.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	646	11.8	111.7
3	Newborn affected by maternal complications of pregnancy (P01)	552	10.1	95.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	315	5.8	54.5
5	Respiratory distress of newborn (P22)	298	5.4	51.5
6	Bacterial sepsis of newborn (P36)	218	4.0	37.7
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	131	2.4	22.7
8	Necrotizing enterocolitis of newborn (P77)	114	2.1	19.7
9	Atelectasis (P28.0-P28.1)	112	2.0	19.4
10	Neonatal hemorrhage (P50-P52,P54)	97	1.8	16.8
11	Birth trauma (P10-P15)	81	1.5	14.0
12	Pulmonary hemorrhage originating in the perinatal period (P26)	61	1.1	10.5
13	Sudden infant death syndrome (R95)	60	1.1	10.4
14	Diseases of the circulatory system (I00-I99)	50	0.9	8.6
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	41	0.7	7.1
...	All other causes (Residual)	916	16.7	158.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	2,958	100.0	1,007.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	993	33.6	338.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	318	10.8	108.3
3	Newborn affected by maternal complications of pregnancy (P01)	314	10.6	106.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	156	5.3	53.1
5	Respiratory distress of newborn (P22)	146	4.9	49.7
6	Bacterial sepsis of newborn (P36)	110	3.7	37.4
7	Atelectasis (P28.0-P28.1)	68	2.3	23.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	66	2.2	22.5
9	Necrotizing enterocolitis of newborn (P77)	63	2.1	21.4
10	Neonatal hemorrhage (P50-P52,P54)	57	1.9	19.4
11	Birth trauma (P10-P15)	51	1.7	17.4
12	Pulmonary hemorrhage originating in the perinatal period (P26)	37	1.3	12.6
13	Diseases of the circulatory system (I00-I99)	27	0.9	9.2
14	Sudden infant death syndrome (R95)	25	0.8	8.5
15	Newborn affected by other complications of labor and delivery (P03)	23	0.8	7.8
...	All other causes (Residual)	504	17.0	171.6
Neonates, black, non-Hispanic, female				
...	All causes	2,512	100.0	882.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	785	31.3	275.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	328	13.1	115.3
3	Newborn affected by maternal complications of pregnancy (P01)	238	9.5	83.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	159	6.3	55.9
5	Respiratory distress of newborn (P22)	152	6.1	53.4
6	Bacterial sepsis of newborn (P36)	108	4.3	38.0
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	65	2.6	22.8
8	Necrotizing enterocolitis of newborn (P77)	51	2.0	17.9
9	Atelectasis (P28.0-P28.1)	44	1.8	15.5
10	Neonatal hemorrhage (P50-P52,P54)	40	1.6	14.1
11	Sudden infant death syndrome (R95)	35	1.4	12.3
12	Birth trauma (P10-P15)	30	1.2	10.5
13	Pulmonary hemorrhage originating in the perinatal period (P26)	24	1.0	8.4
14	Diseases of the circulatory system (I00-I99)	23	0.9	8.1
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	20	0.8	7.0
...	All other causes (Residual)	410	16.3	144.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,612	100.0	183.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	23.9	43.9
2	Sudden infant death syndrome (R95)	279	17.3	31.8
3	Accidents (unintentional injuries) (V01-X59)	116	7.2	13.2
4	Diseases of the circulatory system (I00-I99)	81	5.0	9.2
5	Septicemia (A40-A41)	65	4.0	7.4
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	45	2.8	5.1
7	Assault (homicide) (*U01,X85-Y09)	42	2.6	4.8
8	Influenza and pneumonia (J10-J18)	41	2.5	4.7
9	Chronic respiratory disease originating in the perinatal period (P27)	29	1.8	3.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	20	1.2	2.3
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	1.1	*
12	Malignant neoplasms (C00-C97)	17	1.1	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	15	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D	11	0.7	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.5	*
15	Newborn affected by complications of placenta, cord and membranes (P02)	8	0.5	*
15	Respiratory distress of newborn (P22)	8	0.5	*
15	Bacterial sepsis of newborn (P36)	8	0.5	*
...	All other causes (Residual)	416	25.8	47.5
Postneonates, Hispanic, male				
...	All causes	899	100.0	201.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	22.2	44.7
2	Sudden infant death syndrome (R95)	170	18.9	38.0
3	Accidents (unintentional injuries) (V01-X59)	78	8.7	17.4
4	Diseases of the circulatory system (I00-I99)	39	4.3	8.7
5	Septicemia (A40-A41)	34	3.8	7.6
6	Influenza and pneumonia (J10-J18)	27	3.0	6.0
7	Assault (homicide) (*U01,X85-Y09)	23	2.6	5.1
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	22	2.4	4.9
9	Chronic respiratory disease originating in the perinatal period (P27)	21	2.3	4.7
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	12	1.3	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	1.1	*
12	Malignant neoplasms (C00-C97)	9	1.0	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	9	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D	5	0.6	*
14	Newborn affected by complications of placenta, cord and membranes (P02)	5	0.6	*
...	All other causes (Residual)	235	26.1	52.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	713	100.0	166.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	25.9	43.1
2	Sudden infant death syndrome (R95)	109	15.3	25.4
3	Diseases of the circulatory system (I00-I99)	42	5.9	9.8
4	Accidents (unintentional injuries) (V01-X59)	38	5.3	8.8
5	Septicemia (A40-A41)	31	4.3	7.2
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	23	3.2	5.4
7	Assault (homicide) (*U01,X85-Y09)	19	2.7	*
8	Influenza and pneumonia (J10-J18)	14	2.0	*
9	Malignant neoplasms (C00-C97)	8	1.1	*
9	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	1.1	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	8	1.1	*
9	Chronic respiratory disease originating in the perinatal period (P27)	8	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D	6	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	6	0.8	*
13	Respiratory distress of newborn (P22)	6	0.8	*
13	Bacterial sepsis of newborn (P36)	6	0.8	*
...	All other causes (Residual)	196	27.5	45.6
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,449	100.0	193.6
1	Sudden infant death syndrome (R95)	1,113	25.0	48.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	809	18.2	35.2
3	Accidents (unintentional injuries) (V01-X59)	411	9.2	17.9
4	Diseases of the circulatory system (I00-I99)	185	4.2	8.0
5	Septicemia (A40-A41)	129	2.9	5.6
6	Chronic respiratory disease originating in the perinatal period (P27)	118	2.7	5.1
7	Assault (homicide) (*U01,X85-Y09)	103	2.3	4.5
8	Influenza and pneumonia (J10-J18)	99	2.2	4.3
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	80	1.8	3.5
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	37	0.8	1.6
11	Meningitis (G00,G03)	35	0.8	1.5
12	Malignant neoplasms (C00-C97)	30	0.7	1.3
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	28	0.6	1.2
13	Respiratory distress of newborn (P22)	28	0.6	1.2
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	27	0.6	1.2
...	All other causes (Residual)	1,217	27.4	53.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, male				
...	All causes	2,616	100.0	221.9
1	Sudden infant death syndrome (R95)	667	25.5	56.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	402	15.4	34.1
3	Accidents (unintentional injuries) (V01-X59)	257	9.8	21.8
4	Diseases of the circulatory system (I00-I99)	103	3.9	8.7
5	Chronic respiratory disease originating in the perinatal period (P27)	85	3.2	7.2
6	Septicemia (A40-A41)	67	2.6	5.7
6	Influenza and pneumonia (J10-J18)	67	2.6	5.7
8	Assault (homicide) (*U01,X85-Y09)	61	2.3	5.2
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	51	1.9	4.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	0.8	1.8
11	Malignant neoplasms (C00-C97)	20	0.8	1.7
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	20	0.8	1.7
11	Acute bronchitis and acute bronchiolitis (J20-J21)	20	0.8	1.7
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.7	*
15	Meningitis (G00,G03)	18	0.7	*
...	All other causes (Residual)	738	28.2	62.6
Postneonates, white, non-Hispanic, female				
...	All causes	1,833	100.0	163.8
1	Sudden infant death syndrome (R95)	446	24.3	39.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	407	22.2	36.4
3	Accidents (unintentional injuries) (V01-X59)	154	8.4	13.8
4	Diseases of the circulatory system (I00-I99)	82	4.5	7.3
5	Septicemia (A40-A41)	62	3.4	5.5
6	Assault (homicide) (*U01,X85-Y09)	42	2.3	3.8
7	Chronic respiratory disease originating in the perinatal period (P27)	33	1.8	2.9
8	Influenza and pneumonia (J10-J18)	32	1.7	2.9
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	29	1.6	2.6
10	Meningitis (G00,G03)	17	0.9	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D	12	0.7	*
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	12	0.7	*
14	Respiratory distress of newborn (P22)	11	0.6	*
14	Bacterial sepsis of newborn (P36)	11	0.6	*
...	All other causes (Residual)	467	25.5	41.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,814	100.0	486.6
1	Sudden infant death syndrome (R95)	629	22.4	108.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	369	13.1	63.8
3	Accidents (unintentional injuries) (V01-X59)	295	10.5	51.0
4	Diseases of the circulatory system (I00-I99)	140	5.0	24.2
5	Chronic respiratory disease originating in the perinatal period (P27)	103	3.7	17.8
6	Assault (homicide) (*U01,X85-Y09)	96	3.4	16.6
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	94	3.3	16.3
8	Septicemia (A40-A41)	85	3.0	14.7
9	Influenza and pneumonia (J10-J18)	76	2.7	13.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	41	1.5	7.1
11	Respiratory distress of newborn (P22)	28	1.0	4.8
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	24	0.9	4.1
13	Necrotizing enterocolitis of newborn (P77)	18	0.6	*
14	Meningitis (G00,G03)	17	0.6	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	15	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	15	0.5	*
...	All other causes (Residual)	769	27.3	133.0
Postneonates, black, non-Hispanic, male				
...	All causes	1,562	100.0	531.7
1	Sudden infant death syndrome (R95)	369	23.6	125.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	12.0	64.0
3	Accidents (unintentional injuries) (V01-X59)	155	9.9	52.8
4	Diseases of the circulatory system (I00-I99)	71	4.5	24.2
5	Chronic respiratory disease originating in the perinatal period (P27)	58	3.7	19.7
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	50	3.2	17.0
7	Septicemia (A40-A41)	46	2.9	15.7
8	Assault (homicide) (*U01,X85-Y09)	45	2.9	15.3
9	Influenza and pneumonia (J10-J18)	42	2.7	14.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	1.7	9.2
11	Respiratory distress of newborn (P22)	18	1.2	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	17	1.1	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.7	*
13	Necrotizing enterocolitis of newborn (P77)	11	0.7	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	9	0.6	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	9	0.6	*
...	All other causes (Residual)	436	27.9	148.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2002

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
	Postneonates, black, non-Hispanic, female			
...	All causes	1,252	100.0	439.9
1	Sudden infant death syndrome (R95)	260	20.8	91.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	14.5	63.6
3	Accidents (unintentional injuries) (V01-X59)	140	11.2	49.2
4	Diseases of the circulatory system (I00-I99)	69	5.5	24.2
5	Assault (homicide) (*U01,X85-Y09)	51	4.1	17.9
6	Chronic respiratory disease originating in the perinatal period (P27)	45	3.6	15.8
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	44	3.5	15.5
8	Septicemia (A40-A41)	39	3.1	13.7
9	Influenza and pneumonia (J10-J18)	34	2.7	11.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	1.1	*
11	Respiratory distress of newborn (P22)	10	0.8	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune me	9	0.7	*
12	Meningitis (G00,G03)	9	0.7	*
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	9	0.7	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	0.6	*
...	All other causes (Residual)	330	26.4	116.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable