

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,281	100.0	578.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,324	25.1	145.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	878	16.6	96.2
3	Newborn affected by maternal complications of pregnancy (P01)	309	5.9	33.9
4	Sudden infant death syndrome (R95)	290	5.5	31.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	198	3.7	21.7
6	Respiratory distress of newborn (P22)	148	2.8	16.2
7	Accidents (unintentional injuries) (V01-X59)	131	2.5	14.4
8	Neonatal hemorrhage (P50-P52,P54)	127	2.4	13.9
9	Bacterial sepsis of newborn (P36)	124	2.3	13.6
10	Diseases of the circulatory system (I00-I99)	106	2.0	11.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	88	1.7	9.6
12	Atelectasis (P28.0-P28.1)	85	1.6	9.3
13	Necrotizing enterocolitis of newborn (P77)	81	1.5	8.9
14	Influenza and pneumonia (J10-J18)	66	1.2	7.2
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	63	1.2	6.9
...	All other causes (Residual)	1,263	23.9	138.4
Infants, Hispanic, male				
...	All causes	2,940	100.0	631.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	693	23.6	149.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	470	16.0	101.0
3	Sudden infant death syndrome (R95)	182	6.2	39.1
4	Newborn affected by maternal complications of pregnancy (P01)	173	5.9	37.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	112	3.8	24.1
6	Respiratory distress of newborn (P22)	85	2.9	18.3
7	Accidents (unintentional injuries) (V01-X59)	77	2.6	16.6
8	Bacterial sepsis of newborn (P36)	74	2.5	15.9
9	Neonatal hemorrhage (P50-P52,P54)	72	2.4	15.5
10	Diseases of the circulatory system (I00-I99)	59	2.0	12.7
11	Atelectasis (P28.0-P28.1)	58	2.0	12.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	1.6	10.3
13	Necrotizing enterocolitis of newborn (P77)	45	1.5	9.7
14	Influenza and pneumonia (J10-J18)	40	1.4	8.6
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	40	1.4	8.6
...	All other causes (Residual)	712	24.2	153.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,341	100.0	523.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	631	27.0	141.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	408	17.4	91.3
3	Newborn affected by maternal complications of pregnancy (P01)	136	5.8	30.4
4	Sudden infant death syndrome (R95)	108	4.6	24.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	86	3.7	19.2
6	Respiratory distress of newborn (P22)	63	2.7	14.1
7	Neonatal hemorrhage (P50-P52,P54)	55	2.3	12.3
8	Accidents (unintentional injuries) (V01-X59)	54	2.3	12.1
9	Bacterial sepsis of newborn (P36)	50	2.1	11.2
10	Diseases of the circulatory system (I00-I99)	47	2.0	10.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	1.7	8.9
12	Necrotizing enterocolitis of newborn (P77)	36	1.5	8.1
13	Atelectasis (P28.0-P28.1)	27	1.2	6.0
14	Influenza and pneumonia (J10-J18)	26	1.1	5.8
15	Assault (homicide) (*U01,X85-Y09)	25	1.1	5.6
...	All other causes (Residual)	549	23.5	122.8
Infants, white, non-Hispanic, both sexes				
...	All causes	13,223	100.0	569.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,977	22.5	128.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,890	14.3	81.4
3	Sudden infant death syndrome (R95)	1,104	8.3	47.5
4	Newborn affected by maternal complications of pregnancy (P01)	773	5.8	33.3
5	Newborn affected by complications of placenta, cord and membranes (P02)	545	4.1	23.5
6	Accidents (unintentional injuries) (V01-X59)	496	3.8	21.4
7	Respiratory distress of newborn (P22)	394	3.0	17.0
8	Bacterial sepsis of newborn (P36)	351	2.7	15.1
9	Neonatal hemorrhage (P50-P52,P54)	346	2.6	14.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	302	2.3	13.0
11	Diseases of the circulatory system (I00-I99)	268	2.0	11.5
12	Atelectasis (P28.0-P28.1)	234	1.8	10.1
13	Assault (homicide) (*U01,X85-Y09)	156	1.2	6.7
14	Necrotizing enterocolitis of newborn (P77)	148	1.1	6.4
15	Influenza and pneumonia (J10-J18)	143	1.1	6.2
...	All other causes (Residual)	3,096	23.4	133.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,591	100.0	637.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,596	21.0	134.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,087	14.3	91.3
3	Sudden infant death syndrome (R95)	682	9.0	57.3
4	Newborn affected by maternal complications of pregnancy (P01)	443	5.8	37.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	288	3.8	24.2
6	Accidents (unintentional injuries) (V01-X59)	283	3.7	23.8
7	Respiratory distress of newborn (P22)	239	3.1	20.1
8	Bacterial sepsis of newborn (P36)	218	2.9	18.3
9	Neonatal hemorrhage (P50-P52,P54)	209	2.8	17.5
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	169	2.2	14.2
11	Atelectasis (P28.0-P28.1)	148	1.9	12.4
12	Diseases of the circulatory system (I00-I99)	144	1.9	12.1
13	Assault (homicide) (*U01,X85-Y09)	94	1.2	7.9
14	Influenza and pneumonia (J10-J18)	87	1.1	7.3
15	Necrotizing enterocolitis of newborn (P77)	86	1.1	7.2
...	All other causes (Residual)	1,818	23.9	152.6
Infants, white, non-Hispanic, female				
...	All causes	5,632	100.0	498.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,381	24.5	122.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	803	14.3	71.0
3	Sudden infant death syndrome (R95)	422	7.5	37.3
4	Newborn affected by maternal complications of pregnancy (P01)	330	5.9	29.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	257	4.6	22.7
6	Accidents (unintentional injuries) (V01-X59)	213	3.8	18.8
7	Respiratory distress of newborn (P22)	155	2.8	13.7
8	Neonatal hemorrhage (P50-P52,P54)	137	2.4	12.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	133	2.4	11.8
9	Bacterial sepsis of newborn (P36)	133	2.4	11.8
11	Diseases of the circulatory system (I00-I99)	124	2.2	11.0
12	Atelectasis (P28.0-P28.1)	86	1.5	7.6
13	Necrotizing enterocolitis of newborn (P77)	62	1.1	5.5
13	Assault (homicide) (*U01,X85-Y09)	62	1.1	5.5
15	Influenza and pneumonia (J10-J18)	56	1.0	5.0
...	All other causes (Residual)	1,278	22.7	113.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,159	100.0	1,416.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,849	22.7	321.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,015	12.4	176.2
3	Sudden infant death syndrome (R95)	677	8.3	117.5
4	Newborn affected by maternal complications of pregnancy (P01)	556	6.8	96.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	307	3.8	53.3
6	Accidents (unintentional injuries) (V01-X59)	283	3.5	49.1
7	Bacterial sepsis of newborn (P36)	264	3.2	45.8
8	Respiratory distress of newborn (P22)	250	3.1	43.4
9	Diseases of the circulatory system (I00-I99)	174	2.1	30.2
10	Necrotizing enterocolitis of newborn (P77)	156	1.9	27.1
11	Neonatal hemorrhage (P50-P52,P54)	142	1.7	24.7
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	140	1.7	24.3
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	124	1.5	21.5
14	Septicemia (A40-A41)	119	1.5	20.7
15	Atelectasis (P28.0-P28.1)	109	1.3	18.9
...	All other causes (Residual)	1,994	24.4	346.2
Infants, black, non-Hispanic, male				
...	All causes	4,601	100.0	1,570.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,053	22.9	359.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	558	12.1	190.4
3	Sudden infant death syndrome (R95)	357	7.8	121.8
4	Newborn affected by maternal complications of pregnancy (P01)	306	6.7	104.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	166	3.6	56.6
6	Accidents (unintentional injuries) (V01-X59)	158	3.4	53.9
7	Bacterial sepsis of newborn (P36)	152	3.3	51.9
8	Respiratory distress of newborn (P22)	143	3.1	48.8
9	Diseases of the circulatory system (I00-I99)	107	2.3	36.5
10	Necrotizing enterocolitis of newborn (P77)	88	1.9	30.0
11	Neonatal hemorrhage (P50-P52,P54)	82	1.8	28.0
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	74	1.6	25.3
13	Septicemia (A40-A41)	70	1.5	23.9
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	70	1.5	23.9
13	Chronic respiratory disease originating in the perinatal period (P27)	70	1.5	23.9
...	All other causes (Residual)	1,147	24.9	391.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,558	100.0	1,257.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	796	22.4	281.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	457	12.8	161.5
3	Sudden infant death syndrome (R95)	320	9.0	113.1
4	Newborn affected by maternal complications of pregnancy (P01)	250	7.0	88.3
5	Newborn affected by complications of placenta, cord and membranes (P02)	141	4.0	49.8
6	Accidents (unintentional injuries) (V01-X59)	125	3.5	44.2
7	Bacterial sepsis of newborn (P36)	112	3.1	39.6
8	Respiratory distress of newborn (P22)	107	3.0	37.8
9	Necrotizing enterocolitis of newborn (P77)	68	1.9	24.0
10	Diseases of the circulatory system (I00-I99)	67	1.9	23.7
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	66	1.9	23.3
12	Neonatal hemorrhage (P50-P52,P54)	60	1.7	21.2
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	54	1.5	19.1
14	Septicemia (A40-A41)	49	1.4	17.3
15	Atelectasis (P28.0-P28.1)	48	1.3	17.0
...	All other causes (Residual)	838	23.6	296.1
Neonates, Hispanic, both sexes				
...	All causes	3,601	100.0	394.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	927	25.7	101.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	855	23.7	93.7
3	Newborn affected by maternal complications of pregnancy (P01)	309	8.6	33.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	196	5.4	21.5
5	Respiratory distress of newborn (P22)	134	3.7	14.7
6	Neonatal hemorrhage (P50-P52,P54)	125	3.5	13.7
7	Bacterial sepsis of newborn (P36)	118	3.3	12.9
8	Atelectasis (P28.0-P28.1)	82	2.3	9.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	81	2.2	8.9
10	Necrotizing enterocolitis of newborn (P77)	72	2.0	7.9
11	Diseases of the circulatory system (I00-I99)	37	1.0	4.1
12	Newborn affected by other complications of labor and delivery (P03)	33	0.9	3.6
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	29	0.8	3.2
13	Hydrops fetalis not due to hemolytic disease (P83.2)	29	0.8	3.2
15	Pulmonary hemorrhage originating in the perinatal period (P26)	28	0.8	3.1
...	All other causes (Residual)	546	15.2	59.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,971	100.0	423.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	494	25.1	106.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	452	22.9	97.2
3	Newborn affected by maternal complications of pregnancy (P01)	173	8.8	37.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	110	5.6	23.6
5	Respiratory distress of newborn (P22)	79	4.0	17.0
6	Neonatal hemorrhage (P50-P52,P54)	71	3.6	15.3
7	Bacterial sepsis of newborn (P36)	70	3.6	15.0
8	Atelectasis (P28.0-P28.1)	56	2.8	12.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	44	2.2	9.5
10	Necrotizing enterocolitis of newborn (P77)	39	2.0	8.4
11	Diseases of the circulatory system (I00-I99)	21	1.1	4.5
12	Sudden infant death syndrome (R95)	19	1.0	*
13	Newborn affected by other complications of labor and delivery (P03)	18	0.9	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	17	0.9	*
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	16	0.8	*
...	All other causes (Residual)	292	14.8	62.8
Neonates, Hispanic, female				
...	All causes	1,630	100.0	364.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	433	26.6	96.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	403	24.7	90.1
3	Newborn affected by maternal complications of pregnancy (P01)	136	8.3	30.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	86	5.3	19.2
5	Respiratory distress of newborn (P22)	55	3.4	12.3
6	Neonatal hemorrhage (P50-P52,P54)	54	3.3	12.1
7	Bacterial sepsis of newborn (P36)	48	2.9	10.7
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	37	2.3	8.3
9	Necrotizing enterocolitis of newborn (P77)	33	2.0	7.4
10	Atelectasis (P28.0-P28.1)	26	1.6	5.8
11	Diseases of the circulatory system (I00-I99)	16	1.0	*
11	Pulmonary hemorrhage originating in the perinatal period (P26)	16	1.0	*
13	Newborn affected by other complications of labor and delivery (P03)	15	0.9	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	13	0.8	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	12	0.7	*
...	All other causes (Residual)	247	15.2	55.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	8,906	100.0	383.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,181	24.5	93.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,864	20.9	80.3
3	Newborn affected by maternal complications of pregnancy (P01)	764	8.6	32.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	538	6.0	23.2
5	Respiratory distress of newborn (P22)	371	4.2	16.0
6	Neonatal hemorrhage (P50-P52,P54)	342	3.8	14.7
7	Bacterial sepsis of newborn (P36)	334	3.8	14.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	281	3.2	12.1
9	Atelectasis (P28.0-P28.1)	230	2.6	9.9
10	Necrotizing enterocolitis of newborn (P77)	127	1.4	5.5
11	Hydrops fetalis not due to hemolytic disease (P83.2)	118	1.3	5.1
12	Sudden infant death syndrome (R95)	103	1.2	4.4
13	Diseases of the circulatory system (I00-I99)	97	1.1	4.2
14	Newborn affected by other complications of labor and delivery (P03)	82	0.9	3.5
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	69	0.8	3.0
...	All other causes (Residual)	1,405	15.8	60.5
Neonates, white, non-Hispanic, male				
...	All causes	5,076	100.0	426.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,173	23.1	98.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,077	21.2	90.4
3	Newborn affected by maternal complications of pregnancy (P01)	436	8.6	36.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	285	5.6	23.9
5	Respiratory distress of newborn (P22)	228	4.5	19.1
6	Bacterial sepsis of newborn (P36)	208	4.1	17.5
7	Neonatal hemorrhage (P50-P52,P54)	207	4.1	17.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	157	3.1	13.2
9	Atelectasis (P28.0-P28.1)	146	2.9	12.3
10	Necrotizing enterocolitis of newborn (P77)	75	1.5	6.3
11	Hydrops fetalis not due to hemolytic disease (P83.2)	65	1.3	5.5
12	Diseases of the circulatory system (I00-I99)	55	1.1	4.6
13	Sudden infant death syndrome (R95)	54	1.1	4.5
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	46	0.9	3.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	41	0.8	3.4
...	All other causes (Residual)	823	16.2	69.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,830	100.0	338.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,008	26.3	89.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	787	20.5	69.6
3	Newborn affected by maternal complications of pregnancy (P01)	328	8.6	29.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	253	6.6	22.4
5	Respiratory distress of newborn (P22)	143	3.7	12.6
6	Neonatal hemorrhage (P50-P52,P54)	135	3.5	11.9
7	Bacterial sepsis of newborn (P36)	126	3.3	11.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	124	3.2	11.0
9	Atelectasis (P28.0-P28.1)	84	2.2	7.4
10	Hydrops fetalis not due to hemolytic disease (P83.2)	53	1.4	4.7
11	Necrotizing enterocolitis of newborn (P77)	52	1.4	4.6
12	Sudden infant death syndrome (R95)	49	1.3	4.3
13	Newborn affected by other complications of labor and delivery (P03)	43	1.1	3.8
14	Diseases of the circulatory system (I00-I99)	42	1.1	3.7
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	23	0.6	2.0
...	All other causes (Residual)	580	15.1	51.3
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,452	100.0	946.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,809	33.2	314.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	673	12.3	116.8
3	Newborn affected by maternal complications of pregnancy (P01)	549	10.1	95.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	302	5.5	52.4
5	Bacterial sepsis of newborn (P36)	254	4.7	44.1
6	Respiratory distress of newborn (P22)	231	4.2	40.1
7	Necrotizing enterocolitis of newborn (P77)	140	2.6	24.3
8	Neonatal hemorrhage (P50-P52,P54)	139	2.5	24.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	128	2.3	22.2
10	Atelectasis (P28.0-P28.1)	106	1.9	18.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	63	1.2	10.9
12	Sudden infant death syndrome (R95)	57	1.0	9.9
13	Newborn affected by other complications of labor and delivery (P03)	42	0.8	7.3
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	40	0.7	6.9
15	Diseases of the circulatory system (I00-I99)	39	0.7	6.8
...	All other causes (Residual)	880	16.1	152.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,068	100.0	1,046.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,032	33.6	352.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	365	11.9	124.6
3	Newborn affected by maternal complications of pregnancy (P01)	302	9.8	103.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	164	5.3	56.0
5	Bacterial sepsis of newborn (P36)	146	4.8	49.8
6	Respiratory distress of newborn (P22)	135	4.4	46.1
7	Neonatal hemorrhage (P50-P52,P54)	79	2.6	27.0
7	Necrotizing enterocolitis of newborn (P77)	79	2.6	27.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	67	2.2	22.9
10	Atelectasis (P28.0-P28.1)	59	1.9	20.1
11	Pulmonary hemorrhage originating in the perinatal period (P26)	41	1.3	14.0
12	Sudden infant death syndrome (R95)	30	1.0	10.2
13	Diseases of the circulatory system (I00-I99)	27	0.9	9.2
14	Newborn affected by other complications of labor and delivery (P03)	25	0.8	8.5
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	22	0.7	7.5
...	All other causes (Residual)	495	16.1	168.9
Neonates, black, non-Hispanic, female				
...	All causes	2,384	100.0	842.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	777	32.6	274.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	308	12.9	108.8
3	Newborn affected by maternal complications of pregnancy (P01)	247	10.4	87.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	138	5.8	48.8
5	Bacterial sepsis of newborn (P36)	108	4.5	38.2
6	Respiratory distress of newborn (P22)	96	4.0	33.9
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	61	2.6	21.6
7	Necrotizing enterocolitis of newborn (P77)	61	2.6	21.6
9	Neonatal hemorrhage (P50-P52,P54)	60	2.5	21.2
10	Atelectasis (P28.0-P28.1)	47	2.0	16.6
11	Sudden infant death syndrome (R95)	27	1.1	9.5
12	Pulmonary hemorrhage originating in the perinatal period (P26)	22	0.9	7.8
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	18	0.8	*
14	Newborn affected by other complications of labor and delivery (P03)	17	0.7	*
14	Accidents (unintentional injuries) (V01-X59)	17	0.7	*
...	All other causes (Residual)	380	15.9	134.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,680	100.0	184.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	397	23.6	43.5
2	Sudden infant death syndrome (R95)	265	15.8	29.0
3	Accidents (unintentional injuries) (V01-X59)	119	7.1	13.0
4	Diseases of the circulatory system (I00-I99)	69	4.1	7.6
5	Influenza and pneumonia (J10-J18)	60	3.6	6.6
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	57	3.4	6.2
7	Septicemia (A40-A41)	50	3.0	5.5
7	Assault (homicide) (*U01,X85-Y09)	50	3.0	5.5
9	Chronic respiratory disease originating in the perinatal period (P27)	35	2.1	3.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	1.4	2.5
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	19	1.1	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	17	1.0	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.8	*
13	Respiratory distress of newborn (P22)	14	0.8	*
15	Malignant neoplasms (C00-C97)	13	0.8	*
...	All other causes (Residual)	478	28.5	52.4
Postneonates, Hispanic, male				
...	All causes	969	100.0	208.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	20.5	42.8
2	Sudden infant death syndrome (R95)	163	16.8	35.0
3	Accidents (unintentional injuries) (V01-X59)	72	7.4	15.5
4	Diseases of the circulatory system (I00-I99)	38	3.9	8.2
5	Influenza and pneumonia (J10-J18)	37	3.8	8.0
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	36	3.7	7.7
7	Assault (homicide) (*U01,X85-Y09)	30	3.1	6.4
8	Septicemia (A40-A41)	29	3.0	6.2
9	Chronic respiratory disease originating in the perinatal period (P27)	25	2.6	5.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.9	*
11	Acute bronchitis and acute bronchiolitis (J20-J21)	12	1.2	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	11	1.1	*
13	Meningitis (G00,G03)	6	0.6	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.6	*
13	Respiratory distress of newborn (P22)	6	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	6	0.6	*
...	All other causes (Residual)	275	28.4	59.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	711	100.0	159.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	27.8	44.3
2	Sudden infant death syndrome (R95)	102	14.3	22.8
3	Accidents (unintentional injuries) (V01-X59)	47	6.6	10.5
4	Diseases of the circulatory system (I00-I99)	31	4.4	6.9
5	Influenza and pneumonia (J10-J18)	23	3.2	5.1
6	Septicemia (A40-A41)	21	3.0	4.7
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	21	3.0	4.7
8	Assault (homicide) (*U01,X85-Y09)	20	2.8	4.5
9	Chronic respiratory disease originating in the perinatal period (P27)	10	1.4	*
10	Malignant neoplasms (C00-C97)	9	1.3	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	8	1.1	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	1.1	*
11	Respiratory distress of newborn (P22)	8	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	5	0.7	*
15	Meningitis (G00,G03)	5	0.7	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	5	0.7	*
15	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	5	0.7	*
...	All other causes (Residual)	179	25.2	40.0
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,317	100.0	185.9
1	Sudden infant death syndrome (R95)	1,001	23.2	43.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	796	18.4	34.3
3	Accidents (unintentional injuries) (V01-X59)	435	10.1	18.7
4	Diseases of the circulatory system (I00-I99)	171	4.0	7.4
5	Assault (homicide) (*U01,X85-Y09)	131	3.0	5.6
6	Influenza and pneumonia (J10-J18)	130	3.0	5.6
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	101	2.3	4.3
8	Septicemia (A40-A41)	96	2.2	4.1
9	Chronic respiratory disease originating in the perinatal period (P27)	93	2.2	4.0
10	Malignant neoplasms (C00-C97)	44	1.0	1.9
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	27	0.6	1.2
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	27	0.6	1.2
13	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	26	0.6	1.1
14	Meningitis (G00,G03)	24	0.6	1.0
14	Acute bronchitis and acute bronchiolitis (J20-J21)	24	0.6	1.0
...	All other causes (Residual)	1,191	27.6	51.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
	Postneonates, white, non-Hispanic, male			
...	All causes	2,515	100.0	211.1
1	Sudden infant death syndrome (R95)	628	25.0	52.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	423	16.8	35.5
3	Accidents (unintentional injuries) (V01-X59)	244	9.7	20.5
4	Diseases of the circulatory system (I00-I99)	89	3.5	7.5
5	Assault (homicide) (*U01,X85-Y09)	79	3.1	6.6
6	Influenza and pneumonia (J10-J18)	78	3.1	6.5
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	69	2.7	5.8
8	Chronic respiratory disease originating in the perinatal period (P27)	56	2.2	4.7
9	Septicemia (A40-A41)	46	1.8	3.9
10	Malignant neoplasms (C00-C97)	21	0.8	1.8
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	16	0.6	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	15	0.6	*
13	Meningitis (G00,G03)	14	0.6	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00- D48)	12	0.5	*
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	12	0.5	*
...	All other causes (Residual)	699	27.8	58.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, female				
...	All causes	1,802	100.0	159.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	373	20.7	33.0
1	Sudden infant death syndrome (R95)	373	20.7	33.0
3	Accidents (unintentional injuries) (V01-X59)	191	10.6	16.9
4	Diseases of the circulatory system (I00-I99)	82	4.6	7.3
5	Influenza and pneumonia (J10-J18)	52	2.9	4.6
5	Assault (homicide) (*U01,X85-Y09)	52	2.9	4.6
7	Septicemia (A40-A41)	50	2.8	4.4
8	Chronic respiratory disease originating in the perinatal period (P27)	37	2.1	3.3
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	32	1.8	2.8
10	Malignant neoplasms (C00-C97)	23	1.3	2.0
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	12	0.7	*
12	Respiratory distress of newborn (P22)	12	0.7	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	11	0.6	*
15	Meningitis (G00,G03)	10	0.6	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.6	*
15	Necrotizing enterocolitis of newborn (P77)	10	0.6	*
...	All other causes (Residual)	456	25.3	40.3
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,707	100.0	469.9
1	Sudden infant death syndrome (R95)	620	22.9	107.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	342	12.6	59.4
3	Accidents (unintentional injuries) (V01-X59)	247	9.1	42.9
4	Diseases of the circulatory system (I00-I99)	135	5.0	23.4
5	Septicemia (A40-A41)	119	4.4	20.7
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	116	4.3	20.1
7	Assault (homicide) (*U01,X85-Y09)	94	3.5	16.3
8	Chronic respiratory disease originating in the perinatal period (P27)	87	3.2	15.1
9	Influenza and pneumonia (J10-J18)	83	3.1	14.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	1.5	6.9
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	32	1.2	5.6
12	Anoxic brain damage, not elsewhere classified (G93.1)	21	0.8	3.6
13	Meningitis (G00,G03)	20	0.7	3.5
14	Respiratory distress of newborn (P22)	19	0.7	*
15	Necrotizing enterocolitis of newborn (P77)	16	0.6	*
...	All other causes (Residual)	716	26.4	124.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2003

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, male				
...	All causes	1,533	100.0	523.1
1	Sudden infant death syndrome (R95)	327	21.3	111.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	12.6	65.9
3	Accidents (unintentional injuries) (V01-X59)	139	9.1	47.4
4	Diseases of the circulatory system (I00-I99)	80	5.2	27.3
5	Septicemia (A40-A41)	70	4.6	23.9
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	63	4.1	21.5
7	Chronic respiratory disease originating in the perinatal period (P27)	58	3.8	19.8
8	Assault (homicide) (*U01,X85-Y09)	55	3.6	18.8
9	Influenza and pneumonia (J10-J18)	47	3.1	16.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	1.4	7.2
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	20	1.3	6.8
12	Anoxic brain damage, not elsewhere classified (G93.1)	11	0.7	*
13	Necrotizing enterocolitis of newborn (P77)	9	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	8	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	0.5	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.5	*
14	Respiratory distress of newborn (P22)	8	0.5	*
...	All other causes (Residual)	408	26.6	139.2
Postneonates, black, non-Hispanic, female				
...	All causes	1,174	100.0	414.9
1	Sudden infant death syndrome (R95)	293	25.0	103.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	149	12.7	52.7
3	Accidents (unintentional injuries) (V01-X59)	108	9.2	38.2
4	Diseases of the circulatory system (I00-I99)	55	4.7	19.4
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	53	4.5	18.7
6	Septicemia (A40-A41)	49	4.2	17.3
7	Assault (homicide) (*U01,X85-Y09)	39	3.3	13.8
8	Influenza and pneumonia (J10-J18)	36	3.1	12.7
9	Chronic respiratory disease originating in the perinatal period (P27)	29	2.5	10.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	19	1.6	*
11	Meningitis (G00,G03)	16	1.4	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	12	1.0	*
13	Respiratory distress of newborn (P22)	11	0.9	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	10	0.9	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	7	0.6	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.6	*
15	Necrotizing enterocolitis of newborn (P77)	7	0.6	*
...	All other causes (Residual)	274	23.3	96.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable