

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,724	100.0	580.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,382	24.1	140.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	890	15.5	90.3
3	Newborn affected by maternal complications of pregnancy (P01)	314	5.5	31.9
4	Sudden infant death syndrome (R95)	303	5.3	30.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	221	3.9	22.4
6	Accidents (unintentional injuries) (V01-X59)	172	3.0	17.5
7	Bacterial sepsis of newborn (P36)	171	3.0	17.4
8	Respiratory distress of newborn (P22)	166	2.9	16.8
9	Neonatal hemorrhage (P50-P52,P54)	139	2.4	14.1
10	Necrotizing enterocolitis of newborn (P77)	105	1.8	10.7
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	101	1.8	10.2
12	Diseases of the circulatory system (I00-I99)	100	1.7	10.1
13	Atelectasis (P28.0-P28.1)	82	1.4	8.3
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	76	1.3	7.7
15	Septicemia (A40-A41)	66	1.2	6.7
15	Influenza and pneumonia (J10-J18)	66	1.2	6.7
...	All other causes (Residual)	1,370	23.9	139.0
Infants, Hispanic, male				
...	All causes	3,192	100.0	634.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	720	22.6	143.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	497	15.6	98.7
3	Sudden infant death syndrome (R95)	184	5.8	36.5
4	Newborn affected by maternal complications of pregnancy (P01)	166	5.2	33.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	120	3.8	23.8
6	Respiratory distress of newborn (P22)	107	3.4	21.3
7	Bacterial sepsis of newborn (P36)	95	3.0	18.9
8	Accidents (unintentional injuries) (V01-X59)	87	2.7	17.3
9	Neonatal hemorrhage (P50-P52,P54)	85	2.7	16.9
10	Necrotizing enterocolitis of newborn (P77)	66	2.1	13.1
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	60	1.9	11.9
12	Diseases of the circulatory system (I00-I99)	54	1.7	10.7
12	Atelectasis (P28.0-P28.1)	54	1.7	10.7
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	48	1.5	9.5
15	Septicemia (A40-A41)	37	1.2	7.3
15	Influenza and pneumonia (J10-J18)	37	1.2	7.3
...	All other causes (Residual)	775	24.3	153.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,532	100.0	525.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	662	26.1	137.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	393	15.5	81.5
3	Newborn affected by maternal complications of pregnancy (P01)	148	5.8	30.7
4	Sudden infant death syndrome (R95)	119	4.7	24.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	101	4.0	21.0
6	Accidents (unintentional injuries) (V01-X59)	85	3.4	17.6
7	Bacterial sepsis of newborn (P36)	76	3.0	15.8
8	Respiratory distress of newborn (P22)	59	2.3	12.2
9	Neonatal hemorrhage (P50-P52,P54)	54	2.1	11.2
10	Diseases of the circulatory system (I00-I99)	46	1.8	9.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	41	1.6	8.5
12	Necrotizing enterocolitis of newborn (P77)	39	1.5	8.1
13	Assault (homicide) (*U01,X85-Y09)	30	1.2	6.2
14	Septicemia (A40-A41)	29	1.1	6.0
14	Influenza and pneumonia (J10-J18)	29	1.1	6.0
...	All other causes (Residual)	621	24.5	128.8
Infants, white, non-Hispanic, both sexes				
...	All causes	13,026	100.0	571.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,848	21.9	124.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,783	13.7	78.2
3	Sudden infant death syndrome (R95)	1,198	9.2	52.5
4	Newborn affected by maternal complications of pregnancy (P01)	750	5.8	32.9
5	Accidents (unintentional injuries) (V01-X59)	567	4.4	24.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	546	4.2	23.9
7	Respiratory distress of newborn (P22)	383	2.9	16.8
8	Bacterial sepsis of newborn (P36)	355	2.7	15.6
9	Neonatal hemorrhage (P50-P52,P54)	333	2.6	14.6
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	290	2.2	12.7
11	Diseases of the circulatory system (I00-I99)	246	1.9	10.8
12	Necrotizing enterocolitis of newborn (P77)	214	1.6	9.4
13	Atelectasis (P28.0-P28.1)	185	1.4	8.1
14	Assault (homicide) (*U01,X85-Y09)	139	1.1	6.1
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	120	0.9	5.3
...	All other causes (Residual)	3,069	23.6	134.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,418	100.0	633.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,540	20.8	131.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	985	13.3	84.2
3	Sudden infant death syndrome (R95)	779	10.5	66.6
4	Newborn affected by maternal complications of pregnancy (P01)	398	5.4	34.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	306	4.1	26.1
6	Accidents (unintentional injuries) (V01-X59)	303	4.1	25.9
7	Respiratory distress of newborn (P22)	236	3.2	20.2
8	Bacterial sepsis of newborn (P36)	215	2.9	18.4
9	Neonatal hemorrhage (P50-P52,P54)	195	2.6	16.7
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	166	2.2	14.2
11	Diseases of the circulatory system (I00-I99)	138	1.9	11.8
12	Necrotizing enterocolitis of newborn (P77)	123	1.7	10.5
13	Atelectasis (P28.0-P28.1)	103	1.4	8.8
14	Assault (homicide) (*U01,X85-Y09)	82	1.1	7.0
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	69	0.9	5.9
...	All other causes (Residual)	1,780	24.0	152.1
Infants, white, non-Hispanic, female				
...	All causes	5,608	100.0	505.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,308	23.3	117.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	798	14.2	71.9
3	Sudden infant death syndrome (R95)	419	7.5	37.8
4	Newborn affected by maternal complications of pregnancy (P01)	352	6.3	31.7
5	Accidents (unintentional injuries) (V01-X59)	264	4.7	23.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	240	4.3	21.6
7	Respiratory distress of newborn (P22)	147	2.6	13.3
8	Bacterial sepsis of newborn (P36)	140	2.5	12.6
9	Neonatal hemorrhage (P50-P52,P54)	138	2.5	12.4
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	124	2.2	11.2
11	Diseases of the circulatory system (I00-I99)	108	1.9	9.7
12	Necrotizing enterocolitis of newborn (P77)	91	1.6	8.2
13	Atelectasis (P28.0-P28.1)	82	1.5	7.4
14	Assault (homicide) (*U01,X85-Y09)	57	1.0	5.1
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	51	0.9	4.6
...	All other causes (Residual)	1,289	23.0	116.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,335	100.0	1,427.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,805	21.7	309.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,029	12.3	176.3
3	Sudden infant death syndrome (R95)	650	7.8	111.3
4	Newborn affected by maternal complications of pregnancy (P01)	637	7.6	109.1
5	Accidents (unintentional injuries) (V01-X59)	306	3.7	52.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	305	3.7	52.2
7	Respiratory distress of newborn (P22)	278	3.3	47.6
8	Bacterial sepsis of newborn (P36)	268	3.2	45.9
9	Necrotizing enterocolitis of newborn (P77)	194	2.3	33.2
10	Neonatal hemorrhage (P50-P52,P54)	162	1.9	27.8
11	Diseases of the circulatory system (I00-I99)	152	1.8	26.0
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	130	1.6	22.3
13	Septicemia (A40-A41)	125	1.5	21.4
14	Chronic respiratory disease originating in the perinatal period (P27)	124	1.5	21.2
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	118	1.4	20.2
...	All other causes (Residual)	2,052	24.6	351.5
Infants, black, non-Hispanic, male				
...	All causes	4,665	100.0	1,574.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,033	22.1	348.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	537	11.5	181.3
3	Sudden infant death syndrome (R95)	361	7.7	121.9
4	Newborn affected by maternal complications of pregnancy (P01)	338	7.2	114.1
5	Accidents (unintentional injuries) (V01-X59)	168	3.6	56.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	166	3.6	56.0
7	Respiratory distress of newborn (P22)	158	3.4	53.3
8	Bacterial sepsis of newborn (P36)	147	3.2	49.6
9	Necrotizing enterocolitis of newborn (P77)	109	2.3	36.8
10	Neonatal hemorrhage (P50-P52,P54)	100	2.1	33.8
11	Diseases of the circulatory system (I00-I99)	88	1.9	29.7
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	87	1.9	29.4
13	Chronic respiratory disease originating in the perinatal period (P27)	74	1.6	25.0
14	Septicemia (A40-A41)	65	1.4	21.9
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	62	1.3	20.9
...	All other causes (Residual)	1,172	25.1	395.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,670	100.0	1,276.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	772	21.0	268.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	492	13.4	171.1
3	Newborn affected by maternal complications of pregnancy (P01)	299	8.1	104.0
4	Sudden infant death syndrome (R95)	289	7.9	100.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	139	3.8	48.3
6	Accidents (unintentional injuries) (V01-X59)	138	3.8	48.0
7	Bacterial sepsis of newborn (P36)	121	3.3	42.1
8	Respiratory distress of newborn (P22)	120	3.3	41.7
9	Necrotizing enterocolitis of newborn (P77)	85	2.3	29.6
10	Diseases of the circulatory system (I00-I99)	64	1.7	22.3
11	Neonatal hemorrhage (P50-P52,P54)	62	1.7	21.6
12	Septicemia (A40-A41)	60	1.6	20.9
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	56	1.5	19.5
14	Influenza and pneumonia (J10-J18)	50	1.4	17.4
14	Chronic respiratory disease originating in the perinatal period (P27)	50	1.4	17.4
...	All other causes (Residual)	873	23.8	303.6
Neonates, Hispanic, both sexes				
...	All causes	3,859	100.0	391.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	971	25.2	98.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	869	22.5	88.2
3	Newborn affected by maternal complications of pregnancy (P01)	311	8.1	31.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	218	5.6	22.1
5	Respiratory distress of newborn (P22)	164	4.2	16.6
6	Bacterial sepsis of newborn (P36)	161	4.2	16.3
7	Neonatal hemorrhage (P50-P52,P54)	134	3.5	13.6
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	95	2.5	9.6
9	Necrotizing enterocolitis of newborn (P77)	87	2.3	8.8
10	Atelectasis (P28.0-P28.1)	82	2.1	8.3
11	Hydrops fetalis not due to hemolytic disease (P83.2)	37	1.0	3.8
12	Pulmonary hemorrhage originating in the perinatal period (P26)	34	0.9	3.5
13	Sudden infant death syndrome (R95)	28	0.7	2.8
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	26	0.7	2.6
15	Hematological disorders (P60-P61)	25	0.6	2.5
...	All other causes (Residual)	617	16.0	62.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	2,162	100.0	429.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	519	24.0	103.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	487	22.5	96.7
3	Newborn affected by maternal complications of pregnancy (P01)	164	7.6	32.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	118	5.5	23.4
5	Respiratory distress of newborn (P22)	106	4.9	21.1
6	Bacterial sepsis of newborn (P36)	88	4.1	17.5
7	Neonatal hemorrhage (P50-P52,P54)	82	3.8	16.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	57	2.6	11.3
9	Atelectasis (P28.0-P28.1)	54	2.5	10.7
10	Necrotizing enterocolitis of newborn (P77)	51	2.4	10.1
11	Hydrops fetalis not due to hemolytic disease (P83.2)	21	1.0	4.2
12	Pulmonary hemorrhage originating in the perinatal period (P26)	20	0.9	4.0
13	Hematological disorders (P60-P61)	19	0.9	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	17	0.8	*
15	Sudden infant death syndrome (R95)	15	0.7	*
...	All other causes (Residual)	344	15.9	68.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, female				
...	All causes	1,697	100.0	352.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	452	26.6	93.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	382	22.5	79.2
3	Newborn affected by maternal complications of pregnancy (P01)	147	8.7	30.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	100	5.9	20.7
5	Bacterial sepsis of newborn (P36)	73	4.3	15.1
6	Respiratory distress of newborn (P22)	58	3.4	12.0
7	Neonatal hemorrhage (P50-P52,P54)	52	3.1	10.8
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	2.2	7.9
9	Necrotizing enterocolitis of newborn (P77)	36	2.1	7.5
10	Atelectasis (P28.0-P28.1)	28	1.6	5.8
11	Hydrops fetalis not due to hemolytic disease (P83.2)	16	0.9	*
12	Pulmonary hemorrhage originating in the perinatal period (P26)	14	0.8	*
13	Sudden infant death syndrome (R95)	13	0.8	*
14	Diseases of the circulatory system (I00-I99)	9	0.5	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	0.5	*
14	Newborn affected by other complications of labor and delivery (P03)	9	0.5	*
14	Congenital pneumonia (P23)	9	0.5	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	9	0.5	*
...	All other causes (Residual)	243	14.3	50.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	8,515	100.0	373.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,035	23.9	89.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,746	20.5	76.6
3	Newborn affected by maternal complications of pregnancy (P01)	739	8.7	32.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	538	6.3	23.6
5	Respiratory distress of newborn (P22)	370	4.3	16.2
6	Bacterial sepsis of newborn (P36)	339	4.0	14.9
7	Neonatal hemorrhage (P50-P52,P54)	327	3.8	14.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	279	3.3	12.2
9	Necrotizing enterocolitis of newborn (P77)	191	2.2	8.4
10	Atelectasis (P28.0-P28.1)	180	2.1	7.9
11	Sudden infant death syndrome (R95)	118	1.4	5.2
12	Hydrops fetalis not due to hemolytic disease (P83.2)	90	1.1	3.9
13	Pulmonary hemorrhage originating in the perinatal period (P26)	76	0.9	3.3
14	Accidents (unintentional injuries) (V01-X59)	68	0.8	3.0
15	Newborn affected by other complications of labor and delivery (P03)	60	0.7	2.6
...	All other causes (Residual)	1,359	16.0	59.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, male				
...	All causes	4,721	100.0	403.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,095	23.2	93.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	964	20.4	82.4
3	Newborn affected by maternal complications of pregnancy (P01)	392	8.3	33.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	300	6.4	25.6
5	Respiratory distress of newborn (P22)	229	4.9	19.6
6	Bacterial sepsis of newborn (P36)	204	4.3	17.4
7	Neonatal hemorrhage (P50-P52,P54)	191	4.0	16.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	159	3.4	13.6
9	Necrotizing enterocolitis of newborn (P77)	108	2.3	9.2
10	Atelectasis (P28.0-P28.1)	100	2.1	8.5
11	Sudden infant death syndrome (R95)	70	1.5	6.0
12	Pulmonary hemorrhage originating in the perinatal period (P26)	53	1.1	4.5
13	Hydrops fetalis not due to hemolytic disease (P83.2)	48	1.0	4.1
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	32	0.7	2.7
15	Diseases of the circulatory system (I00-I99)	29	0.6	2.5
15	Newborn affected by other complications of labor and delivery (P03)	29	0.6	2.5
15	Hematological disorders (P60-P61)	29	0.6	2.5
15	Accidents (unintentional injuries) (V01-X59)	29	0.6	2.5
...	All other causes (Residual)	660	14.0	56.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,794	100.0	342.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	940	24.8	84.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	782	20.6	70.5
3	Newborn affected by maternal complications of pregnancy (P01)	347	9.1	31.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	238	6.3	21.5
5	Respiratory distress of newborn (P22)	141	3.7	12.7
6	Neonatal hemorrhage (P50-P52,P54)	136	3.6	12.3
7	Bacterial sepsis of newborn (P36)	135	3.6	12.2
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	120	3.2	10.8
9	Necrotizing enterocolitis of newborn (P77)	83	2.2	7.5
10	Atelectasis (P28.0-P28.1)	80	2.1	7.2
11	Sudden infant death syndrome (R95)	48	1.3	4.3
12	Hydrops fetalis not due to hemolytic disease (P83.2)	42	1.1	3.8
13	Accidents (unintentional injuries) (V01-X59)	39	1.0	3.5
14	Newborn affected by other complications of labor and delivery (P03)	31	0.8	2.8
15	Diseases of the circulatory system (I00-I99)	27	0.7	2.4
...	All other causes (Residual)	605	15.9	54.5
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,488	100.0	940.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,751	31.9	300.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	681	12.4	116.7
3	Newborn affected by maternal complications of pregnancy (P01)	630	11.5	107.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	296	5.4	50.7
5	Respiratory distress of newborn (P22)	266	4.8	45.6
6	Bacterial sepsis of newborn (P36)	250	4.6	42.8
7	Necrotizing enterocolitis of newborn (P77)	162	3.0	27.8
8	Neonatal hemorrhage (P50-P52,P54)	160	2.9	27.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	107	1.9	18.3
10	Atelectasis (P28.0-P28.1)	93	1.7	15.9
11	Pulmonary hemorrhage originating in the perinatal period (P26)	62	1.1	10.6
12	Sudden infant death syndrome (R95)	51	0.9	8.7
13	Newborn affected by other complications of labor and delivery (P03)	44	0.8	7.5
14	Newborn affected by maternal hypertensive disorders (P00.0)	36	0.7	6.2
15	Accidents (unintentional injuries) (V01-X59)	35	0.6	6.0
...	All other causes (Residual)	864	15.7	148.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,061	100.0	1,033.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,002	32.7	338.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	365	11.9	123.2
3	Newborn affected by maternal complications of pregnancy (P01)	333	10.9	112.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	160	5.2	54.0
5	Respiratory distress of newborn (P22)	155	5.1	52.3
6	Bacterial sepsis of newborn (P36)	133	4.3	44.9
7	Neonatal hemorrhage (P50-P52,P54)	98	3.2	33.1
8	Necrotizing enterocolitis of newborn (P77)	91	3.0	30.7
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	57	1.9	19.2
10	Atelectasis (P28.0-P28.1)	53	1.7	17.9
11	Pulmonary hemorrhage originating in the perinatal period (P26)	36	1.2	12.2
12	Sudden infant death syndrome (R95)	27	0.9	9.1
13	Newborn affected by other complications of labor and delivery (P03)	24	0.8	8.1
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	20	0.7	6.8
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.6	*
...	All other causes (Residual)	488	15.9	164.7
Neonates, black, non-Hispanic, female				
...	All causes	2,427	100.0	844.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	749	30.9	260.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	316	13.0	109.9
3	Newborn affected by maternal complications of pregnancy (P01)	297	12.2	103.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	136	5.6	47.3
5	Bacterial sepsis of newborn (P36)	117	4.8	40.7
6	Respiratory distress of newborn (P22)	111	4.6	38.6
7	Necrotizing enterocolitis of newborn (P77)	71	2.9	24.7
8	Neonatal hemorrhage (P50-P52,P54)	62	2.6	21.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	50	2.1	17.4
10	Atelectasis (P28.0-P28.1)	40	1.6	13.9
11	Pulmonary hemorrhage originating in the perinatal period (P26)	26	1.1	9.0
12	Newborn affected by maternal hypertensive disorders (P00.0)	25	1.0	8.7
13	Sudden infant death syndrome (R95)	24	1.0	8.3
14	Newborn affected by other complications of labor and delivery (P03)	20	0.8	7.0
14	Accidents (unintentional injuries) (V01-X59)	20	0.8	7.0
...	All other causes (Residual)	363	15.0	126.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,865	100.0	189.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	411	22.0	41.7
2	Sudden infant death syndrome (R95)	275	14.7	27.9
3	Accidents (unintentional injuries) (V01-X59)	156	8.4	15.8
4	Diseases of the circulatory system (I00-I99)	80	4.3	8.1
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	71	3.8	7.2
6	Septicemia (A40-A41)	66	3.5	6.7
7	Influenza and pneumonia (J10-J18)	65	3.5	6.6
8	Assault (homicide) (*U01,X85-Y09)	52	2.8	5.3
9	Chronic respiratory disease originating in the perinatal period (P27)	32	1.7	3.2
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	21	1.1	2.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	1.1	2.1
12	Necrotizing enterocolitis of newborn (P77)	18	1.0	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	15	0.8	*
14	Malignant neoplasms (C00-C97)	14	0.8	*
15	Bacterial sepsis of newborn (P36)	10	0.5	*
...	All other causes (Residual)	558	29.9	56.6
Postneonates, Hispanic, male				
...	All causes	1,030	100.0	204.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	19.5	39.9
2	Sudden infant death syndrome (R95)	169	16.4	33.6
3	Accidents (unintentional injuries) (V01-X59)	77	7.5	15.3
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	44	4.3	8.7
5	Diseases of the circulatory system (I00-I99)	43	4.2	8.5
6	Septicemia (A40-A41)	37	3.6	7.3
7	Influenza and pneumonia (J10-J18)	36	3.5	7.2
8	Assault (homicide) (*U01,X85-Y09)	26	2.5	5.2
9	Chronic respiratory disease originating in the perinatal period (P27)	17	1.7	*
10	Necrotizing enterocolitis of newborn (P77)	15	1.5	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	1.0	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	8	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.8	*
14	Bacterial sepsis of newborn (P36)	7	0.7	*
15	Whooping cough (A37)	6	0.6	*
...	All other causes (Residual)	326	31.7	64.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	835	100.0	173.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	25.1	43.6
2	Sudden infant death syndrome (R95)	106	12.7	22.0
3	Accidents (unintentional injuries) (V01-X59)	79	9.5	16.4
4	Diseases of the circulatory system (I00-I99)	37	4.4	7.7
5	Septicemia (A40-A41)	29	3.5	6.0
5	Influenza and pneumonia (J10-J18)	29	3.5	6.0
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	27	3.2	5.6
8	Assault (homicide) (*U01,X85-Y09)	26	3.1	5.4
9	Chronic respiratory disease originating in the perinatal period (P27)	15	1.8	*
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	13	1.6	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	11	1.3	*
12	Malignant neoplasms (C00-C97)	9	1.1	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	7	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	4	0.5	*
15	Meningitis (G00,G03)	4	0.5	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	4	0.5	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	4	0.5	*
15	Congenital pneumonia (P23)	4	0.5	*
...	All other causes (Residual)	212	25.4	44.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,511	100.0	197.9
1	Sudden infant death syndrome (R95)	1,080	23.9	47.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	813	18.0	35.7
3	Accidents (unintentional injuries) (V01-X59)	499	11.1	21.9
4	Diseases of the circulatory system (I00-I99)	190	4.2	8.3
5	Assault (homicide) (*U01,X85-Y09)	122	2.7	5.4
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	110	2.4	4.8
7	Septicemia (A40-A41)	100	2.2	4.4
8	Influenza and pneumonia (J10-J18)	93	2.1	4.1
9	Chronic respiratory disease originating in the perinatal period (P27)	92	2.0	4.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	37	0.8	1.6
11	Malignant neoplasms (C00-C97)	36	0.8	1.6
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	29	0.6	1.3
13	Necrotizing enterocolitis of newborn (P77)	23	0.5	1.0
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	22	0.5	1.0
14	Acute bronchitis and acute bronchiolitis (J20-J21)	22	0.5	1.0
...	All other causes (Residual)	1,243	27.6	54.5
Postneonates, white, non-Hispanic, male				
...	All causes	2,697	100.0	230.4
1	Sudden infant death syndrome (R95)	709	26.3	60.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	445	16.5	38.0
3	Accidents (unintentional injuries) (V01-X59)	274	10.2	23.4
4	Diseases of the circulatory system (I00-I99)	109	4.0	9.3
5	Assault (homicide) (*U01,X85-Y09)	74	2.7	6.3
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	64	2.4	5.5
7	Septicemia (A40-A41)	63	2.3	5.4
8	Influenza and pneumonia (J10-J18)	60	2.2	5.1
9	Chronic respiratory disease originating in the perinatal period (P27)	52	1.9	4.4
10	Malignant neoplasms (C00-C97)	26	1.0	2.2
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	0.8	1.8
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	17	0.6	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	15	0.6	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	15	0.6	*
13	Necrotizing enterocolitis of newborn (P77)	15	0.6	*
...	All other causes (Residual)	738	27.4	63.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, female				
...	All causes	1,814	100.0	163.5
1	Sudden infant death syndrome (R95)	371	20.5	33.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	368	20.3	33.2
3	Accidents (unintentional injuries) (V01-X59)	225	12.4	20.3
4	Diseases of the circulatory system (I00-I99)	81	4.5	7.3
5	Assault (homicide) (*U01,X85-Y09)	48	2.6	4.3
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	46	2.5	4.1
7	Chronic respiratory disease originating in the perinatal period (P27)	40	2.2	3.6
8	Septicemia (A40-A41)	37	2.0	3.3
9	Influenza and pneumonia (J10-J18)	33	1.8	3.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	0.9	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	12	0.7	*
12	Malignant neoplasms (C00-C97)	10	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Necrotizing enterocolitis of newborn (P77)	8	0.4	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	7	0.4	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	7	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.4	*
...	All other causes (Residual)	489	27.0	44.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,847	100.0	487.7
1	Sudden infant death syndrome (R95)	599	21.0	102.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	348	12.2	59.6
3	Accidents (unintentional injuries) (V01-X59)	271	9.5	46.4
4	Septicemia (A40-A41)	125	4.4	21.4
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	122	4.3	20.9
6	Diseases of the circulatory system (I00-I99)	121	4.3	20.7
7	Chronic respiratory disease originating in the perinatal period (P27)	108	3.8	18.5
8	Influenza and pneumonia (J10-J18)	95	3.3	16.3
9	Assault (homicide) (*U01,X85-Y09)	85	3.0	14.6
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	54	1.9	9.3
11	Necrotizing enterocolitis of newborn (P77)	32	1.1	5.5
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	28	1.0	4.8
13	Bacterial sepsis of newborn (P36)	18	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	17	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	17	0.6	*
...	All other causes (Residual)	807	28.3	138.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2005

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, male				
...	All causes	1,604	100.0	541.5
1	Sudden infant death syndrome (R95)	334	20.8	112.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	10.7	58.1
3	Accidents (unintentional injuries) (V01-X59)	153	9.5	51.6
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	80	5.0	27.0
5	Diseases of the circulatory system (I00-I99)	70	4.4	23.6
6	Septicemia (A40-A41)	65	4.1	21.9
7	Chronic respiratory disease originating in the perinatal period (P27)	64	4.0	21.6
8	Assault (homicide) (*U01,X85-Y09)	49	3.1	16.5
9	Influenza and pneumonia (J10-J18)	45	2.8	15.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	1.9	10.5
11	Necrotizing enterocolitis of newborn (P77)	18	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	17	1.1	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	15	0.9	*
14	Bacterial sepsis of newborn (P36)	14	0.9	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.6	*
...	All other causes (Residual)	468	29.2	158.0
Postneonates, black, non-Hispanic, female				
...	All causes	1,243	100.0	432.3
1	Sudden infant death syndrome (R95)	265	21.3	92.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	14.2	61.2
3	Accidents (unintentional injuries) (V01-X59)	118	9.5	41.0
4	Septicemia (A40-A41)	60	4.8	20.9
5	Diseases of the circulatory system (I00-I99)	51	4.1	17.7
6	Influenza and pneumonia (J10-J18)	50	4.0	17.4
7	Chronic respiratory disease originating in the perinatal period (P27)	44	3.5	15.3
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	42	3.4	14.6
9	Assault (homicide) (*U01,X85-Y09)	36	2.9	12.5
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	1.9	8.0
11	Necrotizing enterocolitis of newborn (P77)	14	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	11	0.9	*
13	Respiratory distress of newborn (P22)	9	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.6	*
15	Meningitis (G00,G03)	7	0.6	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	7	0.6	*
...	All other causes (Residual)	322	25.9	112.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable