

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,890	100.0	565.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,502	25.5	144.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	939	15.9	90.2
3	Newborn affected by maternal complications of pregnancy (P01)	366	6.2	35.2
4	Sudden infant death syndrome (R95)	331	5.6	31.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	237	4.0	22.8
6	Accidents (unintentional injuries) (V01-X59)	184	3.1	17.7
7	Bacterial sepsis of newborn (P36)	138	2.3	13.3
8	Respiratory distress of newborn (P22)	137	2.3	13.2
9	Diseases of the circulatory system (I00-I99)	132	2.2	12.7
10	Neonatal hemorrhage (P50-P52,P54)	130	2.2	12.5
11	Necrotizing enterocolitis of newborn (P77)	109	1.9	10.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.5	8.3
13	Atelectasis (P28.0-P28.1)	70	1.2	6.7
14	Septicemia (A40-A41)	69	1.2	6.6
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	69	1.2	6.6
...	All other causes (Residual)	1,391	23.6	133.6
Infants, Hispanic, male				
...	All causes	3,279	100.0	616.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	796	24.3	149.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	545	16.6	102.4
3	Newborn affected by maternal complications of pregnancy (P01)	212	6.5	39.8
4	Sudden infant death syndrome (R95)	188	5.7	35.3
5	Newborn affected by complications of placenta, cord and membranes (P02)	125	3.8	23.5
6	Accidents (unintentional injuries) (V01-X59)	111	3.4	20.9
7	Respiratory distress of newborn (P22)	87	2.7	16.4
8	Neonatal hemorrhage (P50-P52,P54)	79	2.4	14.8
9	Bacterial sepsis of newborn (P36)	78	2.4	14.7
10	Diseases of the circulatory system (I00-I99)	74	2.3	13.9
11	Necrotizing enterocolitis of newborn (P77)	62	1.9	11.7
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	47	1.4	8.8
13	Atelectasis (P28.0-P28.1)	41	1.3	7.7
14	Assault (homicide) (*U01,X85-Y09)	40	1.2	7.5
15	Septicemia (A40-A41)	37	1.1	7.0
...	All other causes (Residual)	757	23.1	142.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,611	100.0	512.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	706	27.0	138.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	394	15.1	77.4
3	Newborn affected by maternal complications of pregnancy (P01)	154	5.9	30.2
4	Sudden infant death syndrome (R95)	143	5.5	28.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	112	4.3	22.0
6	Accidents (unintentional injuries) (V01-X59)	73	2.8	14.3
7	Bacterial sepsis of newborn (P36)	60	2.3	11.8
8	Diseases of the circulatory system (I00-I99)	58	2.2	11.4
9	Neonatal hemorrhage (P50-P52,P54)	51	2.0	10.0
10	Respiratory distress of newborn (P22)	50	1.9	9.8
11	Necrotizing enterocolitis of newborn (P77)	47	1.8	9.2
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	39	1.5	7.7
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	38	1.5	7.5
14	Septicemia (A40-A41)	32	1.2	6.3
15	Atelectasis (P28.0-P28.1)	29	1.1	5.7
...	All other causes (Residual)	625	23.9	122.7
Infants, Non-Hispanic, white, both sexes				
...	All causes	12,519	100.0	552.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,831	22.6	124.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,744	13.9	76.9
3	Sudden infant death syndrome (R95)	1,197	9.6	52.8
4	Newborn affected by maternal complications of pregnancy (P01)	767	6.1	33.8
5	Accidents (unintentional injuries) (V01-X59)	675	5.4	29.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	472	3.8	20.8
7	Bacterial sepsis of newborn (P36)	300	2.4	13.2
8	Respiratory distress of newborn (P22)	260	2.1	11.5
9	Diseases of the circulatory system (I00-I99)	257	2.1	11.3
10	Neonatal hemorrhage (P50-P52,P54)	245	2.0	10.8
11	Necrotizing enterocolitis of newborn (P77)	206	1.6	9.1
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	173	1.4	7.6
13	Atelectasis (P28.0-P28.1)	157	1.3	6.9
14	Assault (homicide) (*U01,X85-Y09)	144	1.2	6.3
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	128	1.0	5.6
...	All other causes (Residual)	2,963	23.7	130.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	7,022	100.0	604.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,488	21.2	128.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	973	13.9	83.7
3	Sudden infant death syndrome (R95)	723	10.3	62.2
4	Newborn affected by maternal complications of pregnancy (P01)	448	6.4	38.5
5	Accidents (unintentional injuries) (V01-X59)	380	5.4	32.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	252	3.6	21.7
7	Bacterial sepsis of newborn (P36)	170	2.4	14.6
8	Respiratory distress of newborn (P22)	160	2.3	13.8
9	Neonatal hemorrhage (P50-P52,P54)	144	2.1	12.4
10	Diseases of the circulatory system (I00-I99)	135	1.9	11.6
11	Necrotizing enterocolitis of newborn (P77)	123	1.8	10.6
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	109	1.6	9.4
13	Atelectasis (P28.0-P28.1)	87	1.2	7.5
14	Assault (homicide) (*U01,X85-Y09)	83	1.2	7.1
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	70	1.0	6.0
...	All other causes (Residual)	1,677	23.9	144.2
Infants, Non-Hispanic, white, female				
...	All causes	5,497	100.0	497.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,343	24.4	121.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	771	14.0	69.8
3	Sudden infant death syndrome (R95)	474	8.6	42.9
4	Newborn affected by maternal complications of pregnancy (P01)	319	5.8	28.9
5	Accidents (unintentional injuries) (V01-X59)	295	5.4	26.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	220	4.0	19.9
7	Bacterial sepsis of newborn (P36)	130	2.4	11.8
8	Diseases of the circulatory system (I00-I99)	122	2.2	11.0
9	Neonatal hemorrhage (P50-P52,P54)	101	1.8	9.1
10	Respiratory distress of newborn (P22)	100	1.8	9.0
11	Necrotizing enterocolitis of newborn (P77)	83	1.5	7.5
12	Atelectasis (P28.0-P28.1)	70	1.3	6.3
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	64	1.2	5.8
14	Assault (homicide) (*U01,X85-Y09)	61	1.1	5.5
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	58	1.1	5.2
...	All other causes (Residual)	1,286	23.4	116.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	8,187	100.0	1,314.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,800	22.0	288.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,012	12.4	162.4
3	Sudden infant death syndrome (R95)	731	8.9	117.3
4	Newborn affected by maternal complications of pregnancy (P01)	543	6.6	87.2
5	Accidents (unintentional injuries) (V01-X59)	399	4.9	64.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	312	3.8	50.1
7	Bacterial sepsis of newborn (P36)	218	2.7	35.0
8	Necrotizing enterocolitis of newborn (P77)	211	2.6	33.9
9	Respiratory distress of newborn (P22)	210	2.6	33.7
10	Diseases of the circulatory system (I00-I99)	164	2.0	26.3
11	Neonatal hemorrhage (P50-P52,P54)	157	1.9	25.2
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	137	1.7	22.0
13	Assault (homicide) (*U01,X85-Y09)	109	1.3	17.5
14	Septicemia (A40-A41)	104	1.3	16.7
14	Chronic respiratory disease originating in the perinatal period (P27)	104	1.3	16.7
...	All other causes (Residual)	1,976	24.1	317.2
Infants, Non-Hispanic, black, male				
...	All causes	4,546	100.0	1,436.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,012	22.3	319.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	513	11.3	162.1
3	Sudden infant death syndrome (R95)	420	9.2	132.7
4	Newborn affected by maternal complications of pregnancy (P01)	292	6.4	92.3
5	Accidents (unintentional injuries) (V01-X59)	210	4.6	66.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	169	3.7	53.4
7	Bacterial sepsis of newborn (P36)	127	2.8	40.1
8	Necrotizing enterocolitis of newborn (P77)	118	2.6	37.3
9	Respiratory distress of newborn (P22)	117	2.6	37.0
10	Diseases of the circulatory system (I00-I99)	95	2.1	30.0
11	Neonatal hemorrhage (P50-P52,P54)	91	2.0	28.8
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	72	1.6	22.8
13	Atelectasis (P28.0-P28.1)	70	1.5	22.1
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	59	1.3	18.6
14	Assault (homicide) (*U01,X85-Y09)	59	1.3	18.6
...	All other causes (Residual)	1,122	24.7	354.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	3,641	100.0	1,187.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	788	21.6	257.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	499	13.7	162.8
3	Sudden infant death syndrome (R95)	311	8.5	101.4
4	Newborn affected by maternal complications of pregnancy (P01)	251	6.9	81.9
5	Accidents (unintentional injuries) (V01-X59)	189	5.2	61.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	143	3.9	46.6
7	Respiratory distress of newborn (P22)	93	2.6	30.3
7	Necrotizing enterocolitis of newborn (P77)	93	2.6	30.3
9	Bacterial sepsis of newborn (P36)	91	2.5	29.7
10	Diseases of the circulatory system (I00-I99)	69	1.9	22.5
11	Neonatal hemorrhage (P50-P52,P54)	66	1.8	21.5
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	65	1.8	21.2
13	Septicemia (A40-A41)	50	1.4	16.3
13	Assault (homicide) (*U01,X85-Y09)	50	1.4	16.3
15	Chronic respiratory disease originating in the perinatal period (P27)	48	1.3	15.7
...	All other causes (Residual)	835	22.9	272.4
Neonates, Hispanic, both sexes				
...	All causes	3,970	100.0	381.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,064	26.8	102.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	914	23.0	87.8
3	Newborn affected by maternal complications of pregnancy (P01)	359	9.0	34.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	234	5.9	22.5
5	Bacterial sepsis of newborn (P36)	137	3.5	13.2
6	Respiratory distress of newborn (P22)	135	3.4	13.0
7	Neonatal hemorrhage (P50-P52,P54)	126	3.2	12.1
8	Necrotizing enterocolitis of newborn (P77)	94	2.4	9.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	78	2.0	7.5
10	Atelectasis (P28.0-P28.1)	70	1.8	6.7
11	Hydrops fetalis not due to hemolytic disease (P83.2)	45	1.1	4.3
12	Pulmonary hemorrhage originating in the perinatal period (P26)	41	1.0	3.9
13	Sudden infant death syndrome (R95)	29	0.7	2.8
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	26	0.7	2.5
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	23	0.6	2.2
...	All other causes (Residual)	595	15.0	57.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	2,215	100.0	416.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	574	25.9	107.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	527	23.8	99.1
3	Newborn affected by maternal complications of pregnancy (P01)	209	9.4	39.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	123	5.6	23.1
5	Respiratory distress of newborn (P22)	85	3.8	16.0
6	Bacterial sepsis of newborn (P36)	78	3.5	14.7
7	Neonatal hemorrhage (P50-P52,P54)	76	3.4	14.3
8	Necrotizing enterocolitis of newborn (P77)	51	2.3	9.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	44	2.0	8.3
10	Atelectasis (P28.0-P28.1)	41	1.9	7.7
11	Pulmonary hemorrhage originating in the perinatal period (P26)	26	1.2	4.9
12	Hydrops fetalis not due to hemolytic disease (P83.2)	21	0.9	3.9
13	Sudden infant death syndrome (R95)	18	0.8	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	16	0.7	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	11	0.5	*
15	Accidents (unintentional injuries) (V01-X59)	11	0.5	*
...	All other causes (Residual)	304	13.7	57.1
Neonates, Hispanic, female				
...	All causes	1,755	100.0	344.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	490	27.9	96.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	387	22.1	76.0
3	Newborn affected by maternal complications of pregnancy (P01)	150	8.5	29.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	111	6.3	21.8
5	Bacterial sepsis of newborn (P36)	59	3.4	11.6
6	Respiratory distress of newborn (P22)	50	2.8	9.8
6	Neonatal hemorrhage (P50-P52,P54)	50	2.8	9.8
8	Necrotizing enterocolitis of newborn (P77)	43	2.5	8.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	34	1.9	6.7
10	Atelectasis (P28.0-P28.1)	29	1.7	5.7
11	Hydrops fetalis not due to hemolytic disease (P83.2)	24	1.4	4.7
12	Pulmonary hemorrhage originating in the perinatal period (P26)	15	0.9	*
13	Diseases of the circulatory system (I00-I99)	14	0.8	*
14	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	12	0.7	*
15	Sudden infant death syndrome (R95)	11	0.6	*
...	All other causes (Residual)	276	15.7	54.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	8,011	100.0	353.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,045	25.5	90.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,703	21.3	75.1
3	Newborn affected by maternal complications of pregnancy (P01)	757	9.4	33.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	465	5.8	20.5
5	Bacterial sepsis of newborn (P36)	287	3.6	12.7
6	Respiratory distress of newborn (P22)	255	3.2	11.2
7	Neonatal hemorrhage (P50-P52,P54)	240	3.0	10.6
8	Necrotizing enterocolitis of newborn (P77)	181	2.3	8.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	167	2.1	7.4
10	Atelectasis (P28.0-P28.1)	151	1.9	6.7
11	Sudden infant death syndrome (R95)	116	1.4	5.1
12	Hydrops fetalis not due to hemolytic disease (P83.2)	81	1.0	3.6
13	Accidents (unintentional injuries) (V01-X59)	72	0.9	3.2
14	Pulmonary hemorrhage originating in the perinatal period (P26)	69	0.9	3.0
15	Diseases of the circulatory system (I00-I99)	58	0.7	2.6
...	All other causes (Residual)	1,364	17.0	60.1
Neonates, Non-Hispanic, white, male				
...	All causes	4,460	100.0	383.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,075	24.1	92.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	952	21.3	81.9
3	Newborn affected by maternal complications of pregnancy (P01)	442	9.9	38.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	247	5.5	21.2
5	Bacterial sepsis of newborn (P36)	162	3.6	13.9
6	Respiratory distress of newborn (P22)	156	3.5	13.4
7	Neonatal hemorrhage (P50-P52,P54)	142	3.2	12.2
8	Necrotizing enterocolitis of newborn (P77)	112	2.5	9.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	106	2.4	9.1
10	Atelectasis (P28.0-P28.1)	82	1.8	7.1
11	Sudden infant death syndrome (R95)	61	1.4	5.2
12	Hydrops fetalis not due to hemolytic disease (P83.2)	44	1.0	3.8
13	Pulmonary hemorrhage originating in the perinatal period (P26)	41	0.9	3.5
14	Accidents (unintentional injuries) (V01-X59)	40	0.9	3.4
15	Newborn affected by other complications of labor and delivery (P03)	38	0.9	3.3
...	All other causes (Residual)	760	17.0	65.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,551	100.0	321.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	970	27.3	87.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	751	21.1	68.0
3	Newborn affected by maternal complications of pregnancy (P01)	315	8.9	28.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	218	6.1	19.7
5	Bacterial sepsis of newborn (P36)	125	3.5	11.3
6	Respiratory distress of newborn (P22)	99	2.8	9.0
7	Neonatal hemorrhage (P50-P52,P54)	98	2.8	8.9
8	Atelectasis (P28.0-P28.1)	69	1.9	6.2
8	Necrotizing enterocolitis of newborn (P77)	69	1.9	6.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	61	1.7	5.5
11	Sudden infant death syndrome (R95)	55	1.5	5.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	37	1.0	3.3
13	Diseases of the circulatory system (I00-I99)	32	0.9	2.9
13	Accidents (unintentional injuries) (V01-X59)	32	0.9	2.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	28	0.8	2.5
...	All other causes (Residual)	592	16.7	53.6
Neonates, Non-Hispanic, black, both sexes				
...	All causes	5,268	100.0	845.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,747	33.2	280.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	648	12.3	104.0
3	Newborn affected by maternal complications of pregnancy (P01)	538	10.2	86.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	309	5.9	49.6
5	Bacterial sepsis of newborn (P36)	208	3.9	33.4
6	Respiratory distress of newborn (P22)	205	3.9	32.9
7	Necrotizing enterocolitis of newborn (P77)	169	3.2	27.1
8	Neonatal hemorrhage (P50-P52,P54)	150	2.8	24.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	97	1.8	15.6
10	Atelectasis (P28.0-P28.1)	92	1.7	14.8
11	Pulmonary hemorrhage originating in the perinatal period (P26)	76	1.4	12.2
12	Sudden infant death syndrome (R95)	57	1.1	9.1
13	Accidents (unintentional injuries) (V01-X59)	36	0.7	5.8
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	31	0.6	5.0
15	Newborn affected by maternal hypertensive disorders (P00.0)	28	0.5	4.5
...	All other causes (Residual)	877	16.6	140.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	2,925	100.0	924.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	981	33.5	310.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	342	11.7	108.1
3	Newborn affected by maternal complications of pregnancy (P01)	287	9.8	90.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	168	5.7	53.1
5	Bacterial sepsis of newborn (P36)	119	4.1	37.6
6	Respiratory distress of newborn (P22)	115	3.9	36.3
7	Necrotizing enterocolitis of newborn (P77)	98	3.4	31.0
8	Neonatal hemorrhage (P50-P52,P54)	89	3.0	28.1
9	Atelectasis (P28.0-P28.1)	68	2.3	21.5
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	55	1.9	17.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	43	1.5	13.6
12	Sudden infant death syndrome (R95)	29	1.0	9.2
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	19	0.6	*
14	Slow fetal growth and fetal malnutrition (P05)	17	0.6	*
14	Accidents (unintentional injuries) (V01-X59)	17	0.6	*
...	All other causes (Residual)	478	16.3	151.1
Neonates, Non-Hispanic, black, female				
...	All causes	2,343	100.0	764.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	766	32.7	249.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	13.1	99.8
3	Newborn affected by maternal complications of pregnancy (P01)	251	10.7	81.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	141	6.0	46.0
5	Respiratory distress of newborn (P22)	90	3.8	29.4
6	Bacterial sepsis of newborn (P36)	89	3.8	29.0
7	Necrotizing enterocolitis of newborn (P77)	71	3.0	23.2
8	Neonatal hemorrhage (P50-P52,P54)	61	2.6	19.9
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	42	1.8	13.7
10	Pulmonary hemorrhage originating in the perinatal period (P26)	33	1.4	10.8
11	Sudden infant death syndrome (R95)	28	1.2	9.1
12	Atelectasis (P28.0-P28.1)	24	1.0	7.8
13	Accidents (unintentional injuries) (V01-X59)	19	0.8	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	14	0.6	*
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	12	0.5	*
...	All other causes (Residual)	396	16.9	129.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,920	100.0	184.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	438	22.8	42.1
2	Sudden infant death syndrome (R95)	302	15.7	29.0
3	Accidents (unintentional injuries) (V01-X59)	165	8.6	15.8
4	Diseases of the circulatory system (I00-I99)	111	5.8	10.7
5	Septicemia (A40-A41)	69	3.6	6.6
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	67	3.5	6.4
7	Assault (homicide) (*U01,X85-Y09)	64	3.3	6.1
8	Influenza and pneumonia (J09-J18)	46	2.4	4.4
9	Chronic respiratory disease originating in the perinatal period (P27)	32	1.7	3.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	25	1.3	2.4
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.8	*
12	Necrotizing enterocolitis of newborn (P77)	15	0.8	*
14	Malignant neoplasms (C00-C97)	14	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.7	*
...	All other causes (Residual)	525	27.3	50.4
Postneonates, Hispanic, male				
...	All causes	1,064	100.0	200.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	20.9	41.7
2	Sudden infant death syndrome (R95)	170	16.0	32.0
3	Accidents (unintentional injuries) (V01-X59)	100	9.4	18.8
4	Diseases of the circulatory system (I00-I99)	67	6.3	12.6
5	Assault (homicide) (*U01,X85-Y09)	38	3.6	7.1
6	Septicemia (A40-A41)	37	3.5	7.0
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	31	2.9	5.8
8	Influenza and pneumonia (J09-J18)	23	2.2	4.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.7	*
9	Chronic respiratory disease originating in the perinatal period (P27)	18	1.7	*
11	Necrotizing enterocolitis of newborn (P77)	11	1.0	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.8	*
12	Meningitis (G00,G03)	9	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	0.8	*
15	Malignant neoplasms (C00-C97)	8	0.8	*
...	All other causes (Residual)	294	27.6	55.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	856	100.0	168.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	216	25.2	42.4
2	Sudden infant death syndrome (R95)	132	15.4	25.9
3	Accidents (unintentional injuries) (V01-X59)	65	7.6	12.8
4	Diseases of the circulatory system (I00-I99)	44	5.1	8.6
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	36	4.2	7.1
6	Septicemia (A40-A41)	32	3.7	6.3
7	Assault (homicide) (*U01,X85-Y09)	26	3.0	5.1
8	Influenza and pneumonia (J09-J18)	23	2.7	4.5
9	Chronic respiratory disease originating in the perinatal period (P27)	14	1.6	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	10	1.2	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	1.1	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	7	0.8	*
13	Malignant neoplasms (C00-C97)	6	0.7	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	6	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	5	0.6	*
...	All other causes (Residual)	220	25.7	43.2
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	4,508	100.0	198.8
1	Sudden infant death syndrome (R95)	1,081	24.0	47.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	786	17.4	34.7
3	Accidents (unintentional injuries) (V01-X59)	603	13.4	26.6
4	Diseases of the circulatory system (I00-I99)	199	4.4	8.8
5	Assault (homicide) (*U01,X85-Y09)	128	2.8	5.6
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	121	2.7	5.3
7	Influenza and pneumonia (J09-J18)	106	2.4	4.7
8	Septicemia (A40-A41)	102	2.3	4.5
9	Chronic respiratory disease originating in the perinatal period (P27)	75	1.7	3.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	41	0.9	1.8
11	Malignant neoplasms (C00-C97)	29	0.6	1.3
12	Necrotizing enterocolitis of newborn (P77)	25	0.6	1.1
13	Meningitis (G00,G03)	23	0.5	1.0
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	23	0.5	1.0
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	21	0.5	0.9
15	Anoxic brain damage, not elsewhere classified (G93.1)	21	0.5	0.9
...	All other causes (Residual)	1,124	24.9	49.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	2,562	100.0	220.4
1	Sudden infant death syndrome (R95)	662	25.8	56.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	413	16.1	35.5
3	Accidents (unintentional injuries) (V01-X59)	340	13.3	29.2
4	Diseases of the circulatory system (I00-I99)	109	4.3	9.4
5	Assault (homicide) (*U01,X85-Y09)	71	2.8	6.1
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	68	2.7	5.8
7	Influenza and pneumonia (J09-J18)	60	2.3	5.2
8	Chronic respiratory disease originating in the perinatal period (P27)	50	2.0	4.3
9	Septicemia (A40-A41)	48	1.9	4.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	0.8	1.8
11	Malignant neoplasms (C00-C97)	19	0.7	*
12	Meningitis (G00,G03)	13	0.5	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	13	0.5	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	11	0.4	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	11	0.4	*
14	Necrotizing enterocolitis of newborn (P77)	11	0.4	*
...	All other causes (Residual)	642	25.1	55.2
Postneonates, Non-Hispanic, white, female				
...	All causes	1,946	100.0	176.1
1	Sudden infant death syndrome (R95)	419	21.5	37.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	373	19.2	33.7
3	Accidents (unintentional injuries) (V01-X59)	263	13.5	23.8
4	Diseases of the circulatory system (I00-I99)	90	4.6	8.1
5	Assault (homicide) (*U01,X85-Y09)	57	2.9	5.2
6	Septicemia (A40-A41)	54	2.8	4.9
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	53	2.7	4.8
8	Influenza and pneumonia (J09-J18)	46	2.4	4.2
9	Chronic respiratory disease originating in the perinatal period (P27)	25	1.3	2.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	20	1.0	1.8
11	Necrotizing enterocolitis of newborn (P77)	14	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.7	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	12	0.6	*
14	Malignant neoplasms (C00-C97)	10	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.5	*
14	Meningitis (G00,G03)	10	0.5	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.5	*
...	All other causes (Residual)	467	24.0	42.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	2,919	100.0	468.5
1	Sudden infant death syndrome (R95)	674	23.1	108.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	364	12.5	58.4
3	Accidents (unintentional injuries) (V01-X59)	363	12.4	58.3
4	Diseases of the circulatory system (I00-I99)	144	4.9	23.1
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	127	4.4	20.4
6	Septicemia (A40-A41)	103	3.5	16.5
7	Assault (homicide) (*U01,X85-Y09)	100	3.4	16.1
8	Chronic respiratory disease originating in the perinatal period (P27)	96	3.3	15.4
9	Influenza and pneumonia (J09-J18)	61	2.1	9.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	53	1.8	8.5
11	Necrotizing enterocolitis of newborn (P77)	42	1.4	6.7
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	34	1.2	5.5
13	Anoxic brain damage, not elsewhere classified (G93.1)	23	0.8	3.7
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	17	0.6	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.5	*
...	All other causes (Residual)	703	24.1	112.8
Postneonates, Non-Hispanic, black, male				
...	All causes	1,621	100.0	512.3
1	Sudden infant death syndrome (R95)	391	24.1	123.6
2	Accidents (unintentional injuries) (V01-X59)	193	11.9	61.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	10.5	54.0
4	Diseases of the circulatory system (I00-I99)	84	5.2	26.5
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	65	4.0	20.5
6	Septicemia (A40-A41)	54	3.3	17.1
6	Assault (homicide) (*U01,X85-Y09)	54	3.3	17.1
8	Chronic respiratory disease originating in the perinatal period (P27)	52	3.2	16.4
9	Influenza and pneumonia (J09-J18)	33	2.0	10.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	1.9	9.8
11	Necrotizing enterocolitis of newborn (P77)	20	1.2	6.3
12	Anoxic brain damage, not elsewhere classified (G93.1)	15	0.9	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.9	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.6	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	9	0.6	*
...	All other causes (Residual)	426	26.3	134.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2008

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,298	100.0	423.4
1	Sudden infant death syndrome (R95)	283	21.8	92.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	14.9	63.0
3	Accidents (unintentional injuries) (V01-X59)	170	13.1	55.4
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	62	4.8	20.2
5	Diseases of the circulatory system (I00-I99)	60	4.6	19.6
6	Septicemia (A40-A41)	49	3.8	16.0
7	Assault (homicide) (*U01,X85-Y09)	46	3.5	15.0
8	Chronic respiratory disease originating in the perinatal period (P27)	44	3.4	14.4
9	Influenza and pneumonia (J09-J18)	28	2.2	9.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	1.7	7.2
10	Necrotizing enterocolitis of newborn (P77)	22	1.7	7.2
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	20	1.5	6.5
13	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	0.6	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	8	0.6	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	6	0.5	*
15	Meningitis (G00,G03)	6	0.5	*
...	All other causes (Residual)	271	20.9	88.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable