

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (\*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,772	100.0	522.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,226	25.7	134.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	789	16.5	86.3
3	Newborn affected by maternal complications of pregnancy (P01)	327	6.9	35.8
4	Sudden infant death syndrome (R95)	236	4.9	25.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	201	4.2	22.0
6	Accidents (unintentional injuries) (V01-X59)	150	3.1	16.4
7	Bacterial sepsis of newborn (P36)	131	2.7	14.3
8	Neonatal hemorrhage (P50-P52,P54)	106	2.2	11.6
9	Diseases of the circulatory system (I00-I99)	84	1.8	9.2
10	Necrotizing enterocolitis of newborn (P77)	78	1.6	8.5
11	Respiratory distress of newborn (P22)	77	1.6	8.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	64	1.3	7.0
13	Diarrhea and gastroenteritis of infectious origin (A09)	54	1.1	5.9
14	Atelectasis (P28.0-P28.1)	52	1.1	5.7
15	Assault (homicide) (*U01,X85-Y09)	46	1.0	5.0
...	All other causes (residual)	1,151	24.1	125.9
Infants, Hispanic, male				
...	All causes	2,627	100.0	563.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	609	23.2	130.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	450	17.1	96.5
3	Newborn affected by maternal complications of pregnancy (P01)	196	7.5	42.0
4	Sudden infant death syndrome (R95)	129	4.9	27.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	104	4.0	22.3
6	Accidents (unintentional injuries) (V01-X59)	87	3.3	18.7
7	Bacterial sepsis of newborn (P36)	71	2.7	15.2
8	Neonatal hemorrhage (P50-P52,P54)	65	2.5	13.9
9	Diseases of the circulatory system (I00-I99)	44	1.7	9.4
9	Respiratory distress of newborn (P22)	44	1.7	9.4
9	Necrotizing enterocolitis of newborn (P77)	44	1.7	9.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.2	6.9
13	Diarrhea and gastroenteritis of infectious origin (A09)	31	1.2	6.6
13	Atelectasis (P28.0-P28.1)	31	1.2	6.6
15	Pulmonary hemorrhage originating in the perinatal period (P26)	24	0.9	5.1
...	All other causes (residual)	666	25.4	142.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,145	100.0	479.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	617	28.8	137.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	339	15.8	75.7
3	Newborn affected by maternal complications of pregnancy (P01)	131	6.1	29.3
4	Sudden infant death syndrome (R95)	107	5.0	23.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	97	4.5	21.7
6	Accidents (unintentional injuries) (V01-X59)	63	2.9	14.1
7	Bacterial sepsis of newborn (P36)	60	2.8	13.4
8	Neonatal hemorrhage (P50-P52,P54)	41	1.9	9.2
9	Diseases of the circulatory system (I00-I99)	40	1.9	8.9
10	Necrotizing enterocolitis of newborn (P77)	34	1.6	7.6
11	Respiratory distress of newborn (P22)	33	1.5	7.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.5	7.1
13	Diarrhea and gastroenteritis of infectious origin (A09)	23	1.1	5.1
13	Assault (homicide) (*U01,X85-Y09)	23	1.1	5.1
15	Hydrops fetalis not due to hemolytic disease (P83.2)	22	1.0	4.9
...	All other causes (residual)	483	22.5	107.9
Infants, Non-Hispanic, white, both sexes				
...	All causes	10,341	100.0	481.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,366	22.9	110.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,585	15.3	73.7
3	Sudden infant death syndrome (R95)	786	7.6	36.6
4	Newborn affected by maternal complications of pregnancy (P01)	604	5.8	28.1
5	Accidents (unintentional injuries) (V01-X59)	566	5.5	26.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	406	3.9	18.9
7	Bacterial sepsis of newborn (P36)	220	2.1	10.2
8	Respiratory distress of newborn (P22)	208	2.0	9.7
9	Neonatal hemorrhage (P50-P52,P54)	202	2.0	9.4
10	Diseases of the circulatory system (I00-I99)	187	1.8	8.7
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	163	1.6	7.6
12	Necrotizing enterocolitis of newborn (P77)	148	1.4	6.9
13	Assault (homicide) (*U01,X85-Y09)	113	1.1	5.3
14	Atelectasis (P28.0-P28.1)	103	1.0	4.8
15	Hydrops fetalis not due to hemolytic disease (P83.2)	101	1.0	4.7
...	All other causes (residual)	2,583	25.0	120.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	5,801	100.0	525.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,254	21.6	113.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	919	15.8	83.3
3	Sudden infant death syndrome (R95)	479	8.3	43.4
4	Newborn affected by maternal complications of pregnancy (P01)	341	5.9	30.9
5	Accidents (unintentional injuries) (V01-X59)	329	5.7	29.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	205	3.5	18.6
7	Respiratory distress of newborn (P22)	133	2.3	12.1
8	Neonatal hemorrhage (P50-P52,P54)	130	2.2	11.8
9	Bacterial sepsis of newborn (P36)	126	2.2	11.4
10	Diseases of the circulatory system (I00-I99)	109	1.9	9.9
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.5	7.8
12	Necrotizing enterocolitis of newborn (P77)	83	1.4	7.5
13	Assault (homicide) (*U01,X85-Y09)	69	1.2	6.3
14	Atelectasis (P28.0-P28.1)	66	1.1	6.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	52	0.9	4.7
...	All other causes (residual)	1,420	24.5	128.7
Infants, Non-Hispanic, white, female				
...	All causes	4,540	100.0	434.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,112	24.5	106.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	666	14.7	63.7
3	Sudden infant death syndrome (R95)	307	6.8	29.3
4	Newborn affected by maternal complications of pregnancy (P01)	263	5.8	25.1
5	Accidents (unintentional injuries) (V01-X59)	237	5.2	22.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	201	4.4	19.2
7	Bacterial sepsis of newborn (P36)	94	2.1	9.0
8	Diseases of the circulatory system (I00-I99)	78	1.7	7.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	77	1.7	7.4
10	Respiratory distress of newborn (P22)	75	1.7	7.2
11	Neonatal hemorrhage (P50-P52,P54)	72	1.6	6.9
12	Necrotizing enterocolitis of newborn (P77)	65	1.4	6.2
13	Hydrops fetalis not due to hemolytic disease (P83.2)	49	1.1	4.7
14	Assault (homicide) (*U01,X85-Y09)	44	1.0	4.2
15	Atelectasis (P28.0-P28.1)	37	0.8	3.5
...	All other causes (residual)	1,163	25.6	111.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	6,698	100.0	1,137.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,517	22.6	257.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	879	13.1	149.3
3	Newborn affected by maternal complications of pregnancy (P01)	534	8.0	90.7
4	Sudden infant death syndrome (R95)	451	6.7	76.6
5	Accidents (unintentional injuries) (V01-X59)	403	6.0	68.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	295	4.4	50.1
7	Bacterial sepsis of newborn (P36)	163	2.4	27.7
8	Respiratory distress of newborn (P22)	152	2.3	25.8
9	Diseases of the circulatory system (I00-I99)	141	2.1	23.9
10	Necrotizing enterocolitis of newborn (P77)	138	2.1	23.4
11	Neonatal hemorrhage (P50-P52,P54)	113	1.7	19.2
12	Diarrhea and gastroenteritis of infectious origin (A09)	88	1.3	14.9
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	80	1.2	13.6
14	Assault (homicide) (*U01,X85-Y09)	74	1.1	12.6
15	Influenza and pneumonia (J09-J18)	71	1.1	12.1
...	All other causes (residual)	1,599	23.9	271.5
Infants, Non-Hispanic, black, male				
...	All causes	3,680	100.0	1,233.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	841	22.9	281.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	444	12.1	148.8
3	Newborn affected by maternal complications of pregnancy (P01)	316	8.6	105.9
4	Sudden infant death syndrome (R95)	260	7.1	87.1
5	Accidents (unintentional injuries) (V01-X59)	225	6.1	75.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	160	4.3	53.6
7	Bacterial sepsis of newborn (P36)	84	2.3	28.2
8	Respiratory distress of newborn (P22)	82	2.2	27.5
9	Necrotizing enterocolitis of newborn (P77)	79	2.1	26.5
10	Diseases of the circulatory system (I00-I99)	74	2.0	24.8
11	Neonatal hemorrhage (P50-P52,P54)	64	1.7	21.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	1.3	16.1
13	Diarrhea and gastroenteritis of infectious origin (A09)	46	1.3	15.4
14	Assault (homicide) (*U01,X85-Y09)	43	1.2	14.4
15	Atelectasis (P28.0-P28.1)	40	1.1	13.4
...	All other causes (residual)	874	23.8	292.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	3,018	100.0	1,038.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	676	22.4	232.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	435	14.4	149.7
3	Newborn affected by maternal complications of pregnancy (P01)	218	7.2	75.0
4	Sudden infant death syndrome (R95)	191	6.3	65.7
5	Accidents (unintentional injuries) (V01-X59)	178	5.9	61.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	135	4.5	46.5
7	Bacterial sepsis of newborn (P36)	79	2.6	27.2
8	Respiratory distress of newborn (P22)	70	2.3	24.1
9	Diseases of the circulatory system (I00-I99)	67	2.2	23.1
10	Necrotizing enterocolitis of newborn (P77)	59	2.0	20.3
11	Neonatal hemorrhage (P50-P52,P54)	49	1.6	16.9
12	Influenza and pneumonia (J09-J18)	44	1.5	15.1
13	Diarrhea and gastroenteritis of infectious origin (A09)	42	1.4	14.5
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.1	11.0
15	Assault (homicide) (*U01,X85-Y09)	31	1.0	10.7
...	All other causes (residual)	712	23.6	245.1
Neonates, Hispanic, both sexes				
...	All causes	3,352	100.0	366.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	870	26.0	95.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	771	23.0	84.3
3	Newborn affected by maternal complications of pregnancy (P01)	326	9.7	35.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	198	5.9	21.7
5	Bacterial sepsis of newborn (P36)	125	3.7	13.7
6	Neonatal hemorrhage (P50-P52,P54)	102	3.0	11.2
7	Respiratory distress of newborn (P22)	77	2.3	8.4
8	Necrotizing enterocolitis of newborn (P77)	69	2.1	7.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	61	1.8	6.7
10	Atelectasis (P28.0-P28.1)	49	1.5	5.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	32	1.0	3.5
11	Hydrops fetalis not due to hemolytic disease (P83.2)	32	1.0	3.5
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	20	0.6	2.2
13	Newborn affected by other complications of labor and delivery (P03)	20	0.6	2.2
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	20	0.6	2.2
13	Hematological disorders (P60-P61)	20	0.6	2.2
...	All other causes (residual)	560	16.7	61.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,855	100.0	397.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	441	23.8	94.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	441	23.8	94.6
3	Newborn affected by maternal complications of pregnancy (P01)	195	10.5	41.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	102	5.5	21.9
5	Bacterial sepsis of newborn (P36)	67	3.6	14.4
6	Neonatal hemorrhage (P50-P52,P54)	64	3.5	13.7
7	Respiratory distress of newborn (P22)	44	2.4	9.4
8	Necrotizing enterocolitis of newborn (P77)	38	2.0	8.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	30	1.6	6.4
10	Atelectasis (P28.0-P28.1)	29	1.6	6.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	24	1.3	5.1
12	Hydrops fetalis not due to hemolytic disease (P83.2)	15	0.8	*
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	13	0.7	*
14	Newborn affected by other complications of labor and delivery (P03)	12	0.6	*
15	Diseases of the circulatory system (I00-I99)	11	0.6	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	11	0.6	*
15	Hematological disorders (P60-P61)	11	0.6	*
...	All other causes (residual)	307	16.5	65.8
Neonates, Hispanic, female				
...	All causes	1,497	100.0	334.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	429	28.7	95.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	330	22.0	73.7
3	Newborn affected by maternal complications of pregnancy (P01)	131	8.8	29.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	96	6.4	21.4
5	Bacterial sepsis of newborn (P36)	58	3.9	13.0
6	Neonatal hemorrhage (P50-P52,P54)	38	2.5	8.5
7	Respiratory distress of newborn (P22)	33	2.2	7.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	2.1	6.9
8	Necrotizing enterocolitis of newborn (P77)	31	2.1	6.9
10	Atelectasis (P28.0-P28.1)	20	1.3	4.5
11	Hydrops fetalis not due to hemolytic disease (P83.2)	17	1.1	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	13	0.9	*
13	Slow fetal growth and fetal malnutrition (P05)	11	0.7	*
14	Neonatal aspiration syndromes (P24)	9	0.6	*
14	Hematological disorders (P60-P61)	9	0.6	*
...	All other causes (residual)	241	16.1	53.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	6,944	100.0	323.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,724	24.8	80.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,547	22.3	72.0
3	Newborn affected by maternal complications of pregnancy (P01)	602	8.7	28.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	402	5.8	18.7
5	Bacterial sepsis of newborn (P36)	213	3.1	9.9
6	Respiratory distress of newborn (P22)	203	2.9	9.4
7	Neonatal hemorrhage (P50-P52,P54)	198	2.9	9.2
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	154	2.2	7.2
9	Necrotizing enterocolitis of newborn (P77)	137	2.0	6.4
10	Hydrops fetalis not due to hemolytic disease (P83.2)	99	1.4	4.6
11	Atelectasis (P28.0-P28.1)	94	1.4	4.4
12	Sudden infant death syndrome (R95)	84	1.2	3.9
13	Accidents (unintentional injuries) (V01-X59)	75	1.1	3.5
14	Pulmonary hemorrhage originating in the perinatal period (P26)	65	0.9	3.0
15	Newborn affected by other complications of labor and delivery (P03)	51	0.7	2.4
15	Hematological disorders (P60-P61)	51	0.7	2.4
...	All other causes (residual)	1,245	17.9	57.9
Neonates, Non-Hispanic, white, male				
...	All causes	3,841	100.0	348.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	919	23.9	83.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	898	23.4	81.4
3	Newborn affected by maternal complications of pregnancy (P01)	340	8.9	30.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	205	5.3	18.6
5	Respiratory distress of newborn (P22)	131	3.4	11.9
6	Neonatal hemorrhage (P50-P52,P54)	127	3.3	11.5
7	Bacterial sepsis of newborn (P36)	121	3.2	11.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	82	2.1	7.4
9	Necrotizing enterocolitis of newborn (P77)	78	2.0	7.1
10	Atelectasis (P28.0-P28.1)	58	1.5	5.3
11	Hydrops fetalis not due to hemolytic disease (P83.2)	50	1.3	4.5
12	Sudden infant death syndrome (R95)	47	1.2	4.3
13	Pulmonary hemorrhage originating in the perinatal period (P26)	43	1.1	3.9
14	Accidents (unintentional injuries) (V01-X59)	32	0.8	2.9
15	Newborn affected by other complications of labor and delivery (P03)	30	0.8	2.7
...	All other causes (residual)	680	17.7	61.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,103	100.0	296.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	805	25.9	77.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	649	20.9	62.0
3	Newborn affected by maternal complications of pregnancy (P01)	262	8.4	25.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	197	6.3	18.8
5	Bacterial sepsis of newborn (P36)	92	3.0	8.8
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	72	2.3	6.9
6	Respiratory distress of newborn (P22)	72	2.3	6.9
8	Neonatal hemorrhage (P50-P52,P54)	71	2.3	6.8
9	Necrotizing enterocolitis of newborn (P77)	59	1.9	5.6
10	Hydrops fetalis not due to hemolytic disease (P83.2)	49	1.6	4.7
11	Accidents (unintentional injuries) (V01-X59)	43	1.4	4.1
12	Sudden infant death syndrome (R95)	37	1.2	3.5
13	Atelectasis (P28.0-P28.1)	36	1.2	3.4
14	Hematological disorders (P60-P61)	25	0.8	2.4
15	Diseases of the circulatory system (I00-I99)	24	0.8	2.3
...	All other causes (residual)	610	19.7	58.3
Neonates, Non-Hispanic, black, both sexes				
...	All causes	4,422	100.0	750.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,483	33.5	251.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	556	12.6	94.4
3	Newborn affected by maternal complications of pregnancy (P01)	531	12.0	90.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	293	6.6	49.8
5	Bacterial sepsis of newborn (P36)	159	3.6	27.0
6	Respiratory distress of newborn (P22)	147	3.3	25.0
7	Necrotizing enterocolitis of newborn (P77)	120	2.7	20.4
8	Neonatal hemorrhage (P50-P52,P54)	108	2.4	18.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	76	1.7	12.9
10	Atelectasis (P28.0-P28.1)	66	1.5	11.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	54	1.2	9.2
12	Slow fetal growth and fetal malnutrition (P05)	39	0.9	6.6
13	Sudden infant death syndrome (R95)	36	0.8	6.1
14	Accidents (unintentional injuries) (V01-X59)	32	0.7	5.4
15	Newborn affected by other complications of labor and delivery (P03)	28	0.6	4.8
...	All other causes (residual)	694	15.7	117.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	2,425	100.0	812.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	819	33.8	274.5
2	Newborn affected by maternal complications of pregnancy (P01)	315	13.0	105.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	11.6	94.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	159	6.6	53.3
5	Bacterial sepsis of newborn (P36)	81	3.3	27.1
6	Respiratory distress of newborn (P22)	79	3.3	26.5
7	Necrotizing enterocolitis of newborn (P77)	69	2.8	23.1
8	Neonatal hemorrhage (P50-P52,P54)	61	2.5	20.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	45	1.9	15.1
10	Atelectasis (P28.0-P28.1)	38	1.6	12.7
11	Pulmonary hemorrhage originating in the perinatal period (P26)	34	1.4	11.4
12	Newborn affected by other complications of labor and delivery (P03)	22	0.9	7.4
13	Slow fetal growth and fetal malnutrition (P05)	17	0.7	*
13	Sudden infant death syndrome (R95)	17	0.7	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	16	0.7	*
15	Accidents (unintentional injuries) (V01-X59)	16	0.7	*
...	All other causes (residual)	356	14.7	119.3
Neonates, Non-Hispanic, black, female				
...	All causes	1,997	100.0	687.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	664	33.2	228.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	13.8	94.7
3	Newborn affected by maternal complications of pregnancy (P01)	216	10.8	74.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	134	6.7	46.1
5	Bacterial sepsis of newborn (P36)	78	3.9	26.8
6	Respiratory distress of newborn (P22)	68	3.4	23.4
7	Necrotizing enterocolitis of newborn (P77)	51	2.6	17.6
8	Neonatal hemorrhage (P50-P52,P54)	47	2.4	16.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	1.6	10.7
10	Atelectasis (P28.0-P28.1)	28	1.4	9.6
11	Slow fetal growth and fetal malnutrition (P05)	22	1.1	7.6
12	Pulmonary hemorrhage originating in the perinatal period (P26)	20	1.0	6.9
13	Sudden infant death syndrome (R95)	19	1.0	*
14	Accidents (unintentional injuries) (V01-X59)	16	0.8	*
15	Congenital pneumonia (P23)	13	0.7	*
...	All other causes (residual)	315	15.8	108.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,420	100.0	155.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	356	25.1	38.9
2	Sudden infant death syndrome (R95)	219	15.4	24.0
3	Accidents (unintentional injuries) (V01-X59)	136	9.6	14.9
4	Diseases of the circulatory system (I00-I99)	66	4.6	7.2
5	Diarrhea and gastroenteritis of infectious origin (A09)	52	3.7	5.7
6	Assault (homicide) (*U01,X85-Y09)	41	2.9	4.5
7	Septicemia (A40-A41)	39	2.7	4.3
8	Influenza and pneumonia (J09-J18)	35	2.5	3.8
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.3	*
10	Malignant neoplasms (C00-C97)	10	0.7	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.7	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	9	0.6	*
12	Chronic respiratory disease originating in the perinatal period (P27)	9	0.6	*
12	Necrotizing enterocolitis of newborn (P77)	9	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	0.6	*
...	All other causes (residual)	403	28.4	44.1
Postneonates, Hispanic, male				
...	All causes	772	100.0	165.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	21.8	36.0
2	Sudden infant death syndrome (R95)	120	15.5	25.7
3	Accidents (unintentional injuries) (V01-X59)	79	10.2	16.9
4	Diseases of the circulatory system (I00-I99)	33	4.3	7.1
5	Diarrhea and gastroenteritis of infectious origin (A09)	30	3.9	6.4
6	Septicemia (A40-A41)	23	3.0	4.9
7	Assault (homicide) (*U01,X85-Y09)	22	2.8	4.7
8	Influenza and pneumonia (J09-J18)	20	2.6	4.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.2	*
10	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	6	0.8	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	6	0.8	*
10	Necrotizing enterocolitis of newborn (P77)	6	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	5	0.6	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	5	0.6	*
15	Whooping cough (A37)	4	0.5	*
15	Malignant neoplasms (C00-C97)	4	0.5	*
15	Acute upper respiratory infections (J00-J06)	4	0.5	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	4	0.5	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	4	0.5	*
15	Chronic respiratory disease originating in the perinatal period (P27)	4	0.5	*
15	Bacterial sepsis of newborn (P36)	4	0.5	*
...	All other causes (residual)	212	27.5	45.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	648	100.0	144.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	29.0	42.0
2	Sudden infant death syndrome (R95)	99	15.3	22.1
3	Accidents (unintentional injuries) (V01-X59)	57	8.8	12.7
4	Diseases of the circulatory system (I00-I99)	33	5.1	7.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	22	3.4	4.9
6	Assault (homicide) (*U01,X85-Y09)	19	2.9	*
7	Septicemia (A40-A41)	16	2.5	*
8	Influenza and pneumonia (J09-J18)	15	2.3	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.4	*
10	Malignant neoplasms (C00-C97)	6	0.9	*
11	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	5	0.8	*
11	Chronic respiratory disease originating in the perinatal period (P27)	5	0.8	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	5	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	4	0.6	*
...	All other causes (residual)	161	24.8	36.0
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	3,397	100.0	158.1
1	Sudden infant death syndrome (R95)	702	20.7	32.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	642	18.9	29.9
3	Accidents (unintentional injuries) (V01-X59)	491	14.5	22.8
4	Diseases of the circulatory system (I00-I99)	141	4.2	6.6
5	Assault (homicide) (*U01,X85-Y09)	105	3.1	4.9
6	Influenza and pneumonia (J09-J18)	67	2.0	3.1
7	Diarrhea and gastroenteritis of infectious origin (A09)	60	1.8	2.8
8	Septicemia (A40-A41)	55	1.6	2.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	38	1.1	1.8
9	Chronic respiratory disease originating in the perinatal period (P27)	38	1.1	1.8
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	30	0.9	1.4
12	Malignant neoplasms (C00-C97)	23	0.7	1.1
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.4	*
14	Meningitis (G00,G03)	13	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	11	0.3	*
15	Necrotizing enterocolitis of newborn (P77)	11	0.3	*
...	All other causes (residual)	945	27.8	44.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	1,960	100.0	177.7
1	Sudden infant death syndrome (R95)	432	22.0	39.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	335	17.1	30.4
3	Accidents (unintentional injuries) (V01-X59)	297	15.2	26.9
4	Diseases of the circulatory system (I00-I99)	87	4.4	7.9
5	Assault (homicide) (*U01,X85-Y09)	66	3.4	6.0
6	Influenza and pneumonia (J09-J18)	36	1.8	3.3
7	Septicemia (A40-A41)	31	1.6	2.8
8	Diarrhea and gastroenteritis of infectious origin (A09)	28	1.4	2.5
9	Chronic respiratory disease originating in the perinatal period (P27)	27	1.4	2.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	1.1	1.9
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.8	*
12	Malignant neoplasms (C00-C97)	14	0.7	*
13	Atelectasis (P28.0-P28.1)	8	0.4	*
14	Bronchitis, chronic and unspecified (J40-J42)	7	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	6	0.3	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.3	*
...	All other causes (residual)	544	27.8	49.3
Postneonates, Non-Hispanic, white, female				
...	All causes	1,437	100.0	137.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	307	21.4	29.3
2	Sudden infant death syndrome (R95)	270	18.8	25.8
3	Accidents (unintentional injuries) (V01-X59)	194	13.5	18.5
4	Diseases of the circulatory system (I00-I99)	54	3.8	5.2
5	Assault (homicide) (*U01,X85-Y09)	39	2.7	3.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	32	2.2	3.1
7	Influenza and pneumonia (J09-J18)	31	2.2	3.0
8	Septicemia (A40-A41)	24	1.7	2.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.2	*
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	1.0	*
11	Chronic respiratory disease originating in the perinatal period (P27)	11	0.8	*
12	Malignant neoplasms (C00-C97)	9	0.6	*
13	Meningitis (G00,G03)	8	0.6	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (residual)	411	28.6	39.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	2,276	100.0	386.5
1	Sudden infant death syndrome (R95)	415	18.2	70.5
2	Accidents (unintentional injuries) (V01-X59)	371	16.3	63.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	323	14.2	54.8
4	Diseases of the circulatory system (I00-I99)	119	5.2	20.2
5	Diarrhea and gastroenteritis of infectious origin (A09)	88	3.9	14.9
6	Assault (homicide) (*U01,X85-Y09)	72	3.2	12.2
7	Influenza and pneumonia (J09-J18)	70	3.1	11.9
8	Septicemia (A40-A41)	52	2.3	8.8
9	Chronic respiratory disease originating in the perinatal period (P27)	35	1.5	5.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	34	1.5	5.8
11	Necrotizing enterocolitis of newborn (P77)	18	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.4	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	9	0.4	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	9	0.4	*
...	All other causes (residual)	626	27.5	106.3
Postneonates, Non-Hispanic, black, male				
...	All causes	1,255	100.0	420.6
1	Sudden infant death syndrome (R95)	243	19.4	81.4
2	Accidents (unintentional injuries) (V01-X59)	209	16.7	70.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	13.0	54.6
4	Diseases of the circulatory system (I00-I99)	61	4.9	20.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	46	3.7	15.4
6	Assault (homicide) (*U01,X85-Y09)	42	3.3	14.1
7	Septicemia (A40-A41)	27	2.2	9.0
8	Influenza and pneumonia (J09-J18)	26	2.1	8.7
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	1.8	7.4
9	Chronic respiratory disease originating in the perinatal period (P27)	22	1.8	7.4
11	Necrotizing enterocolitis of newborn (P77)	10	0.8	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.7	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.6	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.5	*
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	5	0.4	*
...	All other causes (residual)	352	28.0	118.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,021	100.0	351.5
1	Sudden infant death syndrome (R95)	172	16.8	59.2
2	Accidents (unintentional injuries) (V01-X59)	162	15.9	55.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	15.7	55.1
4	Diseases of the circulatory system (I00-I99)	58	5.7	20.0
5	Influenza and pneumonia (J09-J18)	44	4.3	15.1
6	Diarrhea and gastroenteritis of infectious origin (A09)	42	4.1	14.5
7	Assault (homicide) (*U01,X85-Y09)	30	2.9	10.3
8	Septicemia (A40-A41)	25	2.4	8.6
9	Chronic respiratory disease originating in the perinatal period (P27)	13	1.3	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	12	1.2	*
11	Meningitis (G00,G03)	9	0.9	*
12	Malignant neoplasms (C00-C97)	8	0.8	*
12	Necrotizing enterocolitis of newborn (P77)	8	0.8	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
...	All other causes (residual)	266	26.1	91.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014