

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,321	100.0	562.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,278	24.0	135.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	816	15.3	86.2
3	Sudden infant death syndrome (R95)	284	5.3	30.0
4	Newborn affected by maternal complications of pregnancy (P01)	265	5.0	28.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	196	3.7	20.7
6	Bacterial sepsis of newborn (P36)	176	3.3	18.6
7	Respiratory distress of newborn (P22)	167	3.1	17.6
8	Accidents (unintentional injuries) (V01-X59)	159	3.0	16.8
9	Neonatal hemorrhage (P50-P52,P54)	133	2.5	14.1
10	Diseases of the circulatory system (I00-I99)	109	2.0	11.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	105	2.0	11.1
12	Atelectasis (P28.0-P28.1)	98	1.8	10.4
13	Necrotizing enterocolitis of newborn (P77)	89	1.7	9.4
14	Assault (homicide) (*U01,X85-Y09)	73	1.4	7.7
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	59	1.1	6.2
...	All other causes (Residual)	1,314	24.7	138.8
Infants, Hispanic, male				
...	All causes	2,948	100.0	610.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	622	21.1	128.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	476	16.1	98.6
3	Sudden infant death syndrome (R95)	169	5.7	35.0
4	Newborn affected by maternal complications of pregnancy (P01)	155	5.3	32.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	106	3.6	21.9
6	Bacterial sepsis of newborn (P36)	101	3.4	20.9
7	Respiratory distress of newborn (P22)	93	3.2	19.3
8	Accidents (unintentional injuries) (V01-X59)	87	3.0	18.0
9	Neonatal hemorrhage (P50-P52,P54)	84	2.8	17.4
10	Atelectasis (P28.0-P28.1)	60	2.0	12.4
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	59	2.0	12.2
12	Diseases of the circulatory system (I00-I99)	57	1.9	11.8
13	Necrotizing enterocolitis of newborn (P77)	50	1.7	10.4
14	Assault (homicide) (*U01,X85-Y09)	42	1.4	8.7
15	Influenza and pneumonia (J10-J18)	37	1.3	7.7
...	All other causes (Residual)	750	25.4	155.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,373	100.0	512.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	656	27.6	141.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	340	14.3	73.4
3	Sudden infant death syndrome (R95)	115	4.8	24.8
4	Newborn affected by maternal complications of pregnancy (P01)	110	4.6	23.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	90	3.8	19.4
6	Bacterial sepsis of newborn (P36)	75	3.2	16.2
7	Respiratory distress of newborn (P22)	74	3.1	16.0
8	Accidents (unintentional injuries) (V01-X59)	72	3.0	15.5
9	Diseases of the circulatory system (I00-I99)	52	2.2	11.2
10	Neonatal hemorrhage (P50-P52,P54)	49	2.1	10.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	46	1.9	9.9
12	Necrotizing enterocolitis of newborn (P77)	39	1.6	8.4
13	Atelectasis (P28.0-P28.1)	38	1.6	8.2
14	Assault (homicide) (*U01,X85-Y09)	31	1.3	6.7
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	27	1.1	5.8
...	All other causes (Residual)	559	23.6	120.6
Infants, white, non-Hispanic, both sexes				
...	All causes	13,046	100.0	568.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,016	23.1	131.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,848	14.2	80.5
3	Sudden infant death syndrome (R95)	1,173	9.0	51.1
4	Newborn affected by maternal complications of pregnancy (P01)	747	5.7	32.5
5	Accidents (unintentional injuries) (V01-X59)	557	4.3	24.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	490	3.8	21.3
7	Respiratory distress of newborn (P22)	398	3.1	17.3
8	Bacterial sepsis of newborn (P36)	349	2.7	15.2
9	Diseases of the circulatory system (I00-I99)	294	2.3	12.8
10	Neonatal hemorrhage (P50-P52,P54)	277	2.1	12.1
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	275	2.1	12.0
12	Atelectasis (P28.0-P28.1)	206	1.6	9.0
13	Necrotizing enterocolitis of newborn (P77)	170	1.3	7.4
14	Assault (homicide) (*U01,X85-Y09)	137	1.1	6.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	122	0.9	5.3
...	All other causes (Residual)	2,987	22.9	130.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,399	100.0	628.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,572	21.2	133.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,052	14.2	89.3
3	Sudden infant death syndrome (R95)	689	9.3	58.5
4	Newborn affected by maternal complications of pregnancy (P01)	439	5.9	37.3
5	Accidents (unintentional injuries) (V01-X59)	335	4.5	28.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	262	3.5	22.2
7	Respiratory distress of newborn (P22)	229	3.1	19.4
8	Bacterial sepsis of newborn (P36)	194	2.6	16.5
9	Neonatal hemorrhage (P50-P52,P54)	174	2.4	14.8
10	Diseases of the circulatory system (I00-I99)	166	2.2	14.1
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	157	2.1	13.3
12	Atelectasis (P28.0-P28.1)	130	1.8	11.0
13	Necrotizing enterocolitis of newborn (P77)	96	1.3	8.1
14	Assault (homicide) (*U01,X85-Y09)	72	1.0	6.1
15	Septicemia (A40-A41)	68	0.9	5.8
...	All other causes (Residual)	1,764	23.8	149.7
Infants, white, non-Hispanic, female				
...	All causes	5,647	100.0	504.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,444	25.6	129.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	796	14.1	71.2
3	Sudden infant death syndrome (R95)	484	8.6	43.3
4	Newborn affected by maternal complications of pregnancy (P01)	308	5.5	27.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	228	4.0	20.4
6	Accidents (unintentional injuries) (V01-X59)	222	3.9	19.8
7	Respiratory distress of newborn (P22)	169	3.0	15.1
8	Bacterial sepsis of newborn (P36)	155	2.7	13.9
9	Diseases of the circulatory system (I00-I99)	128	2.3	11.4
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	118	2.1	10.5
11	Neonatal hemorrhage (P50-P52,P54)	103	1.8	9.2
12	Atelectasis (P28.0-P28.1)	76	1.3	6.8
13	Necrotizing enterocolitis of newborn (P77)	74	1.3	6.6
14	Assault (homicide) (*U01,X85-Y09)	65	1.2	5.8
15	Hydrops fetalis not due to hemolytic disease (P83.2)	59	1.0	5.3
...	All other causes (Residual)	1,218	21.6	108.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,221	100.0	1,420.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,748	21.3	302.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,041	12.7	179.9
3	Sudden infant death syndrome (R95)	715	8.7	123.5
4	Newborn affected by maternal complications of pregnancy (P01)	618	7.5	106.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	307	3.7	53.0
6	Accidents (unintentional injuries) (V01-X59)	298	3.6	51.5
7	Respiratory distress of newborn (P22)	282	3.4	48.7
8	Bacterial sepsis of newborn (P36)	263	3.2	45.4
9	Neonatal hemorrhage (P50-P52,P54)	166	2.0	28.7
10	Necrotizing enterocolitis of newborn (P77)	160	1.9	27.6
11	Diseases of the circulatory system (I00-I99)	149	1.8	25.7
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	130	1.6	22.5
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	124	1.5	21.4
14	Chronic respiratory disease originating in the perinatal period (P27)	113	1.4	19.5
15	Atelectasis (P28.0-P28.1)	103	1.3	17.8
...	All other causes (Residual)	2,004	24.4	346.3
Infants, black, non-Hispanic, male				
...	All causes	4,613	100.0	1,565.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	991	21.5	336.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	551	11.9	187.0
3	Sudden infant death syndrome (R95)	429	9.3	145.6
4	Newborn affected by maternal complications of pregnancy (P01)	334	7.2	113.3
5	Respiratory distress of newborn (P22)	164	3.6	55.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	160	3.5	54.3
7	Accidents (unintentional injuries) (V01-X59)	159	3.4	53.9
8	Bacterial sepsis of newborn (P36)	152	3.3	51.6
9	Neonatal hemorrhage (P50-P52,P54)	103	2.2	34.9
10	Necrotizing enterocolitis of newborn (P77)	92	2.0	31.2
11	Diseases of the circulatory system (I00-I99)	83	1.8	28.2
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	80	1.7	27.1
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	69	1.5	23.4
14	Chronic respiratory disease originating in the perinatal period (P27)	63	1.4	21.4
15	Influenza and pneumonia (J10-J18)	55	1.2	18.7
...	All other causes (Residual)	1,128	24.5	382.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,608	100.0	1,270.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	757	21.0	266.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	490	13.6	172.5
3	Sudden infant death syndrome (R95)	286	7.9	100.7
4	Newborn affected by maternal complications of pregnancy (P01)	284	7.9	100.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	147	4.1	51.8
6	Accidents (unintentional injuries) (V01-X59)	139	3.9	48.9
7	Respiratory distress of newborn (P22)	118	3.3	41.5
8	Bacterial sepsis of newborn (P36)	111	3.1	39.1
9	Necrotizing enterocolitis of newborn (P77)	68	1.9	23.9
10	Diseases of the circulatory system (I00-I99)	66	1.8	23.2
11	Neonatal hemorrhage (P50-P52,P54)	63	1.7	22.2
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	55	1.5	19.4
12	Assault (homicide) (*U01,X85-Y09)	55	1.5	19.4
14	Atelectasis (P28.0-P28.1)	51	1.4	18.0
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	50	1.4	17.6
15	Chronic respiratory disease originating in the perinatal period (P27)	50	1.4	17.6
...	All other causes (Residual)	818	22.7	288.0
Neonates, Hispanic, both sexes				
...	All causes	3,633	100.0	383.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	895	24.6	94.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	789	21.7	83.4
3	Newborn affected by maternal complications of pregnancy (P01)	261	7.2	27.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	194	5.3	20.5
5	Bacterial sepsis of newborn (P36)	163	4.5	17.2
6	Respiratory distress of newborn (P22)	155	4.3	16.4
7	Neonatal hemorrhage (P50-P52,P54)	128	3.5	13.5
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	100	2.8	10.6
9	Atelectasis (P28.0-P28.1)	95	2.6	10.0
10	Necrotizing enterocolitis of newborn (P77)	71	2.0	7.5
11	Hydrops fetalis not due to hemolytic disease (P83.2)	45	1.2	4.8
12	Diseases of the circulatory system (I00-I99)	38	1.0	4.0
13	Pulmonary hemorrhage originating in the perinatal period (P26)	32	0.9	3.4
14	Sudden infant death syndrome (R95)	28	0.8	3.0
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	24	0.7	2.5
...	All other causes (Residual)	615	16.9	65.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	2,015	100.0	417.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	458	22.7	94.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	451	22.4	93.4
3	Newborn affected by maternal complications of pregnancy (P01)	153	7.6	31.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	105	5.2	21.7
5	Bacterial sepsis of newborn (P36)	92	4.6	19.1
6	Respiratory distress of newborn (P22)	90	4.5	18.6
7	Neonatal hemorrhage (P50-P52,P54)	81	4.0	16.8
8	Atelectasis (P28.0-P28.1)	59	2.9	12.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	55	2.7	11.4
10	Necrotizing enterocolitis of newborn (P77)	37	1.8	7.7
11	Pulmonary hemorrhage originating in the perinatal period (P26)	24	1.2	5.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	23	1.1	4.8
13	Diseases of the circulatory system (I00-I99)	19	0.9	*
14	Sudden infant death syndrome (R95)	16	0.8	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.7	*
...	All other causes (Residual)	338	16.8	70.0
Neonates, Hispanic, female				
...	All causes	1,618	100.0	349.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	444	27.4	95.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	331	20.5	71.4
3	Newborn affected by maternal complications of pregnancy (P01)	108	6.7	23.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	89	5.5	19.2
5	Bacterial sepsis of newborn (P36)	71	4.4	15.3
6	Respiratory distress of newborn (P22)	65	4.0	14.0
7	Neonatal hemorrhage (P50-P52,P54)	47	2.9	10.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	45	2.8	9.7
9	Atelectasis (P28.0-P28.1)	36	2.2	7.8
10	Necrotizing enterocolitis of newborn (P77)	34	2.1	7.3
11	Hydrops fetalis not due to hemolytic disease (P83.2)	22	1.4	4.7
12	Diseases of the circulatory system (I00-I99)	19	1.2	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	17	1.1	*
14	Sudden infant death syndrome (R95)	12	0.7	*
15	Newborn affected by other complications of labor and delivery (P03)	11	0.7	*
15	Congenital pneumonia (P23)	11	0.7	*
...	All other causes (Residual)	256	15.8	55.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	8,638	100.0	376.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,215	25.6	96.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,804	20.9	78.5
3	Newborn affected by maternal complications of pregnancy (P01)	740	8.6	32.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	484	5.6	21.1
5	Respiratory distress of newborn (P22)	375	4.3	16.3
6	Bacterial sepsis of newborn (P36)	332	3.8	14.5
7	Neonatal hemorrhage (P50-P52,P54)	268	3.1	11.7
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	261	3.0	11.4
9	Atelectasis (P28.0-P28.1)	202	2.3	8.8
10	Necrotizing enterocolitis of newborn (P77)	147	1.7	6.4
11	Sudden infant death syndrome (R95)	115	1.3	5.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	114	1.3	5.0
13	Diseases of the circulatory system (I00-I99)	108	1.3	4.7
14	Accidents (unintentional injuries) (V01-X59)	74	0.9	3.2
15	Newborn affected by other complications of labor and delivery (P03)	71	0.8	3.1
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	71	0.8	3.1
...	All other causes (Residual)	1,257	14.6	54.7
Neonates, white, non-Hispanic, male				
...	All causes	4,866	100.0	413.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,171	24.1	99.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,022	21.0	86.7
3	Newborn affected by maternal complications of pregnancy (P01)	433	8.9	36.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	257	5.3	21.8
5	Respiratory distress of newborn (P22)	221	4.5	18.8
6	Bacterial sepsis of newborn (P36)	186	3.8	15.8
7	Neonatal hemorrhage (P50-P52,P54)	169	3.5	14.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	149	3.1	12.6
9	Atelectasis (P28.0-P28.1)	129	2.7	10.9
10	Necrotizing enterocolitis of newborn (P77)	81	1.7	6.9
11	Diseases of the circulatory system (I00-I99)	63	1.3	5.3
12	Sudden infant death syndrome (R95)	61	1.3	5.2
13	Hydrops fetalis not due to hemolytic disease (P83.2)	58	1.2	4.9
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	46	0.9	3.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	37	0.8	3.1
...	All other causes (Residual)	783	16.1	66.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,772	100.0	337.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,044	27.7	93.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	782	20.7	69.9
3	Newborn affected by maternal complications of pregnancy (P01)	307	8.1	27.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	227	6.0	20.3
5	Respiratory distress of newborn (P22)	154	4.1	13.8
6	Bacterial sepsis of newborn (P36)	146	3.9	13.1
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	112	3.0	10.0
8	Neonatal hemorrhage (P50-P52,P54)	99	2.6	8.9
9	Atelectasis (P28.0-P28.1)	73	1.9	6.5
10	Necrotizing enterocolitis of newborn (P77)	66	1.7	5.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	56	1.5	5.0
12	Sudden infant death syndrome (R95)	54	1.4	4.8
13	Diseases of the circulatory system (I00-I99)	45	1.2	4.0
14	Accidents (unintentional injuries) (V01-X59)	39	1.0	3.5
15	Newborn affected by other complications of labor and delivery (P03)	36	1.0	3.2
...	All other causes (Residual)	532	14.1	47.6
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,417	100.0	935.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,699	31.4	293.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	672	12.4	116.1
3	Newborn affected by maternal complications of pregnancy (P01)	612	11.3	105.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	304	5.6	52.5
5	Respiratory distress of newborn (P22)	261	4.8	45.1
6	Bacterial sepsis of newborn (P36)	248	4.6	42.8
7	Neonatal hemorrhage (P50-P52,P54)	163	3.0	28.2
8	Necrotizing enterocolitis of newborn (P77)	134	2.5	23.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	111	2.0	19.2
10	Atelectasis (P28.0-P28.1)	101	1.9	17.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	70	1.3	12.1
12	Sudden infant death syndrome (R95)	64	1.2	11.1
13	Diseases of the circulatory system (I00-I99)	40	0.7	6.9
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	37	0.7	6.4
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	35	0.6	6.0
15	Newborn affected by other complications of labor and delivery (P03)	35	0.6	6.0
...	All other causes (Residual)	831	15.3	143.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,008	100.0	1,020.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	963	32.0	326.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	361	12.0	122.5
3	Newborn affected by maternal complications of pregnancy (P01)	330	11.0	112.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	158	5.3	53.6
5	Respiratory distress of newborn (P22)	153	5.1	51.9
6	Bacterial sepsis of newborn (P36)	142	4.7	48.2
7	Neonatal hemorrhage (P50-P52,P54)	102	3.4	34.6
8	Necrotizing enterocolitis of newborn (P77)	77	2.6	26.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	64	2.1	21.7
10	Atelectasis (P28.0-P28.1)	50	1.7	17.0
11	Pulmonary hemorrhage originating in the perinatal period (P26)	36	1.2	12.2
12	Sudden infant death syndrome (R95)	34	1.1	11.5
13	Diseases of the circulatory system (I00-I99)	25	0.8	8.5
14	Newborn affected by other complications of labor and delivery (P03)	22	0.7	7.5
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.6	*
...	All other causes (Residual)	472	15.7	160.1
Neonates, black, non-Hispanic, female				
...	All causes	2,409	100.0	848.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	736	30.6	259.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	311	12.9	109.5
3	Newborn affected by maternal complications of pregnancy (P01)	282	11.7	99.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	146	6.1	51.4
5	Respiratory distress of newborn (P22)	108	4.5	38.0
6	Bacterial sepsis of newborn (P36)	106	4.4	37.3
7	Neonatal hemorrhage (P50-P52,P54)	61	2.5	21.5
8	Necrotizing enterocolitis of newborn (P77)	57	2.4	20.1
9	Atelectasis (P28.0-P28.1)	51	2.1	18.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	47	2.0	16.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	34	1.4	12.0
12	Sudden infant death syndrome (R95)	30	1.2	10.6
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	20	0.8	7.0
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	16	0.7	*
15	Diseases of the circulatory system (I00-I99)	15	0.6	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	15	0.6	*
...	All other causes (Residual)	374	15.5	131.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,688	100.0	178.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	383	22.7	40.5
2	Sudden infant death syndrome (R95)	256	15.2	27.1
3	Accidents (unintentional injuries) (V01-X59)	145	8.6	15.3
4	Diseases of the circulatory system (I00-I99)	71	4.2	7.5
5	Assault (homicide) (*U01,X85-Y09)	63	3.7	6.7
6	Septicemia (A40-A41)	55	3.3	5.8
7	Influenza and pneumonia (J10-J18)	53	3.1	5.6
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	53	3.1	5.6
9	Chronic respiratory disease originating in the perinatal period (P27)	35	2.1	3.7
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	1.6	2.9
11	Malignant neoplasms (C00-C97)	18	1.1	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	1.1	*
11	Necrotizing enterocolitis of newborn (P77)	18	1.1	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	13	0.8	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.8	*
14	Bacterial sepsis of newborn (P36)	13	0.8	*
...	All other causes (Residual)	454	26.9	48.0
Postneonates, Hispanic, male				
...	All causes	933	100.0	193.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	18.3	35.4
2	Sudden infant death syndrome (R95)	153	16.4	31.7
3	Accidents (unintentional injuries) (V01-X59)	81	8.7	16.8
4	Diseases of the circulatory system (I00-I99)	38	4.1	7.9
5	Influenza and pneumonia (J10-J18)	37	4.0	7.7
5	Assault (homicide) (*U01,X85-Y09)	37	4.0	7.7
7	Septicemia (A40-A41)	36	3.9	7.5
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	28	3.0	5.8
9	Chronic respiratory disease originating in the perinatal period (P27)	23	2.5	4.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.9	*
11	Necrotizing enterocolitis of newborn (P77)	13	1.4	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	1.1	*
13	Bacterial sepsis of newborn (P36)	9	1.0	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	8	0.9	*
15	Malignant neoplasms (C00-C97)	7	0.8	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.8	*
...	All other causes (Residual)	257	27.5	53.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	755	100.0	162.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	212	28.1	45.7
2	Sudden infant death syndrome (R95)	103	13.6	22.2
3	Accidents (unintentional injuries) (V01-X59)	64	8.5	13.8
4	Diseases of the circulatory system (I00-I99)	33	4.4	7.1
5	Assault (homicide) (*U01,X85-Y09)	26	3.4	5.6
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	25	3.3	5.4
7	Septicemia (A40-A41)	19	2.5	*
8	Influenza and pneumonia (J10-J18)	16	2.1	*
9	Chronic respiratory disease originating in the perinatal period (P27)	12	1.6	*
10	Malignant neoplasms (C00-C97)	11	1.5	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.2	*
11	Respiratory distress of newborn (P22)	9	1.2	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	8	1.1	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	1.1	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	7	0.9	*
...	All other causes (Residual)	193	25.6	41.6
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,408	100.0	191.9
1	Sudden infant death syndrome (R95)	1,058	24.0	46.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	801	18.2	34.9
3	Accidents (unintentional injuries) (V01-X59)	483	11.0	21.0
4	Diseases of the circulatory system (I00-I99)	186	4.2	8.1
5	Assault (homicide) (*U01,X85-Y09)	122	2.8	5.3
6	Septicemia (A40-A41)	112	2.5	4.9
6	Influenza and pneumonia (J10-J18)	112	2.5	4.9
8	Chronic respiratory disease originating in the perinatal period (P27)	96	2.2	4.2
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	89	2.0	3.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	44	1.0	1.9
11	Malignant neoplasms (C00-C97)	32	0.7	1.4
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	31	0.7	1.3
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	27	0.6	1.2
14	Acute bronchitis and acute bronchiolitis (J20-J21)	26	0.6	1.1
15	Meningitis (G00,G03)	24	0.5	1.0
...	All other causes (Residual)	1,165	26.4	50.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, male				
...	All causes	2,533	100.0	215.0
1	Sudden infant death syndrome (R95)	628	24.8	53.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	401	15.8	34.0
3	Accidents (unintentional injuries) (V01-X59)	300	11.8	25.5
4	Diseases of the circulatory system (I00-I99)	103	4.1	8.7
5	Septicemia (A40-A41)	68	2.7	5.8
6	Assault (homicide) (*U01,X85-Y09)	66	2.6	5.6
7	Influenza and pneumonia (J10-J18)	65	2.6	5.5
8	Chronic respiratory disease originating in the perinatal period (P27)	62	2.4	5.3
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	55	2.2	4.7
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	30	1.2	2.5
11	Malignant neoplasms (C00-C97)	19	0.8	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	18	0.7	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	16	0.6	*
14	Meningitis (G00,G03)	15	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	15	0.6	*
14	Necrotizing enterocolitis of newborn (P77)	15	0.6	*
...	All other causes (Residual)	657	25.9	55.8
Postneonates, white, non-Hispanic, female				
...	All causes	1,875	100.0	167.6
1	Sudden infant death syndrome (R95)	430	22.9	38.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	400	21.3	35.8
3	Accidents (unintentional injuries) (V01-X59)	183	9.8	16.4
4	Diseases of the circulatory system (I00-I99)	83	4.4	7.4
5	Assault (homicide) (*U01,X85-Y09)	56	3.0	5.0
6	Influenza and pneumonia (J10-J18)	47	2.5	4.2
7	Septicemia (A40-A41)	44	2.3	3.9
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	34	1.8	3.0
8	Chronic respiratory disease originating in the perinatal period (P27)	34	1.8	3.0
10	Respiratory distress of newborn (P22)	15	0.8	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	0.7	*
12	Malignant neoplasms (C00-C97)	13	0.7	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	13	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.6	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	11	0.6	*
...	All other causes (Residual)	487	26.0	43.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,804	100.0	484.5
1	Sudden infant death syndrome (R95)	651	23.2	112.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	369	13.2	63.8
3	Accidents (unintentional injuries) (V01-X59)	276	9.8	47.7
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	123	4.4	21.3
5	Diseases of the circulatory system (I00-I99)	109	3.9	18.8
6	Chronic respiratory disease originating in the perinatal period (P27)	105	3.7	18.1
7	Influenza and pneumonia (J10-J18)	92	3.3	15.9
8	Septicemia (A40-A41)	90	3.2	15.6
9	Assault (homicide) (*U01,X85-Y09)	89	3.2	15.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	49	1.7	8.5
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	29	1.0	5.0
12	Necrotizing enterocolitis of newborn (P77)	26	0.9	4.5
13	Respiratory distress of newborn (P22)	21	0.7	3.6
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	17	0.6	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	15	0.5	*
15	Bacterial sepsis of newborn (P36)	15	0.5	*
...	All other causes (Residual)	728	26.0	125.8
Postneonates, black, non-Hispanic, male				
...	All causes	1,605	100.0	544.6
1	Sudden infant death syndrome (R95)	395	24.6	134.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	11.8	64.5
3	Accidents (unintentional injuries) (V01-X59)	150	9.3	50.9
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	77	4.8	26.1
5	Chronic respiratory disease originating in the perinatal period (P27)	59	3.7	20.0
6	Diseases of the circulatory system (I00-I99)	58	3.6	19.7
7	Influenza and pneumonia (J10-J18)	54	3.4	18.3
8	Septicemia (A40-A41)	51	3.2	17.3
9	Assault (homicide) (*U01,X85-Y09)	40	2.5	13.6
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	28	1.7	9.5
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	20	1.2	6.8
12	Necrotizing enterocolitis of newborn (P77)	15	0.9	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	11	0.7	*
13	Respiratory distress of newborn (P22)	11	0.7	*
15	Bacterial sepsis of newborn (P36)	10	0.6	*
...	All other causes (Residual)	436	27.2	147.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2004

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
	Postneonates, black, non-Hispanic, female			
...	All causes	1,199	100.0	422.1
1	Sudden infant death syndrome (R95)	256	21.4	90.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	14.9	63.0
3	Accidents (unintentional injuries) (V01-X59)	126	10.5	44.4
4	Diseases of the circulatory system (I00-I99)	51	4.3	18.0
5	Assault (homicide) (*U01,X85-Y09)	49	4.1	17.3
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	46	3.8	16.2
6	Chronic respiratory disease originating in the perinatal period (P27)	46	3.8	16.2
8	Septicemia (A40-A41)	39	3.3	13.7
9	Influenza and pneumonia (J10-J18)	38	3.2	13.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	1.8	7.4
11	Necrotizing enterocolitis of newborn (P77)	11	0.9	*
12	Respiratory distress of newborn (P22)	10	0.8	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	0.8	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	8	0.7	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.6	*
...	All other causes (Residual)	303	25.3	106.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable