

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

All causes

2007.....	760.2	905.6	643.4	749.4	890.5	634.8	958.0	1,184.4	793.8
2006.....	776.5	924.8	657.8	764.4	908.2	648.2	982.0	1,215.6	813.0
2005.....	798.8	951.1	677.6	785.3	933.2	666.5	1,016.5	1,252.9	845.7
2004.....	800.8	955.7	679.2	786.3	936.9	666.9	1,027.3	1,269.4	855.3
2003.....	832.7	994.3	706.2	817.0	973.9	693.1	1,065.9	1,319.1	885.6
2002.....	845.3	1,013.7	715.2	829.0	992.9	701.3	1,083.3	1,341.4	901.8
2001.....	854.5	1,029.1	721.8	836.5	1,006.1	706.7	1,101.2	1,375.0	912.5
2000.....	869.0	1,053.8	731.4	849.8	1,029.4	715.3	1,121.4	1,403.5	927.6
1999.....	875.6	1,067.0	734.0	854.6	1,040.0	716.6	1,135.7	1,432.6	933.6

Salmonella infections (A01-A02)

2007.....	0.0	*	*	0.0	*	*	*	*	*
2006.....	0.0	0.0	*	0.0	*	*	*	*	*
2005.....	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	0.0	*	0.0	*	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	0.0	0.0	0.0	*	*	*	*	*
2000.....	0.0	*	*	0.0	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	*	*	*	*	*

Shigellosis and amebiasis (A03,A06)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Certain other intestinal infections (A04,A07-A09)

2007.....	2.1	2.1	2.2	2.2	2.2	2.3	1.6	1.6	1.6
2006.....	2.1	2.1	2.1	2.2	2.2	2.2	1.4	1.4	1.4
2005.....	1.8	1.9	1.8	1.9	2.0	1.9	1.2	1.3	1.2
2004.....	1.5	1.4	1.5	1.5	1.5	1.6	0.9	0.9	0.9
2003.....	1.0	1.0	1.1	1.1	1.0	1.1	0.7	0.7	0.8
2002.....	0.8	0.8	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.6	0.5	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Both sexes		Male	Both sexes		Male	Female
	Female			Female		Male	Female		Male	Female

Scarlet fever and erysipelas (A38,A46)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Meningococcal infection (A39)

2007.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2000.....	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*

Septicemia (A40-A41)

2007.....	11.0	12.1	10.2	10.0	11.1	9.3	21.7	24.6	19.9
2006.....	11.0	12.1	10.2	10.1	11.2	9.3	21.6	24.2	19.9
2005.....	11.2	12.3	10.4	10.2	11.2	9.4	22.6	25.8	20.5
2004.....	11.2	12.3	10.3	10.2	11.3	9.4	22.5	25.7	20.4
2003.....	11.6	12.7	10.9	10.5	11.5	9.9	23.8	27.3	21.6
2002.....	11.7	12.9	10.9	10.7	11.7	9.9	24.2	27.8	22.1
2001.....	11.4	12.7	10.5	10.3	11.5	9.5	23.6	27.7	21.2
2000.....	11.3	12.4	10.5	10.2	11.2	9.4	24.6	28.0	22.5
1999.....	11.3	12.5	10.4	10.0	11.1	9.3	24.6	28.9	21.9

Syphilis (A50-A53)

2007.....	0.0	0.0	*	0.0	*	*	0.1	*	*
2006.....	0.0	0.0	*	*	*	*	0.1	*	*
2005.....	0.0	0.0	*	*	*	*	0.1	0.2	*
2004.....	0.0	0.0	0.0	*	*	*	0.1	*	*
2003.....	0.0	0.0	*	*	*	*	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.1	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	0.0	*	0.0	*	*	0.1	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Acute poliomyelitis (A80)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Arthropod-borne viral encephalitis (A83-A84,A85.2)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Measles (B05)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Viral hepatitis (B15-B19)

2007.....	2.3	3.1	1.5	2.2	3.0	1.4	2.9	4.2	1.9
2006.....	2.3	3.1	1.5	2.1	2.9	1.4	3.2	4.5	2.1
2005.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.4	1.9
2004.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.8	1.7
2003.....	1.8	2.5	1.2	1.7	2.4	1.1	2.7	3.8	1.8
2002.....	2.0	2.7	1.3	1.9	2.6	1.2	2.8	3.8	1.9
2001.....	2.0	2.6	1.3	1.8	2.5	1.2	2.7	3.4	2.1
2000.....	1.9	2.6	1.3	1.8	2.4	1.2	2.7	3.8	1.8
1999.....	1.8	2.4	1.2	1.6	2.2	1.1	2.6	3.4	1.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Human immunodeficiency virus (HIV) disease (B20-B24)

2007.....	3.7	5.4	2.1	1.9	3.1	0.7	17.3	24.5	11.3
2006.....	4.0	5.9	2.2	2.1	3.4	0.7	18.6	26.3	12.2
2005.....	4.2	6.2	2.3	2.2	3.6	0.8	19.4	28.2	12.0
2004.....	4.5	6.6	2.4	2.3	3.8	0.9	20.4	29.2	13.0
2003.....	4.7	7.1	2.4	2.5	4.2	0.9	21.3	31.3	12.8
2002.....	4.9	7.4	2.5	2.6	4.3	0.9	22.5	33.3	13.4
2001.....	5.0	7.5	2.5	2.6	4.4	0.9	22.8	33.8	13.4
2000.....	5.2	7.9	2.5	2.8	4.6	1.0	23.3	35.1	13.2
1999.....	5.3	8.2	2.5	2.9	4.9	1.0	23.6	36.1	13.1

Malaria (B50-B54)

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

2007.....	1.8	2.2	1.6	1.8	2.1	1.5	2.4	2.9	2.0
2006.....	1.9	2.3	1.6	1.8	2.2	1.6	2.6	3.1	2.2
2005.....	2.5	3.1	2.0	2.5	3.0	1.9	3.1	3.9	2.5
2004.....	2.5	3.1	2.0	2.4	3.0	2.0	3.2	4.1	2.6
2003.....	2.5	3.1	2.1	2.4	3.0	2.0	3.2	4.0	2.6
2002.....	2.3	2.9	1.9	2.2	2.7	1.8	3.2	4.1	2.5
2001.....	2.0	2.4	1.7	1.9	2.3	1.6	2.8	3.4	2.4
2000.....	2.0	2.4	1.7	1.9	2.3	1.6	2.9	3.7	2.4
1999.....	2.6	3.1	2.3	2.4	2.9	2.1	4.6	5.5	4.0

Malignant neoplasms (C00-C97)

2007.....	178.4	217.5	151.3	177.5	215.1	151.2	215.5	282.3	174.9
2006.....	180.7	220.1	153.6	179.9	217.9	153.6	217.4	284.9	176.1
2005.....	183.8	225.1	155.6	182.6	222.3	155.2	222.7	293.7	179.6
2004.....	185.8	227.7	157.4	184.4	224.4	157.0	227.2	301.2	182.5
2003.....	190.1	233.3	160.9	188.5	230.1	160.2	233.3	308.8	187.7
2002.....	193.5	238.9	163.1	191.7	235.2	162.4	238.8	319.6	190.3
2001.....	196.0	243.7	164.7	193.9	239.2	163.9	243.1	330.9	191.3
2000.....	199.6	248.9	167.6	197.2	243.9	166.9	248.5	340.3	193.8
1999.....	200.8	251.9	167.6	197.9	246.5	166.4	252.5	346.1	196.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

2007.....	2.5	3.9	1.4	2.5	3.7	1.4	3.3	5.7	1.6
2006.....	2.5	3.8	1.4	2.4	3.6	1.4	3.2	5.7	1.5
2005.....	2.5	3.9	1.4	2.4	3.6	1.4	3.5	6.5	1.4
2004.....	2.6	4.0	1.5	2.5	3.7	1.5	3.8	6.9	1.6
2003.....	2.6	4.0	1.5	2.5	3.8	1.4	3.8	6.9	1.6
2002.....	2.7	4.1	1.5	2.5	3.8	1.5	3.6	6.3	1.7
2001.....	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2000.....	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
1999.....	2.7	4.1	1.6	2.5	3.8	1.5	4.3	7.5	2.1

Malignant neoplasm of esophagus (C15)

2007.....	4.3	7.7	1.6	4.3	7.8	1.5	4.7	8.3	2.2
2006.....	4.4	7.8	1.7	4.5	7.9	1.6	4.8	8.3	2.3
2005.....	4.4	7.9	1.7	4.5	8.0	1.6	5.0	8.8	2.3
2004.....	4.3	7.7	1.7	4.3	7.7	1.6	5.4	9.0	2.8
2003.....	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9
2002.....	4.4	7.8	1.7	4.3	7.7	1.6	5.9	10.1	2.9
2001.....	4.4	7.8	1.8	4.3	7.6	1.7	6.4	11.3	3.0
2000.....	4.4	7.7	1.8	4.3	7.6	1.7	6.5	11.0	3.3
1999.....	4.4	7.7	1.8	4.2	7.4	1.6	6.5	11.5	3.1

Malignant neoplasm of stomach (C16)

2007.....	3.6	5.0	2.6	3.1	4.4	2.2	6.9	10.4	4.6
2006.....	3.7	5.0	2.7	3.2	4.4	2.3	6.8	9.8	4.9
2005.....	3.8	5.2	2.7	3.3	4.5	2.4	7.2	10.6	4.9
2004.....	4.0	5.5	2.8	3.5	4.9	2.5	7.5	11.1	5.2
2003.....	4.1	5.7	3.0	3.6	5.0	2.6	7.8	11.3	5.4
2002.....	4.2	5.9	3.0	3.7	5.2	2.5	8.4	11.9	6.1
2001.....	4.4	6.1	3.1	3.8	5.4	2.7	8.3	11.9	5.8
2000.....	4.6	6.4	3.3	4.0	5.6	2.8	9.0	12.7	6.5
1999.....	4.6	6.6	3.3	4.1	5.8	2.8	9.2	13.5	6.4

Malignant neoplasms of colon, rectum and anus (C18-C21)

2007.....	16.9	20.1	14.4	16.4	19.6	13.9	23.5	29.0	19.8
2006.....	17.2	20.5	14.7	16.7	19.9	14.3	24.3	30.6	20.2
2005.....	17.5	20.9	14.8	16.9	20.3	14.3	24.8	30.2	21.2
2004.....	18.0	21.5	15.3	17.6	21.0	14.9	24.7	30.0	21.2
2003.....	19.1	22.9	16.2	18.6	22.4	15.7	26.4	31.9	22.8
2002.....	19.7	23.7	16.7	19.2	23.1	16.2	26.8	33.1	22.8
2001.....	20.1	24.2	17.2	19.6	23.7	16.6	27.6	33.1	24.1
2000.....	20.8	25.1	17.7	20.3	24.5	17.2	28.2	34.9	24.0
1999.....	20.9	25.3	17.8	20.4	24.8	17.3	28.4	34.2	24.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

2007.....	5.4	7.9	3.2	5.0	7.2	3.1	7.1	11.5	3.8
2006.....	5.3	7.7	3.3	4.9	7.0	3.1	7.0	11.1	3.9
2005.....	5.2	7.7	3.2	4.8	7.1	3.0	7.0	11.3	3.8
2004.....	5.1	7.4	3.2	4.7	6.7	3.0	6.9	10.7	4.1
2003.....	5.0	7.4	3.1	4.6	6.7	2.8	6.6	10.2	4.0
2002.....	4.9	7.1	3.0	4.5	6.5	2.8	6.3	9.9	3.6
2001.....	4.7	6.9	3.0	4.3	6.3	2.7	6.2	9.1	4.1
2000.....	4.7	6.8	3.0	4.2	6.2	2.7	6.1	9.5	3.6
1999.....	4.5	6.5	3.0	4.2	5.9	2.7	5.9	8.7	3.8

Malignant neoplasm of pancreas (C25)

2007.....	10.8	12.5	9.4	10.6	12.4	9.2	14.0	15.5	12.7
2006.....	10.8	12.3	9.5	10.6	12.3	9.3	13.6	14.8	12.5
2005.....	10.8	12.3	9.5	10.6	12.2	9.3	13.6	15.3	12.3
2004.....	10.6	12.3	9.3	10.4	12.2	9.1	13.6	15.4	12.1
2003.....	10.5	12.0	9.3	10.3	11.9	9.0	13.7	15.3	12.4
2002.....	10.5	12.1	9.2	10.3	12.0	9.0	13.6	15.3	12.3
2001.....	10.5	12.1	9.3	10.4	12.0	9.1	13.6	15.0	12.5
2000.....	10.6	12.2	9.3	10.4	12.0	9.0	14.1	15.8	12.7
1999.....	10.6	12.3	9.3	10.4	12.0	9.1	14.1	16.4	12.3

Malignant neoplasm of larynx (C32)

2007.....	1.1	2.1	0.4	1.1	1.9	0.4	2.1	4.3	0.6
2006.....	1.2	2.2	0.5	1.2	2.0	0.5	2.1	4.1	0.7
2005.....	1.2	2.2	0.5	1.2	2.1	0.5	2.3	4.6	0.8
2004.....	1.2	2.2	0.4	1.1	2.1	0.4	2.2	4.6	0.7
2003.....	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.1	0.7
2002.....	1.3	2.3	0.5	1.2	2.1	0.5	2.5	4.8	0.8
2001.....	1.3	2.5	0.5	1.2	2.3	0.5	2.5	5.0	0.9
2000.....	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	0.9
1999.....	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.8

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

2007.....	50.6	65.1	40.0	51.2	64.8	41.2	55.6	82.2	38.1
2006.....	51.5	67.0	40.0	52.1	66.8	41.1	56.8	83.7	39.0
2005.....	52.6	69.0	40.5	53.1	68.7	41.5	58.4	86.4	40.0
2004.....	53.2	70.1	40.9	53.6	69.4	41.9	59.8	90.0	39.9
2003.....	54.1	71.7	41.3	54.5	71.1	42.3	60.8	92.4	40.2
2002.....	54.9	73.2	41.6	55.3	72.5	42.6	61.9	95.0	40.1
2001.....	55.3	75.2	41.0	55.6	74.2	42.1	62.5	99.1	38.7
2000.....	56.1	76.7	41.3	56.2	75.7	42.3	64.0	101.1	39.8
1999.....	55.5	76.9	40.2	55.4	75.6	41.0	64.8	103.0	39.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Malignant melanoma of skin (C43)

2007.....	2.7	4.0	1.7	3.0	4.5	1.9	0.5	0.5	0.4
2006.....	2.7	4.0	1.7	3.1	4.6	2.0	0.4	0.6	0.3
2005.....	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.4
2004.....	2.7	3.9	1.7	3.0	4.5	1.9	0.4	0.5	0.4
2003.....	2.7	3.9	1.8	3.0	4.4	2.0	0.4	0.4	0.4
2002.....	2.6	3.8	1.7	2.9	4.2	2.0	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	3.0	4.3	2.0	0.4	0.3	0.4
2000.....	2.7	3.8	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3

Malignant neoplasm of breast (C50)

2007.....	12.9	0.3	22.9	12.5	0.3	22.3	18.6	0.4	31.4
2006.....	13.2	0.3	23.5	12.8	0.3	22.9	18.7	0.5	31.6
2005.....	13.5	0.3	24.1	13.1	0.3	23.4	19.3	0.5	32.8
2004.....	13.8	0.3	24.4	13.4	0.3	23.9	19.0	0.5	32.2
2003.....	14.2	0.3	25.3	13.8	0.3	24.7	20.1	0.5	34.0
2002.....	14.5	0.3	25.6	14.1	0.3	25.0	20.2	0.6	34.0
2001.....	14.8	0.3	26.0	14.4	0.3	25.5	20.4	0.6	34.4
2000.....	15.3	0.4	26.8	14.9	0.4	26.3	20.5	0.6	34.5
1999.....	15.2	0.3	26.6	14.8	0.3	26.0	20.8	0.7	35.1

Malignant neoplasm of cervix uteri (C53)

2007.....	1.3	...	2.4	1.2	...	2.2	2.5	...	4.3
2006.....	1.3	...	2.4	1.1	...	2.2	2.5	...	4.3
2005.....	1.3	...	2.4	1.1	...	2.2	2.5	...	4.4
2004.....	1.3	...	2.4	1.1	...	2.2	2.6	...	4.5
2003.....	1.3	...	2.5	1.2	...	2.2	2.7	...	4.7
2002.....	1.3	...	2.6	1.2	...	2.3	2.9	...	4.9
2001.....	1.4	...	2.7	1.3	...	2.5	2.8	...	4.8
2000.....	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
1999.....	1.5	...	2.8	1.4	...	2.6	3.1	...	5.4

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

2007.....	2.3	...	4.2	2.2	...	3.9	4.5	...	7.5
2006.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3
2005.....	2.3	...	4.1	2.2	...	3.8	4.2	...	7.1
2004.....	2.3	...	4.1	2.2	...	3.9	4.1	...	6.9
2003.....	2.3	...	4.1	2.2	...	3.9	4.2	...	7.0
2002.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3
2001.....	2.4	...	4.2	2.3	...	4.0	4.2	...	7.0
2000.....	2.4	...	4.1	2.2	...	3.9	4.4	...	7.2
1999.....	2.4	...	4.1	2.2	...	3.9	4.2	...	6.8

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Malignant neoplasm of ovary (C56)

2007.....	4.6	...	8.2	4.8	...	8.6	3.9	...	6.5
2006.....	4.8	...	8.5	5.0	...	8.9	4.1	...	6.8
2005.....	4.8	...	8.6	5.0	...	9.0	4.3	...	7.2
2004.....	4.9	...	8.8	5.1	...	9.1	4.5	...	7.6
2003.....	5.0	...	8.8	5.1	...	9.2	4.4	...	7.4
2002.....	5.1	...	9.0	5.3	...	9.4	4.4	...	7.4
2001.....	5.1	...	9.0	5.2	...	9.3	4.5	...	7.5
2000.....	5.1	...	8.9	5.3	...	9.3	4.3	...	7.2
1999.....	5.0	...	8.8	5.1	...	9.1	4.4	...	7.3

Malignant neoplasm of prostate (C61)

2007.....	9.2	23.5	...	8.5	21.6	...	18.6	51.5	...
2006.....	9.2	23.5	...	8.5	21.7	...	18.2	50.5	...
2005.....	9.5	24.5	...	8.8	22.6	...	19.1	53.3	...
2004.....	9.8	25.4	...	9.0	23.4	...	19.7	55.5	...
2003.....	10.1	26.5	...	9.3	24.4	...	20.4	57.4	...
2002.....	10.6	27.9	...	9.8	25.7	...	22.0	62.0	...
2001.....	10.9	29.1	...	10.0	26.6	...	23.0	66.1	...
2000.....	11.3	30.4	...	10.3	27.8	...	23.9	68.1	...
1999.....	11.6	31.3	...	10.7	28.7	...	24.2	69.0	...

Malignant neoplasms of kidney and renal pelvis (C64-C65)

2007.....	4.0	5.8	2.6	4.1	5.9	2.7	4.1	5.9	2.7
2006.....	4.0	5.7	2.6	4.1	5.8	2.7	3.9	5.8	2.5
2005.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	6.0	2.6
2004.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	5.8	2.7
2003.....	4.2	6.0	2.7	4.3	6.2	2.8	4.1	6.0	2.7
2002.....	4.2	6.1	2.8	4.3	6.3	2.8	4.0	5.8	2.8
2001.....	4.3	6.2	2.8	4.3	6.3	2.8	4.2	6.4	2.8
2000.....	4.2	6.2	2.8	4.3	6.3	2.8	4.1	6.3	2.8
1999.....	4.1	5.9	2.7	4.1	5.9	2.7	4.1	6.1	2.8

Malignant neoplasm of bladder (C67)

2007.....	4.4	7.6	2.2	4.6	8.0	2.2	3.8	5.4	2.8
2006.....	4.3	7.5	2.2	4.5	7.9	2.2	3.8	5.5	2.8
2005.....	4.3	7.5	2.2	4.5	7.9	2.2	3.7	5.7	2.6
2004.....	4.4	7.6	2.2	4.5	8.0	2.3	3.6	4.9	2.7
2003.....	4.3	7.3	2.2	4.4	7.7	2.2	3.7	5.4	2.7
2002.....	4.4	7.5	2.4	4.5	7.8	2.4	3.9	5.5	3.0
2001.....	4.3	7.5	2.3	4.5	7.9	2.2	3.7	5.1	2.9
2000.....	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.6	2.7
1999.....	4.4	7.6	2.3	4.5	7.9	2.3	4.0	5.8	3.0

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

2007.....	4.2	5.1	3.5	4.6	5.5	3.8	2.4	2.8	2.1
2006.....	4.2	5.1	3.4	4.5	5.5	3.6	2.3	2.9	1.9
2005.....	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.2	2.0
2004.....	4.3	5.2	3.5	4.7	5.6	3.8	2.5	3.0	2.0
2003.....	4.4	5.3	3.6	4.7	5.7	3.9	2.7	3.4	2.1
2002.....	4.4	5.5	3.6	4.8	5.8	3.9	2.6	3.2	2.2
2001.....	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.1	2.2
2000.....	4.5	5.6	3.7	4.9	6.0	3.9	2.7	3.3	2.3
1999.....	4.6	5.6	3.8	5.0	6.0	4.1	2.7	3.3	2.3

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

2007.....	17.6	22.8	13.7	17.8	23.2	13.8	17.6	21.9	14.8
2006.....	17.9	23.0	14.1	18.2	23.4	14.3	18.0	23.6	14.3
2005.....	18.2	23.5	14.3	18.5	24.0	14.5	17.6	22.5	14.3
2004.....	18.4	23.6	14.6	18.6	24.0	14.7	18.9	23.6	15.8
2003.....	19.1	24.5	15.1	19.4	25.1	15.2	18.8	22.9	16.1
2002.....	19.5	25.1	15.5	19.9	25.6	15.7	19.4	24.4	16.1
2001.....	20.0	25.5	16.0	20.2	25.9	16.2	20.0	25.3	16.5
2000.....	20.4	26.0	16.5	20.7	26.4	16.7	19.9	25.0	16.6
1999.....	20.6	26.1	16.7	20.8	26.5	16.8	20.3	25.6	16.8

Hodgkin's disease (C81)

2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3
2006.....	0.4	0.6	0.3	0.4	0.6	0.4	0.4	0.6	0.3
2005.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
2004.....	0.4	0.5	0.3	0.4	0.6	0.3	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.5	0.3

Non-Hodgkin's lymphoma (C82-C85)

2007.....	6.5	8.3	5.2	6.8	8.6	5.5	4.5	5.7	3.6
2006.....	6.7	8.4	5.4	7.0	8.7	5.6	4.9	6.2	3.9
2005.....	6.9	8.7	5.5	7.2	9.1	5.8	4.7	6.0	3.8
2004.....	7.0	8.8	5.7	7.3	9.2	6.0	5.0	6.2	4.1
2003.....	7.3	9.3	5.9	7.7	9.7	6.1	4.9	5.8	4.2
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001.....	7.9	9.9	6.4	8.2	10.2	6.7	5.6	7.2	4.5
2000.....	8.2	10.2	6.7	8.6	10.7	7.1	5.2	6.6	4.2
1999.....	8.3	10.3	6.9	8.7	10.7	7.2	5.6	7.4	4.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Leukemia (C91-C95)

2007.....	7.0	9.4	5.3	7.2	9.7	5.4	6.1	7.9	5.0
2006.....	7.1	9.5	5.4	7.4	9.8	5.6	6.1	8.6	4.6
2005.....	7.1	9.6	5.4	7.4	10.0	5.6	5.9	7.8	4.7
2004.....	7.2	9.6	5.5	7.4	9.9	5.6	6.7	8.7	5.3
2003.....	7.4	9.9	5.6	7.6	10.2	5.7	6.4	8.3	5.2
2002.....	7.5	10.0	5.7	7.7	10.3	5.8	6.6	8.6	5.3
2001.....	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.7	5.4
2000.....	7.7	10.3	5.9	7.9	10.5	6.1	6.7	9.0	5.2
1999.....	7.6	10.2	6.0	7.8	10.4	6.1	6.7	8.7	5.4

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

2007.....	3.6	4.6	2.9	3.3	4.4	2.6	6.6	7.8	5.9
2006.....	3.6	4.5	2.9	3.4	4.3	2.7	6.5	8.2	5.5
2005.....	3.7	4.7	3.0	3.5	4.4	2.8	6.5	8.1	5.5
2004.....	3.7	4.6	3.1	3.4	4.3	2.8	6.9	8.3	6.0
2003.....	3.8	4.8	3.2	3.6	4.6	2.9	7.2	8.4	6.3
2002.....	4.0	5.0	3.3	3.7	4.8	3.0	6.9	8.3	6.1
2001.....	4.0	4.9	3.3	3.7	4.6	3.0	7.3	8.8	6.4
2000.....	4.0	4.9	3.4	3.7	4.6	3.1	7.7	8.9	6.8
1999.....	4.0	4.9	3.4	3.7	4.7	3.1	7.5	8.9	6.7

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

2007.....	20.0	24.3	16.8	20.0	24.5	16.7	22.0	26.8	18.8
2006.....	20.3	24.5	17.3	20.4	24.6	17.3	22.6	27.4	19.4
2005.....	20.6	24.8	17.5	20.7	24.9	17.5	23.2	28.4	19.6
2004.....	20.8	25.0	17.6	20.7	25.0	17.6	23.9	29.6	19.9
2003.....	21.4	25.7	18.2	21.4	25.7	18.2	24.3	30.0	20.5
2002.....	21.8	26.5	18.5	21.8	26.4	18.4	25.7	31.4	21.9
2001.....	22.2	26.7	18.9	22.0	26.5	18.8	26.2	32.8	21.9
2000.....	22.8	27.5	19.4	22.6	27.3	19.3	26.4	33.0	22.1
1999.....	24.2	29.3	20.6	24.0	29.1	20.4	28.5	35.2	23.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

2007.....	4.5	5.6	3.7	4.6	5.8	3.8	4.1	5.2	3.4
2006.....	4.5	5.6	3.9	4.6	5.7	3.9	4.3	5.3	3.8
2005.....	4.5	5.5	3.8	4.6	5.7	3.8	4.1	4.6	3.9
2004.....	4.5	5.7	3.8	4.6	5.8	3.9	4.4	5.2	3.9
2003.....	4.6	5.7	3.9	4.7	5.9	4.0	4.3	5.1	3.7
2002.....	4.6	5.7	3.9	4.7	5.8	4.0	4.4	5.1	4.1
2001.....	4.9	6.0	4.1	4.9	6.2	4.1	4.6	5.6	4.1
2000.....	4.9	6.0	4.2	5.0	6.1	4.3	4.6	5.4	4.1
1999.....	4.9	6.1	4.1	4.9	6.2	4.1	4.8	5.4	4.4

Anemias (D50-D64)

2007.....	1.5	1.5	1.5	1.3	1.3	1.3	3.0	3.1	2.9
2006.....	1.3	1.3	1.3	1.1	1.1	1.1	2.9	2.8	2.8
2005.....	1.5	1.5	1.5	1.3	1.2	1.3	3.3	3.4	3.2
2004.....	1.5	1.5	1.5	1.3	1.3	1.3	3.3	3.5	3.1
2003.....	1.6	1.6	1.5	1.3	1.3	1.3	3.5	3.9	3.2
2002.....	1.6	1.6	1.6	1.3	1.3	1.4	3.7	3.9	3.5
2001.....	1.6	1.6	1.6	1.4	1.4	1.4	3.5	3.6	3.4
2000.....	1.6	1.6	1.6	1.4	1.4	1.4	3.4	3.4	3.5
1999.....	1.6	1.7	1.6	1.4	1.4	1.4	3.6	3.7	3.5

Diabetes mellitus (E10-E14)

2007.....	22.5	26.4	19.5	20.5	24.6	17.2	42.8	45.9	40.2
2006.....	23.3	27.4	20.1	21.2	25.4	17.9	45.1	49.7	41.6
2005.....	24.6	28.4	21.6	22.5	26.5	19.3	46.9	50.8	43.8
2004.....	24.5	28.2	21.7	22.3	26.2	19.2	48.0	51.3	45.3
2003.....	25.3	28.9	22.5	23.0	27.0	19.9	49.2	50.7	47.5
2002.....	25.4	28.6	23.0	23.1	26.8	20.3	49.5	49.4	48.6
2001.....	25.3	28.1	23.1	23.0	26.2	20.5	49.2	49.9	48.1
2000.....	25.0	27.8	23.0	22.8	26.0	20.4	49.5	48.7	49.4
1999.....	25.0	27.8	23.0	22.6	25.8	20.2	49.7	49.4	49.5

Nutritional deficiencies (E40-E64)

2007.....	0.9	0.9	0.9	0.8	0.8	0.8	1.5	1.6	1.3
2006.....	0.8	0.7	0.8	0.8	0.7	0.8	1.3	1.5	1.1
2005.....	1.0	1.0	1.1	1.0	0.9	1.0	1.7	1.6	1.7
2004.....	1.0	0.9	1.0	0.9	0.9	1.0	1.7	1.9	1.5
2003.....	1.1	1.1	1.2	1.1	1.0	1.1	1.9	2.1	1.7
2002.....	1.3	1.2	1.3	1.2	1.2	1.3	2.1	2.1	2.1
2001.....	1.3	1.2	1.4	1.2	1.1	1.3	2.1	2.4	1.8
2000.....	1.5	1.4	1.6	1.5	1.3	1.5	2.4	2.6	2.2
1999.....	1.6	1.5	1.6	1.5	1.4	1.5	2.7	3.1	2.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Malnutrition (E40-E46)

2007.....	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.6	1.3
2006.....	0.7	0.7	0.8	0.7	0.6	0.7	1.2	1.5	1.0
2005.....	1.0	0.9	1.0	0.9	0.8	1.0	1.6	1.5	1.6
2004.....	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.1	1.0	1.1	1.0	0.9	1.1	1.8	2.1	1.7
2002.....	1.2	1.1	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.4	1.2	1.4	2.2	2.5	2.1
1999.....	1.5	1.4	1.5	1.4	1.3	1.4	2.5	2.9	2.2

Other nutritional deficiencies (E50-E64)

2007.....	0.1	0.1	0.1	0.1	0.1	0.0	*	*	*
2006.....	0.1	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004.....	0.1	0.0	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2

Meningitis (G00,G03)

2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2004.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.5	0.6
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.5	0.7	0.4
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.8	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6

Parkinson's disease (G20-G21)

2007.....	6.4	9.6	4.3	6.8	10.2	4.6	3.2	4.9	2.2
2006.....	6.3	9.6	4.3	6.7	10.2	4.6	2.7	4.3	1.9
2005.....	6.4	9.8	4.4	6.8	10.4	4.6	3.0	4.9	1.9
2004.....	6.1	9.3	4.1	6.4	9.8	4.4	2.7	4.3	1.9
2003.....	6.2	9.3	4.2	6.5	9.8	4.5	2.7	4.4	1.8
2002.....	5.9	9.0	4.1	6.3	9.5	4.3	2.5	4.0	1.7
2001.....	5.9	9.1	4.0	6.2	9.7	4.3	2.5	3.9	1.8
2000.....	5.7	8.8	3.9	6.0	9.3	4.2	2.5	4.2	1.6
1999.....	5.4	8.4	3.7	5.7	8.9	3.9	2.4	3.7	1.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Alzheimer's disease (G30)

2007.....	22.7	18.5	24.9	23.5	19.1	25.9	19.0	15.6	20.6
2006.....	22.6	18.4	24.7	23.4	19.1	25.7	18.3	14.7	19.8
2005.....	22.9	18.5	25.1	23.7	19.0	26.1	19.4	16.4	20.7
2004.....	21.8	17.7	23.8	22.5	18.3	24.7	18.5	15.2	19.9
2003.....	21.4	17.5	23.3	22.2	18.2	24.3	16.6	14.7	17.3
2002.....	20.2	16.6	22.0	21.1	17.3	22.9	15.2	13.2	15.9
2001.....	19.1	16.1	20.5	19.8	16.7	21.3	14.6	13.1	15.1
2000.....	18.1	15.2	19.3	18.8	15.8	20.1	13.0	11.4	13.7
1999.....	16.5	14.4	17.4	17.2	15.0	18.1	11.3	10.2	11.8

Major cardiovascular diseases (I00-I78)

2007.....	249.9	298.9	210.4	244.1	292.7	204.6	335.0	403.5	284.2
2006.....	261.2	312.0	220.4	255.1	305.4	214.3	348.7	420.4	296.0
2005.....	277.3	329.5	235.6	271.0	323.2	229.0	367.9	435.3	317.4
2004.....	286.5	340.0	243.8	279.8	333.6	236.7	381.9	451.1	331.0
2003.....	306.1	363.4	260.7	298.9	356.4	253.0	406.1	479.1	352.4
2002.....	317.4	377.2	270.5	310.4	370.7	262.7	416.1	487.4	363.8
2001.....	326.5	388.1	278.8	319.2	381.1	270.8	427.5	505.3	372.4
2000.....	339.7	407.3	288.4	332.5	400.7	280.3	438.7	517.6	383.3
1999.....	349.3	419.0	296.5	341.9	411.8	288.4	447.6	532.2	388.6

Diseases of heart (I00-I09,I11,I13,I20-I51)

2007.....	190.9	237.7	154.0	187.8	234.8	150.5	247.3	305.9	204.5
2006.....	200.2	248.5	162.2	197.0	245.2	158.6	257.7	320.6	212.5
2005.....	211.1	260.9	172.3	207.8	258.0	168.2	271.3	329.8	228.3
2004.....	217.0	267.9	177.3	213.3	264.6	172.9	280.6	342.1	236.5
2003.....	232.3	286.6	190.3	228.2	282.9	185.4	300.2	364.3	253.8
2002.....	240.8	297.4	197.2	236.7	294.1	192.1	308.4	371.0	263.2
2001.....	247.8	305.4	203.9	243.5	301.8	198.7	316.9	384.5	269.8
2000.....	257.6	320.0	210.9	253.4	316.7	205.6	324.8	392.5	277.6
1999.....	266.5	331.0	218.1	261.9	327.1	212.8	334.3	407.2	283.7

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

2007.....	1.0	0.8	1.2	1.0	0.8	1.2	0.8	0.7	0.9
2006.....	1.1	0.8	1.2	1.1	0.8	1.3	0.8	0.7	0.9
2005.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.7	0.8
2004.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003.....	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2002.....	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.5	1.2	0.9	1.5	1.1	0.7	1.3
2000.....	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.8	1.2

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Hypertensive heart disease (I11)

2007.....	9.5	10.3	8.4	8.1	8.6	7.2	23.3	28.4	19.3
2006.....	9.4	10.1	8.3	8.0	8.4	7.2	23.1	28.5	18.8
2005.....	9.4	10.1	8.5	7.9	8.3	7.2	24.1	28.4	20.4
2004.....	9.5	10.1	8.5	8.0	8.4	7.2	24.2	28.4	20.6
2003.....	9.6	10.1	8.8	8.0	8.3	7.3	25.0	28.7	21.9
2002.....	9.1	9.5	8.3	7.6	7.8	7.0	24.4	28.0	21.2
2001.....	8.7	8.9	8.1	7.3	7.3	6.8	23.2	26.3	20.5
2000.....	8.6	8.7	8.1	7.1	7.1	6.7	23.7	26.3	21.2
1999.....	8.3	8.3	7.9	6.9	6.7	6.6	22.9	25.7	20.5

Hypertensive heart and renal disease (I13)

2007.....	0.9	1.0	0.9	0.7	0.8	0.7	2.8	3.4	2.4
2006.....	0.9	1.0	0.8	0.7	0.8	0.7	2.9	3.5	2.5
2005.....	1.0	1.1	0.9	0.8	0.9	0.8	3.2	3.8	2.8
2004.....	1.0	1.1	0.9	0.8	0.8	0.7	3.2	3.6	2.9
2003.....	1.1	1.1	1.0	0.8	0.9	0.7	3.4	3.7	3.1
2002.....	1.0	1.0	0.9	0.7	0.8	0.7	3.5	3.9	3.2
2001.....	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.6	3.0
2000.....	1.0	1.1	0.9	0.8	0.9	0.7	3.3	3.8	3.1
1999.....	1.2	1.4	1.1	0.9	1.0	0.9	4.2	4.5	3.9

Ischemic heart diseases (I20-I25)

2007.....	126.0	165.4	95.7	125.5	165.6	94.2	150.6	191.6	121.5
2006.....	134.9	176.5	103.1	134.2	176.3	101.5	161.6	206.4	130.0
2005.....	144.4	187.4	111.7	143.8	187.7	110.0	171.3	213.9	140.9
2004.....	150.2	194.1	116.7	149.2	194.2	114.7	179.8	223.9	148.7
2003.....	162.9	209.9	127.2	161.7	209.6	125.0	195.0	243.3	160.9
2002.....	170.8	220.4	133.6	169.8	220.5	131.2	203.0	250.6	169.7
2001.....	177.8	228.5	139.9	176.5	228.4	137.4	211.6	262.0	176.7
2000.....	186.8	241.4	146.5	185.6	241.6	143.9	218.3	269.5	183.2
1999.....	194.6	251.2	152.9	193.4	251.2	150.5	224.9	278.7	188.0

Acute myocardial infarction (I21-I22)

2007.....	41.4	54.0	31.5	41.3	54.3	30.9	49.5	60.9	41.2
2006.....	45.0	58.8	34.2	45.0	59.2	33.7	53.3	66.3	44.0
2005.....	49.1	63.6	37.8	49.1	64.1	37.2	57.6	69.7	48.8
2004.....	52.2	67.8	40.2	52.2	68.3	39.5	62.0	74.8	52.7
2003.....	57.9	75.0	44.8	57.8	75.3	44.0	68.6	84.3	57.6
2002.....	62.1	80.6	48.1	61.9	81.0	47.1	73.9	89.5	63.2
2001.....	65.4	84.9	50.9	65.1	85.4	49.9	76.8	93.1	66.0
2000.....	69.9	91.3	54.1	69.6	91.8	53.1	81.0	98.9	69.2
1999.....	73.2	96.1	56.3	73.1	96.8	55.4	83.1	102.1	70.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Other acute ischemic heart diseases (I24)

2007.....	1.3	1.6	1.0	1.2	1.5	1.0	1.8	2.4	1.4
2006.....	1.2	1.5	1.0	1.2	1.5	1.0	1.8	2.4	1.4
2005.....	1.2	1.4	0.9	1.1	1.3	0.9	2.0	2.6	1.5
2004.....	0.8	1.0	0.6	0.8	1.0	0.6	1.4	2.1	1.0
2003.....	1.1	1.3	0.9	1.0	1.2	0.8	1.6	2.4	1.1
2002.....	1.2	1.5	0.9	1.1	1.4	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.3	1.3
2000.....	1.2	1.5	0.9	1.2	1.5	0.9	1.8	2.5	1.3
1999.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.4	1.3

Other forms of chronic ischemic heart disease (I20,I25)

2007.....	83.3	109.8	63.2	82.9	109.8	62.3	99.3	128.2	78.9
2006.....	88.7	116.1	67.9	88.0	115.7	66.9	106.5	137.7	84.6
2005.....	94.2	122.4	73.0	93.6	122.3	72.0	111.8	141.6	90.5
2004.....	97.1	125.3	75.8	96.3	125.0	74.6	116.4	146.9	95.0
2003.....	103.8	133.5	81.5	102.9	133.0	80.1	124.8	156.7	102.2
2002.....	107.6	138.3	84.5	106.8	138.0	83.2	127.5	158.9	105.2
2001.....	111.2	142.1	88.1	110.2	141.6	86.7	133.0	166.7	109.3
2000.....	115.7	148.6	91.5	114.8	148.3	90.0	135.4	168.1	112.6
1999.....	120.2	153.6	95.6	119.2	153.0	94.2	140.0	174.2	116.2

Atherosclerotic cardiovascular disease, so described (I25.0)

2007.....	18.2	24.3	13.1	17.3	23.0	12.3	30.3	42.3	21.6
2006.....	19.3	25.6	14.0	18.1	24.0	13.1	32.7	46.0	23.3
2005.....	20.3	26.6	15.0	19.1	25.0	14.0	34.5	46.9	25.3
2004.....	21.1	27.5	15.7	19.9	25.9	14.7	36.1	48.9	27.0
2003.....	22.7	29.6	17.1	21.4	27.8	16.1	38.2	52.7	28.0
2002.....	23.5	30.4	17.8	22.1	28.5	16.7	40.3	54.8	30.0
2001.....	24.1	30.8	18.5	22.5	28.8	17.3	41.8	56.0	31.7
2000.....	25.2	32.3	19.4	23.6	30.2	18.2	43.5	58.7	32.9
1999.....	26.4	33.5	20.6	24.7	31.2	19.3	45.8	61.5	34.9

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

2007.....	65.1	85.5	50.1	65.6	86.7	49.9	69.1	85.8	57.4
2006.....	69.4	90.6	53.9	69.9	91.7	53.8	73.8	91.7	61.4
2005.....	73.9	95.8	58.0	74.5	97.3	57.9	77.3	94.7	65.2
2004.....	76.0	97.8	60.1	76.5	99.1	59.9	80.3	98.0	68.0
2003.....	81.1	104.0	64.4	81.5	105.2	64.0	86.6	104.0	74.2
2002.....	84.1	108.0	66.7	84.7	109.5	66.4	87.2	104.1	75.2
2001.....	87.1	111.3	69.6	87.6	112.8	69.4	91.1	110.6	77.6
2000.....	90.5	116.3	72.0	91.2	118.1	71.8	91.9	109.5	79.7
1999.....	93.8	120.1	75.0	94.5	121.8	74.9	94.1	112.7	81.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Other heart diseases (I26-I51)

2007.....	53.4	60.2	48.0	52.6	59.0	47.3	69.7	81.9	60.4
2006.....	53.9	60.0	48.7	53.0	58.8	48.0	69.3	81.6	60.3
2005.....	55.2	61.5	49.9	54.1	60.2	49.0	71.9	83.0	63.4
2004.....	55.2	61.7	49.9	54.1	60.3	49.0	72.5	85.6	63.2
2003.....	57.6	64.5	52.0	56.4	63.1	50.9	75.7	87.7	66.7
2002.....	58.5	65.5	52.8	57.3	64.1	51.7	76.6	87.8	68.1
2001.....	59.1	66.1	53.4	57.7	64.4	52.3	77.8	91.8	68.2
2000.....	60.0	67.9	53.8	58.6	66.3	52.6	78.4	92.2	69.0
1999.....	60.9	69.2	54.5	59.3	67.2	53.1	81.2	97.5	70.1

Acute and subacute endocarditis (I33)

2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	1.0	0.4
2006.....	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2003.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.7
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6

Diseases of pericardium and acute myocarditis (I30-I31,I40)

2007.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.4	0.4
2006.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3
2005.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2004.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4
2003.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2000.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.5

Heart failure (I50)

2007.....	17.3	18.6	16.2	17.3	18.7	16.3	20.1	22.3	18.4
2006.....	18.9	20.0	17.9	19.0	20.2	18.1	20.9	23.0	19.3
2005.....	18.9	19.9	18.0	18.9	20.0	18.1	21.7	23.0	20.4
2004.....	18.9	19.9	18.0	19.0	20.0	18.1	20.9	23.4	19.3
2003.....	19.4	20.6	18.4	19.5	20.7	18.4	21.8	23.6	20.5
2002.....	19.5	20.4	18.6	19.5	20.5	18.6	21.8	23.3	20.5
2001.....	20.2	21.2	19.2	20.2	21.2	19.3	22.1	24.4	20.6
2000.....	20.3	21.5	19.2	20.3	21.6	19.3	22.0	24.2	20.7
1999.....	20.3	21.4	19.3	20.3	21.4	19.3	22.5	25.3	20.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

All other forms of heart disease (I26-I28, I34-I38, I42-I49, I51)

2007.....	35.5	40.7	31.2	34.6	39.5	30.5	48.5	58.1	41.3
2006.....	34.3	39.2	30.2	33.3	37.9	29.4	47.4	57.3	40.2
2005.....	35.6	40.7	31.3	34.5	39.5	30.4	49.0	58.6	41.9
2004.....	35.7	41.0	31.4	34.5	39.5	30.4	50.4	60.9	43.0
2003.....	37.5	43.1	33.0	36.2	41.6	31.9	52.6	62.8	45.1
2002.....	38.4	44.2	33.7	37.1	42.8	32.6	53.6	63.1	46.6
2001.....	38.2	44.1	33.6	36.8	42.4	32.4	54.4	65.8	46.5
2000.....	39.0	45.5	34.0	37.7	43.9	32.8	55.2	66.6	47.2
1999.....	39.9	46.9	34.6	38.4	45.0	33.3	57.4	70.8	48.2

Essential hypertension and hypertensive renal disease (I10, I12, I15)

2007.....	7.4	7.3	7.2	6.5	6.3	6.4	16.3	17.4	15.3
2006.....	7.5	7.5	7.4	6.5	6.4	6.4	17.7	19.1	16.4
2005.....	8.0	7.8	8.0	7.0	6.6	7.1	18.4	19.9	17.1
2004.....	7.7	7.5	7.6	6.6	6.5	6.6	18.2	19.0	17.3
2003.....	7.4	7.1	7.5	6.4	6.1	6.5	17.6	18.6	16.6
2002.....	7.0	6.8	7.0	6.0	5.8	6.0	16.9	17.7	16.1
2001.....	6.8	6.7	6.8	5.8	5.6	5.9	16.6	17.9	15.4
2000.....	6.5	6.5	6.5	5.6	5.5	5.5	16.3	17.1	15.6
1999.....	6.2	6.2	6.1	5.3	5.2	5.2	16.0	17.3	15.0

Cerebrovascular diseases (I60-I69)

2007.....	42.2	42.5	41.3	40.5	40.2	39.9	60.3	67.1	55.0
2006.....	43.6	43.9	42.6	41.7	41.7	41.1	61.6	67.1	57.0
2005.....	46.6	46.9	45.6	44.7	44.7	44.0	65.2	70.5	60.7
2004.....	50.0	50.4	48.9	48.0	48.1	47.2	69.9	74.9	65.5
2003.....	53.5	54.1	52.3	51.4	51.7	50.5	74.3	79.5	69.8
2002.....	56.2	56.5	55.2	54.2	54.2	53.4	76.3	81.7	71.8
2001.....	57.9	59.0	56.4	55.8	56.5	54.5	78.8	85.4	73.7
2000.....	60.9	62.4	59.1	58.8	59.8	57.3	81.9	89.6	76.2
1999.....	61.6	63.2	59.8	59.6	60.8	58.0	81.8	89.6	76.2

Atherosclerosis (I70)

2007.....	2.5	2.6	2.4	2.6	2.7	2.4	2.4	2.6	2.3
2006.....	2.7	2.8	2.6	2.8	2.8	2.6	2.7	3.3	2.4
2005.....	3.8	3.9	3.7	3.9	4.0	3.7	3.4	3.4	3.3
2004.....	3.9	4.1	3.7	4.0	4.2	3.8	3.5	3.5	3.3
2003.....	4.4	4.5	4.3	4.5	4.5	4.4	4.2	4.7	3.8
2002.....	4.7	4.9	4.6	4.8	5.0	4.7	4.4	4.9	4.0
2001.....	5.0	5.1	4.8	5.1	5.2	5.0	4.3	4.6	4.1
2000.....	5.2	5.5	5.0	5.3	5.6	5.1	4.8	5.4	4.4
1999.....	5.5	5.7	5.3	5.6	5.8	5.4	5.0	5.8	4.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Other diseases of circulatory system (I71-I78)

2007.....	6.9	8.8	5.5	6.8	8.8	5.3	8.6	10.6	7.2
2006.....	7.2	9.4	5.6	7.2	9.4	5.5	8.9	10.3	7.8
2005.....	7.8	10.0	6.1	7.6	10.0	5.9	9.6	11.7	8.0
2004.....	7.9	10.2	6.3	7.9	10.2	6.1	9.7	11.6	8.3
2003.....	8.4	11.2	6.4	8.4	11.3	6.3	9.8	11.8	8.3
2002.....	8.7	11.5	6.6	8.6	11.6	6.5	10.0	12.0	8.6
2001.....	9.0	12.0	6.9	8.9	12.0	6.7	10.8	12.8	9.4
2000.....	9.4	12.9	7.1	9.3	13.0	6.8	10.9	13.0	9.5
1999.....	9.5	12.8	7.1	9.4	13.0	7.0	10.5	12.4	9.2

Aortic aneurysm and dissection (I71)

2007.....	4.1	5.7	2.9	4.1	5.8	2.9	4.2	5.5	3.2
2006.....	4.3	6.0	3.0	4.3	6.1	3.0	4.4	5.5	3.5
2005.....	4.6	6.5	3.1	4.6	6.6	3.1	4.6	6.0	3.6
2004.....	4.6	6.7	3.1	4.7	6.8	3.2	4.5	6.1	3.3
2003.....	5.1	7.5	3.3	5.1	7.7	3.3	4.6	6.1	3.5
2002.....	5.2	7.7	3.3	5.2	7.9	3.4	4.6	6.1	3.5
2001.....	5.4	8.0	3.6	5.5	8.2	3.6	5.0	6.4	3.9
2000.....	5.7	8.7	3.7	5.8	9.0	3.6	5.2	6.8	4.1
1999.....	5.8	8.8	3.7	5.9	9.1	3.7	4.9	6.3	3.9

Other diseases of arteries, arterioles and capillaries (I72-I78)

2007.....	2.8	3.1	2.5	2.7	3.0	2.5	4.4	5.1	4.0
2006.....	2.9	3.3	2.7	2.9	3.2	2.6	4.6	4.8	4.3
2005.....	3.2	3.5	2.9	3.1	3.4	2.8	5.0	5.6	4.5
2004.....	3.3	3.5	3.1	3.2	3.4	3.0	5.2	5.6	5.0
2003.....	3.3	3.7	3.1	3.2	3.6	3.0	5.2	5.7	4.9
2002.....	3.5	3.8	3.3	3.4	3.7	3.1	5.4	6.0	5.0
2001.....	3.6	3.9	3.3	3.4	3.8	3.2	5.9	6.4	5.5
2000.....	3.7	4.2	3.4	3.6	4.0	3.2	5.7	6.2	5.3
1999.....	3.7	4.0	3.4	3.5	3.9	3.3	5.6	6.0	5.3

Other disorders of circulatory system (I80-I99)

2007.....	1.3	1.4	1.2	1.2	1.3	1.1	2.2	2.4	1.9
2006.....	1.3	1.3	1.2	1.2	1.2	1.1	2.3	2.4	2.2
2005.....	1.6	1.6	1.5	1.5	1.5	1.4	2.7	3.1	2.3
2004.....	1.5	1.6	1.5	1.5	1.5	1.4	2.7	2.9	2.6
2003.....	1.6	1.6	1.6	1.5	1.4	1.5	2.9	3.2	2.8
2002.....	1.6	1.6	1.6	1.5	1.5	1.6	2.9	3.2	2.6
2001.....	1.6	1.7	1.6	1.5	1.6	1.5	2.7	3.1	2.5
2000.....	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.0	2.8
1999.....	1.5	1.5	1.4	1.4	1.4	1.3	2.4	2.6	2.2

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Influenza and pneumonia (J09-J18)

2007.....	16.2	19.3	14.2	16.0	18.9	14.1	18.4	23.6	15.
2006.....	17.8	21.2	15.5	17.7	20.9	15.5	19.6	24.4	16.
2005.....	20.3	23.9	17.9	20.2	23.6	18.0	21.7	26.9	18.
2004.....	19.8	23.7	17.3	19.6	23.3	17.2	22.3	29.1	18.
2003.....	22.0	26.1	19.4	21.9	25.7	19.5	23.3	30.9	18.
2002.....	22.6	27.0	19.9	22.6	26.7	19.9	24.0	30.8	20.
2001.....	22.0	26.6	19.2	21.7	26.0	19.1	24.1	32.3	19.
2000.....	23.7	28.9	20.7	23.5	28.4	20.7	25.6	33.9	20.
1999.....	23.5	28.5	20.6	23.3	28.1	20.6	25.4	33.4	20.

Influenza (J09-J11)

2007	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*		
2006	0.3	0.3	0.2	0.3	0.3	0.3	0.1	0.2	0.	0.
2005	0.6	0.6	0.6	0.6	0.6	0.6	0.2	0.3	0.	0.
2004	0.4	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.	0.
2003	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.	0.
2002	0.2	0.3	0.2	0.2	0.3	0.2	0.1	*		
2001	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*		
2000	0.6	0.7	0.6	0.6	0.7	0.6	0.3	0.4	0.	0.
1999	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.	0.

Pneumonia (J12-J18)

2007.....	16.1	19.2	14.1	15.9	18.8	14.0	18.4	23.6	15.
2006.....	17.5	20.9	15.3	17.4	20.6	15.2	19.5	24.2	16.
2005.....	19.7	23.3	17.3	19.6	23.0	17.3	21.5	26.6	18.
2004.....	19.4	23.3	16.9	19.2	22.9	16.8	22.1	28.9	18.
2003.....	21.4	25.5	18.8	21.3	25.0	18.9	23.0	30.6	18.
2002.....	22.4	26.7	19.6	22.3	26.4	19.7	23.9	30.7	19.
2001.....	21.9	26.5	19.1	21.6	25.9	19.0	24.1	32.2	19.
2000.....	23.1	28.2	20.1	22.9	27.7	20.0	25.3	33.5	20.
1999.....	22.9	27.9	20.0	22.7	27.5	19.9	25.1	33.2	20.

Other acute lower respiratory infections (J20-J22,U04)

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Acute bronchitis and bronchiolitis (J20-J21)

2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2

Other and unspecified acute lower respiratory infections (J22-U04)

2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*

Chronic lower respiratory diseases (J40-J47)

2007.....	40.8	48.0	36.0	43.0	49.8	38.5	28.1	39.6	21.4
2006.....	40.5	47.6	35.9	42.6	49.2	38.4	28.1	39.5	21.3
2005.....	43.2	51.2	38.1	45.4	52.8	40.7	30.6	44.1	22.8
2004.....	41.1	49.5	36.0	43.2	51.1	38.4	28.2	40.9	20.9
2003.....	43.3	52.3	37.8	45.4	53.8	40.3	30.1	44.4	22.0
2002.....	43.5	53.5	37.4	45.4	54.9	39.7	31.2	46.3	22.6
2001.....	43.7	54.0	37.6	45.6	55.5	39.8	30.9	46.3	22.4
2000.....	44.2	55.8	37.4	46.0	57.2	39.5	31.6	47.5	22.7
1999.....	45.4	58.7	37.7	47.1	60.0	39.7	33.5	51.5	23.4

Bronchitis, chronic and unspecified (J40-J42)

2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.1
2004.....	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.2	0.3	0.2
2001.....	0.3	0.4	0.3	0.3	0.4	0.3	0.2	0.3	0.2
2000.....	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.3	0.3
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Emphysema (J43)

2007.....	4.1	5.1	3.4	4.3	5.3	3.7	2.6	4.2	1.6
2006.....	4.1	5.0	3.5	4.4	5.2	3.8	2.6	4.0	1.7
2005.....	4.6	5.8	3.8	4.9	6.0	4.1	3.1	4.9	1.9
2004.....	4.6	5.8	3.8	4.9	6.1	4.1	2.7	4.6	1.6
2003.....	5.1	6.3	4.3	5.4	6.6	4.6	3.2	5.3	2.0
2002.....	5.4	7.0	4.4	5.7	7.2	4.8	3.4	5.8	2.0
2001.....	5.8	7.4	4.7	6.1	7.8	5.1	3.3	5.3	2.2
2000.....	6.0	7.8	4.9	6.4	8.1	5.3	3.8	6.2	2.4
1999.....	6.5	8.6	5.2	6.9	8.9	5.6	4.0	6.7	2.3

Asthma (J45-J46)

2007.....	1.1	0.9	1.2	0.9	0.7	1.0	2.5	2.5	2.6
2006.....	1.2	0.9	1.3	0.9	0.7	1.1	2.7	2.5	2.8
2005.....	1.3	1.0	1.5	1.0	0.7	1.3	3.0	2.7	3.3
2004.....	1.3	1.0	1.4	1.0	0.8	1.2	3.0	3.0	3.1
2003.....	1.4	1.1	1.6	1.1	0.9	1.3	3.2	2.9	3.3
2002.....	1.5	1.2	1.7	1.2	0.9	1.4	3.4	3.3	3.4
2001.....	1.5	1.2	1.7	1.2	0.9	1.5	3.6	3.2	3.8
2000.....	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.1
1999.....	1.7	1.3	2.0	1.4	1.0	1.7	3.9	3.5	4.1

Other chronic lower respiratory diseases (J44,J47)

2007.....	35.3	41.9	31.2	37.5	43.7	33.6	22.8	32.7	17.1
2006.....	34.9	41.4	30.8	37.0	43.1	33.2	22.5	32.7	16.6
2005.....	37.0	44.1	32.5	39.2	45.8	35.0	24.3	36.3	17.5
2004.....	35.0	42.3	30.5	37.0	43.9	32.8	22.2	33.0	16.0
2003.....	36.5	44.5	31.7	38.6	46.0	34.0	23.4	36.0	16.3
2002.....	36.2	44.9	31.1	38.2	46.4	33.2	24.1	36.9	16.9
2001.....	36.1	45.0	30.8	37.9	46.4	32.9	23.7	37.5	16.2
2000.....	36.1	46.1	30.3	37.8	47.5	32.3	23.7	37.5	15.9
1999.....	36.8	48.3	30.2	38.5	49.6	32.0	25.3	40.7	16.7

Pneumoconioses and chemical effects (J60-J66,J68)

2007.....	0.3	0.7	0.0	0.3	0.8	0.0	0.2	0.4	*
2006.....	0.3	0.7	0.0	0.3	0.8	0.0	0.2	0.5	*
2005.....	0.3	0.8	0.0	0.4	0.9	0.0	0.2	0.5	*
2004.....	0.3	0.9	0.0	0.4	0.9	0.0	0.2	0.6	*
2003.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*
2002.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001.....	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000.....	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.7	*
1999.....	0.4	1.0	0.0	0.4	1.1	0.0	0.3	0.8	*

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Pneumonitis due to solids and liquids (J69)

2007.....	5.2	7.3	4.0	5.3	7.4	4.0	5.3	7.5	4.0
2006.....	5.3	7.5	4.0	5.4	7.6	4.1	5.5	8.0	4.1
2005.....	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9
2004.....	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6
2003.....	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

2007.....	9.0	11.1	7.6	9.1	11.2	7.7	8.6	10.3	7.5
2006.....	8.9	10.9	7.6	9.1	11.0	7.7	8.3	9.9	7.3
2005.....	8.9	10.6	7.7	9.0	10.7	7.8	8.7	10.7	7.5
2004.....	9.2	11.1	7.8	9.3	11.3	7.9	8.6	10.4	7.4
2003.....	8.7	10.5	7.5	8.8	10.6	7.6	8.1	10.0	6.9
2002.....	8.7	10.4	7.6	8.8	10.5	7.6	8.4	9.9	7.4
2001.....	9.3	11.0	8.1	9.3	11.1	8.1	9.4	10.7	8.6
2000.....	9.2	11.3	7.9	9.3	11.3	8.0	9.2	11.2	8.0
1999.....	9.2	11.2	7.8	9.2	11.2	7.8	9.4	11.2	8.4

Peptic ulcer (K25-K28)

2007.....	1.0	1.1	0.8	1.0	1.1	0.8	1.0	1.4	0.7
2006.....	1.1	1.2	0.9	1.0	1.2	0.9	1.2	1.7	0.9
2005.....	1.1	1.4	0.9	1.1	1.3	0.9	1.3	1.8	0.9
2004.....	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1
2003.....	1.3	1.6	1.1	1.3	1.5	1.2	1.4	2.1	1.0
2002.....	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001.....	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000.....	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.3
1999.....	1.7	2.0	1.4	1.7	2.0	1.4	1.9	2.4	1.4

Diseases of appendix (K35-K38)

2007.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2004.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2002.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Hernia (K40-K46)

2007.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.7	0.5
2006.....	0.5	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6
2005.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2004.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2003.....	0.6	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6
2002.....	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5
2000.....	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.6
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6

Chronic liver disease and cirrhosis (K70,K73-K74)

2007.....	9.1	12.7	5.9	9.4	13.1	6.1	7.4	11.0	4.5
2006.....	8.8	12.1	5.8	9.1	12.5	6.0	7.0	10.2	4.4
2005.....	9.0	12.4	5.8	9.2	12.7	6.0	7.7	11.5	4.8
2004.....	9.0	12.5	5.8	9.2	12.8	6.0	7.9	11.7	4.9
2003.....	9.3	13.0	6.0	9.5	13.3	6.1	8.4	12.4	5.2
2002.....	9.4	12.9	6.3	9.6	13.2	6.3	8.5	12.0	5.7
2001.....	9.5	13.2	6.2	9.6	13.3	6.3	9.3	13.8	5.6
2000.....	9.5	13.4	6.2	9.6	13.5	6.1	9.4	13.5	6.3
1999.....	9.6	13.5	6.1	9.6	13.6	6.0	10.1	14.9	6.3

Alcoholic liver disease (K70)

2007.....	4.5	6.8	2.3	4.7	7.1	2.4	3.4	5.5	1.8
2006.....	4.1	6.3	2.2	4.3	6.5	2.3	3.0	4.7	1.7
2005.....	4.2	6.4	2.2	4.3	6.6	2.3	3.5	5.4	1.9
2004.....	4.2	6.3	2.2	4.3	6.5	2.2	3.5	5.6	1.9
2003.....	4.2	6.4	2.1	4.2	6.6	2.1	3.8	6.0	2.0
2002.....	4.2	6.3	2.2	4.2	6.5	2.2	3.8	5.8	2.3
2001.....	4.3	6.6	2.2	4.3	6.6	2.2	4.4	6.9	2.4
2000.....	4.3	6.8	2.2	4.3	6.9	2.0	4.6	7.0	2.8
1999.....	4.3	6.9	2.1	4.3	6.8	2.0	5.1	8.1	2.8

Other chronic liver disease and cirrhosis (K73-K74)

2007.....	4.6	5.9	3.5	4.8	6.0	3.7	4.0	5.5	2.8
2006.....	4.6	5.9	3.6	4.8	6.0	3.7	3.9	5.6	2.7
2005.....	4.7	6.0	3.6	4.9	6.2	3.7	4.2	6.1	2.8
2004.....	4.8	6.1	3.7	4.9	6.2	3.8	4.3	6.2	2.9
2003.....	5.2	6.5	3.9	5.3	6.7	4.0	4.6	6.4	3.2
2002.....	5.2	6.5	4.1	5.4	6.7	4.2	4.6	6.2	3.4
2001.....	5.2	6.6	4.0	5.3	6.7	4.1	4.9	6.9	3.2
2000.....	5.2	6.6	4.0	5.3	6.7	4.0	4.8	6.5	3.5
1999.....	5.2	6.7	4.0	5.3	6.7	4.0	5.0	6.8	3.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Cholelithiasis and other disorders of gallbladder (K80-K82)

2007.....	1.0	1.1	0.9	1.0	1.1	0.9	1.1	1.1	1.0
2006.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.1	1.2
2005.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.2
2004.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.2	1.2
2003.....	1.0	1.2	0.9	1.0	1.1	0.9	1.1	1.3	1.0
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.1	1.2
1999.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.0

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

2007.....	14.5	17.8	12.5	13.2	16.5	11.0	29.4	33.7	26.7
2006.....	14.5	17.8	12.4	13.0	16.3	10.8	30.2	35.0	27.2
2005.....	14.3	17.5	12.3	12.9	16.2	10.8	29.7	34.1	26.9
2004.....	14.2	17.3	12.3	12.8	15.9	10.8	29.6	34.2	26.7
2003.....	14.4	17.8	12.4	13.0	16.3	10.9	30.4	36.1	26.8
2002.....	14.2	17.6	12.1	12.7	16.1	10.7	29.7	35.3	26.3
2001.....	14.0	17.4	12.0	12.5	16.0	10.5	29.5	33.8	27.1
2000.....	13.5	16.9	11.5	12.1	15.6	10.0	28.7	33.1	26.0
1999.....	13.0	16.5	11.1	11.5	15.0	9.6	28.4	33.7	25.4

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2005.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.1	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2000.....	0.1	0.1	0.0	0.0	0.1	0.0	0.1	*	*
1999.....	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	*

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

2007.....	0.9	1.2	0.8	0.8	1.1	0.7	1.9	2.2	1.7
2006.....	0.6	0.7	0.5	0.5	0.6	0.4	1.3	1.6	1.1
2005.....	0.3	0.4	0.2	0.3	0.3	0.2	0.6	0.7	0.5
2004.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
2003.....	0.2	0.2	0.1	0.2	0.2	0.1	0.3	0.4	0.2
2002.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.2
2000.....	0.2	0.3	0.2	0.2	0.2	0.1	0.4	0.4	0.3
1999.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Renal failure (N17-N19)

2007.....	13.6	16.5	11.6	12.3	15.3	10.3	27.4	31.4	24.9
2006.....	13.9	17.0	11.8	12.4	15.6	10.4	28.9	33.3	26.0
2005.....	14.0	17.1	12.0	12.6	15.8	10.5	29.0	33.3	26.3
2004.....	13.9	17.0	12.0	12.5	15.6	10.6	29.1	33.8	26.3
2003.....	14.2	17.5	12.1	12.7	16.1	10.7	30.0	35.6	26.5
2002.....	13.9	17.3	11.9	12.5	15.8	10.5	29.3	34.8	25.9
2001.....	13.7	17.1	11.8	12.3	15.8	10.2	29.2	33.4	26.7
2000.....	13.2	16.6	11.3	11.8	15.2	9.8	28.2	32.6	25.5
1999.....	12.8	16.1	10.8	11.3	14.6	9.4	27.9	33.0	24.9

Other disorders of kidney (N25,N27)

2007.....	0.0	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	0.0	*	*	*	*	*
2005.....	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	*	0.0	0.0	*	*	*	*	*
2003.....	0.0	*	*	0.0	*	*	*	*	*
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	*	*	*	*
2000.....	0.0	0.0	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	0.0	0.0	*	*	*	*	*

Infections of kidney (N10-N12,N13.6,N15.1)

2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2005.....	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.2	0.4
2004.....	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.2	0.3
2003.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.4
2002.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.3
2001.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.3
2000.....	0.3	0.2	0.3	0.3	0.2	0.3	0.4	0.3	0.5
1999.....	0.3	0.2	0.3	0.3	0.3	0.4	0.3	0.3	0.4

Hyperplasia of prostate (N40)

2007.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2006.....	0.2	0.5	...	0.2	0.5	...	0.2	0.4	...
2005.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2004.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
	Male	Female			Male	Female	Male	Female	Male	Female

Inflammatory diseases of female pelvic organs (N70-N76)

2007.....	0.0	...	0.1	0.0	...	0.1	0.0	...	0.1
2006.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2005.....	0.0	...	0.1	0.0	...	0.1	*	...	*
2004.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2003.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

Pregnancy, childbirth and the puerperium (O00-O99)

2007.....	0.3	...	0.5	0.2	...	0.4	0.6	...	1.2
2006.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2005.....	0.3	...	0.5	0.2	...	0.4	0.7	...	1.3
2004.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2003.....	0.2	...	0.4	0.1	...	0.3	0.5	...	0.9
2002.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999.....	0.2	...	0.3	0.1	...	0.2	0.4	...	0.8

Pregnancy with abortive outcome (O00-O07)

2007.....	0.0	...	0.0	*	...	*	*	...	*
2006.....	0.0	...	0.0	*	...	*	*	...	*
2005.....	0.0	...	0.0	*	...	*	*	...	*
2004.....	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	*	...	*

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

2007.....	0.3	...	0.5	0.2	...	0.4	0.5	...	1.1
2006.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2005.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2
2004.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.1
2003.....	0.2	...	0.3	0.1	...	0.3	0.4	...	0.8
2002.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
2000.....	0.1	...	0.3	0.1	...	0.2	0.3	...	0.6
1999.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Certain conditions originating in the perinatal period (P00-P96)

2007.....	4.7	5.2	4.2	3.7	4.1	3.3	9.8	10.8	8.7
2006.....	4.8	5.3	4.3	3.8	4.2	3.3	10.4	11.2	9.6
2005.....	4.9	5.4	4.4	3.8	4.2	3.4	10.5	11.5	9.5
2004.....	4.8	5.4	4.2	3.8	4.3	3.3	10.3	11.3	9.2
2003.....	4.9	5.5	4.3	4.0	4.4	3.5	10.2	11.4	9.0
2002.....	4.9	5.4	4.3	3.9	4.3	3.4	10.1	10.9	9.2
2001.....	4.7	5.3	4.2	3.7	4.1	3.3	10.3	11.3	9.1
2000.....	5.1	5.6	4.5	4.0	4.4	3.5	11.3	12.6	9.9
1999.....	5.2	5.7	4.6	4.0	4.4	3.6	11.8	13.1	10.4

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

2007.....	3.4	3.6	3.2	3.3	3.5	3.1	3.9	4.0	3.8
2006.....	3.5	3.7	3.2	3.4	3.7	3.2	4.0	4.4	3.8
2005.....	3.5	3.7	3.3	3.5	3.6	3.3	3.9	4.1	3.7
2004.....	3.6	3.7	3.4	3.5	3.7	3.4	4.0	4.4	3.7
2003.....	3.6	3.8	3.4	3.6	3.7	3.4	4.0	4.4	3.6
2002.....	3.7	3.8	3.5	3.7	3.8	3.5	4.1	4.4	3.9
2001.....	3.6	3.8	3.4	3.6	3.8	3.4	4.1	4.3	3.9
2000.....	3.8	4.1	3.6	3.7	4.0	3.5	4.4	4.8	4.1
1999.....	3.8	3.9	3.6	3.7	3.9	3.6	4.3	4.7	4.1

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

2007.....	10.4	10.5	9.8	10.0	10.2	9.5	13.9	15.2	12.4
2006.....	10.1	10.5	9.3	9.8	10.1	9.1	13.3	15.4	11.6
2005.....	10.4	11.1	9.4	10.0	10.6	9.1	14.6	17.1	12.5
2004.....	10.0	10.7	9.0	9.6	10.2	8.7	14.0	16.0	12.1
2003.....	10.6	11.4	9.5	10.2	11.0	9.2	14.8	17.4	12.5
2002.....	10.3	11.4	9.0	9.9	10.8	8.7	14.5	18.0	11.8
2001.....	11.3	12.7	9.8	11.0	12.2	9.5	15.1	17.7	12.8
2000.....	11.4	13.1	9.7	11.0	12.6	9.3	16.5	19.7	13.6
1999.....	9.7	11.0	8.4	9.1	10.1	7.9	15.4	19.1	12.4

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

2007.....	73.8	73.6	72.2	73.3	72.9	71.9	89.7	94.3	84.9
2006.....	75.2	75.0	73.4	74.7	74.3	73.0	91.0	96.1	86.2
2005.....	70.5	70.5	68.6	69.7	69.5	67.9	87.7	91.5	83.5
2004.....	67.1	67.3	65.2	66.2	66.1	64.5	84.9	91.8	79.3
2003.....	68.3	68.2	66.6	67.4	66.9	65.8	85.6	91.6	80.3
2002.....	67.1	67.3	65.2	66.1	65.9	64.3	84.7	91.5	79.1
2001.....	65.9	66.1	64.0	64.7	64.5	63.0	84.6	92.0	78.5
2000.....	64.6	65.4	62.4	63.2	63.6	61.2	83.8	92.3	77.2
1999.....	63.0	64.0	60.6	61.6	62.2	59.5	81.0	90.0	74.5

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

2007.....	40.0	55.2	25.8	41.5	56.8	26.9	36.6	54.8	21.6
2006.....	39.8	55.2	25.5	41.0	56.4	26.4	38.3	57.5	22.2
2005.....	39.1	54.2	25.0	40.1	55.2	25.8	38.7	58.1	22.8
2004.....	37.7	52.1	24.5	38.8	53.2	25.3	36.3	54.4	21.7
2003.....	37.3	51.8	24.1	38.2	52.8	24.7	36.1	53.9	21.6
2002.....	36.9	51.5	23.5	37.5	52.0	24.0	36.9	56.2	21.3
2001.....	35.7	50.2	22.5	36.0	50.3	22.8	37.6	56.9	22.0
2000.....	34.9	49.3	22.0	35.1	49.2	22.2	37.7	58.1	21.3
1999.....	35.3	49.8	22.3	35.2	49.3	22.4	40.1	61.3	23.1

Transport accidents (V01-V99,Y85)

2007.....	15.3	22.4	8.6	15.8	22.9	8.9	15.0	24.1	7.3
2006.....	16.0	23.0	9.2	16.4	23.5	9.5	15.5	24.2	8.1
2005.....	16.2	23.4	9.3	16.7	23.9	9.7	15.4	24.1	8.0
2004.....	16.2	23.1	9.7	16.7	23.6	10.0	15.7	24.4	8.4
2003.....	16.4	23.4	9.8	16.8	23.9	10.0	15.8	24.5	8.6
2002.....	16.7	23.9	9.9	17.1	24.2	10.2	16.0	25.0	8.5
2001.....	16.6	23.7	9.8	16.8	24.0	10.0	16.5	25.5	8.9
2000.....	16.6	23.7	10.0	16.8	23.8	10.2	16.7	26.1	8.7
1999.....	16.6	23.6	10.2	16.8	23.6	10.4	17.1	26.5	9.5

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)

2007.....	14.4	20.9	8.2	14.8	21.3	8.5	14.1	22.5	7.0
2006.....	15.0	21.4	8.8	15.4	21.8	9.1	14.6	22.6	7.8
2005.....	15.2	21.7	8.9	15.6	22.2	9.2	14.5	22.5	7.6
2004.....	15.2	21.4	9.3	15.6	21.8	9.5	14.8	22.7	8.1
2003.....	15.3	21.6	9.3	15.7	21.9	9.5	14.9	22.7	8.3
2002.....	15.7	22.1	9.6	16.0	22.4	9.8	15.0	23.2	8.2
2001.....	15.3	21.8	9.3	15.6	22.0	9.5	15.4	23.5	8.5
2000.....	15.4	21.7	9.5	15.6	21.8	9.8	15.7	24.4	8.4
1999.....	15.2	21.3	9.6	15.3	21.3	9.7	15.9	24.3	9.0

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)

2007.....	0.3	0.6	0.1	0.3	0.5	0.1	0.4	0.7	0.2
2006.....	0.4	0.6	0.2	0.4	0.6	0.2	0.4	0.7	0.2
2005.....	0.4	0.7	0.2	0.4	0.6	0.2	0.5	1.0	0.2
2004.....	0.4	0.6	0.2	0.4	0.6	0.2	0.5	0.9	0.2
2003.....	0.4	0.7	0.2	0.5	0.7	0.2	0.5	1.0	0.2
2002.....	0.4	0.6	0.1	0.4	0.6	0.1	0.5	0.9	0.1
2001.....	0.5	0.8	0.2	0.5	0.7	0.2	0.6	1.0	0.3
2000.....	0.5	0.8	0.3	0.5	0.8	0.2	0.6	1.0	0.3
1999.....	0.8	1.1	0.4	0.8	1.1	0.4	0.7	1.2	0.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

2007.....	0.6	1.0	0.2	0.6	1.0	0.2	0.5	0.9	0.1
2006.....	0.6	1.0	0.3	0.7	1.1	0.3	0.5	0.8	0.2
2005.....	0.6	1.0	0.2	0.7	1.1	0.2	0.4	0.7	0.1
2004.....	0.6	1.1	0.3	0.7	1.1	0.3	0.4	0.8	0.1
2003.....	0.7	1.1	0.3	0.7	1.2	0.3	0.4	0.8	0.1
2002.....	0.6	1.1	0.2	0.7	1.2	0.2	0.5	0.9	0.1
2001.....	0.7	1.2	0.3	0.8	1.3	0.3	0.5	1.0	0.1
2000.....	0.7	1.2	0.2	0.7	1.3	0.2	0.4	0.8	*
1999.....	0.7	1.1	0.3	0.7	1.2	0.3	0.5	1.0	0.2

Nontransport accidents (W00-X59,Y86)

2007.....	24.6	32.7	17.2	25.7	34.0	18.0	21.6	30.8	14.2
2006.....	23.8	32.1	16.2	24.6	32.9	16.9	22.8	33.4	14.1
2005.....	22.9	30.9	15.7	23.4	31.4	16.1	23.3	34.0	14.8
2004.....	21.5	29.0	14.9	22.1	29.6	15.3	20.6	30.0	13.3
2003.....	20.9	28.4	14.3	21.4	28.9	14.6	20.2	29.5	13.0
2002.....	20.2	27.6	13.5	20.5	27.8	13.8	20.9	31.2	12.9
2001.....	19.1	26.4	12.7	19.2	26.4	12.8	21.1	31.4	13.1
2000.....	18.4	25.6	12.0	18.3	25.3	12.0	21.0	32.0	12.5
1999.....	18.7	26.3	12.1	18.4	25.7	12.0	22.9	34.8	13.6

Falls (W00-W19)

2007.....	7.0	9.0	5.5	7.5	9.5	5.9	3.4	4.9	2.4
2006.....	6.6	8.6	5.1	7.0	8.9	5.5	3.7	5.4	2.5
2005.....	6.4	8.3	4.9	6.7	8.6	5.2	3.7	5.2	2.6
2004.....	6.3	8.2	4.8	6.6	8.6	5.1	3.6	5.5	2.3
2003.....	5.9	7.6	4.5	6.1	7.9	4.8	3.4	4.7	2.4
2002.....	5.6	7.4	4.3	5.9	7.6	4.6	3.1	4.9	2.0
2001.....	5.3	7.2	3.9	5.6	7.5	4.1	3.3	5.1	2.0
2000.....	4.8	6.6	3.6	5.0	6.8	3.7	2.9	4.6	1.8
1999.....	4.8	6.6	3.5	5.0	6.8	3.7	3.4	5.3	2.1

Accidental discharge of firearms (W32-W34)

2007.....	0.2	0.4	0.0	0.2	0.3	0.1	0.3	0.5	*
2006.....	0.2	0.4	0.0	0.2	0.3	0.0	0.3	0.6	*
2005.....	0.3	0.5	0.1	0.3	0.4	0.1	0.4	0.7	*
2004.....	0.2	0.4	0.0	0.2	0.4	0.1	0.3	0.6	*
2003.....	0.2	0.5	0.1	0.2	0.5	0.1	0.3	0.6	*
2002.....	0.3	0.4	0.1	0.2	0.4	0.1	0.4	0.7	*
2001.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.5	*
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1
1999.....	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Accidental drowning and submersion (W65-W74)

2007.....	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.1	0.4
2006.....	1.2	1.9	0.5	1.2	1.8	0.6	1.3	2.3	0.4
2005.....	1.2	1.9	0.5	1.2	1.8	0.5	1.4	2.5	0.4
2004.....	1.1	1.8	0.5	1.1	1.7	0.5	1.3	2.2	0.5
2003.....	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002.....	1.2	2.0	0.5	1.1	1.8	0.4	1.6	2.7	0.5
2001.....	1.1	1.8	0.5	1.1	1.7	0.5	1.5	2.6	0.5
2000.....	1.2	2.0	0.5	1.2	1.8	0.5	1.5	2.6	0.6
1999.....	1.3	2.0	0.5	1.2	1.9	0.5	1.7	3.0	0.5

Accidental exposure to smoke, fire and flames (X00-X09)

2007.....	1.1	1.3	0.8	0.9	1.2	0.7	2.2	3.0	1.6
2006.....	1.0	1.3	0.8	0.9	1.2	0.7	2.1	2.7	1.6
2005.....	1.1	1.4	0.8	0.9	1.2	0.7	2.2	3.1	1.6
2004.....	1.1	1.4	0.8	0.9	1.2	0.7	2.4	3.3	1.8
2003.....	1.2	1.4	0.9	1.1	1.3	0.8	2.4	3.2	1.9
2002.....	1.1	1.4	0.8	0.9	1.3	0.7	2.4	3.3	1.7
2001.....	1.2	1.5	0.9	1.0	1.3	0.7	2.7	3.8	1.9
2000.....	1.2	1.6	0.9	1.0	1.3	0.7	3.0	4.2	2.2
1999.....	1.2	1.6	0.9	1.0	1.4	0.7	3.0	4.0	2.3

Accidental poisoning and exposure to noxious substances (X40-X49)

2007.....	9.8	13.0	6.6	10.6	14.0	7.2	8.6	12.2	5.5
2006.....	9.1	12.4	5.9	9.7	13.0	6.3	9.4	14.1	5.5
2005.....	7.9	10.7	5.1	8.4	11.3	5.5	8.2	11.9	4.9
2004.....	7.1	9.5	4.7	7.6	10.1	5.0	6.9	9.9	4.5
2003.....	6.7	9.1	4.3	7.1	9.6	4.5	6.8	9.9	4.1
2002.....	6.1	8.4	3.7	6.3	8.7	3.9	6.8	10.1	3.9
2001.....	4.9	7.0	2.9	5.0	7.1	2.9	6.3	9.5	3.7
2000.....	4.5	6.6	2.5	4.5	6.6	2.5	6.0	9.3	3.3
1999.....	4.4	6.4	2.3	4.3	6.3	2.3	6.2	9.8	3.2

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

2007.....	5.4	7.2	3.7	5.4	7.2	3.7	5.9	8.0	4.2
2006.....	5.6	7.6	3.8	5.7	7.7	3.9	5.9	8.3	4.1
2005.....	6.1	8.2	4.2	6.0	8.1	4.1	7.5	10.6	5.2
2004.....	5.7	7.6	4.0	5.7	7.7	4.0	6.0	8.4	4.2
2003.....	5.8	7.9	4.1	5.8	7.9	4.1	6.2	8.9	4.2
2002.....	5.9	7.9	4.2	5.9	7.9	4.2	6.7	9.4	4.7
2001.....	6.3	8.4	4.5	6.2	8.3	4.5	7.0	9.8	4.9
2000.....	6.3	8.5	4.4	6.3	8.4	4.4	7.1	10.8	4.5
1999.....	6.7	9.1	4.8	6.7	8.9	4.8	8.2	12.2	5.4

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

2007.....	11.3	18.4	4.7	12.5	20.2	5.2	5.0	8.8	1.7
2006.....	10.9	18.0	4.5	12.1	19.6	5.1	5.1	9.4	1.4
2005.....	10.9	18.0	4.4	12.0	19.6	4.9	5.2	9.2	1.9
2004.....	10.9	18.0	4.5	12.0	19.6	5.0	5.3	9.6	1.8
2003.....	10.8	18.0	4.2	11.8	19.6	4.6	5.2	9.2	1.9
2002.....	10.9	18.4	4.2	12.0	20.0	4.7	5.3	9.8	1.6
2001 3/.....	10.7	18.2	4.0	11.7	19.6	4.5	5.5	9.8	1.8
2000.....	10.4	17.7	4.0	11.3	19.1	4.3	5.5	10.0	1.8
1999.....	10.5	17.8	4.0	11.3	19.0	4.4	5.6	10.3	1.6

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

2007.....	5.6	10.3	1.4	6.3	11.4	1.6	2.5	4.9	0.5
2006.....	5.5	10.2	1.4	6.2	11.2	1.6	2.6	5.3	0.5
2005.....	5.7	10.5	1.4	6.3	11.5	1.6	2.7	5.3	0.5
2004.....	5.6	10.3	1.5	6.3	11.4	1.7	2.7	5.4	0.5
2003.....	5.8	10.7	1.4	6.4	11.8	1.6	2.7	5.3	0.6
2002.....	5.9	11.1	1.4	6.5	12.1	1.6	3.0	5.9	0.5
2001.....	5.9	11.0	1.4	6.5	12.0	1.6	3.1	6.0	0.6
2000.....	5.9	11.0	1.5	6.5	11.9	1.6	3.1	6.0	0.7
1999.....	6.0	11.1	1.5	6.5	12.0	1.7	3.3	6.5	0.6

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

2007.....	5.6	8.1	3.3	6.2	8.7	3.6	2.5	3.9	1.2
2006.....	5.4	7.8	3.1	5.9	8.4	3.5	2.4	4.1	0.9
2005.....	5.2	7.5	3.0	5.7	8.1	3.3	2.5	3.9	1.4
2004.....	5.3	7.6	3.1	5.7	8.2	3.3	2.6	4.2	1.3
2003.....	5.0	7.3	2.8	5.4	7.8	3.1	2.5	3.9	1.3
2002.....	5.0	7.4	2.8	5.5	7.9	3.1	2.4	3.9	1.1
2001 3/.....	4.8	7.1	2.6	5.2	7.6	2.9	2.4	3.8	1.1
2000.....	4.5	6.7	2.5	4.9	7.1	2.7	2.4	4.0	1.1
1999.....	4.5	6.6	2.5	4.8	7.1	2.7	2.3	3.8	1.0

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

2007.....	6.1	9.6	2.5	3.7	5.4	2.0	21.1	37.1	6.1
2006.....	6.2	9.7	2.5	3.7	5.4	1.9	21.6	37.8	6.4
2005.....	6.1	9.6	2.5	3.7	5.3	1.9	21.1	37.3	6.1
2004.....	5.9	9.2	2.5	3.6	5.3	1.9	20.1	35.1	6.3
2003.....	6.0	9.4	2.6	3.7	5.3	2.0	21.0	36.7	6.4
2002.....	6.1	9.4	2.8	3.7	5.3	2.0	21.0	36.4	6.9
2001 3/.....	7.1	10.8	3.3	4.9	7.1	2.6	21.2	36.2	7.4
2000.....	5.9	9.0	2.8	3.6	5.2	2.1	20.5	35.4	7.1
1999.....	6.0	9.1	2.9	3.8	5.4	2.1	20.1	34.2	7.3

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

2007.....	4.2	7.1	1.2	2.2	3.5	1.0	16.3	30.1	3.2
2006.....	4.3	7.2	1.3	2.2	3.4	0.9	16.7	30.8	3.5
2005.....	4.2	7.0	1.2	2.2	3.5	0.9	16.0	29.8	3.0
2004.....	4.0	6.6	1.1	2.2	3.4	0.8	15.0	28.0	2.9
2003.....	4.1	6.8	1.2	2.2	3.4	0.9	15.7	29.0	3.2
2002.....	4.1	6.8	1.3	2.2	3.4	1.0	15.6	28.7	3.5
2001.....	3.9	6.6	1.3	2.2	3.5	1.0	14.8	27.4	3.0
2000.....	3.8	6.3	1.2	2.1	3.2	0.9	14.6	26.9	3.1
1999.....	3.8	6.3	1.3	2.1	3.3	1.0	14.3	26.1	3.5

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

2007.....	1.9	2.5	1.3	1.4	1.9	1.0	4.8	7.0	2.9
2006.....	1.9	2.6	1.3	1.5	1.9	1.0	4.9	7.0	3.0
2005.....	1.9	2.6	1.3	1.5	1.9	1.0	5.1	7.5	3.1
2004.....	2.0	2.5	1.4	1.5	1.9	1.1	5.1	7.1	3.4
2003.....	2.0	2.6	1.4	1.5	1.9	1.1	5.3	7.7	3.2
2002.....	2.0	2.6	1.4	1.5	1.9	1.1	5.4	7.7	3.4
2001 3/.....	3.1	4.3	2.0	2.6	3.6	1.6	6.4	8.8	4.3
2000.....	2.1	2.7	1.5	1.5	2.0	1.1	6.0	8.5	3.9
1999.....	2.2	2.8	1.6	1.6	2.1	1.2	5.8	8.1	3.8

Legal intervention (Y35,Y89.0)

2007.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2006.....	0.2	0.3	*	0.1	0.2	*	0.3	0.6	*
2005.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2004.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2003.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2002.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2001.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2000.....	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
1999.....	0.1	0.3	*	0.1	0.2	*	0.3	0.7	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

2007.....	1.8	2.2	1.3	1.8	2.2	1.4	2.0	3.0	1.1
2006.....	1.7	2.2	1.2	1.7	2.1	1.3	1.9	2.9	1.0
2005.....	1.6	1.9	1.2	1.6	1.9	1.3	1.8	2.5	1.2
2004.....	1.7	2.2	1.2	1.7	2.2	1.3	1.9	2.7	1.1
2003.....	1.7	2.3	1.2	1.8	2.3	1.3	1.9	2.9	1.0
2002.....	1.7	2.2	1.2	1.7	2.2	1.2	2.1	3.1	1.2
2001.....	1.5	2.0	0.9	1.5	2.0	1.0	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	1.3	1.8	0.8	1.9	3.0	1.0
1999.....	1.4	2.0	0.9	1.4	1.8	0.9	2.0	3.2	0.9

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Discharge of firearms, undetermined intent (Y22-Y24)

2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2006.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

2007.....	1.7	2.1	1.3	1.7	2.1	1.4	1.8	2.7	1.1
2006.....	1.6	2.1	1.2	1.7	2.0	1.3	1.8	2.7	1.0
2005.....	1.5	1.8	1.2	1.5	1.8	1.3	1.7	2.4	1.1
2004.....	1.6	2.0	1.2	1.7	2.1	1.2	1.8	2.6	1.1
2003.....	1.7	2.2	1.2	1.7	2.2	1.3	1.8	2.7	1.0
2002.....	1.6	2.0	1.1	1.6	2.0	1.2	2.0	2.9	1.2
2001.....	1.4	1.9	0.9	1.4	1.8	1.0	1.7	2.7	0.9
2000.....	1.3	1.8	0.8	1.2	1.7	0.8	1.8	2.8	0.9
1999.....	1.3	1.8	0.8	1.2	1.7	0.9	1.8	2.9	0.9

Operations of war and their sequelae (Y36,Y89.1)

2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.0	0.0	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

2007.....	0.8	0.9	0.8	0.8	0.9	0.7	1.4	1.6	1.4
2006.....	0.8	0.8	0.8	0.8	0.8	0.7	1.4	1.4	1.4
2005.....	0.9	0.9	0.8	0.8	0.9	0.8	1.5	1.5	1.5
2004.....	1.0	1.1	0.9	0.9	1.1	0.8	1.5	1.5	1.6
2003.....	1.0	1.1	0.9	0.9	1.0	0.8	1.5	1.5	1.6
2002.....	1.0	1.0	1.0	0.9	1.0	0.9	1.6	1.6	1.6
2001.....	1.1	1.1	1.0	1.0	1.1	0.9	2.0	2.3	1.7
2000.....	1.1	1.2	1.0	1.0	1.2	0.9	1.9	1.9	1.9
1999.....	1.0	1.1	0.9	1.0	1.1	0.9	1.8	2.0	1.7

Worktable 293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/			
	Both sexes		Male	Female	Both sexes		Male	Female	Both sexes	
									Male	Female

Alcohol-induced causes (E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65, Y15) 4/

2007.....	7.3	11.3	3.5	7.5	11.6	3.6	6.3	11.0	2.7
2006.....	7.0	10.9	3.4	7.2	11.2	3.5	6.2	10.4	3.0
2005.....	7.0	11.0	3.4	7.2	11.1	3.4	6.8	11.4	3.3
2004.....	7.0	11.0	3.3	7.1	11.0	3.3	7.2	12.3	3.1
2003.....	7.0	11.0	3.3	7.0	11.0	3.3	7.4	12.8	3.3
2002.....	6.9	11.0	3.3	6.9	10.9	3.2	7.8	13.1	3.6
2001.....	7.0	11.2	3.3	6.9	10.9	3.3	8.9	15.1	3.9
2000.....	7.0	11.4	3.2	6.9	11.1	3.0	9.1	15.3	4.3
1999.....	7.1	11.5	3.2	6.8	11.0	3.0	9.8	16.7	4.5

Drug-induced causes (D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F17.0, F17.3-F17.5, F17.7-F17.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, F19.7-F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, K85.3, L10.5, L27.0-L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2, R78.1-R78.5, X40-X44, X60-X64, X85, Y10-Y14) 4/

2007.....	12.6	15.8	9.3	13.6	16.9	10.2	11.0	15.4	7.3
2006.....	12.7	16.4	9.0	13.5	17.1	9.8	12.9	18.9	7.7
2005.....	11.3	14.4	8.1	11.9	15.1	8.7	11.2	15.8	7.2
2004.....	10.4	13.3	7.6	11.1	13.9	8.1	10.1	14.3	6.6
2003.....	9.9	12.8	7.0	10.4	13.4	7.4	9.9	14.1	6.4
2002.....	9.0	11.7	6.3	9.4	12.1	6.7	9.9	14.2	6.1
2001.....	7.6	10.1	5.1	7.8	10.2	5.3	9.2	13.6	5.4
2000.....	7.0	9.5	4.6	7.1	9.4	4.7	9.0	13.5	5.2
1999.....	6.8	9.4	4.4	6.8	9.2	4.4	9.3	14.3	5.1

Injury by firearms (*U01.4, W32-W34, X72-X74, X93-X95, Y22-Y24, Y35.0) 4/

2007.....	10.2	18.2	2.7	8.9	15.6	2.6	19.5	36.2	3.8
2006.....	10.2	18.1	2.7	8.7	15.3	2.6	20.1	37.4	4.0
2005.....	10.2	18.3	2.7	8.9	15.7	2.6	19.4	36.4	3.6
2004.....	10.0	17.7	2.7	8.8	15.4	2.7	18.4	34.5	3.6
2003.....	10.3	18.4	2.7	9.0	16.0	2.6	19.0	35.6	3.8
2002.....	10.4	18.6	2.8	9.2	16.2	2.7	19.3	36.0	4.1
2001.....	10.3	18.5	2.8	9.2	16.3	2.7	18.4	34.5	3.8
2000.....	10.2	18.1	2.8	9.0	15.9	2.7	18.4	34.2	3.9
1999.....	10.3	18.4	2.9	9.1	16.2	2.7	18.4	34.1	4.3

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Includes races other than white and black.

2/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data--mortality."

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

4/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.