

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,712,630	100.0	844.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	633,842	23.4	197.2
2	Malignant neoplasms (C00-C97)	595,930	22.0	185.4
3	Chronic lower respiratory diseases (J40-J47)	155,041	5.7	48.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146,571	5.4	45.6
5	Cerebrovascular diseases (I60-I69)	140,323	5.2	43.7
6	Alzheimer's disease (G30)	110,561	4.1	34.4
7	Diabetes mellitus (E10-E14)	79,535	2.9	24.7
8	Influenza and pneumonia (J09-J18)	57,062	2.1	17.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49,959	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44,193	1.6	13.7
11	Septicemia (A40-A41)	40,773	1.5	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	40,326	1.5	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32,200	1.2	10.0
14	Parkinson's disease (G20-G21)	27,972	1.0	8.7
15	Pneumonitis due to solids and liquids (J69)	19,803	0.7	6.2
...	All other causes (residual)	538,539	19.9	167.6
All races and origins, both sexes, 1-4 years				
...	All causes	3,965	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,235	31.1	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	435	11.0	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	369	9.3	2.3
4	Malignant neoplasms (C00-C97)	354	8.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	3.7	0.9
6	Influenza and pneumonia (J09-J18)	88	2.2	0.6
7	Septicemia (A40-A41)	54	1.4	0.3
8	Certain conditions originating in the perinatal period (P00-P96)	50	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	42	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	40	1.0	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.9	0.2
12	Anemias (D50-D64)	25	0.6	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	17	0.4	*
15	Meningitis (G00,G03)	16	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
...	All other causes (residual)	1,025	25.9	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-9 years				
...	All causes	2,402	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	755	31.4	3.7
2	Malignant neoplasms (C00-C97)	437	18.2	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	7.5	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	5.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.5	0.4
6	Chronic lower respiratory diseases (J40-J47)	80	3.3	0.4
7	Influenza and pneumonia (J09-J18)	44	1.8	0.2
8	Cerebrovascular diseases (I60-I69)	42	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.6	0.2
10	Septicemia (A40-A41)	31	1.3	0.2
11	Anemias (D50-D64)	22	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	13	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
14	Meningitis (G00,G03)	9	0.4	*
15	Diabetes mellitus (E10-E14)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (residual)	504	21.0	2.5
All races and origins, both sexes, 10-14 years				
...	All causes	3,009	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	25.4	3.7
2	Malignant neoplasms (C00-C97)	428	14.2	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	409	13.6	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	5.3	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	5.2	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	4.2	0.6
7	Chronic lower respiratory diseases (J40-J47)	93	3.1	0.5
8	Cerebrovascular diseases (I60-I69)	42	1.4	0.2
9	Influenza and pneumonia (J09-J18)	39	1.3	0.2
10	Septicemia (A40-A41)	33	1.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.2
12	Diabetes mellitus (E10-E14)	18	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
14	Anemias (D50-D64)	11	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	8	0.3	*
...	All other causes (residual)	680	22.6	3.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 15-19 years				
...	All causes	10,186	100.0	48.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,919	38.5	18.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,061	20.2	9.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,587	15.6	7.5
4	Malignant neoplasms (C00-C97)	583	5.7	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	3.0	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	1.9	0.9
7	Influenza and pneumonia (J09-J18)	72	0.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	63	0.6	0.3
9	Cerebrovascular diseases (I60-I69)	61	0.6	0.3
10	Diabetes mellitus (E10-E14)	52	0.5	0.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	41	0.4	0.2
12	Septicemia (A40-A41)	39	0.4	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	0.2
14	Anemias (D50-D64)	27	0.3	0.1
14	Legal intervention (Y35,Y89.0)	27	0.3	0.1
...	All other causes (residual)	1,114	10.9	5.3
All races and origins, both sexes, 20-24 years				
...	All causes	20,308	100.0	89.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,595	42.3	37.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,430	16.9	15.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,146	15.5	13.8
4	Malignant neoplasms (C00-C97)	886	4.4	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	691	3.4	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.9	0.8
7	Diabetes mellitus (E10-E14)	144	0.7	0.6
8	Chronic lower respiratory diseases (J40-J47)	139	0.7	0.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	120	0.6	0.5
10	Influenza and pneumonia (J09-J18)	112	0.6	0.5
11	Cerebrovascular diseases (I60-I69)	105	0.5	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	96	0.5	0.4
13	Septicemia (A40-A41)	91	0.4	0.4
14	Legal intervention (Y35,Y89.0)	61	0.3	0.3
15	Anemias (D50-D64)	57	0.3	0.3
...	All other causes (residual)	2,444	12.0	10.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-29 years				
...	All causes	23,898	100.0	106.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,933	41.6	44.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,437	14.4	15.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,695	11.3	12.0
4	Malignant neoplasms (C00-C97)	1,334	5.6	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,281	5.4	5.7
6	Diabetes mellitus (E10-E14)	282	1.2	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	221	0.9	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	218	0.9	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	0.9	0.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	204	0.9	0.9
11	Cerebrovascular diseases (I60-I69)	192	0.8	0.9
12	Influenza and pneumonia (J09-J18)	176	0.7	0.8
13	Septicemia (A40-A41)	147	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	128	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.3	0.4
15	Legal intervention (Y35,Y89.0)	83	0.3	0.4
...	All other causes (residual)	3,273	13.7	14.6
All races and origins, both sexes, 30-34 years				
...	All causes	27,619	100.0	127.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,862	35.7	45.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,510	12.7	16.2
3	Malignant neoplasms (C00-C97)	2,370	8.6	10.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,241	8.1	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,168	7.8	10.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	623	2.3	2.9
7	Diabetes mellitus (E10-E14)	516	1.9	2.4
8	Cerebrovascular diseases (I60-I69)	375	1.4	1.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	311	1.1	1.4
10	Septicemia (A40-A41)	232	0.8	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	0.8	1.1
12	Influenza and pneumonia (J09-J18)	221	0.8	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	220	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	160	0.6	0.7
...	All other causes (residual)	4,399	15.9	20.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 35-39 years				
...	All causes	31,417	100.0	154.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,179	29.2	45.1
2	Malignant neoplasms (C00-C97)	3,870	12.3	19.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,666	11.7	18.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,341	10.6	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,640	5.2	8.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,004	3.2	4.9
7	Diabetes mellitus (E10-E14)	718	2.3	3.5
8	Cerebrovascular diseases (I60-I69)	639	2.0	3.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	453	1.4	2.2
10	Septicemia (A40-A41)	292	0.9	1.4
11	Influenza and pneumonia (J09-J18)	282	0.9	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	0.8	1.2
13	Chronic lower respiratory diseases (J40-J47)	238	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	220	0.7	1.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	197	0.6	1.0
...	All other causes (residual)	5,429	17.3	26.6
All races and origins, both sexes, 40-44 years				
...	All causes	41,671	100.0	206.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,639	20.7	42.7
2	Malignant neoplasms (C00-C97)	7,039	16.9	34.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,721	16.1	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,595	8.6	17.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,857	4.5	9.2
6	Diabetes mellitus (E10-E14)	1,268	3.0	6.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,255	3.0	6.2
8	Cerebrovascular diseases (I60-I69)	1,149	2.8	5.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	602	1.4	3.0
10	Septicemia (A40-A41)	537	1.3	2.7
11	Chronic lower respiratory diseases (J40-J47)	464	1.1	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	437	1.0	2.2
13	Influenza and pneumonia (J09-J18)	426	1.0	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	0.7	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	0.6	1.3
...	All other causes (residual)	7,120	17.1	35.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 45-49 years				
...	All causes	64,377	100.0	308.7
1	Malignant neoplasms (C00-C97)	13,769	21.4	66.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,006	18.6	57.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,832	15.3	47.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,011	6.2	19.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,236	5.0	15.5
6	Diabetes mellitus (E10-E14)	2,269	3.5	10.9
7	Cerebrovascular diseases (I60-I69)	1,902	3.0	9.1
8	Chronic lower respiratory diseases (J40-J47)	1,107	1.7	5.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,098	1.7	5.3
10	Septicemia (A40-A41)	929	1.4	4.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	880	1.4	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	774	1.2	3.7
13	Influenza and pneumonia (J09-J18)	752	1.2	3.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	491	0.8	2.4
15	Viral hepatitis (B15-B19)	447	0.7	2.1
...	All other causes (residual)	10,874	16.9	52.1
All races and origins, both sexes, 50-54 years				
...	All causes	110,117	100.0	493.0
1	Malignant neoplasms (C00-C97)	29,285	26.6	131.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,242	20.2	99.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,667	10.6	52.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,638	5.1	25.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,740	4.3	21.2
6	Diabetes mellitus (E10-E14)	3,943	3.6	17.7
7	Cerebrovascular diseases (I60-I69)	3,405	3.1	15.2
8	Chronic lower respiratory diseases (J40-J47)	3,238	2.9	14.5
9	Septicemia (A40-A41)	1,613	1.5	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,350	1.2	6.0
11	Influenza and pneumonia (J09-J18)	1,298	1.2	5.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,147	1.0	5.1
13	Viral hepatitis (B15-B19)	1,049	1.0	4.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,008	0.9	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	986	0.9	4.4
...	All other causes (residual)	17,508	15.9	78.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 55-59 years				
...	All causes	159,589	100.0	731.8
1	Malignant neoplasms (C00-C97)	50,146	31.4	229.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,455	21.0	153.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,952	6.9	50.2
4	Chronic lower respiratory diseases (J40-J47)	6,795	4.3	31.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,789	4.3	31.1
6	Diabetes mellitus (E10-E14)	6,043	3.8	27.7
7	Cerebrovascular diseases (I60-I69)	5,153	3.2	23.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,458	2.8	20.4
9	Septicemia (A40-A41)	2,480	1.6	11.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,225	1.4	10.2
11	Influenza and pneumonia (J09-J18)	1,927	1.2	8.8
12	Viral hepatitis (B15-B19)	1,683	1.1	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,526	1.0	7.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,085	0.7	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	790	0.5	3.6
...	All other causes (residual)	24,082	15.1	110.4
All races and origins, both sexes, 60-64 years				
...	All causes	198,196	100.0	1,039.3
1	Malignant neoplasms (C00-C97)	65,976	33.3	346.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,417	21.9	227.7
3	Chronic lower respiratory diseases (J40-J47)	10,662	5.4	55.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,536	4.3	44.8
5	Diabetes mellitus (E10-E14)	8,123	4.1	42.6
6	Cerebrovascular diseases (I60-I69)	6,963	3.5	36.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,489	3.3	34.0
8	Septicemia (A40-A41)	3,294	1.7	17.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,281	1.7	17.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,227	1.6	16.9
11	Influenza and pneumonia (J09-J18)	2,674	1.3	14.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,069	1.0	10.8
13	Viral hepatitis (B15-B19)	1,876	0.9	9.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	956	0.5	5.0
15	Pneumonitis due to solids and liquids (J69)	869	0.4	4.6
...	All other causes (residual)	29,784	15.0	156.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65-69 years				
...	All causes	235,482	100.0	1,465.6
1	Malignant neoplasms (C00-C97)	80,683	34.3	502.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,925	21.6	316.9
3	Chronic lower respiratory diseases (J40-J47)	15,999	6.8	99.6
4	Diabetes mellitus (E10-E14)	9,711	4.1	60.4
5	Cerebrovascular diseases (I60-I69)	9,002	3.8	56.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,940	2.9	43.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,928	2.1	30.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,467	1.9	27.8
9	Septicemia (A40-A41)	4,116	1.7	25.6
10	Influenza and pneumonia (J09-J18)	3,603	1.5	22.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,442	1.0	15.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,419	1.0	15.1
13	Alzheimer's disease (G30)	1,707	0.7	10.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,332	0.6	8.3
15	Parkinson's disease (G20-G21)	1,225	0.5	7.6
...	All other causes (residual)	35,983	15.3	223.9
All races and origins, both sexes, 70-74 years				
...	All causes	259,534	100.0	2,260.1
1	Malignant neoplasms (C00-C97)	83,045	32.0	723.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,378	21.7	491.0
3	Chronic lower respiratory diseases (J40-J47)	21,643	8.3	188.5
4	Cerebrovascular diseases (I60-I69)	11,791	4.5	102.7
5	Diabetes mellitus (E10-E14)	9,742	3.8	84.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,021	2.3	52.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,195	2.0	45.2
8	Septicemia (A40-A41)	4,539	1.7	39.5
9	Influenza and pneumonia (J09-J18)	4,528	1.7	39.4
10	Alzheimer's disease (G30)	4,460	1.7	38.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,481	1.3	30.3
12	Parkinson's disease (G20-G21)	2,588	1.0	22.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,560	1.0	22.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,824	0.7	15.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,759	0.7	15.3
...	All other causes (residual)	39,980	15.4	348.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 75-79 years				
...	All causes	290,405	100.0	3,574.7
1	Malignant neoplasms (C00-C97)	79,425	27.3	977.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,746	22.3	797.0
3	Chronic lower respiratory diseases (J40-J47)	24,259	8.4	298.6
4	Cerebrovascular diseases (I60-I69)	15,872	5.5	195.4
5	Alzheimer's disease (G30)	9,887	3.4	121.7
6	Diabetes mellitus (E10-E14)	9,786	3.4	120.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,899	2.4	84.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,230	2.1	76.7
9	Influenza and pneumonia (J09-J18)	5,857	2.0	72.1
10	Septicemia (A40-A41)	5,040	1.7	62.0
11	Parkinson's disease (G20-G21)	4,666	1.6	57.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,176	1.1	39.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,591	0.9	31.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,157	0.7	26.6
15	Pneumonitis due to solids and liquids (J69)	2,130	0.7	26.2
...	All other causes (residual)	47,684	16.4	587.0
All races and origins, both sexes, 80-84 years				
...	All causes	347,161	100.0	5,986.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,453	24.3	1,456.3
2	Malignant neoplasms (C00-C97)	73,843	21.3	1,273.3
3	Chronic lower respiratory diseases (J40-J47)	25,573	7.4	441.0
4	Cerebrovascular diseases (I60-I69)	22,140	6.4	381.8
5	Alzheimer's disease (G30)	19,616	5.7	338.2
6	Diabetes mellitus (E10-E14)	10,118	2.9	174.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,619	2.5	148.6
8	Influenza and pneumonia (J09-J18)	8,292	2.4	143.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,651	2.2	131.9
10	Parkinson's disease (G20-G21)	6,807	2.0	117.4
11	Septicemia (A40-A41)	5,441	1.6	93.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,357	1.3	75.1
13	Pneumonitis due to solids and liquids (J69)	3,022	0.9	52.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,680	0.8	46.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,855	0.5	32.0
...	All other causes (residual)	62,694	18.1	1,081.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 85-89 years				
...	All causes	390,747	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	105,700	27.1	---
2	Malignant neoplasms (C00-C97)	60,106	15.4	---
3	Alzheimer's disease (G30)	29,490	7.5	---
4	Cerebrovascular diseases (I60-I69)	27,282	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	23,972	6.1	---
6	Influenza and pneumonia (J09-J18)	10,727	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,523	2.7	---
8	Diabetes mellitus (E10-E14)	9,164	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,444	2.2	---
10	Parkinson's disease (G20-G21)	7,008	1.8	---
11	Septicemia (A40-A41)	5,976	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,483	1.4	---
13	Pneumonitis due to solids and liquids (J69)	4,141	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,773	0.7	---
15	Aortic aneurysm and dissection (I71)	1,449	0.4	---
...	All other causes (residual)	78,509	20.1	---
All races and origins, both sexes, 90-94 years				
...	All causes	313,796	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94,357	30.1	---
2	Malignant neoplasms (C00-C97)	32,626	10.4	---
3	Alzheimer's disease (G30)	28,810	9.2	---
4	Cerebrovascular diseases (I60-I69)	22,693	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	14,841	4.7	---
6	Influenza and pneumonia (J09-J18)	10,010	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,381	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,461	2.1	---
9	Diabetes mellitus (E10-E14)	5,633	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,304	1.7	---
11	Septicemia (A40-A41)	4,083	1.3	---
12	Parkinson's disease (G20-G21)	3,951	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,731	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,893	0.6	---
15	Atherosclerosis (I70)	1,291	0.4	---
...	All other causes (residual)	69,731	22.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 95-99 years				
...	All causes	127,628	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,150	32.2	---
2	Alzheimer's disease (G30)	12,838	10.1	---
3	Cerebrovascular diseases (I60-I69)	9,476	7.4	---
4	Malignant neoplasms (C00-C97)	8,532	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	4,743	3.7	---
6	Influenza and pneumonia (J09-J18)	4,585	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,396	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,534	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,325	1.8	---
10	Diabetes mellitus (E10-E14)	1,707	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,540	1.2	---
12	Septicemia (A40-A41)	1,364	1.1	---
13	Parkinson's disease (G20-G21)	935	0.7	---
14	Atherosclerosis (I70)	632	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	599	0.5	---
...	All other causes (residual)	31,272	24.5	---
All races and origins, both sexes, 100 years and over				
...	All causes	27,530	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,429	34.2	---
2	Alzheimer's disease (G30)	2,687	9.8	---
3	Cerebrovascular diseases (I60-I69)	1,900	6.9	---
4	Influenza and pneumonia (J09-J18)	1,172	4.3	---
5	Malignant neoplasms (C00-C97)	1,129	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	774	2.8	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	682	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	616	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	485	1.8	---
10	Pneumonitis due to solids and liquids (J69)	324	1.2	---
11	Diabetes mellitus (E10-E14)	281	1.0	---
12	Septicemia (A40-A41)	258	0.9	---
13	Atherosclerosis (I70)	207	0.8	---
14	Nutritional deficiencies (E40-E64)	165	0.6	---
15	Parkinson's disease (G20-G21)	89	0.3	---
...	All other causes (residual)	7,332	26.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, all ages				
...	All causes	1,373,404	100.0	868.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	335,002	24.4	211.7
2	Malignant neoplasms (C00-C97)	313,818	22.8	198.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92,919	6.8	58.7
4	Chronic lower respiratory diseases (J40-J47)	72,498	5.3	45.8
5	Cerebrovascular diseases (I60-I69)	58,288	4.2	36.8
6	Diabetes mellitus (E10-E14)	43,123	3.1	27.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,994	2.5	21.5
8	Alzheimer's disease (G30)	33,690	2.5	21.3
9	Influenza and pneumonia (J09-J18)	26,903	2.0	17.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	25,666	1.9	16.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,441	1.9	16.1
12	Septicemia (A40-A41)	19,385	1.4	12.3
13	Parkinson's disease (G20-G21)	16,867	1.2	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,274	1.0	9.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,934	1.0	8.8
...	All other causes (residual)	247,602	18.0	156.5
All races and origins, male, 1-4 years				
...	All causes	2,281	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	33.7	9.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	10.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	10.0	2.8
4	Malignant neoplasms (C00-C97)	196	8.6	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	3.4	1.0
6	Influenza and pneumonia (J09-J18)	49	2.1	0.6
7	Certain conditions originating in the perinatal period (P00-P96)	34	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	27	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
10	Septicemia (A40-A41)	24	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
12	Anemias (D50-D64)	12	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	11	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (residual)	561	24.6	6.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 5-9 years				
...	All causes	1,377	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	441	32.0	4.2
2	Malignant neoplasms (C00-C97)	242	17.6	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	7.6	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	6.2	0.8
5	Chronic lower respiratory diseases (J40-J47)	45	3.3	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.0	0.4
7	Influenza and pneumonia (J09-J18)	21	1.5	0.2
8	Cerebrovascular diseases (I60-I69)	19	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.3	*
10	Septicemia (A40-A41)	14	1.0	*
11	Anemias (D50-D64)	10	0.7	*
12	Meningitis (G00,G03)	7	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
...	All other causes (residual)	312	22.7	3.0
All races and origins, male, 10-14 years				
...	All causes	1,776	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	498	28.0	4.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	14.1	2.4
3	Malignant neoplasms (C00-C97)	227	12.8	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.1	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	5.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.8	0.6
7	Chronic lower respiratory diseases (J40-J47)	59	3.3	0.6
8	Cerebrovascular diseases (I60-I69)	26	1.5	0.2
9	Influenza and pneumonia (J09-J18)	20	1.1	0.2
10	Septicemia (A40-A41)	19	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.8	*
12	Diabetes mellitus (E10-E14)	11	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Anemias (D50-D64)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (residual)	367	20.7	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-19 years				
...	All causes	7,187	100.0	66.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,735	38.1	25.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,537	21.4	14.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,373	19.1	12.7
4	Malignant neoplasms (C00-C97)	346	4.8	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	1.5	1.0
7	Influenza and pneumonia (J09-J18)	41	0.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	36	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	32	0.4	0.3
10	Diabetes mellitus (E10-E14)	31	0.4	0.3
11	Legal intervention (Y35,Y89.0)	25	0.3	0.2
12	Septicemia (A40-A41)	22	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.3	*
14	Anemias (D50-D64)	10	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.1	*
...	All other causes (residual)	662	9.2	6.1
All races and origins, male, 20-24 years				
...	All causes	15,159	100.0	129.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,414	42.3	55.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,822	18.6	24.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,761	18.2	23.7
4	Malignant neoplasms (C00-C97)	534	3.5	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	468	3.1	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	0.7	1.0
7	Chronic lower respiratory diseases (J40-J47)	94	0.6	0.8
8	Diabetes mellitus (E10-E14)	87	0.6	0.7
9	Influenza and pneumonia (J09-J18)	66	0.4	0.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.4	0.5
11	Legal intervention (Y35,Y89.0)	59	0.4	0.5
12	Cerebrovascular diseases (I60-I69)	57	0.4	0.5
13	Septicemia (A40-A41)	49	0.3	0.4
14	Anemias (D50-D64)	32	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.2	0.2
...	All other causes (residual)	1,516	10.0	13.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 25-29 years				
...	All causes	17,173	100.0	150.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,505	43.7	65.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,774	16.2	24.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,298	13.4	20.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	831	4.8	7.3
5	Malignant neoplasms (C00-C97)	711	4.1	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	173	1.0	1.5
7	Diabetes mellitus (E10-E14)	163	0.9	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.8	1.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	118	0.7	1.0
10	Cerebrovascular diseases (I60-I69)	104	0.6	0.9
11	Influenza and pneumonia (J09-J18)	103	0.6	0.9
12	Legal intervention (Y35,Y89.0)	79	0.5	0.7
13	Septicemia (A40-A41)	77	0.4	0.7
14	Chronic lower respiratory diseases (J40-J47)	72	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.4
...	All other causes (residual)	1,995	11.6	17.5
All races and origins, male, 30-34 years				
...	All causes	18,608	100.0	170.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,154	38.4	65.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,729	14.7	25.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,799	9.7	16.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,466	7.9	13.5
5	Malignant neoplasms (C00-C97)	1,107	5.9	10.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	402	2.2	3.7
7	Diabetes mellitus (E10-E14)	301	1.6	2.8
8	Cerebrovascular diseases (I60-I69)	216	1.2	2.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	207	1.1	1.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.7	1.2
11	Influenza and pneumonia (J09-J18)	131	0.7	1.2
12	Septicemia (A40-A41)	116	0.6	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.5	0.9
14	Chronic lower respiratory diseases (J40-J47)	90	0.5	0.8
15	Legal intervention (Y35,Y89.0)	78	0.4	0.7
...	All other causes (residual)	2,580	13.9	23.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 35-39 years				
...	All causes	20,190	100.0	198.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,619	32.8	65.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,537	12.6	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,485	12.3	24.4
4	Malignant neoplasms (C00-C97)	1,688	8.4	16.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,363	6.8	13.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	637	3.2	6.3
7	Diabetes mellitus (E10-E14)	430	2.1	4.2
8	Cerebrovascular diseases (I60-I69)	333	1.6	3.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	287	1.4	2.8
10	Influenza and pneumonia (J09-J18)	167	0.8	1.6
11	Septicemia (A40-A41)	149	0.7	1.5
12	Chronic lower respiratory diseases (J40-J47)	130	0.6	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	0.6	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	0.6	1.1
...	All other causes (residual)	3,001	14.9	29.5
All races and origins, male, 40-44 years				
...	All causes	25,480	100.0	254.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,947	23.3	59.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,609	18.1	46.0
3	Malignant neoplasms (C00-C97)	2,968	11.6	29.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,686	10.5	26.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,174	4.6	11.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	970	3.8	9.7
7	Diabetes mellitus (E10-E14)	812	3.2	8.1
8	Cerebrovascular diseases (I60-I69)	657	2.6	6.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	404	1.6	4.0
10	Septicemia (A40-A41)	249	1.0	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	1.0	2.4
12	Influenza and pneumonia (J09-J18)	235	0.9	2.3
13	Chronic lower respiratory diseases (J40-J47)	204	0.8	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	198	0.8	2.0
15	Aortic aneurysm and dissection (I71)	168	0.7	1.7
...	All other causes (residual)	3,956	15.5	39.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-49 years				
...	All causes	38,807	100.0	375.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,279	21.3	80.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,645	17.1	64.3
3	Malignant neoplasms (C00-C97)	6,372	16.4	61.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,939	7.6	28.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,022	5.2	19.6
6	Diabetes mellitus (E10-E14)	1,383	3.6	13.4
7	Cerebrovascular diseases (I60-I69)	1,027	2.6	9.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	835	2.2	8.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	646	1.7	6.3
10	Chronic lower respiratory diseases (J40-J47)	492	1.3	4.8
11	Septicemia (A40-A41)	479	1.2	4.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	452	1.2	4.4
13	Influenza and pneumonia (J09-J18)	423	1.1	4.1
14	Viral hepatitis (B15-B19)	325	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	318	0.8	3.1
...	All other causes (residual)	6,170	15.9	59.7
All races and origins, male, 50-54 years				
...	All causes	66,740	100.0	608.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,592	23.4	142.2
2	Malignant neoplasms (C00-C97)	14,799	22.2	135.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,719	11.6	70.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,743	5.6	34.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,472	5.2	31.7
6	Diabetes mellitus (E10-E14)	2,501	3.7	22.8
7	Cerebrovascular diseases (I60-I69)	1,873	2.8	17.1
8	Chronic lower respiratory diseases (J40-J47)	1,512	2.3	13.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	869	1.3	7.9
10	Septicemia (A40-A41)	846	1.3	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	813	1.2	7.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	791	1.2	7.2
13	Influenza and pneumonia (J09-J18)	746	1.1	6.8
14	Viral hepatitis (B15-B19)	702	1.1	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	627	0.9	5.7
...	All other causes (residual)	10,135	15.2	92.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 55-59 years				
...	All causes	97,172	100.0	916.9
1	Malignant neoplasms (C00-C97)	27,105	27.9	255.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,330	24.0	220.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,468	7.7	70.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,611	4.7	43.5
5	Diabetes mellitus (E10-E14)	3,755	3.9	35.4
6	Chronic lower respiratory diseases (J40-J47)	3,333	3.4	31.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,260	3.4	30.8
8	Cerebrovascular diseases (I60-I69)	2,973	3.1	28.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,332	1.4	12.6
10	Septicemia (A40-A41)	1,312	1.4	12.4
11	Viral hepatitis (B15-B19)	1,171	1.2	11.0
12	Influenza and pneumonia (J09-J18)	1,076	1.1	10.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	999	1.0	9.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	844	0.9	8.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	582	0.6	5.5
...	All other causes (residual)	14,021	14.4	132.3
All races and origins, male, 60-64 years				
...	All causes	120,454	100.0	1,321.2
1	Malignant neoplasms (C00-C97)	37,075	30.8	406.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,778	24.7	326.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,965	5.0	65.4
4	Chronic lower respiratory diseases (J40-J47)	5,672	4.7	62.2
5	Diabetes mellitus (E10-E14)	4,931	4.1	54.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,465	3.7	49.0
7	Cerebrovascular diseases (I60-I69)	3,997	3.3	43.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,436	2.0	26.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,851	1.5	20.3
10	Septicemia (A40-A41)	1,791	1.5	19.6
11	Influenza and pneumonia (J09-J18)	1,507	1.3	16.5
12	Viral hepatitis (B15-B19)	1,340	1.1	14.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.1	14.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	576	0.5	6.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	555	0.5	6.1
...	All other causes (residual)	17,235	14.3	189.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 65-69 years				
...	All causes	137,630	100.0	1,811.8
1	Malignant neoplasms (C00-C97)	44,927	32.6	591.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,460	24.3	440.5
3	Chronic lower respiratory diseases (J40-J47)	8,340	6.1	109.8
4	Diabetes mellitus (E10-E14)	5,794	4.2	76.3
5	Cerebrovascular diseases (I60-I69)	4,990	3.6	65.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,648	3.4	61.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,248	2.4	42.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,431	1.8	32.0
9	Septicemia (A40-A41)	2,159	1.6	28.4
10	Influenza and pneumonia (J09-J18)	1,974	1.4	26.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,914	1.4	25.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,391	1.0	18.3
13	Parkinson's disease (G20-G21)	831	0.6	10.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	791	0.6	10.4
15	Alzheimer's disease (G30)	739	0.5	9.7
...	All other causes (residual)	19,993	14.5	263.2
All races and origins, male, 70-74 years				
...	All causes	144,717	100.0	2,732.5
1	Malignant neoplasms (C00-C97)	45,541	31.5	859.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,825	24.1	657.6
3	Chronic lower respiratory diseases (J40-J47)	10,789	7.5	203.7
4	Cerebrovascular diseases (I60-I69)	6,130	4.2	115.7
5	Diabetes mellitus (E10-E14)	5,564	3.8	105.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,740	2.6	70.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,847	2.0	53.8
8	Influenza and pneumonia (J09-J18)	2,439	1.7	46.1
9	Septicemia (A40-A41)	2,341	1.6	44.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,100	1.5	39.7
11	Alzheimer's disease (G30)	1,853	1.3	35.0
12	Parkinson's disease (G20-G21)	1,739	1.2	32.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,458	1.0	27.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,340	0.9	25.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,072	0.7	20.2
...	All other causes (residual)	20,939	14.5	395.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 75-79 years				
...	All causes	153,719	100.0	4,257.1
1	Malignant neoplasms (C00-C97)	42,788	27.8	1,185.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,459	24.4	1,037.4
3	Chronic lower respiratory diseases (J40-J47)	11,778	7.7	326.2
4	Cerebrovascular diseases (I60-I69)	7,580	4.9	209.9
5	Diabetes mellitus (E10-E14)	5,189	3.4	143.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,023	2.6	111.4
7	Alzheimer's disease (G30)	3,922	2.6	108.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,228	2.1	89.4
9	Influenza and pneumonia (J09-J18)	3,079	2.0	85.3
10	Parkinson's disease (G20-G21)	3,054	2.0	84.6
11	Septicemia (A40-A41)	2,465	1.6	68.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,521	1.0	42.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,445	0.9	40.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,251	0.8	34.6
15	Pneumonitis due to solids and liquids (J69)	1,241	0.8	34.4
...	All other causes (residual)	23,696	15.4	656.2
All races and origins, male, 80-84 years				
...	All causes	170,127	100.0	7,051.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,127	25.9	1,829.0
2	Malignant neoplasms (C00-C97)	38,881	22.9	1,611.5
3	Chronic lower respiratory diseases (J40-J47)	11,874	7.0	492.2
4	Cerebrovascular diseases (I60-I69)	9,420	5.5	390.4
5	Alzheimer's disease (G30)	7,182	4.2	297.7
6	Diabetes mellitus (E10-E14)	5,148	3.0	213.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,501	2.6	186.6
8	Parkinson's disease (G20-G21)	4,302	2.5	178.3
9	Influenza and pneumonia (J09-J18)	4,184	2.5	173.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,953	2.3	163.8
11	Septicemia (A40-A41)	2,590	1.5	107.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,792	1.1	74.3
13	Pneumonitis due to solids and liquids (J69)	1,732	1.0	71.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,456	0.9	60.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	947	0.6	39.3
...	All other causes (residual)	28,038	16.5	1,162.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 85-89 years				
...	All causes	168,932	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,106	28.5	---
2	Malignant neoplasms (C00-C97)	29,901	17.7	---
3	Chronic lower respiratory diseases (J40-J47)	10,542	6.2	---
4	Cerebrovascular diseases (I60-I69)	9,778	5.8	---
5	Alzheimer's disease (G30)	9,445	5.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,890	2.9	---
7	Influenza and pneumonia (J09-J18)	4,828	2.9	---
8	Diabetes mellitus (E10-E14)	4,148	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,131	2.4	---
10	Parkinson's disease (G20-G21)	4,092	2.4	---
11	Septicemia (A40-A41)	2,559	1.5	---
12	Pneumonitis due to solids and liquids (J69)	2,258	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,937	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,388	0.8	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	700	0.4	---
...	All other causes (residual)	30,229	17.9	---
All races and origins, male, 90-94 years				
...	All causes	112,676	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,629	31.6	---
2	Malignant neoplasms (C00-C97)	14,764	13.1	---
3	Alzheimer's disease (G30)	7,402	6.6	---
4	Cerebrovascular diseases (I60-I69)	6,652	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	5,611	5.0	---
6	Influenza and pneumonia (J09-J18)	3,935	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,279	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,799	2.5	---
9	Diabetes mellitus (E10-E14)	2,285	2.0	---
10	Parkinson's disease (G20-G21)	1,952	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,902	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,629	1.4	---
13	Septicemia (A40-A41)	1,530	1.4	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	809	0.7	---
15	Atherosclerosis (I70)	412	0.4	---
...	All other causes (residual)	22,086	19.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 95-99 years				
...	All causes	35,157	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,185	34.7	---
2	Malignant neoplasms (C00-C97)	3,301	9.4	---
3	Alzheimer's disease (G30)	2,371	6.7	---
4	Cerebrovascular diseases (I60-I69)	2,025	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	1,590	4.5	---
6	Influenza and pneumonia (J09-J18)	1,501	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,064	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	875	2.5	---
9	Pneumonitis due to solids and liquids (J69)	675	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	602	1.7	---
11	Diabetes mellitus (E10-E14)	536	1.5	---
12	Septicemia (A40-A41)	433	1.2	---
13	Parkinson's disease (G20-G21)	395	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	210	0.6	---
15	Atherosclerosis (I70)	166	0.5	---
...	All other causes (residual)	7,228	20.6	---
All races and origins, male, 100 years and over				
...	All causes	4,939	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,816	36.8	---
2	Cerebrovascular diseases (I60-I69)	315	6.4	---
3	Alzheimer's disease (G30)	311	6.3	---
4	Malignant neoplasms (C00-C97)	308	6.2	---
5	Influenza and pneumonia (J09-J18)	264	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	190	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	2.6	---
9	Pneumonitis due to solids and liquids (J69)	104	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	103	2.1	---
11	Septicemia (A40-A41)	62	1.3	---
12	Diabetes mellitus (E10-E14)	47	1.0	---
13	Atherosclerosis (I70)	44	0.9	---
14	Nutritional deficiencies (E40-E64)	29	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	---
...	All other causes (residual)	1,062	21.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,339,226	100.0	820.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	298,840	22.3	183.1
2	Malignant neoplasms (C00-C97)	282,112	21.1	172.9
3	Chronic lower respiratory diseases (J40-J47)	82,543	6.2	50.6
4	Cerebrovascular diseases (I60-I69)	82,035	6.1	50.3
5	Alzheimer's disease (G30)	76,871	5.7	47.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,652	4.0	32.9
7	Diabetes mellitus (E10-E14)	36,412	2.7	22.3
8	Influenza and pneumonia (J09-J18)	30,159	2.3	18.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,518	1.8	15.0
10	Septicemia (A40-A41)	21,388	1.6	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,266	1.4	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14,660	1.1	9.0
13	Parkinson's disease (G20-G21)	11,105	0.8	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,199	0.8	6.2
15	Pneumonitis due to solids and liquids (J69)	8,892	0.7	5.4
...	All other causes (residual)	287,574	21.5	176.2
All races and origins, female, 1-4 years				
...	All causes	1,684	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	27.7	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	12.4	2.7
3	Malignant neoplasms (C00-C97)	158	9.4	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	8.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	4.1	0.9
6	Influenza and pneumonia (J09-J18)	39	2.3	0.5
7	Septicemia (A40-A41)	30	1.8	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.0	*
9	Cerebrovascular diseases (I60-I69)	16	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
11	Anemias (D50-D64)	13	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.8	*
13	Meningitis (G00,G03)	11	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (residual)	469	27.9	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-9 years				
...	All causes	1,025	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	30.6	3.1
2	Malignant neoplasms (C00-C97)	195	19.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	7.4	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	5.4	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.3	0.4
6	Chronic lower respiratory diseases (J40-J47)	35	3.4	0.3
7	Cerebrovascular diseases (I60-I69)	23	2.2	0.2
7	Influenza and pneumonia (J09-J18)	23	2.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.0	0.2
10	Septicemia (A40-A41)	17	1.7	*
11	Anemias (D50-D64)	12	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
13	Diabetes mellitus (E10-E14)	5	0.5	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (residual)	191	18.6	1.9
All races and origins, female, 10-14 years				
...	All causes	1,233	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	21.5	2.6
2	Malignant neoplasms (C00-C97)	201	16.3	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	12.8	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	5.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	4.7	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	34	2.8	0.3
8	Influenza and pneumonia (J09-J18)	19	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
10	Cerebrovascular diseases (I60-I69)	16	1.3	*
11	Septicemia (A40-A41)	14	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
13	Anemias (D50-D64)	7	0.6	*
13	Diabetes mellitus (E10-E14)	7	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.2	*
...	All other causes (residual)	306	24.8	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 15-19 years				
...	All causes	2,999	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	39.5	11.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	17.5	5.1
3	Malignant neoplasms (C00-C97)	237	7.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	2.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	41	1.4	0.4
8	Influenza and pneumonia (J09-J18)	31	1.0	0.3
9	Cerebrovascular diseases (I60-I69)	29	1.0	0.3
10	Chronic lower respiratory diseases (J40-J47)	27	0.9	0.3
11	Diabetes mellitus (E10-E14)	21	0.7	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	0.2
13	Septicemia (A40-A41)	17	0.6	*
13	Anemias (D50-D64)	17	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (residual)	433	14.4	4.2
All races and origins, female, 20-24 years				
...	All causes	5,149	100.0	46.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,181	42.4	19.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	608	11.8	5.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	7.5	3.5
4	Malignant neoplasms (C00-C97)	352	6.8	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	223	4.3	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	120	2.3	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.6	0.7
8	Diabetes mellitus (E10-E14)	57	1.1	0.5
9	Cerebrovascular diseases (I60-I69)	48	0.9	0.4
10	Influenza and pneumonia (J09-J18)	46	0.9	0.4
11	Chronic lower respiratory diseases (J40-J47)	45	0.9	0.4
12	Septicemia (A40-A41)	42	0.8	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.7	0.3
14	Anemias (D50-D64)	25	0.5	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.4	0.2
...	All other causes (residual)	881	17.1	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-29 years				
...	All causes	6,725	100.0	60.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,428	36.1	22.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	663	9.9	6.0
3	Malignant neoplasms (C00-C97)	623	9.3	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	450	6.7	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	397	5.9	3.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	204	3.0	1.8
7	Diabetes mellitus (E10-E14)	119	1.8	1.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	103	1.5	0.9
9	Cerebrovascular diseases (I60-I69)	88	1.3	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	0.7
11	Influenza and pneumonia (J09-J18)	73	1.1	0.7
12	Septicemia (A40-A41)	70	1.0	0.6
13	Chronic lower respiratory diseases (J40-J47)	56	0.8	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.7	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.6	0.4
...	All other causes (residual)	1,282	19.1	11.6
All races and origins, female, 30-34 years				
...	All causes	9,011	100.0	83.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,708	30.1	25.1
2	Malignant neoplasms (C00-C97)	1,263	14.0	11.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	781	8.7	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	775	8.6	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	369	4.1	3.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	221	2.5	2.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	220	2.4	2.0
8	Diabetes mellitus (E10-E14)	215	2.4	2.0
9	Cerebrovascular diseases (I60-I69)	159	1.8	1.5
10	Septicemia (A40-A41)	116	1.3	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	104	1.2	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	1.1	0.9
13	Influenza and pneumonia (J09-J18)	90	1.0	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.9	0.7
15	Chronic lower respiratory diseases (J40-J47)	70	0.8	0.6
...	All other causes (residual)	1,741	19.3	16.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 35-39 years				
...	All causes	11,227	100.0	110.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,560	22.8	25.1
2	Malignant neoplasms (C00-C97)	2,182	19.4	21.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,181	10.5	11.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	804	7.2	7.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	367	3.3	3.6
6	Cerebrovascular diseases (I60-I69)	306	2.7	3.0
7	Diabetes mellitus (E10-E14)	288	2.6	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	2.5	2.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	197	1.8	1.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	166	1.5	1.6
11	Septicemia (A40-A41)	143	1.3	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	1.1	1.2
13	Influenza and pneumonia (J09-J18)	115	1.0	1.1
14	Chronic lower respiratory diseases (J40-J47)	108	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	0.9	1.0
...	All other causes (residual)	2,314	20.6	22.7
All races and origins, female, 40-44 years				
...	All causes	16,191	100.0	159.0
1	Malignant neoplasms (C00-C97)	4,071	25.1	40.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,692	16.6	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,112	13.0	20.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	909	5.6	8.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	683	4.2	6.7
6	Cerebrovascular diseases (I60-I69)	492	3.0	4.8
7	Diabetes mellitus (E10-E14)	456	2.8	4.5
8	Septicemia (A40-A41)	288	1.8	2.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	285	1.8	2.8
10	Chronic lower respiratory diseases (J40-J47)	260	1.6	2.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	198	1.2	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	1.2	1.9
13	Influenza and pneumonia (J09-J18)	191	1.2	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.7	1.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	113	0.7	1.1
...	All other causes (residual)	3,126	19.3	30.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 45-49 years				
...	All causes	25,570	100.0	243.1
1	Malignant neoplasms (C00-C97)	7,397	28.9	70.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,727	14.6	35.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,187	12.5	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,214	4.7	11.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,072	4.2	10.2
6	Diabetes mellitus (E10-E14)	886	3.5	8.4
7	Cerebrovascular diseases (I60-I69)	875	3.4	8.3
8	Chronic lower respiratory diseases (J40-J47)	615	2.4	5.8
9	Septicemia (A40-A41)	450	1.8	4.3
10	Influenza and pneumonia (J09-J18)	329	1.3	3.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	322	1.3	3.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.0	2.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	234	0.9	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	173	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	138	0.5	1.3
...	All other causes (residual)	4,688	18.3	44.6
All races and origins, female, 50-54 years				
...	All causes	43,377	100.0	381.5
1	Malignant neoplasms (C00-C97)	14,486	33.4	127.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,650	15.3	58.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,948	9.1	34.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,895	4.4	16.7
5	Chronic lower respiratory diseases (J40-J47)	1,726	4.0	15.2
6	Cerebrovascular diseases (I60-I69)	1,532	3.5	13.5
7	Diabetes mellitus (E10-E14)	1,442	3.3	12.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,268	2.9	11.2
9	Septicemia (A40-A41)	767	1.8	6.7
10	Influenza and pneumonia (J09-J18)	552	1.3	4.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	537	1.2	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	359	0.8	3.2
13	Viral hepatitis (B15-B19)	347	0.8	3.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	278	0.6	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	0.5	2.0
...	All other causes (residual)	7,365	17.0	64.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 55-59 years				
...	All causes	62,417	100.0	556.8
1	Malignant neoplasms (C00-C97)	23,041	36.9	205.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,125	16.2	90.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,484	5.6	31.1
4	Chronic lower respiratory diseases (J40-J47)	3,462	5.5	30.9
5	Diabetes mellitus (E10-E14)	2,288	3.7	20.4
6	Cerebrovascular diseases (I60-I69)	2,180	3.5	19.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,178	3.5	19.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,198	1.9	10.7
9	Septicemia (A40-A41)	1,168	1.9	10.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	893	1.4	8.0
11	Influenza and pneumonia (J09-J18)	851	1.4	7.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	527	0.8	4.7
13	Viral hepatitis (B15-B19)	512	0.8	4.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	291	0.5	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	283	0.5	2.5
...	All other causes (residual)	9,936	15.9	88.6
All races and origins, female, 60-64 years				
...	All causes	77,742	100.0	781.1
1	Malignant neoplasms (C00-C97)	28,901	37.2	290.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,639	17.5	137.0
3	Chronic lower respiratory diseases (J40-J47)	4,990	6.4	50.1
4	Diabetes mellitus (E10-E14)	3,192	4.1	32.1
5	Cerebrovascular diseases (I60-I69)	2,966	3.8	29.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,571	3.3	25.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,024	2.6	20.3
8	Septicemia (A40-A41)	1,503	1.9	15.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,376	1.8	13.8
10	Influenza and pneumonia (J09-J18)	1,167	1.5	11.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	845	1.1	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	789	1.0	7.9
13	Viral hepatitis (B15-B19)	536	0.7	5.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	401	0.5	4.0
15	Alzheimer's disease (G30)	400	0.5	4.0
...	All other causes (residual)	12,442	16.0	125.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65-69 years				
...	All causes	97,852	100.0	1,155.1
1	Malignant neoplasms (C00-C97)	35,756	36.5	422.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,465	17.8	206.2
3	Chronic lower respiratory diseases (J40-J47)	7,659	7.8	90.4
4	Cerebrovascular diseases (I60-I69)	4,012	4.1	47.4
5	Diabetes mellitus (E10-E14)	3,917	4.0	46.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,292	2.3	27.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,036	2.1	24.0
8	Septicemia (A40-A41)	1,957	2.0	23.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,680	1.7	19.8
10	Influenza and pneumonia (J09-J18)	1,629	1.7	19.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,028	1.1	12.1
12	Alzheimer's disease (G30)	968	1.0	11.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	541	0.6	6.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	528	0.5	6.2
15	Pneumonitis due to solids and liquids (J69)	446	0.5	5.3
...	All other causes (residual)	15,938	16.3	188.1
All races and origins, female, 70-74 years				
...	All causes	114,817	100.0	1,855.8
1	Malignant neoplasms (C00-C97)	37,504	32.7	606.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,553	18.8	348.4
3	Chronic lower respiratory diseases (J40-J47)	10,854	9.5	175.4
4	Cerebrovascular diseases (I60-I69)	5,661	4.9	91.5
5	Diabetes mellitus (E10-E14)	4,178	3.6	67.5
6	Alzheimer's disease (G30)	2,607	2.3	42.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,348	2.0	38.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,281	2.0	36.9
9	Septicemia (A40-A41)	2,198	1.9	35.5
10	Influenza and pneumonia (J09-J18)	2,089	1.8	33.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,381	1.2	22.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,220	1.1	19.7
13	Parkinson's disease (G20-G21)	849	0.7	13.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	752	0.7	12.2
15	Pneumonitis due to solids and liquids (J69)	578	0.5	9.3
...	All other causes (residual)	18,764	16.3	303.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 75-79 years				
...	All causes	136,686	100.0	3,028.8
1	Malignant neoplasms (C00-C97)	36,637	26.8	811.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,287	20.0	604.6
3	Chronic lower respiratory diseases (J40-J47)	12,481	9.1	276.6
4	Cerebrovascular diseases (I60-I69)	8,292	6.1	183.7
5	Alzheimer's disease (G30)	5,965	4.4	132.2
6	Diabetes mellitus (E10-E14)	4,597	3.4	101.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,002	2.2	66.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,876	2.1	63.7
9	Influenza and pneumonia (J09-J18)	2,778	2.0	61.6
10	Septicemia (A40-A41)	2,575	1.9	57.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,655	1.2	36.7
12	Parkinson's disease (G20-G21)	1,612	1.2	35.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,146	0.8	25.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	906	0.7	20.1
15	Pneumonitis due to solids and liquids (J69)	889	0.7	19.7
...	All other causes (residual)	23,988	17.5	531.5
All races and origins, female, 80-84 years				
...	All causes	177,034	100.0	5,227.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,326	22.8	1,190.7
2	Malignant neoplasms (C00-C97)	34,962	19.7	1,032.3
3	Chronic lower respiratory diseases (J40-J47)	13,699	7.7	404.5
4	Cerebrovascular diseases (I60-I69)	12,720	7.2	375.6
5	Alzheimer's disease (G30)	12,434	7.0	367.1
6	Diabetes mellitus (E10-E14)	4,970	2.8	146.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,118	2.3	121.6
8	Influenza and pneumonia (J09-J18)	4,108	2.3	121.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,698	2.1	109.2
10	Septicemia (A40-A41)	2,851	1.6	84.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,565	1.4	75.7
12	Parkinson's disease (G20-G21)	2,505	1.4	74.0
13	Pneumonitis due to solids and liquids (J69)	1,290	0.7	38.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,224	0.7	36.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	922	0.5	27.2
...	All other causes (residual)	34,642	19.6	1,022.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 85-89 years				
...	All causes	221,815	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,594	26.0	---
2	Malignant neoplasms (C00-C97)	30,205	13.6	---
3	Alzheimer's disease (G30)	20,045	9.0	---
4	Cerebrovascular diseases (I60-I69)	17,504	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	13,430	6.1	---
6	Influenza and pneumonia (J09-J18)	5,899	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,633	2.5	---
8	Diabetes mellitus (E10-E14)	5,016	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,313	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,546	1.6	---
11	Septicemia (A40-A41)	3,417	1.5	---
12	Parkinson's disease (G20-G21)	2,916	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,883	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,385	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	887	0.4	---
...	All other causes (residual)	48,142	21.7	---
All races and origins, female, 90-94 years				
...	All causes	201,120	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,728	29.2	---
2	Alzheimer's disease (G30)	21,408	10.6	---
3	Malignant neoplasms (C00-C97)	17,862	8.9	---
4	Cerebrovascular diseases (I60-I69)	16,041	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	9,230	4.6	---
6	Influenza and pneumonia (J09-J18)	6,075	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,102	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,675	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,662	1.8	---
10	Diabetes mellitus (E10-E14)	3,348	1.7	---
11	Septicemia (A40-A41)	2,553	1.3	---
12	Parkinson's disease (G20-G21)	1,999	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,829	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,084	0.5	---
15	Atherosclerosis (I70)	879	0.4	---
...	All other causes (residual)	47,645	23.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 95-99 years				
...	All causes	92,471	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,965	31.3	---
2	Alzheimer's disease (G30)	10,467	11.3	---
3	Cerebrovascular diseases (I60-I69)	7,451	8.1	---
4	Malignant neoplasms (C00-C97)	5,231	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,153	3.4	---
6	Influenza and pneumonia (J09-J18)	3,084	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,332	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,932	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,450	1.6	---
10	Diabetes mellitus (E10-E14)	1,171	1.3	---
11	Septicemia (A40-A41)	931	1.0	---
12	Pneumonitis due to solids and liquids (J69)	865	0.9	---
13	Parkinson's disease (G20-G21)	540	0.6	---
14	Atherosclerosis (I70)	466	0.5	---
15	Nutritional deficiencies (E40-E64)	403	0.4	---
...	All other causes (residual)	24,030	26.0	---
All races and origins, female, 100 years and over				
...	All causes	22,591	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,613	33.7	---
2	Alzheimer's disease (G30)	2,376	10.5	---
3	Cerebrovascular diseases (I60-I69)	1,585	7.0	---
4	Influenza and pneumonia (J09-J18)	908	4.0	---
5	Malignant neoplasms (C00-C97)	821	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	584	2.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	579	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	489	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	1.6	---
10	Diabetes mellitus (E10-E14)	234	1.0	---
11	Pneumonitis due to solids and liquids (J69)	220	1.0	---
12	Septicemia (A40-A41)	196	0.9	---
13	Atherosclerosis (I70)	163	0.7	---
14	Nutritional deficiencies (E40-E64)	136	0.6	---
15	Anemias (D50-D64)	66	0.3	---
15	Parkinson's disease (G20-G21)	66	0.3	---
...	All other causes (residual)	6,202	27.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	179,457	100.0	317.1
1	Malignant neoplasms (C00-C97)	37,804	21.1	66.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,401	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,806	7.7	24.4
4	Cerebrovascular diseases (I60-I69)	9,795	5.5	17.3
5	Diabetes mellitus (E10-E14)	8,278	4.6	14.6
6	Alzheimer's disease (G30)	6,444	3.6	11.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,018	3.4	10.6
8	Chronic lower respiratory diseases (J40-J47)	5,159	2.9	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,581	2.0	6.3
10	Influenza and pneumonia (J09-J18)	3,497	1.9	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,303	1.8	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,886	1.6	5.1
13	Septicemia (A40-A41)	2,692	1.5	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,454	1.4	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,384	1.3	4.2
...	All other causes (residual)	34,955	19.5	61.8
Hispanic, both sexes, 1-4 years				
...	All causes	815	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	31.7	6.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	14.1	2.8
3	Malignant neoplasms (C00-C97)	75	9.2	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	8.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.6	0.7
6	Influenza and pneumonia (J09-J18)	18	2.2	*
7	Septicemia (A40-A41)	17	2.1	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
11	Anemias (D50-D64)	4	0.5	*
11	Meningitis (G00,G03)	4	0.5	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	192	23.6	4.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	495	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	32.1	3.0
2	Malignant neoplasms (C00-C97)	102	20.6	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.5	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.4	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	Chronic lower respiratory diseases (J40-J47)	12	2.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.8	*
7	Cerebrovascular diseases (I60-I69)	9	1.8	*
9	Septicemia (A40-A41)	7	1.4	*
10	Influenza and pneumonia (J09-J18)	6	1.2	*
11	Meningitis (G00,G03)	4	0.8	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
13	Anemias (D50-D64)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	100	20.2	1.9
Hispanic, both sexes, 10-14 years				
...	All causes	568	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	26.2	3.0
2	Malignant neoplasms (C00-C97)	109	19.2	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	8.1	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	5.6	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	5.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	Septicemia (A40-A41)	9	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
9	Influenza and pneumonia (J09-J18)	8	1.4	*
11	Cerebrovascular diseases (I60-I69)	7	1.2	*
12	Anemias (D50-D64)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Aortic aneurysm and dissection (I71)	2	0.4	*
14	Peptic ulcer (K25-K28)	2	0.4	*
...	All other causes (residual)	130	22.9	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,868	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	742	39.7	15.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	316	16.9	6.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	299	16.0	6.3
4	Malignant neoplasms (C00-C97)	137	7.3	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.7	0.7
7	Cerebrovascular diseases (I60-I69)	18	1.0	*
8	Influenza and pneumonia (J09-J18)	14	0.7	*
9	Septicemia (A40-A41)	9	0.5	*
9	Diabetes mellitus (E10-E14)	9	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.4	*
12	Legal intervention (Y35,Y89.0)	7	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (residual)	201	10.8	4.3
Hispanic, both sexes, 20-24 years				
...	All causes	3,485	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,519	43.6	31.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	605	17.4	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	508	14.6	10.6
4	Malignant neoplasms (C00-C97)	195	5.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.5	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.1	0.8
7	Influenza and pneumonia (J09-J18)	22	0.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	22	0.6	0.5
9	Septicemia (A40-A41)	21	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.6	0.4
11	Cerebrovascular diseases (I60-I69)	14	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
13	Diabetes mellitus (E10-E14)	11	0.3	*
14	Legal intervention (Y35,Y89.0)	10	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (residual)	387	11.1	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,396	100.0	74.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,362	40.1	30.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	429	12.6	9.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	394	11.6	8.7
4	Malignant neoplasms (C00-C97)	287	8.5	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	4.7	3.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.4	1.0
7	Cerebrovascular diseases (I60-I69)	39	1.1	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.0	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.0	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.8	0.6
11	Diabetes mellitus (E10-E14)	27	0.8	0.6
12	Influenza and pneumonia (J09-J18)	26	0.8	0.6
13	Legal intervention (Y35,Y89.0)	21	0.6	0.5
14	Septicemia (A40-A41)	20	0.6	0.4
14	Chronic lower respiratory diseases (J40-J47)	20	0.6	0.4
...	All other causes (residual)	470	13.8	10.4
Hispanic, both sexes, 30-34 years				
...	All causes	3,786	100.0	85.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,291	34.1	29.0
2	Malignant neoplasms (C00-C97)	427	11.3	9.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	370	9.8	8.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	362	9.6	8.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	7.4	6.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	123	3.2	2.8
7	Cerebrovascular diseases (I60-I69)	82	2.2	1.8
8	Diabetes mellitus (E10-E14)	79	2.1	1.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.6	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.8	0.7
11	Influenza and pneumonia (J09-J18)	30	0.8	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.7	0.6
13	Septicemia (A40-A41)	26	0.7	0.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.7	0.6
15	Chronic lower respiratory diseases (J40-J47)	19	0.5	*
...	All other causes (residual)	551	14.6	12.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	4,202	100.0	99.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,133	27.0	26.7
2	Malignant neoplasms (C00-C97)	671	16.0	15.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	414	9.9	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	7.5	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	6.8	6.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	218	5.2	5.1
7	Cerebrovascular diseases (I60-I69)	118	2.8	2.8
8	Diabetes mellitus (E10-E14)	92	2.2	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	86	2.0	2.0
10	Influenza and pneumonia (J09-J18)	42	1.0	1.0
11	Septicemia (A40-A41)	40	1.0	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.9	0.9
13	Pregnancy, childbirth and the puerperium (O00-O99)	34	0.8	0.8
14	Aortic aneurysm and dissection (I71)	20	0.5	0.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.4	*
...	All other causes (residual)	675	16.1	15.9
Hispanic, both sexes, 40-44 years				
...	All causes	5,288	100.0	134.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,057	20.0	26.9
2	Malignant neoplasms (C00-C97)	1,047	19.8	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	668	12.6	17.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	376	7.1	9.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	5.1	6.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	4.1	5.6
7	Cerebrovascular diseases (I60-I69)	203	3.8	5.2
8	Diabetes mellitus (E10-E14)	182	3.4	4.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	105	2.0	2.7
10	Septicemia (A40-A41)	65	1.2	1.7
11	Influenza and pneumonia (J09-J18)	60	1.1	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.0	1.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.7	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	32	0.6	0.8
...	All other causes (residual)	876	16.6	22.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	7,159	100.0	204.8
1	Malignant neoplasms (C00-C97)	1,682	23.5	48.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,085	15.2	31.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,040	14.5	29.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	596	8.3	17.1
5	Diabetes mellitus (E10-E14)	282	3.9	8.1
6	Cerebrovascular diseases (I60-I69)	262	3.7	7.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	3.6	7.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	2.4	4.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	140	2.0	4.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	1.7	3.5
11	Viral hepatitis (B15-B19)	113	1.6	3.2
12	Septicemia (A40-A41)	107	1.5	3.1
13	Influenza and pneumonia (J09-J18)	79	1.1	2.3
14	Chronic lower respiratory diseases (J40-J47)	55	0.8	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	0.7	1.5
...	All other causes (residual)	1,120	15.6	32.0
Hispanic, both sexes, 50-54 years				
...	All causes	10,030	100.0	329.5
1	Malignant neoplasms (C00-C97)	2,604	26.0	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,759	17.5	57.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,050	10.5	34.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	881	8.8	28.9
5	Diabetes mellitus (E10-E14)	490	4.9	16.1
6	Cerebrovascular diseases (I60-I69)	403	4.0	13.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	2.5	8.2
8	Viral hepatitis (B15-B19)	167	1.7	5.5
9	Septicemia (A40-A41)	162	1.6	5.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	156	1.6	5.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	1.5	4.9
12	Influenza and pneumonia (J09-J18)	130	1.3	4.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	1.3	4.2
14	Chronic lower respiratory diseases (J40-J47)	112	1.1	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.8	2.6
...	All other causes (residual)	1,510	15.1	49.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	12,128	100.0	494.7
1	Malignant neoplasms (C00-C97)	3,602	29.7	146.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,287	18.9	93.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	913	7.5	37.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	845	7.0	34.5
5	Diabetes mellitus (E10-E14)	659	5.4	26.9
6	Cerebrovascular diseases (I60-I69)	513	4.2	20.9
7	Viral hepatitis (B15-B19)	262	2.2	10.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	1.8	8.9
9	Chronic lower respiratory diseases (J40-J47)	203	1.7	8.3
10	Septicemia (A40-A41)	188	1.6	7.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	1.5	7.3
12	Influenza and pneumonia (J09-J18)	166	1.4	6.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	1.0	5.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	1.0	5.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	0.7	3.4
...	All other causes (residual)	1,759	14.5	71.7
Hispanic, both sexes, 60-64 years				
...	All causes	13,859	100.0	753.9
1	Malignant neoplasms (C00-C97)	4,306	31.1	234.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,809	20.3	152.8
3	Diabetes mellitus (E10-E14)	957	6.9	52.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	854	6.2	46.5
5	Cerebrovascular diseases (I60-I69)	594	4.3	32.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	592	4.3	32.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	2.1	15.9
8	Chronic lower respiratory diseases (J40-J47)	289	2.1	15.7
9	Septicemia (A40-A41)	233	1.7	12.7
10	Viral hepatitis (B15-B19)	231	1.7	12.6
11	Influenza and pneumonia (J09-J18)	186	1.3	10.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.3	9.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	1.0	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	84	0.6	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	3.1
...	All other causes (residual)	2,059	14.9	112.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	14,920	100.0	1,099.0
1	Malignant neoplasms (C00-C97)	4,611	30.9	339.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,132	21.0	230.7
3	Diabetes mellitus (E10-E14)	1,054	7.1	77.6
4	Cerebrovascular diseases (I60-I69)	750	5.0	55.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	661	4.4	48.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	441	3.0	32.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.8	30.3
8	Chronic lower respiratory diseases (J40-J47)	381	2.6	28.1
9	Influenza and pneumonia (J09-J18)	272	1.8	20.0
10	Septicemia (A40-A41)	268	1.8	19.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	216	1.4	15.9
12	Viral hepatitis (B15-B19)	144	1.0	10.6
13	Alzheimer's disease (G30)	117	0.8	8.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	0.7	7.3
15	Parkinson's disease (G20-G21)	80	0.5	5.9
...	All other causes (residual)	2,282	15.3	168.1
Hispanic, both sexes, 70-74 years				
...	All causes	15,355	100.0	1,678.8
1	Malignant neoplasms (C00-C97)	4,522	29.4	494.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,385	22.0	370.1
3	Diabetes mellitus (E10-E14)	1,002	6.5	109.6
4	Cerebrovascular diseases (I60-I69)	811	5.3	88.7
5	Chronic lower respiratory diseases (J40-J47)	576	3.8	63.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	492	3.2	53.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	421	2.7	46.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	2.3	39.3
9	Septicemia (A40-A41)	301	2.0	32.9
10	Alzheimer's disease (G30)	284	1.8	31.1
10	Influenza and pneumonia (J09-J18)	284	1.8	31.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	1.4	23.4
13	Parkinson's disease (G20-G21)	162	1.1	17.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.7	11.0
15	Viral hepatitis (B15-B19)	81	0.5	8.9
...	All other causes (residual)	2,360	15.4	258.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	17,242	100.0	2,670.2
1	Malignant neoplasms (C00-C97)	4,459	25.9	690.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,950	22.9	611.7
3	Cerebrovascular diseases (I60-I69)	1,126	6.5	174.4
4	Diabetes mellitus (E10-E14)	1,032	6.0	159.8
5	Chronic lower respiratory diseases (J40-J47)	694	4.0	107.5
6	Alzheimer's disease (G30)	608	3.5	94.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	447	2.6	69.2
8	Influenza and pneumonia (J09-J18)	415	2.4	64.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	406	2.4	62.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	2.0	53.3
11	Septicemia (A40-A41)	318	1.8	49.2
12	Parkinson's disease (G20-G21)	265	1.5	41.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	263	1.5	40.7
14	Pneumonitis due to solids and liquids (J69)	113	0.7	17.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	16.4
...	All other causes (residual)	2,696	15.6	417.5
Hispanic, both sexes, 80-84 years				
...	All causes	19,847	100.0	4,556.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,885	24.6	1,121.5
2	Malignant neoplasms (C00-C97)	4,031	20.3	925.5
3	Cerebrovascular diseases (I60-I69)	1,522	7.7	349.4
4	Alzheimer's disease (G30)	1,247	6.3	286.3
5	Diabetes mellitus (E10-E14)	1,021	5.1	234.4
6	Chronic lower respiratory diseases (J40-J47)	932	4.7	214.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	530	2.7	121.7
8	Influenza and pneumonia (J09-J18)	524	2.6	120.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	2.1	94.6
10	Parkinson's disease (G20-G21)	368	1.9	84.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	1.7	78.5
12	Septicemia (A40-A41)	300	1.5	68.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	248	1.2	56.9
14	Pneumonitis due to solids and liquids (J69)	167	0.8	38.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.6	29.6
...	All other causes (residual)	3,189	16.1	732.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	20,040	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,465	27.3	---
2	Malignant neoplasms (C00-C97)	3,050	15.2	---
3	Alzheimer's disease (G30)	1,728	8.6	---
4	Cerebrovascular diseases (I60-I69)	1,571	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	913	4.6	---
6	Diabetes mellitus (E10-E14)	819	4.1	---
7	Influenza and pneumonia (J09-J18)	542	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.9	---
11	Parkinson's disease (G20-G21)	352	1.8	---
12	Septicemia (A40-A41)	315	1.6	---
13	Pneumonitis due to solids and liquids (J69)	188	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	142	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	112	0.6	---
...	All other causes (residual)	3,587	17.9	---
Hispanic, both sexes, 90-94 years				
...	All causes	13,807	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,899	28.2	---
2	Alzheimer's disease (G30)	1,590	11.5	---
3	Malignant neoplasms (C00-C97)	1,491	10.8	---
4	Cerebrovascular diseases (I60-I69)	1,179	8.5	---
5	Chronic lower respiratory diseases (J40-J47)	583	4.2	---
6	Diabetes mellitus (E10-E14)	415	3.0	---
7	Influenza and pneumonia (J09-J18)	413	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	277	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	1.9	---
11	Parkinson's disease (G20-G21)	208	1.5	---
12	Septicemia (A40-A41)	181	1.3	---
13	Pneumonitis due to solids and liquids (J69)	150	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	---
15	Nutritional deficiencies (E40-E64)	53	0.4	---
...	All other causes (residual)	2,747	19.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	5,074	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,523	30.0	---
2	Alzheimer's disease (G30)	622	12.3	---
3	Cerebrovascular diseases (I60-I69)	462	9.1	---
4	Malignant neoplasms (C00-C97)	334	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	236	4.7	---
6	Influenza and pneumonia (J09-J18)	173	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	2.7	---
8	Diabetes mellitus (E10-E14)	116	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	1.8	---
11	Pneumonitis due to solids and liquids (J69)	57	1.1	---
12	Septicemia (A40-A41)	52	1.0	---
13	Parkinson's disease (G20-G21)	42	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	---
15	Nutritional deficiencies (E40-E64)	24	0.5	---
...	All other causes (residual)	1,068	21.0	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,282	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	447	34.9	---
2	Alzheimer's disease (G30)	161	12.6	---
3	Cerebrovascular diseases (I60-I69)	85	6.6	---
4	Influenza and pneumonia (J09-J18)	57	4.4	---
5	Malignant neoplasms (C00-C97)	49	3.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	38	3.0	---
8	Diabetes mellitus (E10-E14)	25	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.3	---
11	Pneumonitis due to solids and liquids (J69)	15	1.2	---
12	Septicemia (A40-A41)	12	0.9	---
13	Nutritional deficiencies (E40-E64)	10	0.8	---
14	Atherosclerosis (I70)	8	0.6	---
15	Parkinson's disease (G20-G21)	7	0.5	---
...	All other causes (residual)	292	22.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	98,170	100.0	343.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,225	20.6	70.7
2	Malignant neoplasms (C00-C97)	19,847	20.2	69.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,067	10.3	35.2
4	Cerebrovascular diseases (I60-I69)	4,544	4.6	15.9
5	Diabetes mellitus (E10-E14)	4,426	4.5	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,109	4.2	14.4
7	Chronic lower respiratory diseases (J40-J47)	2,606	2.7	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,587	2.6	9.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,391	2.4	8.4
10	Alzheimer's disease (G30)	2,035	2.1	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,837	1.9	6.4
12	Influenza and pneumonia (J09-J18)	1,735	1.8	6.1
13	Septicemia (A40-A41)	1,370	1.4	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,368	1.4	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,139	1.2	4.0
...	All other causes (residual)	17,884	18.2	62.5
Hispanic, male, 1-4 years				
...	All causes	461	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	33.4	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	9.8	2.1
4	Malignant neoplasms (C00-C97)	40	8.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.6	1.0
6	Influenza and pneumonia (J09-J18)	12	2.6	*
7	Septicemia (A40-A41)	8	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Meningitis (G00,G03)	2	0.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Salmonella infections (A01-A02)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	99	21.5	4.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	301	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	33.9	3.8
2	Malignant neoplasms (C00-C97)	60	19.9	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	8.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.0	*
5	Chronic lower respiratory diseases (J40-J47)	7	2.3	*
6	Septicemia (A40-A41)	5	1.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.7	*
8	Cerebrovascular diseases (I60-I69)	4	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.0	*
9	Influenza and pneumonia (J09-J18)	3	1.0	*
11	Meningitis (G00,G03)	2	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
13	Viral hepatitis (B15-B19)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
...	All other causes (residual)	63	20.9	2.4
Hispanic, male, 10-14 years				
...	All causes	332	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	31.3	4.2
2	Malignant neoplasms (C00-C97)	61	18.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	7.2	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	6.3	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	5.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.7	*
7	Septicemia (A40-A41)	7	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.8	*
9	Cerebrovascular diseases (I60-I69)	4	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
11	Aortic aneurysm and dissection (I71)	2	0.6	*
11	Influenza and pneumonia (J09-J18)	2	0.6	*
11	Peptic ulcer (K25-K28)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
...	All other causes (residual)	64	19.3	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,341	100.0	55.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	549	40.9	22.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	20.2	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	216	16.1	8.9
4	Malignant neoplasms (C00-C97)	81	6.0	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.7	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.3	*
7	Cerebrovascular diseases (I60-I69)	12	0.9	*
8	Influenza and pneumonia (J09-J18)	9	0.7	*
9	Legal intervention (Y35,Y89.0)	7	0.5	*
10	Diabetes mellitus (E10-E14)	6	0.4	*
11	Septicemia (A40-A41)	5	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
12	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.1	*
...	All other causes (residual)	119	8.9	4.9
Hispanic, male, 20-24 years				
...	All causes	2,698	100.0	108.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,199	44.4	48.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	538	19.9	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	15.3	16.5
4	Malignant neoplasms (C00-C97)	127	4.7	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	2.3	2.4
6	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
8	Septicemia (A40-A41)	14	0.5	*
9	Influenza and pneumonia (J09-J18)	12	0.4	*
10	Legal intervention (Y35,Y89.0)	10	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.3	*
11	Diabetes mellitus (E10-E14)	9	0.3	*
13	Cerebrovascular diseases (I60-I69)	7	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.1	*
14	Pneumonitis due to solids and liquids (J69)	4	0.1	*
...	All other causes (residual)	254	9.4	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,592	100.0	108.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,117	43.1	46.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	374	14.4	15.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	325	12.5	13.6
4	Malignant neoplasms (C00-C97)	167	6.4	7.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	4.7	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	31	1.2	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.2	1.3
8	Cerebrovascular diseases (I60-I69)	26	1.0	1.1
9	Diabetes mellitus (E10-E14)	21	0.8	0.9
10	Legal intervention (Y35,Y89.0)	20	0.8	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
12	Influenza and pneumonia (J09-J18)	15	0.6	*
13	Septicemia (A40-A41)	13	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	277	10.7	11.6
Hispanic, male, 30-34 years				
...	All causes	2,676	100.0	115.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	37.8	43.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	300	11.2	12.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	10.7	12.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	7.3	8.4
5	Malignant neoplasms (C00-C97)	194	7.2	8.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	96	3.6	4.1
7	Cerebrovascular diseases (I60-I69)	58	2.2	2.5
8	Diabetes mellitus (E10-E14)	48	1.8	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.8	2.0
10	Influenza and pneumonia (J09-J18)	23	0.9	1.0
11	Legal intervention (Y35,Y89.0)	18	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.6	*
13	Septicemia (A40-A41)	14	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
...	All other causes (residual)	345	12.9	14.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,890	100.0	132.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	31.1	41.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	11.2	14.9
3	Malignant neoplasms (C00-C97)	288	10.0	13.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	8.6	11.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	8.2	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	158	5.5	7.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	73	2.5	3.3
8	Cerebrovascular diseases (I60-I69)	70	2.4	3.2
9	Diabetes mellitus (E10-E14)	56	1.9	2.6
10	Influenza and pneumonia (J09-J18)	28	1.0	1.3
11	Septicemia (A40-A41)	25	0.9	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.6	*
14	Aortic aneurysm and dissection (I71)	14	0.5	*
15	Legal intervention (Y35,Y89.0)	11	0.4	*
...	All other causes (residual)	426	14.7	19.5
Hispanic, male, 40-44 years				
...	All causes	3,495	100.0	175.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	23.8	41.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	13.9	24.3
3	Malignant neoplasms (C00-C97)	424	12.1	21.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	276	7.9	13.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	6.4	11.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	5.1	8.9
7	Diabetes mellitus (E10-E14)	133	3.8	6.7
7	Cerebrovascular diseases (I60-I69)	133	3.8	6.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	91	2.6	4.6
10	Septicemia (A40-A41)	41	1.2	2.1
11	Influenza and pneumonia (J09-J18)	35	1.0	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	1.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.7	1.3
14	Aortic aneurysm and dissection (I71)	23	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	1.2
...	All other causes (residual)	538	15.4	27.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,672	100.0	262.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	828	17.7	46.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	17.3	45.4
3	Malignant neoplasms (C00-C97)	778	16.7	43.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	459	9.8	25.8
5	Diabetes mellitus (E10-E14)	191	4.1	10.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	3.9	10.4
7	Cerebrovascular diseases (I60-I69)	147	3.1	8.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	2.6	6.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	118	2.5	6.6
10	Viral hepatitis (B15-B19)	90	1.9	5.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.6	4.1
12	Septicemia (A40-A41)	61	1.3	3.4
13	Influenza and pneumonia (J09-J18)	46	1.0	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	0.7	1.9
15	Chronic lower respiratory diseases (J40-J47)	28	0.6	1.6
...	All other causes (residual)	706	15.1	39.7
Hispanic, male, 50-54 years				
...	All causes	6,558	100.0	430.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,311	20.0	86.0
2	Malignant neoplasms (C00-C97)	1,310	20.0	85.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	12.1	52.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	689	10.5	45.2
5	Diabetes mellitus (E10-E14)	317	4.8	20.8
6	Cerebrovascular diseases (I60-I69)	232	3.5	15.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	2.9	12.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	128	2.0	8.4
9	Viral hepatitis (B15-B19)	126	1.9	8.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	1.6	7.0
11	Septicemia (A40-A41)	106	1.6	7.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.4	6.1
13	Influenza and pneumonia (J09-J18)	74	1.1	4.9
14	Chronic lower respiratory diseases (J40-J47)	57	0.9	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.7	3.1
...	All other causes (residual)	971	14.8	63.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	7,826	100.0	651.1
1	Malignant neoplasms (C00-C97)	1,964	25.1	163.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	21.5	139.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	665	8.5	55.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	640	8.2	53.2
5	Diabetes mellitus (E10-E14)	424	5.4	35.3
6	Cerebrovascular diseases (I60-I69)	330	4.2	27.5
7	Viral hepatitis (B15-B19)	184	2.4	15.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	1.9	12.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.7	11.2
10	Septicemia (A40-A41)	107	1.4	8.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	102	1.3	8.5
12	Chronic lower respiratory diseases (J40-J47)	97	1.2	8.1
13	Influenza and pneumonia (J09-J18)	92	1.2	7.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.1	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	0.9	6.0
...	All other causes (residual)	1,093	14.0	90.9
Hispanic, male, 60-64 years				
...	All causes	8,540	100.0	976.4
1	Malignant neoplasms (C00-C97)	2,372	27.8	271.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,983	23.2	226.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	607	7.1	69.4
4	Diabetes mellitus (E10-E14)	570	6.7	65.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	5.2	50.5
6	Cerebrovascular diseases (I60-I69)	350	4.1	40.0
7	Viral hepatitis (B15-B19)	168	2.0	19.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	1.9	18.4
9	Chronic lower respiratory diseases (J40-J47)	153	1.8	17.5
10	Septicemia (A40-A41)	120	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	1.3	12.7
12	Influenza and pneumonia (J09-J18)	104	1.2	11.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	1.2	11.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.8	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	0.5	4.5
...	All other causes (residual)	1,192	14.0	136.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	8,776	100.0	1,401.2
1	Malignant neoplasms (C00-C97)	2,599	29.6	415.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,010	22.9	320.9
3	Diabetes mellitus (E10-E14)	593	6.8	94.7
4	Cerebrovascular diseases (I60-I69)	449	5.1	71.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	420	4.8	67.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	3.5	48.5
7	Chronic lower respiratory diseases (J40-J47)	231	2.6	36.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	2.6	36.6
9	Influenza and pneumonia (J09-J18)	153	1.7	24.4
10	Septicemia (A40-A41)	151	1.7	24.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.4	20.3
12	Viral hepatitis (B15-B19)	95	1.1	15.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	0.9	13.1
14	Alzheimer's disease (G30)	54	0.6	8.6
15	Parkinson's disease (G20-G21)	50	0.6	8.0
...	All other causes (residual)	1,229	14.0	196.2
Hispanic, male, 70-74 years				
...	All causes	8,477	100.0	2,097.7
1	Malignant neoplasms (C00-C97)	2,429	28.7	601.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,066	24.4	511.2
3	Diabetes mellitus (E10-E14)	527	6.2	130.4
4	Cerebrovascular diseases (I60-I69)	441	5.2	109.1
5	Chronic lower respiratory diseases (J40-J47)	323	3.8	79.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	285	3.4	70.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.6	53.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	2.5	52.7
9	Septicemia (A40-A41)	174	2.1	43.1
10	Influenza and pneumonia (J09-J18)	153	1.8	37.9
11	Alzheimer's disease (G30)	125	1.5	30.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	1.4	29.0
13	Parkinson's disease (G20-G21)	95	1.1	23.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	12.4
15	Viral hepatitis (B15-B19)	48	0.6	11.9
...	All other causes (residual)	1,213	14.3	300.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	8,926	100.0	3,264.0
1	Malignant neoplasms (C00-C97)	2,438	27.3	891.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,189	24.5	800.5
3	Cerebrovascular diseases (I60-I69)	542	6.1	198.2
4	Diabetes mellitus (E10-E14)	494	5.5	180.6
5	Chronic lower respiratory diseases (J40-J47)	357	4.0	130.5
6	Alzheimer's disease (G30)	228	2.6	83.4
7	Influenza and pneumonia (J09-J18)	222	2.5	81.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	208	2.3	76.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	2.3	73.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	2.2	70.2
11	Parkinson's disease (G20-G21)	164	1.8	60.0
12	Septicemia (A40-A41)	150	1.7	54.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.6	53.0
14	Pneumonitis due to solids and liquids (J69)	57	0.6	20.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.6	19.0
...	All other causes (residual)	1,287	14.4	470.6
Hispanic, male, 80-84 years				
...	All causes	9,557	100.0	5,480.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,498	26.1	1,432.5
2	Malignant neoplasms (C00-C97)	2,114	22.1	1,212.3
3	Cerebrovascular diseases (I60-I69)	649	6.8	372.2
4	Chronic lower respiratory diseases (J40-J47)	473	4.9	271.3
5	Diabetes mellitus (E10-E14)	469	4.9	269.0
6	Alzheimer's disease (G30)	460	4.8	263.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	263	2.8	150.8
8	Influenza and pneumonia (J09-J18)	261	2.7	149.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	2.2	119.3
10	Parkinson's disease (G20-G21)	206	2.2	118.1
11	Septicemia (A40-A41)	139	1.5	79.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	138	1.4	79.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	113	1.2	64.8
14	Pneumonitis due to solids and liquids (J69)	90	0.9	51.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.7	39.6
...	All other causes (residual)	1,407	14.7	806.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	8,574	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,367	27.6	---
2	Malignant neoplasms (C00-C97)	1,539	17.9	---
3	Cerebrovascular diseases (I60-I69)	585	6.8	---
4	Alzheimer's disease (G30)	559	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	426	5.0	---
6	Diabetes mellitus (E10-E14)	344	4.0	---
7	Influenza and pneumonia (J09-J18)	248	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	2.7	---
9	Parkinson's disease (G20-G21)	202	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.3	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	1.5	---
12	Septicemia (A40-A41)	123	1.4	---
13	Pneumonitis due to solids and liquids (J69)	103	1.2	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	---
...	All other causes (residual)	1,405	16.4	---
Hispanic, male, 90-94 years				
...	All causes	5,078	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,443	28.4	---
2	Malignant neoplasms (C00-C97)	704	13.9	---
3	Alzheimer's disease (G30)	429	8.4	---
4	Cerebrovascular diseases (I60-I69)	352	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	272	5.4	---
6	Diabetes mellitus (E10-E14)	178	3.5	---
7	Influenza and pneumonia (J09-J18)	151	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.9	---
11	Parkinson's disease (G20-G21)	89	1.8	---
12	Septicemia (A40-A41)	69	1.4	---
13	Pneumonitis due to solids and liquids (J69)	65	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	---
15	Nutritional deficiencies (E40-E64)	19	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.4	---
...	All other causes (residual)	934	18.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,474	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	461	31.3	---
2	Malignant neoplasms (C00-C97)	137	9.3	---
3	Alzheimer's disease (G30)	118	8.0	---
4	Cerebrovascular diseases (I60-I69)	116	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	93	6.3	---
6	Influenza and pneumonia (J09-J18)	58	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	2.7	---
8	Diabetes mellitus (E10-E14)	35	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	2.4	---
11	Pneumonitis due to solids and liquids (J69)	27	1.8	---
12	Septicemia (A40-A41)	13	0.9	---
12	Parkinson's disease (G20-G21)	13	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	---
15	Atherosclerosis (I70)	7	0.5	---
...	All other causes (residual)	275	18.7	---
Hispanic, male, 100 years and over				
...	All causes	302	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	37.7	---
2	Alzheimer's disease (G30)	26	8.6	---
3	Cerebrovascular diseases (I60-I69)	22	7.3	---
4	Influenza and pneumonia (J09-J18)	14	4.6	---
5	Malignant neoplasms (C00-C97)	13	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	13	4.3	---
7	Diabetes mellitus (E10-E14)	9	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.3	---
11	Septicemia (A40-A41)	5	1.7	---
12	Pneumonitis due to solids and liquids (J69)	4	1.3	---
13	Atherosclerosis (I70)	3	1.0	---
14	Parkinson's disease (G20-G21)	2	0.7	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	---
15	Hyperplasia of prostate (N40)	1	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	---
...	All other causes (residual)	49	16.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	81,287	100.0	290.4
1	Malignant neoplasms (C00-C97)	17,957	22.1	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,176	19.9	57.8
3	Cerebrovascular diseases (I60-I69)	5,251	6.5	18.8
4	Alzheimer's disease (G30)	4,409	5.4	15.8
5	Diabetes mellitus (E10-E14)	3,852	4.7	13.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,739	4.6	13.4
7	Chronic lower respiratory diseases (J40-J47)	2,553	3.1	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,909	2.3	6.8
9	Influenza and pneumonia (J09-J18)	1,762	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,744	2.1	6.2
11	Septicemia (A40-A41)	1,322	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,245	1.5	4.4
13	Certain conditions originating in the perinatal period (P00-P96)	1,086	1.3	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	837	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	0.9	2.6
...	All other causes (residual)	16,729	20.6	59.8
Hispanic, female, 1-4 years				
...	All causes	354	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	29.4	5.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.1	2.8
3	Malignant neoplasms (C00-C97)	35	9.9	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.8	1.2
5	Septicemia (A40-A41)	9	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.3	*
7	Influenza and pneumonia (J09-J18)	6	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.4	*
9	Cerebrovascular diseases (I60-I69)	4	1.1	*
10	Anemias (D50-D64)	3	0.8	*
11	Meningitis (G00,G03)	2	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	86	24.3	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	194	100.0	7.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	29.4	2.2
2	Malignant neoplasms (C00-C97)	42	21.6	1.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	6.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	6.2	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	3.1	*
6	Cerebrovascular diseases (I60-I69)	5	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	5	2.6	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.1	*
9	Influenza and pneumonia (J09-J18)	3	1.5	*
9	Pneumonitis due to solids and liquids (J69)	3	1.5	*
11	Septicemia (A40-A41)	2	1.0	*
11	Diabetes mellitus (E10-E14)	2	1.0	*
11	Meningitis (G00,G03)	2	1.0	*
14	Meningococcal infection (A39)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (residual)	34	17.5	1.3
Hispanic, female, 10-14 years				
...	All causes	236	100.0	9.7
1	Malignant neoplasms (C00-C97)	48	20.3	2.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	19.1	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	10.6	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.0	*
7	Influenza and pneumonia (J09-J18)	6	2.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.1	*
10	Cerebrovascular diseases (I60-I69)	3	1.3	*
11	Septicemia (A40-A41)	2	0.8	*
11	Anemias (D50-D64)	2	0.8	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	60	25.4	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	527	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	36.6	8.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	15.7	3.6
3	Malignant neoplasms (C00-C97)	56	10.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.4	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.7	*
7	Cerebrovascular diseases (I60-I69)	6	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
9	Influenza and pneumonia (J09-J18)	5	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
12	Septicemia (A40-A41)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
14	Meningitis (G00,G03)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	75	14.2	3.3
Hispanic, female, 20-24 years				
...	All causes	787	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	40.7	13.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	12.1	4.1
3	Malignant neoplasms (C00-C97)	68	8.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	8.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.2	1.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	21	2.7	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.4	*
8	Influenza and pneumonia (J09-J18)	10	1.3	*
9	Septicemia (A40-A41)	7	0.9	*
9	Cerebrovascular diseases (I60-I69)	7	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	127	16.1	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	804	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	30.5	11.4
2	Malignant neoplasms (C00-C97)	120	14.9	5.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	8.6	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.8	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.5	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	28	3.5	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	2.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.9	*
9	Cerebrovascular diseases (I60-I69)	13	1.6	*
10	Influenza and pneumonia (J09-J18)	11	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.0	*
13	Septicemia (A40-A41)	7	0.9	*
14	Diabetes mellitus (E10-E14)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
...	All other causes (residual)	162	20.1	7.5
Hispanic, female, 30-34 years				
...	All causes	1,110	100.0	52.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	25.1	13.1
2	Malignant neoplasms (C00-C97)	233	21.0	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	7.7	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	7.5	3.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	2.9
6	Diabetes mellitus (E10-E14)	31	2.8	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.8	1.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.4	1.3
9	Cerebrovascular diseases (I60-I69)	24	2.2	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.3	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.3	*
12	Septicemia (A40-A41)	12	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Influenza and pneumonia (J09-J18)	7	0.6	*
...	All other causes (residual)	188	16.9	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,312	100.0	63.8
1	Malignant neoplasms (C00-C97)	383	29.2	18.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	17.9	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	6.8	4.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	5.1	3.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	60	4.6	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	3.9	2.5
7	Cerebrovascular diseases (I60-I69)	48	3.7	2.3
8	Diabetes mellitus (E10-E14)	36	2.7	1.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	34	2.6	1.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.6	1.0
11	Septicemia (A40-A41)	15	1.1	*
12	Influenza and pneumonia (J09-J18)	14	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.0	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (residual)	231	17.6	11.2
Hispanic, female, 40-44 years				
...	All causes	1,793	100.0	92.4
1	Malignant neoplasms (C00-C97)	623	34.7	32.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	12.6	11.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	10.2	9.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	100	5.6	5.2
5	Cerebrovascular diseases (I60-I69)	70	3.9	3.6
6	Diabetes mellitus (E10-E14)	49	2.7	2.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	2.6	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	2.3	2.2
9	Influenza and pneumonia (J09-J18)	25	1.4	1.3
10	Septicemia (A40-A41)	24	1.3	1.2
11	Chronic lower respiratory diseases (J40-J47)	21	1.2	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.8	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.7	*
...	All other causes (residual)	322	18.0	16.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,487	100.0	144.7
1	Malignant neoplasms (C00-C97)	904	36.3	52.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	257	10.3	15.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	9.4	13.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	137	5.5	8.0
5	Cerebrovascular diseases (I60-I69)	115	4.6	6.7
6	Diabetes mellitus (E10-E14)	91	3.7	5.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	2.9	4.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.0	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.9	2.7
10	Septicemia (A40-A41)	46	1.8	2.7
11	Influenza and pneumonia (J09-J18)	33	1.3	1.9
12	Chronic lower respiratory diseases (J40-J47)	27	1.1	1.6
13	Viral hepatitis (B15-B19)	23	0.9	1.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.9	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.8	*
...	All other causes (residual)	412	16.6	24.0
Hispanic, female, 50-54 years				
...	All causes	3,472	100.0	228.4
1	Malignant neoplasms (C00-C97)	1,294	37.3	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	448	12.9	29.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	7.3	16.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	192	5.5	12.6
5	Diabetes mellitus (E10-E14)	173	5.0	11.4
6	Cerebrovascular diseases (I60-I69)	171	4.9	11.2
7	Septicemia (A40-A41)	56	1.6	3.7
7	Influenza and pneumonia (J09-J18)	56	1.6	3.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	3.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	1.6	3.7
11	Chronic lower respiratory diseases (J40-J47)	55	1.6	3.6
12	Viral hepatitis (B15-B19)	41	1.2	2.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.9	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.8	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.6	1.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.6	1.4
...	All other causes (residual)	518	14.9	34.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	4,302	100.0	344.2
1	Malignant neoplasms (C00-C97)	1,638	38.1	131.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	607	14.1	48.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	248	5.8	19.8
4	Diabetes mellitus (E10-E14)	235	5.5	18.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	4.8	16.4
6	Cerebrovascular diseases (I60-I69)	183	4.3	14.6
7	Chronic lower respiratory diseases (J40-J47)	106	2.5	8.5
8	Septicemia (A40-A41)	81	1.9	6.5
9	Viral hepatitis (B15-B19)	78	1.8	6.2
10	Influenza and pneumonia (J09-J18)	74	1.7	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	1.5	5.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	1.0	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	0.9	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	*
...	All other causes (residual)	658	15.3	52.6
Hispanic, female, 60-64 years				
...	All causes	5,319	100.0	551.9
1	Malignant neoplasms (C00-C97)	1,934	36.4	200.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	826	15.5	85.7
3	Diabetes mellitus (E10-E14)	387	7.3	40.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	247	4.6	25.6
5	Cerebrovascular diseases (I60-I69)	244	4.6	25.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	2.8	15.6
7	Chronic lower respiratory diseases (J40-J47)	136	2.6	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.5	13.7
9	Septicemia (A40-A41)	113	2.1	11.7
10	Influenza and pneumonia (J09-J18)	82	1.5	8.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.2	6.8
12	Viral hepatitis (B15-B19)	63	1.2	6.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.7	3.6
14	Alzheimer's disease (G30)	31	0.6	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	3.0
...	All other causes (residual)	844	15.9	87.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	6,144	100.0	840.2
1	Malignant neoplasms (C00-C97)	2,012	32.7	275.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,122	18.3	153.4
3	Diabetes mellitus (E10-E14)	461	7.5	63.0
4	Cerebrovascular diseases (I60-I69)	301	4.9	41.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	241	3.9	33.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	3.0	25.0
7	Chronic lower respiratory diseases (J40-J47)	150	2.4	20.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	2.2	18.7
9	Influenza and pneumonia (J09-J18)	119	1.9	16.3
10	Septicemia (A40-A41)	117	1.9	16.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.4	12.2
12	Alzheimer's disease (G30)	63	1.0	8.6
13	Viral hepatitis (B15-B19)	49	0.8	6.7
14	Parkinson's disease (G20-G21)	30	0.5	4.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	26	0.4	3.6
...	All other causes (residual)	1,044	17.0	142.8
Hispanic, female, 70-74 years				
...	All causes	6,878	100.0	1,347.2
1	Malignant neoplasms (C00-C97)	2,093	30.4	410.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,319	19.2	258.4
3	Diabetes mellitus (E10-E14)	475	6.9	93.0
4	Cerebrovascular diseases (I60-I69)	370	5.4	72.5
5	Chronic lower respiratory diseases (J40-J47)	253	3.7	49.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	207	3.0	40.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	3.0	39.8
8	Alzheimer's disease (G30)	159	2.3	31.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	2.1	28.6
10	Influenza and pneumonia (J09-J18)	131	1.9	25.7
11	Septicemia (A40-A41)	127	1.8	24.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.4	19.0
13	Parkinson's disease (G20-G21)	67	1.0	13.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.7	10.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	34	0.5	6.7
...	All other causes (residual)	1,146	16.7	224.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	8,316	100.0	2,234.0
1	Malignant neoplasms (C00-C97)	2,021	24.3	542.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,761	21.2	473.1
3	Cerebrovascular diseases (I60-I69)	584	7.0	156.9
4	Diabetes mellitus (E10-E14)	538	6.5	144.5
5	Alzheimer's disease (G30)	380	4.6	102.1
6	Chronic lower respiratory diseases (J40-J47)	337	4.1	90.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	3.0	66.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	198	2.4	53.2
9	Influenza and pneumonia (J09-J18)	193	2.3	51.8
10	Septicemia (A40-A41)	168	2.0	45.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	1.8	40.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	1.4	31.7
13	Parkinson's disease (G20-G21)	101	1.2	27.1
14	Pneumonitis due to solids and liquids (J69)	56	0.7	15.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.6	14.5
...	All other causes (residual)	1,409	16.9	378.5
Hispanic, female, 80-84 years				
...	All causes	10,290	100.0	3,939.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,387	23.2	913.9
2	Malignant neoplasms (C00-C97)	1,917	18.6	734.0
3	Cerebrovascular diseases (I60-I69)	873	8.5	334.2
4	Alzheimer's disease (G30)	787	7.6	301.3
5	Diabetes mellitus (E10-E14)	552	5.4	211.3
6	Chronic lower respiratory diseases (J40-J47)	459	4.5	175.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	2.6	102.2
8	Influenza and pneumonia (J09-J18)	263	2.6	100.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	204	2.0	78.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	2.0	78.1
11	Parkinson's disease (G20-G21)	162	1.6	62.0
12	Septicemia (A40-A41)	161	1.6	61.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.3	51.7
14	Pneumonitis due to solids and liquids (J69)	77	0.7	29.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.6	23.0
...	All other causes (residual)	1,782	17.3	682.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	11,466	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,098	27.0	---
2	Malignant neoplasms (C00-C97)	1,511	13.2	---
3	Alzheimer's disease (G30)	1,169	10.2	---
4	Cerebrovascular diseases (I60-I69)	986	8.6	---
5	Chronic lower respiratory diseases (J40-J47)	487	4.2	---
6	Diabetes mellitus (E10-E14)	475	4.1	---
7	Influenza and pneumonia (J09-J18)	294	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.8	---
11	Septicemia (A40-A41)	192	1.7	---
12	Parkinson's disease (G20-G21)	150	1.3	---
13	Pneumonitis due to solids and liquids (J69)	85	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.6	---
...	All other causes (residual)	2,182	19.0	---
Hispanic, female, 90-94 years				
...	All causes	8,729	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,456	28.1	---
2	Alzheimer's disease (G30)	1,161	13.3	---
3	Cerebrovascular diseases (I60-I69)	827	9.5	---
4	Malignant neoplasms (C00-C97)	787	9.0	---
5	Chronic lower respiratory diseases (J40-J47)	311	3.6	---
6	Influenza and pneumonia (J09-J18)	262	3.0	---
7	Diabetes mellitus (E10-E14)	237	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	147	1.7	---
11	Parkinson's disease (G20-G21)	119	1.4	---
12	Septicemia (A40-A41)	112	1.3	---
13	Pneumonitis due to solids and liquids (J69)	85	1.0	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	35	0.4	---
15	Nutritional deficiencies (E40-E64)	34	0.4	---
...	All other causes (residual)	1,790	20.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	3,600	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,062	29.5	---
2	Alzheimer's disease (G30)	504	14.0	---
3	Cerebrovascular diseases (I60-I69)	346	9.6	---
4	Malignant neoplasms (C00-C97)	197	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	143	4.0	---
6	Influenza and pneumonia (J09-J18)	115	3.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	2.7	---
8	Diabetes mellitus (E10-E14)	81	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	---
11	Septicemia (A40-A41)	39	1.1	---
12	Pneumonitis due to solids and liquids (J69)	30	0.8	---
13	Parkinson's disease (G20-G21)	29	0.8	---
14	Nutritional deficiencies (E40-E64)	18	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	---
...	All other causes (residual)	792	22.0	---
Hispanic, female, 100 years and over				
...	All causes	980	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	34.0	---
2	Alzheimer's disease (G30)	135	13.8	---
3	Cerebrovascular diseases (I60-I69)	63	6.4	---
4	Influenza and pneumonia (J09-J18)	43	4.4	---
5	Malignant neoplasms (C00-C97)	36	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	25	2.6	---
8	Diabetes mellitus (E10-E14)	16	1.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.3	---
10	Pneumonitis due to solids and liquids (J69)	11	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.0	---
12	Nutritional deficiencies (E40-E64)	9	0.9	---
13	Septicemia (A40-A41)	7	0.7	---
14	Anemias (D50-D64)	6	0.6	---
15	Parkinson's disease (G20-G21)	5	0.5	---
15	Atherosclerosis (I70)	5	0.5	---
...	All other causes (residual)	232	23.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,123,631	100.0	1,055.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,172	23.7	250.0
2	Malignant neoplasms (C00-C97)	467,208	22.0	232.2
3	Chronic lower respiratory diseases (J40-J47)	136,228	6.4	67.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111,827	5.3	55.6
5	Cerebrovascular diseases (I60-I69)	106,830	5.0	53.1
6	Alzheimer's disease (G30)	93,325	4.4	46.4
7	Diabetes mellitus (E10-E14)	53,590	2.5	26.6
8	Influenza and pneumonia (J09-J18)	45,242	2.1	22.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,465	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,481	1.7	17.6
11	Septicemia (A40-A41)	30,240	1.4	15.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,343	1.4	14.6
13	Parkinson's disease (G20-G21)	24,339	1.1	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,628	1.1	11.2
15	Pneumonitis due to solids and liquids (J69)	16,430	0.8	8.2
...	All other causes (residual)	411,283	19.4	204.4
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,895	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	631	33.3	7.6
2	Malignant neoplasms (C00-C97)	206	10.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	10.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	7.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.9	0.7
6	Influenza and pneumonia (J09-J18)	37	2.0	0.4
7	Septicemia (A40-A41)	20	1.1	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Cerebrovascular diseases (I60-I69)	17	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	10	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Anemias (D50-D64)	7	0.4	*
...	All other causes (residual)	511	27.0	6.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,192	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	31.4	3.5
2	Malignant neoplasms (C00-C97)	241	20.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	5.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.6	0.3
6	Influenza and pneumonia (J09-J18)	25	2.1	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.8	0.2
8	Cerebrovascular diseases (I60-I69)	20	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	17	1.4	*
10	Septicemia (A40-A41)	12	1.0	*
11	Anemias (D50-D64)	6	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Meningitis (G00,G03)	2	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
14	Peptic ulcer (K25-K28)	2	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	259	21.7	2.4
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,643	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	424	25.8	3.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	289	17.6	2.6
3	Malignant neoplasms (C00-C97)	235	14.3	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	5.6	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	4.3	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	2.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.2
8	Influenza and pneumonia (J09-J18)	22	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	16	1.0	*
12	Diabetes mellitus (E10-E14)	10	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.2	*
...	All other causes (residual)	365	22.2	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,456	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,449	44.9	20.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,411	25.9	12.0
3	Malignant neoplasms (C00-C97)	307	5.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	4.3	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	2.0	0.9
7	Influenza and pneumonia (J09-J18)	44	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	29	0.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	0.2
10	Diabetes mellitus (E10-E14)	22	0.4	0.2
11	Septicemia (A40-A41)	18	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	18	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	Legal intervention (Y35,Y89.0)	9	0.2	*
...	All other causes (residual)	602	11.0	5.1
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,080	100.0	87.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,630	50.8	44.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,266	20.5	17.8
3	Malignant neoplasms (C00-C97)	474	4.3	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	446	4.0	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	3.1	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	1.0	0.9
7	Diabetes mellitus (E10-E14)	82	0.7	0.6
8	Influenza and pneumonia (J09-J18)	63	0.6	0.5
9	Cerebrovascular diseases (I60-I69)	54	0.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	51	0.5	0.4
12	Septicemia (A40-A41)	43	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	25	0.2	0.2
15	Legal intervention (Y35,Y89.0)	23	0.2	0.2
...	All other causes (residual)	1,382	12.5	10.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	14,349	100.0	111.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,014	48.9	54.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,495	17.4	19.4
3	Malignant neoplasms (C00-C97)	757	5.3	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	689	4.8	5.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	519	3.6	4.0
6	Diabetes mellitus (E10-E14)	152	1.1	1.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	131	0.9	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	0.9	1.0
9	Influenza and pneumonia (J09-J18)	103	0.7	0.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	103	0.7	0.8
11	Cerebrovascular diseases (I60-I69)	84	0.6	0.7
12	Septicemia (A40-A41)	76	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	47	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.2	0.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.2	0.3
...	All other causes (residual)	1,988	13.9	15.5
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	17,143	100.0	136.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,129	41.6	56.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,682	15.6	21.4
3	Malignant neoplasms (C00-C97)	1,374	8.0	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,239	7.2	9.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	516	3.0	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	392	2.3	3.1
7	Diabetes mellitus (E10-E14)	256	1.5	2.0
8	Cerebrovascular diseases (I60-I69)	172	1.0	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	0.9	1.2
10	Influenza and pneumonia (J09-J18)	123	0.7	1.0
11	Septicemia (A40-A41)	122	0.7	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	99	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	80	0.5	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	0.5
...	All other causes (residual)	2,680	15.6	21.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	19,392	100.0	165.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,632	34.2	56.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,654	13.7	22.6
3	Malignant neoplasms (C00-C97)	2,196	11.3	18.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	10.5	17.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	638	3.3	5.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	429	2.2	3.6
7	Diabetes mellitus (E10-E14)	378	1.9	3.2
8	Cerebrovascular diseases (I60-I69)	266	1.4	2.3
9	Influenza and pneumonia (J09-J18)	173	0.9	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.8	1.4
11	Septicemia (A40-A41)	159	0.8	1.4
12	Chronic lower respiratory diseases (J40-J47)	129	0.7	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	0.6	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	81	0.4	0.7
...	All other causes (residual)	3,265	16.8	27.8
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,245	100.0	218.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,259	23.8	52.2
2	Malignant neoplasms (C00-C97)	4,340	16.5	36.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,950	15.1	32.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,969	11.3	24.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,194	4.5	10.0
6	Diabetes mellitus (E10-E14)	690	2.6	5.8
7	Cerebrovascular diseases (I60-I69)	530	2.0	4.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	1.7	3.6
9	Chronic lower respiratory diseases (J40-J47)	315	1.2	2.6
10	Septicemia (A40-A41)	304	1.2	2.5
11	Influenza and pneumonia (J09-J18)	251	1.0	2.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	177	0.7	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	0.7	1.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	140	0.5	1.2
15	Viral hepatitis (B15-B19)	117	0.4	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	0.4	1.0
...	All other causes (residual)	4,282	16.3	35.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	42,315	100.0	320.2
1	Malignant neoplasms (C00-C97)	9,062	21.4	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,678	18.1	58.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,184	17.0	54.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,388	8.0	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,181	5.2	16.5
6	Diabetes mellitus (E10-E14)	1,301	3.1	9.8
7	Cerebrovascular diseases (I60-I69)	960	2.3	7.3
8	Chronic lower respiratory diseases (J40-J47)	827	2.0	6.3
9	Septicemia (A40-A41)	547	1.3	4.1
10	Influenza and pneumonia (J09-J18)	494	1.2	3.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	444	1.0	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	0.8	2.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	280	0.7	2.1
14	Viral hepatitis (B15-B19)	270	0.6	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.6	1.8
...	All other causes (residual)	7,110	16.8	53.8
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	76,263	100.0	504.2
1	Malignant neoplasms (C00-C97)	20,504	26.9	135.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,969	19.6	99.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,693	11.4	57.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,158	5.5	27.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,033	5.3	26.7
6	Chronic lower respiratory diseases (J40-J47)	2,636	3.5	17.4
7	Diabetes mellitus (E10-E14)	2,334	3.1	15.4
8	Cerebrovascular diseases (I60-I69)	1,915	2.5	12.7
9	Septicemia (A40-A41)	1,003	1.3	6.6
10	Influenza and pneumonia (J09-J18)	835	1.1	5.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	724	0.9	4.8
12	Viral hepatitis (B15-B19)	712	0.9	4.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	0.7	3.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	480	0.6	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	423	0.6	2.8
...	All other causes (residual)	12,294	16.1	81.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	113,442	100.0	733.0
1	Malignant neoplasms (C00-C97)	36,430	32.1	235.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,105	20.4	149.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,239	7.3	53.2
4	Chronic lower respiratory diseases (J40-J47)	5,597	4.9	36.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,961	4.4	32.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,003	3.5	25.9
7	Diabetes mellitus (E10-E14)	3,755	3.3	24.3
8	Cerebrovascular diseases (I60-I69)	3,040	2.7	19.6
9	Septicemia (A40-A41)	1,610	1.4	10.4
10	Influenza and pneumonia (J09-J18)	1,323	1.2	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,174	1.0	7.6
12	Viral hepatitis (B15-B19)	1,120	1.0	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	865	0.8	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	504	0.4	3.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	449	0.4	2.9
...	All other causes (residual)	17,267	15.2	111.6
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	144,070	100.0	1,025.1
1	Malignant neoplasms (C00-C97)	49,264	34.2	350.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,801	21.4	219.2
3	Chronic lower respiratory diseases (J40-J47)	9,060	6.3	64.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,336	4.4	45.1
5	Diabetes mellitus (E10-E14)	5,072	3.5	36.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,767	3.3	33.9
7	Cerebrovascular diseases (I60-I69)	4,308	3.0	30.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,926	2.0	20.8
9	Septicemia (A40-A41)	2,262	1.6	16.1
10	Influenza and pneumonia (J09-J18)	1,879	1.3	13.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,871	1.3	13.3
12	Viral hepatitis (B15-B19)	1,237	0.9	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,210	0.8	8.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	733	0.5	5.2
15	Pneumonitis due to solids and liquids (J69)	647	0.4	4.6
...	All other causes (residual)	21,697	15.1	154.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	179,608	100.0	1,464.5
1	Malignant neoplasms (C00-C97)	63,108	35.1	514.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,997	21.2	309.8
3	Chronic lower respiratory diseases (J40-J47)	14,033	7.8	114.4
4	Diabetes mellitus (E10-E14)	6,470	3.6	52.8
5	Cerebrovascular diseases (I60-I69)	6,040	3.4	49.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,425	3.0	44.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,686	2.1	30.1
8	Septicemia (A40-A41)	2,959	1.6	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,790	1.6	22.7
10	Influenza and pneumonia (J09-J18)	2,631	1.5	21.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,209	1.2	18.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,533	0.9	12.5
13	Alzheimer's disease (G30)	1,384	0.8	11.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,061	0.6	8.7
15	Parkinson's disease (G20-G21)	1,004	0.6	8.2
...	All other causes (residual)	27,278	15.2	222.4
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	204,486	100.0	2,286.1
1	Malignant neoplasms (C00-C97)	66,699	32.6	745.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,294	21.2	484.0
3	Chronic lower respiratory diseases (J40-J47)	19,219	9.4	214.9
4	Cerebrovascular diseases (I60-I69)	8,526	4.2	95.3
5	Diabetes mellitus (E10-E14)	6,643	3.2	74.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,886	2.4	54.6
7	Alzheimer's disease (G30)	3,621	1.8	40.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,538	1.7	39.6
9	Influenza and pneumonia (J09-J18)	3,481	1.7	38.9
10	Septicemia (A40-A41)	3,352	1.6	37.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,641	1.3	29.5
12	Parkinson's disease (G20-G21)	2,199	1.1	24.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,648	0.8	18.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,618	0.8	18.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,499	0.7	16.8
...	All other causes (residual)	31,622	15.5	353.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	232,080	100.0	3,655.3
1	Malignant neoplasms (C00-C97)	64,236	27.7	1,011.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,751	21.9	799.3
3	Chronic lower respiratory diseases (J40-J47)	21,692	9.3	341.7
4	Cerebrovascular diseases (I60-I69)	11,827	5.1	186.3
5	Alzheimer's disease (G30)	8,134	3.5	128.1
6	Diabetes mellitus (E10-E14)	6,590	2.8	103.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,799	2.5	91.3
8	Influenza and pneumonia (J09-J18)	4,527	2.0	71.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,382	1.9	69.0
10	Parkinson's disease (G20-G21)	4,018	1.7	63.3
11	Septicemia (A40-A41)	3,812	1.6	60.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,164	0.9	34.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,979	0.9	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,817	0.8	28.6
15	Pneumonitis due to solids and liquids (J69)	1,722	0.7	27.1
...	All other causes (residual)	38,630	16.6	608.4
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	285,606	100.0	6,176.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,159	24.2	1,495.7
2	Malignant neoplasms (C00-C97)	61,031	21.4	1,319.9
3	Chronic lower respiratory diseases (J40-J47)	22,781	8.0	492.7
4	Cerebrovascular diseases (I60-I69)	17,335	6.1	374.9
5	Alzheimer's disease (G30)	16,388	5.7	354.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,434	2.6	160.8
7	Diabetes mellitus (E10-E14)	7,174	2.5	155.2
8	Influenza and pneumonia (J09-J18)	6,638	2.3	143.6
9	Parkinson's disease (G20-G21)	5,966	2.1	129.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,787	2.0	125.2
11	Septicemia (A40-A41)	4,264	1.5	92.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,091	1.1	66.8
13	Pneumonitis due to solids and liquids (J69)	2,519	0.9	54.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,270	0.8	49.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,461	0.5	31.6
...	All other causes (residual)	52,308	18.3	1,131.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	331,643	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,934	27.1	---
2	Malignant neoplasms (C00-C97)	50,636	15.3	---
3	Alzheimer's disease (G30)	25,114	7.6	---
4	Cerebrovascular diseases (I60-I69)	22,643	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	21,486	6.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,371	2.8	---
7	Influenza and pneumonia (J09-J18)	9,040	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,768	2.0	---
9	Diabetes mellitus (E10-E14)	6,767	2.0	---
10	Parkinson's disease (G20-G21)	6,181	1.9	---
11	Septicemia (A40-A41)	4,860	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,239	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,579	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,396	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,256	0.4	---
...	All other causes (residual)	67,373	20.3	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	271,961	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,632	30.4	---
2	Malignant neoplasms (C00-C97)	27,914	10.3	---
3	Alzheimer's disease (G30)	24,731	9.1	---
4	Cerebrovascular diseases (I60-I69)	19,330	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	13,345	4.9	---
6	Influenza and pneumonia (J09-J18)	8,529	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,631	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,343	2.0	---
9	Diabetes mellitus (E10-E14)	4,380	1.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,267	1.6	---
11	Parkinson's disease (G20-G21)	3,519	1.3	---
12	Septicemia (A40-A41)	3,386	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,259	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,688	0.6	---
15	Atherosclerosis (I70)	1,140	0.4	---
...	All other causes (residual)	60,867	22.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	110,707	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,193	32.7	---
2	Alzheimer's disease (G30)	10,996	9.9	---
3	Cerebrovascular diseases (I60-I69)	8,109	7.3	---
4	Malignant neoplasms (C00-C97)	7,253	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	4,183	3.8	---
6	Influenza and pneumonia (J09-J18)	3,989	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,092	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,072	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,925	1.7	---
10	Pneumonitis due to solids and liquids (J69)	1,344	1.2	---
11	Diabetes mellitus (E10-E14)	1,310	1.2	---
12	Septicemia (A40-A41)	1,153	1.0	---
13	Parkinson's disease (G20-G21)	818	0.7	---
14	Atherosclerosis (I70)	561	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	510	0.5	---
...	All other causes (residual)	27,199	24.6	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	22,721	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,945	35.0	---
2	Alzheimer's disease (G30)	2,125	9.4	---
3	Cerebrovascular diseases (I60-I69)	1,566	6.9	---
4	Influenza and pneumonia (J09-J18)	957	4.2	---
5	Malignant neoplasms (C00-C97)	906	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	658	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	554	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	519	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	1.6	---
10	Pneumonitis due to solids and liquids (J69)	266	1.2	---
11	Diabetes mellitus (E10-E14)	200	0.9	---
12	Septicemia (A40-A41)	190	0.8	---
13	Atherosclerosis (I70)	181	0.8	---
14	Nutritional deficiencies (E40-E64)	126	0.6	---
15	Parkinson's disease (G20-G21)	73	0.3	---
...	All other causes (residual)	6,091	26.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,063,705	100.0	1,072.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,784	24.9	267.0
2	Malignant neoplasms (C00-C97)	247,645	23.3	249.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,715	6.5	69.3
4	Chronic lower respiratory diseases (J40-J47)	62,857	5.9	63.4
5	Cerebrovascular diseases (I60-I69)	43,100	4.1	43.5
6	Diabetes mellitus (E10-E14)	29,813	2.8	30.1
7	Alzheimer's disease (G30)	28,631	2.7	28.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,045	2.6	28.3
9	Influenza and pneumonia (J09-J18)	20,995	2.0	21.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,546	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,351	1.7	18.5
12	Parkinson's disease (G20-G21)	14,778	1.4	14.9
13	Septicemia (A40-A41)	14,309	1.3	14.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,526	0.9	9.6
15	Pneumonitis due to solids and liquids (J69)	9,085	0.9	9.2
...	All other causes (residual)	184,525	17.3	186.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,083	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	37.0	9.4
2	Malignant neoplasms (C00-C97)	116	10.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
6	Influenza and pneumonia (J09-J18)	17	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
8	Cerebrovascular diseases (I60-I69)	10	0.9	*
9	Septicemia (A40-A41)	9	0.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
13	Anemias (D50-D64)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	267	24.7	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	669	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	31.7	3.8
2	Malignant neoplasms (C00-C97)	129	19.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	8.1	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.0	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.7	*
6	Influenza and pneumonia (J09-J18)	13	1.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
8	Cerebrovascular diseases (I60-I69)	9	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
10	Septicemia (A40-A41)	5	0.7	*
11	Anemias (D50-D64)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	154	23.0	2.8
Non-Hispanic, white, male, 10-14 years				
...	All causes	965	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	29.1	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	19.2	3.2
3	Malignant neoplasms (C00-C97)	118	12.2	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	5.4	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	12	1.2	*
8	Influenza and pneumonia (J09-J18)	12	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Septicemia (A40-A41)	7	0.7	*
12	Diabetes mellitus (E10-E14)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (residual)	195	20.2	3.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,692	100.0	61.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,664	45.1	27.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,063	28.8	17.6
3	Malignant neoplasms (C00-C97)	179	4.8	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	4.6	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.7	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.6	1.0
7	Influenza and pneumonia (J09-J18)	25	0.7	0.4
8	Septicemia (A40-A41)	12	0.3	*
8	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
11	Diabetes mellitus (E10-E14)	10	0.3	*
11	Cerebrovascular diseases (I60-I69)	10	0.3	*
13	Legal intervention (Y35,Y89.0)	8	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
...	All other causes (residual)	359	9.7	5.9
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,061	100.0	123.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,141	51.4	63.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,885	23.4	28.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	3.9	4.8
4	Malignant neoplasms (C00-C97)	289	3.6	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	2.9	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.8	1.0
7	Diabetes mellitus (E10-E14)	48	0.6	0.7
8	Influenza and pneumonia (J09-J18)	37	0.5	0.6
9	Chronic lower respiratory diseases (J40-J47)	34	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	33	0.4	0.5
11	Legal intervention (Y35,Y89.0)	23	0.3	0.4
12	Septicemia (A40-A41)	20	0.2	0.3
13	Pneumonitis due to solids and liquids (J69)	17	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.2	*
...	All other causes (residual)	891	11.1	13.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	10,097	100.0	154.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,223	51.7	80.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,016	20.0	30.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	435	4.3	6.7
4	Malignant neoplasms (C00-C97)	401	4.0	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	3.5	5.5
6	Diabetes mellitus (E10-E14)	87	0.9	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	0.8	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.7	1.1
9	Influenza and pneumonia (J09-J18)	53	0.5	0.8
10	Cerebrovascular diseases (I60-I69)	38	0.4	0.6
11	Septicemia (A40-A41)	36	0.4	0.6
12	Legal intervention (Y35,Y89.0)	32	0.3	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.3	0.4
14	Aortic aneurysm and dissection (I71)	23	0.2	0.4
14	Chronic lower respiratory diseases (J40-J47)	23	0.2	0.4
...	All other causes (residual)	1,197	11.9	18.3
Non-Hispanic, white, male, 30-34 years				
...	All causes	11,499	100.0	181.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,083	44.2	80.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,091	18.2	33.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	821	7.1	13.0
4	Malignant neoplasms (C00-C97)	676	5.9	10.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	354	3.1	5.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	240	2.1	3.8
7	Diabetes mellitus (E10-E14)	158	1.4	2.5
8	Cerebrovascular diseases (I60-I69)	102	0.9	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.8	1.5
10	Influenza and pneumonia (J09-J18)	65	0.6	1.0
11	Septicemia (A40-A41)	62	0.5	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.4	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.6
14	Legal intervention (Y35,Y89.0)	40	0.3	0.6
...	All other causes (residual)	1,591	13.8	25.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	12,398	100.0	209.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,682	37.8	79.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,994	16.1	33.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,364	11.0	23.0
4	Malignant neoplasms (C00-C97)	1,006	8.1	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	405	3.3	6.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	306	2.5	5.2
7	Diabetes mellitus (E10-E14)	225	1.8	3.8
8	Cerebrovascular diseases (I60-I69)	124	1.0	2.1
9	Influenza and pneumonia (J09-J18)	95	0.8	1.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	0.7	1.5
11	Septicemia (A40-A41)	77	0.6	1.3
12	Chronic lower respiratory diseases (J40-J47)	71	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	0.5	1.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.4	0.9
15	Aortic aneurysm and dissection (I71)	41	0.3	0.7
...	All other causes (residual)	1,804	14.6	30.5
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,039	100.0	266.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,191	26.1	69.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,747	17.1	45.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,194	13.7	36.4
4	Malignant neoplasms (C00-C97)	1,919	12.0	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	741	4.6	12.3
6	Diabetes mellitus (E10-E14)	442	2.8	7.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	1.8	4.8
8	Cerebrovascular diseases (I60-I69)	288	1.8	4.8
9	Septicemia (A40-A41)	138	0.9	2.3
10	Influenza and pneumonia (J09-J18)	134	0.8	2.2
11	Chronic lower respiratory diseases (J40-J47)	131	0.8	2.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	107	0.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.6	1.6
14	Aortic aneurysm and dissection (I71)	90	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	0.5	1.5
...	All other causes (residual)	2,446	15.3	40.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	25,542	100.0	386.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,370	21.0	81.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,722	18.5	71.4
3	Malignant neoplasms (C00-C97)	4,276	16.7	64.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,494	9.8	37.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,320	5.2	20.0
6	Diabetes mellitus (E10-E14)	798	3.1	12.1
7	Cerebrovascular diseases (I60-I69)	506	2.0	7.7
8	Chronic lower respiratory diseases (J40-J47)	358	1.4	5.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	1.2	4.7
10	Septicemia (A40-A41)	277	1.1	4.2
11	Influenza and pneumonia (J09-J18)	260	1.0	3.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	247	1.0	3.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	0.8	3.1
14	Viral hepatitis (B15-B19)	191	0.7	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	154	0.6	2.3
...	All other causes (residual)	4,058	15.9	61.4
Non-Hispanic, white, male, 50-54 years				
...	All causes	46,369	100.0	619.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,682	23.0	142.6
2	Malignant neoplasms (C00-C97)	10,528	22.7	140.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,591	12.1	74.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,034	6.5	40.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,616	5.6	34.9
6	Diabetes mellitus (E10-E14)	1,527	3.3	20.4
7	Chronic lower respiratory diseases (J40-J47)	1,220	2.6	16.3
8	Cerebrovascular diseases (I60-I69)	1,032	2.2	13.8
9	Septicemia (A40-A41)	518	1.1	6.9
10	Influenza and pneumonia (J09-J18)	475	1.0	6.3
11	Viral hepatitis (B15-B19)	464	1.0	6.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	0.9	5.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	369	0.8	4.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	353	0.8	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	0.8	4.7
...	All other causes (residual)	7,175	15.5	95.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	69,459	100.0	913.5
1	Malignant neoplasms (C00-C97)	19,920	28.7	262.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,461	23.7	216.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,463	7.9	71.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,346	4.8	44.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,921	4.2	38.4
6	Chronic lower respiratory diseases (J40-J47)	2,770	4.0	36.4
7	Diabetes mellitus (E10-E14)	2,400	3.5	31.6
8	Cerebrovascular diseases (I60-I69)	1,709	2.5	22.5
9	Septicemia (A40-A41)	870	1.3	11.4
10	Viral hepatitis (B15-B19)	785	1.1	10.3
11	Influenza and pneumonia (J09-J18)	725	1.0	9.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	705	1.0	9.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	578	0.8	7.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	322	0.5	4.2
15	Aortic aneurysm and dissection (I71)	309	0.4	4.1
...	All other causes (residual)	10,175	14.6	133.8
Non-Hispanic, white, male, 60-64 years				
...	All causes	88,164	100.0	1,293.8
1	Malignant neoplasms (C00-C97)	27,909	31.7	409.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,507	24.4	315.6
3	Chronic lower respiratory diseases (J40-J47)	4,816	5.5	70.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,347	4.9	63.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,261	3.7	47.9
6	Diabetes mellitus (E10-E14)	3,159	3.6	46.4
7	Cerebrovascular diseases (I60-I69)	2,461	2.8	36.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,173	2.5	31.9
9	Septicemia (A40-A41)	1,251	1.4	18.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,098	1.2	16.1
11	Influenza and pneumonia (J09-J18)	1,043	1.2	15.3
12	Viral hepatitis (B15-B19)	880	1.0	12.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	790	0.9	11.6
14	Aortic aneurysm and dissection (I71)	441	0.5	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	435	0.5	6.4
...	All other causes (residual)	12,593	14.3	184.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	105,474	100.0	1,790.6
1	Malignant neoplasms (C00-C97)	35,234	33.4	598.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,392	24.1	431.1
3	Chronic lower respiratory diseases (J40-J47)	7,194	6.8	122.1
4	Diabetes mellitus (E10-E14)	3,956	3.8	67.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,625	3.4	61.5
6	Cerebrovascular diseases (I60-I69)	3,313	3.1	56.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,441	2.3	41.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,729	1.6	29.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,557	1.5	26.4
10	Septicemia (A40-A41)	1,519	1.4	25.8
11	Influenza and pneumonia (J09-J18)	1,445	1.4	24.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	916	0.9	15.6
13	Parkinson's disease (G20-G21)	684	0.6	11.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	637	0.6	10.8
15	Alzheimer's disease (G30)	591	0.6	10.0
...	All other causes (residual)	15,241	14.5	258.7
Non-Hispanic, white, male, 70-74 years				
...	All causes	114,786	100.0	2,736.6
1	Malignant neoplasms (C00-C97)	36,680	32.0	874.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,273	23.8	650.2
3	Chronic lower respiratory diseases (J40-J47)	9,439	8.2	225.0
4	Cerebrovascular diseases (I60-I69)	4,407	3.8	105.1
5	Diabetes mellitus (E10-E14)	3,981	3.5	94.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,038	2.6	72.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,990	1.7	47.4
8	Influenza and pneumonia (J09-J18)	1,878	1.6	44.8
9	Septicemia (A40-A41)	1,703	1.5	40.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,609	1.4	38.4
11	Parkinson's disease (G20-G21)	1,503	1.3	35.8
12	Alzheimer's disease (G30)	1,498	1.3	35.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,342	1.2	32.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	892	0.8	21.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	873	0.8	20.8
...	All other causes (residual)	16,680	14.5	397.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	123,883	100.0	4,316.7
1	Malignant neoplasms (C00-C97)	34,739	28.0	1,210.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,866	24.1	1,040.7
3	Chronic lower respiratory diseases (J40-J47)	10,416	8.4	362.9
4	Cerebrovascular diseases (I60-I69)	5,646	4.6	196.7
5	Diabetes mellitus (E10-E14)	3,669	3.0	127.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,385	2.7	118.0
7	Alzheimer's disease (G30)	3,247	2.6	113.1
8	Parkinson's disease (G20-G21)	2,656	2.1	92.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,353	1.9	82.0
10	Influenza and pneumonia (J09-J18)	2,343	1.9	81.6
11	Septicemia (A40-A41)	1,883	1.5	65.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,126	0.9	39.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,084	0.9	37.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,068	0.9	37.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,044	0.8	36.4
...	All other causes (residual)	19,358	15.6	674.5
Non-Hispanic, white, male, 80-84 years				
...	All causes	141,645	100.0	7,236.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,724	25.9	1,876.1
2	Malignant neoplasms (C00-C97)	32,416	22.9	1,656.0
3	Chronic lower respiratory diseases (J40-J47)	10,474	7.4	535.1
4	Cerebrovascular diseases (I60-I69)	7,484	5.3	382.3
5	Alzheimer's disease (G30)	6,049	4.3	309.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,905	2.8	199.5
7	Diabetes mellitus (E10-E14)	3,842	2.7	196.3
8	Parkinson's disease (G20-G21)	3,791	2.7	193.7
9	Influenza and pneumonia (J09-J18)	3,359	2.4	171.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,106	2.2	158.7
11	Septicemia (A40-A41)	2,067	1.5	105.6
12	Pneumonitis due to solids and liquids (J69)	1,465	1.0	74.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,293	0.9	66.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,253	0.9	64.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	874	0.6	44.6
...	All other causes (residual)	23,543	16.6	1,202.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	145,196	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,725	28.7	---
2	Malignant neoplasms (C00-C97)	25,381	17.5	---
3	Chronic lower respiratory diseases (J40-J47)	9,363	6.4	---
4	Cerebrovascular diseases (I60-I69)	8,234	5.7	---
5	Alzheimer's disease (G30)	8,183	5.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,327	3.0	---
7	Influenza and pneumonia (J09-J18)	4,069	2.8	---
8	Parkinson's disease (G20-G21)	3,628	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,441	2.4	---
10	Diabetes mellitus (E10-E14)	3,207	2.2	---
11	Septicemia (A40-A41)	2,115	1.5	---
12	Pneumonitis due to solids and liquids (J69)	1,948	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,492	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,244	0.9	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	649	0.4	---
...	All other causes (residual)	26,190	18.0	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	98,276	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,584	32.1	---
2	Malignant neoplasms (C00-C97)	12,659	12.9	---
3	Alzheimer's disease (G30)	6,426	6.5	---
4	Cerebrovascular diseases (I60-I69)	5,683	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	4,947	5.0	---
6	Influenza and pneumonia (J09-J18)	3,370	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,968	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,359	2.4	---
9	Diabetes mellitus (E10-E14)	1,843	1.9	---
10	Parkinson's disease (G20-G21)	1,763	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,687	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,304	1.3	---
13	Septicemia (A40-A41)	1,288	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	737	0.7	---
15	Atherosclerosis (I70)	364	0.4	---
...	All other causes (residual)	19,294	19.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	30,605	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,813	35.3	---
2	Malignant neoplasms (C00-C97)	2,805	9.2	---
3	Alzheimer's disease (G30)	2,033	6.6	---
4	Cerebrovascular diseases (I60-I69)	1,718	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,362	4.5	---
6	Influenza and pneumonia (J09-J18)	1,319	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	955	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	747	2.4	---
9	Pneumonitis due to solids and liquids (J69)	600	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	488	1.6	---
11	Diabetes mellitus (E10-E14)	429	1.4	---
12	Septicemia (A40-A41)	366	1.2	---
13	Parkinson's disease (G20-G21)	350	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	176	0.6	---
15	Atherosclerosis (I70)	148	0.5	---
...	All other causes (residual)	6,296	20.6	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	4,008	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,513	37.7	---
2	Cerebrovascular diseases (I60-I69)	257	6.4	---
3	Malignant neoplasms (C00-C97)	245	6.1	---
4	Alzheimer's disease (G30)	242	6.0	---
5	Influenza and pneumonia (J09-J18)	210	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	144	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.7	---
9	Pneumonitis due to solids and liquids (J69)	89	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.8	---
11	Septicemia (A40-A41)	47	1.2	---
12	Atherosclerosis (I70)	37	0.9	---
13	Diabetes mellitus (E10-E14)	26	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	---
15	Nutritional deficiencies (E40-E64)	22	0.5	---
...	All other causes (residual)	864	21.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,059,926	100.0	1,038.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,388	22.5	233.6
2	Malignant neoplasms (C00-C97)	219,563	20.7	215.1
3	Chronic lower respiratory diseases (J40-J47)	73,371	6.9	71.9
4	Alzheimer's disease (G30)	64,694	6.1	63.4
5	Cerebrovascular diseases (I60-I69)	63,730	6.0	62.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,112	4.1	42.2
7	Influenza and pneumonia (J09-J18)	24,247	2.3	23.8
8	Diabetes mellitus (E10-E14)	23,777	2.2	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,130	1.6	16.8
10	Septicemia (A40-A41)	15,931	1.5	15.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,102	1.2	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,797	1.0	10.6
13	Parkinson's disease (G20-G21)	9,561	0.9	9.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,420	0.8	8.3
15	Pneumonitis due to solids and liquids (J69)	7,345	0.7	7.2
...	All other causes (residual)	226,758	21.4	222.2
Non-Hispanic, white, female, 1-4 years				
...	All causes	812	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	28.3	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	11.3	2.3
3	Malignant neoplasms (C00-C97)	90	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	7.0	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.9	0.8
6	Influenza and pneumonia (J09-J18)	20	2.5	0.5
7	Septicemia (A40-A41)	11	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
10	Meningitis (G00,G03)	6	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	237	29.2	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	523	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	31.0	3.1
2	Malignant neoplasms (C00-C97)	112	21.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.3	*
6	Influenza and pneumonia (J09-J18)	12	2.3	*
8	Cerebrovascular diseases (I60-I69)	11	2.1	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.9	*
10	Septicemia (A40-A41)	7	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Anemias (D50-D64)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
...	All other causes (residual)	104	19.9	2.0
Non-Hispanic, white, female, 10-14 years				
...	All causes	678	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	21.1	2.6
2	Malignant neoplasms (C00-C97)	117	17.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	15.3	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	4.9	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.7	*
7	Influenza and pneumonia (J09-J18)	10	1.5	*
8	Septicemia (A40-A41)	9	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
8	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	8	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Anemias (D50-D64)	3	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	166	24.5	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,764	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	785	44.5	13.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	19.7	6.1
3	Malignant neoplasms (C00-C97)	128	7.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	3.9	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	2.8	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	2.8	0.9
7	Cerebrovascular diseases (I60-I69)	19	1.1	*
7	Influenza and pneumonia (J09-J18)	19	1.1	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
10	Diabetes mellitus (E10-E14)	12	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
15	Meningitis (G00,G03)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (residual)	233	13.2	4.1
Non-Hispanic, white, female, 20-24 years				
...	All causes	3,019	100.0	48.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,489	49.3	24.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	381	12.6	6.2
3	Malignant neoplasms (C00-C97)	185	6.1	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	4.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	3.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.7	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.7	0.8
8	Diabetes mellitus (E10-E14)	34	1.1	0.5
9	Influenza and pneumonia (J09-J18)	26	0.9	0.4
10	Septicemia (A40-A41)	23	0.8	0.4
11	Cerebrovascular diseases (I60-I69)	21	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
...	All other causes (residual)	464	15.4	7.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	4,252	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,791	42.1	28.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	11.3	7.6
3	Malignant neoplasms (C00-C97)	356	8.4	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	254	6.0	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	3.8	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	103	2.4	1.6
7	Diabetes mellitus (E10-E14)	65	1.5	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.4	0.9
9	Influenza and pneumonia (J09-J18)	50	1.2	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.1	0.7
11	Cerebrovascular diseases (I60-I69)	46	1.1	0.7
12	Septicemia (A40-A41)	40	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	24	0.6	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (residual)	747	17.6	11.8
Non-Hispanic, white, female, 30-34 years				
...	All causes	5,644	100.0	90.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,046	36.3	32.9
2	Malignant neoplasms (C00-C97)	698	12.4	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	591	10.5	9.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	418	7.4	6.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	2.9	2.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	152	2.7	2.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	99	1.8	1.6
8	Diabetes mellitus (E10-E14)	98	1.7	1.6
9	Cerebrovascular diseases (I60-I69)	70	1.2	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.1	1.0
11	Septicemia (A40-A41)	60	1.1	1.0
12	Influenza and pneumonia (J09-J18)	58	1.0	0.9
13	Chronic lower respiratory diseases (J40-J47)	40	0.7	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.4	0.4
...	All other causes (residual)	1,038	18.4	16.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,994	100.0	119.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,950	27.9	33.4
2	Malignant neoplasms (C00-C97)	1,190	17.0	20.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	9.6	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	660	9.4	11.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	233	3.3	4.0
6	Diabetes mellitus (E10-E14)	153	2.2	2.6
7	Cerebrovascular diseases (I60-I69)	142	2.0	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	1.8	2.1
9	Septicemia (A40-A41)	82	1.2	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	81	1.2	1.4
11	Influenza and pneumonia (J09-J18)	78	1.1	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.0	1.2
13	Chronic lower respiratory diseases (J40-J47)	58	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.7	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.5	0.6
...	All other causes (residual)	1,420	20.3	24.3
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,206	100.0	170.9
1	Malignant neoplasms (C00-C97)	2,421	23.7	40.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,068	20.3	34.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,203	11.8	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	775	7.6	13.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	453	4.4	7.6
6	Diabetes mellitus (E10-E14)	248	2.4	4.2
7	Cerebrovascular diseases (I60-I69)	242	2.4	4.1
8	Chronic lower respiratory diseases (J40-J47)	184	1.8	3.1
9	Septicemia (A40-A41)	166	1.6	2.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	146	1.4	2.4
11	Influenza and pneumonia (J09-J18)	117	1.1	2.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	0.9	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	0.8	1.4
14	Viral hepatitis (B15-B19)	51	0.5	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	49	0.5	0.8
...	All other causes (residual)	1,913	18.7	32.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	16,773	100.0	253.9
1	Malignant neoplasms (C00-C97)	4,786	28.5	72.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,462	14.7	37.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,308	13.8	34.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	894	5.3	13.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	861	5.1	13.0
6	Diabetes mellitus (E10-E14)	503	3.0	7.6
7	Chronic lower respiratory diseases (J40-J47)	469	2.8	7.1
8	Cerebrovascular diseases (I60-I69)	454	2.7	6.9
9	Septicemia (A40-A41)	270	1.6	4.1
10	Influenza and pneumonia (J09-J18)	234	1.4	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	0.9	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	0.8	2.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.6	1.6
14	Viral hepatitis (B15-B19)	79	0.5	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	0.5	1.2
...	All other causes (residual)	2,983	17.8	45.1
Non-Hispanic, white, female, 50-54 years				
...	All causes	29,894	100.0	391.5
1	Malignant neoplasms (C00-C97)	9,976	33.4	130.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,287	14.3	56.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,102	10.4	40.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,417	4.7	18.6
5	Chronic lower respiratory diseases (J40-J47)	1,416	4.7	18.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,124	3.8	14.7
7	Cerebrovascular diseases (I60-I69)	883	3.0	11.6
8	Diabetes mellitus (E10-E14)	807	2.7	10.6
9	Septicemia (A40-A41)	485	1.6	6.4
10	Influenza and pneumonia (J09-J18)	360	1.2	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	1.0	3.8
12	Viral hepatitis (B15-B19)	248	0.8	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	0.6	2.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	0.6	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	0.4	1.7
...	All other causes (residual)	5,011	16.8	65.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	43,983	100.0	558.7
1	Malignant neoplasms (C00-C97)	16,510	37.5	209.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,644	15.1	84.4
3	Chronic lower respiratory diseases (J40-J47)	2,827	6.4	35.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,776	6.3	35.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,615	3.7	20.5
6	Diabetes mellitus (E10-E14)	1,355	3.1	17.2
7	Cerebrovascular diseases (I60-I69)	1,331	3.0	16.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,082	2.5	13.7
9	Septicemia (A40-A41)	740	1.7	9.4
10	Influenza and pneumonia (J09-J18)	598	1.4	7.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	1.1	6.0
12	Viral hepatitis (B15-B19)	335	0.8	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	0.7	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	0.5	3.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	215	0.5	2.7
...	All other causes (residual)	6,963	15.8	88.4
Non-Hispanic, white, female, 60-64 years				
...	All causes	55,906	100.0	772.2
1	Malignant neoplasms (C00-C97)	21,355	38.2	295.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,294	16.6	128.4
3	Chronic lower respiratory diseases (J40-J47)	4,244	7.6	58.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,989	3.6	27.5
5	Diabetes mellitus (E10-E14)	1,913	3.4	26.4
6	Cerebrovascular diseases (I60-I69)	1,847	3.3	25.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,506	2.7	20.8
8	Septicemia (A40-A41)	1,011	1.8	14.0
9	Influenza and pneumonia (J09-J18)	836	1.5	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	773	1.4	10.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	753	1.3	10.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	420	0.8	5.8
13	Viral hepatitis (B15-B19)	357	0.6	4.9
14	Alzheimer's disease (G30)	314	0.6	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	298	0.5	4.1
...	All other causes (residual)	8,996	16.1	124.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	74,134	100.0	1,163.1
1	Malignant neoplasms (C00-C97)	27,874	37.6	437.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,605	17.0	197.8
3	Chronic lower respiratory diseases (J40-J47)	6,839	9.2	107.3
4	Cerebrovascular diseases (I60-I69)	2,727	3.7	42.8
5	Diabetes mellitus (E10-E14)	2,514	3.4	39.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,800	2.4	28.2
7	Septicemia (A40-A41)	1,440	1.9	22.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,245	1.7	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,233	1.7	19.3
10	Influenza and pneumonia (J09-J18)	1,186	1.6	18.6
11	Alzheimer's disease (G30)	793	1.1	12.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	617	0.8	9.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	480	0.6	7.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	424	0.6	6.7
15	Pneumonitis due to solids and liquids (J69)	332	0.4	5.2
...	All other causes (residual)	12,025	16.2	188.7
Non-Hispanic, white, female, 70-74 years				
...	All causes	89,700	100.0	1,888.3
1	Malignant neoplasms (C00-C97)	30,019	33.5	631.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,021	17.9	337.3
3	Chronic lower respiratory diseases (J40-J47)	9,780	10.9	205.9
4	Cerebrovascular diseases (I60-I69)	4,119	4.6	86.7
5	Diabetes mellitus (E10-E14)	2,662	3.0	56.0
6	Alzheimer's disease (G30)	2,123	2.4	44.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,848	2.1	38.9
8	Septicemia (A40-A41)	1,649	1.8	34.7
9	Influenza and pneumonia (J09-J18)	1,603	1.8	33.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,548	1.7	32.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,032	1.2	21.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	775	0.9	16.3
13	Parkinson's disease (G20-G21)	696	0.8	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	607	0.7	12.8
15	Pneumonitis due to solids and liquids (J69)	460	0.5	9.7
...	All other causes (residual)	14,758	16.5	310.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	108,197	100.0	3,109.8
1	Malignant neoplasms (C00-C97)	29,497	27.3	847.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,885	19.3	600.3
3	Chronic lower respiratory diseases (J40-J47)	11,276	10.4	324.1
4	Cerebrovascular diseases (I60-I69)	6,181	5.7	177.7
5	Alzheimer's disease (G30)	4,887	4.5	140.5
6	Diabetes mellitus (E10-E14)	2,921	2.7	84.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,414	2.2	69.4
8	Influenza and pneumonia (J09-J18)	2,184	2.0	62.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,029	1.9	58.3
10	Septicemia (A40-A41)	1,929	1.8	55.4
11	Parkinson's disease (G20-G21)	1,362	1.3	39.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,120	1.0	32.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	853	0.8	24.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	733	0.7	21.1
15	Pneumonitis due to solids and liquids (J69)	707	0.7	20.3
...	All other causes (residual)	19,219	17.8	552.4
Non-Hispanic, white, female, 80-84 years				
...	All causes	143,961	100.0	5,399.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,435	22.5	1,216.5
2	Malignant neoplasms (C00-C97)	28,615	19.9	1,073.2
3	Chronic lower respiratory diseases (J40-J47)	12,307	8.5	461.6
4	Alzheimer's disease (G30)	10,339	7.2	387.8
5	Cerebrovascular diseases (I60-I69)	9,851	6.8	369.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,529	2.5	132.4
7	Diabetes mellitus (E10-E14)	3,332	2.3	125.0
8	Influenza and pneumonia (J09-J18)	3,279	2.3	123.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,681	1.9	100.6
10	Septicemia (A40-A41)	2,197	1.5	82.4
11	Parkinson's disease (G20-G21)	2,175	1.5	81.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,798	1.2	67.4
13	Pneumonitis due to solids and liquids (J69)	1,054	0.7	39.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,017	0.7	38.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	699	0.5	26.2
...	All other causes (residual)	28,653	19.9	1,074.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	186,447	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,209	25.9	---
2	Malignant neoplasms (C00-C97)	25,255	13.5	---
3	Alzheimer's disease (G30)	16,931	9.1	---
4	Cerebrovascular diseases (I60-I69)	14,409	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	12,123	6.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,044	2.7	---
7	Influenza and pneumonia (J09-J18)	4,971	2.7	---
8	Diabetes mellitus (E10-E14)	3,560	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,327	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,747	1.5	---
11	Septicemia (A40-A41)	2,745	1.5	---
12	Parkinson's disease (G20-G21)	2,553	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,631	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,152	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	773	0.4	---
...	All other causes (residual)	41,017	22.0	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	173,685	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,048	29.4	---
2	Alzheimer's disease (G30)	18,305	10.5	---
3	Malignant neoplasms (C00-C97)	15,255	8.8	---
4	Cerebrovascular diseases (I60-I69)	13,647	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	8,398	4.8	---
6	Influenza and pneumonia (J09-J18)	5,159	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,663	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,984	1.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,963	1.7	---
10	Diabetes mellitus (E10-E14)	2,537	1.5	---
11	Septicemia (A40-A41)	2,098	1.2	---
12	Parkinson's disease (G20-G21)	1,756	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,572	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	951	0.5	---
15	Atherosclerosis (I70)	776	0.4	---
...	All other causes (residual)	41,573	23.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	80,102	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,380	31.7	---
2	Alzheimer's disease (G30)	8,963	11.2	---
3	Cerebrovascular diseases (I60-I69)	6,391	8.0	---
4	Malignant neoplasms (C00-C97)	4,448	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	2,821	3.5	---
6	Influenza and pneumonia (J09-J18)	2,670	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,137	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,584	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,178	1.5	---
10	Diabetes mellitus (E10-E14)	881	1.1	---
11	Septicemia (A40-A41)	787	1.0	---
12	Pneumonitis due to solids and liquids (J69)	744	0.9	---
13	Parkinson's disease (G20-G21)	468	0.6	---
14	Atherosclerosis (I70)	413	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	334	0.4	---
...	All other causes (residual)	20,903	26.1	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	18,713	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,432	34.4	---
2	Alzheimer's disease (G30)	1,883	10.1	---
3	Cerebrovascular diseases (I60-I69)	1,309	7.0	---
4	Influenza and pneumonia (J09-J18)	747	4.0	---
5	Malignant neoplasms (C00-C97)	661	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	514	2.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	446	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	445	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	1.4	---
10	Pneumonitis due to solids and liquids (J69)	177	0.9	---
11	Diabetes mellitus (E10-E14)	174	0.9	---
12	Atherosclerosis (I70)	144	0.8	---
13	Septicemia (A40-A41)	143	0.8	---
14	Nutritional deficiencies (E40-E64)	104	0.6	---
15	Parkinson's disease (G20-G21)	55	0.3	---
...	All other causes (residual)	5,222	27.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	315,254	100.0	754.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,093	23.5	177.4
2	Malignant neoplasms (C00-C97)	68,523	21.7	164.0
3	Cerebrovascular diseases (I60-I69)	17,760	5.6	42.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,366	4.9	36.8
5	Diabetes mellitus (E10-E14)	13,693	4.3	32.8
6	Chronic lower respiratory diseases (J40-J47)	10,327	3.3	24.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,083	2.9	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,038	2.9	21.6
9	Alzheimer's disease (G30)	8,072	2.6	19.3
10	Septicemia (A40-A41)	6,576	2.1	15.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,740	1.8	13.7
12	Influenza and pneumonia (J09-J18)	5,554	1.8	13.3
13	Certain conditions originating in the perinatal period (P00-P96)	3,827	1.2	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,379	1.1	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,185	1.0	7.6
...	All other causes (residual)	61,038	19.4	146.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.1	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	9.2	3.8
4	Malignant neoplasms (C00-C97)	53	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	27	2.7	1.1
6	Certain conditions originating in the perinatal period (P00-P96)	27	2.7	1.1
8	Chronic lower respiratory diseases (J40-J47)	23	2.3	0.9
9	Cerebrovascular diseases (I60-I69)	17	1.7	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (residual)	250	24.6	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	553	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	32.5	5.8
2	Malignant neoplasms (C00-C97)	61	11.0	2.0
3	Chronic lower respiratory diseases (J40-J47)	49	8.9	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.4	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.9	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.1	0.9
7	Anemias (D50-D64)	12	2.2	*
8	Influenza and pneumonia (J09-J18)	11	2.0	*
9	Cerebrovascular diseases (I60-I69)	10	1.8	*
10	Septicemia (A40-A41)	9	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
13	Diseases of appendix (K35-K38)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	90	16.3	2.9
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	634	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	23.2	4.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	12.6	2.6
3	Malignant neoplasms (C00-C97)	65	10.3	2.1
4	Chronic lower respiratory diseases (J40-J47)	53	8.4	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	7.1	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.7	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.4	0.9
8	Cerebrovascular diseases (I60-I69)	12	1.9	*
9	Influenza and pneumonia (J09-J18)	8	1.3	*
10	Septicemia (A40-A41)	7	1.1	*
10	Diabetes mellitus (E10-E14)	7	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Anemias (D50-D64)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (residual)	135	21.3	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,336	100.0	72.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	999	42.8	31.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	23.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	8.5	6.2
4	Malignant neoplasms (C00-C97)	91	3.9	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.6	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.7	1.2
7	Chronic lower respiratory diseases (J40-J47)	30	1.3	0.9
8	Anemias (D50-D64)	24	1.0	0.7
9	Diabetes mellitus (E10-E14)	19	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.7	*
11	Septicemia (A40-A41)	11	0.5	*
11	Influenza and pneumonia (J09-J18)	11	0.5	*
13	Cerebrovascular diseases (I60-I69)	10	0.4	*
14	Legal intervention (Y35,Y89.0)	9	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
...	All other causes (residual)	228	9.8	7.1
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,760	100.0	133.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,014	42.3	56.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,092	22.9	30.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	7.9	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	218	4.6	6.1
5	Malignant neoplasms (C00-C97)	161	3.4	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.5	2.0
7	Chronic lower respiratory diseases (J40-J47)	56	1.2	1.6
8	Anemias (D50-D64)	46	1.0	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	46	1.0	1.3
10	Diabetes mellitus (E10-E14)	45	0.9	1.3
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.6	0.8
13	Legal intervention (Y35,Y89.0)	25	0.5	0.7
14	Influenza and pneumonia (J09-J18)	24	0.5	0.7
15	Septicemia (A40-A41)	23	0.5	0.6
...	All other causes (residual)	509	10.7	14.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	5,006	100.0	154.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,670	33.4	51.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,122	22.4	34.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	7.2	11.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	6.7	10.4
5	Malignant neoplasms (C00-C97)	212	4.2	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	2.9	4.4
7	Diabetes mellitus (E10-E14)	90	1.8	2.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.3	2.0
9	Cerebrovascular diseases (I60-I69)	54	1.1	1.7
10	Anemias (D50-D64)	50	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	48	1.0	1.5
12	Septicemia (A40-A41)	46	0.9	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.9	1.4
14	Influenza and pneumonia (J09-J18)	42	0.8	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.6	1.0
...	All other causes (residual)	689	13.8	21.3
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	5,310	100.0	185.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	22.8	42.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	19.4	36.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	616	11.6	21.5
4	Malignant neoplasms (C00-C97)	419	7.9	14.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	4.8	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	3.4	6.4
7	Diabetes mellitus (E10-E14)	162	3.1	5.7
8	Cerebrovascular diseases (I60-I69)	97	1.8	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.4	2.6
10	Septicemia (A40-A41)	70	1.3	2.4
11	Anemias (D50-D64)	69	1.3	2.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.3	2.4
13	Influenza and pneumonia (J09-J18)	56	1.1	2.0
14	Chronic lower respiratory diseases (J40-J47)	54	1.0	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.8	1.5
...	All other causes (residual)	902	17.0	31.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,273	100.0	233.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,080	17.2	40.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,042	16.6	38.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	859	13.7	32.0
4	Malignant neoplasms (C00-C97)	727	11.6	27.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	260	4.1	9.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	3.5	8.3
7	Diabetes mellitus (E10-E14)	217	3.5	8.1
8	Cerebrovascular diseases (I60-I69)	195	3.1	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.6	3.7
10	Chronic lower respiratory diseases (J40-J47)	83	1.3	3.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	1.3	2.9
12	Septicemia (A40-A41)	74	1.2	2.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	66	1.1	2.5
14	Anemias (D50-D64)	57	0.9	2.1
15	Influenza and pneumonia (J09-J18)	54	0.9	2.0
...	All other causes (residual)	1,160	18.5	43.2
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,029	100.0	307.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,779	22.2	68.1
2	Malignant neoplasms (C00-C97)	1,181	14.7	45.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,003	12.5	38.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	6.8	21.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	333	4.1	12.7
6	Cerebrovascular diseases (I60-I69)	329	4.1	12.6
7	Diabetes mellitus (E10-E14)	323	4.0	12.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	206	2.6	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	2.3	7.1
10	Septicemia (A40-A41)	143	1.8	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.7	5.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	133	1.7	5.1
13	Chronic lower respiratory diseases (J40-J47)	103	1.3	3.9
14	Influenza and pneumonia (J09-J18)	88	1.1	3.4
15	Aortic aneurysm and dissection (I71)	60	0.7	2.3
...	All other causes (residual)	1,481	18.4	56.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	11,774	100.0	446.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,754	23.4	104.5
2	Malignant neoplasms (C00-C97)	2,219	18.8	84.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,196	10.2	45.4
4	Diabetes mellitus (E10-E14)	563	4.8	21.4
5	Cerebrovascular diseases (I60-I69)	547	4.6	20.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	3.7	16.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	425	3.6	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	2.3	10.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	243	2.1	9.2
10	Septicemia (A40-A41)	240	2.0	9.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	188	1.6	7.1
12	Chronic lower respiratory diseases (J40-J47)	187	1.6	7.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	184	1.6	7.0
14	Influenza and pneumonia (J09-J18)	137	1.2	5.2
15	Aortic aneurysm and dissection (I71)	70	0.6	2.7
...	All other causes (residual)	2,125	18.0	80.6
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,398	100.0	703.1
1	Malignant neoplasms (C00-C97)	4,818	24.8	174.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,663	24.0	169.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,522	7.8	55.2
4	Diabetes mellitus (E10-E14)	923	4.8	33.5
5	Cerebrovascular diseases (I60-I69)	887	4.6	32.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	529	2.7	19.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	486	2.5	17.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	2.2	15.2
9	Chronic lower respiratory diseases (J40-J47)	418	2.2	15.2
10	Septicemia (A40-A41)	394	2.0	14.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	1.9	13.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	320	1.6	11.6
13	Influenza and pneumonia (J09-J18)	284	1.5	10.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	0.9	6.5
15	Viral hepatitis (B15-B19)	119	0.6	4.3
...	All other causes (residual)	3,078	15.9	111.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	28,015	100.0	1,078.9
1	Malignant neoplasms (C00-C97)	8,047	28.7	309.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,897	24.6	265.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,452	5.2	55.9
4	Diabetes mellitus (E10-E14)	1,376	4.9	53.0
5	Cerebrovascular diseases (I60-I69)	1,316	4.7	50.7
6	Chronic lower respiratory diseases (J40-J47)	836	3.0	32.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	734	2.6	28.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	637	2.3	24.5
9	Septicemia (A40-A41)	603	2.2	23.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	560	2.0	21.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	472	1.7	18.2
12	Influenza and pneumonia (J09-J18)	366	1.3	14.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	250	0.9	9.6
14	Viral hepatitis (B15-B19)	246	0.9	9.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	0.5	5.5
...	All other causes (residual)	4,079	14.6	157.1
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	32,871	100.0	1,571.1
1	Malignant neoplasms (C00-C97)	9,923	30.2	474.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,171	24.9	390.5
3	Diabetes mellitus (E10-E14)	1,682	5.1	80.4
3	Cerebrovascular diseases (I60-I69)	1,682	5.1	80.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,255	3.8	60.0
6	Chronic lower respiratory diseases (J40-J47)	1,085	3.3	51.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	2.8	44.6
8	Septicemia (A40-A41)	688	2.1	32.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	636	1.9	30.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	594	1.8	28.4
11	Influenza and pneumonia (J09-J18)	494	1.5	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	418	1.3	20.0
13	Viral hepatitis (B15-B19)	333	1.0	15.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	0.5	7.8
15	Pneumonitis due to solids and liquids (J69)	128	0.4	6.1
...	All other causes (residual)	4,686	14.3	224.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	32,541	100.0	2,069.7
1	Malignant neoplasms (C00-C97)	10,023	30.8	637.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,978	24.5	507.4
3	Cerebrovascular diseases (I60-I69)	1,758	5.4	111.8
4	Diabetes mellitus (E10-E14)	1,702	5.2	108.3
5	Chronic lower respiratory diseases (J40-J47)	1,281	3.9	81.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,108	3.4	70.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	783	2.4	49.8
8	Septicemia (A40-A41)	762	2.3	48.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	567	1.7	36.1
10	Influenza and pneumonia (J09-J18)	534	1.6	34.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	439	1.3	27.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	240	0.7	15.3
13	Viral hepatitis (B15-B19)	206	0.6	13.1
14	Alzheimer's disease (G30)	168	0.5	10.7
15	Pneumonitis due to solids and liquids (J69)	165	0.5	10.5
...	All other causes (residual)	4,827	14.8	307.0
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	30,777	100.0	2,964.9
1	Malignant neoplasms (C00-C97)	8,905	28.9	857.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,793	25.3	750.8
3	Cerebrovascular diseases (I60-I69)	1,874	6.1	180.5
4	Diabetes mellitus (E10-E14)	1,596	5.2	153.8
5	Chronic lower respiratory diseases (J40-J47)	1,421	4.6	136.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,040	3.4	100.2
7	Septicemia (A40-A41)	732	2.4	70.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	584	1.9	56.3
9	Influenza and pneumonia (J09-J18)	535	1.7	51.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	1.7	49.9
11	Alzheimer's disease (G30)	467	1.5	45.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	236	0.8	22.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	169	0.5	16.3
14	Pneumonitis due to solids and liquids (J69)	159	0.5	15.3
15	Parkinson's disease (G20-G21)	142	0.5	13.7
...	All other causes (residual)	4,606	15.0	443.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	30,976	100.0	4,311.4
1	Malignant neoplasms (C00-C97)	7,875	25.4	1,096.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,799	25.2	1,085.5
3	Cerebrovascular diseases (I60-I69)	2,165	7.0	301.3
4	Diabetes mellitus (E10-E14)	1,615	5.2	224.8
5	Chronic lower respiratory diseases (J40-J47)	1,390	4.5	193.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,145	3.7	159.4
7	Alzheimer's disease (G30)	909	2.9	126.5
8	Septicemia (A40-A41)	766	2.5	106.6
9	Influenza and pneumonia (J09-J18)	605	2.0	84.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	602	1.9	83.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	498	1.6	69.3
12	Parkinson's disease (G20-G21)	223	0.7	31.0
13	Pneumonitis due to solids and liquids (J69)	222	0.7	30.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.5	23.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	132	0.4	18.4
...	All other causes (residual)	4,862	15.7	676.7
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	30,485	100.0	6,427.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,763	25.5	1,636.7
2	Malignant neoplasms (C00-C97)	6,238	20.5	1,315.2
3	Cerebrovascular diseases (I60-I69)	2,354	7.7	496.3
4	Alzheimer's disease (G30)	1,526	5.0	321.7
5	Diabetes mellitus (E10-E14)	1,409	4.6	297.1
6	Chronic lower respiratory diseases (J40-J47)	1,339	4.4	282.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,076	3.5	226.9
8	Septicemia (A40-A41)	719	2.4	151.6
9	Influenza and pneumonia (J09-J18)	712	2.3	150.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	700	2.3	147.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	452	1.5	95.3
12	Parkinson's disease (G20-G21)	272	0.9	57.3
13	Pneumonitis due to solids and liquids (J69)	235	0.8	49.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	204	0.7	43.0
15	Aortic aneurysm and dissection (I71)	98	0.3	20.7
...	All other causes (residual)	5,388	17.7	1,136.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	27,456	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,395	26.9	---
2	Malignant neoplasms (C00-C97)	4,467	16.3	---
3	Cerebrovascular diseases (I60-I69)	2,124	7.7	---
4	Alzheimer's disease (G30)	1,914	7.0	---
5	Diabetes mellitus (E10-E14)	1,122	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,051	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	942	3.4	---
8	Septicemia (A40-A41)	635	2.3	---
9	Influenza and pneumonia (J09-J18)	629	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	601	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	1.4	---
12	Pneumonitis due to solids and liquids (J69)	245	0.9	---
13	Parkinson's disease (G20-G21)	240	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	171	0.6	---
15	Atherosclerosis (I70)	98	0.4	---
...	All other causes (residual)	5,424	19.8	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	19,189	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,395	28.1	---
2	Malignant neoplasms (C00-C97)	2,219	11.6	---
3	Alzheimer's disease (G30)	1,767	9.2	---
4	Cerebrovascular diseases (I60-I69)	1,473	7.7	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	661	3.4	---
6	Diabetes mellitus (E10-E14)	586	3.1	---
7	Influenza and pneumonia (J09-J18)	566	2.9	---
8	Chronic lower respiratory diseases (J40-J47)	552	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	530	2.8	---
10	Septicemia (A40-A41)	416	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	1.2	---
12	Pneumonitis due to solids and liquids (J69)	192	1.0	---
13	Parkinson's disease (G20-G21)	111	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	---
15	Nutritional deficiencies (E40-E64)	84	0.4	---
...	All other causes (residual)	4,299	22.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	8,333	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,419	29.0	---
2	Alzheimer's disease (G30)	890	10.7	---
3	Malignant neoplasms (C00-C97)	682	8.2	---
4	Cerebrovascular diseases (I60-I69)	624	7.5	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	2.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	2.7	---
7	Influenza and pneumonia (J09-J18)	219	2.6	---
8	Chronic lower respiratory diseases (J40-J47)	208	2.5	---
9	Diabetes mellitus (E10-E14)	201	2.4	---
10	Septicemia (A40-A41)	121	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	1.4	---
12	Pneumonitis due to solids and liquids (J69)	82	1.0	---
13	Nutritional deficiencies (E40-E64)	54	0.6	---
14	Parkinson's disease (G20-G21)	45	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	---
...	All other causes (residual)	2,157	25.9	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,598	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	762	29.3	---
2	Alzheimer's disease (G30)	314	12.1	---
3	Cerebrovascular diseases (I60-I69)	176	6.8	---
4	Malignant neoplasms (C00-C97)	128	4.9	---
5	Influenza and pneumonia (J09-J18)	95	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.9	---
8	Diabetes mellitus (E10-E14)	48	1.8	---
8	Chronic lower respiratory diseases (J40-J47)	48	1.8	---
10	Septicemia (A40-A41)	44	1.7	---
11	Pneumonitis due to solids and liquids (J69)	28	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.1	---
13	Nutritional deficiencies (E40-E64)	26	1.0	---
14	Atherosclerosis (I70)	14	0.5	---
15	Anemias (D50-D64)	8	0.3	---
...	All other causes (residual)	711	27.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	161,850	100.0	809.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,638	23.9	193.2
2	Malignant neoplasms (C00-C97)	34,787	21.5	174.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,516	6.5	52.6
4	Cerebrovascular diseases (I60-I69)	7,962	4.9	39.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,909	4.9	39.6
6	Diabetes mellitus (E10-E14)	6,806	4.2	34.0
7	Chronic lower respiratory diseases (J40-J47)	5,236	3.2	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,351	2.7	21.8
9	Septicemia (A40-A41)	3,061	1.9	15.3
10	Influenza and pneumonia (J09-J18)	2,729	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,599	1.6	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,230	1.4	11.2
13	Alzheimer's disease (G30)	2,154	1.3	10.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,130	1.3	10.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,951	1.2	9.8
...	All other causes (residual)	28,791	17.8	144.0
Non-Hispanic, black, male, 1-4 years				
...	All causes	608	100.0	48.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	29.6	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.8	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	8.6	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.3
5	Malignant neoplasms (C00-C97)	26	4.3	2.1
6	Certain conditions originating in the perinatal period (P00-P96)	19	3.1	*
7	Influenza and pneumonia (J09-J18)	17	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
9	Cerebrovascular diseases (I60-I69)	12	2.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.3	*
11	Septicemia (A40-A41)	7	1.2	*
11	Anemias (D50-D64)	7	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (residual)	137	22.5	11.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	317	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	31.9	6.4
2	Malignant neoplasms (C00-C97)	34	10.7	2.1
3	Chronic lower respiratory diseases (J40-J47)	30	9.5	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	7.3	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	7.3	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.7	*
7	Anemias (D50-D64)	5	1.6	*
7	Cerebrovascular diseases (I60-I69)	5	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
9	Influenza and pneumonia (J09-J18)	4	1.3	*
11	Septicemia (A40-A41)	3	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Meningitis (G00,G03)	2	0.6	*
13	Diseases of appendix (K35-K38)	2	0.6	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
...	All other causes (residual)	61	19.2	3.9
Non-Hispanic, black, male, 10-14 years				
...	All causes	386	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	22.3	5.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	15.5	3.9
3	Malignant neoplasms (C00-C97)	34	8.8	2.2
4	Chronic lower respiratory diseases (J40-J47)	33	8.5	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	8.5	2.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.9	*
8	Cerebrovascular diseases (I60-I69)	8	2.1	*
9	Diabetes mellitus (E10-E14)	6	1.6	*
10	Influenza and pneumonia (J09-J18)	5	1.3	*
11	Septicemia (A40-A41)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Anemias (D50-D64)	2	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Meningococcal infection (A39)	1	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	76	19.7	4.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,795	100.0	110.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	901	50.2	55.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	22.6	24.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	8.4	9.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.1	3.4
5	Malignant neoplasms (C00-C97)	50	2.8	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.3	1.4
7	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
8	Diabetes mellitus (E10-E14)	13	0.7	*
9	Legal intervention (Y35,Y89.0)	9	0.5	*
10	Anemias (D50-D64)	8	0.4	*
11	Cerebrovascular diseases (I60-I69)	6	0.3	*
11	Influenza and pneumonia (J09-J18)	6	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.1	*
...	All other causes (residual)	137	7.6	8.4
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,691	100.0	203.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,843	49.9	101.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	820	22.2	45.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	8.2	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	3.8	7.7
5	Malignant neoplasms (C00-C97)	88	2.4	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	43	1.2	2.4
7	Chronic lower respiratory diseases (J40-J47)	37	1.0	2.0
8	Anemias (D50-D64)	27	0.7	1.5
9	Diabetes mellitus (E10-E14)	26	0.7	1.4
10	Legal intervention (Y35,Y89.0)	23	0.6	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.6	1.2
12	Influenza and pneumonia (J09-J18)	15	0.4	*
13	Cerebrovascular diseases (I60-I69)	14	0.4	*
14	Septicemia (A40-A41)	11	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (residual)	268	7.3	14.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,680	100.0	230.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,506	40.9	94.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	22.6	52.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	280	7.6	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	6.1	14.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	109	3.0	6.8
6	Malignant neoplasms (C00-C97)	99	2.7	6.2
7	Diabetes mellitus (E10-E14)	44	1.2	2.8
8	Influenza and pneumonia (J09-J18)	32	0.9	2.0
8	Chronic lower respiratory diseases (J40-J47)	32	0.9	2.0
10	Cerebrovascular diseases (I60-I69)	30	0.8	1.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	1.8
12	Anemias (D50-D64)	27	0.7	1.7
13	Septicemia (A40-A41)	25	0.7	1.6
14	Legal intervention (Y35,Y89.0)	24	0.7	1.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
...	All other causes (residual)	372	10.1	23.3
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,549	100.0	258.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,086	30.6	79.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	769	21.7	55.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	10.5	27.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	5.9	15.3
5	Malignant neoplasms (C00-C97)	169	4.8	12.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	111	3.1	8.1
7	Diabetes mellitus (E10-E14)	87	2.5	6.3
8	Cerebrovascular diseases (I60-I69)	46	1.3	3.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.0	2.7
10	Septicemia (A40-A41)	36	1.0	2.6
10	Chronic lower respiratory diseases (J40-J47)	36	1.0	2.6
12	Influenza and pneumonia (J09-J18)	35	1.0	2.5
13	Anemias (D50-D64)	32	0.9	2.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.6	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.6	1.5
...	All other causes (residual)	458	12.9	33.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,942	100.0	310.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	20.2	62.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	771	19.6	60.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	672	17.0	52.9
4	Malignant neoplasms (C00-C97)	276	7.0	21.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	4.7	14.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	3.7	11.3
7	Diabetes mellitus (E10-E14)	129	3.3	10.1
8	Cerebrovascular diseases (I60-I69)	107	2.7	8.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.3	4.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.2	3.8
11	Chronic lower respiratory diseases (J40-J47)	43	1.1	3.4
12	Septicemia (A40-A41)	36	0.9	2.8
13	Influenza and pneumonia (J09-J18)	34	0.9	2.7
14	Anemias (D50-D64)	30	0.8	2.4
15	Aortic aneurysm and dissection (I71)	27	0.7	2.1
...	All other causes (residual)	592	15.0	46.6
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,668	100.0	380.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,130	24.2	92.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	15.0	57.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	466	10.0	38.0
4	Malignant neoplasms (C00-C97)	421	9.0	34.3
5	Cerebrovascular diseases (I60-I69)	192	4.1	15.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	189	4.0	15.4
7	Diabetes mellitus (E10-E14)	188	4.0	15.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	172	3.7	14.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	2.2	8.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.7	6.6
11	Septicemia (A40-A41)	58	1.2	4.7
12	Chronic lower respiratory diseases (J40-J47)	57	1.2	4.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.2	4.6
14	Aortic aneurysm and dissection (I71)	48	1.0	3.9
15	Influenza and pneumonia (J09-J18)	46	1.0	3.7
...	All other causes (residual)	759	16.3	61.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	6,686	100.0	539.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,700	25.4	137.0
2	Malignant neoplasms (C00-C97)	969	14.5	78.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	832	12.4	67.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	5.4	28.9
5	Diabetes mellitus (E10-E14)	315	4.7	25.4
6	Cerebrovascular diseases (I60-I69)	299	4.5	24.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	255	3.8	20.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.4	12.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	2.1	11.4
10	Septicemia (A40-A41)	123	1.8	9.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	116	1.7	9.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.7	9.0
13	Chronic lower respiratory diseases (J40-J47)	91	1.4	7.3
14	Influenza and pneumonia (J09-J18)	87	1.3	7.0
15	Aortic aneurysm and dissection (I71)	58	0.9	4.7
...	All other causes (residual)	1,069	16.0	86.2
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,126	100.0	861.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,960	26.6	229.3
2	Malignant neoplasms (C00-C97)	2,297	20.6	177.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,068	9.6	82.7
4	Diabetes mellitus (E10-E14)	543	4.9	42.1
5	Cerebrovascular diseases (I60-I69)	496	4.5	38.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	353	3.2	27.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	2.7	23.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	287	2.6	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.2	19.1
10	Chronic lower respiratory diseases (J40-J47)	195	1.8	15.1
11	Septicemia (A40-A41)	194	1.7	15.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.6	14.0
13	Influenza and pneumonia (J09-J18)	169	1.5	13.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.2	10.5
15	Viral hepatitis (B15-B19)	77	0.7	6.0
...	All other causes (residual)	1,623	14.6	125.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	16,180	100.0	1,347.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,335	26.8	361.0
2	Malignant neoplasms (C00-C97)	4,157	25.7	346.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,061	6.6	88.4
4	Diabetes mellitus (E10-E14)	772	4.8	64.3
5	Cerebrovascular diseases (I60-I69)	770	4.8	64.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.5	34.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	402	2.5	33.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	397	2.5	33.1
9	Chronic lower respiratory diseases (J40-J47)	377	2.3	31.4
10	Septicemia (A40-A41)	294	1.8	24.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	1.8	23.9
12	Influenza and pneumonia (J09-J18)	215	1.3	17.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	1.2	16.7
14	Viral hepatitis (B15-B19)	163	1.0	13.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	0.7	9.7
...	All other causes (residual)	2,223	13.7	185.1
Non-Hispanic, black, male, 60-64 years				
...	All causes	19,263	100.0	2,052.3
1	Malignant neoplasms (C00-C97)	5,489	28.5	584.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,116	26.6	545.1
3	Cerebrovascular diseases (I60-I69)	960	5.0	102.3
4	Diabetes mellitus (E10-E14)	952	4.9	101.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	926	4.8	98.7
6	Chronic lower respiratory diseases (J40-J47)	579	3.0	61.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	521	2.7	55.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	440	2.3	46.9
9	Septicemia (A40-A41)	365	1.9	38.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	321	1.7	34.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	310	1.6	33.0
12	Influenza and pneumonia (J09-J18)	289	1.5	30.8
13	Viral hepatitis (B15-B19)	242	1.3	25.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	0.7	14.1
15	Pneumonitis due to solids and liquids (J69)	75	0.4	8.0
...	All other causes (residual)	2,546	13.2	271.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	18,438	100.0	2,684.5
1	Malignant neoplasms (C00-C97)	5,487	29.8	798.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,833	26.2	703.7
3	Cerebrovascular diseases (I60-I69)	981	5.3	142.8
4	Diabetes mellitus (E10-E14)	958	5.2	139.5
5	Chronic lower respiratory diseases (J40-J47)	732	4.0	106.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	558	3.0	81.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	522	2.8	76.0
8	Septicemia (A40-A41)	406	2.2	59.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	313	1.7	45.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	292	1.6	42.5
11	Influenza and pneumonia (J09-J18)	276	1.5	40.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	166	0.9	24.2
13	Viral hepatitis (B15-B19)	145	0.8	21.1
14	Pneumonitis due to solids and liquids (J69)	88	0.5	12.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.4	11.8
...	All other causes (residual)	2,600	14.1	378.6
Non-Hispanic, black, male, 70-74 years				
...	All causes	16,429	100.0	3,780.6
1	Malignant neoplasms (C00-C97)	4,789	29.1	1,102.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,327	26.3	995.7
3	Cerebrovascular diseases (I60-I69)	976	5.9	224.6
4	Diabetes mellitus (E10-E14)	793	4.8	182.5
5	Chronic lower respiratory diseases (J40-J47)	776	4.7	178.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	541	3.3	124.5
7	Septicemia (A40-A41)	378	2.3	87.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	2.0	75.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	293	1.8	67.4
10	Influenza and pneumonia (J09-J18)	277	1.7	63.7
11	Alzheimer's disease (G30)	188	1.1	43.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.9	33.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.6	21.9
14	Parkinson's disease (G20-G21)	90	0.5	20.7
15	Pneumonitis due to solids and liquids (J69)	88	0.5	20.3
...	All other causes (residual)	2,342	14.3	538.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	15,455	100.0	5,424.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,107	26.6	1,441.4
2	Malignant neoplasms (C00-C97)	4,076	26.4	1,430.5
3	Cerebrovascular diseases (I60-I69)	1,013	6.6	355.5
4	Diabetes mellitus (E10-E14)	750	4.9	263.2
5	Chronic lower respiratory diseases (J40-J47)	742	4.8	260.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	543	3.5	190.6
7	Septicemia (A40-A41)	362	2.3	127.0
8	Alzheimer's disease (G30)	339	2.2	119.0
9	Influenza and pneumonia (J09-J18)	318	2.1	111.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	2.0	107.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	260	1.7	91.2
12	Parkinson's disease (G20-G21)	132	0.9	46.3
13	Pneumonitis due to solids and liquids (J69)	127	0.8	44.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	74	0.5	26.0
...	All other causes (residual)	2,226	14.4	781.2
Non-Hispanic, black, male, 80-84 years				
...	All causes	13,485	100.0	7,879.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,540	26.3	2,068.6
2	Malignant neoplasms (C00-C97)	3,060	22.7	1,788.1
3	Cerebrovascular diseases (I60-I69)	902	6.7	527.1
4	Chronic lower respiratory diseases (J40-J47)	649	4.8	379.2
5	Diabetes mellitus (E10-E14)	600	4.4	350.6
6	Alzheimer's disease (G30)	492	3.6	287.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	3.4	265.9
8	Influenza and pneumonia (J09-J18)	352	2.6	205.7
9	Septicemia (A40-A41)	302	2.2	176.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	276	2.0	161.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	1.7	133.2
12	Parkinson's disease (G20-G21)	169	1.3	98.8
13	Pneumonitis due to solids and liquids (J69)	123	0.9	71.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.7	54.9
15	Aortic aneurysm and dissection (I71)	39	0.3	22.8
...	All other causes (residual)	2,204	16.3	1,287.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,140	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,711	26.7	---
2	Malignant neoplasms (C00-C97)	2,039	20.1	---
3	Cerebrovascular diseases (I60-I69)	625	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	480	4.7	---
5	Alzheimer's disease (G30)	475	4.7	---
6	Diabetes mellitus (E10-E14)	407	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	366	3.6	---
8	Influenza and pneumonia (J09-J18)	251	2.5	---
9	Septicemia (A40-A41)	243	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	2.0	---
12	Parkinson's disease (G20-G21)	127	1.3	---
12	Pneumonitis due to solids and liquids (J69)	127	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.6	---
15	Nutritional deficiencies (E40-E64)	44	0.4	---
...	All other causes (residual)	1,774	17.5	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	5,884	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,634	27.8	---
2	Malignant neoplasms (C00-C97)	957	16.3	---
3	Cerebrovascular diseases (I60-I69)	373	6.3	---
4	Alzheimer's disease (G30)	353	6.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	218	3.7	---
7	Influenza and pneumonia (J09-J18)	193	3.3	---
8	Diabetes mellitus (E10-E14)	172	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	2.5	---
10	Septicemia (A40-A41)	134	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	1.4	---
12	Pneumonitis due to solids and liquids (J69)	78	1.3	---
13	Parkinson's disease (G20-G21)	46	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	---
15	Nutritional deficiencies (E40-E64)	25	0.4	---
...	All other causes (residual)	1,190	20.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,918	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	562	29.3	---
2	Malignant neoplasms (C00-C97)	232	12.1	---
3	Alzheimer's disease (G30)	150	7.8	---
4	Cerebrovascular diseases (I60-I69)	112	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	77	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	3.6	---
7	Influenza and pneumonia (J09-J18)	55	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	2.1	---
10	Diabetes mellitus (E10-E14)	39	2.0	---
11	Septicemia (A40-A41)	33	1.7	---
12	Pneumonitis due to solids and liquids (J69)	24	1.3	---
13	Parkinson's disease (G20-G21)	20	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	---
14	Nutritional deficiencies (E40-E64)	14	0.7	---
...	All other causes (residual)	434	22.6	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	383	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	29.5	---
2	Malignant neoplasms (C00-C97)	33	8.6	---
3	Alzheimer's disease (G30)	30	7.8	---
4	Cerebrovascular diseases (I60-I69)	21	5.5	---
5	Influenza and pneumonia (J09-J18)	16	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	14	3.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	3.1	---
9	Septicemia (A40-A41)	9	2.3	---
9	Diabetes mellitus (E10-E14)	9	2.3	---
11	Nutritional deficiencies (E40-E64)	5	1.3	---
11	Pneumonitis due to solids and liquids (J69)	5	1.3	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.0	---
14	Atherosclerosis (I70)	2	0.5	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.5	---
...	All other causes (residual)	93	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	153,404	100.0	704.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,455	23.1	162.8
2	Malignant neoplasms (C00-C97)	33,736	22.0	154.9
3	Cerebrovascular diseases (I60-I69)	9,798	6.4	45.0
4	Diabetes mellitus (E10-E14)	6,887	4.5	31.6
5	Alzheimer's disease (G30)	5,918	3.9	27.2
6	Chronic lower respiratory diseases (J40-J47)	5,091	3.3	23.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,850	3.2	22.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,732	3.1	21.7
9	Septicemia (A40-A41)	3,515	2.3	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,141	2.0	14.4
11	Influenza and pneumonia (J09-J18)	2,825	1.8	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,697	1.1	7.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,252	0.8	5.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,149	0.7	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,129	0.7	5.2
...	All other causes (residual)	32,229	21.0	148.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	408	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	12.0	4.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.0	3.4
4	Malignant neoplasms (C00-C97)	27	6.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.9	2.0
6	Influenza and pneumonia (J09-J18)	10	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	2.0	*
9	Septicemia (A40-A41)	7	1.7	*
10	Anemias (D50-D64)	6	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Cerebrovascular diseases (I60-I69)	5	1.2	*
13	Meningitis (G00,G03)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	101	24.8	8.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	236	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	33.5	5.1
2	Malignant neoplasms (C00-C97)	27	11.4	1.8
3	Chronic lower respiratory diseases (J40-J47)	19	8.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	7.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	6.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	5.5	*
7	Anemias (D50-D64)	7	3.0	*
7	Influenza and pneumonia (J09-J18)	7	3.0	*
9	Septicemia (A40-A41)	6	2.5	*
10	Cerebrovascular diseases (I60-I69)	5	2.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
14	Pneumonitis due to solids and liquids (J69)	1	0.4	*
14	Diseases of appendix (K35-K38)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	30	12.7	2.0
Non-Hispanic, black, female, 10-14 years				
...	All causes	248	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	24.6	4.0
2	Malignant neoplasms (C00-C97)	31	12.5	2.1
3	Chronic lower respiratory diseases (J40-J47)	20	8.1	1.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	8.1	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	5.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.2	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.8	*
8	Cerebrovascular diseases (I60-I69)	4	1.6	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.6	*
10	Septicemia (A40-A41)	3	1.2	*
10	Influenza and pneumonia (J09-J18)	3	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Anemias (D50-D64)	2	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
15	Meningococcal infection (A39)	1	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Hernia (K40-K46)	1	0.4	*
...	All other causes (residual)	53	21.4	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	541	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	27.9	9.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	18.1	6.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	8.9	3.0
4	Malignant neoplasms (C00-C97)	41	7.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.4	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	17	3.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	3.1	*
8	Anemias (D50-D64)	16	3.0	*
9	Chronic lower respiratory diseases (J40-J47)	12	2.2	*
10	Septicemia (A40-A41)	7	1.3	*
11	Diabetes mellitus (E10-E14)	6	1.1	*
12	Influenza and pneumonia (J09-J18)	5	0.9	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	83	15.3	5.3
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,069	100.0	60.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	25.4	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	16.0	9.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	7.4	4.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	6.9	4.2
5	Malignant neoplasms (C00-C97)	73	6.8	4.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	46	4.3	2.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	27	2.5	1.5
8	Anemias (D50-D64)	19	1.8	*
8	Diabetes mellitus (E10-E14)	19	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	19	1.8	*
11	Cerebrovascular diseases (I60-I69)	15	1.4	*
12	Septicemia (A40-A41)	12	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	Influenza and pneumonia (J09-J18)	9	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
...	All other causes (residual)	217	20.3	12.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,326	100.0	80.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	21.8	17.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	12.4	10.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	10.4	8.4
4	Malignant neoplasms (C00-C97)	113	8.5	6.9
5	Pregnancy, childbirth and the puerperium (O00-O99)	64	4.8	3.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	4.2	3.4
7	Diabetes mellitus (E10-E14)	46	3.5	2.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	2.6	2.1
9	Cerebrovascular diseases (I60-I69)	24	1.8	1.5
10	Anemias (D50-D64)	23	1.7	1.4
11	Septicemia (A40-A41)	21	1.6	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.1	*
...	All other causes (residual)	289	21.8	17.6
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,761	100.0	118.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	14.9	17.7
2	Malignant neoplasms (C00-C97)	250	14.2	16.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	13.9	16.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	126	7.2	8.5
5	Diabetes mellitus (E10-E14)	75	4.3	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	71	4.0	4.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	69	3.9	4.6
8	Cerebrovascular diseases (I60-I69)	51	2.9	3.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	2.4	2.8
10	Anemias (D50-D64)	37	2.1	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.1	2.5
12	Septicemia (A40-A41)	34	1.9	2.3
13	Influenza and pneumonia (J09-J18)	21	1.2	1.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.1	*
...	All other causes (residual)	403	22.9	27.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,331	100.0	164.9
1	Malignant neoplasms (C00-C97)	451	19.3	31.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	15.9	26.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	12.2	20.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	116	5.0	8.2
5	Diabetes mellitus (E10-E14)	88	3.8	6.2
5	Cerebrovascular diseases (I60-I69)	88	3.8	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	3.8	6.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	66	2.8	4.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.1	3.5
10	Chronic lower respiratory diseases (J40-J47)	40	1.7	2.8
11	Septicemia (A40-A41)	38	1.6	2.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	1.6	2.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.2	2.1
14	Anemias (D50-D64)	27	1.2	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.2	1.9
...	All other causes (residual)	532	22.8	37.6
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,361	100.0	242.4
1	Malignant neoplasms (C00-C97)	760	22.6	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	649	19.3	46.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	9.0	21.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	144	4.3	10.4
5	Cerebrovascular diseases (I60-I69)	137	4.1	9.9
6	Diabetes mellitus (E10-E14)	135	4.0	9.7
7	Septicemia (A40-A41)	85	2.5	6.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	2.5	6.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	2.5	6.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	76	2.3	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.5	3.8
12	Chronic lower respiratory diseases (J40-J47)	46	1.4	3.3
13	Influenza and pneumonia (J09-J18)	42	1.2	3.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.2	2.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.0	2.5
...	All other causes (residual)	695	20.7	50.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,088	100.0	364.5
1	Malignant neoplasms (C00-C97)	1,250	24.6	89.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,054	20.7	75.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	7.2	26.1
4	Diabetes mellitus (E10-E14)	248	4.9	17.8
4	Cerebrovascular diseases (I60-I69)	248	4.9	17.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	170	3.3	12.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	127	2.5	9.1
8	Septicemia (A40-A41)	117	2.3	8.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.1	7.7
10	Chronic lower respiratory diseases (J40-J47)	96	1.9	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.4	5.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	1.4	5.1
13	Influenza and pneumonia (J09-J18)	50	1.0	3.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	0.9	3.3
15	Anemias (D50-D64)	39	0.8	2.8
...	All other causes (residual)	1,029	20.2	73.7
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,272	100.0	563.6
1	Malignant neoplasms (C00-C97)	2,521	30.5	171.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,703	20.6	116.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	5.5	30.9
4	Cerebrovascular diseases (I60-I69)	391	4.7	26.6
5	Diabetes mellitus (E10-E14)	380	4.6	25.9
6	Chronic lower respiratory diseases (J40-J47)	223	2.7	15.2
7	Septicemia (A40-A41)	200	2.4	13.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	199	2.4	13.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	176	2.1	12.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.1	11.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	139	1.7	9.5
12	Influenza and pneumonia (J09-J18)	115	1.4	7.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.7	4.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	0.5	2.9
15	Viral hepatitis (B15-B19)	42	0.5	2.9
...	All other causes (residual)	1,455	17.6	99.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	11,835	100.0	847.8
1	Malignant neoplasms (C00-C97)	3,890	32.9	278.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,562	21.6	183.5
3	Diabetes mellitus (E10-E14)	604	5.1	43.3
4	Cerebrovascular diseases (I60-I69)	546	4.6	39.1
5	Chronic lower respiratory diseases (J40-J47)	459	3.9	32.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	3.3	28.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	2.7	23.1
8	Septicemia (A40-A41)	309	2.6	22.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	235	2.0	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.6	13.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	163	1.4	11.7
12	Influenza and pneumonia (J09-J18)	151	1.3	10.8
13	Viral hepatitis (B15-B19)	83	0.7	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	0.4	3.6
15	Pneumonitis due to solids and liquids (J69)	47	0.4	3.4
...	All other causes (residual)	1,837	15.5	131.6
Non-Hispanic, black, female, 60-64 years				
...	All causes	13,608	100.0	1,179.6
1	Malignant neoplasms (C00-C97)	4,434	32.6	384.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,055	22.5	264.8
3	Diabetes mellitus (E10-E14)	730	5.4	63.3
4	Cerebrovascular diseases (I60-I69)	722	5.3	62.6
5	Chronic lower respiratory diseases (J40-J47)	506	3.7	43.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	3.0	35.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	2.4	28.5
8	Septicemia (A40-A41)	323	2.4	28.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	2.0	23.7
10	Influenza and pneumonia (J09-J18)	205	1.5	17.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	196	1.4	17.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	108	0.8	9.4
13	Viral hepatitis (B15-B19)	91	0.7	7.9
14	Pneumonitis due to solids and liquids (J69)	53	0.4	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.4	4.5
...	All other causes (residual)	2,119	15.6	183.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	14,103	100.0	1,592.8
1	Malignant neoplasms (C00-C97)	4,536	32.2	512.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,145	22.3	355.2
3	Cerebrovascular diseases (I60-I69)	777	5.5	87.8
4	Diabetes mellitus (E10-E14)	744	5.3	84.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	550	3.9	62.1
6	Chronic lower respiratory diseases (J40-J47)	549	3.9	62.0
7	Septicemia (A40-A41)	356	2.5	40.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	275	1.9	31.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	1.9	29.5
10	Influenza and pneumonia (J09-J18)	258	1.8	29.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	126	0.9	14.2
12	Alzheimer's disease (G30)	91	0.6	10.3
13	Pneumonitis due to solids and liquids (J69)	77	0.5	8.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	74	0.5	8.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	7.8
...	All other causes (residual)	2,215	15.7	250.2
Non-Hispanic, black, female, 70-74 years				
...	All causes	14,348	100.0	2,377.6
1	Malignant neoplasms (C00-C97)	4,116	28.7	682.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,466	24.2	574.3
3	Cerebrovascular diseases (I60-I69)	898	6.3	148.8
4	Diabetes mellitus (E10-E14)	803	5.6	133.1
5	Chronic lower respiratory diseases (J40-J47)	645	4.5	106.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	499	3.5	82.7
7	Septicemia (A40-A41)	354	2.5	58.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	291	2.0	48.2
9	Alzheimer's disease (G30)	279	1.9	46.2
10	Influenza and pneumonia (J09-J18)	258	1.8	42.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	1.3	31.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.6	14.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	12.3
14	Pneumonitis due to solids and liquids (J69)	71	0.5	11.8
15	Parkinson's disease (G20-G21)	52	0.4	8.6
...	All other causes (residual)	2,264	15.8	375.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	15,521	100.0	3,580.2
1	Malignant neoplasms (C00-C97)	3,799	24.5	876.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,692	23.8	851.6
3	Cerebrovascular diseases (I60-I69)	1,152	7.4	265.7
4	Diabetes mellitus (E10-E14)	865	5.6	199.5
5	Chronic lower respiratory diseases (J40-J47)	648	4.2	149.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	602	3.9	138.9
7	Alzheimer's disease (G30)	570	3.7	131.5
8	Septicemia (A40-A41)	404	2.6	93.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	2.2	78.9
10	Influenza and pneumonia (J09-J18)	287	1.8	66.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	1.2	44.5
12	Pneumonitis due to solids and liquids (J69)	95	0.6	21.9
13	Parkinson's disease (G20-G21)	91	0.6	21.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.6	20.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.4	13.4
...	All other causes (residual)	2,636	17.0	608.0
Non-Hispanic, black, female, 80-84 years				
...	All causes	17,000	100.0	5,607.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,223	24.8	1,393.0
2	Malignant neoplasms (C00-C97)	3,178	18.7	1,048.3
3	Cerebrovascular diseases (I60-I69)	1,452	8.5	478.9
4	Alzheimer's disease (G30)	1,034	6.1	341.1
5	Diabetes mellitus (E10-E14)	809	4.8	266.9
6	Chronic lower respiratory diseases (J40-J47)	690	4.1	227.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	621	3.7	204.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	2.5	139.9
9	Septicemia (A40-A41)	417	2.5	137.5
10	Influenza and pneumonia (J09-J18)	360	2.1	118.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	1.3	73.9
12	Pneumonitis due to solids and liquids (J69)	112	0.7	36.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.6	36.3
14	Parkinson's disease (G20-G21)	103	0.6	34.0
15	Aortic aneurysm and dissection (I71)	59	0.3	19.5
...	All other causes (residual)	3,184	18.7	1,050.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	17,316	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,684	27.1	---
2	Malignant neoplasms (C00-C97)	2,428	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,499	8.7	---
4	Alzheimer's disease (G30)	1,439	8.3	---
5	Diabetes mellitus (E10-E14)	715	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	576	3.3	---
7	Chronic lower respiratory diseases (J40-J47)	571	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	393	2.3	---
9	Septicemia (A40-A41)	392	2.3	---
10	Influenza and pneumonia (J09-J18)	378	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	1.2	---
12	Pneumonitis due to solids and liquids (J69)	118	0.7	---
13	Parkinson's disease (G20-G21)	113	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	---
15	Atherosclerosis (I70)	60	0.3	---
...	All other causes (residual)	3,644	21.0	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	13,305	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,761	28.3	---
2	Alzheimer's disease (G30)	1,414	10.6	---
3	Malignant neoplasms (C00-C97)	1,262	9.5	---
4	Cerebrovascular diseases (I60-I69)	1,100	8.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	3.1	---
6	Diabetes mellitus (E10-E14)	414	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	380	2.9	---
8	Influenza and pneumonia (J09-J18)	373	2.8	---
9	Chronic lower respiratory diseases (J40-J47)	334	2.5	---
10	Septicemia (A40-A41)	282	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.1	---
12	Pneumonitis due to solids and liquids (J69)	114	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	---
14	Parkinson's disease (G20-G21)	65	0.5	---
15	Nutritional deficiencies (E40-E64)	59	0.4	---
...	All other causes (residual)	3,109	23.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	6,415	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,857	28.9	---
2	Alzheimer's disease (G30)	740	11.5	---
3	Cerebrovascular diseases (I60-I69)	512	8.0	---
4	Malignant neoplasms (C00-C97)	450	7.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	2.9	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	2.7	---
7	Influenza and pneumonia (J09-J18)	164	2.6	---
8	Diabetes mellitus (E10-E14)	162	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	131	2.0	---
10	Septicemia (A40-A41)	88	1.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.2	---
12	Pneumonitis due to solids and liquids (J69)	58	0.9	---
13	Nutritional deficiencies (E40-E64)	40	0.6	---
14	Atherosclerosis (I70)	32	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	---
...	All other causes (residual)	1,716	26.7	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,215	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	649	29.3	---
2	Alzheimer's disease (G30)	284	12.8	---
3	Cerebrovascular diseases (I60-I69)	155	7.0	---
4	Malignant neoplasms (C00-C97)	95	4.3	---
5	Influenza and pneumonia (J09-J18)	79	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.8	---
8	Diabetes mellitus (E10-E14)	39	1.8	---
9	Septicemia (A40-A41)	35	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.1	---
12	Pneumonitis due to solids and liquids (J69)	23	1.0	---
13	Nutritional deficiencies (E40-E64)	21	0.9	---
14	Atherosclerosis (I70)	12	0.5	---
15	Anemias (D50-D64)	8	0.4	---
...	All other causes (residual)	616	27.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	18,039	100.0	670.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,303	18.3	122.8
2	Malignant neoplasms (C00-C97)	3,192	17.7	118.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,953	10.8	72.6
4	Diabetes mellitus (E10-E14)	1,034	5.7	38.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,006	5.6	37.4
6	Chronic lower respiratory diseases (J40-J47)	859	4.8	31.9
7	Cerebrovascular diseases (I60-I69)	646	3.6	24.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	535	3.0	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	340	1.9	12.6
10	Influenza and pneumonia (J09-J18)	330	1.8	12.3
11	Alzheimer's disease (G30)	307	1.7	11.4
12	Septicemia (A40-A41)	296	1.6	11.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	1.5	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.1	7.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	0.6	4.1
...	All other causes (residual)	3,675	20.4	136.6
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	82	100.0	51.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	36.6	18.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	12.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	9.8	*
4	Malignant neoplasms (C00-C97)	3	3.7	*
5	Meningitis (G00,G03)	1	1.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.2	*
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.2	*
5	Cerebrovascular diseases (I60-I69)	1	1.2	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.2	*
5	Pneumonitis due to solids and liquids (J69)	1	1.2	*
...	All other causes (residual)	25	30.5	15.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-9 years				
...	All causes	47	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	25.5	*
2	Malignant neoplasms (C00-C97)	8	17.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	12.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.5	*
5	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (residual)	14	29.8	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 10-14 years				
...	All causes	52	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	36.5	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	26.9	*
3	Malignant neoplasms (C00-C97)	3	5.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.9	*
6	Influenza and pneumonia (J09-J18)	1	1.9	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.9	*
...	All other causes (residual)	6	11.5	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-19 years				
...	All causes	174	100.0	82.4
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	37.9	31.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	33.9	27.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	8.0	*
4	Malignant neoplasms (C00-C97)	5	2.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.3	*
6	Chronic lower respiratory diseases (J40-J47)	3	1.7	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
9	Cerebrovascular diseases (I60-I69)	1	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (residual)	16	9.2	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 20-24 years				
...	All causes	351	100.0	153.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	41.6	64.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	23.9	36.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	10.8	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.4	*
5	Malignant neoplasms (C00-C97)	11	3.1	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.4	*
8	Cerebrovascular diseases (I60-I69)	3	0.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
9	Diabetes mellitus (E10-E14)	2	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Meningitis (G00,G03)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Influenza and pneumonia (J09-J18)	1	0.3	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.3	*
12	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (residual)	34	9.7	14.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-29 years				
...	All causes	454	100.0	219.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	44.5	97.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	15.2	33.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.3	16.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	5.9	13.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	25	5.5	12.1
6	Malignant neoplasms (C00-C97)	16	3.5	*
7	Septicemia (A40-A41)	4	0.9	*
7	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
10	Diabetes mellitus (E10-E14)	3	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Cerebrovascular diseases (I60-I69)	1	0.2	*
14	Influenza and pneumonia (J09-J18)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	53	11.7	25.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 30-34 years				
...	All causes	543	100.0	296.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	38.7	114.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	58	10.7	31.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	10.5	31.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	19.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.2	15.3
6	Malignant neoplasms (C00-C97)	25	4.6	13.6
7	Septicemia (A40-A41)	7	1.3	*
7	Diabetes mellitus (E10-E14)	7	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
9	Cerebrovascular diseases (I60-I69)	6	1.1	*
11	Influenza and pneumonia (J09-J18)	4	0.7	*
12	Viral hepatitis (B15-B19)	3	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (residual)	85	15.7	46.3
Non-Hispanic, American Indian or Alaska Native, both sexes, 35-39 years				
...	All causes	589	100.0	354.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	30.2	107.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	67	11.4	40.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	8.0	28.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	8.0	28.3
5	Malignant neoplasms (C00-C97)	44	7.5	26.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.1	18.0
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
8	Diabetes mellitus (E10-E14)	9	1.5	*
9	Septicemia (A40-A41)	7	1.2	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
13	Influenza and pneumonia (J09-J18)	3	0.5	*
13	Legal intervention (Y35,Y89.0)	3	0.5	*
...	All other causes (residual)	126	21.4	75.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 40-44 years				
...	All causes	688	100.0	430.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	22.4	96.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	113	16.4	70.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	13.4	57.5
4	Malignant neoplasms (C00-C97)	46	6.7	28.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	6.1	26.3
6	Diabetes mellitus (E10-E14)	38	5.5	23.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	3.3	14.4
8	Septicemia (A40-A41)	13	1.9	*
8	Cerebrovascular diseases (I60-I69)	13	1.9	*
10	Influenza and pneumonia (J09-J18)	10	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.5	*
12	Viral hepatitis (B15-B19)	5	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.7	*
...	All other causes (residual)	114	16.6	71.3
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-49 years				
...	All causes	997	100.0	609.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	19.9	121.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	156	15.6	95.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	13.1	80.1
4	Malignant neoplasms (C00-C97)	95	9.5	58.1
5	Diabetes mellitus (E10-E14)	54	5.4	33.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	4.3	26.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	2.4	14.7
8	Cerebrovascular diseases (I60-I69)	23	2.3	14.1
9	Septicemia (A40-A41)	18	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	15	1.5	*
11	Influenza and pneumonia (J09-J18)	12	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.1	*
14	Viral hepatitis (B15-B19)	10	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.5	*
...	All other causes (residual)	190	19.1	116.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 50-54 years				
...	All causes	1,368	100.0	760.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	18.4	140.0
2	Malignant neoplasms (C00-C97)	250	18.3	138.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	12.4	93.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	154	11.3	85.6
5	Diabetes mellitus (E10-E14)	88	6.4	48.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.2	24.4
7	Cerebrovascular diseases (I60-I69)	32	2.3	17.8
8	Chronic lower respiratory diseases (J40-J47)	27	2.0	15.0
9	Septicemia (A40-A41)	25	1.8	13.9
10	Viral hepatitis (B15-B19)	20	1.5	11.1
11	Influenza and pneumonia (J09-J18)	18	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.0	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
14	Pneumonitis due to solids and liquids (J69)	10	0.7	*
...	All other causes (residual)	242	17.7	134.5
Non-Hispanic, American Indian or Alaska Native, both sexes, 55-59 years				
...	All causes	1,628	100.0	956.6
1	Malignant neoplasms (C00-C97)	352	21.6	206.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	19.2	183.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	161	9.9	94.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	9.3	88.7
5	Diabetes mellitus (E10-E14)	96	5.9	56.4
6	Chronic lower respiratory diseases (J40-J47)	66	4.1	38.8
7	Cerebrovascular diseases (I60-I69)	45	2.8	26.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.8	17.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.7	16.5
10	Septicemia (A40-A41)	24	1.5	14.1
11	Influenza and pneumonia (J09-J18)	23	1.4	13.5
12	Viral hepatitis (B15-B19)	17	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.9	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.9	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.4	*
...	All other causes (residual)	286	17.6	168.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 60-64 years				
...	All causes	1,700	100.0	1,235.8
1	Malignant neoplasms (C00-C97)	397	23.4	288.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	358	21.1	260.3
3	Diabetes mellitus (E10-E14)	146	8.6	106.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	7.0	86.5
5	Chronic lower respiratory diseases (J40-J47)	104	6.1	75.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	103	6.1	74.9
7	Cerebrovascular diseases (I60-I69)	48	2.8	34.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.3	28.4
9	Septicemia (A40-A41)	37	2.2	26.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.5	18.9
11	Viral hepatitis (B15-B19)	22	1.3	16.0
11	Influenza and pneumonia (J09-J18)	22	1.3	16.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.9	*
14	Pneumonitis due to solids and liquids (J69)	12	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.4	*
...	All other causes (residual)	238	14.0	173.0
Non-Hispanic, American Indian or Alaska Native, both sexes, 65-69 years				
...	All causes	1,727	100.0	1,627.1
1	Malignant neoplasms (C00-C97)	455	26.3	428.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	355	20.6	334.5
3	Diabetes mellitus (E10-E14)	155	9.0	146.0
4	Chronic lower respiratory diseases (J40-J47)	107	6.2	100.8
5	Cerebrovascular diseases (I60-I69)	68	3.9	64.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	65	3.8	61.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	3.5	56.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.6	42.4
9	Septicemia (A40-A41)	37	2.1	34.9
10	Influenza and pneumonia (J09-J18)	30	1.7	28.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.2	18.8
12	Viral hepatitis (B15-B19)	11	0.6	*
13	Pneumonitis due to solids and liquids (J69)	8	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
14	Peptic ulcer (K25-K28)	7	0.4	*
...	All other causes (residual)	297	17.2	279.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 70-74 years				
...	All causes	1,771	100.0	2,578.8
1	Malignant neoplasms (C00-C97)	472	26.7	687.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	353	19.9	514.0
3	Chronic lower respiratory diseases (J40-J47)	152	8.6	221.3
4	Diabetes mellitus (E10-E14)	144	8.1	209.7
5	Cerebrovascular diseases (I60-I69)	80	4.5	116.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	3.0	77.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	47	2.7	68.4
8	Influenza and pneumonia (J09-J18)	46	2.6	67.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.4	62.6
10	Septicemia (A40-A41)	38	2.1	55.3
11	Alzheimer's disease (G30)	19	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.7	*
13	Parkinson's disease (G20-G21)	10	0.6	*
14	Aortic aneurysm and dissection (I71)	9	0.5	*
14	Pneumonitis due to solids and liquids (J69)	9	0.5	*
...	All other causes (residual)	283	16.0	412.1
Non-Hispanic, American Indian or Alaska Native, both sexes, 75-79 years				
...	All causes	1,692	100.0	3,794.0
1	Malignant neoplasms (C00-C97)	427	25.2	957.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	371	21.9	831.9
3	Chronic lower respiratory diseases (J40-J47)	136	8.0	305.0
4	Diabetes mellitus (E10-E14)	115	6.8	257.9
5	Cerebrovascular diseases (I60-I69)	75	4.4	168.2
6	Alzheimer's disease (G30)	50	3.0	112.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	2.6	98.7
8	Influenza and pneumonia (J09-J18)	42	2.5	94.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.5	94.2
10	Septicemia (A40-A41)	35	2.1	78.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.5	58.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.0	*
13	Parkinson's disease (G20-G21)	15	0.9	*
14	Pneumonitis due to solids and liquids (J69)	10	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
...	All other causes (residual)	278	16.4	623.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 80-84 years				
...	All causes	1,532	100.0	5,770.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	377	24.6	1,419.9
2	Malignant neoplasms (C00-C97)	278	18.1	1,047.0
3	Chronic lower respiratory diseases (J40-J47)	101	6.6	380.4
4	Diabetes mellitus (E10-E14)	91	5.9	342.7
5	Cerebrovascular diseases (I60-I69)	86	5.6	323.9
6	Alzheimer's disease (G30)	70	4.6	263.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.7	154.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	2.6	150.7
9	Influenza and pneumonia (J09-J18)	35	2.3	131.8
10	Parkinson's disease (G20-G21)	28	1.8	105.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	1.8	105.5
12	Septicemia (A40-A41)	23	1.5	86.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.8	*
14	Pneumonitis due to solids and liquids (J69)	12	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
...	All other causes (residual)	300	19.6	1,129.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 85-89 years				
...	All causes	1,216	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279	22.9	---
2	Malignant neoplasms (C00-C97)	209	17.2	---
3	Chronic lower respiratory diseases (J40-J47)	85	7.0	---
4	Cerebrovascular diseases (I60-I69)	77	6.3	---
5	Alzheimer's disease (G30)	66	5.4	---
6	Diabetes mellitus (E10-E14)	46	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.8	---
9	Influenza and pneumonia (J09-J18)	31	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	2.0	---
11	Parkinson's disease (G20-G21)	18	1.5	---
12	Septicemia (A40-A41)	14	1.2	---
13	Pneumonitis due to solids and liquids (J69)	12	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.7	---
...	All other causes (residual)	265	21.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 90-94 years				
...	All causes	803	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	26.8	---
2	Malignant neoplasms (C00-C97)	79	9.8	---
3	Alzheimer's disease (G30)	70	8.7	---
4	Cerebrovascular diseases (I60-I69)	53	6.6	---
5	Diabetes mellitus (E10-E14)	32	4.0	---
5	Chronic lower respiratory diseases (J40-J47)	32	4.0	---
7	Influenza and pneumonia (J09-J18)	30	3.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.9	---
11	Nutritional deficiencies (E40-E64)	9	1.1	---
11	Parkinson's disease (G20-G21)	9	1.1	---
13	Septicemia (A40-A41)	8	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	---
14	Pneumonitis due to solids and liquids (J69)	6	0.7	---
...	All other causes (residual)	191	23.8	---
Non-Hispanic, American Indian or Alaska Native, both sexes, 95-99 years				
...	All causes	257	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	24.5	---
2	Alzheimer's disease (G30)	20	7.8	---
3	Cerebrovascular diseases (I60-I69)	15	5.8	---
4	Malignant neoplasms (C00-C97)	14	5.4	---
4	Influenza and pneumonia (J09-J18)	14	5.4	---
6	Chronic lower respiratory diseases (J40-J47)	8	3.1	---
7	Diabetes mellitus (E10-E14)	7	2.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	2.7	---
9	Pneumonitis due to solids and liquids (J69)	6	2.3	---
10	Septicemia (A40-A41)	4	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.6	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.2	---
14	Parkinson's disease (G20-G21)	2	0.8	---
14	Atherosclerosis (I70)	2	0.8	---
...	All other causes (residual)	85	33.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 100 years and over				
...	All causes	68	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	32.4	---
2	Cerebrovascular diseases (I60-I69)	6	8.8	---
3	Alzheimer's disease (G30)	4	5.9	---
3	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	5.9	---
3	Chronic lower respiratory diseases (J40-J47)	4	5.9	---
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	5.9	---
7	Influenza and pneumonia (J09-J18)	3	4.4	---
8	Malignant neoplasms (C00-C97)	2	2.9	---
9	Anemias (D50-D64)	1	1.5	---
9	Diabetes mellitus (E10-E14)	1	1.5	---
9	Pneumonitis due to solids and liquids (J69)	1	1.5	---
9	Hernia (K40-K46)	1	1.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	---
...	All other causes (residual)	14	20.6	---
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	9,869	100.0	747.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,916	19.4	145.1
2	Malignant neoplasms (C00-C97)	1,699	17.2	128.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,282	13.0	97.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	563	5.7	42.6
5	Diabetes mellitus (E10-E14)	537	5.4	40.7
6	Chronic lower respiratory diseases (J40-J47)	398	4.0	30.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	4.0	29.9
8	Cerebrovascular diseases (I60-I69)	291	2.9	22.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.1	15.8
10	Influenza and pneumonia (J09-J18)	181	1.8	13.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	1.6	12.0
12	Septicemia (A40-A41)	130	1.3	9.8
13	Alzheimer's disease (G30)	100	1.0	7.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	0.9	6.6
15	Viral hepatitis (B15-B19)	70	0.7	5.3
...	All other causes (residual)	1,852	18.8	140.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	45	100.0	55.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	31.1	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.9	*
3	Malignant neoplasms (C00-C97)	3	6.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.2	*
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	2.2	*
5	Cerebrovascular diseases (I60-I69)	1	2.2	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (residual)	17	37.8	*
Non-Hispanic, American Indian or Alaska Native, male, 5-9 years				
...	All causes	28	100.0	26.9
1	Malignant neoplasms (C00-C97)	6	21.4	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	21.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	7.1	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	7.1	*
5	Chronic lower respiratory diseases (J40-J47)	1	3.6	*
...	All other causes (residual)	11	39.3	*
Non-Hispanic, American Indian or Alaska Native, male, 10-14 years				
...	All causes	23	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	52.2	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	17.4	*
3	Malignant neoplasms (C00-C97)	2	8.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	8.7	*
5	Influenza and pneumonia (J09-J18)	1	4.3	*
...	All other causes (residual)	2	8.7	*
Non-Hispanic, American Indian or Alaska Native, male, 15-19 years				
...	All causes	112	100.0	104.5
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	37.5	39.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	34.8	36.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	12.5	*
4	Malignant neoplasms (C00-C97)	5	4.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
6	Cerebrovascular diseases (I60-I69)	1	0.9	*
6	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
...	All other causes (residual)	7	6.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 20-24 years				
...	All causes	250	100.0	214.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	39.6	85.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	27.2	58.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	14.0	30.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.8	*
5	Malignant neoplasms (C00-C97)	5	2.0	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5	2.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.2	*
8	Diabetes mellitus (E10-E14)	2	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
9	Cerebrovascular diseases (I60-I69)	1	0.4	*
9	Influenza and pneumonia (J09-J18)	1	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
9	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (residual)	21	8.4	18.0
Non-Hispanic, American Indian or Alaska Native, male, 25-29 years				
...	All causes	317	100.0	303.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	45.7	138.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	17.0	51.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	8.5	25.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	6.0	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	13	4.1	*
6	Malignant neoplasms (C00-C97)	7	2.2	*
7	Septicemia (A40-A41)	3	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
9	Tuberculosis (A16-A19)	1	0.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
9	Diabetes mellitus (E10-E14)	1	0.3	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
9	Cerebrovascular diseases (I60-I69)	1	0.3	*
9	Influenza and pneumonia (J09-J18)	1	0.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
9	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (residual)	38	12.0	36.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 30-34 years				
...	All causes	345	100.0	380.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	39.7	151.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	13.0	49.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	39	11.3	43.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	7.2	27.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.9	*
6	Malignant neoplasms (C00-C97)	13	3.8	*
7	Viral hepatitis (B15-B19)	3	0.9	*
8	Septicemia (A40-A41)	2	0.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
8	Diabetes mellitus (E10-E14)	2	0.6	*
8	Cerebrovascular diseases (I60-I69)	2	0.6	*
8	Influenza and pneumonia (J09-J18)	2	0.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
8	Legal intervention (Y35,Y89.0)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Aortic aneurysm and dissection (I71)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	47	13.6	51.8
Non-Hispanic, American Indian or Alaska Native, male, 35-39 years				
...	All causes	370	100.0	450.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	35.1	158.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	9.7	43.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	30	8.1	36.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	6.8	30.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	6.5	29.2
6	Malignant neoplasms (C00-C97)	20	5.4	24.4
7	Cerebrovascular diseases (I60-I69)	7	1.9	*
8	Diabetes mellitus (E10-E14)	6	1.6	*
9	Septicemia (A40-A41)	5	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.1	*
11	Legal intervention (Y35,Y89.0)	3	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
12	Influenza and pneumonia (J09-J18)	2	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (residual)	70	18.9	85.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 40-44 years				
...	All causes	439	100.0	560.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	24.8	139.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	69	15.7	88.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	13.9	77.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	6.6	37.0
5	Diabetes mellitus (E10-E14)	25	5.7	31.9
6	Malignant neoplasms (C00-C97)	24	5.5	30.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.3	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	Septicemia (A40-A41)	7	1.6	*
9	Influenza and pneumonia (J09-J18)	7	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
12	Viral hepatitis (B15-B19)	3	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
...	All other causes (residual)	65	14.8	83.0
Non-Hispanic, American Indian or Alaska Native, male, 45-49 years				
...	All causes	609	100.0	766.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	21.2	162.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	15.4	118.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	80	13.1	100.6
4	Malignant neoplasms (C00-C97)	45	7.4	56.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	5.4	41.5
6	Diabetes mellitus (E10-E14)	29	4.8	36.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	3.3	25.2
8	Septicemia (A40-A41)	11	1.8	*
9	Cerebrovascular diseases (I60-I69)	9	1.5	*
10	Viral hepatitis (B15-B19)	8	1.3	*
11	Influenza and pneumonia (J09-J18)	7	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
...	All other causes (residual)	123	20.2	154.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 50-54 years				
...	All causes	812	100.0	940.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	21.3	200.4
2	Malignant neoplasms (C00-C97)	137	16.9	158.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	12.2	114.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	87	10.7	100.8
5	Diabetes mellitus (E10-E14)	47	5.8	54.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	4.1	38.2
7	Viral hepatitis (B15-B19)	16	2.0	*
7	Cerebrovascular diseases (I60-I69)	16	2.0	*
9	Septicemia (A40-A41)	13	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.6	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	1.5	*
12	Influenza and pneumonia (J09-J18)	10	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.9	*
15	Pneumonitis due to solids and liquids (J69)	6	0.7	*
...	All other causes (residual)	135	16.6	156.4
Non-Hispanic, American Indian or Alaska Native, male, 55-59 years				
...	All causes	995	100.0	1,242.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	22.0	273.5
2	Malignant neoplasms (C00-C97)	201	20.2	251.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	10.5	129.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	100	10.1	124.9
5	Diabetes mellitus (E10-E14)	56	5.6	69.9
6	Chronic lower respiratory diseases (J40-J47)	27	2.7	33.7
7	Cerebrovascular diseases (I60-I69)	26	2.6	32.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.4	*
10	Influenza and pneumonia (J09-J18)	13	1.3	*
11	Viral hepatitis (B15-B19)	12	1.2	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	1.1	*
13	Septicemia (A40-A41)	10	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (residual)	166	16.7	207.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 60-64 years				
...	All causes	987	100.0	1,532.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251	25.4	389.7
2	Malignant neoplasms (C00-C97)	204	20.7	316.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	9.0	138.2
4	Diabetes mellitus (E10-E14)	85	8.6	132.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	60	6.1	93.2
6	Chronic lower respiratory diseases (J40-J47)	45	4.6	69.9
7	Cerebrovascular diseases (I60-I69)	24	2.4	37.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.4	37.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.6	*
10	Septicemia (A40-A41)	13	1.3	*
11	Viral hepatitis (B15-B19)	12	1.2	*
11	Influenza and pneumonia (J09-J18)	12	1.2	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.2	*
14	Pneumonitis due to solids and liquids (J69)	8	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.4	*
...	All other causes (residual)	128	13.0	198.7
Non-Hispanic, American Indian or Alaska Native, male, 65-69 years				
...	All causes	946	100.0	1,891.8
1	Malignant neoplasms (C00-C97)	237	25.1	473.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	23.5	443.9
3	Diabetes mellitus (E10-E14)	81	8.6	162.0
4	Chronic lower respiratory diseases (J40-J47)	50	5.3	100.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	5.0	94.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	36	3.8	72.0
7	Cerebrovascular diseases (I60-I69)	34	3.6	68.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.2	42.0
9	Septicemia (A40-A41)	20	2.1	40.0
10	Influenza and pneumonia (J09-J18)	16	1.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
12	Pneumonitis due to solids and liquids (J69)	6	0.6	*
13	Viral hepatitis (B15-B19)	5	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
14	Alzheimer's disease (G30)	4	0.4	*
14	Aortic aneurysm and dissection (I71)	4	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.4	*
...	All other causes (residual)	142	15.0	284.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 70-74 years				
...	All causes	962	100.0	3,032.6
1	Malignant neoplasms (C00-C97)	283	29.4	892.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	20.9	633.6
3	Chronic lower respiratory diseases (J40-J47)	78	8.1	245.9
4	Diabetes mellitus (E10-E14)	75	7.8	236.4
5	Cerebrovascular diseases (I60-I69)	45	4.7	141.9
6	Influenza and pneumonia (J09-J18)	30	3.1	94.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	3.1	94.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.2	66.2
9	Septicemia (A40-A41)	18	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.8	*
12	Alzheimer's disease (G30)	7	0.7	*
12	Aortic aneurysm and dissection (I71)	7	0.7	*
12	Pneumonitis due to solids and liquids (J69)	7	0.7	*
15	Parkinson's disease (G20-G21)	6	0.6	*
...	All other causes (residual)	130	13.5	409.8
Non-Hispanic, American Indian or Alaska Native, male, 75-79 years				
...	All causes	888	100.0	4,501.4
1	Malignant neoplasms (C00-C97)	237	26.7	1,201.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	24.1	1,084.8
3	Chronic lower respiratory diseases (J40-J47)	75	8.4	380.2
4	Diabetes mellitus (E10-E14)	54	6.1	273.7
5	Cerebrovascular diseases (I60-I69)	38	4.3	192.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	2.9	131.8
7	Influenza and pneumonia (J09-J18)	22	2.5	111.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.5	111.5
9	Alzheimer's disease (G30)	21	2.4	106.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.6	*
11	Septicemia (A40-A41)	11	1.2	*
12	Parkinson's disease (G20-G21)	7	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
15	Aortic aneurysm and dissection (I71)	5	0.6	*
15	Pneumonitis due to solids and liquids (J69)	5	0.6	*
...	All other causes (residual)	118	13.3	598.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 80-84 years				
...	All causes	705	100.0	6,503.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	26.2	1,706.5
2	Malignant neoplasms (C00-C97)	136	19.3	1,254.5
3	Diabetes mellitus (E10-E14)	44	6.2	405.9
4	Chronic lower respiratory diseases (J40-J47)	38	5.4	350.5
5	Cerebrovascular diseases (I60-I69)	32	4.5	295.2
6	Alzheimer's disease (G30)	28	4.0	258.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.3	212.2
8	Influenza and pneumonia (J09-J18)	20	2.8	184.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.6	*
10	Parkinson's disease (G20-G21)	17	2.4	*
11	Septicemia (A40-A41)	9	1.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.6	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.6	*
...	All other causes (residual)	126	17.9	1,162.3
Non-Hispanic, American Indian or Alaska Native, male, 85-89 years				
...	All causes	525	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	23.4	---
2	Malignant neoplasms (C00-C97)	100	19.0	---
3	Chronic lower respiratory diseases (J40-J47)	38	7.2	---
4	Cerebrovascular diseases (I60-I69)	26	5.0	---
5	Alzheimer's disease (G30)	22	4.2	---
6	Influenza and pneumonia (J09-J18)	18	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	3.0	---
9	Parkinson's disease (G20-G21)	15	2.9	---
10	Diabetes mellitus (E10-E14)	14	2.7	---
11	Pneumonitis due to solids and liquids (J69)	9	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.1	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.8	---
15	Anemias (D50-D64)	3	0.6	---
15	Nutritional deficiencies (E40-E64)	3	0.6	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.6	---
...	All other causes (residual)	104	19.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 90-94 years				
...	All causes	279	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	30.1	---
2	Malignant neoplasms (C00-C97)	30	10.8	---
3	Chronic lower respiratory diseases (J40-J47)	17	6.1	---
4	Cerebrovascular diseases (I60-I69)	16	5.7	---
5	Diabetes mellitus (E10-E14)	13	4.7	---
6	Influenza and pneumonia (J09-J18)	12	4.3	---
7	Alzheimer's disease (G30)	11	3.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.8	---
11	Septicemia (A40-A41)	4	1.4	---
11	Pneumonitis due to solids and liquids (J69)	4	1.4	---
13	Parkinson's disease (G20-G21)	3	1.1	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	1.1	---
15	Viral hepatitis (B15-B19)	2	0.7	---
15	Nutritional deficiencies (E40-E64)	2	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.7	---
...	All other causes (residual)	55	19.7	---
Non-Hispanic, American Indian or Alaska Native, male, 95-99 years				
...	All causes	62	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	27.4	---
2	Alzheimer's disease (G30)	6	9.7	---
3	Influenza and pneumonia (J09-J18)	5	8.1	---
4	Malignant neoplasms (C00-C97)	3	4.8	---
4	Diabetes mellitus (E10-E14)	3	4.8	---
4	Cerebrovascular diseases (I60-I69)	3	4.8	---
4	Pneumonitis due to solids and liquids (J69)	3	4.8	---
8	Chronic lower respiratory diseases (J40-J47)	2	3.2	---
9	Septicemia (A40-A41)	1	1.6	---
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	---
9	Parkinson's disease (G20-G21)	1	1.6	---
9	Atherosclerosis (I70)	1	1.6	---
9	Pneumoconioses and chemical effects (J60-J66,J68)	1	1.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.6	---
...	All other causes (residual)	14	22.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 100 years and over				
...	All causes	9	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	33.3	---
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	22.2	---
3	Chronic lower respiratory diseases (J40-J47)	1	11.1	---
3	Pneumonitis due to solids and liquids (J69)	1	11.1	---
3	Hernia (K40-K46)	1	11.1	---
...	All other causes (residual)	1	11.1	---
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,170	100.0	596.7
1	Malignant neoplasms (C00-C97)	1,493	18.3	109.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,387	17.0	101.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	671	8.2	49.0
4	Diabetes mellitus (E10-E14)	497	6.1	36.3
5	Chronic lower respiratory diseases (J40-J47)	461	5.6	33.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	443	5.4	32.4
7	Cerebrovascular diseases (I60-I69)	355	4.3	25.9
8	Alzheimer's disease (G30)	207	2.5	15.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.2	13.2
10	Septicemia (A40-A41)	166	2.0	12.1
11	Influenza and pneumonia (J09-J18)	149	1.8	10.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	140	1.7	10.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.3	7.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.6	3.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.6	3.8
...	All other causes (residual)	1,810	22.2	132.2
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	37	100.0	47.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	43.2	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	16.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	13.5	*
4	Meningitis (G00,G03)	1	2.7	*
4	Pneumonitis due to solids and liquids (J69)	1	2.7	*
...	All other causes (residual)	8	21.6	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 5-9 years				
...	All causes	19	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	31.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	21.1	*
3	Malignant neoplasms (C00-C97)	2	10.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	10.5	*
5	Chronic lower respiratory diseases (J40-J47)	1	5.3	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	5.3	*
...	All other causes (residual)	3	15.8	*
Non-Hispanic, American Indian or Alaska Native, female, 10-14 years				
...	All causes	29	100.0	28.7
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	34.5	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	24.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	6.9	*
4	Malignant neoplasms (C00-C97)	1	3.4	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.4	*
4	Chronic lower respiratory diseases (J40-J47)	1	3.4	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	3.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	3.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	3.4	*
...	All other causes (residual)	4	13.8	*
Non-Hispanic, American Indian or Alaska Native, female, 15-19 years				
...	All causes	62	100.0	59.6
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	38.7	23.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	32.3	19.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.2	*
3	Chronic lower respiratory diseases (J40-J47)	2	3.2	*
3	Pregnancy, childbirth and the puerperium (O00-O99)	2	3.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.2	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	*
...	All other causes (residual)	9	14.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 20-24 years				
...	All causes	101	100.0	90.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	46.5	42.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	15.8	*
3	Malignant neoplasms (C00-C97)	6	5.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.0	*
6	Cerebrovascular diseases (I60-I69)	2	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.0	*
9	Meningitis (G00,G03)	1	1.0	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	1.0	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.0	*
...	All other causes (residual)	13	12.9	*
Non-Hispanic, American Indian or Alaska Native, female, 25-29 years				
...	All causes	137	100.0	133.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	41.6	55.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	10.9	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	12	8.8	*
4	Malignant neoplasms (C00-C97)	9	6.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.4	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.9	*
8	Diabetes mellitus (E10-E14)	2	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.5	*
10	Septicemia (A40-A41)	1	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.7	*
10	Anemias (D50-D64)	1	0.7	*
10	Pneumonitis due to solids and liquids (J69)	1	0.7	*
10	Hernia (K40-K46)	1	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	*
...	All other causes (residual)	15	10.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 30-34 years				
...	All causes	198	100.0	213.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	36.9	78.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	19	9.6	*
3	Malignant neoplasms (C00-C97)	12	6.1	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	6.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	5.1	*
7	Septicemia (A40-A41)	5	2.5	*
7	Diabetes mellitus (E10-E14)	5	2.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	2.0	*
9	Cerebrovascular diseases (I60-I69)	4	2.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.5	*
12	Influenza and pneumonia (J09-J18)	2	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
14	Meningitis (G00,G03)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (residual)	33	16.7	35.6
Non-Hispanic, American Indian or Alaska Native, female, 35-39 years				
...	All causes	219	100.0	260.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	21.9	57.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	37	16.9	43.9
3	Malignant neoplasms (C00-C97)	24	11.0	28.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	10.5	27.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	5.0	*
6	Cerebrovascular diseases (I60-I69)	5	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	5	2.3	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	2.3	*
9	Diabetes mellitus (E10-E14)	3	1.4	*
10	Septicemia (A40-A41)	2	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	*
13	Aortic aneurysm and dissection (I71)	1	0.5	*
13	Influenza and pneumonia (J09-J18)	1	0.5	*
13	Peptic ulcer (K25-K28)	1	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	*
...	All other causes (residual)	46	21.0	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 40-44 years				
...	All causes	249	100.0	305.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	18.1	55.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	44	17.7	53.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	12.4	38.0
4	Malignant neoplasms (C00-C97)	22	8.8	26.9
5	Diabetes mellitus (E10-E14)	13	5.2	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	5.2	*
7	Septicemia (A40-A41)	6	2.4	*
8	Cerebrovascular diseases (I60-I69)	5	2.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.0	*
10	Chronic lower respiratory diseases (J40-J47)	4	1.6	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.6	*
12	Influenza and pneumonia (J09-J18)	3	1.2	*
13	Viral hepatitis (B15-B19)	2	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.8	*
...	All other causes (residual)	46	18.5	56.3
Non-Hispanic, American Indian or Alaska Native, female, 45-49 years				
...	All causes	388	100.0	461.3
1	Chronic liver disease and cirrhosis (K70,K73-K74)	76	19.6	90.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	17.8	82.0
3	Malignant neoplasms (C00-C97)	50	12.9	59.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	9.5	44.0
5	Diabetes mellitus (E10-E14)	25	6.4	29.7
6	Cerebrovascular diseases (I60-I69)	14	3.6	*
7	Chronic lower respiratory diseases (J40-J47)	10	2.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	2.6	*
9	Septicemia (A40-A41)	7	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.5	*
11	Influenza and pneumonia (J09-J18)	5	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.3	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.0	*
14	Viral hepatitis (B15-B19)	2	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.5	*
...	All other causes (residual)	64	16.5	76.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 50-54 years				
...	All causes	556	100.0	593.7
1	Malignant neoplasms (C00-C97)	113	20.3	120.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	14.2	84.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	12.6	74.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	67	12.1	71.5
5	Diabetes mellitus (E10-E14)	41	7.4	43.8
6	Cerebrovascular diseases (I60-I69)	16	2.9	*
7	Chronic lower respiratory diseases (J40-J47)	14	2.5	*
8	Septicemia (A40-A41)	12	2.2	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.6	*
11	Influenza and pneumonia (J09-J18)	8	1.4	*
12	Viral hepatitis (B15-B19)	4	0.7	*
12	Pneumonitis due to solids and liquids (J69)	4	0.7	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.5	*
15	Anemias (D50-D64)	2	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Hernia (K40-K46)	2	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.4	*
...	All other causes (residual)	97	17.4	103.6
Non-Hispanic, American Indian or Alaska Native, female, 55-59 years				
...	All causes	633	100.0	702.6
1	Malignant neoplasms (C00-C97)	151	23.9	167.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	14.8	104.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	61	9.6	67.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	7.4	52.2
5	Diabetes mellitus (E10-E14)	40	6.3	44.4
6	Chronic lower respiratory diseases (J40-J47)	39	6.2	43.3
7	Cerebrovascular diseases (I60-I69)	19	3.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.4	*
9	Septicemia (A40-A41)	14	2.2	*
10	Influenza and pneumonia (J09-J18)	10	1.6	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.9	*
13	Viral hepatitis (B15-B19)	5	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	113	17.9	125.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 60-64 years				
...	All causes	713	100.0	974.6
1	Malignant neoplasms (C00-C97)	193	27.1	263.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	15.0	146.3
3	Diabetes mellitus (E10-E14)	61	8.6	83.4
4	Chronic lower respiratory diseases (J40-J47)	59	8.3	80.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	43	6.0	58.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	4.2	41.0
7	Septicemia (A40-A41)	24	3.4	32.8
7	Cerebrovascular diseases (I60-I69)	24	3.4	32.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.1	*
10	Viral hepatitis (B15-B19)	10	1.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.4	*
10	Influenza and pneumonia (J09-J18)	10	1.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Pneumonitis due to solids and liquids (J69)	4	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.4	*
...	All other causes (residual)	113	15.8	154.5
Non-Hispanic, American Indian or Alaska Native, female, 65-69 years				
...	All causes	781	100.0	1,391.2
1	Malignant neoplasms (C00-C97)	218	27.9	388.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	17.0	236.9
3	Diabetes mellitus (E10-E14)	74	9.5	131.8
4	Chronic lower respiratory diseases (J40-J47)	57	7.3	101.5
5	Cerebrovascular diseases (I60-I69)	34	4.4	60.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	29	3.7	51.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	3.1	42.8
8	Septicemia (A40-A41)	17	2.2	*
9	Influenza and pneumonia (J09-J18)	14	1.8	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.4	*
12	Viral hepatitis (B15-B19)	6	0.8	*
13	Peptic ulcer (K25-K28)	5	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
14	Anemias (D50-D64)	3	0.4	*
...	All other causes (residual)	140	17.9	249.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 70-74 years				
...	All causes	809	100.0	2,189.2
1	Malignant neoplasms (C00-C97)	189	23.4	511.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	18.8	411.3
3	Chronic lower respiratory diseases (J40-J47)	74	9.1	200.2
4	Diabetes mellitus (E10-E14)	69	8.5	186.7
5	Cerebrovascular diseases (I60-I69)	35	4.3	94.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	3.3	73.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	26	3.2	70.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.8	62.2
9	Septicemia (A40-A41)	20	2.5	54.1
10	Influenza and pneumonia (J09-J18)	16	2.0	*
11	Alzheimer's disease (G30)	12	1.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
13	Parkinson's disease (G20-G21)	4	0.5	*
15	Viral hepatitis (B15-B19)	3	0.4	*
15	Hernia (K40-K46)	3	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
...	All other causes (residual)	144	17.8	389.7
Non-Hispanic, American Indian or Alaska Native, female, 75-79 years				
...	All causes	804	100.0	3,232.8
1	Malignant neoplasms (C00-C97)	190	23.6	764.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	19.5	631.3
3	Diabetes mellitus (E10-E14)	61	7.6	245.3
3	Chronic lower respiratory diseases (J40-J47)	61	7.6	245.3
5	Cerebrovascular diseases (I60-I69)	37	4.6	148.8
6	Alzheimer's disease (G30)	29	3.6	116.6
7	Septicemia (A40-A41)	24	3.0	96.5
8	Influenza and pneumonia (J09-J18)	20	2.5	80.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.5	80.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.2	*
13	Parkinson's disease (G20-G21)	8	1.0	*
14	Pneumonitis due to solids and liquids (J69)	5	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
...	All other causes (residual)	148	18.4	595.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 80-84 years				
...	All causes	827	100.0	5,264.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	23.2	1,222.2
2	Malignant neoplasms (C00-C97)	142	17.2	903.9
3	Chronic lower respiratory diseases (J40-J47)	63	7.6	401.0
4	Cerebrovascular diseases (I60-I69)	54	6.5	343.7
5	Diabetes mellitus (E10-E14)	47	5.7	299.2
6	Alzheimer's disease (G30)	42	5.1	267.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.7	140.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	2.3	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.2	*
10	Influenza and pneumonia (J09-J18)	15	1.8	*
11	Septicemia (A40-A41)	14	1.7	*
12	Parkinson's disease (G20-G21)	11	1.3	*
13	Pneumonitis due to solids and liquids (J69)	9	1.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.0	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
...	All other causes (residual)	163	19.7	1,037.6
Non-Hispanic, American Indian or Alaska Native, female, 85-89 years				
...	All causes	691	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	22.6	---
2	Malignant neoplasms (C00-C97)	109	15.8	---
3	Cerebrovascular diseases (I60-I69)	51	7.4	---
4	Chronic lower respiratory diseases (J40-J47)	47	6.8	---
5	Alzheimer's disease (G30)	44	6.4	---
6	Diabetes mellitus (E10-E14)	32	4.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.6	---
10	Influenza and pneumonia (J09-J18)	13	1.9	---
11	Septicemia (A40-A41)	12	1.7	---
12	Nutritional deficiencies (E40-E64)	4	0.6	---
12	Atherosclerosis (I70)	4	0.6	---
12	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	---
15	Parkinson's disease (G20-G21)	3	0.4	---
15	Pneumonitis due to solids and liquids (J69)	3	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	---
...	All other causes (residual)	143	20.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 90-94 years				
...	All causes	524	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	25.0	---
2	Alzheimer's disease (G30)	59	11.3	---
3	Malignant neoplasms (C00-C97)	49	9.4	---
4	Cerebrovascular diseases (I60-I69)	37	7.1	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	3.8	---
6	Diabetes mellitus (E10-E14)	19	3.6	---
7	Influenza and pneumonia (J09-J18)	18	3.4	---
8	Chronic lower respiratory diseases (J40-J47)	15	2.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.9	---
11	Nutritional deficiencies (E40-E64)	7	1.3	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	---
12	Parkinson's disease (G20-G21)	6	1.1	---
14	Septicemia (A40-A41)	4	0.8	---
15	Atherosclerosis (I70)	3	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.6	---
...	All other causes (residual)	125	23.9	---
Non-Hispanic, American Indian or Alaska Native, female, 95-99 years				
...	All causes	195	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	23.6	---
2	Alzheimer's disease (G30)	14	7.2	---
3	Cerebrovascular diseases (I60-I69)	12	6.2	---
4	Malignant neoplasms (C00-C97)	11	5.6	---
5	Influenza and pneumonia (J09-J18)	9	4.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	3.6	---
7	Chronic lower respiratory diseases (J40-J47)	6	3.1	---
8	Diabetes mellitus (E10-E14)	4	2.1	---
9	Septicemia (A40-A41)	3	1.5	---
9	Pneumonitis due to solids and liquids (J69)	3	1.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.5	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	---
14	Anemias (D50-D64)	1	0.5	---
14	Nutritional deficiencies (E40-E64)	1	0.5	---
14	Parkinson's disease (G20-G21)	1	0.5	---
14	Atherosclerosis (I70)	1	0.5	---
14	Aortic aneurysm and dissection (I71)	1	0.5	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	---
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	---
...	All other causes (residual)	64	32.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 100 years and over				
...	All causes	59	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	32.2	---
2	Cerebrovascular diseases (I60-I69)	6	10.2	---
3	Alzheimer's disease (G30)	4	6.8	---
3	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	6.8	---
5	Influenza and pneumonia (J09-J18)	3	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3	5.1	---
7	Malignant neoplasms (C00-C97)	2	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.4	---
9	Anemias (D50-D64)	1	1.7	---
9	Diabetes mellitus (E10-E14)	1	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.7	---
...	All other causes (residual)	13	22.0	---
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	65,277	100.0	---
1	Malignant neoplasms (C00-C97)	17,306	26.5	---
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,974	21.4	---
3	Cerebrovascular diseases (I60-I69)	4,798	7.4	---
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,839	4.3	---
5	Diabetes mellitus (E10-E14)	2,599	4.0	---
6	Alzheimer's disease (G30)	2,180	3.3	---
7	Influenza and pneumonia (J09-J18)	2,161	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	1,874	2.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,318	2.0	---
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,273	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,136	1.7	---
12	Septicemia (A40-A41)	798	1.2	---
13	Parkinson's disease (G20-G21)	747	1.1	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	598	0.9	---
15	Pneumonitis due to solids and liquids (J69)	500	0.8	---
...	All other causes (residual)	11,176	17.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	141	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	18.4	2.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	14.2	2.2
3	Malignant neoplasms (C00-C97)	13	9.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	7.1	*
5	Influenza and pneumonia (J09-J18)	6	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.3	*
7	Septicemia (A40-A41)	3	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	1.4	*
10	Tuberculosis (A16-A19)	1	0.7	*
10	Anemias (D50-D64)	1	0.7	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	47	33.3	5.1
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	108	100.0	9.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	25.0	2.3
2	Malignant neoplasms (C00-C97)	25	23.1	2.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	4.6	*
5	Septicemia (A40-A41)	3	2.8	*
5	Cerebrovascular diseases (I60-I69)	3	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.9	*
7	Anemias (D50-D64)	2	1.9	*
7	Influenza and pneumonia (J09-J18)	2	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.9	*
11	Meningitis (G00,G03)	1	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (residual)	26	24.1	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	104	100.0	8.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	21.2	1.9
2	Malignant neoplasms (C00-C97)	15	14.4	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	14.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.8	*
6	Cerebrovascular diseases (I60-I69)	2	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.9	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.9	*
9	Septicemia (A40-A41)	1	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
...	All other causes (residual)	34	32.7	2.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	321	100.0	27.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	31.2	8.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	25.5	7.0
3	Malignant neoplasms (C00-C97)	41	12.8	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	5.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	3.4	*
7	Cerebrovascular diseases (I60-I69)	3	0.9	*
7	Influenza and pneumonia (J09-J18)	3	0.9	*
9	Legal intervention (Y35,Y89.0)	2	0.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
11	Septicemia (A40-A41)	1	0.3	*
11	Diabetes mellitus (E10-E14)	1	0.3	*
11	Meningitis (G00,G03)	1	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (residual)	45	14.0	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	552	100.0	39.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	33.2	13.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	32.1	12.6
3	Malignant neoplasms (C00-C97)	43	7.8	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.3	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.9	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
9	Aortic aneurysm and dissection (I71)	3	0.5	*
9	Pneumonitis due to solids and liquids (J69)	3	0.5	*
11	Septicemia (A40-A41)	2	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	59	10.7	4.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	610	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	32.3	12.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	22.1	8.3
3	Malignant neoplasms (C00-C97)	60	9.8	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	6.4	2.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	5.6	2.1
6	Cerebrovascular diseases (I60-I69)	12	2.0	*
7	Diabetes mellitus (E10-E14)	9	1.5	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
9	Aortic aneurysm and dissection (I71)	4	0.7	*
9	Influenza and pneumonia (J09-J18)	4	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
14	Legal intervention (Y35,Y89.0)	2	0.3	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Septicemia (A40-A41)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	87	14.3	5.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	722	100.0	44.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	22.3	9.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	18.1	8.0
3	Malignant neoplasms (C00-C97)	119	16.5	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	9.0	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.7	2.1
6	Cerebrovascular diseases (I60-I69)	17	2.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	17	2.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
9	Diabetes mellitus (E10-E14)	8	1.1	*
9	Influenza and pneumonia (J09-J18)	8	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
12	Septicemia (A40-A41)	7	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
...	All other causes (residual)	124	17.2	7.6
Non-Hispanic, Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	824	100.0	54.0
1	Malignant neoplasms (C00-C97)	223	27.1	14.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	13.7	7.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	13.5	7.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	10.7	5.8
5	Cerebrovascular diseases (I60-I69)	45	5.5	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	3.8	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.8	1.5
8	Diabetes mellitus (E10-E14)	18	2.2	*
9	Septicemia (A40-A41)	11	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.3	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.2	*
12	Influenza and pneumonia (J09-J18)	9	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.5	*
...	All other causes (residual)	113	13.7	7.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,223	100.0	80.8
1	Malignant neoplasms (C00-C97)	404	33.0	26.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	15.8	12.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	9.6	7.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	7.5	6.1
5	Cerebrovascular diseases (I60-I69)	69	5.6	4.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	33	2.7	2.2
7	Diabetes mellitus (E10-E14)	28	2.3	1.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	2.0	1.7
9	Influenza and pneumonia (J09-J18)	14	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	10	0.8	*
14	Septicemia (A40-A41)	9	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.7	*
...	All other causes (residual)	189	15.5	12.5
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,763	100.0	131.4
1	Malignant neoplasms (C00-C97)	663	37.6	49.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	300	17.0	22.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	7.6	10.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	6.5	8.6
5	Cerebrovascular diseases (I60-I69)	96	5.4	7.2
6	Diabetes mellitus (E10-E14)	60	3.4	4.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	41	2.3	3.1
8	Influenza and pneumonia (J09-J18)	25	1.4	1.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	1.3	1.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.1	1.5
11	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
13	Septicemia (A40-A41)	14	0.8	*
14	Viral hepatitis (B15-B19)	12	0.7	*
15	Tuberculosis (A16-A19)	11	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.6	*
...	All other causes (residual)	205	11.6	15.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,449	100.0	199.7
1	Malignant neoplasms (C00-C97)	1,015	41.4	82.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	18.7	37.4
3	Cerebrovascular diseases (I60-I69)	148	6.0	12.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	5.8	11.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	3.6	7.3
6	Diabetes mellitus (E10-E14)	81	3.3	6.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	60	2.4	4.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.6	3.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.0	2.0
10	Viral hepatitis (B15-B19)	22	0.9	1.8
11	Septicemia (A40-A41)	21	0.9	1.7
12	Influenza and pneumonia (J09-J18)	20	0.8	1.6
13	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.6	*
...	All other causes (residual)	282	11.5	23.0
Non-Hispanic, Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,462	100.0	311.0
1	Malignant neoplasms (C00-C97)	1,549	44.7	139.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	630	18.2	56.6
3	Cerebrovascular diseases (I60-I69)	218	6.3	19.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	4.5	14.0
5	Diabetes mellitus (E10-E14)	120	3.5	10.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	2.3	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	78	2.3	7.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.6	5.1
9	Septicemia (A40-A41)	44	1.3	4.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	1.1	3.4
11	Influenza and pneumonia (J09-J18)	37	1.1	3.3
12	Chronic lower respiratory diseases (J40-J47)	34	1.0	3.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	0.8	2.4
14	Aortic aneurysm and dissection (I71)	22	0.6	2.0
15	Viral hepatitis (B15-B19)	21	0.6	1.9
...	All other causes (residual)	351	10.1	31.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,547	100.0	479.7
1	Malignant neoplasms (C00-C97)	1,835	40.4	193.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	952	20.9	100.4
3	Cerebrovascular diseases (I60-I69)	296	6.5	31.2
4	Diabetes mellitus (E10-E14)	224	4.9	23.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	3.2	15.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	85	1.9	9.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	1.8	8.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	1.7	8.1
9	Chronic lower respiratory diseases (J40-J47)	71	1.6	7.5
10	Influenza and pneumonia (J09-J18)	67	1.5	7.1
11	Septicemia (A40-A41)	60	1.3	6.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	1.1	5.1
13	Viral hepatitis (B15-B19)	36	0.8	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.8	3.7
15	Aortic aneurysm and dissection (I71)	24	0.5	2.5
...	All other causes (residual)	512	11.3	54.0
Non-Hispanic, Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	5,463	100.0	712.2
1	Malignant neoplasms (C00-C97)	2,193	40.1	285.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,096	20.1	142.9
3	Cerebrovascular diseases (I60-I69)	336	6.2	43.8
4	Diabetes mellitus (E10-E14)	290	5.3	37.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	3.2	22.7
6	Chronic lower respiratory diseases (J40-J47)	120	2.2	15.6
7	Influenza and pneumonia (J09-J18)	108	2.0	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.6	11.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.3	9.3
10	Septicemia (A40-A41)	68	1.2	8.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	65	1.2	8.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	0.9	6.3
13	Parkinson's disease (G20-G21)	45	0.8	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.8	5.6
15	Viral hepatitis (B15-B19)	29	0.5	3.8
15	Aortic aneurysm and dissection (I71)	29	0.5	3.8
...	All other causes (residual)	662	12.1	86.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	6,033	100.0	1,166.9
1	Malignant neoplasms (C00-C97)	2,176	36.1	420.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,218	20.2	235.6
3	Cerebrovascular diseases (I60-I69)	451	7.5	87.2
4	Diabetes mellitus (E10-E14)	314	5.2	60.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	3.0	35.0
6	Chronic lower respiratory diseases (J40-J47)	180	3.0	34.8
7	Influenza and pneumonia (J09-J18)	153	2.5	29.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	138	2.3	26.7
9	Septicemia (A40-A41)	87	1.4	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	1.4	16.4
11	Parkinson's disease (G20-G21)	68	1.1	13.2
12	Alzheimer's disease (G30)	59	1.0	11.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.9	10.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	0.7	8.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	7.7
...	All other causes (residual)	785	13.0	151.8
Non-Hispanic, Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	7,323	100.0	2,001.1
1	Malignant neoplasms (C00-C97)	2,177	29.7	594.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,561	21.3	426.6
3	Cerebrovascular diseases (I60-I69)	609	8.3	166.4
4	Diabetes mellitus (E10-E14)	399	5.4	109.0
5	Chronic lower respiratory diseases (J40-J47)	255	3.5	69.7
6	Influenza and pneumonia (J09-J18)	240	3.3	65.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	2.8	55.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.7	53.3
9	Alzheimer's disease (G30)	159	2.2	43.4
10	Parkinson's disease (G20-G21)	135	1.8	36.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.7	33.3
12	Septicemia (A40-A41)	94	1.3	25.7
13	Pneumonitis due to solids and liquids (J69)	55	0.8	15.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.7	13.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	45	0.6	12.3
...	All other causes (residual)	1,025	14.0	280.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	8,592	100.0	3,592.7
1	Malignant neoplasms (C00-C97)	2,070	24.1	865.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,938	22.6	810.4
3	Cerebrovascular diseases (I60-I69)	784	9.1	327.8
4	Diabetes mellitus (E10-E14)	389	4.5	162.7
5	Chronic lower respiratory diseases (J40-J47)	359	4.2	150.1
6	Alzheimer's disease (G30)	349	4.1	145.9
7	Influenza and pneumonia (J09-J18)	345	4.0	144.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	2.9	103.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	2.3	82.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	2.1	74.0
11	Parkinson's disease (G20-G21)	157	1.8	65.6
12	Septicemia (A40-A41)	116	1.4	48.5
13	Pneumonitis due to solids and liquids (J69)	76	0.9	31.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.7	26.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.7	23.4
...	All other causes (residual)	1,267	14.7	529.8
Non-Hispanic, Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	9,279	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,306	24.9	---
2	Malignant neoplasms (C00-C97)	1,571	16.9	---
3	Cerebrovascular diseases (I60-I69)	807	8.7	---
4	Alzheimer's disease (G30)	597	6.4	---
5	Influenza and pneumonia (J09-J18)	441	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	382	4.1	---
7	Diabetes mellitus (E10-E14)	376	4.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	238	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	2.4	---
11	Parkinson's disease (G20-G21)	198	2.1	---
12	Septicemia (A40-A41)	127	1.4	---
13	Pneumonitis due to solids and liquids (J69)	106	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.8	---
15	Aortic aneurysm and dissection (I71)	48	0.5	---
...	All other causes (residual)	1,522	16.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	7,180	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,971	27.5	---
2	Malignant neoplasms (C00-C97)	831	11.6	---
3	Alzheimer's disease (G30)	599	8.3	---
4	Cerebrovascular diseases (I60-I69)	593	8.3	---
5	Influenza and pneumonia (J09-J18)	438	6.1	---
6	Chronic lower respiratory diseases (J40-J47)	286	4.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	210	2.9	---
8	Diabetes mellitus (E10-E14)	204	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	2.2	---
11	Pneumonitis due to solids and liquids (J69)	111	1.5	---
12	Parkinson's disease (G20-G21)	98	1.4	---
13	Septicemia (A40-A41)	82	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	---
15	Atherosclerosis (I70)	31	0.4	---
15	Aortic aneurysm and dissection (I71)	31	0.4	---
...	All other causes (residual)	1,340	18.7	---
Non-Hispanic, Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	2,909	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	830	28.5	---
2	Alzheimer's disease (G30)	288	9.9	---
2	Malignant neoplasms (C00-C97)	236	8.1	---
3	Cerebrovascular diseases (I60-I69)	236	8.1	---
5	Influenza and pneumonia (J09-J18)	178	6.1	---
6	Chronic lower respiratory diseases (J40-J47)	101	3.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	2.7	---
8	Diabetes mellitus (E10-E14)	69	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.2	---
11	Pneumonitis due to solids and liquids (J69)	45	1.5	---
12	Septicemia (A40-A41)	31	1.1	---
13	Parkinson's disease (G20-G21)	28	1.0	---
14	Nutritional deficiencies (E40-E64)	16	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	---
...	All other causes (residual)	631	21.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	779	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	28.9	---
2	Alzheimer's disease (G30)	81	10.4	---
3	Cerebrovascular diseases (I60-I69)	64	8.2	---
4	Influenza and pneumonia (J09-J18)	56	7.2	---
5	Malignant neoplasms (C00-C97)	42	5.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	3.3	---
7	Chronic lower respiratory diseases (J40-J47)	23	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.8	---
9	Pneumonitis due to solids and liquids (J69)	14	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.3	---
11	Septicemia (A40-A41)	9	1.2	---
12	Diabetes mellitus (E10-E14)	7	0.9	---
13	Atherosclerosis (I70)	4	0.5	---
14	Nutritional deficiencies (E40-E64)	3	0.4	---
14	Parkinson's disease (G20-G21)	3	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	---
...	All other causes (residual)	187	24.0	---
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	33,306	100.0	364.9
1	Malignant neoplasms (C00-C97)	8,774	26.3	96.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,612	22.9	83.4
3	Cerebrovascular diseases (I60-I69)	2,153	6.5	23.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,788	5.4	19.6
5	Diabetes mellitus (E10-E14)	1,322	4.0	14.5
6	Influenza and pneumonia (J09-J18)	1,109	3.3	12.2
7	Chronic lower respiratory diseases (J40-J47)	1,061	3.2	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	855	2.6	9.4
9	Alzheimer's disease (G30)	675	2.0	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	651	2.0	7.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	515	1.5	5.6
12	Parkinson's disease (G20-G21)	448	1.3	4.9
13	Septicemia (A40-A41)	418	1.3	4.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	384	1.2	4.2
15	Pneumonitis due to solids and liquids (J69)	285	0.9	3.1
...	All other causes (residual)	5,256	15.8	57.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	72	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	22.2	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	12.5	*
3	Malignant neoplasms (C00-C97)	7	9.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.2	*
5	Influenza and pneumonia (J09-J18)	3	4.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.2	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	22	30.6	4.7
Non-Hispanic, Asian or Pacific Islander, male, 5-9 years				
...	All causes	58	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	29.3	*
2	Malignant neoplasms (C00-C97)	13	22.4	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.4	*
6	Septicemia (A40-A41)	1	1.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
6	Meningitis (G00,G03)	1	1.7	*
6	Cerebrovascular diseases (I60-I69)	1	1.7	*
6	Influenza and pneumonia (J09-J18)	1	1.7	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.7	*
...	All other causes (residual)	15	25.9	*
Non-Hispanic, Asian or Pacific Islander, male, 10-14 years				
...	All causes	66	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	21.2	*
2	Malignant neoplasms (C00-C97)	12	18.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	12.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.1	*
5	Cerebrovascular diseases (I60-I69)	2	3.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.0	*
8	Septicemia (A40-A41)	1	1.5	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.5	*
...	All other causes (residual)	20	30.3	3.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 15-19 years				
...	All causes	225	100.0	38.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	30.7	11.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	27.1	10.3
3	Malignant neoplasms (C00-C97)	29	12.9	4.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	3.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.1	*
7	Cerebrovascular diseases (I60-I69)	3	1.3	*
8	Septicemia (A40-A41)	1	0.4	*
8	Diabetes mellitus (E10-E14)	1	0.4	*
8	Influenza and pneumonia (J09-J18)	1	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
8	Legal intervention (Y35,Y89.0)	1	0.4	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	28	12.4	4.7
Non-Hispanic, Asian or Pacific Islander, male, 20-24 years				
...	All causes	405	100.0	56.7
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	35.1	19.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	34.3	19.5
3	Malignant neoplasms (C00-C97)	25	6.2	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.9	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.9	2.8
6	Aortic aneurysm and dissection (I71)	3	0.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
8	Septicemia (A40-A41)	2	0.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
8	Cerebrovascular diseases (I60-I69)	2	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	32	7.9	4.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 25-29 years				
...	All causes	435	100.0	54.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	36.1	19.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	22.3	12.2
3	Malignant neoplasms (C00-C97)	36	8.3	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.4	3.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.2	3.4
6	Diabetes mellitus (E10-E14)	9	2.1	*
6	Cerebrovascular diseases (I60-I69)	9	2.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
9	Aortic aneurysm and dissection (I71)	4	0.9	*
11	Influenza and pneumonia (J09-J18)	2	0.5	*
11	Legal intervention (Y35,Y89.0)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Viral hepatitis (B15-B19)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
...	All other causes (residual)	51	11.7	6.4
Non-Hispanic, Asian or Pacific Islander, male, 30-34 years				
...	All causes	452	100.0	58.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	27.0	15.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	17.9	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	11.7	6.8
4	Malignant neoplasms (C00-C97)	49	10.8	6.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.2	3.6
6	Cerebrovascular diseases (I60-I69)	8	1.8	*
7	Influenza and pneumonia (J09-J18)	6	1.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Diabetes mellitus (E10-E14)	4	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
13	Septicemia (A40-A41)	2	0.4	*
13	Aortic aneurysm and dissection (I71)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
13	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (residual)	75	16.6	9.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 35-39 years				
...	All causes	503	100.0	70.5
1	Malignant neoplasms (C00-C97)	93	18.5	13.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	17.5	12.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	16.7	11.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	12.1	8.6
5	Cerebrovascular diseases (I60-I69)	24	4.8	3.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.6	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	17	3.4	*
8	Diabetes mellitus (E10-E14)	12	2.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.8	*
10	Influenza and pneumonia (J09-J18)	8	1.6	*
11	Septicemia (A40-A41)	6	1.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
15	Legal intervention (Y35,Y89.0)	3	0.6	*
...	All other causes (residual)	62	12.3	8.7
Non-Hispanic, Asian or Pacific Islander, male, 40-44 years				
...	All causes	709	100.0	99.9
1	Malignant neoplasms (C00-C97)	172	24.3	24.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	22.0	22.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	11.6	11.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	7.8	7.7
5	Cerebrovascular diseases (I60-I69)	31	4.4	4.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	25	3.5	3.5
7	Diabetes mellitus (E10-E14)	21	3.0	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.1	*
9	Influenza and pneumonia (J09-J18)	11	1.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.8	*
12	Aortic aneurysm and dissection (I71)	6	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.7	*
...	All other causes (residual)	103	14.5	14.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,032	100.0	164.3
1	Malignant neoplasms (C00-C97)	280	27.1	44.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	23.1	37.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	9.7	15.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	6.6	10.8
5	Cerebrovascular diseases (I60-I69)	55	5.3	8.8
6	Diabetes mellitus (E10-E14)	42	4.1	6.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	32	3.1	5.1
8	Influenza and pneumonia (J09-J18)	19	1.8	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	1.7	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.4	*
11	Viral hepatitis (B15-B19)	10	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
15	Septicemia (A40-A41)	6	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
15	Aortic aneurysm and dissection (I71)	6	0.6	*
...	All other causes (residual)	110	10.7	17.5
Non-Hispanic, Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,429	100.0	249.6
1	Malignant neoplasms (C00-C97)	468	32.8	81.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	24.9	62.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	6.5	16.2
4	Cerebrovascular diseases (I60-I69)	85	5.9	14.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	4.2	10.5
6	Diabetes mellitus (E10-E14)	45	3.1	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	44	3.1	7.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.9	4.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.3	*
10	Viral hepatitis (B15-B19)	13	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.8	*
11	Influenza and pneumonia (J09-J18)	12	0.8	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.8	*
14	Septicemia (A40-A41)	10	0.7	*
14	Aortic aneurysm and dissection (I71)	10	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.7	*
...	All other causes (residual)	153	10.7	26.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 55-59 years				
...	All causes	2,067	100.0	404.0
1	Malignant neoplasms (C00-C97)	766	37.1	149.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	464	22.4	90.7
3	Cerebrovascular diseases (I60-I69)	126	6.1	24.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	5.7	23.1
5	Diabetes mellitus (E10-E14)	80	3.9	15.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	70	3.4	13.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.4	9.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.9	7.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	1.4	5.5
10	Septicemia (A40-A41)	26	1.3	5.1
11	Influenza and pneumonia (J09-J18)	25	1.2	4.9
12	Chronic lower respiratory diseases (J40-J47)	23	1.1	4.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	0.9	*
14	Viral hepatitis (B15-B19)	15	0.7	*
15	Aortic aneurysm and dissection (I71)	10	0.5	*
...	All other causes (residual)	209	10.1	40.8
Non-Hispanic, Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,679	100.0	629.6
1	Malignant neoplasms (C00-C97)	943	35.2	221.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	665	24.8	156.3
3	Cerebrovascular diseases (I60-I69)	177	6.6	41.6
4	Diabetes mellitus (E10-E14)	135	5.0	31.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	3.3	20.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.5	15.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	2.0	12.7
8	Chronic lower respiratory diseases (J40-J47)	48	1.8	11.3
9	Influenza and pneumonia (J09-J18)	42	1.6	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.4	8.9
11	Septicemia (A40-A41)	36	1.3	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.2	7.8
13	Viral hepatitis (B15-B19)	23	0.9	5.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
15	Aortic aneurysm and dissection (I71)	17	0.6	*
...	All other causes (residual)	294	11.0	69.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 65-69 years				
...	All causes	3,152	100.0	920.3
1	Malignant neoplasms (C00-C97)	1,183	37.5	345.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	726	23.0	212.0
3	Cerebrovascular diseases (I60-I69)	187	5.9	54.6
4	Diabetes mellitus (E10-E14)	177	5.6	51.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	3.2	29.8
6	Chronic lower respiratory diseases (J40-J47)	81	2.6	23.7
7	Influenza and pneumonia (J09-J18)	62	2.0	18.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.6	14.6
9	Septicemia (A40-A41)	46	1.5	13.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.3	12.0
11	Parkinson's disease (G20-G21)	34	1.1	9.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.0	9.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.9	8.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	6.1
15	Aortic aneurysm and dissection (I71)	18	0.6	*
...	All other causes (residual)	362	11.5	105.7
Non-Hispanic, Asian or Pacific Islander, male, 70-74 years				
...	All causes	3,347	100.0	1,446.7
1	Malignant neoplasms (C00-C97)	1,196	35.7	516.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	729	21.8	315.1
3	Cerebrovascular diseases (I60-I69)	235	7.0	101.6
4	Diabetes mellitus (E10-E14)	163	4.9	70.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	3.4	48.8
6	Chronic lower respiratory diseases (J40-J47)	109	3.3	47.1
7	Influenza and pneumonia (J09-J18)	84	2.5	36.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	2.2	31.1
9	Septicemia (A40-A41)	48	1.4	20.7
10	Parkinson's disease (G20-G21)	40	1.2	17.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.2	17.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.0	14.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	30	0.9	13.0
14	Alzheimer's disease (G30)	27	0.8	11.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.7	10.4
...	All other causes (residual)	403	12.0	174.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 75-79 years				
...	All causes	3,934	100.0	2,414.7
1	Malignant neoplasms (C00-C97)	1,169	29.7	717.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	888	22.6	545.1
3	Cerebrovascular diseases (I60-I69)	303	7.7	186.0
4	Diabetes mellitus (E10-E14)	196	5.0	120.3
5	Influenza and pneumonia (J09-J18)	153	3.9	93.9
6	Chronic lower respiratory diseases (J40-J47)	134	3.4	82.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	2.7	65.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	2.5	61.4
9	Parkinson's disease (G20-G21)	89	2.3	54.6
10	Alzheimer's disease (G30)	74	1.9	45.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.5	36.8
12	Septicemia (A40-A41)	51	1.3	31.3
13	Pneumonitis due to solids and liquids (J69)	31	0.8	19.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	0.7	16.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.6	15.3
...	All other causes (residual)	527	13.4	323.5
Non-Hispanic, Asian or Pacific Islander, male, 80-84 years				
...	All causes	4,171	100.0	4,221.3
1	Malignant neoplasms (C00-C97)	1,049	25.1	1,061.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	988	23.7	999.9
3	Cerebrovascular diseases (I60-I69)	331	7.9	335.0
4	Chronic lower respiratory diseases (J40-J47)	209	5.0	211.5
5	Influenza and pneumonia (J09-J18)	174	4.2	176.1
6	Diabetes mellitus (E10-E14)	172	4.1	174.1
7	Alzheimer's disease (G30)	137	3.3	138.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	3.0	127.5
9	Parkinson's disease (G20-G21)	108	2.6	109.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	2.4	100.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.7	69.8
12	Septicemia (A40-A41)	62	1.5	62.7
13	Pneumonitis due to solids and liquids (J69)	42	1.0	42.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.7	30.4
15	Aortic aneurysm and dissection (I71)	27	0.6	27.3
...	All other causes (residual)	548	13.1	554.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 85-89 years				
...	All causes	4,007	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,026	25.6	---
2	Malignant neoplasms (C00-C97)	764	19.1	---
3	Cerebrovascular diseases (I60-I69)	285	7.1	---
4	Influenza and pneumonia (J09-J18)	222	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	213	5.3	---
6	Alzheimer's disease (G30)	179	4.5	---
7	Diabetes mellitus (E10-E14)	158	3.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	2.8	---
9	Parkinson's disease (G20-G21)	108	2.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	2.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	2.4	---
12	Pneumonitis due to solids and liquids (J69)	64	1.6	---
13	Septicemia (A40-A41)	62	1.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.4	---
...	All other causes (residual)	575	14.3	---
Non-Hispanic, Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,852	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	795	27.9	---
2	Malignant neoplasms (C00-C97)	380	13.3	---
3	Cerebrovascular diseases (I60-I69)	202	7.1	---
4	Influenza and pneumonia (J09-J18)	198	6.9	---
5	Alzheimer's disease (G30)	168	5.9	---
6	Chronic lower respiratory diseases (J40-J47)	140	4.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.9	---
8	Diabetes mellitus (E10-E14)	74	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	2.4	---
11	Pneumonitis due to solids and liquids (J69)	65	2.3	---
12	Parkinson's disease (G20-G21)	47	1.6	---
13	Septicemia (A40-A41)	31	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	---
15	Nutritional deficiencies (E40-E64)	12	0.4	---
15	Atherosclerosis (I70)	12	0.4	---
15	Aortic aneurysm and dissection (I71)	12	0.4	---
...	All other causes (residual)	479	16.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 95-99 years				
...	All causes	992	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	293	29.5	---
2	Malignant neoplasms (C00-C97)	121	12.2	---
3	Cerebrovascular diseases (I60-I69)	70	7.1	---
4	Influenza and pneumonia (J09-J18)	59	5.9	---
5	Alzheimer's disease (G30)	56	5.6	---
6	Chronic lower respiratory diseases (J40-J47)	55	5.5	---
7	Diabetes mellitus (E10-E14)	29	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	2.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.2	---
11	Septicemia (A40-A41)	18	1.8	---
12	Pneumonitis due to solids and liquids (J69)	17	1.7	---
13	Parkinson's disease (G20-G21)	11	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	---
15	Nutritional deficiencies (E40-E64)	6	0.6	---
...	All other causes (residual)	173	17.4	---
Non-Hispanic, Asian or Pacific Islander, male, 100 years and over				
...	All causes	218	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	30.7	---
2	Influenza and pneumonia (J09-J18)	22	10.1	---
3	Malignant neoplasms (C00-C97)	17	7.8	---
4	Chronic lower respiratory diseases (J40-J47)	16	7.3	---
5	Cerebrovascular diseases (I60-I69)	14	6.4	---
6	Alzheimer's disease (G30)	13	6.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	3.2	---
8	Pneumonitis due to solids and liquids (J69)	5	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.3	---
10	Diabetes mellitus (E10-E14)	3	1.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.4	---
12	Parkinson's disease (G20-G21)	2	0.9	---
12	Atherosclerosis (I70)	2	0.9	---
14	Septicemia (A40-A41)	1	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
14	Nutritional deficiencies (E40-E64)	1	0.5	---
14	Aortic aneurysm and dissection (I71)	1	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	---
...	All other causes (residual)	37	17.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	31,971	100.0	320.1
1	Malignant neoplasms (C00-C97)	8,532	26.7	85.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,362	19.9	63.7
3	Cerebrovascular diseases (I60-I69)	2,645	8.3	26.5
4	Alzheimer's disease (G30)	1,505	4.7	15.1
5	Diabetes mellitus (E10-E14)	1,277	4.0	12.8
6	Influenza and pneumonia (J09-J18)	1,052	3.3	10.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,051	3.3	10.5
8	Chronic lower respiratory diseases (J40-J47)	813	2.5	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	667	2.1	6.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	621	1.9	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	1.3	4.2
12	Septicemia (A40-A41)	380	1.2	3.8
13	Parkinson's disease (G20-G21)	299	0.9	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	222	0.7	2.2
15	Pneumonitis due to solids and liquids (J69)	215	0.7	2.2
...	All other causes (residual)	5,912	18.5	59.2
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	69	100.0	15.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	14.5	*
3	Malignant neoplasms (C00-C97)	6	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	7.2	*
5	Septicemia (A40-A41)	3	4.3	*
5	Influenza and pneumonia (J09-J18)	3	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.3	*
8	Tuberculosis (A16-A19)	1	1.4	*
8	Anemias (D50-D64)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	25	36.2	5.6
Non-Hispanic, Asian or Pacific Islander, female, 5-9 years				
...	All causes	50	100.0	8.7
1	Malignant neoplasms (C00-C97)	12	24.0	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	20.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	12.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.0	*
5	Septicemia (A40-A41)	2	4.0	*
5	Anemias (D50-D64)	2	4.0	*
5	Cerebrovascular diseases (I60-I69)	2	4.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.0	*
8	Influenza and pneumonia (J09-J18)	1	2.0	*
...	All other causes (residual)	11	22.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 10-14 years				
...	All causes	38	100.0	6.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	21.1	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	18.4	*
3	Malignant neoplasms (C00-C97)	3	7.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (residual)	14	36.8	*
Non-Hispanic, Asian or Pacific Islander, female, 15-19 years				
...	All causes	96	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	32.3	5.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	21.9	3.6
3	Malignant neoplasms (C00-C97)	12	12.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.2	*
6	Influenza and pneumonia (J09-J18)	2	2.1	*
7	Meningitis (G00,G03)	1	1.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.0	*
7	Legal intervention (Y35,Y89.0)	1	1.0	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.0	*
...	All other causes (residual)	17	17.7	*
Non-Hispanic, Asian or Pacific Islander, female, 20-24 years				
...	All causes	147	100.0	21.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	27.9	5.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	25.9	5.5
3	Malignant neoplasms (C00-C97)	18	12.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	6.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.4	*
6	Cerebrovascular diseases (I60-I69)	2	1.4	*
6	Chronic lower respiratory diseases (J40-J47)	2	1.4	*
6	Pneumonitis due to solids and liquids (J69)	2	1.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.4	*
10	Tuberculosis (A16-A19)	1	0.7	*
10	Anemias (D50-D64)	1	0.7	*
10	Diabetes mellitus (E10-E14)	1	0.7	*
10	Influenza and pneumonia (J09-J18)	1	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.7	*
10	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (residual)	20	13.6	2.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 25-29 years				
...	All causes	175	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	22.9	4.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	21.7	4.6
3	Malignant neoplasms (C00-C97)	24	13.7	2.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.0	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.3	*
7	Cerebrovascular diseases (I60-I69)	3	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.7	*
9	Influenza and pneumonia (J09-J18)	2	1.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
11	Septicemia (A40-A41)	1	0.6	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
11	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (residual)	36	20.6	4.3
Non-Hispanic, Asian or Pacific Islander, female, 30-34 years				
...	All causes	270	100.0	31.4
1	Malignant neoplasms (C00-C97)	70	25.9	8.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	18.5	5.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	14.4	4.5
4	Pregnancy, childbirth and the puerperium (O00-O99)	17	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.4	*
6	Cerebrovascular diseases (I60-I69)	9	3.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.2	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	2.2	*
9	Septicemia (A40-A41)	5	1.9	*
10	Diabetes mellitus (E10-E14)	4	1.5	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.5	*
12	Influenza and pneumonia (J09-J18)	2	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	39	14.4	4.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 35-39 years				
...	All causes	321	100.0	39.5
1	Malignant neoplasms (C00-C97)	130	40.5	16.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	8.4	3.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	8.4	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	7.8	3.1
5	Cerebrovascular diseases (I60-I69)	21	6.5	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	11	3.4	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	2.5	*
8	Diabetes mellitus (E10-E14)	6	1.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.9	*
10	Septicemia (A40-A41)	5	1.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Anemias (D50-D64)	1	0.3	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Influenza and pneumonia (J09-J18)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	43	13.4	5.3
Non-Hispanic, Asian or Pacific Islander, female, 40-44 years				
...	All causes	514	100.0	63.9
1	Malignant neoplasms (C00-C97)	232	45.1	28.8
2	Cerebrovascular diseases (I60-I69)	38	7.4	4.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	7.2	4.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	7.2	4.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	7.0	4.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	10	1.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.6	*
9	Diabetes mellitus (E10-E14)	7	1.4	*
10	Septicemia (A40-A41)	5	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
13	Anemias (D50-D64)	3	0.6	*
13	Influenza and pneumonia (J09-J18)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (residual)	76	14.8	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 45-49 years				
...	All causes	731	100.0	102.5
1	Malignant neoplasms (C00-C97)	383	52.4	53.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	8.5	8.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	6.4	6.6
4	Cerebrovascular diseases (I60-I69)	41	5.6	5.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	4.7	4.8
6	Diabetes mellitus (E10-E14)	18	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.2	*
9	Septicemia (A40-A41)	8	1.1	*
10	Tuberculosis (A16-A19)	6	0.8	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.8	*
10	Influenza and pneumonia (J09-J18)	6	0.8	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.7	*
...	All other causes (residual)	85	11.6	11.9
Non-Hispanic, Asian or Pacific Islander, female, 50-54 years				
...	All causes	1,020	100.0	156.0
1	Malignant neoplasms (C00-C97)	547	53.6	83.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	10.1	15.7
3	Cerebrovascular diseases (I60-I69)	63	6.2	9.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	4.9	7.6
5	Diabetes mellitus (E10-E14)	36	3.5	5.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.8	4.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.6	*
8	Septicemia (A40-A41)	11	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
10	Viral hepatitis (B15-B19)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
11	Influenza and pneumonia (J09-J18)	8	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.5	*
15	Anemias (D50-D64)	4	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.4	*
...	All other causes (residual)	110	10.8	16.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,395	100.0	231.9
1	Malignant neoplasms (C00-C97)	783	56.1	130.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	11.9	27.6
3	Cerebrovascular diseases (I60-I69)	92	6.6	15.3
4	Diabetes mellitus (E10-E14)	40	2.9	6.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	2.7	6.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	2.2	5.0
7	Septicemia (A40-A41)	18	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.3	*
9	Aortic aneurysm and dissection (I71)	12	0.9	*
9	Influenza and pneumonia (J09-J18)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.6	*
...	All other causes (residual)	140	10.0	23.3
Non-Hispanic, Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,868	100.0	357.6
1	Malignant neoplasms (C00-C97)	892	47.8	170.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	15.4	54.9
3	Cerebrovascular diseases (I60-I69)	119	6.4	22.8
4	Diabetes mellitus (E10-E14)	89	4.8	17.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	2.9	10.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.1	7.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	1.4	5.2
8	Influenza and pneumonia (J09-J18)	25	1.3	4.8
9	Septicemia (A40-A41)	24	1.3	4.6
10	Chronic lower respiratory diseases (J40-J47)	23	1.2	4.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.8	*
14	Viral hepatitis (B15-B19)	13	0.7	*
15	Aortic aneurysm and dissection (I71)	7	0.4	*
15	Pneumonitis due to solids and liquids (J69)	7	0.4	*
...	All other causes (residual)	211	11.3	40.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 65-69 years				
...	All causes	2,311	100.0	544.3
1	Malignant neoplasms (C00-C97)	1,010	43.7	237.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	16.0	87.1
3	Cerebrovascular diseases (I60-I69)	149	6.4	35.1
4	Diabetes mellitus (E10-E14)	113	4.9	26.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	3.1	17.0
6	Influenza and pneumonia (J09-J18)	46	2.0	10.8
7	Chronic lower respiratory diseases (J40-J47)	39	1.7	9.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	36	1.6	8.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.6	8.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.3	7.1
11	Septicemia (A40-A41)	22	1.0	5.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.0	5.2
13	Alzheimer's disease (G30)	17	0.7	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.6	*
15	Viral hepatitis (B15-B19)	12	0.5	*
...	All other causes (residual)	322	13.9	75.8
Non-Hispanic, Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,686	100.0	940.3
1	Malignant neoplasms (C00-C97)	980	36.5	343.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	489	18.2	171.2
3	Cerebrovascular diseases (I60-I69)	216	8.0	75.6
4	Diabetes mellitus (E10-E14)	151	5.6	52.9
5	Chronic lower respiratory diseases (J40-J47)	71	2.6	24.9
6	Influenza and pneumonia (J09-J18)	69	2.6	24.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.5	23.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	2.5	23.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.7	15.8
10	Septicemia (A40-A41)	39	1.5	13.7
11	Alzheimer's disease (G30)	32	1.2	11.2
12	Parkinson's disease (G20-G21)	28	1.0	9.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.9	8.8
14	Aortic aneurysm and dissection (I71)	19	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
...	All other causes (residual)	372	13.8	130.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 75-79 years				
...	All causes	3,389	100.0	1,669.3
1	Malignant neoplasms (C00-C97)	1,008	29.7	496.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	673	19.9	331.5
3	Cerebrovascular diseases (I60-I69)	306	9.0	150.7
4	Diabetes mellitus (E10-E14)	203	6.0	100.0
5	Chronic lower respiratory diseases (J40-J47)	121	3.6	59.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	2.8	46.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.8	46.8
8	Influenza and pneumonia (J09-J18)	87	2.6	42.9
9	Alzheimer's disease (G30)	85	2.5	41.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.8	30.5
11	Parkinson's disease (G20-G21)	46	1.4	22.7
12	Septicemia (A40-A41)	43	1.3	21.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	12.3
14	Pneumonitis due to solids and liquids (J69)	24	0.7	11.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.7	11.8
...	All other causes (residual)	492	14.5	242.3
Non-Hispanic, Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,421	100.0	3,150.2
1	Malignant neoplasms (C00-C97)	1,021	23.1	727.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	950	21.5	676.9
3	Cerebrovascular diseases (I60-I69)	453	10.2	322.8
4	Diabetes mellitus (E10-E14)	217	4.9	154.6
5	Alzheimer's disease (G30)	212	4.8	151.1
6	Influenza and pneumonia (J09-J18)	171	3.9	121.8
7	Chronic lower respiratory diseases (J40-J47)	150	3.4	106.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	2.8	86.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	2.4	77.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	2.2	70.5
11	Septicemia (A40-A41)	54	1.2	38.5
12	Parkinson's disease (G20-G21)	49	1.1	34.9
13	Pneumonitis due to solids and liquids (J69)	34	0.8	24.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	34	0.8	24.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.7	23.5
...	All other causes (residual)	714	16.2	508.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 85-89 years				
...	All causes	5,272	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,280	24.3	---
2	Malignant neoplasms (C00-C97)	807	15.3	---
3	Cerebrovascular diseases (I60-I69)	522	9.9	---
4	Alzheimer's disease (G30)	418	7.9	---
5	Influenza and pneumonia (J09-J18)	219	4.2	---
6	Diabetes mellitus (E10-E14)	218	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	169	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	2.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	2.4	---
11	Parkinson's disease (G20-G21)	90	1.7	---
12	Septicemia (A40-A41)	65	1.2	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.9	---
14	Pneumonitis due to solids and liquids (J69)	42	0.8	---
15	Aortic aneurysm and dissection (I71)	33	0.6	---
...	All other causes (residual)	946	17.9	---
Non-Hispanic, Asian or Pacific Islander, female, 90-94 years				
...	All causes	4,328	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,176	27.2	---
2	Malignant neoplasms (C00-C97)	451	10.4	---
3	Alzheimer's disease (G30)	431	10.0	---
4	Cerebrovascular diseases (I60-I69)	391	9.0	---
5	Influenza and pneumonia (J09-J18)	240	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	146	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	3.2	---
8	Diabetes mellitus (E10-E14)	130	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	1.7	---
11	Septicemia (A40-A41)	51	1.2	---
11	Parkinson's disease (G20-G21)	51	1.2	---
13	Pneumonitis due to solids and liquids (J69)	46	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	---
15	Atherosclerosis (I70)	19	0.4	---
15	Aortic aneurysm and dissection (I71)	19	0.4	---
...	All other causes (residual)	849	19.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,917	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	537	28.0	---
2	Alzheimer's disease (G30)	232	12.1	---
3	Cerebrovascular diseases (I60-I69)	166	8.7	---
4	Influenza and pneumonia (J09-J18)	119	6.2	---
5	Malignant neoplasms (C00-C97)	115	6.0	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	46	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	2.1	---
9	Diabetes mellitus (E10-E14)	40	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	2.0	---
11	Pneumonitis due to solids and liquids (J69)	28	1.5	---
12	Parkinson's disease (G20-G21)	17	0.9	---
13	Septicemia (A40-A41)	13	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	7	0.4	---
15	Atherosclerosis (I70)	7	0.4	---
...	All other causes (residual)	450	23.5	---
Non-Hispanic, Asian or Pacific Islander, female, 100 years and over				
...	All causes	561	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	158	28.2	---
2	Alzheimer's disease (G30)	68	12.1	---
3	Cerebrovascular diseases (I60-I69)	50	8.9	---
4	Influenza and pneumonia (J09-J18)	34	6.1	---
5	Malignant neoplasms (C00-C97)	25	4.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	3.0	---
8	Pneumonitis due to solids and liquids (J69)	9	1.6	---
9	Septicemia (A40-A41)	8	1.4	---
10	Chronic lower respiratory diseases (J40-J47)	7	1.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.2	---
12	Diabetes mellitus (E10-E14)	4	0.7	---
13	Anemias (D50-D64)	2	0.4	---
13	Nutritional deficiencies (E40-E64)	2	0.4	---
13	Atherosclerosis (I70)	2	0.4	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.4	---
...	All other causes (residual)	145	25.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015