

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,813,503	100.0	863.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	647,457	23.0	198.8
2	Malignant neoplasms (C00-C97)	599,108	21.3	183.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169,936	6.0	52.2
4	Chronic lower respiratory diseases (J40-J47)	160,201	5.7	49.2
5	Cerebrovascular diseases (I60-I69)	146,383	5.2	44.9
6	Alzheimer disease (G30)	121,404	4.3	37.3
7	Diabetes mellitus (E10-E14)	83,564	3.0	25.7
8	Influenza and pneumonia (J09-J18)	55,672	2.0	17.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,633	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47,173	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41,743	1.5	12.8
12	Septicemia (A40-A41)	40,922	1.5	12.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35,316	1.3	10.8
14	Parkinson disease (G20-G21)	31,963	1.1	9.8
15	Pneumonitis due to solids and liquids (J69)	20,108	0.7	6.2
...	All other causes (residual)	561,920	20.0	172.5
All races and origins, both sexes, 1-4 years				
...	All causes	3,880	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	32.7	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	424	10.9	2.7
3	Malignant neoplasms (C00-C97)	325	8.4	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	7.8	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.3	0.8
6	Influenza and pneumonia (J09-J18)	104	2.7	0.7
7	Cerebrovascular diseases (I60-I69)	66	1.7	0.4
8	Septicemia (A40-A41)	48	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	42	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.2
12	Anemias (D50-D64)	24	0.6	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	24	0.6	0.2
14	Acute bronchitis and bronchiolitis (J20-J21)	22	0.6	0.1
15	Meningitis (G00,G03)	19	0.5	*
...	All other causes (residual)	1,010	26.0	6.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-9 years				
...	All causes	2,354	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	718	30.5	3.5
2	Malignant neoplasms (C00-C97)	418	17.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	8.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	6.5	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.2	0.4
6	Influenza and pneumonia (J09-J18)	62	2.6	0.3
7	Chronic lower respiratory diseases (J40-J47)	59	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	41	1.7	0.2
9	Septicemia (A40-A41)	33	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.3	0.2
11	Anemias (D50-D64)	10	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
14	Diabetes mellitus (E10-E14)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (residual)	530	22.5	2.6
All races and origins, both sexes, 10-14 years				
...	All causes	3,217	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	860	26.7	4.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	517	16.1	2.5
3	Malignant neoplasms (C00-C97)	437	13.6	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	5.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	5.5	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	75	2.3	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.7	0.3
9	Influenza and pneumonia (J09-J18)	51	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.0	0.1
11	Diabetes mellitus (E10-E14)	26	0.8	0.1
12	Septicemia (A40-A41)	24	0.7	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
14	Anemias (D50-D64)	14	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (residual)	627	19.5	3.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 15-19 years				
...	All causes	10,886	100.0	51.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,948	36.3	18.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,491	22.9	11.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,835	16.9	8.7
4	Malignant neoplasms (C00-C97)	560	5.1	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	2.8	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	1.5	0.8
7	Diabetes mellitus (E10-E14)	76	0.7	0.4
8	Influenza and pneumonia (J09-J18)	69	0.6	0.3
9	Cerebrovascular diseases (I60-I69)	66	0.6	0.3
10	Chronic lower respiratory diseases (J40-J47)	65	0.6	0.3
11	Septicemia (A40-A41)	38	0.3	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.3	0.2
12	Legal intervention (Y35,Y89.0)	33	0.3	0.2
14	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.3	0.1
15	Anemias (D50-D64)	21	0.2	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	21	0.2	0.1
...	All other causes (residual)	1,127	10.4	5.3
All races and origins, both sexes, 20-24 years				
...	All causes	21,139	100.0	95.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,493	44.9	42.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,761	17.8	17.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,070	14.5	13.9
4	Malignant neoplasms (C00-C97)	814	3.9	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	606	2.9	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	187	0.9	0.8
7	Diabetes mellitus (E10-E14)	172	0.8	0.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	140	0.7	0.6
9	Chronic lower respiratory diseases (J40-J47)	123	0.6	0.6
10	Influenza and pneumonia (J09-J18)	121	0.6	0.5
11	Cerebrovascular diseases (I60-I69)	89	0.4	0.4
12	Septicemia (A40-A41)	85	0.4	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	66	0.3	0.3
14	Legal intervention (Y35,Y89.0)	64	0.3	0.3
15	Anemias (D50-D64)	48	0.2	0.2
...	All other causes (residual)	2,300	10.9	10.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-29 years				
...	All causes	28,276	100.0	121.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,734	45.0	54.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,069	14.4	17.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,083	10.9	13.2
4	Malignant neoplasms (C00-C97)	1,301	4.6	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,266	4.5	5.4
6	Diabetes mellitus (E10-E14)	344	1.2	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	257	0.9	1.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	253	0.9	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	0.8	0.9
10	Cerebrovascular diseases (I60-I69)	216	0.8	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	215	0.8	0.9
12	Septicemia (A40-A41)	172	0.6	0.7
13	Influenza and pneumonia (J09-J18)	156	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	153	0.5	0.7
15	Legal intervention (Y35,Y89.0)	109	0.4	0.5
...	All other causes (residual)	3,731	13.2	16.0
All races and origins, both sexes, 30-34 years				
...	All causes	31,939	100.0	145.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,935	40.5	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,879	12.1	17.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,415	7.6	11.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,405	7.5	10.9
5	Malignant neoplasms (C00-C97)	2,315	7.2	10.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	661	2.1	3.0
7	Diabetes mellitus (E10-E14)	479	1.5	2.2
8	Cerebrovascular diseases (I60-I69)	377	1.2	1.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	298	0.9	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	259	0.8	1.2
11	Influenza and pneumonia (J09-J18)	249	0.8	1.1
12	Septicemia (A40-A41)	220	0.7	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	220	0.7	1.0
14	Chronic lower respiratory diseases (J40-J47)	179	0.6	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	0.5	0.7
...	All other causes (residual)	4,895	15.3	22.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 35-39 years				
...	All causes	36,901	100.0	173.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,351	33.5	58.2
2	Malignant neoplasms (C00-C97)	4,104	11.1	19.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,909	10.6	18.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,722	10.1	17.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,984	5.4	9.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,227	3.3	5.8
7	Diabetes mellitus (E10-E14)	834	2.3	3.9
8	Cerebrovascular diseases (I60-I69)	685	1.9	3.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	404	1.1	1.9
10	Septicemia (A40-A41)	366	1.0	1.7
11	Influenza and pneumonia (J09-J18)	325	0.9	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	249	0.7	1.2
14	Pregnancy, childbirth and the puerperium (O00-O99)	217	0.6	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	0.6	1.0
...	All other causes (residual)	6,051	16.4	28.5
All races and origins, both sexes, 40-44 years				
...	All causes	42,895	100.0	218.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,477	24.4	53.3
2	Malignant neoplasms (C00-C97)	6,796	15.8	34.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,492	15.1	33.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,613	8.4	18.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,773	4.1	9.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,367	3.2	7.0
7	Diabetes mellitus (E10-E14)	1,284	3.0	6.5
8	Cerebrovascular diseases (I60-I69)	1,126	2.6	5.7
9	Septicemia (A40-A41)	488	1.1	2.5
10	Chronic lower respiratory diseases (J40-J47)	464	1.1	2.4
11	Influenza and pneumonia (J09-J18)	457	1.1	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	1.0	2.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	427	1.0	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	305	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	202	0.5	1.0
...	All other causes (residual)	7,188	16.8	36.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 45-49 years				
...	All causes	65,698	100.0	313.2
1	Malignant neoplasms (C00-C97)	13,108	20.0	62.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,647	17.7	55.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,633	17.7	55.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,230	6.4	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,222	4.9	15.4
6	Diabetes mellitus (E10-E14)	2,398	3.7	11.4
7	Cerebrovascular diseases (I60-I69)	1,946	3.0	9.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	1.8	5.8
9	Chronic lower respiratory diseases (J40-J47)	1,017	1.5	4.8
10	Septicemia (A40-A41)	876	1.3	4.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	806	1.2	3.8
12	Influenza and pneumonia (J09-J18)	765	1.2	3.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	666	1.0	3.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	585	0.9	2.8
15	Aortic aneurysm and dissection (I71)	301	0.5	1.4
...	All other causes (residual)	11,286	17.2	53.8
All races and origins, both sexes, 50-54 years				
...	All causes	104,444	100.0	488.0
1	Malignant neoplasms (C00-C97)	26,158	25.0	122.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,011	20.1	98.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,828	12.3	59.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,090	4.9	23.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,331	4.1	20.2
6	Diabetes mellitus (E10-E14)	4,011	3.8	18.7
7	Cerebrovascular diseases (I60-I69)	3,252	3.1	15.2
8	Chronic lower respiratory diseases (J40-J47)	2,958	2.8	13.8
9	Septicemia (A40-A41)	1,565	1.5	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,391	1.3	6.5
11	Influenza and pneumonia (J09-J18)	1,248	1.2	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,063	1.0	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,057	1.0	4.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	943	0.9	4.4
15	Viral hepatitis (B15-B19)	653	0.6	3.1
...	All other causes (residual)	16,885	16.2	78.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 55-59 years				
...	All causes	162,098	100.0	736.5
1	Malignant neoplasms (C00-C97)	47,646	29.4	216.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,228	21.1	155.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,142	8.1	59.7
4	Chronic lower respiratory diseases (J40-J47)	6,963	4.3	31.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,851	4.2	31.1
6	Diabetes mellitus (E10-E14)	6,356	3.9	28.9
7	Cerebrovascular diseases (I60-I69)	5,243	3.2	23.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,419	2.7	20.1
9	Septicemia (A40-A41)	2,414	1.5	11.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,303	1.4	10.5
11	Influenza and pneumonia (J09-J18)	2,079	1.3	9.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,684	1.0	7.7
13	Viral hepatitis (B15-B19)	1,169	0.7	5.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	975	0.6	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	921	0.6	4.2
...	All other causes (residual)	25,705	15.9	116.8
All races and origins, both sexes, 60-64 years				
...	All causes	209,908	100.0	1,050.2
1	Malignant neoplasms (C00-C97)	67,164	32.0	336.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,874	21.9	229.5
3	Chronic lower respiratory diseases (J40-J47)	11,704	5.6	58.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,266	4.9	51.4
5	Diabetes mellitus (E10-E14)	8,548	4.1	42.8
6	Cerebrovascular diseases (I60-I69)	7,465	3.6	37.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,886	3.3	34.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,563	1.7	17.8
9	Septicemia (A40-A41)	3,424	1.6	17.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,368	1.6	16.9
11	Influenza and pneumonia (J09-J18)	2,966	1.4	14.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,307	1.1	11.5
13	Viral hepatitis (B15-B19)	1,348	0.6	6.7
14	Pneumonitis due to solids and liquids (J69)	944	0.4	4.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	921	0.4	4.6
...	All other causes (residual)	33,160	15.8	165.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65-69 years				
...	All causes	248,087	100.0	1,473.5
1	Malignant neoplasms (C00-C97)	81,606	32.9	484.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,469	22.0	323.5
3	Chronic lower respiratory diseases (J40-J47)	16,531	6.7	98.2
4	Diabetes mellitus (E10-E14)	10,275	4.1	61.0
5	Cerebrovascular diseases (I60-I69)	9,701	3.9	57.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,982	3.2	47.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,380	2.2	32.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,611	1.9	27.4
9	Septicemia (A40-A41)	4,298	1.7	25.5
10	Influenza and pneumonia (J09-J18)	3,840	1.5	22.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,754	1.1	16.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,557	1.0	15.2
13	Alzheimer disease (G30)	2,116	0.9	12.6
14	Parkinson disease (G20-G21)	1,435	0.6	8.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,333	0.5	7.9
...	All other causes (residual)	39,199	15.8	232.8
All races and origins, both sexes, 70-74 years				
...	All causes	283,523	100.0	2,206.9
1	Malignant neoplasms (C00-C97)	86,852	30.6	676.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,164	21.9	483.9
3	Chronic lower respiratory diseases (J40-J47)	23,173	8.2	180.4
4	Cerebrovascular diseases (I60-I69)	12,989	4.6	101.1
5	Diabetes mellitus (E10-E14)	11,069	3.9	86.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,056	2.5	54.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,691	2.0	44.3
8	Alzheimer disease (G30)	5,165	1.8	40.2
9	Influenza and pneumonia (J09-J18)	4,944	1.7	38.5
10	Septicemia (A40-A41)	4,866	1.7	37.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	4,020	1.4	31.3
12	Parkinson disease (G20-G21)	3,226	1.1	25.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,102	1.1	24.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,063	0.7	16.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,775	0.6	13.8
...	All other causes (residual)	45,368	16.0	353.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 75-79 years				
...	All causes	307,498	100.0	3,517.8
1	Malignant neoplasms (C00-C97)	81,955	26.7	937.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,322	22.2	781.6
3	Chronic lower respiratory diseases (J40-J47)	25,285	8.2	289.3
4	Cerebrovascular diseases (I60-I69)	16,625	5.4	190.2
5	Alzheimer disease (G30)	11,408	3.7	130.5
6	Diabetes mellitus (E10-E14)	10,599	3.4	121.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,569	2.5	86.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,421	2.1	73.5
9	Influenza and pneumonia (J09-J18)	5,952	1.9	68.1
10	Parkinson disease (G20-G21)	5,443	1.8	62.3
11	Septicemia (A40-A41)	5,194	1.7	59.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,542	1.2	40.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,701	0.9	30.9
14	Pneumonitis due to solids and liquids (J69)	2,235	0.7	25.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,215	0.7	25.3
...	All other causes (residual)	52,032	16.9	595.2
All races and origins, both sexes, 80-84 years				
...	All causes	350,261	100.0	5,871.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,927	23.7	1,390.2
2	Malignant neoplasms (C00-C97)	73,962	21.1	1,239.9
3	Chronic lower respiratory diseases (J40-J47)	25,831	7.4	433.0
4	Cerebrovascular diseases (I60-I69)	22,072	6.3	370.0
5	Alzheimer disease (G30)	20,901	6.0	350.4
6	Diabetes mellitus (E10-E14)	10,104	2.9	169.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,092	2.6	152.4
8	Influenza and pneumonia (J09-J18)	7,850	2.2	131.6
9	Parkinson disease (G20-G21)	7,729	2.2	129.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,672	2.2	128.6
11	Septicemia (A40-A41)	5,454	1.6	91.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,669	1.3	78.3
13	Pneumonitis due to solids and liquids (J69)	3,001	0.9	50.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,520	0.7	42.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,909	0.5	32.0
...	All other causes (residual)	64,568	18.4	1,082.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 85-89 years				
...	All causes	388,121	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	102,358	26.4	---
2	Malignant neoplasms (C00-C97)	59,677	15.4	---
3	Alzheimer disease (G30)	31,483	8.1	---
4	Cerebrovascular diseases (I60-I69)	27,551	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	23,584	6.1	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,817	2.8	---
7	Influenza and pneumonia (J09-J18)	9,692	2.5	---
8	Diabetes mellitus (E10-E14)	8,986	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,992	2.1	---
10	Parkinson disease (G20-G21)	7,802	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,689	1.5	---
12	Septicemia (A40-A41)	5,552	1.4	---
13	Pneumonitis due to solids and liquids (J69)	3,971	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,638	0.7	---
15	Nutritional deficiencies (E40-E64)	1,521	0.4	---
...	All other causes (residual)	78,808	20.3	---
All races and origins, both sexes, 90-94 years				
...	All causes	319,443	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94,363	29.5	---
2	Malignant neoplasms (C00-C97)	33,148	10.4	---
3	Alzheimer disease (G30)	30,676	9.6	---
4	Cerebrovascular diseases (I60-I69)	24,047	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	15,558	4.9	---
6	Influenza and pneumonia (J09-J18)	9,070	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,931	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,246	2.0	---
9	Diabetes mellitus (E10-E14)	5,754	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,573	1.7	---
11	Parkinson disease (G20-G21)	4,320	1.4	---
12	Septicemia (A40-A41)	3,897	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,502	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,886	0.6	---
15	Nutritional deficiencies (E40-E64)	1,550	0.5	---
...	All other causes (residual)	70,922	22.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 95-99 years				
...	All causes	141,108	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,628	31.6	---
2	Alzheimer disease (G30)	15,120	10.7	---
3	Cerebrovascular diseases (I60-I69)	10,571	7.5	---
4	Malignant neoplasms (C00-C97)	9,518	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	5,370	3.8	---
6	Influenza and pneumonia (J09-J18)	4,445	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,786	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,892	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,572	1.8	---
10	Diabetes mellitus (E10-E14)	1,940	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,595	1.1	---
12	Septicemia (A40-A41)	1,500	1.1	---
13	Parkinson disease (G20-G21)	1,095	0.8	---
14	Nutritional deficiencies (E40-E64)	967	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	677	0.5	---
...	All other causes (residual)	34,432	24.4	---
All races and origins, both sexes, 100 years and over				
...	All causes	29,363	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,821	33.4	---
2	Alzheimer disease (G30)	3,238	11.0	---
3	Cerebrovascular diseases (I60-I69)	2,097	7.1	---
4	Malignant neoplasms (C00-C97)	1,178	4.0	---
5	Influenza and pneumonia (J09-J18)	1,069	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	815	2.8	---
7	Chronic lower respiratory diseases (J40-J47)	807	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	718	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	465	1.6	---
10	Diabetes mellitus (E10-E14)	293	1.0	---
10	Pneumonitis due to solids and liquids (J69)	293	1.0	---
12	Septicemia (A40-A41)	261	0.9	---
13	Nutritional deficiencies (E40-E64)	256	0.9	---
14	Atherosclerosis (I70)	163	0.6	---
15	Parkinson disease (G20-G21)	127	0.4	---
...	All other causes (residual)	7,762	26.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, all ages				
...	All causes	1,439,111	100.0	897.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	347,879	24.2	216.9
2	Malignant neoplasms (C00-C97)	315,147	21.9	196.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,722	7.6	68.4
4	Chronic lower respiratory diseases (J40-J47)	75,005	5.2	46.8
5	Cerebrovascular diseases (I60-I69)	61,645	4.3	38.4
6	Diabetes mellitus (E10-E14)	46,302	3.2	28.9
7	Alzheimer disease (G30)	37,325	2.6	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,782	2.6	22.9
9	Influenza and pneumonia (J09-J18)	26,558	1.8	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	26,451	1.8	16.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,744	1.8	16.0
12	Septicemia (A40-A41)	19,603	1.4	12.2
13	Parkinson disease (G20-G21)	19,388	1.3	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,749	1.1	9.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,524	1.1	9.7
...	All other causes (residual)	260,287	18.1	162.3
All races and origins, male, 1-4 years				
...	All causes	2,232	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	770	34.5	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	10.1	2.8
3	Malignant neoplasms (C00-C97)	184	8.2	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.0	0.8
6	Influenza and pneumonia (J09-J18)	63	2.8	0.8
7	Cerebrovascular diseases (I60-I69)	42	1.9	0.5
8	Septicemia (A40-A41)	26	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	19	0.9	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	15	0.7	*
14	Meningitis (G00,G03)	14	0.6	*
15	Anemias (D50-D64)	8	0.4	*
...	All other causes (residual)	579	25.9	7.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 5-9 years				
...	All causes	1,299	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	32.4	4.1
2	Malignant neoplasms (C00-C97)	199	15.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	8.0	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	6.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.4
6	Chronic lower respiratory diseases (J40-J47)	37	2.8	0.4
7	Influenza and pneumonia (J09-J18)	30	2.3	0.3
8	Cerebrovascular diseases (I60-I69)	20	1.5	0.2
9	Septicemia (A40-A41)	16	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.2	*
11	Anemias (D50-D64)	6	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
14	Meningitis (G00,G03)	4	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (residual)	294	22.6	2.8
All races and origins, male, 10-14 years				
...	All causes	1,970	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	28.6	5.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	17.7	3.3
3	Malignant neoplasms (C00-C97)	256	13.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	5.7	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	5.3	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	2.7	0.5
7	Chronic lower respiratory diseases (J40-J47)	47	2.4	0.4
8	Cerebrovascular diseases (I60-I69)	33	1.7	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Influenza and pneumonia (J09-J18)	19	1.0	*
11	Diabetes mellitus (E10-E14)	16	0.8	*
12	Septicemia (A40-A41)	15	0.8	*
13	Anemias (D50-D64)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
...	All other causes (residual)	359	18.2	3.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-19 years				
...	All causes	7,849	100.0	72.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,755	35.1	25.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,938	24.7	17.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,592	20.3	14.7
4	Malignant neoplasms (C00-C97)	332	4.2	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	2.7	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.2	0.8
7	Chronic lower respiratory diseases (J40-J47)	44	0.6	0.4
8	Diabetes mellitus (E10-E14)	41	0.5	0.4
9	Cerebrovascular diseases (I60-I69)	37	0.5	0.3
10	Influenza and pneumonia (J09-J18)	36	0.5	0.3
11	Legal intervention (Y35,Y89.0)	31	0.4	0.3
12	Septicemia (A40-A41)	21	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
13	Aortic aneurysm and dissection (I71)	13	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	10	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.1	*
...	All other causes (residual)	676	8.6	6.3
All races and origins, male, 20-24 years				
...	All causes	15,654	100.0	137.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,991	44.7	61.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,089	19.7	27.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,642	16.9	23.3
4	Malignant neoplasms (C00-C97)	501	3.2	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	411	2.6	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.7	1.0
7	Diabetes mellitus (E10-E14)	103	0.7	0.9
8	Influenza and pneumonia (J09-J18)	73	0.5	0.6
9	Chronic lower respiratory diseases (J40-J47)	71	0.5	0.6
10	Legal intervention (Y35,Y89.0)	59	0.4	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.3	0.4
12	Cerebrovascular diseases (I60-I69)	46	0.3	0.4
13	Septicemia (A40-A41)	41	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.2	0.3
15	Pneumonitis due to solids and liquids (J69)	24	0.2	0.2
...	All other causes (residual)	1,404	9.0	12.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 25-29 years				
...	All causes	20,385	100.0	171.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,520	46.7	80.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,302	16.2	27.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,643	13.0	22.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	830	4.1	7.0
5	Malignant neoplasms (C00-C97)	681	3.3	5.7
6	Diabetes mellitus (E10-E14)	198	1.0	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	157	0.8	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	153	0.8	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.7	1.1
10	Cerebrovascular diseases (I60-I69)	116	0.6	1.0
11	Legal intervention (Y35,Y89.0)	104	0.5	0.9
12	Chronic lower respiratory diseases (J40-J47)	94	0.5	0.8
13	Influenza and pneumonia (J09-J18)	91	0.4	0.8
14	Septicemia (A40-A41)	85	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.3	0.5
...	All other causes (residual)	2,218	10.9	18.6
All races and origins, male, 30-34 years				
...	All causes	21,764	100.0	196.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,481	43.6	85.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,046	14.0	27.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,964	9.0	17.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,619	7.4	14.6
5	Malignant neoplasms (C00-C97)	1,009	4.6	9.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	398	1.8	3.6
7	Diabetes mellitus (E10-E14)	273	1.3	2.5
8	Cerebrovascular diseases (I60-I69)	223	1.0	2.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	212	1.0	1.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	0.7	1.3
11	Influenza and pneumonia (J09-J18)	124	0.6	1.1
12	Septicemia (A40-A41)	115	0.5	1.0
13	Chronic lower respiratory diseases (J40-J47)	98	0.5	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.4	0.7
15	Legal intervention (Y35,Y89.0)	77	0.4	0.7
...	All other causes (residual)	2,900	13.3	26.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 35-39 years				
...	All causes	24,105	100.0	227.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,911	37.0	83.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,887	12.0	27.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,614	10.8	24.6
4	Malignant neoplasms (C00-C97)	1,780	7.4	16.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,583	6.6	14.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	751	3.1	7.1
7	Diabetes mellitus (E10-E14)	508	2.1	4.8
8	Cerebrovascular diseases (I60-I69)	398	1.7	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	281	1.2	2.6
10	Influenza and pneumonia (J09-J18)	195	0.8	1.8
11	Septicemia (A40-A41)	192	0.8	1.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	0.6	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	0.6	1.3
14	Chronic lower respiratory diseases (J40-J47)	123	0.5	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	0.5	1.0
...	All other causes (residual)	3,487	14.5	32.8
All races and origins, male, 40-44 years				
...	All causes	26,687	100.0	273.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,332	27.5	75.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,460	16.7	45.7
3	Malignant neoplasms (C00-C97)	2,945	11.0	30.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,709	10.2	27.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,102	4.1	11.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,093	4.1	11.2
7	Diabetes mellitus (E10-E14)	790	3.0	8.1
8	Cerebrovascular diseases (I60-I69)	636	2.4	6.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	286	1.1	2.9
10	Septicemia (A40-A41)	264	1.0	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	251	0.9	2.6
12	Influenza and pneumonia (J09-J18)	242	0.9	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	212	0.8	2.2
14	Chronic lower respiratory diseases (J40-J47)	200	0.7	2.1
15	Aortic aneurysm and dissection (I71)	139	0.5	1.4
...	All other causes (residual)	4,026	15.1	41.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-49 years				
...	All causes	40,223	100.0	387.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,081	20.1	77.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,040	20.0	77.4
3	Malignant neoplasms (C00-C97)	5,986	14.9	57.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,122	7.8	30.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,094	5.2	20.2
6	Diabetes mellitus (E10-E14)	1,485	3.7	14.3
7	Cerebrovascular diseases (I60-I69)	1,110	2.8	10.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	935	2.3	9.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	474	1.2	4.6
10	Chronic lower respiratory diseases (J40-J47)	467	1.2	4.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	456	1.1	4.4
12	Septicemia (A40-A41)	454	1.1	4.4
13	Influenza and pneumonia (J09-J18)	401	1.0	3.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	394	1.0	3.8
15	Aortic aneurysm and dissection (I71)	235	0.6	2.3
...	All other causes (residual)	6,489	16.1	62.5
All races and origins, male, 50-54 years				
...	All causes	63,581	100.0	604.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,727	23.2	140.0
2	Malignant neoplasms (C00-C97)	12,912	20.3	122.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,837	13.9	84.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,295	5.2	31.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,209	5.0	30.5
6	Diabetes mellitus (E10-E14)	2,535	4.0	24.1
7	Cerebrovascular diseases (I60-I69)	1,900	3.0	18.1
8	Chronic lower respiratory diseases (J40-J47)	1,365	2.1	13.0
9	Septicemia (A40-A41)	869	1.4	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	814	1.3	7.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	802	1.3	7.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	711	1.1	6.8
13	Influenza and pneumonia (J09-J18)	675	1.1	6.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	667	1.0	6.3
15	Viral hepatitis (B15-B19)	438	0.7	4.2
...	All other causes (residual)	9,825	15.5	93.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 55-59 years				
...	All causes	98,387	100.0	919.5
1	Malignant neoplasms (C00-C97)	25,165	25.6	235.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,803	24.2	222.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,159	9.3	85.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,572	4.6	42.7
5	Diabetes mellitus (E10-E14)	3,985	4.1	37.2
6	Chronic lower respiratory diseases (J40-J47)	3,345	3.4	31.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,327	3.4	31.1
8	Cerebrovascular diseases (I60-I69)	2,981	3.0	27.9
9	Septicemia (A40-A41)	1,337	1.4	12.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,323	1.3	12.4
11	Influenza and pneumonia (J09-J18)	1,161	1.2	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,057	1.1	9.9
13	Viral hepatitis (B15-B19)	805	0.8	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	733	0.7	6.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	680	0.7	6.4
...	All other causes (residual)	14,954	15.2	139.8
All races and origins, male, 60-64 years				
...	All causes	126,948	100.0	1,328.3
1	Malignant neoplasms (C00-C97)	37,215	29.3	389.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,532	24.8	329.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,101	5.6	74.3
4	Chronic lower respiratory diseases (J40-J47)	6,079	4.8	63.6
5	Diabetes mellitus (E10-E14)	5,297	4.2	55.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,651	3.7	48.7
7	Cerebrovascular diseases (I60-I69)	4,351	3.4	45.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,713	2.1	28.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,876	1.5	19.6
10	Septicemia (A40-A41)	1,823	1.4	19.1
11	Influenza and pneumonia (J09-J18)	1,665	1.3	17.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,416	1.1	14.8
13	Viral hepatitis (B15-B19)	934	0.7	9.8
14	Pneumonitis due to solids and liquids (J69)	604	0.5	6.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	581	0.5	6.1
...	All other causes (residual)	19,110	15.1	200.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 65-69 years				
...	All causes	145,258	100.0	1,831.8
1	Malignant neoplasms (C00-C97)	45,349	31.2	571.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,958	24.8	453.5
3	Chronic lower respiratory diseases (J40-J47)	8,587	5.9	108.3
4	Diabetes mellitus (E10-E14)	6,143	4.2	77.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,402	3.7	68.1
6	Cerebrovascular diseases (I60-I69)	5,298	3.6	66.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,652	2.5	46.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,563	1.8	32.3
9	Septicemia (A40-A41)	2,225	1.5	28.1
10	Influenza and pneumonia (J09-J18)	2,138	1.5	27.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,967	1.4	24.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,585	1.1	20.0
13	Alzheimer disease (G30)	921	0.6	11.6
14	Parkinson disease (G20-G21)	906	0.6	11.4
15	Pneumonitis due to solids and liquids (J69)	822	0.6	10.4
...	All other causes (residual)	21,742	15.0	274.2
All races and origins, male, 70-74 years				
...	All causes	158,673	100.0	2,668.0
1	Malignant neoplasms (C00-C97)	47,733	30.1	802.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,765	24.4	651.8
3	Chronic lower respiratory diseases (J40-J47)	11,596	7.3	195.0
4	Cerebrovascular diseases (I60-I69)	6,734	4.2	113.2
5	Diabetes mellitus (E10-E14)	6,459	4.1	108.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,395	2.8	73.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,079	1.9	51.8
8	Influenza and pneumonia (J09-J18)	2,644	1.7	44.5
9	Septicemia (A40-A41)	2,410	1.5	40.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,407	1.5	40.5
11	Parkinson disease (G20-G21)	2,166	1.4	36.4
12	Alzheimer disease (G30)	2,164	1.4	36.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,671	1.1	28.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,651	1.0	27.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,055	0.7	17.7
...	All other causes (residual)	23,744	15.0	399.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 75-79 years				
...	All causes	163,505	100.0	4,193.7
1	Malignant neoplasms (C00-C97)	44,293	27.1	1,136.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,719	24.3	1,018.7
3	Chronic lower respiratory diseases (J40-J47)	12,278	7.5	314.9
4	Cerebrovascular diseases (I60-I69)	8,008	4.9	205.4
5	Diabetes mellitus (E10-E14)	5,881	3.6	150.8
6	Alzheimer disease (G30)	4,426	2.7	113.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,403	2.7	112.9
8	Parkinson disease (G20-G21)	3,542	2.2	90.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,319	2.0	85.1
10	Influenza and pneumonia (J09-J18)	3,138	1.9	80.5
11	Septicemia (A40-A41)	2,508	1.5	64.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,749	1.1	44.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,557	1.0	39.9
14	Pneumonitis due to solids and liquids (J69)	1,360	0.8	34.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,303	0.8	33.4
...	All other causes (residual)	26,021	15.9	667.4
All races and origins, male, 80-84 years				
...	All causes	173,166	100.0	6,901.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,019	25.4	1,754.4
2	Malignant neoplasms (C00-C97)	39,056	22.6	1,556.6
3	Chronic lower respiratory diseases (J40-J47)	11,861	6.8	472.7
4	Cerebrovascular diseases (I60-I69)	9,516	5.5	379.3
5	Alzheimer disease (G30)	7,672	4.4	305.8
6	Diabetes mellitus (E10-E14)	5,330	3.1	212.4
7	Parkinson disease (G20-G21)	4,919	2.8	196.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,864	2.8	193.9
9	Influenza and pneumonia (J09-J18)	3,918	2.3	156.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,904	2.3	155.6
11	Septicemia (A40-A41)	2,560	1.5	102.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,025	1.2	80.7
13	Pneumonitis due to solids and liquids (J69)	1,748	1.0	69.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,356	0.8	54.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	0.6	39.5
...	All other causes (residual)	29,428	17.0	1,172.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 85-89 years				
...	All causes	171,951	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,935	27.9	---
2	Malignant neoplasms (C00-C97)	29,984	17.4	---
3	Chronic lower respiratory diseases (J40-J47)	10,462	6.1	---
4	Alzheimer disease (G30)	10,311	6.0	---
5	Cerebrovascular diseases (I60-I69)	10,270	6.0	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,219	3.0	---
7	Parkinson disease (G20-G21)	4,588	2.7	---
8	Influenza and pneumonia (J09-J18)	4,434	2.6	---
9	Diabetes mellitus (E10-E14)	4,239	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,952	2.3	---
11	Septicemia (A40-A41)	2,480	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,228	1.3	---
13	Pneumonitis due to solids and liquids (J69)	2,222	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,401	0.8	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	753	0.4	---
...	All other causes (residual)	31,473	18.3	---
All races and origins, male, 90-94 years				
...	All causes	117,273	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,033	31.6	---
2	Malignant neoplasms (C00-C97)	15,372	13.1	---
3	Alzheimer disease (G30)	8,030	6.8	---
4	Cerebrovascular diseases (I60-I69)	7,099	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	6,304	5.4	---
6	Influenza and pneumonia (J09-J18)	3,652	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,459	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,738	2.3	---
9	Diabetes mellitus (E10-E14)	2,305	2.0	---
10	Parkinson disease (G20-G21)	2,236	1.9	---
11	Pneumonitis due to solids and liquids (J69)	1,794	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,733	1.5	---
13	Septicemia (A40-A41)	1,522	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	821	0.7	---
15	Nutritional deficiencies (E40-E64)	481	0.4	---
...	All other causes (residual)	22,694	19.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 95-99 years				
...	All causes	40,205	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,826	34.4	---
2	Malignant neoplasms (C00-C97)	3,786	9.4	---
3	Alzheimer disease (G30)	2,850	7.1	---
4	Cerebrovascular diseases (I60-I69)	2,446	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	1,737	4.3	---
6	Influenza and pneumonia (J09-J18)	1,498	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,167	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	955	2.4	---
9	Pneumonitis due to solids and liquids (J69)	709	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	686	1.7	---
11	Diabetes mellitus (E10-E14)	636	1.6	---
12	Septicemia (A40-A41)	495	1.2	---
13	Parkinson disease (G20-G21)	463	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	258	0.6	---
15	Nutritional deficiencies (E40-E64)	257	0.6	---
...	All other causes (residual)	8,436	21.0	---
All races and origins, male, 100 years and over				
...	All causes	5,437	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,993	36.7	---
2	Alzheimer disease (G30)	383	7.0	---
3	Malignant neoplasms (C00-C97)	377	6.9	---
4	Cerebrovascular diseases (I60-I69)	322	5.9	---
5	Influenza and pneumonia (J09-J18)	262	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	179	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	2.5	---
9	Pneumonitis due to solids and liquids (J69)	108	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.9	---
11	Diabetes mellitus (E10-E14)	71	1.3	---
12	Septicemia (A40-A41)	66	1.2	---
13	Parkinson disease (G20-G21)	52	1.0	---
14	Nutritional deficiencies (E40-E64)	35	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	---
...	All other causes (residual)	1,163	21.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,374,392	100.0	831.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	299,578	21.8	181.2
2	Malignant neoplasms (C00-C97)	283,961	20.7	171.8
3	Chronic lower respiratory diseases (J40-J47)	85,196	6.2	51.5
4	Cerebrovascular diseases (I60-I69)	84,738	6.2	51.3
5	Alzheimer disease (G30)	84,079	6.1	50.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,214	4.4	36.4
7	Diabetes mellitus (E10-E14)	37,262	2.7	22.5
8	Influenza and pneumonia (J09-J18)	29,114	2.1	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,889	1.8	15.1
10	Septicemia (A40-A41)	21,319	1.6	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,567	1.4	11.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	15,292	1.1	9.3
13	Parkinson disease (G20-G21)	12,575	0.9	7.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,391	0.8	6.3
15	Pneumonitis due to solids and liquids (J69)	8,758	0.6	5.3
...	All other causes (residual)	297,459	21.6	179.9
All races and origins, female, 1-4 years				
...	All causes	1,648	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	30.2	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	12.1	2.5
3	Malignant neoplasms (C00-C97)	141	8.6	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	8.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.6	0.8
6	Influenza and pneumonia (J09-J18)	41	2.5	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	24	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	23	1.4	0.3
10	Septicemia (A40-A41)	22	1.3	0.3
11	Anemias (D50-D64)	16	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Meningitis (G00,G03)	5	0.3	*
...	All other causes (residual)	431	26.2	5.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-9 years				
...	All causes	1,055	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	297	28.2	3.0
2	Malignant neoplasms (C00-C97)	219	20.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	8.0	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	6.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.1	0.3
6	Influenza and pneumonia (J09-J18)	32	3.0	0.3
7	Chronic lower respiratory diseases (J40-J47)	22	2.1	0.2
8	Cerebrovascular diseases (I60-I69)	21	2.0	0.2
9	Septicemia (A40-A41)	17	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
12	Anemias (D50-D64)	4	0.4	*
12	Diabetes mellitus (E10-E14)	4	0.4	*
12	Diseases of appendix (K35-K38)	4	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
...	All other causes (residual)	222	21.0	2.2
All races and origins, female, 10-14 years				
...	All causes	1,247	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	23.7	2.9
2	Malignant neoplasms (C00-C97)	181	14.5	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	13.6	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	7.0	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	5.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.0	0.5
7	Influenza and pneumonia (J09-J18)	32	2.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	28	2.2	0.3
9	Cerebrovascular diseases (I60-I69)	23	1.8	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
11	Diabetes mellitus (E10-E14)	10	0.8	*
12	Septicemia (A40-A41)	9	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	Anemias (D50-D64)	5	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (residual)	268	21.5	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 15-19 years				
...	All causes	3,037	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,193	39.3	11.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	553	18.2	5.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	243	8.0	2.4
4	Malignant neoplasms (C00-C97)	228	7.5	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.2	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	2.5	0.7
7	Diabetes mellitus (E10-E14)	35	1.2	0.3
8	Influenza and pneumonia (J09-J18)	33	1.1	0.3
9	Cerebrovascular diseases (I60-I69)	29	1.0	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.9	0.3
11	Chronic lower respiratory diseases (J40-J47)	21	0.7	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	0.2
13	Septicemia (A40-A41)	17	0.6	*
14	Anemias (D50-D64)	14	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
...	All other causes (residual)	437	14.4	4.2
All races and origins, female, 20-24 years				
...	All causes	5,485	100.0	50.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,502	45.6	23.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	672	12.3	6.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	7.8	4.0
4	Malignant neoplasms (C00-C97)	313	5.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	3.6	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	140	2.6	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	1.3	0.6
8	Diabetes mellitus (E10-E14)	69	1.3	0.6
9	Chronic lower respiratory diseases (J40-J47)	52	0.9	0.5
10	Influenza and pneumonia (J09-J18)	48	0.9	0.4
11	Septicemia (A40-A41)	44	0.8	0.4
12	Cerebrovascular diseases (I60-I69)	43	0.8	0.4
13	Anemias (D50-D64)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.3	*
...	All other causes (residual)	850	15.5	7.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-29 years				
...	All causes	7,891	100.0	68.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,214	40.7	28.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	767	9.7	6.7
3	Malignant neoplasms (C00-C97)	620	7.9	5.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	440	5.6	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	5.5	3.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	253	3.2	2.2
7	Diabetes mellitus (E10-E14)	146	1.9	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.3	0.9
9	Cerebrovascular diseases (I60-I69)	100	1.3	0.9
10	Septicemia (A40-A41)	87	1.1	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	1.1	0.7
12	Influenza and pneumonia (J09-J18)	65	0.8	0.6
13	Chronic lower respiratory diseases (J40-J47)	59	0.7	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	58	0.7	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.5	0.4
...	All other causes (residual)	1,416	17.9	12.3
All races and origins, female, 30-34 years				
...	All causes	10,175	100.0	93.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,454	33.9	31.7
2	Malignant neoplasms (C00-C97)	1,306	12.8	12.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	833	8.2	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	796	7.8	7.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	441	4.3	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	263	2.6	2.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	259	2.5	2.4
8	Diabetes mellitus (E10-E14)	206	2.0	1.9
9	Cerebrovascular diseases (I60-I69)	154	1.5	1.4
10	Influenza and pneumonia (J09-J18)	125	1.2	1.1
11	Septicemia (A40-A41)	105	1.0	1.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	86	0.8	0.8
13	Chronic lower respiratory diseases (J40-J47)	81	0.8	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.7	0.7
...	All other causes (residual)	1,918	18.9	17.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 35-39 years				
...	All causes	12,796	100.0	120.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,440	26.9	32.4
2	Malignant neoplasms (C00-C97)	2,324	18.2	21.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,295	10.1	12.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	835	6.5	7.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	476	3.7	4.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	3.1	3.8
7	Diabetes mellitus (E10-E14)	326	2.5	3.1
8	Cerebrovascular diseases (I60-I69)	287	2.2	2.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	217	1.7	2.0
10	Septicemia (A40-A41)	174	1.4	1.6
11	Influenza and pneumonia (J09-J18)	130	1.0	1.2
12	Chronic lower respiratory diseases (J40-J47)	126	1.0	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.0	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	0.9	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	0.8	1.0
...	All other causes (residual)	2,422	18.9	22.8
All races and origins, female, 40-44 years				
...	All causes	16,208	100.0	163.9
1	Malignant neoplasms (C00-C97)	3,851	23.8	38.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,145	19.4	31.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,032	12.5	20.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	904	5.6	9.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	671	4.1	6.8
6	Diabetes mellitus (E10-E14)	494	3.0	5.0
7	Cerebrovascular diseases (I60-I69)	490	3.0	5.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	274	1.7	2.8
9	Chronic lower respiratory diseases (J40-J47)	264	1.6	2.7
10	Septicemia (A40-A41)	224	1.4	2.3
11	Influenza and pneumonia (J09-J18)	215	1.3	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	1.1	1.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	141	0.9	1.4
14	Pregnancy, childbirth and the puerperium (O00-O99)	111	0.7	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	0.6	0.9
...	All other causes (residual)	3,114	19.2	31.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 45-49 years				
...	All causes	25,475	100.0	240.6
1	Malignant neoplasms (C00-C97)	7,122	28.0	67.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,593	14.1	33.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,566	14.0	33.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,128	4.4	10.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,108	4.3	10.5
6	Diabetes mellitus (E10-E14)	913	3.6	8.6
7	Cerebrovascular diseases (I60-I69)	836	3.3	7.9
8	Chronic lower respiratory diseases (J40-J47)	550	2.2	5.2
9	Septicemia (A40-A41)	422	1.7	4.0
10	Influenza and pneumonia (J09-J18)	364	1.4	3.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	1.3	3.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	1.1	2.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	210	0.8	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	0.6	1.4
...	All other causes (residual)	4,720	18.5	44.6
All races and origins, female, 50-54 years				
...	All causes	40,863	100.0	375.5
1	Malignant neoplasms (C00-C97)	13,246	32.4	121.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,284	15.4	57.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,991	9.8	36.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,795	4.4	16.5
5	Chronic lower respiratory diseases (J40-J47)	1,593	3.9	14.6
6	Diabetes mellitus (E10-E14)	1,476	3.6	13.6
7	Cerebrovascular diseases (I60-I69)	1,352	3.3	12.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,122	2.7	10.3
9	Septicemia (A40-A41)	696	1.7	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	577	1.4	5.3
11	Influenza and pneumonia (J09-J18)	573	1.4	5.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	390	1.0	3.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	0.6	2.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	232	0.6	2.1
15	Viral hepatitis (B15-B19)	215	0.5	2.0
...	All other causes (residual)	7,060	17.3	64.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 55-59 years				
...	All causes	63,711	100.0	563.4
1	Malignant neoplasms (C00-C97)	22,481	35.3	198.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,425	16.4	92.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,983	6.3	35.2
4	Chronic lower respiratory diseases (J40-J47)	3,618	5.7	32.0
5	Diabetes mellitus (E10-E14)	2,371	3.7	21.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,279	3.6	20.2
7	Cerebrovascular diseases (I60-I69)	2,262	3.6	20.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,092	1.7	9.7
9	Septicemia (A40-A41)	1,077	1.7	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	980	1.5	8.7
11	Influenza and pneumonia (J09-J18)	918	1.4	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	627	1.0	5.5
13	Viral hepatitis (B15-B19)	364	0.6	3.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	295	0.5	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	260	0.4	2.3
...	All other causes (residual)	10,679	16.8	94.4
All races and origins, female, 60-64 years				
...	All causes	82,960	100.0	795.4
1	Malignant neoplasms (C00-C97)	29,949	36.1	287.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,342	17.3	137.5
3	Chronic lower respiratory diseases (J40-J47)	5,625	6.8	53.9
4	Diabetes mellitus (E10-E14)	3,251	3.9	31.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,165	3.8	30.3
6	Cerebrovascular diseases (I60-I69)	3,114	3.8	29.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,235	2.7	21.4
8	Septicemia (A40-A41)	1,601	1.9	15.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,492	1.8	14.3
10	Influenza and pneumonia (J09-J18)	1,301	1.6	12.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	891	1.1	8.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	850	1.0	8.1
13	Alzheimer disease (G30)	482	0.6	4.6
14	Viral hepatitis (B15-B19)	414	0.5	4.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	373	0.4	3.6
...	All other causes (residual)	13,875	16.7	133.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65-69 years				
...	All causes	102,829	100.0	1,154.5
1	Malignant neoplasms (C00-C97)	36,257	35.3	407.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,511	18.0	207.8
3	Chronic lower respiratory diseases (J40-J47)	7,944	7.7	89.2
4	Cerebrovascular diseases (I60-I69)	4,403	4.3	49.4
5	Diabetes mellitus (E10-E14)	4,132	4.0	46.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,580	2.5	29.0
7	Septicemia (A40-A41)	2,073	2.0	23.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,048	2.0	23.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,728	1.7	19.4
10	Influenza and pneumonia (J09-J18)	1,702	1.7	19.1
11	Alzheimer disease (G30)	1,195	1.2	13.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,169	1.1	13.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	590	0.6	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	531	0.5	6.0
15	Parkinson disease (G20-G21)	529	0.5	5.9
...	All other causes (residual)	17,437	17.0	195.8
All races and origins, female, 70-74 years				
...	All causes	124,850	100.0	1,809.5
1	Malignant neoplasms (C00-C97)	39,119	31.3	567.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,399	18.7	339.1
3	Chronic lower respiratory diseases (J40-J47)	11,577	9.3	167.8
4	Cerebrovascular diseases (I60-I69)	6,255	5.0	90.7
5	Diabetes mellitus (E10-E14)	4,610	3.7	66.8
6	Alzheimer disease (G30)	3,001	2.4	43.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,661	2.1	38.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,612	2.1	37.9
9	Septicemia (A40-A41)	2,456	2.0	35.6
10	Influenza and pneumonia (J09-J18)	2,300	1.8	33.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,613	1.3	23.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,451	1.2	21.0
13	Parkinson disease (G20-G21)	1,060	0.8	15.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	720	0.6	10.4
15	Pneumonitis due to solids and liquids (J69)	642	0.5	9.3
...	All other causes (residual)	21,374	17.1	309.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 75-79 years				
...	All causes	143,993	100.0	2,973.6
1	Malignant neoplasms (C00-C97)	37,662	26.2	777.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,603	19.9	590.7
3	Chronic lower respiratory diseases (J40-J47)	13,007	9.0	268.6
4	Cerebrovascular diseases (I60-I69)	8,617	6.0	177.9
5	Alzheimer disease (G30)	6,982	4.8	144.2
6	Diabetes mellitus (E10-E14)	4,718	3.3	97.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,166	2.2	65.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,102	2.2	64.1
9	Influenza and pneumonia (J09-J18)	2,814	2.0	58.1
10	Septicemia (A40-A41)	2,686	1.9	55.5
11	Parkinson disease (G20-G21)	1,901	1.3	39.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,793	1.2	37.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,144	0.8	23.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	952	0.7	19.7
15	Pneumonitis due to solids and liquids (J69)	875	0.6	18.1
...	All other causes (residual)	25,971	18.0	536.3
All races and origins, female, 80-84 years				
...	All causes	177,095	100.0	5,123.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,908	22.0	1,125.7
2	Malignant neoplasms (C00-C97)	34,906	19.7	1,009.9
3	Chronic lower respiratory diseases (J40-J47)	13,970	7.9	404.2
4	Alzheimer disease (G30)	13,229	7.5	382.8
5	Cerebrovascular diseases (I60-I69)	12,556	7.1	363.3
6	Diabetes mellitus (E10-E14)	4,774	2.7	138.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,228	2.4	122.3
8	Influenza and pneumonia (J09-J18)	3,932	2.2	113.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,768	2.1	109.0
10	Septicemia (A40-A41)	2,894	1.6	83.7
11	Parkinson disease (G20-G21)	2,810	1.6	81.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,644	1.5	76.5
13	Pneumonitis due to solids and liquids (J69)	1,253	0.7	36.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,164	0.7	33.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	920	0.5	26.6
...	All other causes (residual)	35,139	19.8	1,016.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 85-89 years				
...	All causes	216,170	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,423	25.2	---
2	Malignant neoplasms (C00-C97)	29,693	13.7	---
3	Alzheimer disease (G30)	21,172	9.8	---
4	Cerebrovascular diseases (I60-I69)	17,281	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	13,122	6.1	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,598	2.6	---
7	Influenza and pneumonia (J09-J18)	5,258	2.4	---
8	Diabetes mellitus (E10-E14)	4,747	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,040	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,461	1.6	---
11	Parkinson disease (G20-G21)	3,214	1.5	---
12	Septicemia (A40-A41)	3,072	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,749	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,237	0.6	---
15	Nutritional deficiencies (E40-E64)	993	0.5	---
...	All other causes (residual)	47,110	21.8	---
All races and origins, female, 90-94 years				
...	All causes	202,170	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,330	28.4	---
2	Alzheimer disease (G30)	22,646	11.2	---
3	Malignant neoplasms (C00-C97)	17,776	8.8	---
4	Cerebrovascular diseases (I60-I69)	16,948	8.4	---
5	Chronic lower respiratory diseases (J40-J47)	9,254	4.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,472	2.7	---
7	Influenza and pneumonia (J09-J18)	5,418	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,840	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,508	1.7	---
10	Diabetes mellitus (E10-E14)	3,449	1.7	---
11	Septicemia (A40-A41)	2,375	1.2	---
12	Parkinson disease (G20-G21)	2,084	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,708	0.8	---
14	Nutritional deficiencies (E40-E64)	1,069	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,065	0.5	---
...	All other causes (residual)	48,228	23.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 95-99 years				
...	All causes	100,903	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,802	30.5	---
2	Alzheimer disease (G30)	12,270	12.2	---
3	Cerebrovascular diseases (I60-I69)	8,125	8.1	---
4	Malignant neoplasms (C00-C97)	5,732	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,633	3.6	---
6	Influenza and pneumonia (J09-J18)	2,947	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,619	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,206	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,617	1.6	---
10	Diabetes mellitus (E10-E14)	1,304	1.3	---
11	Septicemia (A40-A41)	1,005	1.0	---
12	Pneumonitis due to solids and liquids (J69)	886	0.9	---
13	Nutritional deficiencies (E40-E64)	710	0.7	---
14	Parkinson disease (G20-G21)	632	0.6	---
15	Atherosclerosis (I70)	476	0.5	---
...	All other causes (residual)	25,939	25.7	---
All races and origins, female, 100 years and over				
...	All causes	23,926	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,828	32.7	---
2	Alzheimer disease (G30)	2,855	11.9	---
3	Cerebrovascular diseases (I60-I69)	1,775	7.4	---
4	Influenza and pneumonia (J09-J18)	807	3.4	---
5	Malignant neoplasms (C00-C97)	801	3.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	710	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	628	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	562	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	331	1.4	---
10	Diabetes mellitus (E10-E14)	222	0.9	---
11	Nutritional deficiencies (E40-E64)	221	0.9	---
12	Septicemia (A40-A41)	195	0.8	---
13	Pneumonitis due to solids and liquids (J69)	185	0.8	---
14	Atherosclerosis (I70)	140	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.3	---
...	All other causes (residual)	6,586	27.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	197,249	100.0	334.6
1	Malignant neoplasms (C00-C97)	40,550	20.6	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,402	20.0	66.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,757	8.5	28.4
4	Cerebrovascular diseases (I60-I69)	10,775	5.5	18.3
5	Diabetes mellitus (E10-E14)	9,216	4.7	15.6
6	Alzheimer disease (G30)	7,383	3.7	12.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,286	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,599	2.8	9.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,933	2.0	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,928	2.0	6.7
11	Influenza and pneumonia (J09-J18)	3,851	2.0	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,186	1.6	5.4
13	Septicemia (A40-A41)	2,869	1.5	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,799	1.4	4.7
15	Certain conditions originating in the perinatal period (P00-P96)	2,409	1.2	4.1
...	All other causes (residual)	38,306	19.4	65.0
Hispanic, both sexes, 1-4 years				
...	All causes	805	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	29.1	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	13.9	2.7
3	Malignant neoplasms (C00-C97)	92	11.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	6.2	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.2	0.6
6	Influenza and pneumonia (J09-J18)	24	3.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Cerebrovascular diseases (I60-I69)	12	1.5	*
9	Septicemia (A40-A41)	11	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	6	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.6	*
14	Meningitis (G00,G03)	4	0.5	*
15	Anemias (D50-D64)	3	0.4	*
...	All other causes (residual)	196	24.3	4.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	521	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	28.4	2.8
2	Malignant neoplasms (C00-C97)	100	19.2	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	8.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	5.0	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.0	0.4
6	Septicemia (A40-A41)	15	2.9	*
7	Influenza and pneumonia (J09-J18)	13	2.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	8	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Diseases of appendix (K35-K38)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
...	All other causes (residual)	115	22.1	2.2
Hispanic, both sexes, 10-14 years				
...	All causes	644	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	23.9	3.0
2	Malignant neoplasms (C00-C97)	122	18.9	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	14.0	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	5.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	5.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.6	*
7	Cerebrovascular diseases (I60-I69)	16	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	13	2.0	*
9	Influenza and pneumonia (J09-J18)	8	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
11	Septicemia (A40-A41)	3	0.5	*
11	Diabetes mellitus (E10-E14)	3	0.5	*
13	Anemias (D50-D64)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	132	20.5	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,175	100.0	44.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	37.6	16.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	422	19.4	8.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	381	17.5	7.8
4	Malignant neoplasms (C00-C97)	152	7.0	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	2.0	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.6	0.7
7	Cerebrovascular diseases (I60-I69)	17	0.8	*
8	Influenza and pneumonia (J09-J18)	16	0.7	*
9	Legal intervention (Y35,Y89.0)	13	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
12	Diabetes mellitus (E10-E14)	10	0.5	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.4	*
14	Septicemia (A40-A41)	6	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (residual)	226	10.4	4.6
Hispanic, both sexes, 20-24 years				
...	All causes	3,679	100.0	76.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,668	45.3	34.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	590	16.0	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	558	15.2	11.5
4	Malignant neoplasms (C00-C97)	219	6.0	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.0	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	26	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	24	0.7	0.5
9	Diabetes mellitus (E10-E14)	19	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
11	Septicemia (A40-A41)	17	0.5	*
12	Influenza and pneumonia (J09-J18)	16	0.4	*
13	Legal intervention (Y35,Y89.0)	14	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.3	*
15	Anemias (D50-D64)	7	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.2	*
...	All other causes (residual)	362	9.8	7.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	4,166	100.0	86.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,803	43.3	37.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	525	12.6	11.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	12.5	10.8
4	Malignant neoplasms (C00-C97)	272	6.5	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	4.1	3.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.4	1.2
7	Cerebrovascular diseases (I60-I69)	49	1.2	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	1.0	0.9
9	Diabetes mellitus (E10-E14)	40	1.0	0.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	38	0.9	0.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.9	0.8
12	Septicemia (A40-A41)	29	0.7	0.6
13	Legal intervention (Y35,Y89.0)	25	0.6	0.5
14	Influenza and pneumonia (J09-J18)	24	0.6	0.5
14	Chronic lower respiratory diseases (J40-J47)	24	0.6	0.5
...	All other causes (residual)	511	12.3	10.7
Hispanic, both sexes, 30-34 years				
...	All causes	4,326	100.0	95.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,576	36.4	34.8
2	Malignant neoplasms (C00-C97)	456	10.5	10.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	439	10.1	9.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	9.0	8.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	7.5	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	116	2.7	2.6
7	Cerebrovascular diseases (I60-I69)	76	1.8	1.7
8	Diabetes mellitus (E10-E14)	73	1.7	1.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	48	1.1	1.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.0	0.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.7	0.7
12	Influenza and pneumonia (J09-J18)	31	0.7	0.7
13	Septicemia (A40-A41)	30	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	24	0.6	0.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.6	0.5
...	All other causes (residual)	645	14.9	14.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	5,014	100.0	113.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,568	31.3	35.5
2	Malignant neoplasms (C00-C97)	716	14.3	16.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	451	9.0	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	7.7	8.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	344	6.9	7.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	263	5.2	6.0
7	Cerebrovascular diseases (I60-I69)	131	2.6	3.0
8	Diabetes mellitus (E10-E14)	94	1.9	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	77	1.5	1.7
10	Septicemia (A40-A41)	55	1.1	1.2
11	Influenza and pneumonia (J09-J18)	43	0.9	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	35	0.7	0.8
13	Chronic lower respiratory diseases (J40-J47)	30	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.5	0.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	0.5	0.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.5	0.6
...	All other causes (residual)	745	14.9	16.9
Hispanic, both sexes, 40-44 years				
...	All causes	5,724	100.0	141.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,336	23.3	32.9
2	Malignant neoplasms (C00-C97)	1,061	18.5	26.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	714	12.5	17.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	364	6.4	9.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	318	5.6	7.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	4.1	5.8
7	Cerebrovascular diseases (I60-I69)	220	3.8	5.4
8	Diabetes mellitus (E10-E14)	186	3.2	4.6
9	Septicemia (A40-A41)	69	1.2	1.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	68	1.2	1.7
11	Influenza and pneumonia (J09-J18)	64	1.1	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.0	1.4
13	Chronic lower respiratory diseases (J40-J47)	37	0.6	0.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.6	0.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.5	0.6
...	All other causes (residual)	937	16.4	23.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	7,738	100.0	209.4
1	Malignant neoplasms (C00-C97)	1,670	21.6	45.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,361	17.6	36.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,106	14.3	29.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	623	8.1	16.9
5	Diabetes mellitus (E10-E14)	383	4.9	10.4
6	Cerebrovascular diseases (I60-I69)	307	4.0	8.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	299	3.9	8.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	186	2.4	5.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	1.5	3.1
10	Influenza and pneumonia (J09-J18)	105	1.4	2.8
11	Septicemia (A40-A41)	104	1.3	2.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	92	1.2	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	0.9	1.8
14	Chronic lower respiratory diseases (J40-J47)	58	0.7	1.6
15	Viral hepatitis (B15-B19)	56	0.7	1.5
...	All other causes (residual)	1,209	15.6	32.7
Hispanic, both sexes, 50-54 years				
...	All causes	10,671	100.0	333.4
1	Malignant neoplasms (C00-C97)	2,739	25.7	85.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,845	17.3	57.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,282	12.0	40.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	842	7.9	26.3
5	Diabetes mellitus (E10-E14)	551	5.2	17.2
6	Cerebrovascular diseases (I60-I69)	426	4.0	13.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	2.7	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	1.7	5.6
9	Septicemia (A40-A41)	171	1.6	5.3
10	Influenza and pneumonia (J09-J18)	148	1.4	4.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	1.2	4.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	131	1.2	4.1
13	Viral hepatitis (B15-B19)	125	1.2	3.9
14	Chronic lower respiratory diseases (J40-J47)	110	1.0	3.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	0.9	3.2
...	All other causes (residual)	1,605	15.0	50.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	13,056	100.0	492.5
1	Malignant neoplasms (C00-C97)	3,649	27.9	137.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,491	19.1	94.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,005	7.7	37.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	957	7.3	36.1
5	Diabetes mellitus (E10-E14)	763	5.8	28.8
6	Cerebrovascular diseases (I60-I69)	575	4.4	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	1.8	9.1
8	Chronic lower respiratory diseases (J40-J47)	222	1.7	8.4
9	Septicemia (A40-A41)	206	1.6	7.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	198	1.5	7.5
11	Viral hepatitis (B15-B19)	182	1.4	6.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	169	1.3	6.4
13	Influenza and pneumonia (J09-J18)	167	1.3	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	127	1.0	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	0.8	3.8
...	All other causes (residual)	2,004	15.3	75.6
Hispanic, both sexes, 60-64 years				
...	All causes	15,477	100.0	754.5
1	Malignant neoplasms (C00-C97)	4,604	29.7	224.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,092	20.0	150.7
3	Diabetes mellitus (E10-E14)	979	6.3	47.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	940	6.1	45.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	779	5.0	38.0
6	Cerebrovascular diseases (I60-I69)	736	4.8	35.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	2.3	17.4
8	Chronic lower respiratory diseases (J40-J47)	324	2.1	15.8
9	Influenza and pneumonia (J09-J18)	268	1.7	13.1
10	Septicemia (A40-A41)	259	1.7	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	1.5	11.0
12	Viral hepatitis (B15-B19)	192	1.2	9.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	0.8	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	93	0.6	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.5	3.8
...	All other causes (residual)	2,420	15.6	118.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	16,280	100.0	1,085.1
1	Malignant neoplasms (C00-C97)	5,050	31.0	336.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,389	20.8	225.9
3	Diabetes mellitus (E10-E14)	1,089	6.7	72.6
4	Cerebrovascular diseases (I60-I69)	794	4.9	52.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	711	4.4	47.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	549	3.4	36.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	2.8	30.1
8	Chronic lower respiratory diseases (J40-J47)	412	2.5	27.5
9	Influenza and pneumonia (J09-J18)	295	1.8	19.7
10	Septicemia (A40-A41)	287	1.8	19.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	248	1.5	16.5
12	Alzheimer disease (G30)	168	1.0	11.2
13	Viral hepatitis (B15-B19)	117	0.7	7.8
14	Parkinson disease (G20-G21)	92	0.6	6.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	0.5	5.9
...	All other causes (residual)	2,539	15.6	169.2
Hispanic, both sexes, 70-74 years				
...	All causes	17,088	100.0	1,628.5
1	Malignant neoplasms (C00-C97)	4,901	28.7	467.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,758	22.0	358.1
3	Diabetes mellitus (E10-E14)	1,174	6.9	111.9
4	Cerebrovascular diseases (I60-I69)	920	5.4	87.7
5	Chronic lower respiratory diseases (J40-J47)	622	3.6	59.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	538	3.1	51.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	2.6	42.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	440	2.6	41.9
9	Influenza and pneumonia (J09-J18)	340	2.0	32.4
10	Alzheimer disease (G30)	312	1.8	29.7
11	Septicemia (A40-A41)	296	1.7	28.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	262	1.5	25.0
13	Parkinson disease (G20-G21)	178	1.0	17.0
14	Pneumonitis due to solids and liquids (J69)	94	0.6	9.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.4	7.0
...	All other causes (residual)	2,732	16.0	260.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	18,699	100.0	2,652.3
1	Malignant neoplasms (C00-C97)	4,795	25.6	680.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,249	22.7	602.7
3	Cerebrovascular diseases (I60-I69)	1,231	6.6	174.6
4	Diabetes mellitus (E10-E14)	1,209	6.5	171.5
5	Chronic lower respiratory diseases (J40-J47)	758	4.1	107.5
6	Alzheimer disease (G30)	731	3.9	103.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	482	2.6	68.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	444	2.4	63.0
9	Influenza and pneumonia (J09-J18)	399	2.1	56.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	355	1.9	50.4
11	Septicemia (A40-A41)	319	1.7	45.2
12	Parkinson disease (G20-G21)	302	1.6	42.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	277	1.5	39.3
14	Pneumonitis due to solids and liquids (J69)	121	0.6	17.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.6	15.3
...	All other causes (residual)	2,919	15.6	414.0
Hispanic, both sexes, 80-84 years				
...	All causes	21,131	100.0	4,462.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,130	24.3	1,083.4
2	Malignant neoplasms (C00-C97)	4,301	20.4	908.3
3	Cerebrovascular diseases (I60-I69)	1,526	7.2	322.3
4	Alzheimer disease (G30)	1,277	6.0	269.7
5	Diabetes mellitus (E10-E14)	1,103	5.2	232.9
6	Chronic lower respiratory diseases (J40-J47)	928	4.4	196.0
7	Influenza and pneumonia (J09-J18)	560	2.7	118.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	526	2.5	111.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	2.2	98.6
10	Parkinson disease (G20-G21)	453	2.1	95.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	384	1.8	81.1
12	Septicemia (A40-A41)	345	1.6	72.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	283	1.3	59.8
14	Pneumonitis due to solids and liquids (J69)	174	0.8	36.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.6	26.2
...	All other causes (residual)	3,550	16.8	749.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	21,683	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,595	25.8	---
2	Malignant neoplasms (C00-C97)	3,465	16.0	---
3	Alzheimer disease (G30)	1,889	8.7	---
4	Cerebrovascular diseases (I60-I69)	1,714	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	984	4.5	---
6	Diabetes mellitus (E10-E14)	852	3.9	---
7	Influenza and pneumonia (J09-J18)	612	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	494	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	478	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	412	1.9	---
11	Parkinson disease (G20-G21)	405	1.9	---
12	Septicemia (A40-A41)	317	1.5	---
13	Pneumonitis due to solids and liquids (J69)	192	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	153	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.6	---
...	All other causes (residual)	3,990	18.4	---
Hispanic, both sexes, 90-94 years				
...	All causes	15,820	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,519	28.6	---
2	Alzheimer disease (G30)	1,864	11.8	---
3	Malignant neoplasms (C00-C97)	1,666	10.5	---
4	Cerebrovascular diseases (I60-I69)	1,285	8.1	---
5	Chronic lower respiratory diseases (J40-J47)	691	4.4	---
6	Diabetes mellitus (E10-E14)	529	3.3	---
7	Influenza and pneumonia (J09-J18)	466	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	349	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	1.9	---
11	Parkinson disease (G20-G21)	242	1.5	---
12	Septicemia (A40-A41)	214	1.4	---
13	Pneumonitis due to solids and liquids (J69)	134	0.8	---
14	Nutritional deficiencies (E40-E64)	80	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.4	---
...	All other causes (residual)	3,070	19.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	6,368	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,884	29.6	---
2	Alzheimer disease (G30)	837	13.1	---
3	Cerebrovascular diseases (I60-I69)	578	9.1	---
4	Malignant neoplasms (C00-C97)	446	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	258	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	2.9	---
7	Influenza and pneumonia (J09-J18)	171	2.7	---
8	Diabetes mellitus (E10-E14)	136	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	1.8	---
11	Septicemia (A40-A41)	75	1.2	---
12	Parkinson disease (G20-G21)	71	1.1	---
13	Pneumonitis due to solids and liquids (J69)	53	0.8	---
14	Nutritional deficiencies (E40-E64)	52	0.8	---
15	Atherosclerosis (I70)	30	0.5	---
...	All other causes (residual)	1,357	21.3	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,365	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	432	31.6	---
2	Alzheimer disease (G30)	188	13.8	---
3	Cerebrovascular diseases (I60-I69)	118	8.6	---
4	Chronic lower respiratory diseases (J40-J47)	58	4.2	---
5	Malignant neoplasms (C00-C97)	57	4.2	---
6	Influenza and pneumonia (J09-J18)	46	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	3.2	---
8	Diabetes mellitus (E10-E14)	20	1.5	---
9	Nutritional deficiencies (E40-E64)	19	1.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.1	---
12	Atherosclerosis (I70)	11	0.8	---
13	Septicemia (A40-A41)	10	0.7	---
14	Pneumonitis due to solids and liquids (J69)	9	0.7	---
15	Parkinson disease (G20-G21)	7	0.5	---
...	All other causes (residual)	313	22.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	108,579	100.0	364.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,009	20.3	73.9
2	Malignant neoplasms (C00-C97)	21,034	19.4	70.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,455	11.5	41.8
4	Cerebrovascular diseases (I60-I69)	5,073	4.7	17.0
5	Diabetes mellitus (E10-E14)	5,054	4.7	17.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,369	4.0	14.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,175	2.9	10.7
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.5	9.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,588	2.4	8.7
10	Alzheimer disease (G30)	2,297	2.1	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,988	1.8	6.7
12	Influenza and pneumonia (J09-J18)	1,911	1.8	6.4
13	Septicemia (A40-A41)	1,473	1.4	4.9
14	Certain conditions originating in the perinatal period (P00-P96)	1,356	1.2	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,354	1.2	4.5
...	All other causes (residual)	19,684	18.1	66.1
Hispanic, male, 1-4 years				
...	All causes	455	100.0	21.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	31.4	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	12.1	2.6
3	Malignant neoplasms (C00-C97)	50	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.6	1.4
5	Influenza and pneumonia (J09-J18)	15	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.1	*
7	Cerebrovascular diseases (I60-I69)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
11	Acute bronchitis and bronchiolitis (J20-J21)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
14	Meningitis (G00,G03)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (residual)	102	22.4	4.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	282	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	28.4	3.0
2	Malignant neoplasms (C00-C97)	42	14.9	1.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	9.6	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	5.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	5.0	*
6	Septicemia (A40-A41)	7	2.5	*
7	Influenza and pneumonia (J09-J18)	6	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.8	*
9	Cerebrovascular diseases (I60-I69)	3	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
11	Diabetes mellitus (E10-E14)	2	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
14	Tuberculosis (A16-A19)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
...	All other causes (residual)	69	24.5	2.6
Hispanic, male, 10-14 years				
...	All causes	395	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	25.6	3.8
2	Malignant neoplasms (C00-C97)	71	18.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	15.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.6	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	5.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.0	*
8	Cerebrovascular diseases (I60-I69)	7	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
9	Influenza and pneumonia (J09-J18)	4	1.0	*
11	Diabetes mellitus (E10-E14)	3	0.8	*
12	Diseases of appendix (K35-K38)	2	0.5	*
13	Meningococcal infection (A39)	1	0.3	*
13	Septicemia (A40-A41)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	72	18.2	2.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,547	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	37.4	23.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	319	20.6	12.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	20.2	12.5
4	Malignant neoplasms (C00-C97)	90	5.8	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.1	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.2	*
7	Legal intervention (Y35,Y89.0)	12	0.8	*
8	Cerebrovascular diseases (I60-I69)	9	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
10	Diabetes mellitus (E10-E14)	7	0.5	*
11	Influenza and pneumonia (J09-J18)	5	0.3	*
12	Septicemia (A40-A41)	4	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
15	Salmonella infections (A01-A02)	1	0.1	*
15	Anemias (D50-D64)	1	0.1	*
15	Parkinson disease (G20-G21)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Diseases of appendix (K35-K38)	1	0.1	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.1	*
...	All other causes (residual)	138	8.9	5.5
Hispanic, male, 20-24 years				
...	All causes	2,840	100.0	113.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,311	46.2	52.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	18.3	20.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	468	16.5	18.7
4	Malignant neoplasms (C00-C97)	142	5.0	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	2.4	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.7	0.8
7	Cerebrovascular diseases (I60-I69)	17	0.6	*
8	Legal intervention (Y35,Y89.0)	13	0.5	*
9	Diabetes mellitus (E10-E14)	12	0.4	*
10	Septicemia (A40-A41)	11	0.4	*
10	Influenza and pneumonia (J09-J18)	11	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
14	Anemias (D50-D64)	5	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
...	All other causes (residual)	220	7.7	8.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	3,181	100.0	126.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,450	45.6	57.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	443	13.9	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	440	13.8	17.5
4	Malignant neoplasms (C00-C97)	159	5.0	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	4.1	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	37	1.2	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.9	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.9	1.2
9	Cerebrovascular diseases (I60-I69)	24	0.8	1.0
9	Legal intervention (Y35,Y89.0)	24	0.8	1.0
11	Diabetes mellitus (E10-E14)	20	0.6	0.8
12	Septicemia (A40-A41)	19	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	19	0.6	*
14	Influenza and pneumonia (J09-J18)	16	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (residual)	330	10.4	13.1
Hispanic, male, 30-34 years				
...	All causes	3,128	100.0	131.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,277	40.8	53.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	356	11.4	15.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	316	10.1	13.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	253	8.1	10.6
5	Malignant neoplasms (C00-C97)	192	6.1	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	84	2.7	3.5
7	Cerebrovascular diseases (I60-I69)	51	1.6	2.1
8	Diabetes mellitus (E10-E14)	49	1.6	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	40	1.3	1.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	0.8
11	Influenza and pneumonia (J09-J18)	19	0.6	*
12	Septicemia (A40-A41)	16	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.5	*
12	Legal intervention (Y35,Y89.0)	16	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
...	All other causes (residual)	410	13.1	17.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	3,516	100.0	153.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,276	36.3	55.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	9.4	14.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	9.0	13.8
4	Malignant neoplasms (C00-C97)	293	8.3	12.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	7.9	12.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	192	5.5	8.4
7	Cerebrovascular diseases (I60-I69)	95	2.7	4.2
8	Diabetes mellitus (E10-E14)	64	1.8	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.7	2.7
10	Septicemia (A40-A41)	32	0.9	1.4
11	Influenza and pneumonia (J09-J18)	27	0.8	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	0.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.5	*
13	Legal intervention (Y35,Y89.0)	19	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	17	0.5	*
...	All other causes (residual)	478	13.6	20.9
Hispanic, male, 40-44 years				
...	All causes	3,842	100.0	186.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,059	27.6	51.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	525	13.7	25.5
3	Malignant neoplasms (C00-C97)	455	11.8	22.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	273	7.1	13.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	6.6	12.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	186	4.8	9.0
7	Cerebrovascular diseases (I60-I69)	146	3.8	7.1
8	Diabetes mellitus (E10-E14)	123	3.2	6.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	56	1.5	2.7
10	Septicemia (A40-A41)	43	1.1	2.1
11	Influenza and pneumonia (J09-J18)	39	1.0	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.9	1.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	21	0.5	1.0
15	Viral hepatitis (B15-B19)	20	0.5	1.0
...	All other causes (residual)	586	15.3	28.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	5,072	100.0	271.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,080	21.3	57.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	825	16.3	44.2
3	Malignant neoplasms (C00-C97)	751	14.8	40.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	473	9.3	25.3
5	Diabetes mellitus (E10-E14)	252	5.0	13.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	4.5	12.2
7	Cerebrovascular diseases (I60-I69)	185	3.6	9.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	2.8	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.5	4.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	76	1.5	4.1
11	Septicemia (A40-A41)	59	1.2	3.2
11	Influenza and pneumonia (J09-J18)	59	1.2	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	0.9	2.4
14	Viral hepatitis (B15-B19)	44	0.9	2.4
15	Chronic lower respiratory diseases (J40-J47)	20	0.4	1.1
...	All other causes (residual)	754	14.9	40.4
Hispanic, male, 50-54 years				
...	All causes	7,073	100.0	439.8
1	Malignant neoplasms (C00-C97)	1,414	20.0	87.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,357	19.2	84.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,029	14.5	64.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	650	9.2	40.4
5	Diabetes mellitus (E10-E14)	354	5.0	22.0
6	Cerebrovascular diseases (I60-I69)	270	3.8	16.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	229	3.2	14.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	108	1.5	6.7
9	Septicemia (A40-A41)	107	1.5	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	1.4	6.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	1.4	6.3
12	Viral hepatitis (B15-B19)	94	1.3	5.8
13	Influenza and pneumonia (J09-J18)	84	1.2	5.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.1	4.7
15	Chronic lower respiratory diseases (J40-J47)	53	0.7	3.3
...	All other causes (residual)	1,044	14.8	64.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	8,392	100.0	643.2
1	Malignant neoplasms (C00-C97)	1,948	23.2	149.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,811	21.6	138.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	785	9.4	60.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	726	8.7	55.6
5	Diabetes mellitus (E10-E14)	470	5.6	36.0
6	Cerebrovascular diseases (I60-I69)	368	4.4	28.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	1.9	12.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.5	10.0
9	Chronic lower respiratory diseases (J40-J47)	125	1.5	9.6
10	Septicemia (A40-A41)	121	1.4	9.3
11	Viral hepatitis (B15-B19)	120	1.4	9.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	113	1.3	8.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	102	1.2	7.8
14	Influenza and pneumonia (J09-J18)	98	1.2	7.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	0.9	6.0
...	All other causes (residual)	1,237	14.7	94.8
Hispanic, male, 60-64 years				
...	All causes	9,582	100.0	979.9
1	Malignant neoplasms (C00-C97)	2,488	26.0	254.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,180	22.8	222.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	695	7.3	71.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	581	6.1	59.4
5	Diabetes mellitus (E10-E14)	568	5.9	58.1
6	Cerebrovascular diseases (I60-I69)	463	4.8	47.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.2	21.3
8	Chronic lower respiratory diseases (J40-J47)	158	1.6	16.2
9	Influenza and pneumonia (J09-J18)	153	1.6	15.6
10	Septicemia (A40-A41)	151	1.6	15.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.4	14.0
12	Viral hepatitis (B15-B19)	133	1.4	13.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	1.1	11.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	73	0.8	7.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.6	5.6
...	All other causes (residual)	1,429	14.9	146.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	9,533	100.0	1,372.9
1	Malignant neoplasms (C00-C97)	2,737	28.7	394.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,238	23.5	322.3
3	Diabetes mellitus (E10-E14)	625	6.6	90.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	492	5.2	70.9
5	Cerebrovascular diseases (I60-I69)	434	4.6	62.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	376	3.9	54.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	245	2.6	35.3
8	Chronic lower respiratory diseases (J40-J47)	240	2.5	34.6
9	Influenza and pneumonia (J09-J18)	168	1.8	24.2
10	Septicemia (A40-A41)	143	1.5	20.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.4	19.3
12	Viral hepatitis (B15-B19)	83	0.9	12.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	0.7	10.2
14	Alzheimer disease (G30)	66	0.7	9.5
15	Parkinson disease (G20-G21)	59	0.6	8.5
...	All other causes (residual)	1,422	14.9	204.8
Hispanic, male, 70-74 years				
...	All causes	9,499	100.0	2,037.5
1	Malignant neoplasms (C00-C97)	2,634	27.7	565.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,277	24.0	488.4
3	Diabetes mellitus (E10-E14)	676	7.1	145.0
4	Cerebrovascular diseases (I60-I69)	496	5.2	106.4
5	Chronic lower respiratory diseases (J40-J47)	320	3.4	68.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	302	3.2	64.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	3.1	62.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	2.4	48.7
9	Influenza and pneumonia (J09-J18)	189	2.0	40.5
10	Septicemia (A40-A41)	159	1.7	34.1
11	Alzheimer disease (G30)	135	1.4	29.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.4	27.9
13	Parkinson disease (G20-G21)	120	1.3	25.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	0.7	13.3
15	Pneumonitis due to solids and liquids (J69)	56	0.6	12.0
...	All other causes (residual)	1,426	15.0	305.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	9,834	100.0	3,288.8
1	Malignant neoplasms (C00-C97)	2,601	26.4	869.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,361	24.0	789.6
3	Cerebrovascular diseases (I60-I69)	623	6.3	208.4
4	Diabetes mellitus (E10-E14)	610	6.2	204.0
5	Chronic lower respiratory diseases (J40-J47)	411	4.2	137.5
6	Alzheimer disease (G30)	285	2.9	95.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	2.7	89.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	2.3	76.9
9	Influenza and pneumonia (J09-J18)	223	2.3	74.6
10	Parkinson disease (G20-G21)	192	2.0	64.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	183	1.9	61.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	1.5	48.2
13	Septicemia (A40-A41)	141	1.4	47.2
14	Pneumonitis due to solids and liquids (J69)	68	0.7	22.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.6	20.1
...	All other causes (residual)	1,435	14.6	479.9
Hispanic, male, 80-84 years				
...	All causes	10,177	100.0	5,338.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,631	25.9	1,380.2
2	Malignant neoplasms (C00-C97)	2,256	22.2	1,183.5
3	Cerebrovascular diseases (I60-I69)	639	6.3	335.2
4	Diabetes mellitus (E10-E14)	568	5.6	298.0
5	Chronic lower respiratory diseases (J40-J47)	469	4.6	246.0
6	Alzheimer disease (G30)	420	4.1	220.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	2.6	136.4
8	Parkinson disease (G20-G21)	254	2.5	133.2
9	Influenza and pneumonia (J09-J18)	251	2.5	131.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	2.5	131.1
11	Septicemia (A40-A41)	161	1.6	84.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.6	84.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	137	1.3	71.9
14	Pneumonitis due to solids and liquids (J69)	105	1.0	55.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.5	28.9
...	All other causes (residual)	1,560	15.3	818.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	9,402	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,472	26.3	---
2	Malignant neoplasms (C00-C97)	1,717	18.3	---
3	Cerebrovascular diseases (I60-I69)	652	6.9	---
4	Alzheimer disease (G30)	611	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	467	5.0	---
6	Diabetes mellitus (E10-E14)	375	4.0	---
7	Influenza and pneumonia (J09-J18)	263	2.8	---
8	Parkinson disease (G20-G21)	236	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	2.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	1.9	---
12	Septicemia (A40-A41)	148	1.6	---
13	Pneumonitis due to solids and liquids (J69)	93	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	75	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.7	---
...	All other causes (residual)	1,601	17.0	---
Hispanic, male, 90-94 years				
...	All causes	5,910	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,717	29.1	---
2	Malignant neoplasms (C00-C97)	781	13.2	---
3	Alzheimer disease (G30)	521	8.8	---
4	Cerebrovascular diseases (I60-I69)	413	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	300	5.1	---
6	Diabetes mellitus (E10-E14)	223	3.8	---
7	Influenza and pneumonia (J09-J18)	194	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	2.4	---
9	Parkinson disease (G20-G21)	129	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	2.1	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	119	2.0	---
12	Septicemia (A40-A41)	97	1.6	---
13	Pneumonitis due to solids and liquids (J69)	61	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	---
...	All other causes (residual)	1,029	17.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,955	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	618	31.6	---
2	Malignant neoplasms (C00-C97)	182	9.3	---
3	Alzheimer disease (G30)	178	9.1	---
4	Cerebrovascular diseases (I60-I69)	145	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	76	3.9	---
6	Influenza and pneumonia (J09-J18)	56	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	2.4	---
9	Diabetes mellitus (E10-E14)	46	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.3	---
11	Septicemia (A40-A41)	28	1.4	---
11	Parkinson disease (G20-G21)	28	1.4	---
13	Pneumonitis due to solids and liquids (J69)	24	1.2	---
14	Nutritional deficiencies (E40-E64)	20	1.0	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.5	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	10	0.5	---
...	All other causes (residual)	387	19.8	---
Hispanic, male, 100 years and over				
...	All causes	321	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	34.9	---
2	Alzheimer disease (G30)	34	10.6	---
3	Malignant neoplasms (C00-C97)	25	7.8	---
4	Cerebrovascular diseases (I60-I69)	19	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	17	5.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	2.8	---
7	Influenza and pneumonia (J09-J18)	8	2.5	---
8	Nutritional deficiencies (E40-E64)	7	2.2	---
9	Diabetes mellitus (E10-E14)	6	1.9	---
9	Pneumonitis due to solids and liquids (J69)	6	1.9	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.6	---
13	Parkinson disease (G20-G21)	4	1.2	---
13	Atherosclerosis (I70)	4	1.2	---
15	Septicemia (A40-A41)	2	0.6	---
15	Anemias (D50-D64)	2	0.6	---
...	All other causes (residual)	56	17.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	88,670	100.0	304.0
1	Malignant neoplasms (C00-C97)	19,516	22.0	66.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,393	19.6	59.6
3	Cerebrovascular diseases (I60-I69)	5,702	6.4	19.5
4	Alzheimer disease (G30)	5,086	5.7	17.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,302	4.9	14.7
6	Diabetes mellitus (E10-E14)	4,162	4.7	14.3
7	Chronic lower respiratory diseases (J40-J47)	2,840	3.2	9.7
8	Influenza and pneumonia (J09-J18)	1,940	2.2	6.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,940	2.2	6.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,917	2.2	6.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,445	1.6	5.0
12	Septicemia (A40-A41)	1,396	1.6	4.8
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.2	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.0	3.0
15	Parkinson disease (G20-G21)	758	0.9	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	758	0.9	2.6
...	All other causes (residual)	17,589	19.8	60.3
Hispanic, female, 1-4 years				
...	All causes	350	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	26.0	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.3	2.8
3	Malignant neoplasms (C00-C97)	42	12.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	5.7	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.4	*
6	Influenza and pneumonia (J09-J18)	9	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.3	*
8	Septicemia (A40-A41)	4	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.1	*
10	Cerebrovascular diseases (I60-I69)	3	0.9	*
11	Anemias (D50-D64)	2	0.6	*
11	Meningitis (G00,G03)	2	0.6	*
13	Meningococcal infection (A39)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	91	26.0	4.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	239	100.0	9.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	28.5	2.6
2	Malignant neoplasms (C00-C97)	58	24.3	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	7.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	4.2	*
5	Septicemia (A40-A41)	8	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.9	*
6	Influenza and pneumonia (J09-J18)	7	2.9	*
8	Cerebrovascular diseases (I60-I69)	5	2.1	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
11	Diseases of appendix (K35-K38)	2	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	46	19.2	1.8
Hispanic, female, 10-14 years				
...	All causes	249	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	21.3	2.1
2	Malignant neoplasms (C00-C97)	51	20.5	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	12.0	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	4.8	*
6	Cerebrovascular diseases (I60-I69)	9	3.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	5	2.0	*
9	Influenza and pneumonia (J09-J18)	4	1.6	*
10	Septicemia (A40-A41)	2	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Anemias (D50-D64)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	55	22.1	2.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	628	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	38.1	10.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	17.5	4.6
3	Malignant neoplasms (C00-C97)	62	9.9	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	9.9	2.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	1.9	*
7	Influenza and pneumonia (J09-J18)	11	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	*
8	Cerebrovascular diseases (I60-I69)	8	1.3	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
12	Diabetes mellitus (E10-E14)	3	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Tuberculosis (A16-A19)	2	0.3	*
14	Septicemia (A40-A41)	2	0.3	*
...	All other causes (residual)	78	12.4	3.3
Hispanic, female, 20-24 years				
...	All causes	839	100.0	35.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	42.6	15.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	10.7	3.8
3	Malignant neoplasms (C00-C97)	77	9.2	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	8.3	3.0
5	Pregnancy, childbirth and the puerperium (O00-O99)	26	3.1	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	2.4	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.2	*
9	Diabetes mellitus (E10-E14)	7	0.8	*
9	Cerebrovascular diseases (I60-I69)	7	0.8	*
11	Septicemia (A40-A41)	6	0.7	*
12	Influenza and pneumonia (J09-J18)	5	0.6	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
14	Anemias (D50-D64)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (residual)	135	16.1	5.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	985	100.0	43.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	35.8	15.5
2	Malignant neoplasms (C00-C97)	113	11.5	5.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	8.6	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	7.8	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.2	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	38	3.9	1.7
7	Cerebrovascular diseases (I60-I69)	25	2.5	1.1
8	Diabetes mellitus (E10-E14)	20	2.0	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	20	2.0	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.3	*
11	Septicemia (A40-A41)	10	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
13	Influenza and pneumonia (J09-J18)	8	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (residual)	157	15.9	6.9
Hispanic, female, 30-34 years				
...	All causes	1,198	100.0	55.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	25.0	13.9
2	Malignant neoplasms (C00-C97)	264	22.0	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	6.9	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	6.1	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	6.0	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	42	3.5	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.7	1.5
8	Cerebrovascular diseases (I60-I69)	25	2.1	1.2
9	Diabetes mellitus (E10-E14)	24	2.0	1.1
10	Septicemia (A40-A41)	14	1.2	*
11	Influenza and pneumonia (J09-J18)	12	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (residual)	219	18.3	10.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,498	100.0	70.3
1	Malignant neoplasms (C00-C97)	423	28.2	19.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	19.5	13.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	8.0	5.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	71	4.7	3.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	4.7	3.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	4.5	3.1
7	Cerebrovascular diseases (I60-I69)	36	2.4	1.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	35	2.3	1.6
9	Diabetes mellitus (E10-E14)	30	2.0	1.4
10	Septicemia (A40-A41)	23	1.5	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.1	*
11	Influenza and pneumonia (J09-J18)	16	1.1	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
...	All other causes (residual)	259	17.3	12.2
Hispanic, female, 40-44 years				
...	All causes	1,882	100.0	94.4
1	Malignant neoplasms (C00-C97)	606	32.2	30.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	14.7	13.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	10.0	9.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	91	4.8	4.6
5	Cerebrovascular diseases (I60-I69)	74	3.9	3.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	3.4	3.2
7	Diabetes mellitus (E10-E14)	63	3.3	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	2.7	2.5
9	Septicemia (A40-A41)	26	1.4	1.3
10	Influenza and pneumonia (J09-J18)	25	1.3	1.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.2	1.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	21	1.1	1.1
13	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.5	*
...	All other causes (residual)	327	17.4	16.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,666	100.0	146.0
1	Malignant neoplasms (C00-C97)	919	34.5	50.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	281	10.5	15.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	10.5	15.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	150	5.6	8.2
5	Diabetes mellitus (E10-E14)	131	4.9	7.2
6	Cerebrovascular diseases (I60-I69)	122	4.6	6.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	2.7	3.9
8	Influenza and pneumonia (J09-J18)	46	1.7	2.5
9	Septicemia (A40-A41)	45	1.7	2.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	1.6	2.3
11	Chronic lower respiratory diseases (J40-J47)	38	1.4	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.3	1.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.8	1.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.6	*
...	All other causes (residual)	449	16.8	24.6
Hispanic, female, 50-54 years				
...	All causes	3,598	100.0	225.9
1	Malignant neoplasms (C00-C97)	1,325	36.8	83.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	488	13.6	30.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	7.0	15.9
4	Diabetes mellitus (E10-E14)	197	5.5	12.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	192	5.3	12.1
6	Cerebrovascular diseases (I60-I69)	156	4.3	9.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	2.2	4.9
8	Septicemia (A40-A41)	64	1.8	4.0
8	Influenza and pneumonia (J09-J18)	64	1.8	4.0
10	Chronic lower respiratory diseases (J40-J47)	57	1.6	3.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	1.5	3.4
12	Viral hepatitis (B15-B19)	31	0.9	1.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	0.8	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	0.7	1.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	25	0.7	1.6
...	All other causes (residual)	559	15.5	35.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	4,664	100.0	346.5
1	Malignant neoplasms (C00-C97)	1,701	36.5	126.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	680	14.6	50.5
3	Diabetes mellitus (E10-E14)	293	6.3	21.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	231	5.0	17.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	4.7	16.3
6	Cerebrovascular diseases (I60-I69)	207	4.4	15.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.4	8.2
8	Chronic lower respiratory diseases (J40-J47)	97	2.1	7.2
9	Septicemia (A40-A41)	85	1.8	6.3
10	Influenza and pneumonia (J09-J18)	69	1.5	5.1
11	Viral hepatitis (B15-B19)	62	1.3	4.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.2	4.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.8	2.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	1.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	0.5	1.6
...	All other causes (residual)	767	16.4	57.0
Hispanic, female, 60-64 years				
...	All causes	5,895	100.0	549.2
1	Malignant neoplasms (C00-C97)	2,116	35.9	197.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	912	15.5	85.0
3	Diabetes mellitus (E10-E14)	411	7.0	38.3
4	Cerebrovascular diseases (I60-I69)	273	4.6	25.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	245	4.2	22.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	3.4	18.4
7	Chronic lower respiratory diseases (J40-J47)	166	2.8	15.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	2.5	13.9
9	Influenza and pneumonia (J09-J18)	115	2.0	10.7
10	Septicemia (A40-A41)	108	1.8	10.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.5	8.3
12	Viral hepatitis (B15-B19)	59	1.0	5.5
13	Alzheimer disease (G30)	45	0.8	4.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	2.5
15	Pneumonitis due to solids and liquids (J69)	23	0.4	2.1
...	All other causes (residual)	959	16.3	89.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	6,747	100.0	837.1
1	Malignant neoplasms (C00-C97)	2,313	34.3	287.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,151	17.1	142.8
3	Diabetes mellitus (E10-E14)	464	6.9	57.6
4	Cerebrovascular diseases (I60-I69)	360	5.3	44.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	219	3.2	27.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	3.1	25.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	2.6	21.5
8	Chronic lower respiratory diseases (J40-J47)	172	2.5	21.3
9	Septicemia (A40-A41)	144	2.1	17.9
10	Influenza and pneumonia (J09-J18)	127	1.9	15.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.7	14.1
12	Alzheimer disease (G30)	102	1.5	12.7
13	Viral hepatitis (B15-B19)	34	0.5	4.2
14	Parkinson disease (G20-G21)	33	0.5	4.1
15	Pneumonitis due to solids and liquids (J69)	30	0.4	3.7
...	All other causes (residual)	1,105	16.4	137.1
Hispanic, female, 70-74 years				
...	All causes	7,589	100.0	1,301.5
1	Malignant neoplasms (C00-C97)	2,267	29.9	388.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,481	19.5	254.0
3	Diabetes mellitus (E10-E14)	498	6.6	85.4
4	Cerebrovascular diseases (I60-I69)	424	5.6	72.7
5	Chronic lower respiratory diseases (J40-J47)	302	4.0	51.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	236	3.1	40.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	221	2.9	37.9
8	Alzheimer disease (G30)	177	2.3	30.4
9	Influenza and pneumonia (J09-J18)	151	2.0	25.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	2.0	25.7
11	Septicemia (A40-A41)	137	1.8	23.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	1.7	22.6
13	Parkinson disease (G20-G21)	58	0.8	9.9
14	Pneumonitis due to solids and liquids (J69)	38	0.5	6.5
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	29	0.4	5.0
...	All other causes (residual)	1,288	17.0	220.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	8,865	100.0	2,183.5
1	Malignant neoplasms (C00-C97)	2,194	24.7	540.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,888	21.3	465.0
3	Cerebrovascular diseases (I60-I69)	608	6.9	149.8
4	Diabetes mellitus (E10-E14)	599	6.8	147.5
5	Alzheimer disease (G30)	446	5.0	109.9
6	Chronic lower respiratory diseases (J40-J47)	347	3.9	85.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	252	2.8	62.1
8	Septicemia (A40-A41)	178	2.0	43.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	2.0	43.6
10	Influenza and pneumonia (J09-J18)	176	2.0	43.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	172	1.9	42.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.5	32.8
13	Parkinson disease (G20-G21)	110	1.2	27.1
14	Pneumonitis due to solids and liquids (J69)	53	0.6	13.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.5	11.8
...	All other causes (residual)	1,484	16.7	365.5
Hispanic, female, 80-84 years				
...	All causes	10,954	100.0	3,872.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,499	22.8	883.5
2	Malignant neoplasms (C00-C97)	2,045	18.7	723.0
3	Cerebrovascular diseases (I60-I69)	887	8.1	313.6
4	Alzheimer disease (G30)	857	7.8	303.0
5	Diabetes mellitus (E10-E14)	535	4.9	189.1
6	Chronic lower respiratory diseases (J40-J47)	459	4.2	162.3
7	Influenza and pneumonia (J09-J18)	309	2.8	109.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	2.5	97.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	2.0	78.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	1.9	73.2
11	Parkinson disease (G20-G21)	199	1.8	70.4
12	Septicemia (A40-A41)	184	1.7	65.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.3	51.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	24.4
14	Pneumonitis due to solids and liquids (J69)	69	0.6	24.4
...	All other causes (residual)	1,990	18.2	703.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	12,281	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,123	25.4	---
2	Malignant neoplasms (C00-C97)	1,748	14.2	---
3	Alzheimer disease (G30)	1,278	10.4	---
4	Cerebrovascular diseases (I60-I69)	1,062	8.6	---
5	Chronic lower respiratory diseases (J40-J47)	517	4.2	---
6	Diabetes mellitus (E10-E14)	477	3.9	---
7	Influenza and pneumonia (J09-J18)	349	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	1.9	---
11	Septicemia (A40-A41)	169	1.4	---
11	Parkinson disease (G20-G21)	169	1.4	---
13	Pneumonitis due to solids and liquids (J69)	99	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	78	0.6	---
15	Nutritional deficiencies (E40-E64)	73	0.6	---
...	All other causes (residual)	2,379	19.4	---
Hispanic, female, 90-94 years				
...	All causes	9,910	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,802	28.3	---
2	Alzheimer disease (G30)	1,343	13.6	---
3	Malignant neoplasms (C00-C97)	885	8.9	---
4	Cerebrovascular diseases (I60-I69)	872	8.8	---
5	Chronic lower respiratory diseases (J40-J47)	391	3.9	---
6	Diabetes mellitus (E10-E14)	306	3.1	---
7	Influenza and pneumonia (J09-J18)	272	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.8	---
11	Septicemia (A40-A41)	117	1.2	---
12	Parkinson disease (G20-G21)	113	1.1	---
13	Pneumonitis due to solids and liquids (J69)	73	0.7	---
14	Nutritional deficiencies (E40-E64)	62	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	---
...	All other causes (residual)	2,022	20.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	4,413	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,266	28.7	---
2	Alzheimer disease (G30)	659	14.9	---
3	Cerebrovascular diseases (I60-I69)	433	9.8	---
4	Malignant neoplasms (C00-C97)	264	6.0	---
5	Chronic lower respiratory diseases (J40-J47)	182	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	3.0	---
7	Influenza and pneumonia (J09-J18)	115	2.6	---
8	Diabetes mellitus (E10-E14)	90	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.6	---
11	Septicemia (A40-A41)	47	1.1	---
12	Parkinson disease (G20-G21)	43	1.0	---
13	Nutritional deficiencies (E40-E64)	32	0.7	---
14	Pneumonitis due to solids and liquids (J69)	29	0.7	---
15	Atherosclerosis (I70)	21	0.5	---
...	All other causes (residual)	959	21.7	---
Hispanic, female, 100 years and over				
...	All causes	1,044	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	30.7	---
2	Alzheimer disease (G30)	154	14.8	---
3	Cerebrovascular diseases (I60-I69)	99	9.5	---
4	Chronic lower respiratory diseases (J40-J47)	41	3.9	---
5	Influenza and pneumonia (J09-J18)	38	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	3.3	---
7	Malignant neoplasms (C00-C97)	32	3.1	---
8	Diabetes mellitus (E10-E14)	14	1.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.3	---
10	Nutritional deficiencies (E40-E64)	12	1.1	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	---
12	Septicemia (A40-A41)	8	0.8	---
13	Atherosclerosis (I70)	7	0.7	---
14	Anemias (D50-D64)	4	0.4	---
15	Parkinson disease (G20-G21)	3	0.3	---
15	Pneumonitis due to solids and liquids (J69)	3	0.3	---
...	All other causes (residual)	251	24.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,179,857	100.0	1,083.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	508,485	23.3	252.7
2	Malignant neoplasms (C00-C97)	465,679	21.4	231.4
3	Chronic lower respiratory diseases (J40-J47)	139,833	6.4	69.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,029	5.8	63.1
5	Cerebrovascular diseases (I60-I69)	110,038	5.0	54.7
6	Alzheimer disease (G30)	101,876	4.7	50.6
7	Diabetes mellitus (E10-E14)	55,116	2.5	27.4
8	Influenza and pneumonia (J09-J18)	43,397	2.0	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,106	1.7	18.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,191	1.6	17.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,223	1.4	15.0
12	Septicemia (A40-A41)	30,198	1.4	15.0
13	Parkinson disease (G20-G21)	27,582	1.3	13.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,465	1.1	12.2
15	Pneumonitis due to solids and liquids (J69)	16,449	0.8	8.2
...	All other causes (residual)	426,190	19.6	211.8
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,858	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	36.4	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	9.8	2.2
3	Malignant neoplasms (C00-C97)	164	8.8	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.1	0.7
6	Influenza and pneumonia (J09-J18)	44	2.4	0.5
7	Cerebrovascular diseases (I60-I69)	29	1.6	0.3
8	Septicemia (A40-A41)	24	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.2	0.3
10	Anemias (D50-D64)	12	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
15	Meningitis (G00,G03)	7	0.4	*
...	All other causes (residual)	490	26.4	5.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,115	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	31.2	3.3
2	Malignant neoplasms (C00-C97)	227	20.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	8.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.4	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.0	0.3
5	Influenza and pneumonia (J09-J18)	33	3.0	0.3
7	Cerebrovascular diseases (I60-I69)	22	2.0	0.2
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.3	*
10	Septicemia (A40-A41)	8	0.7	*
11	Pneumonitis due to solids and liquids (J69)	4	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
13	Meningitis (G00,G03)	3	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
...	All other causes (residual)	242	21.7	2.3
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,667	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	471	28.3	4.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	327	19.6	2.9
3	Malignant neoplasms (C00-C97)	206	12.4	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	6.2	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	3.5	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	3.1	0.5
7	Influenza and pneumonia (J09-J18)	32	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.8	*
11	Septicemia (A40-A41)	11	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.7	*
13	Diabetes mellitus (E10-E14)	8	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Anemias (D50-D64)	4	0.2	*
15	Meningitis (G00,G03)	4	0.2	*
...	All other causes (residual)	318	19.1	2.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,592	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,364	42.3	20.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,636	29.3	14.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	281	5.0	2.4
4	Malignant neoplasms (C00-C97)	276	4.9	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.6	0.8
7	Diabetes mellitus (E10-E14)	36	0.6	0.3
8	Influenza and pneumonia (J09-J18)	34	0.6	0.3
9	Cerebrovascular diseases (I60-I69)	29	0.5	0.3
10	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
11	Septicemia (A40-A41)	18	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.2	*
15	Aortic aneurysm and dissection (I71)	10	0.2	*
...	All other causes (residual)	606	10.8	5.2
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,615	100.0	95.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,191	53.3	50.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,438	21.0	20.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	482	4.1	3.9
4	Malignant neoplasms (C00-C97)	410	3.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	291	2.5	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.8	0.8
7	Diabetes mellitus (E10-E14)	93	0.8	0.8
8	Influenza and pneumonia (J09-J18)	66	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.5	0.5
10	Septicemia (A40-A41)	47	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	45	0.4	0.4
12	Cerebrovascular diseases (I60-I69)	37	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.3	0.3
14	Legal intervention (Y35,Y89.0)	19	0.2	*
15	Pneumonitis due to solids and liquids (J69)	17	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.1	*
...	All other causes (residual)	1,271	10.9	10.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	16,736	100.0	127.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,810	52.6	67.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,828	16.9	21.6
3	Malignant neoplasms (C00-C97)	695	4.2	5.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	659	3.9	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	565	3.4	4.3
6	Diabetes mellitus (E10-E14)	164	1.0	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	156	0.9	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	0.8	1.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	118	0.7	0.9
10	Cerebrovascular diseases (I60-I69)	103	0.6	0.8
11	Influenza and pneumonia (J09-J18)	85	0.5	0.6
12	Septicemia (A40-A41)	83	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	74	0.4	0.6
14	Legal intervention (Y35,Y89.0)	46	0.3	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.2	0.3
...	All other causes (residual)	2,188	13.1	16.7
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	20,028	100.0	159.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,433	47.1	75.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,858	14.3	22.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,303	6.5	10.4
4	Malignant neoplasms (C00-C97)	1,297	6.5	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	628	3.1	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	426	2.1	3.4
7	Diabetes mellitus (E10-E14)	229	1.1	1.8
8	Cerebrovascular diseases (I60-I69)	177	0.9	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	149	0.7	1.2
10	Influenza and pneumonia (J09-J18)	139	0.7	1.1
11	Septicemia (A40-A41)	130	0.6	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	126	0.6	1.0
13	Chronic lower respiratory diseases (J40-J47)	72	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.4
...	All other causes (residual)	2,955	14.8	23.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	22,872	100.0	187.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,783	38.4	72.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,843	12.4	23.3
3	Malignant neoplasms (C00-C97)	2,378	10.4	19.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,173	9.5	17.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	749	3.3	6.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	558	2.4	4.6
7	Diabetes mellitus (E10-E14)	416	1.8	3.4
8	Cerebrovascular diseases (I60-I69)	322	1.4	2.6
9	Septicemia (A40-A41)	204	0.9	1.7
10	Influenza and pneumonia (J09-J18)	198	0.9	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	0.7	1.3
12	Chronic lower respiratory diseases (J40-J47)	142	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	0.5	1.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	108	0.5	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	106	0.5	0.9
...	All other causes (residual)	3,614	15.8	29.6
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,524	100.0	233.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,346	27.7	64.8
2	Malignant neoplasms (C00-C97)	4,095	15.4	36.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,742	14.1	33.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,921	11.0	25.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,139	4.3	10.0
6	Diabetes mellitus (E10-E14)	638	2.4	5.6
7	Cerebrovascular diseases (I60-I69)	510	1.9	4.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	1.6	3.8
9	Chronic lower respiratory diseases (J40-J47)	317	1.2	2.8
10	Septicemia (A40-A41)	272	1.0	2.4
11	Influenza and pneumonia (J09-J18)	261	1.0	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	184	0.7	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	0.5	1.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	0.5	1.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	104	0.4	0.9
...	All other causes (residual)	4,291	16.2	37.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	42,453	100.0	326.7
1	Malignant neoplasms (C00-C97)	8,523	20.1	65.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,213	19.3	63.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,196	17.0	55.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,550	8.4	27.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,122	5.0	16.3
6	Diabetes mellitus (E10-E14)	1,306	3.1	10.0
7	Cerebrovascular diseases (I60-I69)	979	2.3	7.5
8	Chronic lower respiratory diseases (J40-J47)	747	1.8	5.7
9	Septicemia (A40-A41)	506	1.2	3.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	484	1.1	3.7
11	Influenza and pneumonia (J09-J18)	452	1.1	3.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	362	0.9	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	271	0.6	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	228	0.5	1.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	213	0.5	1.6
...	All other causes (residual)	7,301	17.2	56.2
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	70,312	100.0	499.8
1	Malignant neoplasms (C00-C97)	17,878	25.4	127.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,744	19.5	97.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,143	13.0	65.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,711	5.3	26.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,591	5.1	25.5
6	Chronic lower respiratory diseases (J40-J47)	2,385	3.4	17.0
7	Diabetes mellitus (E10-E14)	2,318	3.3	16.5
8	Cerebrovascular diseases (I60-I69)	1,734	2.5	12.3
9	Septicemia (A40-A41)	971	1.4	6.9
10	Influenza and pneumonia (J09-J18)	806	1.1	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	686	1.0	4.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	576	0.8	4.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	510	0.7	3.6
14	Viral hepatitis (B15-B19)	436	0.6	3.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	345	0.5	2.5
...	All other causes (residual)	11,478	16.3	81.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	114,084	100.0	741.6
1	Malignant neoplasms (C00-C97)	34,379	30.1	223.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,392	20.5	152.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,596	8.4	62.4
4	Chronic lower respiratory diseases (J40-J47)	5,790	5.1	37.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,948	4.3	32.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,943	3.5	25.6
7	Diabetes mellitus (E10-E14)	3,860	3.4	25.1
8	Cerebrovascular diseases (I60-I69)	3,026	2.7	19.7
9	Septicemia (A40-A41)	1,591	1.4	10.3
10	Influenza and pneumonia (J09-J18)	1,432	1.3	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,219	1.1	7.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	960	0.8	6.2
13	Viral hepatitis (B15-B19)	794	0.7	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	541	0.5	3.5
15	Pneumonitis due to solids and liquids (J69)	479	0.4	3.1
...	All other causes (residual)	18,134	15.9	117.9
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	150,915	100.0	1,040.5
1	Malignant neoplasms (C00-C97)	49,705	32.9	342.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,123	21.3	221.5
3	Chronic lower respiratory diseases (J40-J47)	9,837	6.5	67.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,461	4.9	51.4
5	Diabetes mellitus (E10-E14)	5,352	3.5	36.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,035	3.3	34.7
7	Cerebrovascular diseases (I60-I69)	4,565	3.0	31.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,219	2.1	22.2
9	Septicemia (A40-A41)	2,288	1.5	15.8
10	Influenza and pneumonia (J09-J18)	2,067	1.4	14.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,820	1.2	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,319	0.9	9.1
13	Viral hepatitis (B15-B19)	847	0.6	5.8
14	Alzheimer disease (G30)	700	0.5	4.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	692	0.5	4.8
...	All other causes (residual)	23,885	15.8	164.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	184,996	100.0	1,463.6
1	Malignant neoplasms (C00-C97)	62,242	33.6	492.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,696	21.5	314.1
3	Chronic lower respiratory diseases (J40-J47)	14,274	7.7	112.9
4	Diabetes mellitus (E10-E14)	6,660	3.6	52.7
5	Cerebrovascular diseases (I60-I69)	6,396	3.5	50.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,934	3.2	46.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,004	2.2	31.7
8	Septicemia (A40-A41)	3,026	1.6	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,791	1.5	22.1
10	Influenza and pneumonia (J09-J18)	2,757	1.5	21.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,288	1.2	18.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,733	0.9	13.7
13	Alzheimer disease (G30)	1,665	0.9	13.2
14	Parkinson disease (G20-G21)	1,171	0.6	9.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,058	0.6	8.4
...	All other causes (residual)	29,301	15.8	231.8
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	223,431	100.0	2,240.1
1	Malignant neoplasms (C00-C97)	69,633	31.2	698.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,952	21.5	480.8
3	Chronic lower respiratory diseases (J40-J47)	20,522	9.2	205.8
4	Cerebrovascular diseases (I60-I69)	9,316	4.2	93.4
5	Diabetes mellitus (E10-E14)	7,570	3.4	75.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,715	2.6	57.3
7	Alzheimer disease (G30)	4,260	1.9	42.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,843	1.7	38.5
9	Influenza and pneumonia (J09-J18)	3,815	1.7	38.2
10	Septicemia (A40-A41)	3,634	1.6	36.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,086	1.4	30.9
12	Parkinson disease (G20-G21)	2,721	1.2	27.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,044	0.9	20.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,889	0.8	18.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,486	0.7	14.9
...	All other causes (residual)	35,945	16.1	360.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	245,250	100.0	3,603.5
1	Malignant neoplasms (C00-C97)	66,110	27.0	971.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,570	21.8	787.1
3	Chronic lower respiratory diseases (J40-J47)	22,472	9.2	330.2
4	Cerebrovascular diseases (I60-I69)	12,375	5.0	181.8
5	Alzheimer disease (G30)	9,358	3.8	137.5
6	Diabetes mellitus (E10-E14)	7,090	2.9	104.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,298	2.6	92.5
8	Parkinson disease (G20-G21)	4,663	1.9	68.5
9	Influenza and pneumonia (J09-J18)	4,611	1.9	67.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,492	1.8	66.0
11	Septicemia (A40-A41)	3,925	1.6	57.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,466	1.0	36.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,110	0.9	31.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,855	0.8	27.3
15	Pneumonitis due to solids and liquids (J69)	1,796	0.7	26.4
...	All other causes (residual)	42,059	17.1	618.0
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	284,866	100.0	6,078.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,865	23.5	1,426.8
2	Malignant neoplasms (C00-C97)	60,470	21.2	1,290.4
3	Chronic lower respiratory diseases (J40-J47)	22,944	8.1	489.6
4	Alzheimer disease (G30)	17,415	6.1	371.6
5	Cerebrovascular diseases (I60-I69)	17,116	6.0	365.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,777	2.7	166.0
7	Diabetes mellitus (E10-E14)	6,944	2.4	148.2
8	Parkinson disease (G20-G21)	6,676	2.3	142.5
9	Influenza and pneumonia (J09-J18)	6,191	2.2	132.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,648	2.0	120.5
11	Septicemia (A40-A41)	4,244	1.5	90.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,293	1.2	70.3
13	Pneumonitis due to solids and liquids (J69)	2,465	0.9	52.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,152	0.8	45.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,476	0.5	31.5
...	All other causes (residual)	53,190	18.7	1,135.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	326,484	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,501	26.5	---
2	Malignant neoplasms (C00-C97)	49,845	15.3	---
3	Alzheimer disease (G30)	26,706	8.2	---
4	Cerebrovascular diseases (I60-I69)	22,438	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	20,969	6.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,632	3.0	---
7	Influenza and pneumonia (J09-J18)	7,982	2.4	---
8	Parkinson disease (G20-G21)	6,861	2.1	---
9	Diabetes mellitus (E10-E14)	6,516	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,248	1.9	---
11	Septicemia (A40-A41)	4,512	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,254	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,342	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,301	0.7	---
15	Nutritional deficiencies (E40-E64)	1,232	0.4	---
...	All other causes (residual)	67,145	20.6	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	274,073	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,763	29.8	---
2	Malignant neoplasms (C00-C97)	28,122	10.3	---
3	Alzheimer disease (G30)	26,048	9.5	---
4	Cerebrovascular diseases (I60-I69)	20,192	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	13,763	5.0	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,133	3.0	---
7	Influenza and pneumonia (J09-J18)	7,642	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,147	1.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,364	1.6	---
10	Diabetes mellitus (E10-E14)	4,248	1.5	---
11	Parkinson disease (G20-G21)	3,802	1.4	---
12	Septicemia (A40-A41)	3,240	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,053	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,691	0.6	---
15	Nutritional deficiencies (E40-E64)	1,281	0.5	---
...	All other causes (residual)	61,584	22.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	121,331	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,815	32.0	---
2	Alzheimer disease (G30)	12,806	10.6	---
3	Cerebrovascular diseases (I60-I69)	8,905	7.3	---
4	Malignant neoplasms (C00-C97)	8,052	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	4,732	3.9	---
6	Influenza and pneumonia (J09-J18)	3,797	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,456	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,305	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,104	1.7	---
10	Diabetes mellitus (E10-E14)	1,461	1.2	---
11	Pneumonitis due to solids and liquids (J69)	1,418	1.2	---
12	Septicemia (A40-A41)	1,207	1.0	---
13	Parkinson disease (G20-G21)	959	0.8	---
14	Nutritional deficiencies (E40-E64)	808	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	591	0.5	---
...	All other causes (residual)	29,915	24.7	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	24,236	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,267	34.1	---
2	Alzheimer disease (G30)	2,592	10.7	---
3	Cerebrovascular diseases (I60-I69)	1,695	7.0	---
4	Malignant neoplasms (C00-C97)	937	3.9	---
5	Influenza and pneumonia (J09-J18)	894	3.7	---
6	Chronic lower respiratory diseases (J40-J47)	671	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	628	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	370	1.5	---
10	Pneumonitis due to solids and liquids (J69)	247	1.0	---
11	Septicemia (A40-A41)	204	0.8	---
12	Nutritional deficiencies (E40-E64)	202	0.8	---
13	Diabetes mellitus (E10-E14)	200	0.8	---
14	Atherosclerosis (I70)	136	0.6	---
15	Parkinson disease (G20-G21)	110	0.5	---
...	All other causes (residual)	6,441	26.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,102,838	100.0	1,111.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,428	24.7	274.5
2	Malignant neoplasms (C00-C97)	247,022	22.4	248.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,280	7.2	79.9
4	Chronic lower respiratory diseases (J40-J47)	64,673	5.9	65.2
5	Cerebrovascular diseases (I60-I69)	45,078	4.1	45.4
6	Alzheimer disease (G30)	31,577	2.9	31.8
7	Diabetes mellitus (E10-E14)	31,343	2.8	31.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,708	2.7	29.9
9	Influenza and pneumonia (J09-J18)	20,515	1.9	20.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,866	1.7	19.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,187	1.6	18.3
12	Parkinson disease (G20-G21)	16,841	1.5	17.0
13	Septicemia (A40-A41)	14,419	1.3	14.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,696	1.0	10.8
15	Pneumonitis due to solids and liquids (J69)	9,352	0.8	9.4
...	All other causes (residual)	192,853	17.5	194.3
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,107	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	38.4	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.1	2.4
3	Malignant neoplasms (C00-C97)	93	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.8	1.5
5	Influenza and pneumonia (J09-J18)	30	2.7	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	18	1.6	*
8	Septicemia (A40-A41)	12	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	295	26.6	6.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	623	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	33.5	3.9
2	Malignant neoplasms (C00-C97)	119	19.1	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	5.0	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.7	*
6	Influenza and pneumonia (J09-J18)	14	2.2	*
7	Cerebrovascular diseases (I60-I69)	12	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Septicemia (A40-A41)	5	0.8	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
12	Nutritional deficiencies (E40-E64)	2	0.3	*
12	Meningitis (G00,G03)	2	0.3	*
12	Pneumonitis due to solids and liquids (J69)	2	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (residual)	139	22.3	2.6
Non-Hispanic, white, male, 10-14 years				
...	All causes	1,005	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	30.3	5.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	22.5	4.0
3	Malignant neoplasms (C00-C97)	116	11.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	5.4	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	3.1	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.4	0.4
7	Cerebrovascular diseases (I60-I69)	15	1.5	*
8	Influenza and pneumonia (J09-J18)	11	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
10	Septicemia (A40-A41)	8	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Diabetes mellitus (E10-E14)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
...	All other causes (residual)	176	17.5	3.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,894	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,609	41.3	27.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,309	33.6	22.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	5.6	3.7
4	Malignant neoplasms (C00-C97)	165	4.2	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.5	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.2	0.8
7	Influenza and pneumonia (J09-J18)	18	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
9	Diabetes mellitus (E10-E14)	15	0.4	*
10	Cerebrovascular diseases (I60-I69)	13	0.3	*
11	Aortic aneurysm and dissection (I71)	10	0.3	*
12	Septicemia (A40-A41)	7	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
14	Legal intervention (Y35,Y89.0)	6	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (residual)	350	9.0	5.9
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,391	100.0	133.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,511	53.8	71.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,019	24.1	32.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	3.8	5.1
4	Malignant neoplasms (C00-C97)	248	3.0	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	2.3	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.8	1.0
7	Diabetes mellitus (E10-E14)	59	0.7	0.9
8	Influenza and pneumonia (J09-J18)	38	0.5	0.6
9	Chronic lower respiratory diseases (J40-J47)	27	0.3	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.3	0.4
11	Septicemia (A40-A41)	20	0.2	0.3
12	Cerebrovascular diseases (I60-I69)	19	0.2	*
13	Legal intervention (Y35,Y89.0)	16	0.2	*
14	Pneumonitis due to solids and liquids (J69)	14	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.1	*
...	All other causes (residual)	801	9.5	12.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	11,865	100.0	178.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,485	54.7	97.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,294	19.3	34.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	423	3.6	6.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	3.5	6.2
5	Malignant neoplasms (C00-C97)	368	3.1	5.5
6	Diabetes mellitus (E10-E14)	103	0.9	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	92	0.8	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	0.6	1.1
9	Cerebrovascular diseases (I60-I69)	65	0.5	1.0
10	Influenza and pneumonia (J09-J18)	52	0.4	0.8
11	Legal intervention (Y35,Y89.0)	43	0.4	0.6
12	Chronic lower respiratory diseases (J40-J47)	38	0.3	0.6
13	Septicemia (A40-A41)	34	0.3	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.2	*
15	Aortic aneurysm and dissection (I71)	19	0.2	*
...	All other causes (residual)	1,319	11.1	19.8
Non-Hispanic, white, male, 30-34 years				
...	All causes	13,495	100.0	212.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,796	50.4	107.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,233	16.5	35.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	850	6.3	13.4
4	Malignant neoplasms (C00-C97)	581	4.3	9.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	438	3.2	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	248	1.8	3.9
7	Diabetes mellitus (E10-E14)	131	1.0	2.1
8	Cerebrovascular diseases (I60-I69)	106	0.8	1.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	0.7	1.6
10	Septicemia (A40-A41)	68	0.5	1.1
11	Influenza and pneumonia (J09-J18)	65	0.5	1.0
12	Legal intervention (Y35,Y89.0)	42	0.3	0.7
13	Chronic lower respiratory diseases (J40-J47)	37	0.3	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.3	0.6
15	Aortic aneurysm and dissection (I71)	31	0.2	0.5
...	All other causes (residual)	1,732	12.8	27.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	14,806	100.0	241.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,171	41.7	100.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,197	14.8	35.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,477	10.0	24.0
4	Malignant neoplasms (C00-C97)	1,076	7.3	17.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	439	3.0	7.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	2.4	5.9
7	Diabetes mellitus (E10-E14)	247	1.7	4.0
8	Cerebrovascular diseases (I60-I69)	174	1.2	2.8
9	Influenza and pneumonia (J09-J18)	111	0.7	1.8
10	Septicemia (A40-A41)	108	0.7	1.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	82	0.6	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.5	1.3
13	Chronic lower respiratory diseases (J40-J47)	68	0.5	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.4	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	0.4	1.0
...	All other causes (residual)	2,089	14.1	34.0
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,468	100.0	289.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,954	30.1	87.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,599	15.8	45.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,188	13.3	38.4
4	Malignant neoplasms (C00-C97)	1,849	11.2	32.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	685	4.2	12.0
6	Diabetes mellitus (E10-E14)	402	2.4	7.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	310	1.9	5.4
8	Cerebrovascular diseases (I60-I69)	273	1.7	4.8
9	Septicemia (A40-A41)	145	0.9	2.5
10	Influenza and pneumonia (J09-J18)	133	0.8	2.3
11	Chronic lower respiratory diseases (J40-J47)	127	0.8	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	0.7	2.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	0.5	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.5	1.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	77	0.5	1.4
...	All other causes (residual)	2,439	14.8	42.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	25,965	100.0	399.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,494	21.2	84.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,074	19.5	78.0
3	Malignant neoplasms (C00-C97)	4,015	15.5	61.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,620	10.1	40.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,335	5.1	20.5
6	Diabetes mellitus (E10-E14)	801	3.1	12.3
7	Cerebrovascular diseases (I60-I69)	551	2.1	8.5
8	Chronic lower respiratory diseases (J40-J47)	339	1.3	5.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	1.3	5.0
10	Septicemia (A40-A41)	257	1.0	3.9
11	Influenza and pneumonia (J09-J18)	225	0.9	3.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	0.8	3.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	0.7	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	164	0.6	2.5
15	Aortic aneurysm and dissection (I71)	129	0.5	2.0
...	All other causes (residual)	4,241	16.3	65.2
Non-Hispanic, white, male, 50-54 years				
...	All causes	42,816	100.0	613.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,811	22.9	140.6
2	Malignant neoplasms (C00-C97)	8,960	20.9	128.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,148	14.4	88.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,711	6.3	38.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,259	5.3	32.4
6	Diabetes mellitus (E10-E14)	1,510	3.5	21.6
7	Chronic lower respiratory diseases (J40-J47)	1,107	2.6	15.9
8	Cerebrovascular diseases (I60-I69)	1,002	2.3	14.4
9	Septicemia (A40-A41)	539	1.3	7.7
10	Influenza and pneumonia (J09-J18)	431	1.0	6.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	1.0	5.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	360	0.8	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	0.8	4.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	297	0.7	4.3
15	Viral hepatitis (B15-B19)	273	0.6	3.9
...	All other causes (residual)	6,654	15.5	95.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	69,506	100.0	919.5
1	Malignant neoplasms (C00-C97)	18,397	26.5	243.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,568	23.8	219.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,494	9.3	85.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,250	4.7	43.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,957	4.3	39.1
6	Chronic lower respiratory diseases (J40-J47)	2,766	4.0	36.6
7	Diabetes mellitus (E10-E14)	2,477	3.6	32.8
8	Cerebrovascular diseases (I60-I69)	1,698	2.4	22.5
9	Septicemia (A40-A41)	875	1.3	11.6
10	Influenza and pneumonia (J09-J18)	802	1.2	10.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	728	1.0	9.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	619	0.9	8.2
13	Viral hepatitis (B15-B19)	547	0.8	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	0.5	4.3
15	Pneumonitis due to solids and liquids (J69)	298	0.4	3.9
...	All other causes (residual)	10,708	15.4	141.7
Non-Hispanic, white, male, 60-64 years				
...	All causes	91,792	100.0	1,305.4
1	Malignant neoplasms (C00-C97)	27,765	30.2	394.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,532	24.5	320.4
3	Chronic lower respiratory diseases (J40-J47)	5,130	5.6	73.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,039	5.5	71.7
5	Diabetes mellitus (E10-E14)	3,434	3.7	48.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,331	3.6	47.4
7	Cerebrovascular diseases (I60-I69)	2,622	2.9	37.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,446	2.7	34.8
9	Septicemia (A40-A41)	1,227	1.3	17.4
10	Influenza and pneumonia (J09-J18)	1,162	1.3	16.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	1.1	14.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	837	0.9	11.9
13	Viral hepatitis (B15-B19)	585	0.6	8.3
14	Pneumonitis due to solids and liquids (J69)	445	0.5	6.3
15	Aortic aneurysm and dissection (I71)	419	0.5	6.0
...	All other causes (residual)	13,774	15.0	195.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	108,920	100.0	1,799.6
1	Malignant neoplasms (C00-C97)	34,791	31.9	574.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,628	24.4	440.0
3	Chronic lower respiratory diseases (J40-J47)	7,339	6.7	121.3
4	Diabetes mellitus (E10-E14)	4,080	3.7	67.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,941	3.6	65.1
6	Cerebrovascular diseases (I60-I69)	3,505	3.2	57.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,696	2.5	44.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,767	1.6	29.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,600	1.5	26.4
10	Septicemia (A40-A41)	1,581	1.5	26.1
11	Influenza and pneumonia (J09-J18)	1,524	1.4	25.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,039	1.0	17.2
13	Parkinson disease (G20-G21)	748	0.7	12.4
14	Alzheimer disease (G30)	733	0.7	12.1
15	Pneumonitis due to solids and liquids (J69)	644	0.6	10.6
...	All other causes (residual)	16,304	15.0	269.4
Non-Hispanic, white, male, 70-74 years				
...	All causes	125,660	100.0	2,675.8
1	Malignant neoplasms (C00-C97)	38,394	30.6	817.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,436	24.2	648.1
3	Chronic lower respiratory diseases (J40-J47)	10,115	8.0	215.4
4	Cerebrovascular diseases (I60-I69)	4,820	3.8	102.6
5	Diabetes mellitus (E10-E14)	4,543	3.6	96.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,537	2.8	75.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,120	1.7	45.1
8	Influenza and pneumonia (J09-J18)	2,015	1.6	42.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,864	1.5	39.7
10	Parkinson disease (G20-G21)	1,841	1.5	39.2
11	Septicemia (A40-A41)	1,798	1.4	38.3
12	Alzheimer disease (G30)	1,793	1.4	38.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,530	1.2	32.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,118	0.9	23.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	878	0.7	18.7
...	All other causes (residual)	18,858	15.0	401.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	131,567	100.0	4,259.2
1	Malignant neoplasms (C00-C97)	35,860	27.3	1,160.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,791	24.2	1,029.2
3	Chronic lower respiratory diseases (J40-J47)	10,769	8.2	348.6
4	Cerebrovascular diseases (I60-I69)	6,010	4.6	194.6
5	Diabetes mellitus (E10-E14)	4,131	3.1	133.7
6	Alzheimer disease (G30)	3,654	2.8	118.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,646	2.8	118.0
8	Parkinson disease (G20-G21)	3,059	2.3	99.0
9	Influenza and pneumonia (J09-J18)	2,425	1.8	78.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,386	1.8	77.2
11	Septicemia (A40-A41)	1,930	1.5	62.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,247	0.9	40.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,239	0.9	40.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,204	0.9	39.0
15	Pneumonitis due to solids and liquids (J69)	1,115	0.8	36.1
...	All other causes (residual)	21,101	16.0	683.1
Non-Hispanic, white, male, 80-84 years				
...	All causes	142,527	100.0	7,102.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,165	25.4	1,802.2
2	Malignant neoplasms (C00-C97)	32,242	22.6	1,606.7
3	Chronic lower respiratory diseases (J40-J47)	10,397	7.3	518.1
4	Cerebrovascular diseases (I60-I69)	7,451	5.2	371.3
5	Alzheimer disease (G30)	6,474	4.5	322.6
6	Parkinson disease (G20-G21)	4,310	3.0	214.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,135	2.9	206.1
8	Diabetes mellitus (E10-E14)	3,829	2.7	190.8
9	Influenza and pneumonia (J09-J18)	3,124	2.2	155.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,005	2.1	149.7
11	Septicemia (A40-A41)	2,000	1.4	99.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,459	1.0	72.7
13	Pneumonitis due to solids and liquids (J69)	1,433	1.0	71.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,173	0.8	58.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	918	0.6	45.7
...	All other causes (residual)	24,412	17.1	1,216.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	146,555	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,321	28.2	---
2	Malignant neoplasms (C00-C97)	25,314	17.3	---
3	Chronic lower respiratory diseases (J40-J47)	9,195	6.3	---
4	Alzheimer disease (G30)	8,872	6.1	---
5	Cerebrovascular diseases (I60-I69)	8,432	5.8	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,691	3.2	---
7	Parkinson disease (G20-G21)	4,047	2.8	---
8	Influenza and pneumonia (J09-J18)	3,687	2.5	---
9	Diabetes mellitus (E10-E14)	3,235	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,197	2.2	---
11	Septicemia (A40-A41)	2,052	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,909	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,680	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,233	0.8	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	714	0.5	---
...	All other causes (residual)	26,976	18.4	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	101,497	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,521	32.0	---
2	Malignant neoplasms (C00-C97)	13,146	13.0	---
3	Alzheimer disease (G30)	6,870	6.8	---
4	Cerebrovascular diseases (I60-I69)	5,963	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	5,532	5.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,133	3.1	---
7	Influenza and pneumonia (J09-J18)	3,107	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,326	2.3	---
9	Parkinson disease (G20-G21)	1,972	1.9	---
10	Diabetes mellitus (E10-E14)	1,797	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,596	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,364	1.3	---
13	Septicemia (A40-A41)	1,274	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	746	0.7	---
15	Nutritional deficiencies (E40-E64)	403	0.4	---
...	All other causes (residual)	19,747	19.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	34,589	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,101	35.0	---
2	Malignant neoplasms (C00-C97)	3,213	9.3	---
3	Alzheimer disease (G30)	2,419	7.0	---
4	Cerebrovascular diseases (I60-I69)	2,043	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	1,499	4.3	---
6	Influenza and pneumonia (J09-J18)	1,281	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,055	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	800	2.3	---
9	Pneumonitis due to solids and liquids (J69)	636	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	532	1.5	---
11	Diabetes mellitus (E10-E14)	492	1.4	---
12	Parkinson disease (G20-G21)	402	1.2	---
13	Septicemia (A40-A41)	393	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	231	0.7	---
15	Nutritional deficiencies (E40-E64)	212	0.6	---
...	All other causes (residual)	7,280	21.0	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	4,462	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,693	37.9	---
2	Alzheimer disease (G30)	309	6.9	---
3	Malignant neoplasms (C00-C97)	290	6.5	---
4	Cerebrovascular diseases (I60-I69)	257	5.8	---
5	Influenza and pneumonia (J09-J18)	224	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	145	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.4	---
9	Pneumonitis due to solids and liquids (J69)	86	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.6	---
11	Septicemia (A40-A41)	56	1.3	---
12	Diabetes mellitus (E10-E14)	49	1.1	---
13	Parkinson disease (G20-G21)	44	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	---
15	Nutritional deficiencies (E40-E64)	23	0.5	---
...	All other causes (residual)	949	21.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,077,019	100.0	1,055.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,057	21.9	231.4
2	Malignant neoplasms (C00-C97)	218,657	20.3	214.3
3	Chronic lower respiratory diseases (J40-J47)	75,160	7.0	73.7
4	Alzheimer disease (G30)	70,299	6.5	68.9
5	Cerebrovascular diseases (I60-I69)	64,960	6.0	63.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,749	4.4	46.8
7	Diabetes mellitus (E10-E14)	23,773	2.2	23.3
8	Influenza and pneumonia (J09-J18)	22,882	2.1	22.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,004	1.6	16.7
10	Septicemia (A40-A41)	15,779	1.5	15.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,769	1.3	13.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,357	1.1	11.1
13	Parkinson disease (G20-G21)	10,741	1.0	10.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,398	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,097	0.7	7.0
...	All other causes (residual)	233,337	21.7	228.7
Non-Hispanic, white, female, 1-4 years				
...	All causes	751	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	33.6	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	10.9	2.0
3	Malignant neoplasms (C00-C97)	71	9.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	6.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.9	0.7
6	Influenza and pneumonia (J09-J18)	14	1.9	*
7	Septicemia (A40-A41)	12	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.5	*
8	Cerebrovascular diseases (I60-I69)	11	1.5	*
10	Anemias (D50-D64)	8	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
14	Hernia (K40-K46)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	184	24.5	4.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	492	100.0	9.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	28.3	2.7
2	Malignant neoplasms (C00-C97)	108	22.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	8.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	5.9	0.6
5	Influenza and pneumonia (J09-J18)	19	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.3	*
7	Cerebrovascular diseases (I60-I69)	10	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
10	Septicemia (A40-A41)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
12	Anemias (D50-D64)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (residual)	104	21.1	2.0
Non-Hispanic, white, female, 10-14 years				
...	All causes	662	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	25.1	3.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	15.3	1.9
3	Malignant neoplasms (C00-C97)	90	13.6	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	7.4	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.1	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	4.1	0.5
7	Influenza and pneumonia (J09-J18)	21	3.2	0.4
8	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
9	Cerebrovascular diseases (I60-I69)	8	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
12	Septicemia (A40-A41)	3	0.5	*
12	Diabetes mellitus (E10-E14)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	139	21.0	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,698	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	755	44.5	13.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	327	19.3	5.8
3	Malignant neoplasms (C00-C97)	111	6.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	3.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	3.0	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	2.7	0.8
7	Diabetes mellitus (E10-E14)	21	1.2	0.4
8	Cerebrovascular diseases (I60-I69)	16	0.9	*
8	Influenza and pneumonia (J09-J18)	16	0.9	*
10	Septicemia (A40-A41)	11	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
15	Anemias (D50-D64)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	239	14.1	4.2
Non-Hispanic, white, female, 20-24 years				
...	All causes	3,224	100.0	54.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,680	52.1	28.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	419	13.0	7.1
3	Malignant neoplasms (C00-C97)	162	5.0	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	5.0	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	3.0	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	58	1.8	1.0
7	Diabetes mellitus (E10-E14)	34	1.1	0.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.1	0.6
9	Influenza and pneumonia (J09-J18)	28	0.9	0.5
10	Septicemia (A40-A41)	27	0.8	0.5
11	Cerebrovascular diseases (I60-I69)	18	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (residual)	460	14.3	7.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	4,871	100.0	75.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,325	47.7	36.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	534	11.0	8.3
3	Malignant neoplasms (C00-C97)	327	6.7	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	4.8	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	3.2	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	118	2.4	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	64	1.3	1.0
8	Diabetes mellitus (E10-E14)	61	1.3	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	1.0	0.8
10	Septicemia (A40-A41)	49	1.0	0.8
11	Cerebrovascular diseases (I60-I69)	38	0.8	0.6
12	Chronic lower respiratory diseases (J40-J47)	36	0.7	0.6
13	Influenza and pneumonia (J09-J18)	33	0.7	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (residual)	820	16.8	12.8
Non-Hispanic, white, female, 30-34 years				
...	All causes	6,533	100.0	105.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,637	40.4	42.4
2	Malignant neoplasms (C00-C97)	716	11.0	11.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	625	9.6	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	453	6.9	7.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	2.9	3.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	178	2.7	2.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	126	1.9	2.0
8	Diabetes mellitus (E10-E14)	98	1.5	1.6
9	Influenza and pneumonia (J09-J18)	74	1.1	1.2
10	Cerebrovascular diseases (I60-I69)	71	1.1	1.1
11	Septicemia (A40-A41)	62	0.9	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	0.7	0.8
13	Chronic lower respiratory diseases (J40-J47)	35	0.5	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.4	0.4
...	All other causes (residual)	1,168	17.9	18.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	8,066	100.0	133.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,612	32.4	43.1
2	Malignant neoplasms (C00-C97)	1,302	16.1	21.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	696	8.6	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	646	8.0	10.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	310	3.8	5.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	198	2.5	3.3
7	Diabetes mellitus (E10-E14)	169	2.1	2.8
8	Cerebrovascular diseases (I60-I69)	148	1.8	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	108	1.3	1.8
10	Septicemia (A40-A41)	96	1.2	1.6
11	Influenza and pneumonia (J09-J18)	87	1.1	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	0.9	1.2
13	Chronic lower respiratory diseases (J40-J47)	74	0.9	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	0.7	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.3	0.4
...	All other causes (residual)	1,464	18.2	24.2
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,056	100.0	178.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,392	23.8	42.4
2	Malignant neoplasms (C00-C97)	2,246	22.3	39.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,143	11.4	20.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	733	7.3	13.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	454	4.5	8.0
6	Cerebrovascular diseases (I60-I69)	237	2.4	4.2
7	Diabetes mellitus (E10-E14)	236	2.3	4.2
8	Chronic lower respiratory diseases (J40-J47)	190	1.9	3.4
9	Influenza and pneumonia (J09-J18)	128	1.3	2.3
10	Septicemia (A40-A41)	127	1.3	2.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	1.2	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	0.7	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.6	1.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	44	0.4	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.4	0.7
...	All other causes (residual)	1,835	18.2	32.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	16,488	100.0	254.1
1	Malignant neoplasms (C00-C97)	4,508	27.3	69.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,719	16.5	41.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,122	12.9	32.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	930	5.6	14.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	787	4.8	12.1
6	Diabetes mellitus (E10-E14)	505	3.1	7.8
7	Cerebrovascular diseases (I60-I69)	428	2.6	6.6
8	Chronic lower respiratory diseases (J40-J47)	408	2.5	6.3
9	Septicemia (A40-A41)	249	1.5	3.8
10	Influenza and pneumonia (J09-J18)	227	1.4	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	1.0	2.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	0.9	2.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	0.7	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	0.5	1.2
15	Pneumonitis due to solids and liquids (J69)	72	0.4	1.1
...	All other causes (residual)	3,028	18.4	46.7
Non-Hispanic, white, female, 50-54 years				
...	All causes	27,496	100.0	387.8
1	Malignant neoplasms (C00-C97)	8,918	32.4	125.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,933	14.3	55.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,995	10.9	42.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,332	4.8	18.8
5	Chronic lower respiratory diseases (J40-J47)	1,278	4.6	18.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,000	3.6	14.1
7	Diabetes mellitus (E10-E14)	808	2.9	11.4
8	Cerebrovascular diseases (I60-I69)	732	2.7	10.3
9	Septicemia (A40-A41)	432	1.6	6.1
10	Influenza and pneumonia (J09-J18)	375	1.4	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	275	1.0	3.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	216	0.8	3.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	167	0.6	2.4
14	Viral hepatitis (B15-B19)	163	0.6	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	0.6	2.2
...	All other causes (residual)	4,718	17.2	66.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	44,578	100.0	569.7
1	Malignant neoplasms (C00-C97)	15,982	35.9	204.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,824	15.3	87.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,102	7.0	39.6
4	Chronic lower respiratory diseases (J40-J47)	3,024	6.8	38.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,698	3.8	21.7
6	Diabetes mellitus (E10-E14)	1,383	3.1	17.7
7	Cerebrovascular diseases (I60-I69)	1,328	3.0	17.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	986	2.2	12.6
9	Septicemia (A40-A41)	716	1.6	9.1
10	Influenza and pneumonia (J09-J18)	630	1.4	8.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	491	1.1	6.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	341	0.8	4.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	257	0.6	3.3
14	Viral hepatitis (B15-B19)	247	0.6	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.4	2.5
...	All other causes (residual)	7,371	16.5	94.2
Non-Hispanic, white, female, 60-64 years				
...	All causes	59,123	100.0	791.3
1	Malignant neoplasms (C00-C97)	21,940	37.1	293.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,591	16.2	128.4
3	Chronic lower respiratory diseases (J40-J47)	4,707	8.0	63.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,422	4.1	32.4
5	Cerebrovascular diseases (I60-I69)	1,943	3.3	26.0
6	Diabetes mellitus (E10-E14)	1,918	3.2	25.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,704	2.9	22.8
8	Septicemia (A40-A41)	1,061	1.8	14.2
9	Influenza and pneumonia (J09-J18)	905	1.5	12.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	776	1.3	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	773	1.3	10.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	482	0.8	6.5
13	Alzheimer disease (G30)	378	0.6	5.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	276	0.5	3.7
15	Viral hepatitis (B15-B19)	262	0.4	3.5
...	All other causes (residual)	9,985	16.9	133.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	76,076	100.0	1,154.9
1	Malignant neoplasms (C00-C97)	27,451	36.1	416.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,068	17.2	198.4
3	Chronic lower respiratory diseases (J40-J47)	6,935	9.1	105.3
4	Cerebrovascular diseases (I60-I69)	2,891	3.8	43.9
5	Diabetes mellitus (E10-E14)	2,580	3.4	39.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,993	2.6	30.3
7	Septicemia (A40-A41)	1,445	1.9	21.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,308	1.7	19.9
9	Influenza and pneumonia (J09-J18)	1,233	1.6	18.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,191	1.6	18.1
11	Alzheimer disease (G30)	932	1.2	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	694	0.9	10.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	521	0.7	7.9
14	Parkinson disease (G20-G21)	423	0.6	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	416	0.5	6.3
...	All other causes (residual)	12,995	17.1	197.3
Non-Hispanic, white, female, 70-74 years				
...	All causes	97,771	100.0	1,852.4
1	Malignant neoplasms (C00-C97)	31,239	32.0	591.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,516	17.9	331.9
3	Chronic lower respiratory diseases (J40-J47)	10,407	10.6	197.2
4	Cerebrovascular diseases (I60-I69)	4,496	4.6	85.2
5	Diabetes mellitus (E10-E14)	3,027	3.1	57.4
6	Alzheimer disease (G30)	2,467	2.5	46.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,178	2.2	41.3
8	Septicemia (A40-A41)	1,836	1.9	34.8
9	Influenza and pneumonia (J09-J18)	1,800	1.8	34.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,723	1.8	32.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,222	1.2	23.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	926	0.9	17.5
13	Parkinson disease (G20-G21)	880	0.9	16.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	608	0.6	11.5
15	Pneumonitis due to solids and liquids (J69)	488	0.5	9.2
...	All other causes (residual)	16,958	17.3	321.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	113,683	100.0	3,058.6
1	Malignant neoplasms (C00-C97)	30,250	26.6	813.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,779	19.2	586.0
3	Chronic lower respiratory diseases (J40-J47)	11,703	10.3	314.9
4	Cerebrovascular diseases (I60-I69)	6,365	5.6	171.2
5	Alzheimer disease (G30)	5,704	5.0	153.5
6	Diabetes mellitus (E10-E14)	2,959	2.6	79.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,652	2.3	71.4
8	Influenza and pneumonia (J09-J18)	2,186	1.9	58.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,106	1.9	56.7
10	Septicemia (A40-A41)	1,995	1.8	53.7
11	Parkinson disease (G20-G21)	1,604	1.4	43.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,219	1.1	32.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	871	0.8	23.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	778	0.7	20.9
15	Pneumonitis due to solids and liquids (J69)	681	0.6	18.3
...	All other causes (residual)	20,831	18.3	560.5
Non-Hispanic, white, female, 80-84 years				
...	All causes	142,339	100.0	5,312.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,700	21.6	1,145.7
2	Malignant neoplasms (C00-C97)	28,228	19.8	1,053.5
3	Chronic lower respiratory diseases (J40-J47)	12,547	8.8	468.3
4	Alzheimer disease (G30)	10,941	7.7	408.3
5	Cerebrovascular diseases (I60-I69)	9,665	6.8	360.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,642	2.6	135.9
7	Diabetes mellitus (E10-E14)	3,115	2.2	116.3
8	Influenza and pneumonia (J09-J18)	3,067	2.2	114.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,643	1.9	98.6
10	Parkinson disease (G20-G21)	2,366	1.7	88.3
11	Septicemia (A40-A41)	2,244	1.6	83.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,834	1.3	68.4
13	Pneumonitis due to solids and liquids (J69)	1,032	0.7	38.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	979	0.7	36.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	694	0.5	25.9
...	All other causes (residual)	28,642	20.1	1,068.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	179,929	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,180	25.1	---
2	Malignant neoplasms (C00-C97)	24,531	13.6	---
3	Alzheimer disease (G30)	17,834	9.9	---
4	Cerebrovascular diseases (I60-I69)	14,006	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	11,774	6.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,941	2.7	---
7	Influenza and pneumonia (J09-J18)	4,295	2.4	---
8	Diabetes mellitus (E10-E14)	3,281	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,051	1.7	---
10	Parkinson disease (G20-G21)	2,814	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,574	1.4	---
12	Septicemia (A40-A41)	2,460	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,433	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,068	0.6	---
15	Nutritional deficiencies (E40-E64)	807	0.4	---
...	All other causes (residual)	39,880	22.2	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	172,576	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,242	28.5	---
2	Alzheimer disease (G30)	19,178	11.1	---
3	Malignant neoplasms (C00-C97)	14,976	8.7	---
4	Cerebrovascular diseases (I60-I69)	14,229	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	8,231	4.8	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,000	2.9	---
7	Influenza and pneumonia (J09-J18)	4,535	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,000	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,821	1.6	---
10	Diabetes mellitus (E10-E14)	2,451	1.4	---
11	Septicemia (A40-A41)	1,966	1.1	---
12	Parkinson disease (G20-G21)	1,830	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,457	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	945	0.5	---
15	Nutritional deficiencies (E40-E64)	878	0.5	---
...	All other causes (residual)	41,837	24.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	86,742	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,714	30.8	---
2	Alzheimer disease (G30)	10,387	12.0	---
3	Cerebrovascular diseases (I60-I69)	6,862	7.9	---
4	Malignant neoplasms (C00-C97)	4,839	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	3,233	3.7	---
6	Influenza and pneumonia (J09-J18)	2,516	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,401	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,773	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,304	1.5	---
10	Diabetes mellitus (E10-E14)	969	1.1	---
11	Septicemia (A40-A41)	814	0.9	---
12	Pneumonitis due to solids and liquids (J69)	782	0.9	---
13	Nutritional deficiencies (E40-E64)	596	0.7	---
14	Parkinson disease (G20-G21)	557	0.6	---
15	Atherosclerosis (I70)	416	0.5	---
...	All other causes (residual)	22,579	26.0	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	19,774	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,574	33.2	---
2	Alzheimer disease (G30)	2,283	11.5	---
3	Cerebrovascular diseases (I60-I69)	1,438	7.3	---
4	Influenza and pneumonia (J09-J18)	670	3.4	---
5	Malignant neoplasms (C00-C97)	647	3.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	556	2.8	---
7	Chronic lower respiratory diseases (J40-J47)	526	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	510	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.3	---
10	Nutritional deficiencies (E40-E64)	179	0.9	---
11	Pneumonitis due to solids and liquids (J69)	161	0.8	---
12	Diabetes mellitus (E10-E14)	151	0.8	---
13	Septicemia (A40-A41)	148	0.7	---
14	Atherosclerosis (I70)	119	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.3	---
...	All other causes (residual)	5,478	27.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	335,667	100.0	787.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,161	23.3	183.4
2	Malignant neoplasms (C00-C97)	69,872	20.8	163.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,869	5.9	46.6
4	Cerebrovascular diseases (I60-I69)	19,088	5.7	44.8
5	Diabetes mellitus (E10-E14)	14,798	4.4	34.7
6	Chronic lower respiratory diseases (J40-J47)	11,217	3.3	26.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,908	3.0	23.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,542	2.8	22.4
9	Alzheimer disease (G30)	8,991	2.7	21.1
10	Septicemia (A40-A41)	6,568	2.0	15.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,273	1.9	14.7
12	Influenza and pneumonia (J09-J18)	5,556	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,637	1.1	8.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,290	1.0	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,965	0.9	7.0
...	All other causes (residual)	65,932	19.6	154.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	985	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	29.9	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	12.7	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	10.3	4.1
4	Malignant neoplasms (C00-C97)	42	4.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.8	1.5
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Cerebrovascular diseases (I60-I69)	21	2.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	18	1.8	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
11	Septicemia (A40-A41)	9	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Meningitis (G00,G03)	6	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
...	All other causes (residual)	259	26.3	10.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	573	100.0	18.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	31.6	5.8
2	Malignant neoplasms (C00-C97)	66	11.5	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	10.3	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	6.8	1.3
5	Chronic lower respiratory diseases (J40-J47)	37	6.5	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.0	*
7	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Cerebrovascular diseases (I60-I69)	11	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
10	Septicemia (A40-A41)	7	1.2	*
11	Anemias (D50-D64)	5	0.9	*
12	Diabetes mellitus (E10-E14)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	117	20.4	3.8
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	723	100.0	23.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	27.1	6.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	11.5	2.7
3	Malignant neoplasms (C00-C97)	70	9.7	2.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	8.7	2.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	5.7	1.3
6	Chronic lower respiratory diseases (J40-J47)	40	5.5	1.3
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.1	1.0
8	Cerebrovascular diseases (I60-I69)	14	1.9	*
9	Diabetes mellitus (E10-E14)	13	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
11	Influenza and pneumonia (J09-J18)	9	1.2	*
12	Septicemia (A40-A41)	7	1.0	*
12	Anemias (D50-D64)	7	1.0	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	134	18.5	4.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,560	100.0	80.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,135	44.3	35.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	585	22.9	18.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	9.9	7.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.6	2.9
5	Malignant neoplasms (C00-C97)	86	3.4	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.4	1.1
7	Diabetes mellitus (E10-E14)	29	1.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	26	1.0	0.8
9	Cerebrovascular diseases (I60-I69)	15	0.6	*
10	Influenza and pneumonia (J09-J18)	14	0.5	*
11	Legal intervention (Y35,Y89.0)	13	0.5	*
12	Septicemia (A40-A41)	12	0.5	*
12	Anemias (D50-D64)	12	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.3	*
...	All other causes (residual)	237	9.3	7.4
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,804	100.0	140.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,920	40.0	56.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,245	25.9	36.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	457	9.5	13.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	4.2	5.9
5	Malignant neoplasms (C00-C97)	133	2.8	3.9
6	Chronic lower respiratory diseases (J40-J47)	56	1.2	1.6
7	Diabetes mellitus (E10-E14)	54	1.1	1.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.1	1.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.9	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.9	1.3
11	Anemias (D50-D64)	32	0.7	0.9
12	Influenza and pneumonia (J09-J18)	27	0.6	0.8
13	Legal intervention (Y35,Y89.0)	24	0.5	0.7
14	Cerebrovascular diseases (I60-I69)	23	0.5	0.7
15	Septicemia (A40-A41)	17	0.4	*
...	All other causes (residual)	476	9.9	14.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	6,066	100.0	171.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,918	31.6	54.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,623	26.8	45.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	439	7.2	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	379	6.2	10.7
5	Malignant neoplasms (C00-C97)	245	4.0	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	142	2.3	4.0
7	Diabetes mellitus (E10-E14)	125	2.1	3.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	88	1.5	2.5
9	Anemias (D50-D64)	55	0.9	1.6
9	Cerebrovascular diseases (I60-I69)	55	0.9	1.6
11	Septicemia (A40-A41)	52	0.9	1.5
11	Chronic lower respiratory diseases (J40-J47)	52	0.9	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.7	1.2
14	Influenza and pneumonia (J09-J18)	40	0.7	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.7	1.1
...	All other causes (residual)	770	12.7	21.8
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	6,018	100.0	203.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,447	24.0	49.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,278	21.2	43.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	679	11.3	23.0
4	Malignant neoplasms (C00-C97)	408	6.8	13.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	354	5.9	12.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	193	3.2	6.5
7	Diabetes mellitus (E10-E14)	156	2.6	5.3
8	Cerebrovascular diseases (I60-I69)	87	1.4	2.9
9	Chronic lower respiratory diseases (J40-J47)	76	1.3	2.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.2	2.5
11	Anemias (D50-D64)	71	1.2	2.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.0	2.0
13	Influenza and pneumonia (J09-J18)	56	0.9	1.9
14	Septicemia (A40-A41)	46	0.8	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	43	0.7	1.5
...	All other causes (residual)	992	16.5	33.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	7,251	100.0	257.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,557	21.5	55.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,098	15.1	39.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,025	14.1	36.5
4	Malignant neoplasms (C00-C97)	738	10.2	26.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	297	4.1	10.6
6	Diabetes mellitus (E10-E14)	271	3.7	9.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	212	2.9	7.5
8	Cerebrovascular diseases (I60-I69)	185	2.6	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.3	3.3
10	Septicemia (A40-A41)	92	1.3	3.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.2	3.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	83	1.1	3.0
13	Chronic lower respiratory diseases (J40-J47)	72	1.0	2.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.9	2.4
15	Influenza and pneumonia (J09-J18)	65	0.9	2.3
...	All other causes (residual)	1,308	18.0	46.5
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,475	100.0	332.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,741	20.5	68.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,394	16.4	54.7
3	Malignant neoplasms (C00-C97)	1,188	14.0	46.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	637	7.5	25.0
5	Diabetes mellitus (E10-E14)	375	4.4	14.7
6	Cerebrovascular diseases (I60-I69)	290	3.4	11.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	237	2.8	9.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	2.6	8.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.0	6.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	136	1.6	5.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.6	5.3
12	Septicemia (A40-A41)	123	1.5	4.8
13	Influenza and pneumonia (J09-J18)	105	1.2	4.1
14	Chronic lower respiratory diseases (J40-J47)	90	1.1	3.5
15	Anemias (D50-D64)	67	0.8	2.6
...	All other causes (residual)	1,568	18.5	61.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	12,288	100.0	461.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,858	23.3	107.3
2	Malignant neoplasms (C00-C97)	2,061	16.8	77.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,605	13.1	60.3
4	Diabetes mellitus (E10-E14)	577	4.7	21.7
5	Cerebrovascular diseases (I60-I69)	527	4.3	19.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	4.0	18.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	336	2.7	12.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	2.3	10.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	254	2.1	9.5
10	Septicemia (A40-A41)	229	1.9	8.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.8	8.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	1.7	8.0
13	Chronic lower respiratory diseases (J40-J47)	183	1.5	6.9
14	Influenza and pneumonia (J09-J18)	159	1.3	6.0
15	Aortic aneurysm and dissection (I71)	93	0.8	3.5
...	All other causes (residual)	2,191	17.8	82.3
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	18,938	100.0	703.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,594	24.3	170.7
2	Malignant neoplasms (C00-C97)	4,199	22.2	156.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,952	10.3	72.5
4	Diabetes mellitus (E10-E14)	919	4.9	34.1
5	Cerebrovascular diseases (I60-I69)	875	4.6	32.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	461	2.4	17.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	437	2.3	16.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	405	2.1	15.0
9	Chronic lower respiratory diseases (J40-J47)	394	2.1	14.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	2.0	13.9
11	Septicemia (A40-A41)	365	1.9	13.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	1.7	12.2
13	Influenza and pneumonia (J09-J18)	245	1.3	9.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	186	1.0	6.9
15	Aortic aneurysm and dissection (I71)	69	0.4	2.6
...	All other causes (residual)	3,134	16.5	116.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	28,790	100.0	1,088.2
1	Malignant neoplasms (C00-C97)	7,601	26.4	287.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,109	24.7	268.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,077	7.2	78.5
4	Diabetes mellitus (E10-E14)	1,399	4.9	52.9
5	Cerebrovascular diseases (I60-I69)	1,367	4.7	51.7
6	Chronic lower respiratory diseases (J40-J47)	794	2.8	30.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	759	2.6	28.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	677	2.4	25.6
9	Septicemia (A40-A41)	533	1.9	20.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	503	1.7	19.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	478	1.7	18.1
12	Influenza and pneumonia (J09-J18)	402	1.4	15.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	1.1	11.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	0.5	5.6
15	Viral hepatitis (B15-B19)	144	0.5	5.4
...	All other causes (residual)	4,487	15.6	169.6
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	35,708	100.0	1,583.5
1	Malignant neoplasms (C00-C97)	10,149	28.4	450.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,019	25.3	400.0
3	Cerebrovascular diseases (I60-I69)	1,795	5.0	79.6
4	Diabetes mellitus (E10-E14)	1,794	5.0	79.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,642	4.6	72.8
6	Chronic lower respiratory diseases (J40-J47)	1,276	3.6	56.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,039	2.9	46.1
8	Septicemia (A40-A41)	768	2.2	34.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	667	1.9	29.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	662	1.9	29.4
11	Influenza and pneumonia (J09-J18)	486	1.4	21.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	408	1.1	18.1
13	Viral hepatitis (B15-B19)	236	0.7	10.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	198	0.6	8.8
15	Pneumonitis due to solids and liquids (J69)	159	0.4	7.1
...	All other causes (residual)	5,410	15.2	239.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	37,568	100.0	2,168.8
1	Malignant neoplasms (C00-C97)	11,181	29.8	645.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,343	24.9	539.4
3	Cerebrovascular diseases (I60-I69)	2,043	5.4	117.9
4	Diabetes mellitus (E10-E14)	2,024	5.4	116.8
5	Chronic lower respiratory diseases (J40-J47)	1,519	4.0	87.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,181	3.1	68.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,145	3.0	66.1
8	Septicemia (A40-A41)	855	2.3	49.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	661	1.8	38.2
10	Influenza and pneumonia (J09-J18)	605	1.6	34.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	472	1.3	27.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	242	0.6	14.0
13	Alzheimer disease (G30)	239	0.6	13.8
14	Pneumonitis due to solids and liquids (J69)	200	0.5	11.5
15	Viral hepatitis (B15-B19)	198	0.5	11.4
...	All other causes (residual)	5,660	15.1	326.7
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	33,453	100.0	2,899.5
1	Malignant neoplasms (C00-C97)	9,429	28.2	817.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,350	25.0	723.7
3	Cerebrovascular diseases (I60-I69)	2,126	6.4	184.3
4	Diabetes mellitus (E10-E14)	1,770	5.3	153.4
5	Chronic lower respiratory diseases (J40-J47)	1,562	4.7	135.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,172	3.5	101.6
7	Septicemia (A40-A41)	788	2.4	68.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	643	1.9	55.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	623	1.9	54.0
10	Influenza and pneumonia (J09-J18)	564	1.7	48.9
11	Alzheimer disease (G30)	499	1.5	43.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	255	0.8	22.1
13	Pneumonitis due to solids and liquids (J69)	218	0.7	18.9
14	Parkinson disease (G20-G21)	205	0.6	17.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	13.6
...	All other causes (residual)	5,092	15.2	441.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	32,828	100.0	4,258.0
1	Malignant neoplasms (C00-C97)	8,156	24.8	1,057.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,106	24.7	1,051.4
3	Cerebrovascular diseases (I60-I69)	2,270	6.9	294.4
4	Diabetes mellitus (E10-E14)	1,712	5.2	222.1
5	Chronic lower respiratory diseases (J40-J47)	1,528	4.7	198.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,190	3.6	154.3
7	Alzheimer disease (G30)	1,055	3.2	136.8
8	Septicemia (A40-A41)	789	2.4	102.3
9	Influenza and pneumonia (J09-J18)	634	1.9	82.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	629	1.9	81.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	516	1.6	66.9
12	Parkinson disease (G20-G21)	302	0.9	39.2
13	Pneumonitis due to solids and liquids (J69)	240	0.7	31.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	183	0.6	23.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.4	17.8
...	All other causes (residual)	5,381	16.4	697.9
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	32,248	100.0	6,363.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,128	25.2	1,603.9
2	Malignant neoplasms (C00-C97)	6,611	20.5	1,304.6
3	Cerebrovascular diseases (I60-I69)	2,440	7.6	481.5
4	Alzheimer disease (G30)	1,713	5.3	338.0
5	Diabetes mellitus (E10-E14)	1,469	4.6	289.9
6	Chronic lower respiratory diseases (J40-J47)	1,384	4.3	273.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,190	3.7	234.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	712	2.2	140.5
9	Septicemia (A40-A41)	696	2.2	137.3
10	Influenza and pneumonia (J09-J18)	685	2.1	135.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	499	1.5	98.5
12	Parkinson disease (G20-G21)	343	1.1	67.7
13	Pneumonitis due to solids and liquids (J69)	241	0.7	47.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	163	0.5	32.2
15	Nutritional deficiencies (E40-E64)	129	0.4	25.5
...	All other causes (residual)	5,845	18.1	1,153.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	27,936	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,360	26.3	---
2	Malignant neoplasms (C00-C97)	4,408	15.8	---
3	Cerebrovascular diseases (I60-I69)	2,300	8.2	---
4	Alzheimer disease (G30)	2,093	7.5	---
5	Diabetes mellitus (E10-E14)	1,132	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,113	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	950	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	711	2.5	---
9	Influenza and pneumonia (J09-J18)	610	2.2	---
10	Septicemia (A40-A41)	582	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	1.5	---
12	Pneumonitis due to solids and liquids (J69)	291	1.0	---
13	Parkinson disease (G20-G21)	283	1.0	---
14	Nutritional deficiencies (E40-E64)	138	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.5	---
...	All other causes (residual)	5,430	19.4	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	19,883	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,434	27.3	---
2	Malignant neoplasms (C00-C97)	2,276	11.4	---
3	Alzheimer disease (G30)	1,941	9.8	---
4	Cerebrovascular diseases (I60-I69)	1,663	8.4	---
5	Chronic lower respiratory diseases (J40-J47)	703	3.5	---
6	Diabetes mellitus (E10-E14)	685	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	581	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	577	2.9	---
9	Influenza and pneumonia (J09-J18)	475	2.4	---
10	Septicemia (A40-A41)	347	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	1.3	---
12	Pneumonitis due to solids and liquids (J69)	186	0.9	---
13	Parkinson disease (G20-G21)	152	0.8	---
14	Nutritional deficiencies (E40-E64)	131	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.5	---
...	All other causes (residual)	4,384	22.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	9,110	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,657	29.2	---
2	Alzheimer disease (G30)	1,000	11.0	---
3	Cerebrovascular diseases (I60-I69)	725	8.0	---
4	Malignant neoplasms (C00-C97)	686	7.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	275	3.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	261	2.9	---
7	Diabetes mellitus (E10-E14)	239	2.6	---
8	Chronic lower respiratory diseases (J40-J47)	237	2.6	---
9	Influenza and pneumonia (J09-J18)	208	2.3	---
10	Septicemia (A40-A41)	160	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	1.4	---
12	Nutritional deficiencies (E40-E64)	86	0.9	---
13	Pneumonitis due to solids and liquids (J69)	78	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	---
15	Atherosclerosis (I70)	40	0.4	---
...	All other causes (residual)	2,290	25.1	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,716	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	830	30.6	---
2	Alzheimer disease (G30)	324	11.9	---
3	Cerebrovascular diseases (I60-I69)	219	8.1	---
4	Malignant neoplasms (C00-C97)	130	4.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	3.5	---
6	Influenza and pneumonia (J09-J18)	73	2.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.2	---
8	Chronic lower respiratory diseases (J40-J47)	52	1.9	---
9	Diabetes mellitus (E10-E14)	49	1.8	---
10	Septicemia (A40-A41)	40	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	1.4	---
12	Nutritional deficiencies (E40-E64)	28	1.0	---
13	Pneumonitis due to solids and liquids (J69)	26	1.0	---
14	Atherosclerosis (I70)	10	0.4	---
15	Anemias (D50-D64)	8	0.3	---
...	All other causes (residual)	735	27.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	174,403	100.0	854.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,411	23.7	202.8
2	Malignant neoplasms (C00-C97)	35,261	20.2	172.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,844	7.9	67.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,643	5.0	42.3
5	Cerebrovascular diseases (I60-I69)	8,566	4.9	42.0
6	Diabetes mellitus (E10-E14)	7,494	4.3	36.7
7	Chronic lower respiratory diseases (J40-J47)	5,605	3.2	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,555	2.6	22.3
9	Septicemia (A40-A41)	3,052	1.7	14.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,874	1.6	14.1
11	Influenza and pneumonia (J09-J18)	2,682	1.5	13.1
12	Alzheimer disease (G30)	2,476	1.4	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,324	1.3	11.4
14	Certain conditions originating in the perinatal period (P00-P96)	2,080	1.2	10.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,064	1.2	10.1
...	All other causes (residual)	31,472	18.0	154.2
Non-Hispanic, black, male, 1-4 years				
...	All causes	535	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.8	13.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	12.9	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Malignant neoplasms (C00-C97)	23	4.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.9	1.7
6	Influenza and pneumonia (J09-J18)	13	2.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
11	Septicemia (A40-A41)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Anemias (D50-D64)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (residual)	144	26.9	11.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	319	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	34.8	7.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	9.4	1.9
3	Malignant neoplasms (C00-C97)	29	9.1	1.8
4	Chronic lower respiratory diseases (J40-J47)	25	7.8	1.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	6.6	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.5	*
6	Influenza and pneumonia (J09-J18)	8	2.5	*
8	Cerebrovascular diseases (I60-I69)	5	1.6	*
9	Septicemia (A40-A41)	4	1.3	*
9	Anemias (D50-D64)	4	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Diabetes mellitus (E10-E14)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	63	19.7	4.0
Non-Hispanic, black, male, 10-14 years				
...	All causes	465	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	29.2	8.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	12.3	3.6
3	Malignant neoplasms (C00-C97)	44	9.5	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	9.5	2.8
5	Chronic lower respiratory diseases (J40-J47)	29	6.2	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	5.2	1.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.2	*
8	Cerebrovascular diseases (I60-I69)	10	2.2	*
9	Diabetes mellitus (E10-E14)	8	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
11	Septicemia (A40-A41)	4	0.9	*
12	Anemias (D50-D64)	3	0.6	*
12	Influenza and pneumonia (J09-J18)	3	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (residual)	78	16.8	5.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	2,019	100.0	124.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,022	50.6	62.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	439	21.7	27.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	9.4	11.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.3	4.1
5	Malignant neoplasms (C00-C97)	47	2.3	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.1	1.4
7	Diabetes mellitus (E10-E14)	19	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	18	0.9	*
9	Legal intervention (Y35,Y89.0)	13	0.6	*
10	Cerebrovascular diseases (I60-I69)	11	0.5	*
11	Influenza and pneumonia (J09-J18)	9	0.4	*
12	Septicemia (A40-A41)	8	0.4	*
13	Anemias (D50-D64)	4	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.1	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.1	*
...	All other causes (residual)	144	7.1	8.9
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,692	100.0	213.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,739	47.1	100.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	889	24.1	51.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	10.2	21.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	3.6	7.6
5	Malignant neoplasms (C00-C97)	80	2.2	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.9	2.0
7	Chronic lower respiratory diseases (J40-J47)	34	0.9	2.0
8	Diabetes mellitus (E10-E14)	28	0.8	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	1.4
10	Legal intervention (Y35,Y89.0)	23	0.6	1.3
11	Influenza and pneumonia (J09-J18)	16	0.4	*
12	Anemias (D50-D64)	13	0.4	*
13	Septicemia (A40-A41)	8	0.2	*
13	Cerebrovascular diseases (I60-I69)	8	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (residual)	280	7.6	16.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	4,454	100.0	252.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,730	38.8	98.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,223	27.5	69.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	366	8.2	20.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	5.4	13.6
5	Malignant neoplasms (C00-C97)	115	2.6	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	101	2.3	5.7
7	Diabetes mellitus (E10-E14)	65	1.5	3.7
8	Chronic lower respiratory diseases (J40-J47)	36	0.8	2.0
9	Anemias (D50-D64)	30	0.7	1.7
10	Legal intervention (Y35,Y89.0)	28	0.6	1.6
11	Septicemia (A40-A41)	26	0.6	1.5
12	Cerebrovascular diseases (I60-I69)	23	0.5	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.5	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.5	1.2
15	Influenza and pneumonia (J09-J18)	19	0.4	*
...	All other causes (residual)	407	9.1	23.1
Non-Hispanic, black, male, 30-34 years				
...	All causes	4,098	100.0	286.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,135	27.7	79.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,048	25.6	73.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	433	10.6	30.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	286	7.0	20.0
5	Malignant neoplasms (C00-C97)	179	4.4	12.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	130	3.2	9.1
7	Diabetes mellitus (E10-E14)	76	1.9	5.3
8	Cerebrovascular diseases (I60-I69)	44	1.1	3.1
8	Chronic lower respiratory diseases (J40-J47)	44	1.1	3.1
10	Anemias (D50-D64)	40	1.0	2.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.8	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.7	2.1
13	Influenza and pneumonia (J09-J18)	27	0.7	1.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.6	1.7
15	Septicemia (A40-A41)	23	0.6	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	1.6
...	All other causes (residual)	524	12.8	36.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	4,663	100.0	349.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,147	24.6	85.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	910	19.5	68.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	14.3	50.1
4	Malignant neoplasms (C00-C97)	301	6.5	22.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	4.9	17.2
6	Diabetes mellitus (E10-E14)	164	3.5	12.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	131	2.8	9.8
8	Cerebrovascular diseases (I60-I69)	99	2.1	7.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.1	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.1	3.7
11	Septicemia (A40-A41)	46	1.0	3.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	45	1.0	3.4
13	Influenza and pneumonia (J09-J18)	43	0.9	3.2
14	Anemias (D50-D64)	36	0.8	2.7
14	Chronic lower respiratory diseases (J40-J47)	36	0.8	2.7
...	All other causes (residual)	706	15.1	52.8
Non-Hispanic, black, male, 40-44 years				
...	All causes	5,058	100.0	422.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,118	22.1	93.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,041	20.6	86.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	10.9	45.8
4	Malignant neoplasms (C00-C97)	450	8.9	37.6
5	Diabetes mellitus (E10-E14)	216	4.3	18.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	3.2	13.4
7	Cerebrovascular diseases (I60-I69)	154	3.0	12.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	136	2.7	11.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.8	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.8	7.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.4	5.8
12	Septicemia (A40-A41)	67	1.3	5.6
13	Influenza and pneumonia (J09-J18)	53	1.0	4.4
14	Aortic aneurysm and dissection (I71)	50	1.0	4.2
15	Chronic lower respiratory diseases (J40-J47)	46	0.9	3.8
...	All other causes (residual)	768	15.2	64.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,229	100.0	578.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,814	25.1	145.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,151	15.9	92.1
3	Malignant neoplasms (C00-C97)	849	11.7	67.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	429	5.9	34.3
5	Diabetes mellitus (E10-E14)	347	4.8	27.8
6	Cerebrovascular diseases (I60-I69)	293	4.1	23.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	193	2.7	15.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	2.3	13.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	2.2	13.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	146	2.0	11.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	1.9	10.8
12	Septicemia (A40-A41)	120	1.7	9.6
13	Chronic lower respiratory diseases (J40-J47)	94	1.3	7.5
14	Influenza and pneumonia (J09-J18)	85	1.2	6.8
15	Aortic aneurysm and dissection (I71)	72	1.0	5.8
...	All other causes (residual)	1,171	16.2	93.7
Non-Hispanic, black, male, 50-54 years				
...	All causes	10,952	100.0	869.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,946	26.9	234.0
2	Malignant neoplasms (C00-C97)	1,918	17.5	152.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,348	12.3	107.1
4	Diabetes mellitus (E10-E14)	534	4.9	42.4
5	Cerebrovascular diseases (I60-I69)	508	4.6	40.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	2.9	25.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	282	2.6	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	2.4	20.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	232	2.1	18.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.8	15.3
11	Septicemia (A40-A41)	187	1.7	14.9
12	Chronic lower respiratory diseases (J40-J47)	166	1.5	13.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	1.4	12.3
14	Influenza and pneumonia (J09-J18)	133	1.2	10.6
15	Viral hepatitis (B15-B19)	46	0.4	3.7
...	All other causes (residual)	1,724	15.7	136.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	16,723	100.0	1,365.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,535	27.1	370.4
2	Malignant neoplasms (C00-C97)	3,816	22.8	311.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,532	9.2	125.1
4	Diabetes mellitus (E10-E14)	813	4.9	66.4
5	Cerebrovascular diseases (I60-I69)	758	4.5	61.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	423	2.5	34.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.5	33.6
8	Chronic lower respiratory diseases (J40-J47)	371	2.2	30.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	332	2.0	27.1
10	Septicemia (A40-A41)	288	1.7	23.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	277	1.7	22.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	253	1.5	20.7
13	Influenza and pneumonia (J09-J18)	212	1.3	17.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	0.6	8.7
15	Viral hepatitis (B15-B19)	100	0.6	8.2
...	All other causes (residual)	2,496	14.9	203.8
Non-Hispanic, black, male, 60-64 years				
...	All causes	20,829	100.0	2,051.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,658	27.2	557.3
2	Malignant neoplasms (C00-C97)	5,505	26.4	542.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,209	5.8	119.1
4	Cerebrovascular diseases (I60-I69)	1,049	5.0	103.3
5	Diabetes mellitus (E10-E14)	1,020	4.9	100.5
6	Chronic lower respiratory diseases (J40-J47)	645	3.1	63.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	536	2.6	52.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	466	2.2	45.9
9	Septicemia (A40-A41)	390	1.9	38.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.8	37.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	285	1.4	28.1
12	Influenza and pneumonia (J09-J18)	271	1.3	26.7
13	Viral hepatitis (B15-B19)	171	0.8	16.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	0.8	16.2
15	Pneumonitis due to solids and liquids (J69)	97	0.5	9.6
...	All other causes (residual)	2,986	14.3	294.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	21,346	100.0	2,831.9
1	Malignant neoplasms (C00-C97)	6,109	28.6	810.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,702	26.7	756.5
3	Diabetes mellitus (E10-E14)	1,133	5.3	150.3
4	Cerebrovascular diseases (I60-I69)	1,102	5.2	146.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	841	3.9	111.6
6	Chronic lower respiratory diseases (J40-J47)	807	3.8	107.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	2.9	81.3
8	Septicemia (A40-A41)	428	2.0	56.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	351	1.6	46.6
10	Influenza and pneumonia (J09-J18)	348	1.6	46.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	337	1.6	44.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	177	0.8	23.5
13	Viral hepatitis (B15-B19)	141	0.7	18.7
14	Pneumonitis due to solids and liquids (J69)	123	0.6	16.3
15	Alzheimer disease (G30)	101	0.5	13.4
...	All other causes (residual)	3,033	14.2	402.4
Non-Hispanic, black, male, 70-74 years				
...	All causes	18,050	100.0	3,717.9
1	Malignant neoplasms (C00-C97)	5,115	28.3	1,053.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,724	26.2	973.0
3	Cerebrovascular diseases (I60-I69)	1,093	6.1	225.1
4	Diabetes mellitus (E10-E14)	933	5.2	192.2
5	Chronic lower respiratory diseases (J40-J47)	872	4.8	179.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	591	3.3	121.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	415	2.3	85.5
8	Septicemia (A40-A41)	377	2.1	77.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	321	1.8	66.1
10	Influenza and pneumonia (J09-J18)	297	1.6	61.2
11	Alzheimer disease (G30)	203	1.1	41.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	166	0.9	34.2
13	Parkinson disease (G20-G21)	129	0.7	26.6
14	Pneumonitis due to solids and liquids (J69)	124	0.7	25.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.5	18.7
...	All other causes (residual)	2,599	14.4	535.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	16,251	100.0	5,298.3
1	Malignant neoplasms (C00-C97)	4,254	26.2	1,386.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,159	25.6	1,356.0
3	Cerebrovascular diseases (I60-I69)	1,029	6.3	335.5
4	Diabetes mellitus (E10-E14)	814	5.0	265.4
5	Chronic lower respiratory diseases (J40-J47)	790	4.9	257.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	575	3.5	187.5
7	Alzheimer disease (G30)	381	2.3	124.2
8	Septicemia (A40-A41)	348	2.1	113.5
9	Influenza and pneumonia (J09-J18)	320	2.0	104.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	1.9	100.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	1.7	88.7
12	Parkinson disease (G20-G21)	182	1.1	59.3
13	Pneumonitis due to solids and liquids (J69)	128	0.8	41.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	84	0.5	27.4
...	All other causes (residual)	2,521	15.5	821.9
Non-Hispanic, black, male, 80-84 years				
...	All causes	14,488	100.0	7,821.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,768	26.0	2,034.2
2	Malignant neoplasms (C00-C97)	3,193	22.0	1,723.7
3	Cerebrovascular diseases (I60-I69)	995	6.9	537.2
4	Chronic lower respiratory diseases (J40-J47)	684	4.7	369.3
5	Diabetes mellitus (E10-E14)	642	4.4	346.6
6	Alzheimer disease (G30)	585	4.0	315.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	514	3.5	277.5
8	Influenza and pneumonia (J09-J18)	321	2.2	173.3
9	Septicemia (A40-A41)	317	2.2	171.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	289	2.0	156.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	1.9	147.9
12	Parkinson disease (G20-G21)	205	1.4	110.7
13	Pneumonitis due to solids and liquids (J69)	133	0.9	71.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.6	44.3
15	Nutritional deficiencies (E40-E64)	55	0.4	29.7
...	All other causes (residual)	2,431	16.8	1,312.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,672	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,789	26.1	---
2	Malignant neoplasms (C00-C97)	2,001	18.8	---
3	Cerebrovascular diseases (I60-I69)	752	7.0	---
4	Alzheimer disease (G30)	570	5.3	---
5	Chronic lower respiratory diseases (J40-J47)	514	4.8	---
6	Diabetes mellitus (E10-E14)	432	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	3.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	2.3	---
8	Influenza and pneumonia (J09-J18)	247	2.3	---
10	Septicemia (A40-A41)	215	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.7	---
12	Parkinson disease (G20-G21)	161	1.5	---
13	Pneumonitis due to solids and liquids (J69)	135	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.6	---
15	Nutritional deficiencies (E40-E64)	52	0.5	---
...	All other causes (residual)	1,933	18.1	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	6,220	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,768	28.4	---
2	Malignant neoplasms (C00-C97)	950	15.3	---
3	Cerebrovascular diseases (I60-I69)	422	6.8	---
4	Alzheimer disease (G30)	407	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	286	4.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	3.2	---
7	Diabetes mellitus (E10-E14)	179	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	2.5	---
9	Influenza and pneumonia (J09-J18)	151	2.4	---
10	Septicemia (A40-A41)	111	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	1.5	---
12	Parkinson disease (G20-G21)	79	1.3	---
13	Pneumonitis due to solids and liquids (J69)	71	1.1	---
14	Nutritional deficiencies (E40-E64)	37	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.5	---
...	All other causes (residual)	1,277	20.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	2,181	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	668	30.6	---
2	Malignant neoplasms (C00-C97)	240	11.0	---
3	Alzheimer disease (G30)	152	7.0	---
4	Cerebrovascular diseases (I60-I69)	150	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	85	3.9	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	3.3	---
7	Diabetes mellitus (E10-E14)	60	2.8	---
8	Influenza and pneumonia (J09-J18)	59	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	2.6	---
10	Septicemia (A40-A41)	49	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.7	---
12	Pneumonitis due to solids and liquids (J69)	27	1.2	---
13	Nutritional deficiencies (E40-E64)	17	0.8	---
14	Parkinson disease (G20-G21)	16	0.7	---
15	Atherosclerosis (I70)	12	0.6	---
...	All other causes (residual)	480	22.0	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	389	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	32.6	---
2	Malignant neoplasms (C00-C97)	38	9.8	---
3	Cerebrovascular diseases (I60-I69)	26	6.7	---
4	Alzheimer disease (G30)	21	5.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	4.6	---
6	Influenza and pneumonia (J09-J18)	14	3.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.1	---
7	Pneumonitis due to solids and liquids (J69)	12	3.1	---
9	Diabetes mellitus (E10-E14)	10	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	8	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.1	---
12	Septicemia (A40-A41)	5	1.3	---
12	Nutritional deficiencies (E40-E64)	5	1.3	---
14	Parkinson disease (G20-G21)	2	0.5	---
15	Syphilis (A50-A53)	1	0.3	---
15	Anemias (D50-D64)	1	0.3	---
15	Atherosclerosis (I70)	1	0.3	---
...	All other causes (residual)	80	20.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	161,264	100.0	726.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,750	22.8	165.5
2	Malignant neoplasms (C00-C97)	34,611	21.5	155.8
3	Cerebrovascular diseases (I60-I69)	10,522	6.5	47.4
4	Diabetes mellitus (E10-E14)	7,304	4.5	32.9
5	Alzheimer disease (G30)	6,515	4.0	29.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,025	3.7	27.1
7	Chronic lower respiratory diseases (J40-J47)	5,612	3.5	25.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,987	3.1	22.5
9	Septicemia (A40-A41)	3,516	2.2	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,399	2.1	15.3
11	Influenza and pneumonia (J09-J18)	2,874	1.8	12.9
12	Certain conditions originating in the perinatal period (P00-P96)	1,557	1.0	7.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,265	0.8	5.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,226	0.8	5.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,002	0.6	4.5
...	All other causes (residual)	34,099	21.1	153.5
Non-Hispanic, black, female, 1-4 years				
...	All causes	450	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	28.9	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	12.4	4.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	11.3	4.2
4	Malignant neoplasms (C00-C97)	19	4.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Influenza and pneumonia (J09-J18)	12	2.7	*
6	Certain conditions originating in the perinatal period (P00-P96)	12	2.7	*
8	Cerebrovascular diseases (I60-I69)	10	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
10	Septicemia (A40-A41)	5	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
13	Anemias (D50-D64)	3	0.7	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (residual)	113	25.1	9.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	254	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	27.6	4.6
2	Malignant neoplasms (C00-C97)	37	14.6	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	11.4	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	7.1	*
5	Chronic lower respiratory diseases (J40-J47)	12	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.5	*
7	Cerebrovascular diseases (I60-I69)	6	2.4	*
7	Influenza and pneumonia (J09-J18)	6	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.0	*
10	Septicemia (A40-A41)	3	1.2	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
13	Viral hepatitis (B15-B19)	1	0.4	*
13	Anemias (D50-D64)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
...	All other causes (residual)	50	19.7	3.3
Non-Hispanic, black, female, 10-14 years				
...	All causes	258	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	23.3	3.9
2	Malignant neoplasms (C00-C97)	26	10.1	1.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	10.1	1.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	7.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	6.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.8	*
7	Chronic lower respiratory diseases (J40-J47)	11	4.3	*
8	Influenza and pneumonia (J09-J18)	6	2.3	*
9	Diabetes mellitus (E10-E14)	5	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.6	*
10	Anemias (D50-D64)	4	1.6	*
10	Cerebrovascular diseases (I60-I69)	4	1.6	*
13	Septicemia (A40-A41)	3	1.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
15	Meningitis (G00,G03)	1	0.4	*
15	Pneumonitis due to solids and liquids (J69)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (residual)	53	20.5	3.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	541	100.0	34.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	27.0	9.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	20.9	7.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	11.6	4.0
4	Malignant neoplasms (C00-C97)	39	7.2	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.8	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.4	*
7	Diabetes mellitus (E10-E14)	10	1.8	*
8	Anemias (D50-D64)	8	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	8	1.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
11	Influenza and pneumonia (J09-J18)	5	0.9	*
13	Septicemia (A40-A41)	4	0.7	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	88	16.3	5.6
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,112	100.0	66.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	32.0	21.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	16.3	10.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	7.4	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	6.2	4.1
5	Malignant neoplasms (C00-C97)	53	4.8	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	4.6	3.0
7	Diabetes mellitus (E10-E14)	26	2.3	1.5
8	Chronic lower respiratory diseases (J40-J47)	22	2.0	1.3
9	Anemias (D50-D64)	19	1.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.6	*
11	Cerebrovascular diseases (I60-I69)	15	1.3	*
12	Influenza and pneumonia (J09-J18)	11	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
14	Septicemia (A40-A41)	9	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
...	All other causes (residual)	178	16.0	10.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,612	100.0	90.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	24.8	22.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	11.7	10.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	8.6	7.8
4	Malignant neoplasms (C00-C97)	130	8.1	7.3
5	Pregnancy, childbirth and the puerperium (O00-O99)	88	5.5	5.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	4.5	4.1
7	Diabetes mellitus (E10-E14)	60	3.7	3.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.5	2.3
9	Cerebrovascular diseases (I60-I69)	32	2.0	1.8
10	Septicemia (A40-A41)	26	1.6	1.5
11	Anemias (D50-D64)	25	1.6	1.4
12	Influenza and pneumonia (J09-J18)	21	1.3	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.2	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.1	*
15	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
...	All other causes (residual)	335	20.8	18.9
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,920	100.0	126.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	20.8	26.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	12.8	16.2
3	Malignant neoplasms (C00-C97)	229	11.9	15.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	7.4	9.4
5	Diabetes mellitus (E10-E14)	80	4.2	5.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	73	3.8	4.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	3.5	4.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	63	3.3	4.1
9	Cerebrovascular diseases (I60-I69)	43	2.2	2.8
10	Chronic lower respiratory diseases (J40-J47)	32	1.7	2.1
11	Anemias (D50-D64)	31	1.6	2.0
12	Influenza and pneumonia (J09-J18)	29	1.5	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.5	1.9
14	Septicemia (A40-A41)	23	1.2	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.0	*
...	All other causes (residual)	413	21.5	27.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,588	100.0	175.4
1	Malignant neoplasms (C00-C97)	437	16.9	29.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	429	16.6	29.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	15.8	27.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	4.4	7.8
5	Diabetes mellitus (E10-E14)	107	4.1	7.3
6	Cerebrovascular diseases (I60-I69)	86	3.3	5.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	81	3.1	5.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	67	2.6	4.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	2.6	4.5
10	Septicemia (A40-A41)	46	1.8	3.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.7	3.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.5	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.4	2.4
13	Chronic lower respiratory diseases (J40-J47)	36	1.4	2.4
15	Anemias (D50-D64)	25	1.0	1.7
...	All other causes (residual)	563	21.8	38.2
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,417	100.0	252.8
1	Malignant neoplasms (C00-C97)	738	21.6	54.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	18.2	46.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	10.3	26.1
4	Diabetes mellitus (E10-E14)	159	4.7	11.8
5	Cerebrovascular diseases (I60-I69)	136	4.0	10.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	101	3.0	7.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	2.6	6.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	2.4	6.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.0	5.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	1.7	4.2
11	Septicemia (A40-A41)	56	1.6	4.1
12	Influenza and pneumonia (J09-J18)	52	1.5	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.3	3.3
13	Chronic lower respiratory diseases (J40-J47)	44	1.3	3.3
15	Pregnancy, childbirth and the puerperium (O00-O99)	37	1.1	2.7
...	All other causes (residual)	780	22.8	57.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,059	100.0	357.9
1	Malignant neoplasms (C00-C97)	1,212	24.0	85.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,044	20.6	73.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	9.0	32.1
4	Cerebrovascular diseases (I60-I69)	234	4.6	16.6
5	Diabetes mellitus (E10-E14)	230	4.5	16.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	143	2.8	10.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	2.4	8.5
8	Septicemia (A40-A41)	109	2.2	7.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	108	2.1	7.6
10	Chronic lower respiratory diseases (J40-J47)	89	1.8	6.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.7	6.1
12	Influenza and pneumonia (J09-J18)	74	1.5	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	1.3	4.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	1.0	3.5
15	Anemias (D50-D64)	33	0.7	2.3
...	All other causes (residual)	1,008	19.9	71.3
Non-Hispanic, black, female, 50-54 years				
...	All causes	7,986	100.0	557.6
1	Malignant neoplasms (C00-C97)	2,281	28.6	159.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,648	20.6	115.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	7.6	42.2
4	Diabetes mellitus (E10-E14)	385	4.8	26.9
5	Cerebrovascular diseases (I60-I69)	367	4.6	25.6
6	Chronic lower respiratory diseases (J40-J47)	228	2.9	15.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.5	14.2
8	Septicemia (A40-A41)	178	2.2	12.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	173	2.2	12.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	155	1.9	10.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	1.7	9.4
12	Influenza and pneumonia (J09-J18)	112	1.4	7.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.7	3.7
14	Anemias (D50-D64)	34	0.4	2.4
15	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.4	2.2
...	All other causes (residual)	1,398	17.5	97.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	12,067	100.0	849.1
1	Malignant neoplasms (C00-C97)	3,785	31.4	266.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,574	21.3	181.1
3	Cerebrovascular diseases (I60-I69)	609	5.0	42.9
4	Diabetes mellitus (E10-E14)	586	4.9	41.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	545	4.5	38.3
6	Chronic lower respiratory diseases (J40-J47)	423	3.5	29.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	2.9	24.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	254	2.1	17.9
9	Septicemia (A40-A41)	245	2.0	17.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	201	1.7	14.1
11	Influenza and pneumonia (J09-J18)	190	1.6	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	171	1.4	12.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.5	4.2
14	Anemias (D50-D64)	48	0.4	3.4
15	Viral hepatitis (B15-B19)	44	0.4	3.1
15	Pneumonitis due to solids and liquids (J69)	44	0.4	3.1
...	All other causes (residual)	1,940	16.1	136.5
Non-Hispanic, black, female, 60-64 years				
...	All causes	14,879	100.0	1,200.3
1	Malignant neoplasms (C00-C97)	4,644	31.2	374.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,361	22.6	271.1
3	Diabetes mellitus (E10-E14)	774	5.2	62.4
4	Cerebrovascular diseases (I60-I69)	746	5.0	60.2
5	Chronic lower respiratory diseases (J40-J47)	631	4.2	50.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	503	3.4	40.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	2.9	34.9
8	Septicemia (A40-A41)	378	2.5	30.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	1.9	23.0
10	Influenza and pneumonia (J09-J18)	215	1.4	17.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	201	1.4	16.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	123	0.8	9.9
13	Viral hepatitis (B15-B19)	65	0.4	5.2
14	Pneumonitis due to solids and liquids (J69)	62	0.4	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.3	4.2
...	All other causes (residual)	2,406	16.2	194.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	16,222	100.0	1,657.9
1	Malignant neoplasms (C00-C97)	5,072	31.3	518.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,641	22.4	372.1
3	Cerebrovascular diseases (I60-I69)	941	5.8	96.2
4	Diabetes mellitus (E10-E14)	891	5.5	91.1
5	Chronic lower respiratory diseases (J40-J47)	712	4.4	72.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	568	3.5	58.1
7	Septicemia (A40-A41)	427	2.6	43.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	310	1.9	31.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	1.9	31.1
10	Influenza and pneumonia (J09-J18)	257	1.6	26.3
11	Alzheimer disease (G30)	138	0.9	14.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	135	0.8	13.8
13	Pneumonitis due to solids and liquids (J69)	77	0.5	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.4	6.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.4	6.6
...	All other causes (residual)	2,617	16.1	267.5
Non-Hispanic, black, female, 70-74 years				
...	All causes	15,403	100.0	2,305.0
1	Malignant neoplasms (C00-C97)	4,314	28.0	645.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,626	23.5	542.6
3	Cerebrovascular diseases (I60-I69)	1,033	6.7	154.6
4	Diabetes mellitus (E10-E14)	837	5.4	125.3
5	Chronic lower respiratory diseases (J40-J47)	690	4.5	103.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	581	3.8	86.9
7	Septicemia (A40-A41)	411	2.7	61.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	322	2.1	48.2
9	Alzheimer disease (G30)	296	1.9	44.3
10	Influenza and pneumonia (J09-J18)	267	1.7	40.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	1.4	31.1
12	Pneumonitis due to solids and liquids (J69)	94	0.6	14.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.6	13.3
14	Parkinson disease (G20-G21)	76	0.5	11.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.4	9.9
...	All other causes (residual)	2,493	16.2	373.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	16,577	100.0	3,570.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,947	23.8	850.2
2	Malignant neoplasms (C00-C97)	3,902	23.5	840.5
3	Cerebrovascular diseases (I60-I69)	1,241	7.5	267.3
4	Diabetes mellitus (E10-E14)	898	5.4	193.4
5	Chronic lower respiratory diseases (J40-J47)	738	4.5	159.0
6	Alzheimer disease (G30)	674	4.1	145.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	615	3.7	132.5
8	Septicemia (A40-A41)	441	2.7	95.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	357	2.2	76.9
10	Influenza and pneumonia (J09-J18)	314	1.9	67.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	1.3	45.0
12	Parkinson disease (G20-G21)	120	0.7	25.8
13	Pneumonitis due to solids and liquids (J69)	112	0.7	24.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.6	20.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.3	11.4
...	All other causes (residual)	2,860	17.3	616.0
Non-Hispanic, black, female, 80-84 years				
...	All causes	17,760	100.0	5,523.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,360	24.5	1,356.0
2	Malignant neoplasms (C00-C97)	3,418	19.2	1,063.1
3	Cerebrovascular diseases (I60-I69)	1,445	8.1	449.4
4	Alzheimer disease (G30)	1,128	6.4	350.8
5	Diabetes mellitus (E10-E14)	827	4.7	257.2
6	Chronic lower respiratory diseases (J40-J47)	700	3.9	217.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	676	3.8	210.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	423	2.4	131.6
9	Septicemia (A40-A41)	379	2.1	117.9
10	Influenza and pneumonia (J09-J18)	364	2.0	113.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	1.3	70.0
12	Parkinson disease (G20-G21)	138	0.8	42.9
13	Pneumonitis due to solids and liquids (J69)	108	0.6	33.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	25.2
15	Nutritional deficiencies (E40-E64)	74	0.4	23.0
...	All other causes (residual)	3,414	19.2	1,061.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	17,264	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,571	26.5	---
2	Malignant neoplasms (C00-C97)	2,407	13.9	---
3	Cerebrovascular diseases (I60-I69)	1,548	9.0	---
4	Alzheimer disease (G30)	1,523	8.8	---
5	Diabetes mellitus (E10-E14)	700	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	599	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	568	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	464	2.7	---
9	Septicemia (A40-A41)	367	2.1	---
10	Influenza and pneumonia (J09-J18)	363	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	1.3	---
12	Pneumonitis due to solids and liquids (J69)	156	0.9	---
13	Parkinson disease (G20-G21)	122	0.7	---
14	Nutritional deficiencies (E40-E64)	86	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.4	---
...	All other causes (residual)	3,497	20.3	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	13,663	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,666	26.8	---
2	Alzheimer disease (G30)	1,534	11.2	---
3	Malignant neoplasms (C00-C97)	1,326	9.7	---
4	Cerebrovascular diseases (I60-I69)	1,241	9.1	---
5	Diabetes mellitus (E10-E14)	506	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	422	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	417	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	2.8	---
9	Influenza and pneumonia (J09-J18)	324	2.4	---
10	Septicemia (A40-A41)	236	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	1.2	---
12	Pneumonitis due to solids and liquids (J69)	115	0.8	---
13	Nutritional deficiencies (E40-E64)	94	0.7	---
14	Parkinson disease (G20-G21)	73	0.5	---
15	Atherosclerosis (I70)	62	0.5	---
...	All other causes (residual)	3,103	22.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	6,929	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,989	28.7	---
2	Alzheimer disease (G30)	848	12.2	---
3	Cerebrovascular diseases (I60-I69)	575	8.3	---
4	Malignant neoplasms (C00-C97)	446	6.4	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	218	3.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	2.7	---
7	Diabetes mellitus (E10-E14)	179	2.6	---
8	Chronic lower respiratory diseases (J40-J47)	152	2.2	---
9	Influenza and pneumonia (J09-J18)	149	2.2	---
10	Septicemia (A40-A41)	111	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	1.3	---
12	Nutritional deficiencies (E40-E64)	69	1.0	---
13	Pneumonitis due to solids and liquids (J69)	51	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.5	---
15	Atherosclerosis (I70)	28	0.4	---
...	All other causes (residual)	1,801	26.0	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,327	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	703	30.2	---
2	Alzheimer disease (G30)	303	13.0	---
3	Cerebrovascular diseases (I60-I69)	193	8.3	---
4	Malignant neoplasms (C00-C97)	92	4.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	3.5	---
6	Influenza and pneumonia (J09-J18)	59	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	44	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.8	---
9	Diabetes mellitus (E10-E14)	39	1.7	---
10	Septicemia (A40-A41)	35	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	1.3	---
12	Nutritional deficiencies (E40-E64)	23	1.0	---
13	Pneumonitis due to solids and liquids (J69)	14	0.6	---
14	Atherosclerosis (I70)	9	0.4	---
15	Anemias (D50-D64)	7	0.3	---
...	All other causes (residual)	652	28.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	19,198	100.0	703.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,472	18.1	127.2
2	Malignant neoplasms (C00-C97)	3,257	17.0	119.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,229	11.6	81.7
4	Diabetes mellitus (E10-E14)	1,114	5.8	40.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,047	5.5	38.4
6	Chronic lower respiratory diseases (J40-J47)	939	4.9	34.4
7	Cerebrovascular diseases (I60-I69)	737	3.8	27.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	602	3.1	22.1
9	Influenza and pneumonia (J09-J18)	391	2.0	14.3
10	Alzheimer disease (G30)	376	2.0	13.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	1.7	12.2
12	Septicemia (A40-A41)	295	1.5	10.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	1.4	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.0	7.0
15	Certain conditions originating in the perinatal period (P00-P96)	117	0.6	4.3
...	All other causes (residual)	3,833	20.0	140.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	72	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	34.7	15.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	11.1	*
3	Influenza and pneumonia (J09-J18)	5	6.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	6.9	*
5	Malignant neoplasms (C00-C97)	4	5.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.8	*
6	Cerebrovascular diseases (I60-I69)	2	2.8	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
8	Hernia (K40-K46)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	18	25.0	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-9 years				
...	All causes	38	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	50.0	*
2	Malignant neoplasms (C00-C97)	4	10.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	7.9	*
4	Certain conditions originating in the perinatal period (P00-P96)	2	5.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
...	All other causes (residual)	5	13.2	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 10-14 years				
...	All causes	64	100.0	31.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	32.8	10.2
2	Malignant neoplasms (C00-C97)	12	18.8	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	18.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.1	*
6	Diabetes mellitus (E10-E14)	1	1.6	*
6	Influenza and pneumonia (J09-J18)	1	1.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.6	*
6	Legal intervention (Y35,Y89.0)	1	1.6	*
...	All other causes (residual)	10	15.6	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-19 years				
...	All causes	208	100.0	99.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	43.8	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	32.7	32.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	7.7	*
4	Malignant neoplasms (C00-C97)	9	4.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	0.5	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
...	All other causes (residual)	15	7.2	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 20-24 years				
...	All causes	354	100.0	163.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	46.6	76.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	25.7	42.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.2	13.4
4	Malignant neoplasms (C00-C97)	7	2.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.7	*
6	Legal intervention (Y35,Y89.0)	5	1.4	*
7	Diabetes mellitus (E10-E14)	3	0.8	*
7	Cerebrovascular diseases (I60-I69)	3	0.8	*
9	Septicemia (A40-A41)	2	0.6	*
9	Influenza and pneumonia (J09-J18)	2	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (residual)	33	9.3	15.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-29 years				
...	All causes	547	100.0	245.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	42.6	104.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	19.9	49.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.5	18.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	23	4.2	10.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.5	*
6	Malignant neoplasms (C00-C97)	13	2.4	*
7	Septicemia (A40-A41)	5	0.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
8	Diabetes mellitus (E10-E14)	4	0.7	*
8	Influenza and pneumonia (J09-J18)	4	0.7	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
14	Cerebrovascular diseases (I60-I69)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	76	13.9	34.1
Non-Hispanic, American Indian or Alaska Native, both sexes, 30-34 years				
...	All causes	647	100.0	343.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	34.9	119.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	11.6	39.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	62	9.6	32.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	8.5	29.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.3	21.7
6	Malignant neoplasms (C00-C97)	15	2.3	*
7	Diabetes mellitus (E10-E14)	10	1.5	*
8	Cerebrovascular diseases (I60-I69)	8	1.2	*
8	Influenza and pneumonia (J09-J18)	8	1.2	*
10	Septicemia (A40-A41)	7	1.1	*
11	Legal intervention (Y35,Y89.0)	5	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
...	All other causes (residual)	115	17.8	61.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 35-39 years				
...	All causes	680	100.0	394.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	33.5	132.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	96	14.1	55.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	9.6	37.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	8.4	33.0
5	Malignant neoplasms (C00-C97)	30	4.4	17.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.3	16.8
7	Diabetes mellitus (E10-E14)	24	3.5	13.9
8	Influenza and pneumonia (J09-J18)	9	1.3	*
9	Cerebrovascular diseases (I60-I69)	7	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
11	Septicemia (A40-A41)	4	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
13	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (residual)	116	17.1	67.3
Non-Hispanic, American Indian or Alaska Native, both sexes, 40-44 years				
...	All causes	729	100.0	465.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	26.5	123.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	93	12.8	59.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	10.7	49.8
4	Malignant neoplasms (C00-C97)	52	7.1	33.2
5	Diabetes mellitus (E10-E14)	51	7.0	32.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	5.8	26.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	3.4	16.0
8	Cerebrovascular diseases (I60-I69)	13	1.8	*
9	Septicemia (A40-A41)	12	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
12	Influenza and pneumonia (J09-J18)	7	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
14	Viral hepatitis (B15-B19)	4	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (residual)	124	17.0	79.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-49 years				
...	All causes	948	100.0	584.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	20.4	118.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	146	15.4	90.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	14.7	85.6
4	Malignant neoplasms (C00-C97)	95	10.0	58.5
5	Diabetes mellitus (E10-E14)	48	5.1	29.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	4.4	25.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.2	12.9
8	Septicemia (A40-A41)	20	2.1	12.3
8	Influenza and pneumonia (J09-J18)	20	2.1	12.3
10	Cerebrovascular diseases (I60-I69)	18	1.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
13	Viral hepatitis (B15-B19)	6	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
...	All other causes (residual)	167	17.6	102.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 50-54 years				
...	All causes	1,338	100.0	784.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	16.4	129.0
2	Malignant neoplasms (C00-C97)	213	15.9	124.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	14.6	114.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	165	12.3	96.7
5	Diabetes mellitus (E10-E14)	88	6.6	51.6
6	Cerebrovascular diseases (I60-I69)	37	2.8	21.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.8	21.7
8	Chronic lower respiratory diseases (J40-J47)	34	2.5	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	1.9	14.7
10	Influenza and pneumonia (J09-J18)	23	1.7	13.5
11	Septicemia (A40-A41)	22	1.6	12.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.2	*
13	Viral hepatitis (B15-B19)	11	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.3	*
15	Anemias (D50-D64)	4	0.3	*
...	All other causes (residual)	232	17.3	136.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 55-59 years				
...	All causes	1,686	100.0	969.7
1	Malignant neoplasms (C00-C97)	374	22.2	215.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	298	17.7	171.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	10.0	97.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	156	9.3	89.7
5	Diabetes mellitus (E10-E14)	120	7.1	69.0
6	Chronic lower respiratory diseases (J40-J47)	66	3.9	38.0
7	Cerebrovascular diseases (I60-I69)	43	2.6	24.7
8	Influenza and pneumonia (J09-J18)	29	1.7	16.7
9	Septicemia (A40-A41)	27	1.6	15.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.3	12.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.2	11.5
12	Viral hepatitis (B15-B19)	17	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.9	*
14	Pneumonitis due to solids and liquids (J69)	13	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.5	*
...	All other causes (residual)	307	18.2	176.6
Non-Hispanic, American Indian or Alaska Native, both sexes, 60-64 years				
...	All causes	1,857	100.0	1,251.2
1	Malignant neoplasms (C00-C97)	501	27.0	337.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	19.4	242.6
3	Diabetes mellitus (E10-E14)	124	6.7	83.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	122	6.6	82.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	6.1	76.8
6	Chronic lower respiratory diseases (J40-J47)	112	6.0	75.5
7	Cerebrovascular diseases (I60-I69)	59	3.2	39.8
8	Influenza and pneumonia (J09-J18)	35	1.9	23.6
9	Septicemia (A40-A41)	34	1.8	22.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.8	22.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	1.0	*
12	Viral hepatitis (B15-B19)	17	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.8	*
14	Pneumonitis due to solids and liquids (J69)	10	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
...	All other causes (residual)	295	15.9	198.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 65-69 years				
...	All causes	1,889	100.0	1,649.4
1	Malignant neoplasms (C00-C97)	492	26.0	429.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	20.0	330.0
3	Diabetes mellitus (E10-E14)	154	8.2	134.5
4	Chronic lower respiratory diseases (J40-J47)	125	6.6	109.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	4.7	77.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	81	4.3	70.7
7	Cerebrovascular diseases (I60-I69)	67	3.5	58.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.4	40.2
9	Influenza and pneumonia (J09-J18)	38	2.0	33.2
10	Septicemia (A40-A41)	34	1.8	29.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.9	*
12	Alzheimer disease (G30)	9	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
13	Pneumonitis due to solids and liquids (J69)	8	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.4	*
...	All other causes (residual)	335	17.7	292.5
Non-Hispanic, American Indian or Alaska Native, both sexes, 70-74 years				
...	All causes	1,855	100.0	2,355.3
1	Malignant neoplasms (C00-C97)	466	25.1	591.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	388	20.9	492.6
3	Chronic lower respiratory diseases (J40-J47)	158	8.5	200.6
4	Diabetes mellitus (E10-E14)	138	7.4	175.2
5	Cerebrovascular diseases (I60-I69)	77	4.2	97.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	60	3.2	76.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	3.2	76.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.6	62.2
9	Septicemia (A40-A41)	36	1.9	45.7
9	Influenza and pneumonia (J09-J18)	36	1.9	45.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.4	33.0
12	Alzheimer disease (G30)	16	0.9	*
13	Parkinson disease (G20-G21)	15	0.8	*
14	Pneumonitis due to solids and liquids (J69)	9	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
...	All other causes (residual)	307	16.5	389.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 75-79 years				
...	All causes	1,798	100.0	3,598.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	394	21.9	788.6
2	Malignant neoplasms (C00-C97)	368	20.5	736.6
3	Chronic lower respiratory diseases (J40-J47)	152	8.5	304.2
4	Diabetes mellitus (E10-E14)	140	7.8	280.2
5	Cerebrovascular diseases (I60-I69)	103	5.7	206.2
6	Alzheimer disease (G30)	60	3.3	120.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	3.2	114.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.4	86.1
9	Influenza and pneumonia (J09-J18)	40	2.2	80.1
10	Septicemia (A40-A41)	34	1.9	68.1
11	Parkinson disease (G20-G21)	31	1.7	62.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.3	46.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.2	44.0
14	Pneumonitis due to solids and liquids (J69)	13	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
...	All other causes (residual)	310	17.2	620.5
Non-Hispanic, American Indian or Alaska Native, both sexes, 80-84 years				
...	All causes	1,731	100.0	5,805.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	409	23.6	1,371.8
2	Malignant neoplasms (C00-C97)	321	18.5	1,076.6
3	Cerebrovascular diseases (I60-I69)	115	6.6	385.7
4	Chronic lower respiratory diseases (J40-J47)	112	6.5	375.6
5	Diabetes mellitus (E10-E14)	106	6.1	355.5
6	Alzheimer disease (G30)	92	5.3	308.6
7	Influenza and pneumonia (J09-J18)	56	3.2	187.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	3.2	187.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.2	127.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.7	100.6
11	Parkinson disease (G20-G21)	28	1.6	93.9
12	Septicemia (A40-A41)	26	1.5	87.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.8	*
14	Pneumonitis due to solids and liquids (J69)	13	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.4	*
...	All other causes (residual)	301	17.4	1,009.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 85-89 years				
...	All causes	1,306	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	317	24.3	---
2	Malignant neoplasms (C00-C97)	178	13.6	---
3	Alzheimer disease (G30)	93	7.1	---
4	Chronic lower respiratory diseases (J40-J47)	89	6.8	---
5	Cerebrovascular diseases (I60-I69)	87	6.7	---
6	Diabetes mellitus (E10-E14)	70	5.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	3.1	---
8	Influenza and pneumonia (J09-J18)	32	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	1.9	---
11	Parkinson disease (G20-G21)	23	1.8	---
12	Septicemia (A40-A41)	16	1.2	---
13	Pneumonitis due to solids and liquids (J69)	11	0.8	---
14	Nutritional deficiencies (E40-E64)	8	0.6	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	8	0.6	---
...	All other causes (residual)	280	21.4	---
Non-Hispanic, American Indian or Alaska Native, both sexes, 90-94 years				
...	All causes	803	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	29.3	---
2	Malignant neoplasms (C00-C97)	78	9.7	---
3	Cerebrovascular diseases (I60-I69)	73	9.1	---
4	Alzheimer disease (G30)	63	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	49	6.1	---
6	Influenza and pneumonia (J09-J18)	27	3.4	---
7	Diabetes mellitus (E10-E14)	25	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.1	---
9	Parkinson disease (G20-G21)	11	1.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.1	---
12	Nutritional deficiencies (E40-E64)	8	1.0	---
13	Septicemia (A40-A41)	7	0.9	---
14	Pneumonitis due to solids and liquids (J69)	5	0.6	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	---
...	All other causes (residual)	182	22.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 95-99 years				
...	All causes	293	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	31.7	---
2	Alzheimer disease (G30)	28	9.6	---
3	Malignant neoplasms (C00-C97)	22	7.5	---
4	Cerebrovascular diseases (I60-I69)	20	6.8	---
5	Influenza and pneumonia (J09-J18)	16	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	13	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	3.8	---
8	Diabetes mellitus (E10-E14)	7	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.0	---
10	Septicemia (A40-A41)	5	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.7	---
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	1.0	---
13	Anemias (D50-D64)	2	0.7	---
13	Nutritional deficiencies (E40-E64)	2	0.7	---
13	Parkinson disease (G20-G21)	2	0.7	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.7	---
...	All other causes (residual)	56	19.1	---
Non-Hispanic, American Indian or Alaska Native, both sexes, 100 years and over				
...	All causes	71	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	29.6	---
2	Alzheimer disease (G30)	9	12.7	---
3	Chronic lower respiratory diseases (J40-J47)	4	5.6	---
4	Malignant neoplasms (C00-C97)	3	4.2	---
5	Nutritional deficiencies (E40-E64)	2	2.8	---
5	Cerebrovascular diseases (I60-I69)	2	2.8	---
5	Influenza and pneumonia (J09-J18)	2	2.8	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.8	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	2.8	---
10	Septicemia (A40-A41)	1	1.4	---
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	---
10	Diabetes mellitus (E10-E14)	1	1.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.4	---
10	Pneumonitis due to solids and liquids (J69)	1	1.4	---
10	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.4	---
...	All other causes (residual)	18	25.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,502	100.0	784.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	19.4	151.8
2	Malignant neoplasms (C00-C97)	1,725	16.4	128.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,448	13.8	108.1
4	Diabetes mellitus (E10-E14)	624	5.9	46.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	559	5.3	41.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	4.3	33.5
7	Chronic lower respiratory diseases (J40-J47)	441	4.2	32.9
8	Cerebrovascular diseases (I60-I69)	322	3.1	24.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	1.9	14.9
10	Influenza and pneumonia (J09-J18)	186	1.8	13.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.5	11.5
12	Septicemia (A40-A41)	130	1.2	9.7
13	Alzheimer disease (G30)	120	1.1	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	0.9	7.2
15	Parkinson disease (G20-G21)	68	0.6	5.1
...	All other causes (residual)	1,948	18.5	145.5
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	33	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	33.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	12.1	*
3	Malignant neoplasms (C00-C97)	3	9.1	*
4	Cerebrovascular diseases (I60-I69)	2	6.1	*
4	Influenza and pneumonia (J09-J18)	2	6.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	6.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	3.0	*
7	Hernia (K40-K46)	1	3.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	3.0	*
...	All other causes (residual)	5	15.2	*
Non-Hispanic, American Indian or Alaska Native, male, 5-9 years				
...	All causes	19	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	52.6	*
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	10.5	*
3	Malignant neoplasms (C00-C97)	1	5.3	*
3	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	5.3	*
3	Certain conditions originating in the perinatal period (P00-P96)	1	5.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	5.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	5.3	*
...	All other causes (residual)	2	10.5	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 10-14 years				
...	All causes	36	100.0	34.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	36.1	*
2	Malignant neoplasms (C00-C97)	8	22.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	13.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.8	*
5	Legal intervention (Y35,Y89.0)	1	2.8	*
...	All other causes (residual)	6	16.7	*
Non-Hispanic, American Indian or Alaska Native, male, 15-19 years				
...	All causes	148	100.0	139.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	41.2	57.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	33.8	47.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	10.1	*
4	Malignant neoplasms (C00-C97)	7	4.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.4	*
...	All other causes (residual)	10	6.8	*
Non-Hispanic, American Indian or Alaska Native, male, 20-24 years				
...	All causes	244	100.0	221.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	47.5	105.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	27.0	60.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	9.4	20.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.0	*
4	Legal intervention (Y35,Y89.0)	5	2.0	*
6	Malignant neoplasms (C00-C97)	4	1.6	*
7	Diabetes mellitus (E10-E14)	2	0.8	*
8	Septicemia (A40-A41)	1	0.4	*
8	Influenza and pneumonia (J09-J18)	1	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (residual)	19	7.8	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 25-29 years				
...	All causes	348	100.0	307.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	44.0	135.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	21.8	67.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.3	25.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	11	3.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.6	*
6	Malignant neoplasms (C00-C97)	5	1.4	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.1	*
7	Legal intervention (Y35,Y89.0)	4	1.1	*
9	Septicemia (A40-A41)	3	0.9	*
9	Diabetes mellitus (E10-E14)	3	0.9	*
11	Influenza and pneumonia (J09-J18)	2	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
...	All other causes (residual)	46	13.2	40.7
Non-Hispanic, American Indian or Alaska Native, male, 30-34 years				
...	All causes	431	100.0	459.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	36.0	165.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	12.5	57.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	9.0	41.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	35	8.1	37.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	7.2	33.1
6	Diabetes mellitus (E10-E14)	7	1.6	*
7	Malignant neoplasms (C00-C97)	6	1.4	*
8	Cerebrovascular diseases (I60-I69)	5	1.2	*
8	Legal intervention (Y35,Y89.0)	5	1.2	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J09-J18)	4	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	76	17.6	81.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 35-39 years				
...	All causes	441	100.0	518.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	35.6	184.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	12.0	62.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	48	10.9	56.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	7.7	39.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.5	23.5
6	Malignant neoplasms (C00-C97)	15	3.4	*
6	Diabetes mellitus (E10-E14)	15	3.4	*
8	Influenza and pneumonia (J09-J18)	7	1.6	*
9	Cerebrovascular diseases (I60-I69)	5	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
11	Septicemia (A40-A41)	3	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
13	Legal intervention (Y35,Y89.0)	2	0.5	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
...	All other causes (residual)	68	15.4	79.9
Non-Hispanic, American Indian or Alaska Native, male, 40-44 years				
...	All causes	430	100.0	561.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	27.2	152.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	50	11.6	65.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	10.5	58.7
4	Diabetes mellitus (E10-E14)	31	7.2	40.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	7.2	40.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.1	28.7
7	Malignant neoplasms (C00-C97)	20	4.7	26.1
8	Cerebrovascular diseases (I60-I69)	9	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.4	*
10	Septicemia (A40-A41)	5	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
10	Influenza and pneumonia (J09-J18)	5	1.2	*
13	Viral hepatitis (B15-B19)	3	0.7	*
13	Aortic aneurysm and dissection (I71)	3	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
15	Nutritional deficiencies (E40-E64)	2	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
...	All other causes (residual)	70	16.3	91.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 45-49 years				
...	All causes	592	100.0	748.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	21.6	161.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	16.0	120.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	83	14.0	104.9
4	Malignant neoplasms (C00-C97)	40	6.8	50.6
5	Diabetes mellitus (E10-E14)	32	5.4	40.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	5.1	37.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	3.0	*
8	Cerebrovascular diseases (I60-I69)	13	2.2	*
9	Influenza and pneumonia (J09-J18)	12	2.0	*
10	Septicemia (A40-A41)	10	1.7	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
14	Viral hepatitis (B15-B19)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Aortic aneurysm and dissection (I71)	3	0.5	*
...	All other causes (residual)	104	17.6	131.5
Non-Hispanic, American Indian or Alaska Native, male, 50-54 years				
...	All causes	775	100.0	945.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	19.9	187.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	15.9	150.0
3	Malignant neoplasms (C00-C97)	103	13.3	125.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	84	10.8	102.5
5	Diabetes mellitus (E10-E14)	51	6.6	62.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.1	39.0
7	Cerebrovascular diseases (I60-I69)	20	2.6	24.4
8	Septicemia (A40-A41)	13	1.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.7	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	11	1.4	*
12	Influenza and pneumonia (J09-J18)	10	1.3	*
13	Viral hepatitis (B15-B19)	9	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
15	Anemias (D50-D64)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (residual)	122	15.7	148.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 55-59 years				
...	All causes	1,000	100.0	1,219.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	20.2	246.4
2	Malignant neoplasms (C00-C97)	195	19.5	237.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	12.6	153.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	89	8.9	108.6
5	Diabetes mellitus (E10-E14)	81	8.1	98.8
6	Chronic lower respiratory diseases (J40-J47)	32	3.2	39.0
7	Cerebrovascular diseases (I60-I69)	21	2.1	25.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	1.8	*
9	Septicemia (A40-A41)	16	1.6	*
10	Influenza and pneumonia (J09-J18)	15	1.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.2	*
12	Viral hepatitis (B15-B19)	10	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
14	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (residual)	155	15.5	189.1
Non-Hispanic, American Indian or Alaska Native, male, 60-64 years				
...	All causes	1,068	100.0	1,549.5
1	Malignant neoplasms (C00-C97)	296	27.7	429.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	20.7	320.6
3	Diabetes mellitus (E10-E14)	83	7.8	120.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	6.8	105.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	64	6.0	92.9
6	Chronic lower respiratory diseases (J40-J47)	49	4.6	71.1
7	Cerebrovascular diseases (I60-I69)	28	2.6	40.6
8	Influenza and pneumonia (J09-J18)	20	1.9	29.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.4	*
10	Septicemia (A40-A41)	13	1.2	*
11	Viral hepatitis (B15-B19)	12	1.1	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.9	*
14	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
...	All other causes (residual)	162	15.2	235.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 65-69 years				
...	All causes	1,087	100.0	2,037.6
1	Malignant neoplasms (C00-C97)	279	25.7	523.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	256	23.6	479.9
3	Diabetes mellitus (E10-E14)	96	8.8	180.0
4	Chronic lower respiratory diseases (J40-J47)	62	5.7	116.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	5.3	108.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	50	4.6	93.7
7	Cerebrovascular diseases (I60-I69)	33	3.0	61.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.1	43.1
9	Influenza and pneumonia (J09-J18)	18	1.7	*
10	Septicemia (A40-A41)	14	1.3	*
11	Pneumonitis due to solids and liquids (J69)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.6	*
15	Alzheimer disease (G30)	5	0.5	*
...	All other causes (residual)	168	15.5	314.9
Non-Hispanic, American Indian or Alaska Native, male, 70-74 years				
...	All causes	997	100.0	2,739.0
1	Malignant neoplasms (C00-C97)	249	25.0	684.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	239	24.0	656.6
3	Chronic lower respiratory diseases (J40-J47)	77	7.7	211.5
4	Diabetes mellitus (E10-E14)	66	6.6	181.3
5	Cerebrovascular diseases (I60-I69)	43	4.3	118.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	35	3.5	96.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	3.2	87.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.7	74.2
9	Influenza and pneumonia (J09-J18)	19	1.9	*
10	Septicemia (A40-A41)	15	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.3	*
12	Parkinson disease (G20-G21)	9	0.9	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.7	*
14	Pneumonitis due to solids and liquids (J69)	6	0.6	*
15	Alzheimer disease (G30)	5	0.5	*
...	All other causes (residual)	155	15.5	425.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 75-79 years				
...	All causes	933	100.0	4,166.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	24.8	1,031.6
2	Malignant neoplasms (C00-C97)	198	21.2	884.2
3	Chronic lower respiratory diseases (J40-J47)	84	9.0	375.1
4	Diabetes mellitus (E10-E14)	71	7.6	317.1
5	Cerebrovascular diseases (I60-I69)	41	4.4	183.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	3.9	160.8
7	Alzheimer disease (G30)	26	2.8	116.1
8	Influenza and pneumonia (J09-J18)	20	2.1	89.3
9	Parkinson disease (G20-G21)	18	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.7	*
11	Septicemia (A40-A41)	15	1.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.8	*
14	Pneumonitis due to solids and liquids (J69)	6	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
...	All other causes (residual)	145	15.5	647.6
Non-Hispanic, American Indian or Alaska Native, male, 80-84 years				
...	All causes	829	100.0	6,639.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212	25.6	1,697.9
2	Malignant neoplasms (C00-C97)	175	21.1	1,401.6
3	Chronic lower respiratory diseases (J40-J47)	51	6.2	408.5
4	Diabetes mellitus (E10-E14)	50	6.0	400.4
5	Cerebrovascular diseases (I60-I69)	42	5.1	336.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	3.9	256.3
7	Alzheimer disease (G30)	30	3.6	240.3
8	Influenza and pneumonia (J09-J18)	22	2.7	176.2
9	Parkinson disease (G20-G21)	19	2.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.6	*
12	Septicemia (A40-A41)	12	1.4	*
13	Pneumonitis due to solids and liquids (J69)	8	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
14	Aortic aneurysm and dissection (I71)	3	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	*
...	All other causes (residual)	132	15.9	1,057.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 85-89 years				
...	All causes	565	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	167	29.6	---
2	Malignant neoplasms (C00-C97)	82	14.5	---
3	Chronic lower respiratory diseases (J40-J47)	37	6.5	---
4	Cerebrovascular diseases (I60-I69)	34	6.0	---
5	Alzheimer disease (G30)	27	4.8	---
6	Diabetes mellitus (E10-E14)	23	4.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	3.2	---
8	Parkinson disease (G20-G21)	15	2.7	---
9	Influenza and pneumonia (J09-J18)	13	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	2.1	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.9	---
12	Pneumonitis due to solids and liquids (J69)	5	0.9	---
13	Nutritional deficiencies (E40-E64)	4	0.7	---
14	Aortic aneurysm and dissection (I71)	3	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.5	---
...	All other causes (residual)	111	19.6	---
Non-Hispanic, American Indian or Alaska Native, male, 90-94 years				
...	All causes	286	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	31.5	---
2	Malignant neoplasms (C00-C97)	31	10.8	---
3	Cerebrovascular diseases (I60-I69)	24	8.4	---
4	Alzheimer disease (G30)	19	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	19	6.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	3.8	---
7	Diabetes mellitus (E10-E14)	10	3.5	---
7	Influenza and pneumonia (J09-J18)	10	3.5	---
9	Nutritional deficiencies (E40-E64)	4	1.4	---
9	Parkinson disease (G20-G21)	4	1.4	---
9	Pneumonitis due to solids and liquids (J69)	4	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.0	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	---
14	Septicemia (A40-A41)	2	0.7	---
14	Pneumoconioses and chemical effects (J60-J66,J68)	2	0.7	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.7	---
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.7	---
...	All other causes (residual)	46	16.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 95-99 years				
...	All causes	76	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	36.8	---
2	Malignant neoplasms (C00-C97)	7	9.2	---
2	Chronic lower respiratory diseases (J40-J47)	7	9.2	---
4	Influenza and pneumonia (J09-J18)	5	6.6	---
5	Alzheimer disease (G30)	4	5.3	---
6	Diabetes mellitus (E10-E14)	3	3.9	---
7	Cerebrovascular diseases (I60-I69)	2	2.6	---
7	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	2.6	---
9	Septicemia (A40-A41)	1	1.3	---
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.3	---
9	Anemias (D50-D64)	1	1.3	---
9	Pneumoconioses and chemical effects (J60-J66,J68)	1	1.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.3	---
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.3	---
...	All other causes (residual)	12	15.8	---
Non-Hispanic, American Indian or Alaska Native, male, 100 years and over				
...	All causes	13	100.0	---
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	15.4	---
2	Malignant neoplasms (C00-C97)	1	7.7	---
2	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	7.7	---
2	Alzheimer disease (G30)	1	7.7	---
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	7.7	---
2	Pneumonitis due to solids and liquids (J69)	1	7.7	---
2	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	7.7	---
...	All other causes (residual)	5	38.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,696	100.0	625.5
1	Malignant neoplasms (C00-C97)	1,532	17.6	110.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,439	16.5	103.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	781	9.0	56.2
4	Chronic lower respiratory diseases (J40-J47)	498	5.7	35.8
5	Diabetes mellitus (E10-E14)	490	5.6	35.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	488	5.6	35.1
7	Cerebrovascular diseases (I60-I69)	415	4.8	29.9
8	Alzheimer disease (G30)	256	2.9	18.4
9	Influenza and pneumonia (J09-J18)	205	2.4	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	2.1	12.9
11	Septicemia (A40-A41)	165	1.9	11.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	1.8	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	1.1	6.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	0.8	4.8
15	Certain conditions originating in the perinatal period (P00-P96)	56	0.6	4.0
...	All other causes (residual)	1,878	21.6	135.1
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	39	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	35.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.4	*
3	Influenza and pneumonia (J09-J18)	3	7.7	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.6	*
...	All other causes (residual)	13	33.3	*
Non-Hispanic, American Indian or Alaska Native, female, 5-9 years				
...	All causes	19	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	47.4	*
2	Malignant neoplasms (C00-C97)	3	15.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	5.3	*
3	Certain conditions originating in the perinatal period (P00-P96)	1	5.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	5.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	5.3	*
...	All other causes (residual)	3	15.8	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 10-14 years				
...	All causes	28	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	28.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	25.0	*
3	Malignant neoplasms (C00-C97)	4	14.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	7.1	*
5	Diabetes mellitus (E10-E14)	1	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.6	*
5	Influenza and pneumonia (J09-J18)	1	3.6	*
...	All other causes (residual)	4	14.3	*
Non-Hispanic, American Indian or Alaska Native, female, 15-19 years				
...	All causes	60	100.0	58.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	50.0	29.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	30.0	*
3	Malignant neoplasms (C00-C97)	2	3.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.7	*
4	Cerebrovascular diseases (I60-I69)	1	1.7	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.7	*
...	All other causes (residual)	5	8.3	*
Non-Hispanic, American Indian or Alaska Native, female, 20-24 years				
...	All causes	110	100.0	103.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	44.5	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	22.7	23.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	5.5	*
4	Malignant neoplasms (C00-C97)	3	2.7	*
4	Cerebrovascular diseases (I60-I69)	3	2.7	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.8	*
7	Septicemia (A40-A41)	1	0.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
7	Diabetes mellitus (E10-E14)	1	0.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.9	*
7	Influenza and pneumonia (J09-J18)	1	0.9	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.9	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.9	*
...	All other causes (residual)	14	12.7	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 25-29 years				
...	All causes	199	100.0	181.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	40.2	73.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	16.6	30.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	12	6.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	6.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	5.0	*
6	Malignant neoplasms (C00-C97)	8	4.0	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.0	*
8	Septicemia (A40-A41)	2	1.0	*
8	Cerebrovascular diseases (I60-I69)	2	1.0	*
8	Influenza and pneumonia (J09-J18)	2	1.0	*
11	Diabetes mellitus (E10-E14)	1	0.5	*
11	Nutritional deficiencies (E40-E64)	1	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (residual)	30	15.1	27.4
Non-Hispanic, American Indian or Alaska Native, female, 30-34 years				
...	All causes	216	100.0	227.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	32.9	74.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	27	12.5	28.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	9.7	22.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	7.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.6	*
6	Malignant neoplasms (C00-C97)	9	4.2	*
7	Influenza and pneumonia (J09-J18)	4	1.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.9	*
9	Septicemia (A40-A41)	3	1.4	*
9	Diabetes mellitus (E10-E14)	3	1.4	*
9	Cerebrovascular diseases (I60-I69)	3	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
13	Nutritional deficiencies (E40-E64)	1	0.5	*
13	Peptic ulcer (K25-K28)	1	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	*
...	All other causes (residual)	39	18.1	41.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 35-39 years				
...	All causes	239	100.0	273.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	29.7	81.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	48	20.1	55.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	9.6	26.3
4	Malignant neoplasms (C00-C97)	15	6.3	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	5.0	*
6	Diabetes mellitus (E10-E14)	9	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	3.8	*
8	Cerebrovascular diseases (I60-I69)	2	0.8	*
8	Influenza and pneumonia (J09-J18)	2	0.8	*
10	Tuberculosis (A16-A19)	1	0.4	*
10	Septicemia (A40-A41)	1	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
10	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (residual)	40	16.7	45.8
Non-Hispanic, American Indian or Alaska Native, female, 40-44 years				
...	All causes	299	100.0	373.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	25.4	95.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	43	14.4	53.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	11.0	41.3
4	Malignant neoplasms (C00-C97)	32	10.7	40.0
5	Diabetes mellitus (E10-E14)	20	6.7	25.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	3.7	*
7	Septicemia (A40-A41)	7	2.3	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.3	*
9	Cerebrovascular diseases (I60-I69)	4	1.3	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.0	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.0	*
12	Influenza and pneumonia (J09-J18)	2	0.7	*
12	Pneumonitis due to solids and liquids (J69)	2	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
...	All other causes (residual)	52	17.4	65.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 45-49 years				
...	All causes	356	100.0	427.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	18.3	78.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	63	17.7	75.7
3	Malignant neoplasms (C00-C97)	55	15.4	66.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	12.4	52.9
5	Diabetes mellitus (E10-E14)	16	4.5	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	3.4	*
7	Septicemia (A40-A41)	10	2.8	*
8	Influenza and pneumonia (J09-J18)	8	2.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.7	*
10	Cerebrovascular diseases (I60-I69)	5	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
12	Viral hepatitis (B15-B19)	3	0.8	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.8	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
15	Anemias (D50-D64)	1	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	55	15.4	66.1
Non-Hispanic, American Indian or Alaska Native, female, 50-54 years				
...	All causes	563	100.0	635.5
1	Malignant neoplasms (C00-C97)	110	19.5	124.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	81	14.4	91.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	13.0	82.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	11.7	74.5
5	Diabetes mellitus (E10-E14)	37	6.6	41.8
6	Chronic lower respiratory diseases (J40-J47)	23	4.1	26.0
7	Cerebrovascular diseases (I60-I69)	17	3.0	*
8	Influenza and pneumonia (J09-J18)	13	2.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.1	*
10	Septicemia (A40-A41)	9	1.6	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.9	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.7	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
14	Viral hepatitis (B15-B19)	2	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	100	17.8	112.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 55-59 years				
...	All causes	686	100.0	746.4
1	Malignant neoplasms (C00-C97)	179	26.1	194.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	14.0	104.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	67	9.8	72.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	6.3	46.8
5	Diabetes mellitus (E10-E14)	39	5.7	42.4
6	Chronic lower respiratory diseases (J40-J47)	34	5.0	37.0
7	Cerebrovascular diseases (I60-I69)	22	3.2	23.9
8	Influenza and pneumonia (J09-J18)	14	2.0	*
9	Septicemia (A40-A41)	11	1.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.2	*
11	Pneumonitis due to solids and liquids (J69)	8	1.2	*
13	Viral hepatitis (B15-B19)	7	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.6	*
15	Nutritional deficiencies (E40-E64)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	132	19.2	143.6
Non-Hispanic, American Indian or Alaska Native, female, 60-64 years				
...	All causes	789	100.0	992.5
1	Malignant neoplasms (C00-C97)	205	26.0	257.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	17.6	174.8
3	Chronic lower respiratory diseases (J40-J47)	63	8.0	79.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	58	7.4	73.0
5	Diabetes mellitus (E10-E14)	41	5.2	51.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	5.2	51.6
7	Cerebrovascular diseases (I60-I69)	31	3.9	39.0
8	Septicemia (A40-A41)	21	2.7	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.4	*
10	Influenza and pneumonia (J09-J18)	15	1.9	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.8	*
13	Viral hepatitis (B15-B19)	5	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.6	*
13	Pneumonitis due to solids and liquids (J69)	5	0.6	*
...	All other causes (residual)	129	16.3	162.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 65-69 years				
...	All causes	802	100.0	1,310.8
1	Malignant neoplasms (C00-C97)	213	26.6	348.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	15.2	199.4
3	Chronic lower respiratory diseases (J40-J47)	63	7.9	103.0
4	Diabetes mellitus (E10-E14)	58	7.2	94.8
5	Cerebrovascular diseases (I60-I69)	34	4.2	55.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.9	50.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	3.9	50.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.9	37.6
9	Septicemia (A40-A41)	20	2.5	32.7
9	Influenza and pneumonia (J09-J18)	20	2.5	32.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.4	*
12	Viral hepatitis (B15-B19)	4	0.5	*
12	Nutritional deficiencies (E40-E64)	4	0.5	*
12	Alzheimer disease (G30)	4	0.5	*
15	Atherosclerosis (I70)	3	0.4	*
...	All other causes (residual)	161	20.1	263.1
Non-Hispanic, American Indian or Alaska Native, female, 70-74 years				
...	All causes	858	100.0	2,025.5
1	Malignant neoplasms (C00-C97)	217	25.3	512.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	17.4	351.8
3	Chronic lower respiratory diseases (J40-J47)	81	9.4	191.2
4	Diabetes mellitus (E10-E14)	72	8.4	170.0
5	Cerebrovascular diseases (I60-I69)	34	4.0	80.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.3	66.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	25	2.9	59.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.6	51.9
9	Septicemia (A40-A41)	21	2.4	49.6
10	Influenza and pneumonia (J09-J18)	17	2.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.5	*
12	Alzheimer disease (G30)	11	1.3	*
13	Parkinson disease (G20-G21)	6	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.3	*
14	Aortic aneurysm and dissection (I71)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Peptic ulcer (K25-K28)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.3	*
...	All other causes (residual)	144	16.8	340.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 75-79 years				
...	All causes	865	100.0	3,137.7
1	Malignant neoplasms (C00-C97)	170	19.7	616.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	18.8	591.3
3	Diabetes mellitus (E10-E14)	69	8.0	250.3
4	Chronic lower respiratory diseases (J40-J47)	68	7.9	246.7
5	Cerebrovascular diseases (I60-I69)	62	7.2	224.9
6	Alzheimer disease (G30)	34	3.9	123.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	3.1	97.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.4	76.2
9	Influenza and pneumonia (J09-J18)	20	2.3	72.5
10	Septicemia (A40-A41)	19	2.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.7	*
12	Parkinson disease (G20-G21)	13	1.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
14	Pneumonitis due to solids and liquids (J69)	7	0.8	*
15	Anemias (D50-D64)	6	0.7	*
...	All other causes (residual)	162	18.7	587.6
Non-Hispanic, American Indian or Alaska Native, female, 80-84 years				
...	All causes	902	100.0	5,205.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	21.8	1,136.8
2	Malignant neoplasms (C00-C97)	146	16.2	842.5
3	Cerebrovascular diseases (I60-I69)	73	8.1	421.3
4	Alzheimer disease (G30)	62	6.9	357.8
5	Chronic lower respiratory diseases (J40-J47)	61	6.8	352.0
6	Diabetes mellitus (E10-E14)	56	6.2	323.2
7	Influenza and pneumonia (J09-J18)	34	3.8	196.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	2.7	138.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.4	127.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.9	*
11	Septicemia (A40-A41)	14	1.6	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.3	*
13	Parkinson disease (G20-G21)	9	1.0	*
14	Pneumonitis due to solids and liquids (J69)	5	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.4	*
...	All other causes (residual)	162	18.0	934.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 85-89 years				
...	All causes	741	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	20.2	---
2	Malignant neoplasms (C00-C97)	96	13.0	---
3	Alzheimer disease (G30)	66	8.9	---
4	Cerebrovascular diseases (I60-I69)	53	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	52	7.0	---
6	Diabetes mellitus (E10-E14)	47	6.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	3.0	---
8	Influenza and pneumonia (J09-J18)	19	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.4	---
10	Septicemia (A40-A41)	14	1.9	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.8	---
12	Parkinson disease (G20-G21)	8	1.1	---
13	Pneumonitis due to solids and liquids (J69)	6	0.8	---
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.8	---
15	Anemias (D50-D64)	4	0.5	---
15	Nutritional deficiencies (E40-E64)	4	0.5	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	---
...	All other causes (residual)	159	21.5	---
Non-Hispanic, American Indian or Alaska Native, female, 90-94 years				
...	All causes	517	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	28.0	---
2	Cerebrovascular diseases (I60-I69)	49	9.5	---
3	Malignant neoplasms (C00-C97)	47	9.1	---
4	Alzheimer disease (G30)	44	8.5	---
5	Chronic lower respiratory diseases (J40-J47)	30	5.8	---
6	Influenza and pneumonia (J09-J18)	17	3.3	---
7	Diabetes mellitus (E10-E14)	15	2.9	---
8	Parkinson disease (G20-G21)	7	1.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.2	---
12	Septicemia (A40-A41)	5	1.0	---
13	Nutritional deficiencies (E40-E64)	4	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.4	---
...	All other causes (residual)	127	24.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 95-99 years				
...	All causes	217	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	30.0	---
2	Alzheimer disease (G30)	24	11.1	---
3	Cerebrovascular diseases (I60-I69)	18	8.3	---
4	Malignant neoplasms (C00-C97)	15	6.9	---
5	Influenza and pneumonia (J09-J18)	11	5.1	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	4.6	---
7	Chronic lower respiratory diseases (J40-J47)	6	2.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	2.3	---
10	Septicemia (A40-A41)	4	1.8	---
10	Diabetes mellitus (E10-E14)	4	1.8	---
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	1.4	---
13	Nutritional deficiencies (E40-E64)	2	0.9	---
13	Parkinson disease (G20-G21)	2	0.9	---
15	Anemias (D50-D64)	1	0.5	---
15	Pneumonitis due to solids and liquids (J69)	1	0.5	---
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	---
...	All other causes (residual)	39	18.0	---
Non-Hispanic, American Indian or Alaska Native, female, 100 years and over				
...	All causes	58	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	34.5	---
2	Alzheimer disease (G30)	8	13.8	---
3	Chronic lower respiratory diseases (J40-J47)	4	6.9	---
4	Malignant neoplasms (C00-C97)	2	3.4	---
4	Nutritional deficiencies (E40-E64)	2	3.4	---
4	Cerebrovascular diseases (I60-I69)	2	3.4	---
4	Influenza and pneumonia (J09-J18)	2	3.4	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.4	---
9	Septicemia (A40-A41)	1	1.7	---
9	Diabetes mellitus (E10-E14)	1	1.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.7	---
...	All other causes (residual)	13	22.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	72,598	100.0	359.8
1	Malignant neoplasms (C00-C97)	18,257	25.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,490	21.3	76.8
3	Cerebrovascular diseases (I60-I69)	5,430	7.5	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,256	4.5	16.1
5	Diabetes mellitus (E10-E14)	3,047	4.2	15.1
6	Alzheimer disease (G30)	2,583	3.6	12.8
7	Influenza and pneumonia (J09-J18)	2,286	3.1	11.3
8	Chronic lower respiratory diseases (J40-J47)	2,063	2.8	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,534	2.1	7.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,472	2.0	7.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,414	1.9	7.0
12	Septicemia (A40-A41)	879	1.2	4.4
13	Parkinson disease (G20-G21)	869	1.2	4.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	725	1.0	3.6
15	Pneumonitis due to solids and liquids (J69)	549	0.8	2.7
...	All other causes (residual)	12,744	17.6	63.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	148	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	24.3	3.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.2	2.2
3	Malignant neoplasms (C00-C97)	20	13.5	2.1
4	Influenza and pneumonia (J09-J18)	6	4.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.7	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.7	*
7	Septicemia (A40-A41)	3	2.0	*
7	Anemias (D50-D64)	3	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.0	*
10	Meningitis (G00,G03)	2	1.4	*
10	Cerebrovascular diseases (I60-I69)	2	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
12	Hernia (K40-K46)	1	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	40	27.0	4.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	100	100.0	8.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	20.0	1.7
2	Malignant neoplasms (C00-C97)	18	18.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	12.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.0	*
5	Septicemia (A40-A41)	3	3.0	*
6	Anemias (D50-D64)	2	2.0	*
6	Influenza and pneumonia (J09-J18)	2	2.0	*
6	Acute bronchitis and bronchiolitis (J20-J21)	2	2.0	*
9	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (residual)	31	31.0	2.6
Non-Hispanic, Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	108	100.0	8.9
1	Malignant neoplasms (C00-C97)	24	22.2	2.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	20.4	1.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	15.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	8.3	*
5	Septicemia (A40-A41)	3	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.8	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.9	*
10	Meningococcal infection (A39)	1	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Anemias (D50-D64)	1	0.9	*
10	Diabetes mellitus (E10-E14)	1	0.9	*
10	Meningitis (G00,G03)	1	0.9	*
10	Influenza and pneumonia (J09-J18)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (residual)	17	15.7	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	325	100.0	26.5
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	32.3	8.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	24.9	6.6
3	Malignant neoplasms (C00-C97)	34	10.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.9	*
6	Influenza and pneumonia (J09-J18)	5	1.5	*
7	Cerebrovascular diseases (I60-I69)	4	1.2	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
9	Septicemia (A40-A41)	2	0.6	*
9	Anemias (D50-D64)	2	0.6	*
9	Aortic aneurysm and dissection (I71)	2	0.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	42	12.9	3.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	635	100.0	44.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	33.1	14.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	31.8	14.1
3	Malignant neoplasms (C00-C97)	45	7.1	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.3	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.3	1.5
6	Influenza and pneumonia (J09-J18)	10	1.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.4	*
8	Diabetes mellitus (E10-E14)	3	0.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
10	Septicemia (A40-A41)	2	0.3	*
10	Cerebrovascular diseases (I60-I69)	2	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
10	Pneumonitis due to solids and liquids (J69)	2	0.3	*
10	Legal intervention (Y35,Y89.0)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	73	11.5	5.1
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	680	100.0	39.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	33.8	13.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	23.1	9.0
3	Malignant neoplasms (C00-C97)	71	10.4	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	5.7	2.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	4.4	1.7
6	Diabetes mellitus (E10-E14)	10	1.5	*
7	Cerebrovascular diseases (I60-I69)	7	1.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
8	Legal intervention (Y35,Y89.0)	5	0.7	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.7	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
13	Septicemia (A40-A41)	3	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
13	Aortic aneurysm and dissection (I71)	3	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
...	All other causes (residual)	98	14.4	5.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	820	100.0	47.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	24.8	11.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	17.3	8.2
3	Malignant neoplasms (C00-C97)	135	16.5	7.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	7.2	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	6.2	2.9
6	Cerebrovascular diseases (I60-I69)	27	3.3	1.6
7	Influenza and pneumonia (J09-J18)	14	1.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	14	1.7	*
10	Diabetes mellitus (E10-E14)	9	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.9	*
12	Septicemia (A40-A41)	6	0.7	*
12	Aortic aneurysm and dissection (I71)	6	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
...	All other causes (residual)	119	14.5	6.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	944	100.0	57.8
1	Malignant neoplasms (C00-C97)	228	24.2	14.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	17.3	10.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	12.7	7.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	113	12.0	6.9
5	Cerebrovascular diseases (I60-I69)	38	4.0	2.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	34	3.6	2.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	2.9	1.7
8	Diabetes mellitus (E10-E14)	25	2.6	1.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.2	*
10	Septicemia (A40-A41)	10	1.1	*
10	Influenza and pneumonia (J09-J18)	10	1.1	*
12	Aortic aneurysm and dissection (I71)	8	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
...	All other causes (residual)	137	14.5	8.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,283	100.0	83.2
1	Malignant neoplasms (C00-C97)	377	29.4	24.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	15.8	13.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	11.5	9.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	7.7	6.4
5	Cerebrovascular diseases (I60-I69)	90	7.0	5.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.7	2.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	2.7	2.2
8	Diabetes mellitus (E10-E14)	32	2.5	2.1
9	Influenza and pneumonia (J09-J18)	15	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
11	Septicemia (A40-A41)	11	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
14	Viral hepatitis (B15-B19)	9	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.7	*
...	All other causes (residual)	189	14.7	12.3
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	2,007	100.0	137.6
1	Malignant neoplasms (C00-C97)	732	36.5	50.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	15.0	20.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	9.7	13.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	5.8	8.0
5	Cerebrovascular diseases (I60-I69)	112	5.6	7.7
6	Diabetes mellitus (E10-E14)	72	3.6	4.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	63	3.1	4.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.3	1.8
9	Influenza and pneumonia (J09-J18)	25	1.2	1.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	1.1	1.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
12	Aortic aneurysm and dissection (I71)	17	0.8	*
13	Septicemia (A40-A41)	14	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	14	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
...	All other causes (residual)	254	12.7	17.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,678	100.0	210.9
1	Malignant neoplasms (C00-C97)	1,053	39.3	82.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	498	18.6	39.2
3	Cerebrovascular diseases (I60-I69)	163	6.1	12.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	5.9	12.4
5	Diabetes mellitus (E10-E14)	110	4.1	8.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	3.4	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.5	5.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.2	2.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.2	2.4
10	Septicemia (A40-A41)	26	1.0	2.0
11	Influenza and pneumonia (J09-J18)	25	0.9	2.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.9	2.0
13	Chronic lower respiratory diseases (J40-J47)	24	0.9	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.6	*
15	Viral hepatitis (B15-B19)	16	0.6	*
...	All other causes (residual)	341	12.7	26.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,611	100.0	313.1
1	Malignant neoplasms (C00-C97)	1,485	41.1	128.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	694	19.2	60.2
3	Cerebrovascular diseases (I60-I69)	217	6.0	18.8
4	Diabetes mellitus (E10-E14)	184	5.1	16.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	4.8	15.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	2.5	7.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	72	2.0	6.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	4.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.4	4.5
10	Septicemia (A40-A41)	48	1.3	4.2
11	Chronic lower respiratory diseases (J40-J47)	42	1.2	3.6
12	Influenza and pneumonia (J09-J18)	36	1.0	3.1
13	Aortic aneurysm and dissection (I71)	27	0.7	2.3
14	Viral hepatitis (B15-B19)	24	0.7	2.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	0.6	1.9
...	All other causes (residual)	388	10.7	33.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,891	100.0	475.1
1	Malignant neoplasms (C00-C97)	1,989	40.7	193.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	976	20.0	94.8
3	Cerebrovascular diseases (I60-I69)	284	5.8	27.6
4	Diabetes mellitus (E10-E14)	258	5.3	25.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	3.6	17.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	2.2	10.7
7	Influenza and pneumonia (J09-J18)	94	1.9	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	82	1.7	8.0
9	Chronic lower respiratory diseases (J40-J47)	73	1.5	7.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.4	6.8
11	Septicemia (A40-A41)	67	1.4	6.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	1.4	6.5
13	Viral hepatitis (B15-B19)	41	0.8	4.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	2.4
15	Aortic aneurysm and dissection (I71)	24	0.5	2.3
...	All other causes (residual)	554	11.3	53.8
Non-Hispanic, Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	6,260	100.0	736.9
1	Malignant neoplasms (C00-C97)	2,399	38.3	282.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,309	20.9	154.1
3	Cerebrovascular diseases (I60-I69)	371	5.9	43.7
4	Diabetes mellitus (E10-E14)	303	4.8	35.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	3.2	23.7
6	Influenza and pneumonia (J09-J18)	127	2.0	14.9
7	Chronic lower respiratory diseases (J40-J47)	126	2.0	14.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.0	14.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.5	11.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.4	10.1
11	Septicemia (A40-A41)	81	1.3	9.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	1.1	7.8
13	Aortic aneurysm and dissection (I71)	48	0.8	5.7
14	Parkinson disease (G20-G21)	45	0.7	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	4.4
...	All other causes (residual)	841	13.4	99.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	6,633	100.0	1,122.0
1	Malignant neoplasms (C00-C97)	2,209	33.3	373.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,401	21.1	237.0
3	Cerebrovascular diseases (I60-I69)	503	7.6	85.1
4	Diabetes mellitus (E10-E14)	371	5.6	62.8
5	Chronic lower respiratory diseases (J40-J47)	215	3.2	36.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	2.7	30.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.5	28.1
8	Influenza and pneumonia (J09-J18)	163	2.5	27.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.7	18.6
10	Septicemia (A40-A41)	97	1.5	16.4
11	Parkinson disease (G20-G21)	95	1.4	16.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.0	11.3
13	Alzheimer disease (G30)	61	0.9	10.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.8	8.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	0.8	8.5
...	All other causes (residual)	893	13.5	151.1
Non-Hispanic, Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	8,035	100.0	1,962.4
1	Malignant neoplasms (C00-C97)	2,348	29.2	573.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,727	21.5	421.8
3	Cerebrovascular diseases (I60-I69)	609	7.6	148.7
4	Diabetes mellitus (E10-E14)	430	5.4	105.0
5	Chronic lower respiratory diseases (J40-J47)	285	3.5	69.6
6	Influenza and pneumonia (J09-J18)	247	3.1	60.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	2.8	55.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.5	49.6
9	Alzheimer disease (G30)	181	2.3	44.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.7	33.5
11	Parkinson disease (G20-G21)	135	1.7	33.0
12	Septicemia (A40-A41)	112	1.4	27.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	70	0.9	17.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.7	14.4
15	Pneumonitis due to solids and liquids (J69)	56	0.7	13.7
...	All other causes (residual)	1,209	15.0	295.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	9,495	100.0	3,529.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,163	22.8	804.1
2	Malignant neoplasms (C00-C97)	2,106	22.2	782.9
3	Cerebrovascular diseases (I60-I69)	841	8.9	312.6
4	Diabetes mellitus (E10-E14)	456	4.8	169.5
5	Chronic lower respiratory diseases (J40-J47)	403	4.2	149.8
6	Alzheimer disease (G30)	377	4.0	140.2
7	Influenza and pneumonia (J09-J18)	332	3.5	123.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	2.9	102.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	261	2.7	97.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	236	2.5	87.7
11	Parkinson disease (G20-G21)	213	2.2	79.2
12	Septicemia (A40-A41)	132	1.4	49.1
13	Pneumonitis due to solids and liquids (J69)	97	1.0	36.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.7	25.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	45	0.5	16.7
...	All other causes (residual)	1,489	15.7	553.5
Non-Hispanic, Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	9,998	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,348	23.5	---
2	Malignant neoplasms (C00-C97)	1,690	16.9	---
3	Cerebrovascular diseases (I60-I69)	965	9.7	---
4	Alzheimer disease (G30)	655	6.6	---
5	Influenza and pneumonia (J09-J18)	437	4.4	---
6	Diabetes mellitus (E10-E14)	406	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	383	3.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	2.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	2.4	---
11	Parkinson disease (G20-G21)	222	2.2	---
12	Pneumonitis due to solids and liquids (J69)	124	1.2	---
13	Septicemia (A40-A41)	116	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	51	0.5	---
...	All other causes (residual)	1,761	17.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	8,272	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,215	26.8	---
2	Malignant neoplasms (C00-C97)	939	11.4	---
3	Cerebrovascular diseases (I60-I69)	803	9.7	---
4	Alzheimer disease (G30)	714	8.6	---
5	Influenza and pneumonia (J09-J18)	433	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	325	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	267	3.2	---
8	Diabetes mellitus (E10-E14)	261	3.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.0	---
11	Pneumonitis due to solids and liquids (J69)	115	1.4	---
12	Parkinson disease (G20-G21)	106	1.3	---
13	Septicemia (A40-A41)	82	1.0	---
14	Nutritional deficiencies (E40-E64)	50	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.4	---
...	All other causes (residual)	1,559	18.8	---
Non-Hispanic, Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	3,787	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,102	29.1	---
2	Alzheimer disease (G30)	429	11.3	---
3	Cerebrovascular diseases (I60-I69)	328	8.7	---
4	Malignant neoplasms (C00-C97)	303	8.0	---
5	Influenza and pneumonia (J09-J18)	242	6.4	---
6	Chronic lower respiratory diseases (J40-J47)	126	3.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	3.1	---
8	Diabetes mellitus (E10-E14)	93	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.0	---
11	Septicemia (A40-A41)	48	1.3	---
12	Pneumonitis due to solids and liquids (J69)	44	1.2	---
13	Parkinson disease (G20-G21)	24	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.5	---
15	Nutritional deficiencies (E40-E64)	18	0.5	---
...	All other causes (residual)	742	19.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	931	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263	28.2	---
2	Alzheimer disease (G30)	118	12.7	---
3	Cerebrovascular diseases (I60-I69)	58	6.2	---
4	Influenza and pneumonia (J09-J18)	52	5.6	---
5	Malignant neoplasms (C00-C97)	48	5.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	5.0	---
7	Diabetes mellitus (E10-E14)	22	2.4	---
8	Chronic lower respiratory diseases (J40-J47)	21	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	1.7	---
11	Pneumonitis due to solids and liquids (J69)	9	1.0	---
12	Septicemia (A40-A41)	6	0.6	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	---
13	Nutritional deficiencies (E40-E64)	5	0.5	---
13	Atherosclerosis (I70)	5	0.5	---
...	All other causes (residual)	238	25.6	---
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	37,236	100.0	386.2
1	Malignant neoplasms (C00-C97)	9,243	24.8	95.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,418	22.6	87.3
3	Cerebrovascular diseases (I60-I69)	2,442	6.6	25.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,074	5.6	21.5
5	Diabetes mellitus (E10-E14)	1,612	4.3	16.7
6	Chronic lower respiratory diseases (J40-J47)	1,198	3.2	12.4
7	Influenza and pneumonia (J09-J18)	1,155	3.1	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	2.7	10.3
9	Alzheimer disease (G30)	790	2.1	8.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	789	2.1	8.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	654	1.8	6.8
12	Parkinson disease (G20-G21)	489	1.3	5.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	468	1.3	4.9
14	Septicemia (A40-A41)	460	1.2	4.8
15	Pneumonitis due to solids and liquids (J69)	317	0.9	3.3
...	All other causes (residual)	6,137	16.5	63.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	14.6	*
3	Malignant neoplasms (C00-C97)	13	13.5	*
4	Influenza and pneumonia (J09-J18)	3	3.1	*
5	Septicemia (A40-A41)	2	2.1	*
5	Meningitis (G00,G03)	2	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
5	Cerebrovascular diseases (I60-I69)	2	2.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.1	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (residual)	28	29.2	5.7
Non-Hispanic, Asian or Pacific Islander, male, 5-9 years				
...	All causes	52	100.0	8.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	19.2	*
2	Malignant neoplasms (C00-C97)	7	13.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	11.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.6	*
5	Influenza and pneumonia (J09-J18)	2	3.8	*
6	Anemias (D50-D64)	1	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.9	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	1.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.9	*
...	All other causes (residual)	18	34.6	*
Non-Hispanic, Asian or Pacific Islander, male, 10-14 years				
...	All causes	62	100.0	10.1
1	Malignant neoplasms (C00-C97)	16	25.8	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	17.7	*
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	12.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	9.7	*
5	Septicemia (A40-A41)	2	3.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.2	*
7	Meningococcal infection (A39)	1	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	*
7	Anemias (D50-D64)	1	1.6	*
7	Meningitis (G00,G03)	1	1.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
7	Influenza and pneumonia (J09-J18)	1	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.6	*
...	All other causes (residual)	9	14.5	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 15-19 years				
...	All causes	222	100.0	35.9
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	32.4	11.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	27.0	9.7
3	Malignant neoplasms (C00-C97)	21	9.5	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.1	*
6	Cerebrovascular diseases (I60-I69)	4	1.8	*
6	Influenza and pneumonia (J09-J18)	4	1.8	*
8	Septicemia (A40-A41)	2	0.9	*
8	Anemias (D50-D64)	2	0.9	*
8	Aortic aneurysm and dissection (I71)	2	0.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	26	11.7	4.2
Non-Hispanic, Asian or Pacific Islander, male, 20-24 years				
...	All causes	445	100.0	61.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	35.1	21.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	32.1	19.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.4	4.5
4	Malignant neoplasms (C00-C97)	27	6.1	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.9	*
6	Influenza and pneumonia (J09-J18)	7	1.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.6	*
8	Diabetes mellitus (E10-E14)	2	0.4	*
8	Cerebrovascular diseases (I60-I69)	2	0.4	*
8	Pneumonitis due to solids and liquids (J69)	2	0.4	*
8	Legal intervention (Y35,Y89.0)	2	0.4	*
12	Septicemia (A40-A41)	1	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Peptic ulcer (K25-K28)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	45	10.1	6.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 25-29 years				
...	All causes	477	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	38.6	21.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	24.3	13.6
3	Malignant neoplasms (C00-C97)	31	6.5	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.1	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.8	2.7
6	Diabetes mellitus (E10-E14)	6	1.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.0	*
7	Legal intervention (Y35,Y89.0)	5	1.0	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
11	Septicemia (A40-A41)	3	0.6	*
11	Aortic aneurysm and dissection (I71)	3	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	55	11.5	6.4
Non-Hispanic, Asian or Pacific Islander, male, 30-34 years				
...	All causes	536	100.0	64.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.8	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	20.3	13.1
3	Malignant neoplasms (C00-C97)	49	9.1	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	8.4	5.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	6.2	4.0
6	Cerebrovascular diseases (I60-I69)	16	3.0	*
7	Diabetes mellitus (E10-E14)	8	1.5	*
7	Influenza and pneumonia (J09-J18)	8	1.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.3	*
10	Aortic aneurysm and dissection (I71)	6	1.1	*
11	Septicemia (A40-A41)	3	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
14	Viral hepatitis (B15-B19)	2	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
14	Peptic ulcer (K25-K28)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	69	12.9	8.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 35-39 years				
...	All causes	584	100.0	76.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	21.1	16.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	16.3	12.4
3	Malignant neoplasms (C00-C97)	91	15.6	11.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	13.2	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	26	4.5	3.4
6	Cerebrovascular diseases (I60-I69)	24	4.1	3.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.7	*
8	Diabetes mellitus (E10-E14)	14	2.4	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.7	*
10	Aortic aneurysm and dissection (I71)	7	1.2	*
10	Influenza and pneumonia (J09-J18)	7	1.2	*
12	Viral hepatitis (B15-B19)	4	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
15	Septicemia (A40-A41)	3	0.5	*
15	Legal intervention (Y35,Y89.0)	3	0.5	*
...	All other causes (residual)	76	13.0	9.9
Non-Hispanic, Asian or Pacific Islander, male, 40-44 years				
...	All causes	790	100.0	109.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	20.8	22.7
2	Malignant neoplasms (C00-C97)	162	20.5	22.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	15.1	16.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	8.1	8.9
5	Cerebrovascular diseases (I60-I69)	51	6.5	7.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.3	3.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.9	3.2
8	Diabetes mellitus (E10-E14)	18	2.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.1	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.1	*
11	Viral hepatitis (B15-B19)	8	1.0	*
11	Influenza and pneumonia (J09-J18)	8	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
...	All other causes (residual)	114	14.4	15.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,181	100.0	173.3
1	Malignant neoplasms (C00-C97)	318	26.9	46.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	20.5	35.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	11.6	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	6.4	11.0
5	Cerebrovascular diseases (I60-I69)	66	5.6	9.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	49	4.1	7.2
7	Diabetes mellitus (E10-E14)	42	3.6	6.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.5	*
9	Influenza and pneumonia (J09-J18)	17	1.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.3	*
11	Aortic aneurysm and dissection (I71)	13	1.1	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	Viral hepatitis (B15-B19)	10	0.8	*
15	Septicemia (A40-A41)	6	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
...	All other causes (residual)	142	12.0	20.8
Non-Hispanic, Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,613	100.0	272.4
1	Malignant neoplasms (C00-C97)	478	29.6	80.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	23.4	63.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	7.1	19.4
4	Cerebrovascular diseases (I60-I69)	88	5.5	14.9
5	Diabetes mellitus (E10-E14)	72	4.5	12.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	3.9	10.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	57	3.5	9.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.4	3.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.4	3.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	1.2	3.4
11	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.1	*
13	Septicemia (A40-A41)	16	1.0	*
13	Influenza and pneumonia (J09-J18)	16	1.0	*
15	Viral hepatitis (B15-B19)	13	0.8	*
15	Aortic aneurysm and dissection (I71)	13	0.8	*
...	All other causes (residual)	203	12.6	34.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 55-59 years				
...	All causes	2,133	100.0	402.2
1	Malignant neoplasms (C00-C97)	713	33.4	134.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	506	23.7	95.4
3	Cerebrovascular diseases (I60-I69)	127	6.0	23.9
4	Diabetes mellitus (E10-E14)	121	5.7	22.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	5.7	22.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	3.1	12.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	52	2.4	9.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.7	6.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.7	6.8
10	Septicemia (A40-A41)	30	1.4	5.7
11	Influenza and pneumonia (J09-J18)	26	1.2	4.9
12	Chronic lower respiratory diseases (J40-J47)	23	1.1	4.3
13	Viral hepatitis (B15-B19)	22	1.0	4.1
14	Aortic aneurysm and dissection (I71)	19	0.9	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.8	*
...	All other causes (residual)	219	10.3	41.3
Non-Hispanic, Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,898	100.0	625.4
1	Malignant neoplasms (C00-C97)	1,021	35.2	220.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	701	24.2	151.3
3	Cerebrovascular diseases (I60-I69)	172	5.9	37.1
4	Diabetes mellitus (E10-E14)	166	5.7	35.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	4.0	25.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	2.3	14.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	60	2.1	12.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	1.7	10.8
9	Influenza and pneumonia (J09-J18)	48	1.7	10.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.5	9.5
11	Chronic lower respiratory diseases (J40-J47)	41	1.4	8.8
12	Septicemia (A40-A41)	39	1.3	8.4
13	Viral hepatitis (B15-B19)	21	0.7	4.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.5	*
15	Aortic aneurysm and dissection (I71)	13	0.4	*
...	All other causes (residual)	324	11.2	69.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 65-69 years				
...	All causes	3,600	100.0	957.5
1	Malignant neoplasms (C00-C97)	1,273	35.4	338.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	861	23.9	229.0
3	Cerebrovascular diseases (I60-I69)	202	5.6	53.7
4	Diabetes mellitus (E10-E14)	182	5.1	48.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	3.6	34.3
6	Chronic lower respiratory diseases (J40-J47)	92	2.6	24.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.9	18.6
8	Influenza and pneumonia (J09-J18)	68	1.9	18.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.7	16.5
10	Septicemia (A40-A41)	51	1.4	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	1.4	13.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	1.2	11.4
13	Aortic aneurysm and dissection (I71)	30	0.8	8.0
14	Viral hepatitis (B15-B19)	21	0.6	5.6
15	Parkinson disease (G20-G21)	20	0.6	5.3
...	All other causes (residual)	447	12.4	118.9
Non-Hispanic, Asian or Pacific Islander, male, 70-74 years				
...	All causes	3,752	100.0	1,426.1
1	Malignant neoplasms (C00-C97)	1,202	32.0	456.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	870	23.2	330.7
3	Cerebrovascular diseases (I60-I69)	253	6.7	96.2
4	Diabetes mellitus (E10-E14)	211	5.6	80.2
5	Chronic lower respiratory diseases (J40-J47)	143	3.8	54.4
6	Influenza and pneumonia (J09-J18)	106	2.8	40.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	2.7	38.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.5	36.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.6	22.8
10	Parkinson disease (G20-G21)	59	1.6	22.4
11	Septicemia (A40-A41)	52	1.4	19.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.9	13.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	0.9	12.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.8	11.0
15	Aortic aneurysm and dissection (I71)	24	0.6	9.1
...	All other causes (residual)	479	12.8	182.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 75-79 years				
...	All causes	4,393	100.0	2,418.3
1	Malignant neoplasms (C00-C97)	1,282	29.2	705.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	995	22.6	547.7
3	Cerebrovascular diseases (I60-I69)	290	6.6	159.6
4	Diabetes mellitus (E10-E14)	243	5.5	133.8
5	Chronic lower respiratory diseases (J40-J47)	181	4.1	99.6
6	Influenza and pneumonia (J09-J18)	139	3.2	76.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	2.8	68.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.4	57.8
9	Parkinson disease (G20-G21)	85	1.9	46.8
10	Alzheimer disease (G30)	70	1.6	38.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	1.5	35.8
12	Septicemia (A40-A41)	63	1.4	34.7
13	Pneumonitis due to solids and liquids (J69)	39	0.9	21.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	39	0.9	21.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	17.6
...	All other causes (residual)	640	14.6	352.3
Non-Hispanic, Asian or Pacific Islander, male, 80-84 years				
...	All causes	4,705	100.0	4,126.1
1	Malignant neoplasms (C00-C97)	1,114	23.7	976.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,107	23.5	970.8
3	Cerebrovascular diseases (I60-I69)	374	7.9	328.0
4	Chronic lower respiratory diseases (J40-J47)	229	4.9	200.8
5	Diabetes mellitus (E10-E14)	226	4.8	198.2
6	Influenza and pneumonia (J09-J18)	181	3.8	158.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	3.2	131.5
8	Alzheimer disease (G30)	149	3.2	130.7
9	Parkinson disease (G20-G21)	121	2.6	106.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	2.4	99.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	2.0	82.4
12	Septicemia (A40-A41)	63	1.3	55.2
13	Pneumonitis due to solids and liquids (J69)	62	1.3	54.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.9	35.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.5	21.9
...	All other causes (residual)	657	14.0	576.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 85-89 years				
...	All causes	4,436	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,078	24.3	---
2	Malignant neoplasms (C00-C97)	824	18.6	---
3	Cerebrovascular diseases (I60-I69)	381	8.6	---
4	Chronic lower respiratory diseases (J40-J47)	224	5.0	---
5	Influenza and pneumonia (J09-J18)	219	4.9	---
6	Alzheimer disease (G30)	217	4.9	---
7	Diabetes mellitus (E10-E14)	170	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.8	---
9	Parkinson disease (G20-G21)	124	2.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	103	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	2.2	---
12	Pneumonitis due to solids and liquids (J69)	72	1.6	---
13	Septicemia (A40-A41)	56	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.8	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	21	0.5	---
...	All other causes (residual)	687	15.5	---
Non-Hispanic, Asian or Pacific Islander, male, 90-94 years				
...	All causes	3,147	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	861	27.4	---
2	Malignant neoplasms (C00-C97)	439	13.9	---
3	Cerebrovascular diseases (I60-I69)	264	8.4	---
4	Alzheimer disease (G30)	200	6.4	---
5	Influenza and pneumonia (J09-J18)	180	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	155	4.9	---
7	Diabetes mellitus (E10-E14)	92	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	2.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.1	---
11	Pneumonitis due to solids and liquids (J69)	59	1.9	---
12	Parkinson disease (G20-G21)	48	1.5	---
13	Septicemia (A40-A41)	37	1.2	---
14	Nutritional deficiencies (E40-E64)	19	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	---
...	All other causes (residual)	530	16.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 95-99 years				
...	All causes	1,344	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	391	29.1	---
2	Malignant neoplasms (C00-C97)	140	10.4	---
3	Cerebrovascular diseases (I60-I69)	103	7.7	---
4	Alzheimer disease (G30)	94	7.0	---
4	Influenza and pneumonia (J09-J18)	94	7.0	---
6	Chronic lower respiratory diseases (J40-J47)	69	5.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.8	---
9	Diabetes mellitus (E10-E14)	33	2.5	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.0	---
11	Septicemia (A40-A41)	22	1.6	---
12	Pneumonitis due to solids and liquids (J69)	21	1.6	---
13	Parkinson disease (G20-G21)	17	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	---
15	Nutritional deficiencies (E40-E64)	8	0.6	---
...	All other causes (residual)	239	17.8	---
Non-Hispanic, Asian or Pacific Islander, male, 100 years and over				
...	All causes	242	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	24.4	---
2	Malignant neoplasms (C00-C97)	22	9.1	---
3	Alzheimer disease (G30)	18	7.4	---
3	Cerebrovascular diseases (I60-I69)	18	7.4	---
5	Influenza and pneumonia (J09-J18)	14	5.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	5.0	---
7	Chronic lower respiratory diseases (J40-J47)	9	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	3.7	---
9	Diabetes mellitus (E10-E14)	6	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.5	---
11	Septicemia (A40-A41)	3	1.2	---
12	Parkinson disease (G20-G21)	2	0.8	---
12	Pneumonitis due to solids and liquids (J69)	2	0.8	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
15	Anemias (D50-D64)	1	0.4	---
15	Atherosclerosis (I70)	1	0.4	---
15	Aortic aneurysm and dissection (I71)	1	0.4	---
15	Peptic ulcer (K25-K28)	1	0.4	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	---
...	All other causes (residual)	54	22.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	35,362	100.0	335.6
1	Malignant neoplasms (C00-C97)	9,014	25.5	85.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,072	20.0	67.1
3	Cerebrovascular diseases (I60-I69)	2,988	8.4	28.4
4	Alzheimer disease (G30)	1,793	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,435	4.1	13.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	3.3	11.2
7	Influenza and pneumonia (J09-J18)	1,131	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	865	2.4	8.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	818	2.3	7.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	745	2.1	7.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	1.2	4.0
12	Septicemia (A40-A41)	419	1.2	4.0
13	Parkinson disease (G20-G21)	380	1.1	3.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	257	0.7	2.4
15	Pneumonitis due to solids and liquids (J69)	232	0.7	2.2
...	All other causes (residual)	6,607	18.7	62.7
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	52	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	19.2	*
2	Malignant neoplasms (C00-C97)	7	13.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	13.5	*
4	Anemias (D50-D64)	3	5.8	*
4	Influenza and pneumonia (J09-J18)	3	5.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.8	*
9	Septicemia (A40-A41)	1	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.9	*
9	Hernia (K40-K46)	1	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.9	*
...	All other causes (residual)	12	23.1	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 5-9 years				
...	All causes	48	100.0	8.3
1	Malignant neoplasms (C00-C97)	11	22.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	20.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	12.5	*
4	Septicemia (A40-A41)	3	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.2	*
6	Anemias (D50-D64)	1	2.1	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
...	All other causes (residual)	13	27.1	*
Non-Hispanic, Asian or Pacific Islander, female, 10-14 years				
...	All causes	46	100.0	7.6
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	23.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	19.6	*
3	Malignant neoplasms (C00-C97)	8	17.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
5	Cerebrovascular diseases (I60-I69)	2	4.3	*
7	Septicemia (A40-A41)	1	2.2	*
7	Diabetes mellitus (E10-E14)	1	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (residual)	8	17.4	*
Non-Hispanic, Asian or Pacific Islander, female, 15-19 years				
...	All causes	103	100.0	16.9
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	32.0	5.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	20.4	3.4
3	Malignant neoplasms (C00-C97)	13	12.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
6	Diabetes mellitus (E10-E14)	1	1.0	*
6	Influenza and pneumonia (J09-J18)	1	1.0	*
6	Pneumonitis due to solids and liquids (J69)	1	1.0	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.0	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.0	*
6	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.0	*
...	All other causes (residual)	16	15.5	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 20-24 years				
...	All causes	190	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	31.1	8.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	28.4	7.6
3	Malignant neoplasms (C00-C97)	18	9.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.7	*
6	Influenza and pneumonia (J09-J18)	3	1.6	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
10	Septicemia (A40-A41)	1	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
10	Diabetes mellitus (E10-E14)	1	0.5	*
10	Meningitis (G00,G03)	1	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (residual)	28	14.7	3.9
Non-Hispanic, Asian or Pacific Islander, female, 25-29 years				
...	All causes	203	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	22.7	5.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	20.2	4.7
3	Malignant neoplasms (C00-C97)	40	19.7	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.4	*
6	Diabetes mellitus (E10-E14)	4	2.0	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.0	*
8	Cerebrovascular diseases (I60-I69)	3	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
11	Viral hepatitis (B15-B19)	1	0.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
11	Anemias (D50-D64)	1	0.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	*
11	Influenza and pneumonia (J09-J18)	1	0.5	*
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
11	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	34	16.7	3.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 30-34 years				
...	All causes	284	100.0	31.6
1	Malignant neoplasms (C00-C97)	86	30.3	9.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	13.4	4.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	11.6	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.9	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	14	4.9	*
7	Cerebrovascular diseases (I60-I69)	11	3.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	2.5	*
9	Influenza and pneumonia (J09-J18)	6	2.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.4	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.4	*
12	Septicemia (A40-A41)	3	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
14	Anemias (D50-D64)	1	0.4	*
14	Diabetes mellitus (E10-E14)	1	0.4	*
14	Nutritional deficiencies (E40-E64)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
14	Peptic ulcer (K25-K28)	1	0.4	*
...	All other causes (residual)	36	12.7	4.0
Non-Hispanic, Asian or Pacific Islander, female, 35-39 years				
...	All causes	360	100.0	41.5
1	Malignant neoplasms (C00-C97)	137	38.1	15.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	11.1	4.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	10.0	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.9	2.9
5	Cerebrovascular diseases (I60-I69)	14	3.9	*
6	Diabetes mellitus (E10-E14)	11	3.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.1	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8	2.2	*
9	Septicemia (A40-A41)	7	1.9	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.1	*
13	Influenza and pneumonia (J09-J18)	3	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
...	All other causes (residual)	47	13.1	5.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 40-44 years				
...	All causes	493	100.0	60.1
1	Malignant neoplasms (C00-C97)	215	43.6	26.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	7.9	4.8
2	Cerebrovascular diseases (I60-I69)	39	7.9	4.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	7.1	4.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	5.7	3.4
6	Diabetes mellitus (E10-E14)	14	2.8	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	11	2.2	*
8	Septicemia (A40-A41)	8	1.6	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.6	*
10	Influenza and pneumonia (J09-J18)	7	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Alzheimer disease (G30)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
...	All other causes (residual)	58	11.8	7.1
Non-Hispanic, Asian or Pacific Islander, female, 45-49 years				
...	All causes	826	100.0	106.4
1	Malignant neoplasms (C00-C97)	414	50.1	53.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	7.3	7.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	7.0	7.5
4	Cerebrovascular diseases (I60-I69)	46	5.6	5.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	5.0	5.3
6	Diabetes mellitus (E10-E14)	30	3.6	3.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.2	*
10	Septicemia (A40-A41)	8	1.0	*
10	Influenza and pneumonia (J09-J18)	8	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.6	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
...	All other causes (residual)	99	12.0	12.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 50-54 years				
...	All causes	1,065	100.0	157.1
1	Malignant neoplasms (C00-C97)	575	54.0	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	11.3	17.7
3	Cerebrovascular diseases (I60-I69)	75	7.0	11.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	4.0	6.3
5	Diabetes mellitus (E10-E14)	38	3.6	5.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	2.6	4.1
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.0	*
8	Septicemia (A40-A41)	10	0.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
10	Influenza and pneumonia (J09-J18)	9	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
...	All other causes (residual)	116	10.9	17.1
Non-Hispanic, Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,478	100.0	237.2
1	Malignant neoplasms (C00-C97)	772	52.2	123.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	12.7	30.2
3	Cerebrovascular diseases (I60-I69)	90	6.1	14.4
4	Diabetes mellitus (E10-E14)	63	4.3	10.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	3.6	8.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.6	3.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.4	3.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.4	3.2
9	Chronic lower respiratory diseases (J40-J47)	19	1.3	*
10	Septicemia (A40-A41)	18	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	*
12	Influenza and pneumonia (J09-J18)	10	0.7	*
14	Aortic aneurysm and dissection (I71)	8	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.4	*
...	All other causes (residual)	155	10.5	24.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,993	100.0	352.0
1	Malignant neoplasms (C00-C97)	968	48.6	171.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	275	13.8	48.6
3	Cerebrovascular diseases (I60-I69)	112	5.6	19.8
4	Diabetes mellitus (E10-E14)	92	4.6	16.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	3.1	10.8
6	Influenza and pneumonia (J09-J18)	46	2.3	8.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.1	7.4
8	Chronic lower respiratory diseases (J40-J47)	32	1.6	5.7
9	Septicemia (A40-A41)	28	1.4	4.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.3	4.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.1	3.9
12	Viral hepatitis (B15-B19)	20	1.0	3.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.7	*
15	Aortic aneurysm and dissection (I71)	11	0.6	*
...	All other causes (residual)	228	11.4	40.3
Non-Hispanic, Asian or Pacific Islander, female, 65-69 years				
...	All causes	2,660	100.0	561.7
1	Malignant neoplasms (C00-C97)	1,126	42.3	237.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	448	16.8	94.6
3	Cerebrovascular diseases (I60-I69)	169	6.4	35.7
4	Diabetes mellitus (E10-E14)	121	4.5	25.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	2.7	15.2
6	Influenza and pneumonia (J09-J18)	59	2.2	12.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.1	11.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.4	7.8
9	Chronic lower respiratory diseases (J40-J47)	34	1.3	7.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.2	6.8
11	Septicemia (A40-A41)	30	1.1	6.3
12	Parkinson disease (G20-G21)	25	0.9	5.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	0.9	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
14	Alzheimer disease (G30)	18	0.7	*
14	Aortic aneurysm and dissection (I71)	18	0.7	*
...	All other causes (residual)	374	14.1	79.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,881	100.0	878.2
1	Malignant neoplasms (C00-C97)	1,007	35.0	306.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	531	18.4	161.9
3	Cerebrovascular diseases (I60-I69)	250	8.7	76.2
4	Diabetes mellitus (E10-E14)	160	5.6	48.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	3.0	26.2
6	Chronic lower respiratory diseases (J40-J47)	72	2.5	21.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.2	19.5
8	Influenza and pneumonia (J09-J18)	57	2.0	17.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.7	15.2
10	Septicemia (A40-A41)	45	1.6	13.7
11	Alzheimer disease (G30)	40	1.4	12.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.3	11.6
13	Parkinson disease (G20-G21)	36	1.2	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
...	All other causes (residual)	411	14.3	125.3
Non-Hispanic, Asian or Pacific Islander, female, 75-79 years				
...	All causes	3,642	100.0	1,598.9
1	Malignant neoplasms (C00-C97)	1,066	29.3	468.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	732	20.1	321.4
3	Cerebrovascular diseases (I60-I69)	319	8.8	140.0
4	Diabetes mellitus (E10-E14)	187	5.1	82.1
5	Alzheimer disease (G30)	111	3.0	48.7
6	Influenza and pneumonia (J09-J18)	108	3.0	47.4
7	Chronic lower respiratory diseases (J40-J47)	104	2.9	45.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.8	44.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.7	43.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	2.0	31.6
11	Parkinson disease (G20-G21)	50	1.4	22.0
12	Septicemia (A40-A41)	49	1.3	21.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.9	13.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.7	11.9
15	Viral hepatitis (B15-B19)	20	0.5	8.8
...	All other causes (residual)	566	15.5	248.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,790	100.0	3,091.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,056	22.0	681.4
2	Malignant neoplasms (C00-C97)	992	20.7	640.1
3	Cerebrovascular diseases (I60-I69)	467	9.7	301.4
4	Diabetes mellitus (E10-E14)	230	4.8	148.4
5	Alzheimer disease (G30)	228	4.8	147.1
6	Chronic lower respiratory diseases (J40-J47)	174	3.6	112.3
7	Influenza and pneumonia (J09-J18)	151	3.2	97.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	3.1	95.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	3.0	91.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	2.6	80.7
11	Parkinson disease (G20-G21)	92	1.9	59.4
12	Septicemia (A40-A41)	69	1.4	44.5
13	Pneumonitis due to solids and liquids (J69)	35	0.7	22.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.6	18.7
15	Aortic aneurysm and dissection (I71)	23	0.5	14.8
...	All other causes (residual)	829	17.3	535.0
Non-Hispanic, Asian or Pacific Islander, female, 85-89 years				
...	All causes	5,562	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,270	22.8	---
2	Malignant neoplasms (C00-C97)	866	15.6	---
3	Cerebrovascular diseases (I60-I69)	584	10.5	---
4	Alzheimer disease (G30)	438	7.9	---
5	Diabetes mellitus (E10-E14)	236	4.2	---
6	Influenza and pneumonia (J09-J18)	218	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	169	3.0	---
8	Chronic lower respiratory diseases (J40-J47)	159	2.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	2.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	2.4	---
11	Parkinson disease (G20-G21)	98	1.8	---
12	Septicemia (A40-A41)	60	1.1	---
13	Pneumonitis due to solids and liquids (J69)	52	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	30	0.5	---
...	All other causes (residual)	1,074	19.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 90-94 years				
...	All causes	5,125	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,354	26.4	---
2	Cerebrovascular diseases (I60-I69)	539	10.5	---
3	Alzheimer disease (G30)	514	10.0	---
4	Malignant neoplasms (C00-C97)	500	9.8	---
5	Influenza and pneumonia (J09-J18)	253	4.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	170	3.3	---
8	Diabetes mellitus (E10-E14)	169	3.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.9	---
11	Parkinson disease (G20-G21)	58	1.1	---
12	Pneumonitis due to solids and liquids (J69)	56	1.1	---
13	Septicemia (A40-A41)	45	0.9	---
14	Nutritional deficiencies (E40-E64)	31	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	---
15	Aortic aneurysm and dissection (I71)	20	0.4	---
...	All other causes (residual)	1,009	19.7	---
Non-Hispanic, Asian or Pacific Islander, female, 95-99 years				
...	All causes	2,443	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	711	29.1	---
2	Alzheimer disease (G30)	335	13.7	---
3	Cerebrovascular diseases (I60-I69)	225	9.2	---
4	Malignant neoplasms (C00-C97)	163	6.7	---
5	Influenza and pneumonia (J09-J18)	148	6.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	3.1	---
7	Diabetes mellitus (E10-E14)	60	2.5	---
8	Chronic lower respiratory diseases (J40-J47)	57	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.7	---
11	Septicemia (A40-A41)	26	1.1	---
12	Pneumonitis due to solids and liquids (J69)	23	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
13	Nutritional deficiencies (E40-E64)	10	0.4	---
13	Atherosclerosis (I70)	10	0.4	---
...	All other causes (residual)	500	20.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 100 years and over				
...	All causes	689	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	29.6	---
2	Alzheimer disease (G30)	100	14.5	---
3	Cerebrovascular diseases (I60-I69)	40	5.8	---
4	Influenza and pneumonia (J09-J18)	38	5.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	5.1	---
6	Malignant neoplasms (C00-C97)	26	3.8	---
7	Diabetes mellitus (E10-E14)	16	2.3	---
8	Chronic lower respiratory diseases (J40-J47)	12	1.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.7	---
10	Pneumonitis due to solids and liquids (J69)	7	1.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.0	---
12	Nutritional deficiencies (E40-E64)	5	0.7	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	---
13	Atherosclerosis (I70)	4	0.6	---
15	Septicemia (A40-A41)	3	0.4	---
...	All other causes (residual)	176	25.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017