Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
All causes  All origins, both sexes.  Male. Female. Hispanic, both sexes. Male. Female. Mexican, both sexes. Male. Female. Puerto Rican, both sexes. Male. Female. Cuban, both sexes. Male. Female. Cuban, both sexes. Male. Female. Contral and South American, both sexes. Male. Female. Other and Unknown Hispanic, both sexes. Male. Female. Non-Hispanic 1/, both sexes. Male. Female. Non-Hispanic white, both sexes.	1,181,668 1,215,947 122,416 68,544 53,872 69,285 40,096 29,189 15,421 8,445 6,976 12,308 6,371 5,937 8,964 4,704 4,260 16,438 8,928 7,510 2,269,583 1,109,848 1,159,735	27,936 15,718 12,218 5,321 2,948 2,373 3,844 2,121 1,723 439 251 188 60 32 28 336 177 159 642 267 275 22,370 12,633 9,737 13,046	4,785 2,649 2,136 947 535 412 724 405 319 64 38 26 9 9 6 3 362 28 88 88 82 28 82 34 28 28 28 28 28 28 28 28 28 28 28 28 28	2,888 1,645 1,243 487 273 214 336 190 146 44 20 10 6 4 34 15 19 63 38 25 2,390 1,364 1,026 1,573	3,946 2,354 1,592 618 383 235 438 269 169 59 38 21 12 27 7 5 35 22 13 74 47 27 3,309 1,963 1,963 1,946 2,247	13,706 9,678 4,028 2,141 1,662 479 1,518 1,200 318 152 111 41 47 37 10 171 122 49 253 192 61 11,522 7,987 3,535 8,346	19,715 14,909 4,806 3,287 2,714 573 2,307 1,942 365 255 199 56 45 34 11 330 262 68 350 277 73 16,374 12,158 4,216 11,338	18,771 13,554 5,217 3,019 2,361 658 2,077 1,674 403 266 189 77 42 31 11 304 229 75 330 238 92 15,695 11,154 4,541 10,452	22,097 14,805 7,292 2,940 2,186 754 1,920 1,440 480 316 226 90 60 44 16 310 228 82 334 248 86 19,079 12,556 12,945	31,944 20,485 11,459 3,686 2,636 1,050 2,245 1,647 598 516 331 185 112 26 350 245 105 463 327 136 28,153 17,774 10,379 20,033	53,418 33,192 20,226 4,860 3,337 1,523 2,919 2,055 864 678 445 233 197 147 50 417 269 148 649 421 228 48,371 29,727 18,644 35,510	77,927 48,514 29,413 5,903 3,984 1,919 3,401 2,349 1,052 921 612 309 270 193 77 501 302 199 810 528 282 71,740 44,331 27,409 53,263
Male  Female  Non-Hispanic black, both sexes	938,143 995,239 283,859	7,399 5,647 8,221	1,365 1,093 1,128	914 659 665	1,350 897 886	5,636 2,710 2,633	8,381 2,957 4,291	7,473 2,979 4,541	8,625 4,320 5,227	12,860 7,173 7,003	22,393 13,117 11,234	33,465 19,798 16,211
Male Female. Origin not stated 2/, both sexes Male Female.	144,022 139,837 5,616 3,276 2,340	4,613 3,608 245 137	623 505 12 8 4	361 304 11 8	509 377 19 8	1,986 647 43 29	3,253 1,038 54 37	3,209 1,332 57 39 18	3,355 1,872 78 63 15	4,199 2,804 105 75 30	6,353 4,881 187 128 59	9,495 6,716 284 199 85

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Salmonella infections (A01-A02) All origins, both sexes..... 30 1 1 1 2 Male.... 2 Female..... 8 Hispanic, both sexes..... 1 1 Male.... 1 Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 1 Male.... 1 Female..... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 2.7 Non-Hispanic 1/, both sexes..... 1 19 Male.... 1 1 8 1 Female..... Non-Hispanic white, both sexes..... 17 Male.... 11 Female..... 6 1 Non-Hispanic black, both sexes..... 4 3 1 Male.... Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Shigellosis and amebiasis (A03,A06)												
All origins, both sexes	3	_	_	_	1	_	1	_	_	_	_	_
Male	2	_	-	_	1	_	-	-	-	-	_	-
Female	1	-	-	_	-	_	1	-	-	-	-	_
Hispanic, both sexes	2	-	-	_	1	_	1	-	-	-	-	_
Male	1	-	_	_	1	-	-	-	-	_	_	-
Female	1	-	_	_	-	-	1	-	-	_	_	-
Mexican, both sexes	2	-	_	_	1	-	1	-	-	_	_	-
Male	1	-	_	_	1	-	-	-	-	_	_	-
Female	1	-	_	_	-	-	1	-	-	_	_	-
Puerto Rican, both sexes	-	-	_	_	-	-	-	-	-	_	_	-
Male	_	-	-	-	-	-	-	-	-	_	-	-
Female	_	-	-	-	-	-	-	-	-	_	-	-
Cuban, both sexes	_	-	-	-	-	-	-	-	-	_	-	-
Male	_	-	-	-	-	-	-	-	-	_	-	-
Female	_	-	-	-	-	-	-	-	-	_	-	-
Central and South American, both sexes.	_	-	-	-	-	-	-	-	-	_	-	-
Male	_	-	-	-	-	-	-	-	-	_	-	_
Female	_	-	-	-	-	-	-	-	-	_	-	_
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	_	-	-	-	-	-	-	-	-	_	-	-
Female	_	-	-	-	-	-	-	-	-	_	-	-
Non-Hispanic 1/, both sexes	1	-	_	_	-	-	-	-	-	_	_	-
Male	1	-	-	-	-	-	-	-	-	-	_	-
Female	-	-	_	_	-	-	-	-	-	_	_	-
Non-Hispanic white, both sexes	1	_	-	-	-	-	-	-	-	-	_	-
Male	1	-	-	-	-	-	-	-	-	_	-	-
Female	_	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	_	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Certain other intestinal infections (A04,A07-A09) All origins, both sexes..... 4,357 Male.... 1.622 Female..... 2.735 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 5.0 Puerto Rican, both sexes..... Male.... 1.0 Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 4,200 1,560 Male.... 2,640 Female..... Non-Hispanic white, both sexes..... 3,919 1,440 Male.... 2,479 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Tuberculosis (A16-A19) All origins, both sexes..... 1.0 2.4 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Ω Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Respiratory tuberculosis (A16) All origins, both sexes..... 2.0 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 5.8 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male..... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other tuberculosis (A17-A19) All origins, both sexes..... 153 1 3 2 3 8 9 Male.... 90 1 2 3 Female..... 63 1 1 1 4 2 Hispanic, both sexes..... 27 1 2 2 1 Male.... Female..... 8 Mexican, both sexes..... 17 1 1 2 1 Male.... 13 1 1 1 1 Female..... 4 1 1 Puerto Rican, both sexes..... 1 Male.... 1 Female..... Cuban, both sexes..... 3 Male Female.... 3 1 1 Central and South American, both sexes. Male 1 Female..... Other and Unknown Hispanic, both sexes. 2 Male.... 2 Female..... 1 Non-Hispanic 1/, both sexes..... 126 1 71 Male.... 1 55 1 2 Female..... 3 Non-Hispanic white, both sexes..... 63 Male..... 33 1 4 30 Female..... Non-Hispanic black, both sexes..... 48 1 1 5 1 4 30 2 Male..... 1 5 1 Female..... 1.8 Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
All origins, both sexes	16	15	_	_	1	_	_	_	_	_	_	_
Male	10	9	-	_	1	_	-	-	-	_	_	-
Female	6	6	-	_	-	_	-	-	-	_	_	-
Hispanic, both sexes	6	5	-	_	1	_	-	-	-	_	_	-
Male	4	3	-	-	1	-	-	-	-	-	-	-
Female	2	2	_	-	-	-	_	-	-	-	-	_
Mexican, both sexes	3	3	_	-	-	-	_	-	-	-	-	_
Male	1	1	_	-	-	-	-	_	-	-	-	-
Female	2	2	_	-	-	-	_	-	-	-	-	_
Puerto Rican, both sexes	2	1	_	-	1	-	-	_	-	-	-	-
Male	2	1	_	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	9	9	-	-	-	-	-	-	-	-	-	-
Male	5	5	-	-	-	-	-	-	-	-	-	-
Female	4	4	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white, both sexes	8	8	-	-	-	-	-	-	-	-	-	-
Male	5	5	-	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
All origins, both sexes	3	_	_	_	_	_	_	_	-	_	_	_
Male	1	-	-	-	-	-	-	-	-	_	_	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Hispanic, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	_	-	-	-	-	-
Mexican, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	1	-	-	-	-	-	_	-	-	-	-	-
Male	1	-	-	-	-	-	_	-	-	-	-	-
Female	-	-	-	-	-	-	_	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	_	-	-	-	-	-
Male	-	-	-	-	-	-	_	-	-	-	-	-
Female	-	-	-	-	-	-	_	-	-	-	-	-
Non-Hispanic 1/, both sexes	2	_	_	_	_	_	-	-	_	-	-	_
Male	-	_	_	_	_	_	-	-	_	-	-	_
Female	2	_	_	_	_	_	-	-	_	-	-	_
Non-Hispanic white, both sexes	2	_	_	_	_	_	-	-	_	-	-	_
Male	-	_	_	_	_	_	-	-	_	-	-	_
Female	2	_	_	_	_	_	-	-	_	-	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	_	-	-	-	-	-
Male	-	_	-	-	_	-	-	-	-	-	-	-
Female	-	_	-	-	_	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	_	-	-	-	-	_	-	-	-	-	-
Male	-	_	-	-	-	-	_	-	-	-	-	-
Female	-	_	-	-	-	-	_	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
All origins, both sexes	138	8	18	8	5	16	15	9	5	6	9	7
Male	79	2	11	5	4	11	8	7	3	3	5	5
Female	59	6	7	3	1	5	7	2	2	3	4	2
Hispanic, both sexes	20	2	5	_	1	2	1	3	_	_	_	2
Male	12	1	2	_	1	_	_	3	_	_	_	2
Female	8	1	3	_	_	2	1	_	_	_	_	_
Mexican, both sexes	13	2	4	_	1	1	1	1	_	_	_	1
Male	7	1	1	_	1	_	_	1	_	_	_	1
Female	6	1	3	_	_	1	1	_	_	_	_	_
Puerto Rican, both sexes	1	_	_	_	_	_	_	1	_	_	_	_
Male	1	_	_	_	_	_	_	1	_	_	_	_
Female	_	-	-	_	-	-	-	-	-	_	_	-
Cuban, both sexes	1	-	1	_	-	-	-	-	-	_	_	-
Male	1	-	1	_	-	-	-	-	-	_	_	-
Female	_	-	-	_	-	-	-	-	-	_	_	-
Central and South American, both sexes.	1	-	-	_	-	-	-	1	-	_	_	-
Male	1	_	_	_	_	_	_	1	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	4	_	_	_	_	1	_	_	_	_	_	1
Male	2	_	_	_	_	_	_	_	_	_	_	1
Female	2	_	_	_	_	1	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	118	6	13	8	4	14	14	6	5	6	9	5
Male	67	1	9	5	3	11	8	4	3	3	5	3
Female	51	5	4	3	1	3	6	2	2	3	4	2
Non-Hispanic white, both sexes	90	5	10	7	4	10	11	5	2	5	5	4
Male	50	1	8	4	3	7	6	3	2	3	3	3
Female	40	4	2	3	1	3	5	2	_	2	2	1
Non-Hispanic black, both sexes	22	_	3	1	_	3	2	1	2	1	3	1
Male	13	-	1	1	-	3	1	1	1	_	1	-
Female	9	-	2	_	-	_	1	-	1	1	2	1
Origin not stated 2/, both sexes	_	-	_	_	-	_	-	-	-	_	_	_
Male	_	-	_	_	-	_	-	-	-	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
All origins, both sexes	33,373	271	84	38	28	48	87	115	213	306	540	898
Male	15,011	167	41	18	13	25	46	51	114	162	271	487
Female	18,362	104	43	20	15	23	41	64	99	144	269	411
Hispanic, both sexes	1,648	55	24	8	1	4	9	19	16	41	47	79
Male	801	36	14	6	_	2	5	6	10	23	30	52
Female	847	19	10	2	1	2	4	13	6	18	17	27
Mexican, both sexes	889	40	20	6	-	3	4	10	11	28	25	44
Male	420	27	11	5	-	2	2	3	6	16	17	29
Female	469	13	9	1	-	1	2	7	5	12	8	15
Puerto Rican, both sexes	273	6	1	1	-	-	3	3	4	5	12	20
Male	139	4	-	1	-	-	1	1	3	2	6	12
Female	134	2	1	-	-	_	2	2	1	3	6	8
Cuban, both sexes	158	1	1	-	-	1	-	1	-	1	3	1
Male	81	1	1	-	-	_	-	-	-	1	2	1
Female	77	-	_	-	-	1	-	1	-	-	1	-
Central and South American, both sexes.	135	4	_	1	1	_	1	2	-	2	1	7
Male	57	2	_	-	-	_	1	2	-	1	1	6
Female	78	2	_	1	1	_	-	-	-	1	-	1
Other and Unknown Hispanic, both sexes.	193	4	2	-	-	_	1	3	1	5	6	7
Male	104	2	2	-	-	_	1	-	1	3	4	4
Female	89	2	_	-	-	_	-	3	-	2	2	3
Non-Hispanic 1/, both sexes	31,656	212	60	30	27	44	78	96	196	265	493	817
Male	14,174	128	27	12	13	23	41	45	103	139	241	433
Female	17,482	84	33	18	14	21	37	51	93	126	252	384
Non-Hispanic white, both sexes	25,041	112	36	23	18	30	40	58	114	178	296	510
Male	11,188	68	15	11	9	17	22	28	57	99	144	267
Female	13,853	44	21	12	9	13	18	30	57	79	152	243
Non-Hispanic black, both sexes	5,945	90	19	5	7	13	32	34	68	73	182	273
Male	2,660	51	9	1	3	6	16	15	37	32	93	146
Female	3,285	39	10	4	4	7	16	19	31	41	89	127
Origin not stated 2/, both sexes	69	4	-	_	-	-	-	-	1	-	-	2
Male	36	3	-	_	-	-	-	-	1	-	-	2
Female	33	1	-	_	-	-	-	_	-	_	_	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
All origins, both sexes	43	_	_	_	_	_	_	_	3	1	2	_
Male	21	_	_	_	_	_	_	_	2	1	1	_
Female	22	_	_	_	_	_	_	_	1	_	1	_
Hispanic, both sexes	1	_	_	_	_	_	_	_	_	_	_	_
Male	1	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	1	-	_	_	-	_	-	_	-	_	-	-
Male	1	-	_	_	-	_	-	-	-	_	-	-
Female	-	-	_	_	-	_	-	-	-	_	-	-
Puerto Rican, both sexes	-	-	_	_	-	_	-	-	-	_	-	-
Male	-	-	_	_	-	_	-	-	-	_	-	-
Female	_	-	-	-	_	-	-	-	-	-	_	_
Cuban, both sexes	_	-	-	-	_	-	-	-	-	-	_	_
Male	_	-	-	-	_	-	-	-	-	-	_	_
Female	-	-	-	-	_	-	-	-	-	-	_	-
Central and South American, both sexes.	-	-	-	-	_	-	-	-	-	-	_	-
Male	-	-	-	-	_	-	-	-	-	-	-	-
Female	-	-	-	-	_	-	-	-	-	-	_	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	42	-	-	-	_	-	-	-	3	1	2	-
Male	20	-	-	-	_	-	-	-	2	1	1	-
Female	22	-	-	-	_	-	-	-	1	-	1	-
Non-Hispanic white, both sexes	16	-	-	-	_	-	-	-	-	1	2	-
Male	6	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	_	-	-	-	-	-	1	-
Non-Hispanic black, both sexes	23	-	-	-	-	-	-	-	3	-	-	-
Male	13	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute poliomyelitis (A80)												
All origins, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	_	-	_	-	_	-	-	_	-	_	_	-
Hispanic, both sexes	-	-	_	_	_	_	-	_	_	_	_	_
Male	-	-	_	_	_	_	-	_	_	_	_	_
Female	-	-	_	_	_	_	-	_	_	_	_	_
Mexican, both sexes	-	-	_	_	-	_	-	_	_	_	_	_
Male	_	_	_	-	_	_	_	-	-	_	_	_
Female	_	_	_	-	_	_	_	-	-	_	_	_
Puerto Rican, both sexes	_	_	_	-	_	_	_	-	-	_	_	_
Male	-	_	-	_	-	-	-	_	-	_	-	-
Female	-	_	-	_	-	-	-	_	-	_	-	-
Cuban, both sexes	-	_	-	_	-	-	-	_	-	_	-	-
Male	-	_	-	_	_	-	_	_	-	_	_	-
Female	-	_	-	_	_	-	_	_	-	_	_	-
Central and South American, both sexes.	-	_	-	_	_	-	_	_	-	_	_	-
Male	-	_	-	_	_	-	_	_	-	_	_	_
Female	-	_	-	_	-	-	-	_	-	_	-	_
Other and Unknown Hispanic, both sexes.	-	_	-	_	-	-	-	_	-	_	-	-
Male	-	_	-	_	-	-	-	_	-	_	-	-
Female	-	_	-	_	-	-	-	_	-	_	-	-
Non-Hispanic 1/, both sexes	_	_	_	-	_	_	_	-	-	_	_	_
Male	_	_	_	-	_	_	_	-	-	_	_	_
Female	_	_	_	-	_	_	_	-	-	_	_	_
Non-Hispanic white, both sexes	_	_	_	-	_	_	_	-	-	_	_	_
Male	_	_	_	-	_	_	_	-	-	_	_	_
Female	_	_	_	-	_	_	_	-	-	_	-	-
Non-Hispanic black, both sexes  Male	_	_	-	_	-	-	-	_	-	-	_	_
	_	_	-	_	-	_	-	_	-	-	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Origin not stated 2/, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	-	_	_	_	-	-	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Arthropod-borne viral encephalitis (A83-A84,A85.2) All origins, both sexes..... 5 1 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1 Female..... 3 Non-Hispanic white, both sexes..... Male.... 3 Female..... Non-Hispanic black, both sexes..... 1 Male..... 1 Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Measles (B05)												
All origins, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Hispanic, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female												
Non-Hispanic 1/, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic white, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	-	_	-	_	_	_
Non-Hispanic black, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	-	_	_	_	_	-
Female	_	_	_	-	_	_	_	_	_	_	_	_
Origin not stated 2/, both sexes	_	_	_	-	_	_	_	_	_	_	_	_
Male	_	_	_	-	_	_	_	_	_	_	_	_
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Viral hepatitis (B15-B19) All origins, both sexes..... 5,382 Male.... 3,513 Female..... 1.869 Hispanic, both sexes..... 2.4 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 4,647 3,031 Male.... 2.44 1,616 Female..... 1.3 Non-Hispanic white, both sexes..... 3,533 2,330 Male.... 1,203 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Human immunodeficiency virus (HIV) disease (B20-B24) All origins, both sexes..... 13.063 1,036 1,995 2,831 2.554 Male.... 9,506 1,411 2,055 1.911 Female..... 3.557 Hispanic, both sexes..... 1.758 Male.... 1,371 Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... 2.8 Central and South American, both sexes. 2.7 Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 11,195 1,667 2,455 2,204 Male.... 8.052 1,154 1,767 1.639 3,143 Female..... Non-Hispanic white, both sexes..... 3,862 2.2 3,227 Male.... Female..... Non-Hispanic black, both sexes..... 7,163 1,030 1,492 1,383 4,693 Male.... Female..... 2,470 Origin not stated 2/, both sexes..... Male.... Female..... 2.7 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malaria (B50-B54)	-5-2	_ 1	7	7	1	1	1	7	2	7	7	7
All origins, both sexes	8	_	_	_		_	1	_	_	1	3	1
Male	8	_	_	_	_	_	1	_	_	1	3	1
Female	-	_	_	_	_	_	_	_	_	_	_	_
Hispanic, both sexes	1	_	_	_	_	_	1	_	_	_	_	_
Male	1	_	_	_	_	_	1	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes								_				
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
·	_	_	_	_	_	_	_	-	-	_	_	_
Male	-	_	-	-	-	-	-	-	-	-	-	_
Female	_	-	_	_	_	-	_	_	-	_	_	-
Central and South American, both sexes.	1	-	_	_	_	-	1	-	-	_	_	-
Male	1	_	_	_	_	-	1	-	-	_	_	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	7	-	-	-	-	-	-	-	-	1	3	1
Male	7	-	-	-	-	-	-	-	-	1	3	1
Female	_	-	_	_	_	-	-	-	-	_	_	-
Non-Hispanic white, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	_	-	-	-	-	-	-	_	-	-	-	-
Non-Hispanic black, both sexes	3	_	_	_	_	-	-	-	-	_	2	1
Male	3	-	-	-	-	-	-	_	-	-	2	1
Female	_	_	-	-	-	_	-	_	_	-	-	-
Origin not stated 2/, both sexes	_	_	-	-	-	_	-	_	_	-	-	-
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	-	-	_	_	_	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) All origins, both sexes..... 7,540 Male.... 4,097 5.8 Female..... 3,443 1.0 Hispanic, both sexes..... 2.0 2.8 Male.... 5.8 Female..... Mexican, both sexes..... Male.... Female..... 1.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω4 a Male.... Female..... 6,797 Non-Hispanic 1/, both sexes..... 3,651 Male.... 2.4 1.0 2.7 3,146 Female..... Non-Hispanic white, both sexes..... 5,545 2,977 Male.... 2,568 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant neoplasms (C00-C97) All origins, both sexes..... 553.888 1,286 2,347 4,518 10,205 19,190 Male.... 1,071 1,909 4,499 9.395 Female..... 267.058 1,276 2,609 5,706 9.795 Hispanic, both sexes..... 24,522 1,253 Male.... 5.8 Female..... 11.520 Mexican, both sexes..... 13.163 Male.... 7.048 Female..... 6,115 2.8 Puerto Rican, both sexes..... 2,998 Male.... 1,597 Female..... 1,401 Cuban, both sexes..... 2,710 Male 1,573 Female..... 1,137 2,216 Central and South American, both sexes. Male 1,021 Female..... 1,195 Other and Unknown Hispanic, both sexes. 3,435 1.0 1 Q Male.... 1,763 Female..... 1,672 Non-Hispanic 1/, both sexes..... 528,395 1,087 2,011 3,988 9,364 17,906 273,282 Male.... 1,680 4,114 8.772 255,113 1,108 2,308 5,250 9,134 Female..... Non-Hispanic white, both sexes..... 453,534 1,475 2,994 7,068 13,736 234,392 3,150 6,763 Male.... 2.4 1,317 219,142 1,677 3,918 6,973 Female..... Non-Hispanic black, both sexes..... 61,842 1,868 3,466 32.089 1.669 Male.... Female..... 29,753 2.7 1,101 1,797 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) All origins, both sexes..... 7.826 Male.... 5,312 Female..... 2,514 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 3.8 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... 2.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 7,473 Non-Hispanic 1/, both sexes..... 5,055 Male.... 2,418 Female..... 2. Non-Hispanic white, both sexes..... 6,106 4,028 Male.... 2,078 Female..... Non-Hispanic black, both sexes..... 1,137 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of esophagus (C15)												
All origins, both sexes	13,023	_	_	_	_	_	5	11	25	69	222	513
Male	10,081	-	_	_	-	-	4	8	20	58	184	440
Female	2,942	_	-	-	-	-	1	3	5	11	38	73
Hispanic, both sexes	440	-	_	_	-	-	-	2	2	9	10	14
Male	346	-	_	_	-	_	-	1	2	7	6	14
Female	94	-	_	_	-	_	-	1	-	2	4	_
Mexican, both sexes	225	-	_	_	-	_	-	1	-	6	7	6
Male	182	-	_	_	-	_	-	-	-	5	4	6
Female	43	_	-	-	-	-	-	1	_	1	3	-
Puerto Rican, both sexes	93	-	-	-	_	-	-	1	1	1	-	2
Male	69	-	-	-	_	-	-	1	1	1	-	2
Female	24	-	-	-	_	-	-	-	-	-	-	-
Cuban, both sexes	38	_	_	_	_	_	-	_	_	_	_	2
Male	30	_	_	_	_	_	_	_	_	_	_	2
Female	8	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	24	_	_	_	_	_	-	_	1	_	2	2
Male	19	_	_	_	_	_	_	_	1	_	2	2
Female	5	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	60	_	_	_	_	_	_	_	_	2	1	2
Male	46	_	_	_	_	_	_	_	_	1	_	2
Female	14	_	_	_	_	_	_	_	_	1	1	_
Non-Hispanic 1/, both sexes	12,557	_	_	_	_	_	5	9	23	60	212	497
Male	9,715	_	_	_	_	_	4	7	18	51	178	424
Female	2,842	_	_	_	_	_	1	2	5	9	34	73
Non-Hispanic white, both sexes	10,809	_	_	_	_	_	3	8	16	53	174	382
Male	8,482	_	_	_	_	_	3	6	13	47	153	339
Female	2,327						_	2	3	- 6	21	43
Non-Hispanic black, both sexes	1,509			_	_		2	1	5	5	34	107
Male	1,047	_	_	_	_	_	1	1	4	3	23	79
Female	462	_	_	_	_	_	1	_	1	2	11	28
Origin not stated 2/, both sexes	26	_	_	_	_	_	1	_	1	2	11	20
Male	20	_	_	_	_	_	_	_	_	_	_	2
Female	20	_	_	_	_	_	_	_	_	_	_	_
r cma±c	0	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of stomach (C16)												
All origins, both sexes	11,859	_	_	_	_	2	10	29	93	145	333	457
Male	6,958	-	_	-	-	1	6	17	46	78	191	280
Female	4,901	-	_	-	-	1	4	12	47	67	142	177
Hispanic, both sexes	1,355	-	_	-	-	-	4	8	29	49	76	82
Male	771	-	_	-	-	-	4	4	18	28	36	44
Female	584	-	_	_	-	-	-	4	11	21	40	38
Mexican, both sexes	763	-	_	_	-	-	2	6	21	25	46	44
Male	434	-	_	_	-	-	2	3	12	15	22	22
Female	329	_	-	-	-	-	-	3	9	10	24	22
Puerto Rican, both sexes	148	-	-	-	-	-	-	-	1	3	8	11
Male	85	-	-	-	-	-	-	-	-	2	3	8
Female	63	-	-	-	-	-	-	-	1	1	5	3
Cuban, both sexes	85	_	_	-	-	-	-	1	2	1	2	1
Male	58	_	_	_	_	_	_	1	2	1	2	1
Female	27	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	164	_	_	_	_	_	2	1	5	12	10	14
Male	88	_	_	_	_	_	2	_	4	4	5	7
Female	76	_	_	_	_	_	_	1	1	8	5	7
Other and Unknown Hispanic, both sexes.	195	_	_	_	_	_	_	_	_	8	10	12
Male	106	_	_	_	_	_	_	_	_	6	4	6
Female	89	_	_	_	_	_	_	_	_	2	6	6
Non-Hispanic 1/, both sexes	10,481	_	_	_	_	2	6	21	63	96	256	374
Male	6,173	_	_	_	_	1	2	13	27	50	154	235
Female	4,308	_	_	_	_	1	4	8	36	46	102	139
Non-Hispanic white, both sexes	7,732	_	_	_	_	2	1	10	36	67	142	227
Male	4,575	_	_	_	_	1	_	5	20	36	92	148
Female	3,157	_	_	_	_	1	1	5	16	31	50	79
Non-Hispanic black, both sexes	2,004	_	_	_	_	_	4	8	18	14	79	105
Male	1,171	_	_	_	_	_	2	5	5	7	43	66
Female	833	_	_	_	_	_	2	3	13	7	36	39
Origin not stated 2/, both sexes	23	_	_	_	_	_	_	-	1	_	1	1
Male	14	_	_	_	_	_	_	_	1	_	1	1
Female	9	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years vears vears vears Malignant neoplasms of colon, rectum and anus (C18-C21) All origins, both sexes..... 53,772 1.637 Male.... 26,930 Female..... 26,842 2,438 Hispanic, both sexes..... 2.7 Male.... 1,327 Female..... 1.111 Mexican, both sexes..... 1.224 Male.... 2.0 Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 Male.... Female..... 51,252 Non-Hispanic 1/, both sexes..... 1,521 25,561 Male.... 2.3 25,691 Female..... Non-Hispanic white, both sexes..... 43,390 1,132 21,715 Male.... 21,675 Female..... Non-Hispanic black, both sexes..... 6.578 3.173 Male.... Female..... 3,405 Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Malignant neoplasms of liver and intrahepatic bile ducts (C22) All origins, both sexes..... 15,321 2.8 Male.... 9,832 2.0 Female..... 5.489 Hispanic, both sexes..... 1.565 Male.... 1,046 Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 13,718 Non-Hispanic 1/, both sexes..... 8,761 Male.... 1.5 2.8 4,957 Female..... Non-Hispanic white, both sexes..... 10,554 6,656 Male.... 3,898 Female..... Non-Hispanic black, both sexes..... 1,988 1,330 1.0 Male.... Female..... Origin not stated 2/, both sexes..... 3.8 Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
All origins, both sexes	31,772	_	_	_	_	1	10	11	51	125	396	926
Male	15,776	_	_	_	_	_	6	4	35	76	247	596
Female	15,996	_	_	_	_	1	4	7	16	49	149	330
Hispanic, both sexes	1,570	_	_	_	_	_	3	1	9	15	30	55
Male	789	_	_	_	_	_	1	_	5	11	20	37
Female	781	_	_	_	_	_	2	1	4	4	10	18
Mexican, both sexes	823	_	_	_	_	_	2	1	4	10	18	33
Male	413	_	_	_	_	_	1	_	2	7	13	21
Female	410	_	_	_	_	_	1	1	2	3	5	12
Puerto Rican, both sexes	176	_	_	_	_	_	_	_	3	1	4	10
Male	101	_	_	_	_	_	_	_	1	1	2	7
Female	75	_	_	_	_	_	_	_	2	_	2	3
Cuban, both sexes	165	_	_	_	_	_	_	_	_	_	_	2
Male	93						_					2
Female	72	_	_	_	_	_	_	_	_	_	_	2
Central and South American, both sexes.	152	_	_	_	_	_	- 1	-	2	3	- 5	_
•		_	-	-	-		_	-	2	3	5	6
Male	64	-	-	-	-	-	-	_	_	2	2	4
Female	88	-	-	_	_	-	1	-	-	1	3	2
Other and Unknown Hispanic, both sexes.	254	-	-	_	_	-	-	-	-	1	3	4
Male	118	-	-	-	-	-	-	-	-	1	3	3
Female	136	-	-	-	-	-	-	-	-	-	-	1
Non-Hispanic 1/, both sexes	30,148	-	-	-	-	1	7	10	42	110	366	871
Male	14,953	-	-	-	-	-	5	4	30	65	227	559
Female	15,195	-	-	-	-	1	2	6	12	45	139	312
Non-Hispanic white, both sexes	25,662	-	-	_	_	1	2	5	32	84	274	683
Male	12,814	-	-	-	-	-	1	3	23	53	166	446
Female	12,848	-	_	_	_	1	1	2	9	31	108	237
Non-Hispanic black, both sexes	3,631	-	-	-	-	-	5	4	6	16	75	160
Male	1,717	_	-	-	-	_	4	_	5	8	47	94
Female	1,914	-	-	-	-	-	1	4	1	8	28	66
Origin not stated 2/, both sexes	54	-	_	-	-	-	-	-	-	-	_	-
Male	34	_	-	-	-	-	-	-	-	-	_	-
Female	20	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Malignant neoplasm of larynx (C32) All origins, both sexes..... 3,668 1 1 3 52 161 Male.... 2,935 1 3 43 134 Female..... 733 1 27 Hispanic, both sexes..... 190 8 Male.... 8 Female..... 12 Mexican, both sexes..... 79 3 74 1 3 Male.... Female..... 5 Puerto Rican, both sexes..... 34 3 Male.... 33 3 - 1 Female.... Cuban, both sexes..... 31 Male 2.7 4 Female..... 14 1 Central and South American, both sexes. Male 14 1 Female..... Other and Unknown Hispanic, both sexes. 32 1 1 Male.... 30 1 1 Female..... 2 Non-Hispanic 1/, both sexes..... 3,463 3 50 153 2,745 Male.... 3 5 41 126 718 1 9 27 Female..... Non-Hispanic white, both sexes..... 2,793 3 35 119 2,192 Male.... 3 30 99 601 1 20 Female..... Non-Hispanic black, both sexes..... 625 2 15 33 518 11 27 Male.... 1 Female..... 107 6 Origin not stated 2/, both sexes...... 15 1 12 1 Male.... Female..... 3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Malignant neoplasms of trachea, bronchus and lung (C33-C34) 1,956 4.564 Male.... 89.630 1,044 2,566 Female..... 68,461 1.998 Hispanic, both sexes..... 4,215 Male.... 2,704 Female..... 1,511 Mexican, both sexes..... 2.133 1.0 Male.... 1.390 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female 153,581 Non-Hispanic 1/, both sexes..... 1,893 4,411 86,740 1,003 Male.... 2.4 2.472 66,841 1,939 Female..... 1.0 Non-Hispanic white, both sexes..... 134,118 1,467 3,451 74,960 Male.... 1.889 59,158 1,562 Female..... Non-Hispanic black, both sexes..... 16,324 9,906 Male.... Female..... 6,418 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant melanoma of skin (C43) All origins, both sexes..... 7,952 Male.... 5,125 Female..... 2,827 2.7 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 3.0 Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... 1.8 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 7,804 Non-Hispanic 1/, both sexes..... 5,038 Male.... 2,766 Female..... Non-Hispanic white, both sexes..... 7,654 4,967 Male.... 1.3 2,687 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Malignant neoplasm of breast (C50)												
All origins, both sexes	41,316	_	_	_	_	_	11	76	320	823	1,688	2,593
Male	362	_	_	_	_	_	1	1	-	3	7	13
Female	40,954	_	_	_	_	_	10	75	320	820	1,681	2,580
Hispanic, both sexes	1,926	_	_	_	_	_	1	12	36	90	135	166
Male	17	_	_	_	_	_	_		_	_		2
Female	1,909	_	_	_	_	_	1	12	36	9.0	135	164
Mexican, both sexes	978	_	_	_	_	_	1	4	24	59	71	80
Male	6	_	_	_	_	_	_	_		_		2
Female	972	_	_	_	_	_	1	4	2.4	59	71	78
Puerto Rican, both sexes	248	_	_	_	_	_	_	3	3	10	20	20
Male	4	_	_	_	_	_	_	_	_			
Female	244	_	_	_	_	_	_	3	3	10	20	20
Cuban, both sexes	202	_	_	_	_	_	_	_	_	3	2	15
Male	2	_	_	_	_	_	_	_	_	_	_	
Female	200	_	_	_	_	_	_	_	_	3	2	15
Central and South American, both sexes.	228	_	_	_	_	_	_	3	6	8	22	29
Male	1	_	_	_	_	_	_	_	_	_		
Female	227	_	_	_	_	_	_	3	6	8	22	29
Other and Unknown Hispanic, both sexes.	270	_	_	_	_	_	_	2	3	1.0	20	22
Male	4	_	_	_	_	_	_	_	_	_	_	_
Female	266	_	_	_	_	_	_	2	3	10	20	22
Non-Hispanic 1/, both sexes	39,329	_	_	_	_	_	10	64	283	732	1,550	2,423
Male	343	_	_	_	_	_	1	1	_	3	7	11
Female	38,986	_	_	_	_	_	9	63	283	729	1,543	2.412
Non-Hispanic white, both sexes	32,833	_	_	_	_	_	6	35	180	481	1,079	1,755
Male	282	_	_	_	_	_	_	-	_	1	5	9
Female	32,551	_	_	_	_	_	6	35	180	480	1,074	1,746
Non-Hispanic black, both sexes	5,546	_	_	_	_	_	2	26	89	222	397	567
Male	56	_	_	_	_	_	_	1	_	2	1	2
Female	5,490	_	_	_	_	_	2	25	89	220	396	565
Origin not stated 2/, both sexes	61	_	_	_	_	_	_	_	1	1	3	4
Male	2	_	_	_	_	_	_	_	_	_	_	_
Female	59	_	_	_	_	_	_	_	1	1	3	4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant neoplasm of cervix uteri (C53) All origins, both sexes..... 3,850 2 17 70 143 232 390 455 Male.... Female..... 3.850 2 17 70 143 232 390 455 Hispanic, both sexes..... 430 2 19 39 58 51 1 13 Male.... Female..... 430 1 2 13 19 39 58 51 Mexican, both sexes..... 286 1 1 12 16 29 35 34 Male.... Female..... 286 1 1 12 29 35 34 16 Puerto Rican, both sexes..... 60 1 7 9 1 8 Male.... 1 1 7 9 Female..... Cuban, both sexes..... 8 1 1 1 Male Female..... 1 Central and South American, both sexes. 39 1 6 Male . . . . . . . . . Female..... 39 Other and Unknown Hispanic, both sexes. 37 2 1 7 5 Male.... Female..... 37 5 Non-Hispanic 1/, both sexes..... 3,415 1 15 57 124 192 331 403 Male.... . . . . . . . . . . . . Female..... 3,415 57 192 331 403 1.5 124 1 Non-Hispanic white, both sexes..... 2,474 1 10 37 91 139 233 292 Male.... 2,474 37 91 139 233 292 Female..... 1.0 1 Non-Hispanic black, both sexes..... 782 4 18 25 44 81 95 Male.... Female..... 4 18 81 95 Origin not stated 2/, both sexes...... 5 1 1 1 Male.... Female..... 5

8

4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55) All origins, both sexes..... 6,990 2 3 6 14 51 117 206 Male.... Female..... 6.990 2 3 6 14 51 117 206 Hispanic, both sexes..... 325 1 2 11 19 6 Male.... Female..... 325 1 2 11 19 Mexican, both sexes..... 166 1 3 9 9 Male.... Female..... 166 9 1 3 Puerto Rican, both sexes..... 41 2 1 1 Male.... 41 1 2 Female..... Cuban, both sexes..... 37 1 Male 37 Female..... 39 Central and South American, both sexes. Male . . . . . . Female..... 39 Other and Unknown Hispanic, both sexes. 42 1 3 1 3 Male.... Female..... 42 3 3 2 Non-Hispanic 1/, both sexes..... 6,654 3 5 12 45 106 186 Male.... . . . . . . . . . . . . Female..... 6,654 3 45 106 186 5 12 Non-Hispanic white, both sexes..... 5,392 1 9 37 78 150 Male.... 5,392 2 9 37 150 Female..... 1 4 Non-Hispanic black, both sexes..... 1,102 2 1 3 7 22 26 Male.... Female..... 1,102 2 26 Origin not stated 2/, both sexes...... 11 1 Male....

Female.....

4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Malignant neoplasm of ovary (C56) All origins, both sexes..... 14,716 2 8 21 31 63 128 326 647 Male.... Female..... 14.716 2 8 21 31 63 128 326 647 Hispanic, both sexes..... 689 1 3 3 8 19 32 57 1 Male.... Female..... 689 1 1 3 3 8 19 32 57 Mexican, both sexes..... 393 2 2 3 15 17 37 Male.... Female..... 393 2 2 15 17 37 3 Puerto Rican, both sexes..... 73 1 5 1 1 Male.... 73 1 1 1 5 Female..... Cuban, both sexes..... 60 1 1 3 Male 60 Female..... 71 1 3 3 7 Central and South American, both sexes. 1 8 Male . . . Female..... 71 3 Other and Unknown Hispanic, both sexes. 92 5 2 5 Male.... Female..... 92 5 7 Non-Hispanic 1/, both sexes..... 14,008 1 18 28 55 108 294 590 Male.... . . . . . . . . . 14,008 28 108 294 590 Female..... 18 55 1 Non-Hispanic white, both sexes..... 12,432 5 11 17 482 Male.... 12,432 5 17 43 83 243 482 Female..... 11 1 Non-Hispanic black, both sexes..... 1,247 2 5 9 8 22 30 82 Male.... Female..... 1,247 5 30 Origin not stated 2/, both sexes...... 19 1 Male.... Female..... 19

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant neoplasm of prostate (C61) All origins, both sexes..... 29,004 1 1 1 1 1 1 22 102 Male.... 29,004 1 1 1 1 1 1 22 102 Female..... Hispanic, both sexes..... 1.239 6 Male.... 1,239 6 Female..... . . . Mexican, both sexes..... 651 3 Male.... 651 3 Female..... Puerto Rican, both sexes..... 140 Male.... 140 Female..... Cuban, both sexes..... 194 2 Male 194 2 Female..... 96 Central and South American, both sexes. 1 Male 96 1 Female..... . . . . . . . . . Other and Unknown Hispanic, both sexes. 158 Male.... 158 Female..... . . . . . . . . . . . . . . . . . . Non-Hispanic 1/, both sexes..... 27,723 1 1 1 22 96 Male.... 27,723 1 1 1 1 1 1 22 96 Female..... . . . Non-Hispanic white, both sexes..... 22,509 17 22,509 Male.... 1 1 1 1 1 17 60 Female..... Non-Hispanic black, both sexes..... 4,768 4 35 4,768 35 Male.... 4 Female..... . . . . . . . . . . . . . . . . . . Origin not stated 2/, both sexes...... 42 42 Male.... Female..... .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant neoplasms of kidney and renal pelvis (C64-C65) All origins, both sexes..... 12,313 2.2 2.7 Male.... 7.640 Female..... 4,673 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. QΩ Male.... Female..... Non-Hispanic 1/, both sexes..... 11,531 Male.... 7,121 4,410 Female..... Non-Hispanic white, both sexes..... 10,218 6,329 Male.... 3,889 Female..... Non-Hispanic black, both sexes..... 1,084 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of bladder (C67)												
All origins, both sexes	13,030	_	_	_	1	_	1	2	10	35	85	199
Male	8,995	-	_	_	1	-	1	1	6	17	52	141
Female	4,035	-	-	-	-	-	-	1	4	18	33	58
Hispanic, both sexes	397	-	_	_	-	-	-	-	1	3	3	9
Male	253	-	_	_	-	-	-	-	-	2	-	7
Female	144	-	_	_	-	-	-	-	1	1	3	2
Mexican, both sexes	157	-	_	_	-	-	-	-	1	3	1	5
Male	106	-	_	_	-	-	-	-	-	2	-	4
Female	51	_	-	-	_	_	-	_	1	1	1	1
Puerto Rican, both sexes	54	_	-	-	-	_	-	-	-	_	2	1
Male	26	_	-	-	-	_	-	-	-	_	_	1
Female	28	_	-	-	-	_	-	-	-	_	2	-
Cuban, both sexes	72	-	_	_	-	_	-	_	_	_	_	-
Male	57	_	_	_	_	_	_	_	_	_	_	_
Female	15	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	36	_	_	_	_	_	-	_	_	_	_	2
Male	1.8	_	_	_	_	_	_	_	_	_	_	1
Female	18	_	_	_	_	_	_	_	_	_	_	1
Other and Unknown Hispanic, both sexes.	78	_	_	_	_	_	_	_	_	_	_	1
Male	46	_	_	_	_	_	_	_	_	_	_	1
Female	32	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	12,608	_	_	_	1	_	1	2	8	32	82	190
Male	8,724	_	_	_	1	_	1	1	5	15	52	134
Female	3,884	_	_	_	_	_	_	1	3	17	30	56
Non-Hispanic white, both sexes	11,544	_	_	_	1	_	_	2	6	21	58	154
Male	8,140	_	_	_	1	_	_	1	0	9	37	111
Female	3,404	_	_	_	1	_	_	1	4	12	21	43
Non-Hispanic black, both sexes	904	_	_	_	_	_	1	_	2	9	18	31
Male	474	_	_	_	_	_	1	_	1	5	13	20
Female	474	_	_	_	-	-	1	_	1	4	13 5	20 11
	430 25	_	-	-	_	_	-	_	1	4	5	11
Origin not stated 2/, both sexes	25 18	_	-	-	_	_	-	_	1	_	_	_
Male	18	_	-	-	_	_	-	_	1	_	_	-
Female	/	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) All origins, both sexes..... 12.829 Male.... 7,074 Female..... 5,755 Hispanic, both sexes..... 2.4 2.4 3.8 Male.... Female..... 2.0 1.0 Mexican, both sexes..... 2.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... 12,137 Male.... 6.699 5,438 Female..... Non-Hispanic white, both sexes..... 11,121 6,139 Male.... 4,982 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) All origins, both sexes..... 54.645 1,477 Male.... 29,522 Female..... 25,123 Hispanic, both sexes..... 2,866 Male.... 1,619 Female..... 1.247 2.0 2.7 Mexican, both sexes..... 1.634 Male.... Female..... 2.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. 3.0 Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 51,695 1,338 27,865 Male.... 2.2.2 23,830 Female..... Non-Hispanic white, both sexes..... 45,383 1,051 24,604 Male.... 20,779 Female..... Non-Hispanic black, both sexes..... 5,163 2.7 2.608 Male.... Female..... 2,555 Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Hodgkin's disease (C81) All origins, both sexes..... 1,276 Male.... Female..... 1.0 Hispanic, both sexes..... Male.... Female..... 3.8 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,176 Male.... 3.0 2.8 Female..... Non-Hispanic white, both sexes..... 1,020 Male.... 2.1 2.8 2.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
All origins, both sexes	20,938	1	2	22	19	56	88	117	139	197	346	611
Male	11,056	_	2	15	13	37	56	83	81	129	238	405
Female	9,882	1	_	7	6	19	32	34	58	68	108	206
Hispanic, both sexes	1,062	-	_	6	3	11	14	25	22	24	51	61
Male	605	-	_	4	2	8	9	16	12	15	43	44
Female	457	_	_	2	1	3	5	9	10	9	8	17
Mexican, both sexes	593	_	_	3	3	6	10	16	16	15	26	28
Male	338	_	_	2	2	5	7	11	7	10	22	19
Female	255	_	_	1	1	1	3	5	9	5	4	9
Puerto Rican, both sexes	128	_	_	3	_	1	2	3	1	2	6	12
Male	75	_	_	2	_	_	1	2	1	1	6	8
Female	53	_	_	1	_	1	1	1	_	1	_	4
Cuban, both sexes	101	-	_	-	_	1	-	-	-	1	6	3
Male	65	-	_	-	_	1	-	-	-	-	5	3
Female	36	-	_	-	_	-	-	-	-	1	1	-
Central and South American, both sexes.	127	-	_	-	_	3	1	4	5	4	7	13
Male	65	_	_	_	_	2	1	2	4	2	5	9
Female	62	_	_	_	_	1	_	2	1	2	2	4
Other and Unknown Hispanic, both sexes.	113	_	_	_	_	_	1	2	_	2	6	5
Male	62	_	_	_	_	_	_	1	_	2	5	5
Female	51	-	_	-	_	-	1	1	-	-	1	_
Non-Hispanic 1/, both sexes	19,842	1	2	16	15	45	74	92	117	172	293	549
Male	10,437	_	2	11	11	29	47	67	69	114	195	360
Female	9,405	1	_	5	4	16	27	25	48	58	98	189
Non-Hispanic white, both sexes	17,970	1	2	12	8	35	50	59	80	133	229	427
Male	9,440	_	2	10	6	22	30	48	47	92	156	278
Female	8,530	1	_	2	2	13	20	11	33	41	73	149
Non-Hispanic black, both sexes	1,391	_	_	4	4	9	21	30	29	34	50	103
Male	728	_	_	1	2	6	14	17	19	19	31	68
Female	663	_	-	3	2	3	7	13	10	15	19	35
Origin not stated 2/, both sexes	34	_	-	_	1	_	-	_	_	1	2	1
Male	14	_	-	-	-	-	-	_	-	_	-	1
Female	20	-	-	-	1	-	-	-	-	1	2	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of												
Diseases, Tenth Revision, 1992),												
Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years	years	years	years	years	years	years	years	years	years
Leukemia (C91-C95)												
All origins, both sexes	21,395	23	120	156	156	210	225	173	243	287	426	555
Male	12,007	9	70	96	86	126	142	95	147	159	239	303
Female	9,388	14	50	60	70	84	83	78	96	128	187	252
Hispanic, both sexes	1,150	8	26	41	57	51	57	37	49	46	49	50
Male	669	3	12	29	33	35	42	21	33	24	33	24
Female	481	5	14	12	24	16	15	16	16	22	16	26
Mexican, both sexes	675	5	19	33	44	40	42	29	30	33	33	27
Male	401	1	9	24	26	27	33	17	20	16	21	14
Female	274	4	10	9	18	13	9	12	10	17	12	13
Puerto Rican, both sexes	107	1	2	4	2	4	4	2	6	1	5	4
Male	63	1	1	3	2	3	2	_	3	1	3	3
Female	44	_	1	1	_	1	2	2	3	_	2	1
Cuban, both sexes	96		_	_	_	_	1	_	2		_	1
Male	54	_	_	_	_	_	Τ.	_	1	_	_	1
	42	_	_	_	_	_	- 1	_	1	_	_	Τ.
Female  Central and South American, both sexes.	118	1	2	_	- 8	3	5	4	±	- 8	- 6	12
	64	1	∠ 1	_	3	3	3	2	5	8	0	3
Male		1	1	-	3 5	2	2	2	5	4	4 2	3
Female	54	-	1	_	5	_	_	_	_	4	2	9
Other and Unknown Hispanic, both sexes.	154	1	3	4	3	4	5	2	6	4	5	6
Male	87	_	1	2	2	4	4	2	4	3	5	3
Female	67	1	2	2	1	-	1	-	2	1	-	3
Non-Hispanic 1/, both sexes	20,214	15	94	115	97	158	167	136	194	241	377	505
Male	11,322	6	58	67	51	90	100	74	114	135	206	279
Female	8,892	9	36	48	46	68	67	62	80	106	171	226
Non-Hispanic white, both sexes	17,936	11	67	90	70	117	127	98	148	173	297	412
Male	10,098	5	39	52	38	69	74	57	88	98	162	230
Female	7,838	6	28	38	32	48	53	41	60	75	135	182
Non-Hispanic black, both sexes	1,831	3	18	14	23	34	27	30	40	49	64	75
Male	959	-	13	10	11	16	18	13	23	28	32	37
Female	872	3	5	4	12	18	9	17	17	21	32	38
Origin not stated 2/, both sexes	31	-	-	-	2	1	1	-	-	-	-	-
Male	16	-	-	-	2	1	-	-	-	-	-	-
Female	15	-	-	-	-	-	1	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Multiple myeloma and immunoproliferative neoplasms (C88,C90) All origins, both sexes..... 10,965 1 1 1 13 31 105 223 Male.... 5,694 1 1 1 10 16 62 120 Female..... 5,271 3 15 43 103 Hispanic, both sexes..... 558 11 17 4 8 Male.... 287 Female..... 271 3 1.0 Mexican, both sexes..... 318 2 3 10 12 Male.... 169 2 5 Female..... 149 1 3 Puerto Rican, both sexes..... 57 1 Male.... 36 1 Female.... Cuban, both sexes..... 54 Male 25 1 29 1 Female..... 56 5 Central and South American, both sexes. Male 30 Female..... 26 Other and Unknown Hispanic, both sexes. 73 2 1 Male.... 42 2 1 Female..... 31 Non-Hispanic 1/, both sexes..... 10,393 9 23 94 206 5,403 Male.... 1 1 1 6 12 54 113 4,990 3 11 40 93 Female..... Non-Hispanic white, both sexes..... 8,397 1 63 145 4,448 38 Male.... 1 1 1 4 8 82 3,949 2 6 25 63 Female..... Non-Hispanic black, both sexes..... 1,800 2 9 30 55 850 16 27 Male.... 1 Female..... 950 14 2.8 Origin not stated 2/, both sexes...... 14 4 Male....

Female.....

2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96) All origins, both sexes..... 71 2 1 1 1 Male.... 2 Female..... 36 Hispanic, both sexes..... 1 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male..... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 70 Non-Hispanic 1/, both sexes..... 34 Male.... 2 1 1 1 36 Female..... Non-Hispanic white, both sexes..... 31 Male.... 1 1 1 1 29 Female..... Non-Hispanic black, both sexes..... 8 Male..... Female..... Origin not stated 2/, both sexes..... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97) All origins, both sexes..... 61,911 1,227 2.117 Male.... 31,654 1,153 Female..... 30,257 Hispanic, both sexes..... 2,973 2.4 Male.... 1,521 5.8 Female..... 1.452 Mexican, both sexes..... 1.651 1.0 2.0 Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 2.0 Male.... Female..... Non-Hispanic 1/, both sexes..... 58,818 1,123 1,965 30,066 1,079 Male.... 28,752 Female..... Non-Hispanic white, both sexes..... 50,810 1,496 26,000 Male.... 24,810 Female..... Non-Hispanic black, both sexes..... 6.582 3,347 Male.... Female..... 3,235 2.0 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) All origins, both sexes..... 13.580 Male.... 6.764 2.0 Female..... 6,816 2.8 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 12,939 Non-Hispanic 1/, both sexes..... 6,440 Male.... 2.4 2.4 6,499 Female..... Non-Hispanic white, both sexes..... 11,469 5,754 Male.... 1.5 2.7 2.2 5,715 Female..... Non-Hispanic black, both sexes..... 1,181 

1.0

2.4 

Male....

Female.....

Male.... Female.....

Origin not stated 2/, both sexes.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
All origins, both sexes	4,575	15	33	25	21	50	78	84	80	84	117	120
Male	1,833	9	18	10	11	31	38	39	51	50	57	56
Female	2,742	6	15	15	10	19	40	45	29	34	60	64
Hispanic, both sexes	193	2	5	2	1	5	4	10	2	4	6	9
Male	75	1	3	1	1	-	1	3	2	1	3	6
Female	118	1	2	1	_	5	3	7	-	3	3	3
Mexican, both sexes	97	2	3	-	1	4	1	5	1	2	2	4
Male	36	1	1	-	1	-	1	2	1	_	1	3
Female	61	1	2	-	_	4	-	3	-	2	1	1
Puerto Rican, both sexes	27	-	-	-	_	-	-	1	-	_	1	2
Male	10	-	-	-	_	-	-	-	-	_	_	1
Female	17	-	-	-	_	-	-	1	-	_	1	1
Cuban, both sexes	26	_	_	_	_	_	2	_	_	_	_	_
Male	8	-	-	-	_	-	-	-	-	_	_	-
Female	18	_	_	_	_	_	2	_	_	_	_	_
Central and South American, both sexes.	18	_	1	1	_	_	1	_	_	2	2	1
Male	9	_	1	_	_	_	_	_	_	1	1	1
Female	9	_	_	1	_	_	1	_	_	1	1	_
Other and Unknown Hispanic, both sexes.	25	_	1	1	_	1	_	4	1	_	1	2
Male	12	_	1	1	_	_	_	1	1	_	1	1
Female	13	_	_	_	_	1	_	3	_	_	_	1
Non-Hispanic 1/, both sexes	4.371	13	28	23	20	45	74	74	78	80	110	111
Male	1,751	8	15	9	10	31	37	36	49	49	53	50
Female	2,620	5	1.3	14	1.0	14	37	38	29	31	57	61
Non-Hispanic white, both sexes	3,269	6	- 8	8	9	16	13	12	12	11	34	41
Male	1,247	3	3	4	2	11	7	4	10		18	22
Female	2,022	3	5	4	7	5	6	8	2	3	16	19
Non-Hispanic black, both sexes	1,013	4	19	12	11	29	61	60	66	67	72	66
Male	465	3	12	4	8	20	30	31	39	40	34	26
Female	548	1	7	8	3	9	31	29	27	27	38	40
Origin not stated 2/, both sexes	11	_	-	_	_	_	_				1	
Male	7	_	_	_	_	_	_	_	_	_	1	_
Female	4	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Diabetes mellitus (E10-E14) All origins, both sexes..... 73,138 1,317 2.217 Male.... 35,267 1,359 Female..... 37.871 Hispanic, both sexes..... 5.999 2.1 Male.... 2,855 2.6 Female..... 3.144 1.0 2.8 Mexican, both sexes..... 3.860 2.8 Male.... 1,862 Female..... 1.998 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... 66,953 Non-Hispanic 1/, both sexes..... 1,191 1,975 32,315 Male.... 1.222 34,638 Female..... Non-Hispanic white, both sexes..... 52,101 1,389 25,773 Male.... 26,328 Female..... Non-Hispanic black, both sexes..... 12,685 5,498 Male.... Female..... 7,187 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Nutritional deficiencies (E40-E64)												
All origins, both sexes	3,019	7	2	2	1	3	3	6	7	16	23	40
Male	1,087	5	1	1	1	2	3	3	3	8	7	16
Female	1,932	2	1	1	_	1	_	3	4	8	16	24
Hispanic, both sexes	107	1	_	_	1	_	_	1	_	_	2	1
Male	36	_	_	_	1	_	_	_	_	_	_	1
Female	71	1	_	_	_	_	_	1	_	_	2	_
Mexican, both sexes	72	1	_	_	_	_	_	_	_	_	2	1
Male	2.4	_	_	_	_	_	_	_	_	_	_	1
Female	48	1	_	_	_	_	_	_	_	_	2	_
Puerto Rican, both sexes	9	_	_	_	1	_	_	_	_	_	_	_
Male	4	_	_	_	1	_	_	_	_	_	_	_
Female	5	_	_	_	_	_	_	-	-	_	_	_
Cuban, both sexes	8	_	_	_	_	_	_	_	_	_	_	_
Male	3	_	_	_	_	_	_	_	_	_	_	_
Female	5	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	5	_	_	_	_	_	_	_	_	_	_	_
Male	1	_	_	_	_	_	_	_	_	_	_	_
Female	4	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	13	_	_	_	_	_	_	1	_	_	_	_
Male	4	_	_	_	_	_	_	_	_	_	_	_
Female	9	_	_	_	_	_	_	1	_	_	_	_
Non-Hispanic 1/, both sexes	2.906	6	2	2	_	3	3	5	7	16	21	39
Male	1,047	5	1	1	_	2	3	3	3	8	7	15
Female	1,859	1	1	1	_	1	_	2	4	8	14	24
Non-Hispanic white, both sexes	2,421	4	1	1	_	1	2	3	6	8	15	26
Male	847	3	_	_	_	1	2	2	3	3	4	7
Female	1,574	1	1	1	_	_	_	1	3	5	11	19
Non-Hispanic black, both sexes	412	1	1	1	_	1	1	2	1	7	5	12
Male	170	1	1	1	_	1	1	1	_	5	2	7
Female	242	_	_	_	_	_	_	1	1	2	3	5
Origin not stated 2/, both sexes	6	_	_	_	_	_	_	_	_	_	_	_
Male	4	_	_	_	_	_	_	_	_	_	_	_
Female	2	_	_	_	_	_	_	_	_	_	_	_
	_											

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Malnutrition (E40-E46) All origins, both sexes..... 2,847 6 1 1 1 2 3 6 5 14 21 36 Male.... 1,026 5 1 3 3 3 16 Female..... 1,821 1 3 2 15 2.0 Hispanic, both sexes..... 104 1 1 2 1 Male.... 36 Female..... 1 1 Mexican, both sexes..... 71 1 2 Male.... 1 Female..... 47 1 2 Puerto Rican, both sexes..... 9 1 Male.... 1 Female..... Cuban, both sexes..... 7 Male 3 4 Female..... Central and South American, both sexes. 5 Male Female..... Other and Unknown Hispanic, both sexes. 12 1 Male.... 4 Female..... 8 2,737 Non-Hispanic 1/, both sexes..... 5 14 19 35 Male.... 986 5 1 1 2 3 3 3 6 15 1,751 13 20 Female..... 2 2. Non-Hispanic white, both sexes..... 2,268 2 2.4 792 Male.... 3 1 2 2 3 3 1,476 11 17 Female..... 1 1 Non-Hispanic black, both sexes..... 398 1 1 1 1 1 2 1 6 4 10 164 1 1 7 Male.... 1 1 1 1 2 Female..... 234 3 Origin not stated 2/, both sexes...... 6 4 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other nutritional deficiencies (E50-E64) All origins, both sexes..... 172 1 1 1 1 2 2 4 Male.... Female..... 111 1 1 1 2 1 1 4 Hispanic, both sexes..... 3 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 Male.... Female..... Non-Hispanic 1/, both sexes..... 169 1 2 Male.... 61 1 1 108 1 1 1 1 2 Female..... 1 1 Non-Hispanic white, both sexes..... 153 1 2 Male..... 5.5 1 98 1 1 2 Female..... 1 Non-Hispanic black, both sexes..... 14 1 1 2 Male..... 6 1 Female..... 2 Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Meningitis (G00,G03) All origins, both sexes..... 2.0 Male.... Female..... 1.0 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.9 1.3 Female..... Non-Hispanic white, both sexes..... Male.... 2.2 1.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Parkinson's disease (G20-G21)												
All origins, both sexes	17,989	_	1	_	_	1	_	2	_	2	6	16
Male	10,366	_	_	_	_	1	_	2	_	2	4	12
Female	7,623	_	1	_	_	_	_	_	_	_	2	4
Hispanic, both sexes	594	_	_	_	_	_	_	1	_	_	_	1
Male	348	_	_	_	_	_	_	1	_	_	_	1
Female	246	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	290	_	_	_	_	_	_	_	_	_	_	_
Male	166	_	_	_	_	_	_	_	_	_	_	_
Female	124	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	51	_	_	_	_	_	_	_	_	_	_	_
Male	26	_	_	_	_	_	_	_	_	_	_	_
Female	25	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	114	_	_	_	_	_	_	_	_	_	_	_
Male	69	_	_	_	_	_	_	_	_	_	_	_
Female	45	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	27	_	_	_	_	_	_	_	_	_	_	_
Male	18	_	_	_	_	_	_	_	_	_	_	_
Female	9	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	112	_	_	_	_	_	_	1	_	_	_	1
Male	69	_	_	_	_	_	_	1	_	_	_	1
Female	43	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	17,368	_	1	_	_	1	_	1	_	2	6	15
Male	10,000	_	_	_	_	1	_	1	_	2	4	11
Female	7,368	_	1	_	_	_	_	_	_	_	2	4
Non-Hispanic white, both sexes	16,428	_	1	_	_	_	_	1	_	1	6	14
Male	9,469	_	_	_	_	_	_	1	_	1	4	11
Female	6,959	_	1	_	_	_	_	_	_	_	2	3
Non-Hispanic black, both sexes	623	_	_	_	_	_	_	_	_	1	_	_
Male	342	_	_	_	_	_	_	_	_	1	_	_
Female	281	_	_	_	_	_	_	_	_	_	_	_
Origin not stated 2/, both sexes	27	_	_	_	_	_	_	_	_	_	_	_
Male	18	_	_	_	_	_	_	_	_	_	_	_
Female	9	_	_	_	_	_	_	_	_	_	_	_
	_											

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alzheimer's disease (G30)												
All origins, both sexes	65,965	_	_	_	_	_	1	_	1	1	6	36
Male	18,974	-	_	_	-	-	-	-	1	1	3	8
Female	46,991	-	-	-	-	-	1	-	-	_	3	28
Hispanic, both sexes	1,897	_	-	-	-	-	1	_	-	_	_	3
Male	606	-	-	-	_	-	-	-	-	-	_	2
Female	1,291	-	-	-	_	-	1	-	-	-	_	1
Mexican, both sexes	839	_	_	_	_	_	1	_	_	_	_	2
Male	283	_	_	_	_	_	_	_	_	_	_	1
Female	556	_	_	_	_	_	1	_	_	_	_	1
Puerto Rican, both sexes	230	_	_	_	_	_	_	_	_	_	_	_
Male	74	_	_	_	_	_	-	_	_	_	_	_
Female	156	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	436	_	_	_	_	_	-	_	_	_	_	_
Male	126	_	_	_	_	_	-	_	_	_	_	_
Female	310	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	82	_	_	_	_	_	_	_	_	_	_	_
Male	23	_	_	_	_	_	_	_	_	_	_	_
Female	59	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	310	_	_	_	_	_	_	_	_	_	_	1
Male	100	_	_	_	_	_	_	_	_	_	_	1
Female	210	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	63,981	_	_	_	_	_	_	_	1	1	6	33
Male	18,345			_	_	_	_	_	1	1	3	6
Female	45,636	_	_	_	_	_	_	_	1	1	3	27
Non-Hispanic white, both sexes	59,153	_	_	_	_	_	_	_	- 1	- 1	3	33
Male	16,989	_	_	_	_	_	_	_	1	1	2	33 6
	42,164	_	_	_	_	_	_	_	1	Τ.	3	27
Female Non-Hispanic black, both sexes	42,164	_	_	_	_	_	_	-	-	_	1	27
Male	1,138	_	_	_	_	_	_	_	_	_	1	_
		_	_	_	_	_	_	-	-	_	1	_
Female	3,058	_	-	-	_	-	_	_	-	_	_	_
Origin not stated 2/, both sexes	87	_	_	_	_	_	_	_	_	-	_	-
Male	23	_	_	_	-	_	_	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Major cardiovascular diseases (I00-I78)												
All origins, both sexes	861,190	564	234	120	216	455	878	1,420	2,586	5,118	11,020	19,015
Male	406,458	307	120	53	117	298	569	916	1,686	3,398	7,408	13,126
Female	454,732	257	114	67	99	157	309	504	900	1,720	3,612	5,889
Hispanic, both sexes	36,900	104	43	20	26	58	121	159	265	464	840	1,218
Male	19,348	54	28	6	15	43	89	123	200	341	607	882
Female	17,552	50	15	14	11	15	32	36	65	123	233	336
Mexican, both sexes	19,639	70	34	12	16	45	84	119	173	300	493	709
Male	10,609	37	24	5	9	36	62	96	127	220	364	524
Female	9,030	33	10	7	7	9	22	23	46	80	129	185
Puerto Rican, both sexes	4,852	11	6	2	2	4	11	10	32	57	104	176
Male	2,485	5	3	_	2	3	8	5	25	39	74	119
Female	2,367	6	3	2	_	1	3	5	7	18	30	57
Cuban, both sexes	4,981	1	_	_	1	1	1	2	7	13	51	69
Male	2,505	_	_	_	1	_	1	1	5	12	45	52
Female	2,476	1	_	_	_	1	_	1	2	1	6	17
Central and South American, both sexes.	2,622	7	2	1	4	2	14	16	27	51	75	104
Male	1,268	2	_	_	2	1	1.0	12	19	41	53	74
Female	1,354	5	2	1	2	1	4	4	8	1.0	22	3.0
Other and Unknown Hispanic, both sexes.	4,806	15	1	- 5	3	6	11	12	26	43	117	160
Male	2,481	10	1	1	1	3	8	9	24	29	71	113
Female	2,325	5	_	4	2	3	3	3	2	14	46	47
Non-Hispanic 1/, both sexes	822,392	457	191	99	189	395	754	1,257	2,318	4,637	10,141	17,734
Male	386,042	252	92	46	101	253	479	790	1,483	3,044	6,778	12,201
Female	436,350	205	99	53	88	142	275	467	835	1,593	3,363	5,533
Non-Hispanic white, both sexes	705,435	277	109	62	120	243	478	730	1,427	3,095	7,018	12,418
Male	330,160	154	54	31	65	154	309	480	941	2,078	4,961	8,911
Female	375,275	123	55	31	55	89	169	250	486	1,017	2,057	3,507
Non-Hispanic black, both sexes	99,408	142	74	35	63	135	237	460	786	1,375	2,841	4,794
Male	46,787	80	37	15	32	91	151	267	458	853	1,614	2,916
Female	52,621	62	37	20	31	44	86	193	328	522	1,014	1,878
Origin not stated 2/, both sexes	1,898	3	3 /	20 1	1	2	3	193	328 3	17	39	1,878
	1,898	1	_	1	1	2	1	3	3	13	23	43
Male	830	2	_	1	1	_	2	3 1	3	13 4	23 16	20
Female	830	2	-	-	-	_	2	1	-	4	16	20

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Diseases of heart (I00-I09,I11,I13,I20-I51) 1,110 2,053 4,085 8,840 15,390 Male.... 321.973 1,408 2,800 6,211 11.078 Female..... 330.513 1,285 2,629 4,312 Hispanic, both sexes..... 27,788 Male.... 2.4 Female..... 12.805 Mexican, both sexes..... 14.389 1.0 Male.... 8,062 2.0 Female..... 6.327 1.0 2.7 Puerto Rican, both sexes..... 3,821 1.0 2.4 Male.... 1,984 Female..... 1,837 Cuban, both sexes..... 4,098 Male 2,067 Female..... 2,031 1.953 2.7 Central and South American, both sexes. 1.8 Male Female..... Other and Unknown Hispanic, both sexes. 3,527 a Male.... 1,916 Female..... 1,611 Non-Hispanic 1/, both sexes..... 623,165 1,859 3,745 8,197 14,447 306,083 2,543 Male.... 1,251 5,720 10.362 317,082 1,202 2,477 4,085 Female..... Non-Hispanic white, both sexes..... 537,512 1,165 2,567 5,844 10,463 263,638 1,789 Male.... 4,291 7.802 273,874 1,553 2,661 Female..... Non-Hispanic black, both sexes..... 73,373 1,053 2,149 3,600 35,749 1,273 2,269 Male.... Female..... 37.624 1,331 Origin not stated 2/, both sexes..... 1.533 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) All origins, both sexes..... 3,254 Male.... 1,009 Female..... 2,245 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 2.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 3,091 Male.... 2. 2,141 Female..... 2. Non-Hispanic white, both sexes..... 2,754 Male.... 2.7 1,914 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),												
Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years	years	years	years	years	years	years	years	years	years
Hypertensive heart disease (II1)												
All origins, both sexes	28,585	_	_	_	2	16	28	100	202	401	916	1,505
Male	12,929	_	_	_	_	14	18	71	152	296	600	1,007
Female	15,656	_	_	_	2	2	10	29	50	105	316	498
Hispanic, both sexes	1,491	_	_	_	1	1	3	9	15	38	67	88
Male	806	_	_	_	_	1	1	6	13	33	53	63
Female	685	_	_	_	1	_	2	3	2	5	14	25
Mexican, both sexes	761	_	_	_	1	1	3	7	8	23	34	47
Male	440	_	_	_	_	1	1	4	7	20	28	37
Female	321	_	_	_	1	_	2	3	1	3	6	10
Puerto Rican, both sexes	206	_	_	_	_	_	_	_	3	5	8	19
Male	106	_	_	_	_	_	_	_	2	5	7	10
Female	100	_	_	_	_	_	_	_	1	_	1	9
Cuban, both sexes	195							_	1	1	6	4
Male	99							_	1	1	6	3
Female	96	_	_	_	_		_	_	1	1	0	1
Central and South American, both sexes.	142	_	_	_	_	_	_	_ 1	2	_	9	<u></u>
		_	_	_	_	_	_	1	2	5	9	5
Male	66	_	_	_	_			1		4 1	3	5 1
Female	76	-	-	-	-	-	_	-	-	1	5	_
Other and Unknown Hispanic, both sexes.	187	_	_	_	_	-	-	1	1	4	10	12
Male	95	_	_	_	_	-	-	1	1	3	6	8
Female	92	_	_	_			_		=	1	4	4
Non-Hispanic 1/, both sexes	26,972	-	-	-	1	14	25	91	187	360	844	1,409
Male	12,051	-	-	-	-	12	17	65	139	261	545	940
Female	14,921	-	-	-	1	2	8	26	48	99	299	469
Non-Hispanic white, both sexes	19,630	-	-	-	-	7	12	33	88	186	466	786
Male	8,401	-	-	-	-	6	10	32	70	141	332	571
Female	11,229	-	-	-	-	1	2	1	18	45	134	215
Non-Hispanic black, both sexes	6,718	-	-	-	1	7	13	58	89	166	358	591
Male	3,336	-	-	-	-	6	7	33	59	113	199	346
Female	3,382	-	-	-	1	1	6	25	30	53	159	245
Origin not stated 2/, both sexes	122	-	-	-	-	1	-	-	-	3	5	8
Male	72	-	_	-	_	1	-	-	-	2	2	4
Female	50	-	_	-	_	-	-	-	-	1	3	4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years vears vears vears Hypertensive heart and renal disease (I13) All origins, both sexes..... 3,046 1 2 5 8 15 38 58 104 Male.... 1,301 4 29 60 Female..... 1.745 1 2 1 1 8 9 18 44 Hispanic, both sexes..... 190 1 7 1.0 1 3 5 Male.... 96 3 5 Female..... 94 2 5 Mexican, both sexes..... 117 1 1 3 4 3 5 57 1 1 3 3 3 Male.... Female..... 60 2 1 2 Puerto Rican, both sexes..... 2.0 2 4 Male.... 2 1.0 2 Female..... Cuban, both sexes..... 19 Male 11 8 Female..... 1.8 2 Central and South American, both sexes. Male 13 2 Female..... 5 Other and Unknown Hispanic, both sexes. 16 1 1 Male.... 5 Female..... 11 7 Non-Hispanic 1/, both sexes..... 2,845 4 12 33 50 93 Male.... 1.199 3 6 4 26 34 55 1,646 1 8 38 Female..... 1 1 16 Non-Hispanic white, both sexes..... 1,878 13 16 748 Male.... 1 2 1 3 7 12 1,130 1 4 Female..... 1 Non-Hispanic black, both sexes..... 877 2 3 5 10 29 36 74 404 2 3 23 26 41 Male.... 4 Female..... 473 33 Origin not stated 2/, both sexes...... 11 1 1 6 1 Male.... Female..... 5

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
All origins, both sexes	451,326	13	10	8	9	24	101	292	800	1,977	5,155	9,731
Male	233,538	8	1	4	5	20	76	206	616	1,467	3,899	7,442
Female	217,788	5	9	4	4	4	25	86	184	510	1,256	2,289
Hispanic, both sexes	20,482	4	2	2	2	2	13	20	75	138	339	562
Male	11,245	2	-	1	2	2	12	15	66	110	277	452
Female	9,237	2	2	1	-	_	1	5	9	28	62	110
Mexican, both sexes	10,481	3	1	2	2	2	10	14	46	84	198	339
Male	6,006	2	_	1	2	2	9	11	40	67	163	275
Female	4,475	1	1	1	-	_	1	3	6	17	35	64
Puerto Rican, both sexes	2,895	-	_	-	-	_	-	2	9	20	43	73
Male	1,513	-	_	-	-	_	-	1	8	15	33	55
Female	1,382	-	_	-	-	_	-	1	1	5	10	18
Cuban, both sexes	3,167	1	_	-	-	_	-	_	2	8	27	36
Male	1,591	-	_	-	-	_	-	_	1	7	26	30
Female	1,576	1	-	-	_	_	_	-	1	1	1	6
Central and South American, both sexes.	1,433	-	1	-	-	_	2	4	8	11	29	39
Male	708	-	-	-	_	-	2	3	7	9	24	33
Female	725	_	1	-	_	_	_	1	1	2	5	6
Other and Unknown Hispanic, both sexes.	2,506	-	-	-	_	-	1	-	10	15	42	75
Male	1,427	-	-	-	_	-	1	-	10	12	31	59
Female	1,079	-	-	-	_	-	-	-	-	3	11	16
Non-Hispanic 1/, both sexes	429,713	9	8	6	7	22	88	271	724	1,830	4,799	9,136
Male	221,607	6	1	3	3	18	64	190	549	1,349	3,611	6,965
Female	208,106	3	7	3	4	4	24	81	175	481	1,188	2,171
Non-Hispanic white, both sexes	374,900	6	5	3	4	12	45	179	511	1,416	3,712	7,088
Male	193,976	4	1	2	2	10	33	133	394	1,055	2,901	5,565
Female	180,924	2	4	1	2	2	12	46	117	361	811	1,523
Non-Hispanic black, both sexes	46,064	2	1	3	3	8	38	78	179	348	967	1,797
Male	22,720	2	_	1	1	7	29	44	125	238	610	1,197
Female	23,344	_	1	2	2	1	9	34	54	110	357	600
Origin not stated 2/, both sexes	1,131	_	_	_	_	_	_	1	1	9	17	33
Male	686	_	_	_	_	_	_	1	1	8	11	25
Female	445	-	-	-	-	-	-	-	-	1	6	8

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
All origins, both sexes	. 156,816	9	2	3	2	7	44	120	327	796	2,042	3,917
Male	. 82,909	7	-	2	2	5	34	80	242	598	1,515	2,989
Female	. 73,907	2	2	1	_	2	10	40	85	198	527	928
Hispanic, both sexes	. 7,132	2	1	1	1	1	4	7	28	51	127	205
Male	. 3,931	2	-	1	1	1	3	5	22	39	102	166
Female	. 3,201	-	1	_	-	-	1	2	6	12	25	39
Mexican, both sexes	. 3,976	2	1	1	1	1	4	4	20	36	77	132
Male	. 2,323	2	-	1	1	1	3	4	16	27	63	109
Female	. 1,653	_	1	_	_	-	1	-	4	9	14	23
Puerto Rican, both sexes	. 841	_	_	_	_	_	_	1	2	7	17	27
Male	. 435	_	_	_	_	_	_	_	1	5	12	19
Female	. 406	_	_	_	_	_	_	1	1	2	5	8
Cuban, both sexes		_	_	_	_	_	_	_	_	3	8	10
Male	. 533	_	-	_	_	-	_	-	_	2	8	9
Female	. 594	_	_	_	_	_	_	_	_	1	_	1
Central and South American, both sexes	. 396	_	_	_	_	_	_	2	3	1	12	11
Male	. 188	_	_	_	_	_	_	1	2	1	12	9
Female		_	_	_	_	_	_	1	1	_	_	2
Other and Unknown Hispanic, both sexes		_	_	_	_	_	_	_	3	4	13	25
Male		_	_	_	_	_	_	_	3	4	7	20
Female	. 340	_	_	_	_	_	_	_	_	_	6	5
Non-Hispanic 1/, both sexes		7	1	2	1	6	40	112	299	741	1,912	3,702
Male		5	_	1	1	4	31	74	220	555	1,410	2,814
Female		2	1	1	_	2	9	38	79	186	502	888
Non-Hispanic white, both sexes		4	1	_	1	2	20	75	228	583	1,506	2,950
Male		3	_	_	1	1	16	51	178	438	1,165	2,301
Female		1	1	_	_	1	4	24	50	145	341	649
Non-Hispanic black, both sexes		2	_	2	_	3	18	31	62	136	361	672
Male		2	_	1	_	3	14	17	36	100	209	452
Female		_	_	1	_	_	4	14	26	36	152	220
Origin not stated 2/, both sexes		_	_	_	_	_	_	1	-	4	3	10
Male												
	. 190	_	_	_	_	_	_	1	_	4	3	9

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other acute ischemic heart diseases (I24) All origins, both sexes..... 2,462 1 2 1 1 4 2 8 15 65 81 Male.... 1,301 1 3 5 13 46 55 Female..... 1,161 2 1 1 3 2 19 26 Hispanic, both sexes..... 35 1 1 2 1 1 1 Male.... Female..... 14 1 1 1 Mexican, both sexes..... 17 1 1 Male.... 13 1 Female..... 4 1 Puerto Rican, both sexes..... 6 Male.... 2 Female..... Cuban, both sexes..... 2 1 Male 1 1 Female..... 3 1 Central and South American, both sexes. Male Female..... 1 Other and Unknown Hispanic, both sexes. 1 Male.... 1 Female..... 4 Non-Hispanic 1/, both sexes..... 2,419 1 15 63 78 1,275 Male.... 1 3 1 4 13 44 53 1,144 1 3 2 19 25 Female..... 1 1 Non-Hispanic white, both sexes..... 1,988 2 12 46 51 Male..... 1,019 33 1 1 2. 11 36 969 13 15 Female..... 1 1 Non-Hispanic black, both sexes..... 389 1 2 5 2 16 25 232 1.0 15 Male.... 1 1 2 2 Female..... 157 10 Origin not stated 2/, both sexes...... 8 2 5 1 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other forms of chronic ischemic heart disease (I20,I25) 1.166 3,048 5.733 Male.... 149.328 2,338 4.398 Female..... 142.720 1.335 Hispanic, both sexes..... 13,315 Male.... Female..... 6,022 Mexican, both sexes..... 6.488 Male.... 3.670 Female..... 2,818 Puerto Rican, both sexes..... 2,048 Male.... 1.074 Female..... Cuban, both sexes..... 2.038 Male 1,057 Female..... 1.034 2.8 Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. 1 707 4 Q Male.... Female..... Non-Hispanic 1/, both sexes..... 277,926 1,074 2,824 5,356 141,544 Male.... 2,157 4.098 136,382 1,258 Female..... Non-Hispanic white, both sexes..... 242,493 2,160 4,087 123,514 Male.... 1,703 3,228 118,979 Female..... Non-Hispanic black, both sexes..... 29,640 1,100 14,767 Male.... Female..... 14,873 Origin not stated 2/, both sexes..... 2.1 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Atherosclerotic cardiovascular disease, so described (I25.0) All origins, both sexes..... 63,661 1 1 8 37 156 351 1.112 2.110 34,574 1 7 30 127 266 877 1,674 Male.... Female..... 29,087 1 1 7 29 85 235 436 Hispanic, both sexes..... 3,277 3 1 16 2.0 82 134 Male.... 2,098 3 15 17 114 Female..... 1,179 1 3 11 2.0 Mexican, both sexes..... 1.747 2 5 7 45 78 Male.... 1,137 2 5 40 69 6 Female..... 610 5 9 1 Puerto Rican, both sexes..... 444 1 3 16 6 Male.... 1 3 4 13 171 2 2 Female..... Cuban, both sexes..... 278 1 6 1 Male 172 1 6 106 1 Female..... 263 Central and South American, both sexes. 4 5 14 Male 168 4 5 6 12 Female..... 95 Other and Unknown Hispanic, both sexes. 545 1 2 1 15 1 Q Male.... 348 1 3 1 12 15 Female..... 197 Non-Hispanic 1/, both sexes..... 60,067 1 5 36 140 329 1,025 1,966 32,271 Male.... 1 4 29 112 248 802 1.553 27,796 1 7 28 81 223 413 Female..... 1 Non-Hispanic white, both sexes..... 49,197 3 26 89 258 778 1,446 26,188 193 Male.... 1 2 2.2 69 633 1.193 23,009 20 65 145 253 Female..... 1 4 Non-Hispanic black, both sexes..... 9,497 1 2 8 41 53 218 464 5.231 2 34 38 144 309 Male.... 5 Female..... 4,266 74 155 Origin not stated 2/, both sexes..... 317 2 5 1.0 205 4 Male.... 1

Female.....

112

3

7

5

2

2

4

7

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) 1,936 3.623 Male.... 114.754 1,461 2.724 Female..... 113.633 Hispanic, both sexes..... 10,038 Male.... 2.8 Female..... 4,843 Mexican, both sexes..... 4.741 2.0 Male.... 2.533 Female..... 2,208 Puerto Rican, both sexes..... 1,604 Male.... Female..... Cuban, both sexes..... 1.760 Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. 1 162 1.0 3 U Male.... Female..... Non-Hispanic 1/, both sexes..... 217,859 1,799 3,390 109,273 Male.... 1,355 2.545 108,586 Female..... Non-Hispanic white, both sexes....... 193,296 1,382 2,641 97,326 1,070 2,035 Male.... 95,970 Female..... Non-Hispanic black, both sexes..... 20,143 9,536 Male.... Female..... 10,607 Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

All origins, both sexes	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Male         73,196         218         90         28         83         204         351         447         628         994         1,648         2,530           Female         93,079         188         84         45         65         117         183         249         395         643         999         1,481         226           Male         2,780         39         23         3         8         29         54         68         71         97         134         158         226           Male         2,687         38         13         10         7         10         15         12         26         41         54         68           Male         1,536         24         19         3         3         23         37         51         44         70         80         91           Pemale         1,536         24         19         3         3         23         37         51         44         70         80         91           Pemale         1,641         25         4         4         6         8         6         20         27         33         3         9	Other heart diseases (I26-I51)												
Remale	All origins, both sexes	166,275	406	174	73	148	321	534	696	1,023	1,637	2,647	3,965
Hispanic, both sexes. 5,467 77 36 13 15 39 69 80 97 138 188 226 Male. 2,780 39 23 3 8 29 54 68 71 197 134 158 Female. 2,780 39 23 3 8 29 54 68 71 197 134 158 Female. 2,687 38 13 10 7 10 15 12 26 41 54 68 Mexican, both sexes. 2,947 49 28 7 7 29 45 57 64 97 113 130 Male. 1,536 24 19 3 3 3 23 37 51 44 70 80 91 Female. 1,1411 25 9 4 4 6 8 6 6 20 27 33 39 Female. 1,1411 25 9 4 4 4 6 8 6 6 20 27 33 39 Female. 333 6 6 3 1 - 1 3 3 2 2 5 5 6 8 8 Cuban, both sexes. 660 10 6 1 2 4 10 6 12 14 29 31 Male. 333 6 6 3 1 - 1 1 3 2 2 2 5 6 6 8 8 Cuban, both sexes. 706 1 1 1 1 1 1 2 2 2 4 10 9 9 23 23 8 8 25 Female. 347 1 1 1 1 1 1 2 2 2 4 4 10 9 9 23 23 23 8 8 29 54 68 8 7 16 Female. 343 3 6 6 3 1 - 1 1 3 1 2 2 2 4 1 10 9 8 23 23 24 2 5 6 6 8 8 8 6 8 7 1 10 9 9 23 23 23 24 1 10 9 9 23 23 23 1 10 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Male	73,196	218	90	28	83	204	351	447	628	994	1,648	2,530
Male         2 780         39         23         3         8         29         54         68         71         97         134         158           Female         2 687         38         13         10         7         10         15         12         26         41         54         68           Mexican, both sexes         2 947         49         28         7         7         29         45         57         64         97         113         130           Male         1,536         24         19         3         3         23         37         51         44         70         80         91           Female         1,111         25         9         4         4         6         8         6         20         27         33         39           Puerto Rican, both sexes         680         10         6         1         2         4         10         6         12         14         29         31           Male         347         4         3         -         -         -         1         1         1         2         2         4         8         5	Female	93,079	188	84	45	65	117	183	249	395	643	999	1,435
Pemale	Hispanic, both sexes	5,467	77	36	13	15	39	69	80	97	138	188	226
Mexican, both sexes.         2,947         49         28         7         7         29         45         57         64         97         113         130           Male.         1,536         24         19         3         3         23         37         51         44         70         80         91           Female.         1,411         25         9         4         4         6         8         6         20         27         33         39           Puerto Rican, both sexes.         680         10         6         1         2         4         10         6         12         14         29         31           Male.         3333         6         3         1         -         1         3         2         2         5         6         8           Cuban, both sexes.         706         -         -         -         1         1         1         2         2         4         10         9           Male.         3359         -         -         -         1         -         1         1         1         2         2         4         10         9	Male	2,780	39	23	3	8	29	54	68	71	97	134	158
Male         1,536         24         19         3         3         23         37         51         44         70         80         91           Female         1,411         25         9         4         4         6         8         6         20         27         33         39           Puerto Rican, both sexes         680         10         6         1         2         4         10         6         12         14         29         31           Male         347         4         3         -         2         3         7         4         10         9         23         23           Cuban, both sexes         706         -         -         -         1         1         1         2         2         4         10         9           Male         359         -         -         -         1         1         1         2         2         4         10         9           Male         359         -         -         -         1         1         1         2         2         4         10         9           Guban, both sexes         349         4 <td>Female</td> <td>2,687</td> <td>38</td> <td>13</td> <td>10</td> <td>7</td> <td>10</td> <td>15</td> <td>12</td> <td>26</td> <td>41</td> <td>54</td> <td>68</td>	Female	2,687	38	13	10	7	10	15	12	26	41	54	68
Female         1,411         25         9         4         4         6         8         6         20         27         33         39           Puerto Rican, both sexes         680         10         6         1         2         4         10         6         12         14         29         31           Male         333         6         3         1         -         1         3         2         2         5         6         8           Cuban, both sexes         706         -         -         1         1         1         2         2         5         6         8         5         6         8         6         3         1         -         1         1         1         2         2         4         10         9           Male         3559         -         -         -         1         -         1         1         1         2         4         8         5           Female         347         4         1         1         3         -         6         5         8         7         16           Female         1         1         4         <	Mexican, both sexes	2,947	49	28	7	7	29	45	57	64	97	113	130
Puerto Rican, both sexes. 680 10 6 1 2 4 10 6 12 14 29 31 Male. 347 4 3 - 2 3 7 4 10 9 23 23 Female. 333 6 3 1 - 1 1 3 2 2 5 5 6 8 8 Cuban, both sexes. 706 1 1 1 1 1 2 2 2 4 10 9 9 23 23 8 8 5 Female. 347 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Male	1,536	24	19	3	3	23	37	51	44	70	80	91
Male       347       4       3       -       2       3       7       4       10       9       23       23         Female       333       6       3       1       -       1       3       2       2       5       6       8         Cuban, both sexes       706       -       -       -       1       1       1       2       2       4       10       9         Male       359       -       -       -       1       -       1       1       1       2       4       8       5         Female       347       -       -       -       -       1       -       1       -       -       2       4       4       8       5         Female       160       163       2       -       -       1       -       5       6       5       8       7       16         Female       160       163       2       -       -       1       -       5       6       5       8       7       16         Female       10       160       5       1       1       1       2       -       1	Female	1,411	25	9	4	4	6	8	6	20	27	33	39
Female	Puerto Rican, both sexes	680	10	6	1	2	4	10	6	12	14	29	31
Cuban, both sexes	Male	347	4	3	-	2	3	7	4	10	9	23	23
Male	Female	333	6	3	1	-	1	3	2	2	5	6	8
Female	Cuban, both sexes	706	-	-	-	1	1	1	2	2	4	10	9
Central and South American, both sexes. 349 4 1 1 3 3 - 6 7 8 10 12 23 Male	Male	359	-	-	-	1	-	1	1	2	4	8	5
Male.       163       2       -       -       1       -       5       6       5       8       7       16         Female.       186       2       1       1       2       -       1       1       3       2       5       7         Other and Unknown Hispanic, both sexes.       785       14       1       4       2       5       7       8       11       13       24       33         Male.       375       9       1       -       1       3       4       6       10       6       16       23         Female.       410       5       -       4       1       2       3       2       1       7       8       10         Non-Hispanic 1/, both sexes.       160,544       327       138       59       132       281       462       614       924       1,496       2,449       3,728         Male.       70,276       178       67       24       74       174       296       377       555       895       1,508       2,365         Female       90,268       149       71       35       58       107       166       237	Female	347	_	_	_	_	1	_	1	_	_	2	4
Male.       163       2       -       -       1       -       5       6       5       8       7       16         Female.       186       2       1       1       2       -       1       1       3       2       5       7         Other and Unknown Hispanic, both sexes.       785       14       1       4       2       5       7       8       11       13       24       33         Male.       375       9       1       -       1       3       4       6       10       6       16       23         Female.       410       5       -       4       1       2       3       2       1       7       8       10         Non-Hispanic 1/, both sexes.       160,544       327       138       59       132       281       462       614       924       1,496       2,449       3,728         Male.       70,276       178       67       24       74       174       296       377       555       895       1,508       2,365         Female       90,268       149       71       35       58       107       166       237	Central and South American, both sexes.	349	4	1	1	3	_	6	7	8	10	12	23
Other and Unknown Hispanic, both sexes. 785 14 1 4 2 5 7 8 11 13 24 33 Male		163	2	_	_	1	_	5	6	5	8	7	16
Other and Unknown Hispanic, both sexes. 785 14 1 4 2 5 7 8 11 13 24 33 Male	Female	186	2	1	1	2	_	1	1	3	2	5	7
Male.         375         9         1         -         1         3         4         6         10         6         16         23           Female.         410         5         -         4         1         2         3         2         1         7         8         10           Non-Hispanic 1/, both sexes.         160,544         327         138         59         132         281         462         614         924         1,496         2,449         3,728           Male.         .         70,276         178         67         24         74         174         296         377         555         895         1,508         2,365           Female.         90,268         149         71         35         58         107         166         237         369         601         941         1,363           Non-Hispanic white, both sexes.         138,350         202         74         34         82         171         302         358         558         948         1,624         2,519           Male.         .         78,673         110         35         15         48         104         197         218         <		785	14	1	4	2	5	7	8	11	13	24	33
Non-Hispanic 1/, both sexes         160,544         327         138         59         132         281         462         614         924         1,496         2,449         3,728           Male         70,276         178         67         24         74         174         296         377         555         895         1,508         2,365           Female         90,268         149         71         35         58         107         166         237         369         601         941         1,363           Non-Hispanic white, both sexes.         138,350         202         74         34         82         171         302         358         558         948         1,624         2,519           Male         59,673         110         35         15         48         104         197         218         343         584         1,624         2,519           Female         78,677         92         39         19         34         67         105         140         215         364         586         892           Non-Hispanic black, both sexes.         19,469         97         59         23         46         97         139<		375	9	1	_	1	3	4	6	10	6	16	23
Non-Hispanic 1/, both sexes.         160,544         327         138         59         132         281         462         614         924         1,496         2,449         3,728           Male.         70,276         178         67         24         74         174         296         377         555         895         1,508         2,365           Female.         90,268         149         71         35         58         107         166         237         369         601         941         1,363           Non-Hispanic white, both sexes.         138,350         202         74         34         82         171         302         358         558         948         1,624         2,519           Male.         59,673         110         35         15         48         104         197         218         343         584         1,038         1,627           Female.         78,677         92         39         19         34         67         105         140         215         364         586         892           Non-Hispanic black, both sexes.         19,469         97         59         23         46         97         139	Female	410	5	-	4	1	2	3	2	1	7	8	10
Male.       70,276       178       67       24       74       174       296       377       555       895       1,508       2,365         Female.       90,268       149       71       35       58       107       166       237       369       601       941       1,363         Non-Hispanic white, both sexes.       138,350       202       74       34       82       171       302       358       558       948       1,624       2,519         Male.       59,673       110       35       15       48       104       197       218       343       584       1,038       1,627         Female.       78,677       92       39       19       34       67       105       140       215       364       586       892         Non-Hispanic black, both sexes.       19,469       97       59       23       46       97       139       219       327       499       766       1,112         Male.       9,206       53       31       9       23       64       86       136       180       283       430       676         Female       10,263       44       28		160,544	327	138	59	132	281	462	614	924	1,496	2,449	3,728
Non-Hispanic white, both sexes. 138,350 202 74 34 82 171 302 358 558 948 1,624 2,519 Male		70,276	178	67	24	74	174	296	377	555	895	1,508	2,365
Non-Hispanic white, both sexes. 138,350 202 74 34 82 171 302 358 558 948 1,624 2,519 Male	Female	90,268	149	71	35	58	107	166	237	369	601		
Male.     59,673     110     35     15     48     104     197     218     343     584     1,038     1,627       Female.     78,677     92     39     19     34     67     105     140     215     364     586     892       Non-Hispanic black, both sexes     19,469     97     59     23     46     97     139     219     327     499     766     1,112       Male.     9,206     53     31     9     23     64     86     136     180     283     430     676       Female.     10,263     44     28     14     23     33     53     83     147     216     336     436       Origin not stated 2/, both sexes     264     2     -     1     1     1     3     2     2     3     10     11       Male.     140     1     -     1     1     1     1     2     2     2     6     7			202	74	34	82	171	302	358	558	948	1,624	
Female     78,677     92     39     19     34     67     105     140     215     364     586     892       Non-Hispanic black, both sexes     19,469     97     59     23     46     97     139     219     327     499     766     1,112       Male     9,206     53     31     9     23     64     86     136     180     283     430     676       Female     10,263     44     28     14     23     33     53     83     147     216     336     436       Origin not stated 2/, both sexes     264     2     -     1     1     1     3     2     2     2     3     10     11       Male     140     1     -     1     1     1     1     2     2     2     2     6     7			110	35	15	48	104	197	218	343	584	1,038	
Non-Hispanic black, both sexes		78,677	92	39	19	34	67	105	140	215	364	586	
Female     10,263     44     28     14     23     33     53     83     147     216     336     436       Origin not stated 2/, both sexes     264     2     -     1     1     1     3     2     2     3     10     11       Male     140     1     -     1     1     1     1     2     2     2     2     6     7		19,469	97	59	23	46	97	139	219	327	499	766	1,112
Origin not stated 2/, both sexes 264 2 - 1 1 1 3 2 2 3 10 11 Male	Male	9,206	53	31	9	23	64	86	136	180	283	430	676
$oxdet{ exttt{Male}}$	Female	10,263	44	28	14	23	33	53	83	147	216	336	436
$oxdet{ exttt{Male}}$	Origin not stated 2/, both sexes		2	_	1	1	1	3	2	2	3	10	11
Female		140	1	_	1	1	1	1	2	2	2	6	7
	Female	124	1	_	_	_	_	2	-	-	1	4	4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Acute and subacute endocarditis (I33) All origins, both sexes..... 1,196 2.7 Male.... 2.4 Female..... Hispanic, both sexes..... Male.... Female..... 3.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,097 Male.... 1.5 Female..... Non-Hispanic white, both sexes..... Male.... 2.7 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Diseases of pericardium and acute myocarditis (I30-I31,I40) All origins, both sexes..... 1.0 Male.... 2.0 Female..... 1.0 1.0 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1.0 2.4 Female..... Non-Hispanic white, both sexes..... Male.... R Female..... 2. Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Heart failure (I50)												
All origins, both sexes	57,120	30	11	3	7	12	21	38	52	110	203	386
Male	22,292	16	4	1	6	6	13	28	31	69	117	237
Female	34,828	14	7	2	1	6	8	10	21	41	86	149
Hispanic, both sexes	1,545	5	5	_	_	2	1	3	2	9	13	23
Male	668	4	3	_	_	1	1	2	1	5	5	16
Female	877	1	2	_	_	1	_	1	1	4	8	7
Mexican, both sexes	905	4	3	_	_	1	_	1	1	8	10	15
Male	392	3	2	_	_	1	_	_	_	4	4	11
Female	513	1	1	_	_	_	_	1	1	4	6	4
Puerto Rican, both sexes	171	_	1	_	_	_	1	_	_	_	2	4
Male	73	_	_	_	_	_	1	_	_	_	1	3
Female	98	_	1	_	_	_	_	_	_	_	1	1
Cuban, both sexes	149	_	_	_	_	1	_	_	_	1	_	_
Male	69	-	-	_	_	_	-	-	-	1	_	-
Female	80	_	-	_	_	1	-	-	-	_	_	-
Central and South American, both sexes.	65	-	-	_	_	_	-	-	-	_	_	-
Male	17	-	-	_	_	_	-	-	-	_	_	-
Female	48	-	-	_	_	_	-	-	-	_	_	-
Other and Unknown Hispanic, both sexes.	255	1	1	_	_	_	-	2	1	_	1	4
Male	117	1	1	_	_	_	-	2	1	_	_	2
Female	138	-	-	_	_	_	-	-	-	_	1	2
Non-Hispanic 1/, both sexes	55,488	25	6	3	7	10	20	35	50	101	190	362
Male	21,580	12	1	1	6	5	12	26	30	64	112	221
Female	33,908	13	5	2	1	5	8	9	20	37	78	141
Non-Hispanic white, both sexes	49,628	15	5	_	5	5	13	19	27	53	118	223
Male	19,125	6	1	_	4	3	8	15	13	39	71	132
Female	30,503	9	4	_	1	2	5	4	14	14	47	91
Non-Hispanic black, both sexes	5,198	8	1	2	2	5	6	15	21	46	65	135
Male	2,154	5	-	1	2	2	4	10	15	23	36	86
Female	3,044	3	1	1	-	3	2	5	6	23	29	49
Origin not stated 2/, both sexes	87	-	-	-	-	-	-	-	-	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	_
Female	43	-	-	-	-	-	-	-	-	-	-	1

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51) 1,461 2,359 3.440 Male.... 1,480 2,208 Female..... 57,339 1.232 Hispanic, both sexes..... 3,760 Male.... 2,006 Female..... 1.754 1.0 1.0 5.8 Mexican, both sexes..... 1.944 2.0 Male.... 1.073 2.0 5.8 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1.0 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... 103,244 1,343 2,185 3,242 47,739 Male.... 1,352 2.071 55,505 1,171 Female..... Non-Hispanic white, both sexes..... 87,330 1,459 2,222 39,812 1,455 Male.... 47,518 Female..... Non-Hispanic black, both sexes..... 13,906 6.860 Male.... Female..... 7.046 Origin not stated 2/, both sexes..... 1.0 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Essential (primary) hypertension and hypertensive renal disease (I10,I12) All origins, both sexes..... 23,076 Male.... 8,869 Female..... 14,207 Hispanic, both sexes..... 1,116 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 21,905 Non-Hispanic 1/, both sexes..... 8,353 Male.... 2.0 13,552 Female..... Non-Hispanic white, both sexes..... 16,569 6,147 Male.... 10,422 Female..... Non-Hispanic black, both sexes..... 4.709 1,945 Male.... Female..... 2,764 2.7 Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
All origins, both sexes	150,074	127	41	34	43	69	142	208	359	719	1,642	2,622
Male	58,800	71	23	18	22	41	75	112	172	389	851	1,394
Female	91,274	56	18	16	21	28	67	96	187	330	791	1,228
Hispanic, both sexes	6,781	20	3	4	6	11	25	37	56	106	198	260
Male	3,207	11	3	1	4	6	14	27	34	75	115	159
Female	3,574	9	-	3	2	5	11	10	22	31	83	101
Mexican, both sexes	3,999	15	3	3	5	8	20	31	40	67	117	157
Male	1,938	9	3	1	3	5	11	24	25	47	73	100
Female	2,061	6	-	2	2	3	9	7	15	20	44	57
Puerto Rican, both sexes	737	1	-	1	-	-	-	1	7	15	20	33
Male	354	1	-	-	-	-	-	-	4	9	8	18
Female	383	_	-	1	-	-	-	1	3	6	12	15
Cuban, both sexes	627	_	-	-	-	-	-	-	2	-	7	14
Male	300	_	-	-	-	-	-	-	1	-	5	11
Female	327	_	-	-	-	-	-	-	1	-	2	3
Central and South American, both sexes.	489	3	-	-	1	2	4	3	6	19	22	26
Male	228	-	-	-	1	1	2	2	3	15	13	14
Female	261	3	-	-	-	1	2	1	3	4	9	12
Other and Unknown Hispanic, both sexes.	929	1	-	-	-	1	1	2	1	5	32	30
Male	387	1	-	-	-	-	1	1	1	4	16	16
Female	542	-	-	-	-	1	-	1	-	1	16	14
Non-Hispanic 1/, both sexes	143,044	106	38	30	37	58	117	170	303	612	1,439	2,355
Male	55,489	60	20	17	18	35	61	85	138	314	734	1,231
Female	87,555	46	18	13	19	23	56	85	165	298	705	1,124
Non-Hispanic white, both sexes	121,065	59	25	22	26	39	78	102	178	388	868	1,400
Male	46,087	34	15	13	11	21	42	58	79	190	463	726
Female	74,978	25	10	9	15	18	36	44	99	198	405	674
Non-Hispanic black, both sexes	17,980	39	12	8	9	18	30	60	113	195	503	845
Male	7,586	23	5	4	6	13	17	23	54	112	232	442
Female	10,394	16	7	4	3	5	13	37	59	83	271	403
Origin not stated 2/, both sexes	249	1	-	_	_	-	-	1	-	1	5	7
Male	104	-	-	_	_	-	-	-	-	-	2	4
Female	145	1	-	-	-	-	-	1	-	1	3	3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
All origins, both sexes	11,861	_	1	_	_	_	2	_	6	12	31	60
Male	4,510	_	_	_	_	_	2	_	3	7	18	42
Female	7,351	_	1	_	_	_	_	_	3	5	13	18
Hispanic, both sexes	369	_	_	_	_	_	_	_	_	2	1	_
Male	172	_	_	_	_	_	_	_	_	1	1	_
Female	197	_	_	_	_	_	_	_	_	1	_	_
Mexican, both sexes	220	_	_	_	_	_	_	_	_	1	1	_
Male	108	_	_	_	_	_	_	_	_	_	1	_
Female	112	_	_	_	_	_	_	_	_	1	_	_
Puerto Rican, both sexes	29	_	_	_	_	_	_	_	_	1	_	_
Male	16	_	_	_	_	_	_	_	_	1	_	_
Female	13	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	43	_	_	_	_	_	_	_	_	_	_	_
Male	1.5	_	_	_	_	_	_	_	_	_	_	_
Female	28	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	14	_	_	_	_	_	_	_	_	_	_	_
Male	- 3	_	_	_	_	_	_	_	_	_	_	_
Female	11	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	63	_	_	_	_	_	_	_	_	_	_	_
Male	30	_	_	_	_	_	_	_	_	_	_	_
Female	33	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	11,478	_	1	_	_	_	2	_	6	1.0	30	60
Male	4,331	_	_	_	_	_	2	_	3	6	17	42
Female	7,147	_	1	_	_	_	_	_	3	4	13	18
Non-Hispanic white, both sexes	10,514	_	1	_	_	_	2	_	5	5	23	52
Male	3,980	_	_	_	_	_	2	_	3	3	15	38
Female	6,534	_	1	_	_	_	_	_	2	2	8	14
Non-Hispanic black, both sexes	823	_	_	_	_	_	_	_	1	4	7	8
Male	296	_	_	_	_	_	_	_	_	2	2	4
Female	527	_	_	_	_	_	_	_	1	2	5	4
Origin not stated 2/, both sexes	14	_	_	_	_	_	_	_	_	_	_	_
Male	7	_	_	_	_	_	_	_	_	_	_	_
Female	7	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other diseases of circulatory system (I71-I78) All origins, both sexes..... 23,693 1.0 Male.... 12.306 Female..... 11.387 Hispanic, both sexes..... 2.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 22,800 11,786 Male.... 2. 2.7 11,014 Female..... Non-Hispanic white, both sexes..... 19,775 10,308 Male.... 9,467 Female..... Non-Hispanic black, both sexes..... 2,523 1,211 Male.... Female..... 1,312 2.0 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Aortic aneurysm and dissection (I71) All origins, both sexes..... 13,753 Male.... 8,140 2.8 Female..... 5,613 Hispanic, both sexes..... 1.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 7.8 Male.... Female..... Non-Hispanic 1/, both sexes..... 13,266 7,807 Male.... 5,459 Female..... Non-Hispanic white, both sexes..... 11,719 6,906 Male.... 4,813 Female..... Non-Hispanic black, both sexes..... 1,215 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other diseases of arteries, arterioles and capillaries (I72-I78) All origins, both sexes..... 9,940 Male.... 4,166 Female..... 5,774 2.8 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... 2.4 Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 9,534 3,979 Male.... 2. 5,555 Female..... Non-Hispanic white, both sexes..... 8,056 3,402 Male.... 2. 4,654 Female..... Non-Hispanic black, both sexes..... 1,308 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other disorders of circulatory system (I80-I99) All origins, both sexes..... 4,673 Male.... 2,083 Female..... 2,590 Hispanic, both sexes..... 3.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... 2.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 4,439 1,971 Male.... 1.5 2.2 2,468 Female..... 2. Non-Hispanic white, both sexes..... 3,571 1,576 Male.... 2.3 1,995 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... 1.0 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Influenza and pneumonia (J10-J18) All origins, both sexes..... 59,664 Male.... 26,861 Female..... 32,803 Hispanic, both sexes..... 2,912 2.4 Male.... 1,403 2.0 Female..... 1.509 Mexican, both sexes..... 1.551 2.0 2.4 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 56,581 Non-Hispanic 1/, both sexes..... 25,359 Male.... 31,222 Female..... Non-Hispanic white, both sexes..... 49,456 21,905 Male.... 27,551 Female..... Non-Hispanic black, both sexes..... 5,620 2,729 Male.... Female..... 2,891 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Influenza (J10-J11) All origins, both sexes..... 1.100 18 1.8 3 1.0 4 3 4 8 14 Male.... 421 10 8 2 6 3 Female..... 679 8 1.0 1 4 3 2 2 2 11 Hispanic, both sexes..... 26 2 2 6 2 1 Male.... 2 Female..... 15 3 2 2 Mexican, both sexes..... 17 6 2 1 1 Male.... 9 3 2 1 Female..... 3 1 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 3 1 Male 2 1 Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 4 2 Male.... Female..... 2 Non-Hispanic 1/, both sexes..... 1,071 12 16 9 3 4 12 Male.... 410 7 8 6 1 2 1 3 661 8 1 2 2 Female..... 3 Non-Hispanic white, both sexes..... 1,004 12 9 Male..... 377 2 5 5 1 1 1 627 2 2 2 8 Female..... 3 Non-Hispanic black, both sexes..... 5.0 4 3 1 1 1 1 2 25 2 Male.... 4 2 1 Female..... Origin not stated 2/, both sexes...... 3 1 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
All origins, both sexes	58,564	255	101	30	39	63	115	107	192	302	573	789
Male	26,440	156	39	17	22	41	75	71	121	178	309	463
Female	32,124	99	62	13	17	22	40	36	71	124	264	326
Hispanic, both sexes	2,886	47	22	3	8	10	21	18	30	37	45	63
Male	1,392	34	9	3	5	7	16	9	22	20	28	43
Female	1,494	13	13	-	3	3	5	9	8	17	17	20
Mexican, both sexes	1,534	34	17	3	5	4	12	9	20	24	23	40
Male	771	2.5	6	3	2	4	10	5	13	12	16	30
Female	763	9	11	-	3	-	2	4	7	12	7	10
Puerto Rican, both sexes	454	4	1	-	_	5	5	4	2	5	12	7
Male	217	4	1	-	-	2	2	2	2	3	7	6
Female	237	-	-	-	-	3	3	2	_	2	5	1
Cuban, both sexes	328	_	-	-	_	-	_	-	-	-	4	-
Male	148	-	-	-	_	-	-	-	-	-	3	-
Female	180	-	-	-	_	-	-	-	-	-	1	-
Central and South American, both sexes.	188	1	2	_	_	1	2	1	3	5	2	5
Male	82	_	1	_	_	1	2	_	2	3	_	3
Female	106	1	1	_	_	_	_	1	1	2	2	2
Other and Unknown Hispanic, both sexes.	382	8	2	_	3	_	2	4	5	3	4	11
Male	174	5	1	_	3	_	2	2	5	2	2	4
Female	208	3	1	_	_	_	_	2	_	1	2	7
Non-Hispanic 1/, both sexes	55,510	208	79	27	31	52	93	89	162	263	522	721
Male	24,949	122	30	14	17	34	58	62	99	157	277	415
Female	30,561	86	49	13	14	18	35	27	63	106	245	306
Non-Hispanic white, both sexes	48,452	106	43	14	21	39	64	64	116	171	367	512
Male	21,528	63	18		11	26	46	44	70	103	191	297
Female	26,924	43	25	8	10	13	18	20	46	68	176	215
Non-Hispanic black, both sexes	5,570	89	31	11	8	13	26	22	40	81	141	187
Male	2,704	51	10		4	8	11	16	24	48	78	105
Female	2,866	38	21	3	4	5	15	- 6	16	33	63	82
Origin not stated 2/, both sexes	168	-		_	_	1	1	_	_	2	6	5
Male	99	_	_	_	_	_	1	_	_	1	4	5
Female	69	_	_	_	_	1	_	_	_	1	2	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other acute lower respiratory infections (J20-J22) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... 2. Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Acute bronchitis and bronchiolitis (J20-J21) All origins, both sexes..... 226 58 16 1 3 1 3 3 3 Male.... 108 33 9 1 Female..... 118 25 1 1 1 2 2 Hispanic, both sexes..... 3.0 14 5 1 1 1 Male.... 7 Female..... 16 7 3 1 Mexican, both sexes..... 21 12 3 1 1 Male.... 9 6 1 Female..... 12 6 3 1 Puerto Rican, both sexes..... 2 1 Male.... 1 Female..... Cuban, both sexes..... Male.... Female.... 1 Central and South American, both sexes. Male 1 Female..... Other and Unknown Hispanic, both sexes. 5 2 1 Male.... 3 1 1 Female..... 2 1 Non-Hispanic 1/, both sexes..... 195 44 11 3 2 3 Male.... 93 26 7 2 1 2. 2 1 102 18 2 Female..... 1 Non-Hispanic white, both sexes..... 159 28 2 1 3 Male..... 72 16 2 1 1 2. 2 87 12 2 2 Female..... 1 Non-Hispanic black, both sexes..... 28 12 6 1 15 1 Male.... 6 4 Female..... 6 Origin not stated 2/, both sexes...... 1 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Unspecified acute lower respiratory infection (J22) All origins, both sexes..... 93 2 1 2 Male.... 29 2 Female..... 1 Hispanic, both sexes..... 2 1 Male.... 1 Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... 1 Male.... 1 Female.... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 91 1 28 Male.... 1 2 63 1 Female..... Non-Hispanic white, both sexes..... 87 2 27 Male.... 1 2 Female..... 60 Non-Hispanic black, both sexes..... 3 1 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Chronic lower respiratory diseases (J40-J47) 1.232 Male.... 3.0 Female..... 63,341 Hispanic, both sexes..... 3.107 Male.... 1,631 Female..... 1.476 Mexican, both sexes..... 1.428 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 118,650 1,178 56,884 Male.... 61,766 Female..... Non-Hispanic white, both sexes..... 109,670 51,909 Male.... 2.0 57,761 Female..... Non-Hispanic black, both sexes..... 7.331 4.010 1.0 Male.... Female..... 3,321 Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Bronchitis, chronic and unspecified (J40-J42) All origins, both sexes..... 773 17 12 3 1 2 5 2 3 2 10 15 Male.... 338 12 9 3 2 2 2 3 10 Female..... 435 5 3 3 1 7 5 Hispanic, both sexes..... 3.8 4 2 1 Male.... Female..... 19 1 1 Mexican, both sexes..... 19 4 1 1 Male.... 9 3 1 Female..... 1.0 1 1 Puerto Rican, both sexes..... 6 1 1 1 Male.... 1 1 Female..... Cuban, both sexes..... 9 Male 3 6 Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 4 Male.... 2 Female..... 2 Non-Hispanic 1/, both sexes..... 733 13 10 3 10 13 Male.... 318 9 7 2 1 2 2 2 2. 2 3 9 415 3 3 Female..... 1 Non-Hispanic white, both sexes..... 653 1 3 11 276 Male.... 3 1 1 1 2 2 2. 377 2 Female..... 1 1 Non-Hispanic black, both sexes..... 64 5 5 1 1 1 2 2 34 Male.... 5 4 1 1 2 1 Female..... 1 Origin not stated 2/, both sexes...... 2 Male.... Female.....

2

4

4

2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Emphysema (J43) All origins, both sexes..... 13,639 3 2 1 3 1 9 23 57 130 Male.... 7,075 3 5 14 Female..... 6,564 1 4 9 22 43 Hispanic, both sexes..... 269 1 1 3 2 1 1 Male.... 152 Female..... 117 1 1 Mexican, both sexes..... 128 1 1 1 1 Male.... 77 1 Female..... 51 1 1 Puerto Rican, both sexes..... 29 1 Male.... 18 1 11 Female..... Cuban, both sexes..... 55 1 Male 26 29 1 Female..... 6 Central and South American, both sexes. Male 3 Female..... Other and Unknown Hispanic, both sexes. 51 1 Male.... 2.8 1 Female..... 23 Non-Hispanic 1/, both sexes..... 13,348 1 22 54 126 6,909 2 Male.... 1 3 4 1.3 32 84 6,439 1 1 9 22 42 Female..... Non-Hispanic white, both sexes..... 12,471 2 19 39 106 6,348 Male.... 1 2 2 1.0 22 68 6,123 4 9 17 38 Female..... Non-Hispanic black, both sexes..... 716 2 1 1 1 3 12 17 458 2 8 14 Male.... 1 1 3 Female..... 258 3 Origin not stated 2/, both sexes...... 22 2 14 2 Male.... Female..... 8

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
All origins, both sexes	3,816	7	29	39	66	76	83	97	100	157	219	293
Male	1,386	7	19	23	41	50	49	52	61	64	94	122
Female	2,430	-	10	16	25	26	34	45	39	93	125	171
Hispanic, both sexes	281	3	7	5	6	5	9	12	11	10	23	20
Male	111	3	4	2	4	3	6	7	8	5	14	7
Female	170	-	3	3	2	2	3	5	3	5	9	13
Mexican, both sexes	98	2	6	3	1	2	5	6	1	3	6	6
Male	46	2	4	1	_	1	3	4	1	2	3	3
Female	52	-	2	2	1	1	2	2	-	1	3	3
Puerto Rican, both sexes	97	1	1	_	3	1	1	1	7	5	9	8
Male	37	1	_	_	2	1	1	1	5	3	8	2
Female	60	-	1	_	1	-	-	-	2	2	1	6
Cuban, both sexes	25	_	_	_	_	_	1	_	_	1	2	1
Male	6	-	-	_	_	-	-	-	-	_	1	1
Female	19	_	_	_	_	-	1	_	_	1	1	_
Central and South American, both sexes.	27	_	_	_	1	_	_	2	2	_	3	2
Male	11	_	_	_	1	_	_	1	1	_	1	_
Female	16	_	_	_	_	_	_	1	1	_	2	2
Other and Unknown Hispanic, both sexes.	34	_	_	2	1	2	2	3	1	1	3	3
Male	11	_	_	1	1	1	2	1	1	_	1	1
Female	23	_	_	1	_	1	_	2	_	1	2	2
Non-Hispanic 1/, both sexes	3,524	4	22	34	60	71	74	85	88	147	194	273
Male	1,272	4	15	21	37	47	43	45	53	59	80	115
Female	2,252	_	7	1.3	23	24	31	40	35	88	114	158
Non-Hispanic white, both sexes	2,377	2	7	9	25	27	36	39	41	87	108	132
Male	743	2	5	3	15	19	20	14	23	32	42	51
Female	1,634	_	2	6	10	- 8	16	25	18	55	66	81
Non-Hispanic black, both sexes	1,001	2	13	23	34	40	34	44	47	57	82	129
Male	452	2	9	17	22	25	20	30	30	2.4	36	59
Female	549	_	4	6	12	15	14	14	17	33	46	70
Origin not stated 2/, both sexes	11	_	_	_					1	_	2	-
Male	3	_	_	_	_	_	_	_	_	_	_	_
Female	8	_	_	_	_	_	_	_	1	_	2	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other chronic lower respiratory diseases (J44,J47) Male.... 2.0 Female..... 53,912 2,519 Hispanic, both sexes..... Male.... 1.349 Female..... 1.170 Mexican, both sexes..... 1.183 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 101,045 48,385 Male.... 52,660 Female..... Non-Hispanic white, both sexes..... 94,169 44,542 Male.... 49,627 Female..... Non-Hispanic black, both sexes..... 5,550 3.066 Male.... Female..... 2,484 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Pneumoconioses and chemical effects (J60-J66,J68) All origins, both sexes..... 1.033 1 2 4 Male.... 993 1 2 Female..... 40 2 Hispanic, both sexes..... 25 Male.... Female..... Mexican, both sexes..... 17 Male.... 17 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 1 Male.... 6 1 1 Female..... Non-Hispanic 1/, both sexes..... 1,007 967 Male.... 1 1 1 2 Female..... Non-Hispanic white, both sexes..... 912 Male.... 1 1 1 39 Female..... Non-Hispanic black, both sexes..... 54 54 Male..... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Pneumonitis due to solids and liquids (J69) All origins, both sexes..... 16,780 Male.... 8,593 Female..... 8.187 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... 1.8 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... 2.7 Female..... 16,213 Non-Hispanic 1/, both sexes..... 8,312 Male.... 7,901 Female..... Non-Hispanic white, both sexes..... 14,411 7,367 Male.... 1.3 7,044 Female..... Non-Hispanic black, both sexes..... 1,488 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other diseases of respiratory system (J00-J06, J30-J39, J67, J70-J98) All origins, both sexes..... 27.310 Male.... 13,434 Female..... 13.876 Hispanic, both sexes..... 1,508 2.4 3.0 Male.... Female..... 2.7 Mexican, both sexes..... Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... \_ 25,769 Non-Hispanic 1/, both sexes..... 12,687 Male.... 2.2 3.0 13,082 Female..... Non-Hispanic white, both sexes..... 22,756 11,218 Male.... 2.4 11,538 Female..... Non-Hispanic black, both sexes..... 2.374 1,147 1.0 Male.... Female..... 1,227 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Peptic ulcer (K25-K28) All origins, both sexes..... 3,682 1.0 2.8 Male.... 1,690 Female..... 1,992 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 2.0 Male.... 1.0 Female..... Cuban, both sexes..... Male Female..... 1.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 3,506 Non-Hispanic 1/, both sexes..... 1,592 Male.... 2. 1,914 Female..... 2. Non-Hispanic white, both sexes..... 3,045 1,356 Male.... 1.0 1,689 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Diseases of appendix (K35-K38) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male..... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Hernia (K40-K46)												
All origins, both sexes	1,602	40	4	_	4	7	2	5	5	11	25	43
Male	652	20	1	_	2	4	1	3	4	6	16	20
Female	950	20	3	_	2	3	1	2	1	5	9	23
Hispanic, both sexes	88	11	_	_	1	_	_	2	_	_	2	5
Male	31	4	_	_	_	_	_	2	_	_	2	2
Female	57	7	_	_	1	_	_	_	_	_	_	3
Mexican, both sexes	51	9	_	_	1	_	_	1	_	_	1	2
Male	18	3	_	_	_	_	_	1	_	_	1	2
Female	33	6	_	_	1	_	_	_	_	_	_	_
Puerto Rican, both sexes	11	_	_	_	_	_	_	_	_	_	_	_
Male	4	_	_	_	_	_	_	_	_	_	_	_
Female	7	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	8	_	_	_	_	_	_	_	_	_	_	_
Male	5	-	_	-	-	-	-	-	-	_	_	-
Female	3	-	_	-	-	-	-	-	-	_	_	-
Central and South American, both sexes.	7	2	_	-	-	-	-	-	-	_	1	2
Male	2	1	_	-	-	-	-	-	-	_	1	-
Female	5	1	_	-	-	-	-	-	-	_	_	2
Other and Unknown Hispanic, both sexes.	11	_	_	_	_	_	_	1	_	_	_	1
Male	2	-	_	-	-	-	-	1	-	_	_	-
Female	9	-	_	-	-	-	-	-	-	_	_	1
Non-Hispanic 1/, both sexes	1,510	29	4	_	3	7	2	3	5	11	23	38
Male	617	16	1	-	2	4	1	1	4	6	14	18
Female	893	13	3	_	1	3	1	2	1	5	9	20
Non-Hispanic white, both sexes	1,335	22	2	_	3	5	2	2	5	6	17	25
Male	540	12	_	_	2	3	1	_	4	2	11	14
Female	795	10	2	_	1	2	1	2	1	4	6	11
Non-Hispanic black, both sexes	148	6	2	-	-	2	-	1	-	3	6	11
Male	65	4	1	_	_	1	-	1	-	2	3	4
Female	83	2	1	_	_	1	-	-	-	1	3	7
Origin not stated 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Chronic liver disease and cirrhosis (K70,K73-K74) All origins, both sexes..... 27,013 2,017 3,351 Male.... 17.545 1,394 2.344 Female..... 9.468 1,007 Hispanic, both sexes..... 3,383 Male.... 2,376 Female..... 1,007 Mexican, both sexes..... 2.245 Male.... 1,601 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male 1.0 Female..... Other and Unknown Hispanic, both sexes. Ω 1 Q Male.... Female 23,542 Non-Hispanic 1/, both sexes..... 1,682 2,869 15,104 Male.... 1,122 1.952 8,438 Female..... Non-Hispanic white, both sexes..... 20,144 1,360 2,368 12,931 Male.... 1.612 7,213 Female..... Non-Hispanic black, both sexes..... 2,506 1,650 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
All origins, both sexes	12,548	_	_	_	_	_	11	42	186	533	1,338	2,023
Male	9,155	-	-	_	-	-	8	28	139	346	921	1,404
Female	3,393	-	-	_	-	-	3	14	47	187	417	619
Hispanic, both sexes	1,685	-	-	_	-	-	2	8	47	100	225	299
Male	1,414	-	-	_	-	-	2	6	42	80	186	262
Female	271	-	-	_	-	-	-	2	5	20	39	37
Mexican, both sexes	1,171	-	-	-	-	_	2	5	33	75	166	204
Male	998	-	-	_	-	-	2	3	30	63	142	179
Female	173	-	-	_	-	-	-	2	3	12	24	25
Puerto Rican, both sexes	162	-	-	_	-	-	-	-	1	4	17	25
Male	125	-	-	_	-	-	-	-	-	2	12	20
Female	37	-	-	_	-	-	-	-	1	2	5	5
Cuban, both sexes	31	_	-	-	-	-	_	_	_	1	1	4
Male	29	_	-	-	-	-	_	_	_	1	1	4
Female	2	-	-	-	_	-	-	-	-	-	-	-
Central and South American, both sexes.	64	-	-	-	_	-	-	-	6	3	10	10
Male	51	-	-	-	_	-	-	-	5	3	7	9
Female	13	_	_	_	_	_	_	_	1	_	3	1
Other and Unknown Hispanic, both sexes.	257	_	_	_	_	_	_	3	7	17	31	56
Male	211	_	_	_	_	_	_	3	7	11	24	50
Female	46	_	_	_	_	_	_	_	_	6	7	6
Non-Hispanic 1/, both sexes	10,814	_	_	_	_	_	9	34	139	433	1,112	1.711
Male	7,700	_	_	_	_	_	6	22	97	266	734	1,133
Female	3,114	_	_	_	_	_	3	12	42	167	378	578
Non-Hispanic white, both sexes	9,153	_	_	_	_	_	5	25	109	341	920	1,426
Male	6,558	_	_	_	_	_	4	14	78	204	614	945
Female	2,595	_	_	_	_	_	1	11	31	137	306	481
Non-Hispanic black, both sexes	1,160	_	_	_	_	_	2	1	8	43	121	198
Male	805	_	-	_	_	-	_	1	6	29	77	132
Female	355	_	_	_	_	_	2	_	2	14	44	66
Origin not stated 2/, both sexes	49	_	_	_	_	_	_	_	_	_	1	13
Male	41	_	_	_	_	_	_	_	_	_	1	9
Female	8	_	_	_	-	-	-	-	-	_	_	4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Other chronic liver disease and cirrhosis (K73-K74) All origins, both sexes..... 14,465 1.0 1.328 Male.... 8,390 Female..... 6.075 Hispanic, both sexes..... 1,698 Male.... Female..... Mexican, both sexes..... 1.074 Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 12,728 Non-Hispanic 1/, both sexes..... 1,158 7,404 Male.... 5,324 Female..... Non-Hispanic white, both sexes..... 10,991 Male..... 6,373 4,618 Female..... Non-Hispanic black, both sexes..... 1,346 2.4 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Cholelithiasis and other disorders of gallbladder (K80-K82) All origins, both sexes..... 3,086 1 7 8 9 17 38 61 Male.... 1,370 3 4 6 11 19 32 Female..... 1,716 1 4 4 3 6 19 29 Hispanic, both sexes..... 215 1 3 7 8 1 3 Male.... 97 Female..... 118 1 5 Mexican, both sexes..... 127 1 2 1 2 1 Male.... 59 1 1 1 Female..... 68 1 1 1 Puerto Rican, both sexes..... 23 1 1 Male.... 7 1 16 1 Female..... Cuban, both sexes..... 16 1 Male 7 1 9 Female..... 16 Central and South American, both sexes. Male 10 Female..... 6 Other and Unknown Hispanic, both sexes. 33 1 4 1 3 Male.... 14 1 1 1 1 Female..... 19 3 2 Non-Hispanic 1/, both sexes..... 2,867 4 8 8 14 31 53 1,272 Male.... 1 4 5 9 16 29 1,595 3 3 15 24 Female..... Non-Hispanic white, both sexes..... 2,460 2 3 10 25 36 1,110 Male.... 1 2. 6 13 2.0 1,350 12 16 Female..... 1 2 1 Non-Hispanic black, both sexes..... 313 4 3 6 14 120 8 Male.... 2 2 3 Female..... 193 Origin not stated 2/, both sexes...... 4 Male.... Female..... 3

3

2

2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) All origins, both sexes..... 42,480 2.7 Male.... 20,370 Female..... 22,110 Hispanic, both sexes..... 2,196 2.7 Male.... Female..... 1.095 Mexican, both sexes..... 1.311 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 40,192 19,220 Male.... 20,972 Female..... Non-Hispanic white, both sexes..... 31,508 2.4 15,247 Male.... 1.3 2.4 2.3 16,261 Female..... Non-Hispanic black, both sexes..... 7.759 3,505 Male.... Female..... 4,254 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04) All origins, both sexes..... 157 7 5 1 2 1 1 1 Male.... 73 4 2 Female..... 84 5 1 Hispanic, both sexes..... 15 1 1 1 1 Male.... 1 Female..... 1 1 Mexican, both sexes..... 1.0 1 1 1 1 1 Male.... 4 1 1 1 Female..... 6 1 1 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male 2 Female.... 1 Central and South American, both sexes. Male 1 Female..... Other and Unknown Hispanic, both sexes. 1 Male.... Female..... Non-Hispanic 1/, both sexes..... 142 3 Male.... 6.5 2 2. 1 1 77 1 Female..... 1 Non-Hispanic white, both sexes..... 113 Male..... 52 2. 61 2 Female..... Non-Hispanic black, both sexes..... 24 2 1 1 1 1 1.0 Male..... 1 1 1 Female..... Origin not stated 2/, both sexes..... Male....

Female.....

6

4

2

4

2

2

3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26) All origins, both sexes..... 553 1 1 1 1 4 9 12 14 16 12 Male.... 275 1 3 3 6 11 Female..... 278 1 3 6 9 8 5 Hispanic, both sexes..... 3.0 1 1 1 1 Male.... 1 Female..... 12 1 Mexican, both sexes..... 20 1 1 1 Male.... 12 1 Female..... 8 1 1 Puerto Rican, both sexes..... 4 1 Male.... 1 Female..... Cuban, both sexes..... Male Female.... 2 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 3 Male.... 2 Female..... 1 Non-Hispanic 1/, both sexes..... 522 12 13 16 12 Male.... 256 1 2 3 6 11 6 266 1 3 Female..... 1 Non-Hispanic white, both sexes..... 416 10 Male.... 202 2 2 5 7 214 1 1 2 6 5 Female..... 2 Non-Hispanic black, both sexes..... 90 2 4 4 3 7 42 Male.... 1 1 1 4 Female..... Origin not stated 2/, both sexes..... 1 Male....

Female.....

6

6

6

4

5

2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
All origins, both sexes	41,732	165	9	7	11	26	53	83	140	277	490	801
Male	20,006	106	6	3	8	13	33	42	63	149	284	465
Female	21,726	59	3	4	3	13	20	41	77	128	206	336
Hispanic, both sexes	2,149	35	1	3	6	2	8	9	11	26	45	59
Male	1,073	24	-	-	4	1	5	3	3	12	33	37
Female	1,076	11	1	3	2	1	3	6	8	14	12	22
Mexican, both sexes	1,280	24	1	3	3	1	5	6	9	17	32	37
Male	621	14	-	-	2	1	3	2	2	7	24	23
Female	659	10	1	3	1	-	2	4	7	10	8	14
Puerto Rican, both sexes	301	2	_	_	2	1	3	-	-	5	7	11
Male	150	2	_	_	1	_	2	_	_	3	4	7
Female	151	_	_	_	1	1	1	-	-	2	3	4
Cuban, both sexes	175	3	_	_	_	_	_	_	_	_	_	1
Male	103	2	_	_	_	_	_	_	_	_	_	1
Female	72	1	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	134	4	_	_	_	_	_	1	_	3	1	2
Male	71	4	_	_	_	_	_	_	_	2	1	1
Female	63	_	_	_	_	_	_	1	_	1	_	1
Other and Unknown Hispanic, both sexes.	259	2	_	_	1	_	_	2	2	1	5	8
Male	128	2	_	_	1	_	_	1	1	_	4	5
Female	131	_	_	_	_	_	_	1	1	1	1	3
Non-Hispanic 1/, both sexes	39,492	130	7	4	5	24	45	73	129	251	445	737
Male	18,885	82	5	3	4	12	28	38	60	137	251	425
Female	20,607	48	2	1	1	12	17	35	69	114	194	312
Non-Hispanic white, both sexes	30,948	58	4	2	4	16	22	38	57	134	240	390
Male	14,981	36	2	1	3	8	13	21	21	67	149	223
Female	15,967	22	2	1	1	8	9	17	36	67	91	167
Non-Hispanic black, both sexes	7,640	64	3	2	_	7	21	32	63	110	193	327
Male	3,451	39	3	2	_	4	13	14	36	66	94	186
Female	4,189	25	_	_	_	3	8	18	27	44	99	141
Origin not stated 2/, both sexes	91	23	1	_	_	_	-	1			<i></i>	5
Male	48	_	1	_	_	_	_	1	_	_	_	3
Female	43	_	_	_	_	_	_	_	_	_	_	2
r cmarc	43	_	_	_	_	_	_	_	_	_	_	2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of kidney (N25,N27)												
All origins, both sexes	38	1	_	_	_	_	1	1	_	_	2	3
Male	16	1	_	_	_	_	1	1	_	_	_	3
Female	22	_	_	_	_	_	_	_	_	_	2	_
Hispanic, both sexes	2	_	_	_	_	_	1	_	_	_	_	_
Male	2	_	_	_	_	_	1	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	1	_	_	_	_	_	_	_	_	_	_	_
Male	1	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	1	_	_	_	_	_	1	_	_	_	_	_
Male	1	_	_	_	_	_	1	_	_	_	_	_
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	_	_	_	_	_	-	-	-	-	_	_	-
Non-Hispanic 1/, both sexes	36	1	-	-	-	-	-	1	-	-	2	3
Male	14	1	-	-	-	-	-	1	-	-	-	3
Female	22	-	-	-	-	-	-	-	-	-	2	-
Non-Hispanic white, both sexes	31	1	-	-	-	-	-	1	-	-	1	1
Male	12	1	-	-	-	-	-	1	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	_	1	-
Non-Hispanic black, both sexes	5	_	_	_	_	-	-	-	-	_	1	2
Male	2	_	_	_	_	-	-	-	-	_	_	2
Female	3	-	-	-	-	-	-		-	-	1	-
Origin not stated 2/, both sexes	_	-	-	-	-	-	-		-	-	-	-
Male	_	-	-	-	-	-	-		-	-	-	-
Female	_	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Infections of kidney (N10-N12,N13.6,N15.1) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female.... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male..... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Hyperplasia of prostate (N40) All origins, both sexes..... 507 1 Male.... 507 1 Female..... . . . . . . . . . . . . . . . . . . Hispanic, both sexes..... 19 Male.... Female..... . . . Mexican, both sexes..... 1.0 Male.... 1.0 Female..... Puerto Rican, both sexes..... Male.... 2 Female.... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... . . . . . . . . . . . . Other and Unknown Hispanic, both sexes. Male.... 7 Female..... Non-Hispanic 1/, both sexes..... 486 Male.... 486 1 Female..... . . . . . . Non-Hispanic white, both sexes..... 438 438 Male.... 1 Female..... Non-Hispanic black, both sexes..... 34 34 Male.... Female..... . . . . . . . . . . . . Origin not stated 2/, both sexes...... 2 2 Male.... Female..... .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Inflammatory diseases of female pelvic organs (N70-N76) All origins, both sexes..... 112 1 1 1 2 2 4 4 8 Male.... Female..... 112 1 1 1 2 2 4 4 8 Hispanic, both sexes..... 9 1 1 1 Male.... Female..... 1 1 1 Mexican, both sexes..... 4 1 1 1 Male.... Female..... 1 1 1 Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... 1 Male Female..... 2 Central and South American, both sexes. Male . . . . . . . . . . . . Female..... 2 Other and Unknown Hispanic, both sexes. 2 Male.... Female..... 2 2 Non-Hispanic 1/, both sexes..... 102 1 1 4 4 8 Male.... . . . . . . . . . . . . Female..... 102 2 8 1 1 Non-Hispanic white, both sexes..... 83 1 4 Male.... 83 4 4 Female..... 2. 1 Non-Hispanic black, both sexes..... 19 1 2 4 Male.... Female..... 19 4 Origin not stated 2/, both sexes...... 1 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Pregnancy, childbirth and the puerperium (000-099) All origins, both sexes..... 697 2 37 131 142 145 96 61 39 Male.... . . . . . . . . . Female..... 697 2 37 131 142 145 96 61 39 . . . . . . . . . Hispanic, both sexes..... 115 11 18 27 22 16 10 6 Male.... . . . . . . . . . Female..... 115 11 18 2.7 22 16 1.0 6 . . . . . . . . . Mexican, both sexes..... 78 8 14 19 12 11 6 4 Male.... . . . . . . . . . Female..... 78 14 19 12 11 8 6 4 . . . . . . . . . Puerto Rican, both sexes..... 2 2 2 2 8 . . . Male.... . . . . . . . . . 2 2 2 2 Female..... . . . . . . . . . Cuban, both sexes..... Male . . . . . . . . . . . . Female..... . . . . . . . . . 1 2 5 5 3 1 Central and South American, both sexes. 2.0 . . . . . . 3 Male . . . . . . . . . . . . . . . Female..... 2 5 . . . . . . . . . Other and Unknown Hispanic, both sexes. 9 . . . . . . 2 1 3 1 1 . . . Male.... . . . . . . . . . Female..... 9 2 . . . . . . . . . Non-Hispanic 1/, both sexes..... 581 2 26 113 115 123 79 51 33 . . . . . . . . . Male.... . . . . . . Female..... 581 26 113 115 123 51 33 2 . . . . . . . . . Non-Hispanic white, both sexes..... 302 13 47 52 70 45 29 16 . . . . . . . . . Male.... . . . . . . . . . 302 13 47 52 70 45 29 16 Female..... . . . . . . . . . Non-Hispanic black, both sexes..... 245 12 61 59 47 29 15 15 . . . . . . . . . Male.... . . . . . . . . . Female..... 61 59 15 . . . . . . . . . Origin not stated 2/, both sexes...... 1 1 . . . . . . . . . Male.... Female.....

. . .

. . .

. . .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Pregnancy with abortive outcome (000-007) All origins, both sexes..... 32 1 13 6 5 4 2 Male.... . . . . . . . . . Female..... 32 1 13 6 5 4 2 . . . . . . . . . Hispanic, both sexes..... 1 1 Male.... . . . . . . . . . Female..... 1 . . . . . . . . . Mexican, both sexes..... 1 1 Male.... . . . . . . . . . Female..... . . . 1 . . . . . . Puerto Rican, both sexes..... Male.... . . . . . . . . . Female.... . . . . . . . . . Cuban, both sexes..... Male . . . . . . . . . Female..... . . . . . . . . . Central and South American, both sexes. . . . . . . Male . . . . . . . . . . . . . . . . . . Female..... . . . . . . . . . Other and Unknown Hispanic, both sexes. . . . . . . . . . Male.... . . . . . . . . . Female..... . . . . . . . . . Non-Hispanic 1/, both sexes..... 31 1 13 5 5 4 2 . . . . . . . . . Male.... . . . . . . 31 13 1 5 Female..... . . . . . . . . . Non-Hispanic white, both sexes..... 1.0 5 2 . . . . . . . . . Male.... . . . . . . . . . 10 Female..... 5 2. 2. . . . . . . . . . Non-Hispanic black, both sexes..... 19 1 7 3 3 4 1 . . . . . . . . . Male.... . . . . . . . . . Female..... 7 . . . . . . . . . Origin not stated 2/, both sexes...... \_ . . . . . . . . . Male.... Female..... . . . . . . . . .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other complications of pregnancy, childbirth and the puerperium (010-099) All origins, both sexes..... 665 2 36 118 136 140 92 59 39 Male.... . . . . . . . . . Female..... 665 2 36 118 136 140 92 59 39 . . . . . . . . . Hispanic, both sexes..... 114 11 18 26 22 16 10 6 Male.... . . . . . . . . . Female..... 114 11 18 26 22 16 1.0 . . . . . . . . . Mexican, both sexes..... 77 8 14 18 12 11 6 Male.... . . . . . . . . . Female..... 14 18 12 11 . . . 8 6 . . . . . . Puerto Rican, both sexes..... 2 2 2 2 8 . . . Male.... . . . . . . . . . 2 2 2 2 Female..... . . . . . . . . . Cuban, both sexes..... Male . . . . . . . . . . . . Female..... . . . . . . . . . 1 2 5 5 3 1 Central and South American, both sexes. 2.0 . . . . . . 3 Male . . . . . . . . . . . . . . . Female..... 2 5 . . . . . . . . . Other and Unknown Hispanic, both sexes. 9 . . . . . . . . . 2 1 3 1 1 Male.... . . . . . . . . . Female..... 9 2 . . . . . . . . . Non-Hispanic 1/, both sexes..... 550 2 25 100 110 118 75 49 33 . . . . . . . . . Male.... . . . . . . Female..... 25 100 110 75 49 33 550 2 118 . . . . . . . . . Non-Hispanic white, both sexes..... 292 13 42 50 68 45 29 16 . . . . . . . . . Male.... . . . . . . . . . 292 13 42 50 45 29 16 Female..... 68 . . . . . . . . . Non-Hispanic black, both sexes..... 226 11 54 56 44 25 14 15 . . . . . . . . . Male.... . . . . . . . . . Female..... 54 56 14 15 . . . . . . . . . Origin not stated 2/, both sexes...... - 1 1 . . . . . . . . . Male....

. . .

. . .

. . .

Female.....

6

4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Certain conditions originating in the perinatal period (P00-P96) All origins, both sexes..... 14,213 14,079 61 25 12 4 7 6 4 3 3 Male.... 8,093 8,013 38 15 8 3 2 3 Female..... 6,120 6,066 23 1.0 4 1 5 5 2 1 1 Hispanic, both sexes..... 2,681 2,660 1.0 5 3 1 1 1 Male.... 1,538 1.527 6 Female..... 1.143 1.133 4 2 1 1 1 Mexican, both sexes..... 1.892 1.880 8 1 1 1 1 Male.... 1,085 1.079 5 1 Female..... 807 801 3 1 1 Puerto Rican, both sexes..... 252 250 1 1 Male.... 136 134 1 1 116 116 Female..... Cuban, both sexes..... 35 35 Male 2.0 20 15 15 Female..... 174 170 2 1 1 Central and South American, both sexes. Male 94 92 1 1 Female..... 80 78 1 Other and Unknown Hispanic, both sexes. 328 325 1 2 Male.... 203 202 1 Female..... 125 123 1 1 Non-Hispanic 1/, both sexes..... 11,371 11,258 51 20 9 3 6 6 4 Male.... 6,469 6,400 32 12 6 3 2. 1 2. 3 2 1 4,902 4,858 19 8 4 5 2 Female..... 3 1 Non-Hispanic white, both sexes..... 6,118 6,049 24 13 3 5 4 3 Male..... 3,485 3,529 17 8 3 3 2 1 2 3 1 2,589 2,564 5 3 3 2 Female..... 2. Non-Hispanic black, both sexes..... 4,740 4,703 23 6 3 1 2 2 2.653 2.633 13 Male.... 3 2 2 Female..... 2,087 2,070 1.0 1 Origin not stated 2/, both sexes...... 161 161 86 86 Male....

Female.....

75

75

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) 10,498 All origins, both sexes..... 5,622 Male.... 5,423 2,898 Female..... 5.075 2.724 Hispanic, both sexes..... 1,765 1.278 2.4 Male.... Female..... 1.0 Mexican, both sexes..... 1.288 Male.... 5.8 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 8,686 4,309 4,493 Male.... 2.256 4,193 2,053 Female..... Non-Hispanic white, both sexes..... 6,598 3,016 3,375 1,572 Male.... 3,223 1,444 Female..... Non-Hispanic black, both sexes..... 1,697 1.041 Male.... Female..... Origin not stated 2/, both sexes..... 2.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) All origins, both sexes..... 30.126 3,361 1,132 1.270 Male.... 13,712 2.021 Female..... 16,414 1,340 Hispanic, both sexes..... 1,621 Male.... Female..... 2.6 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 28,297 2,881 1,017 1,163 Male.... 12,607 1.730 15,690 1,151 Female..... Non-Hispanic white, both sexes..... 23,344 1,654 Male.... 10,064 13,280 Female..... Non-Hispanic black, both sexes..... 4.376 1,106 Male..... 2,244 Female..... 2,132 2.4 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13,0-N13,5,N13,7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98) 1,175 1,393 1.839 3.013 5,059 7.428 Male.... 1,045 1,755 2,984 4,432 Female..... 119.189 1,258 2,075 2,996 Hispanic, both sexes..... 9,217 Male.... 4,541 Female..... 4,676 Mexican, both sexes..... 5.287 Male.... 2.714 Female..... 2.573 2.1 Puerto Rican, both sexes..... 1.133 2.4 Male.... Female..... Cuban, both sexes..... Male Female..... 2.0 2.2 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1,365 a Ω a 1 4 Male.... Female..... Non-Hispanic 1/, both sexes..... 192,252 1,018 1,219 1,592 2,680 4,558 6,843 77 963 1,513 Male.... 2,653 4.031 114,289 1,167 1,905 2,812 Female..... Non-Hispanic white, both sexes..... 165,312 1,029 1,858 3,353 5,087 66,039 2,054 Male.... 1,102 3.104 99,273 1,299 1.983 Female..... Non-Hispanic black, both sexes..... 23,295 1,033 1,588 10,222 Male.... Female..... 13,073 Origin not stated 2/, both sexes..... 3.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 1 - 4non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Accidents (unintentional injuries) (V01-X59,Y85-Y86) 1,052 1,641 1,126 1,540 6,825 8,624 6.771 6,261 7.319 9,152 9.426 Male.... 72,050 4,744 6,679 5,222 4,649 5,252 6,419 6,602 Female..... 39.962 2,081 1,945 1,549 1,612 2,067 2,733 2,824 Hispanic, both sexes..... 10,408 1,416 1,164 Male.... 7.796 1,186 Female..... 2,612 Mexican, both sexes..... 6.767 Male.... 5.185 Female..... 1,582 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 2.4 2.7 Male 2.0 Female..... Central and South American, both sexes. 2.2 Male Female..... Other and Unknown Hispanic, both sexes. 1 241 1 Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 101,235 1,323 1,320 5,855 7,190 5,585 5,316 6,360 8,237 8,654 Male.... 63.982 3,995 5,481 4,228 3,875 4,473 5.706 5.992 37,253 1,860 1,709 1,357 1,441 1,887 2,531 2,662 Female..... Non-Hispanic white, both sexes..... 85,466 4,911 5,936 4,461 4,251 5,159 6,866 7,124 53,465 4,563 3,645 Male.... 3,322 3,374 3,084 4,767 4,953 32,001 1,589 1,373 1,087 1,167 1,514 2,099 2,171 Female..... Non-Hispanic black, both sexes..... 12,446 1,114 1,305 8.430 Male.... Female..... 4,016 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years	years	years	years	years	years	years	years	years	years
Transport accidents (V01-V99,Y85)												
All origins, both sexes	48,053	145	659	681	1,074	5,394	5,983	4,094	3,427	3,460	3,787	3,775
Male	33,311	81	359	392	647	3,577	4,536	3,117	2,524	2,539	2,758	2,717
Female	14,742	64	300	289	427	1,817	1,447	977	903	921	1,029	1,058
Hispanic, both sexes	6,010	40	164	136	166	793	1,073	800	581	493	422	328
Male	4,551	18	95	77	104	605	887	659	464	397	323	251
Female	1,459	22	69	59	62	188	186	141	117	96	99	77
Mexican, both sexes	4,170	31	142	97	129	557	778	566	400	342	301	215
Male	3,196	15	84	52	81	429	656	466	323	276	225	175
Female	974	16	58	45	48	128	122	100	77	66	76	40
Puerto Rican, both sexes	362	2	8	9	9	47	51	34	30	40	29	28
Male	272	2	4	5	6	36	35	29	23	35	26	22
Female	90	2	4	4	3	11	16	29 5	23 7	5	3	6
	229	_	2	3	5	24	21	10	13	20	16	13
Cuban, both sexes		_	1	3	3			8	13			10
Male	172					21	16	-		14	14	
Female	57	-	1	-	2	3	5	2	1	6	2	3
Central and South American, both sexes.	603	1	6	14	3	66	126	106	71	41	42	31
Male	459	1	3	9	2	52	105	89	55	34	35	20
Female	144	-	3	5	1	14	21	17	16	7	7	11
Other and Unknown Hispanic, both sexes.	646	6	6	13	20	99	97	84	67	50	34	41
Male	452	2	3	8	12	67	75	67	51	38	23	24
Female	194	4	3	5	8	32	22	17	16	12	11	17
Non-Hispanic 1/, both sexes	41,892	104	494	542	902	4,591	4,897	3,281	2,836	2,961	3,351	3,430
Male	28,652	63	263	312	542	2,967	3,641	2,450	2,051	2,137	2,427	2,452
Female	13,240	41	231	230	360	1,624	1,256	831	785	824	924	978
Non-Hispanic white, both sexes	34,315	66	338	390	695	3,867	3,945	2,496	2,156	2,338	2,727	2,802
Male	23,403	39	177	229	427	2,476	2,961	1,858	1,538	1,694	1,983	2,014
Female	10,912	27	161	161	268	1,391	984	638	618	644	744	788
Non-Hispanic black, both sexes	5,744	32	129	120	164	535	693	603	531	480	476	511
Male	4,109	20	68	65	93	375	509	468	412	357	347	367
Female	1,635	12	61	55	71	160	184	135	119	123	129	144
Origin not stated 2/, both sexes	151	1	1	3	6	10	13	13	10	6	14	17
Male	108	_	1	3	1	5	8	8	9	5	8	14
Female	43	1	-	_	5	5	5	5	1	1	6	3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2) All origins, both sexes..... 44,933 1,013 5,224 5,763 3,849 3,187 3.190 3,473 3.422 Male.... 30.837 3,453 4,361 2,924 2,321 2,318 2,508 2,431 Female..... 14.096 1,771 1,402 Hispanic, both sexes..... 5,739 1.048 Male.... 4,318 Female..... 1,421 Mexican, both sexes..... 4.007 Male.... 3.050 Female..... Puerto Rican, both sexes..... Male.... 2.7 Female..... Cuban, both sexes..... 2.0 Male Female..... Central and South American, both sexes. 3.0 Male Female..... Other and Unknown Hispanic, both sexes. 1 2 Male.... Female..... Non-Hispanic 1/, both sexes..... 39,059 4,442 4,704 3,071 2,635 2,722 3,063 3,106 26,427 2,858 1,883 Male.... 3,490 2,289 1,942 2.199 2.193 12,632 1,584 1,214 Female..... Non-Hispanic white, both sexes..... 31,931 3,749 3,786 2,330 1,991 2,135 2,487 2,535 21,540 2,393 1,403 1,795 1,797 Male.... 2,842 1,734 1,530 10,391 1,356 Female..... Non-Hispanic black, both sexes..... 5,395 3,821 Male.... Female..... 1.574 Origin not stated 2/, both sexes..... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9) All origins, both sexes..... 1,199 Male.... Female..... Hispanic, both sexes..... 2.0 2.0 Male.... Female..... 2.0 Mexican, both sexes..... 1.0 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 1,028 Non-Hispanic 1/, both sexes..... Male.... 8.5 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... 1 0 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85) All origins, both sexes..... 1,921 Male.... 1.548 Female..... Hispanic, both sexes..... Male.... 1.0 Female..... Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,805 1,450 Male.... 8.5 Female..... Non-Hispanic white, both sexes..... 1,602 1,283 Male.... 2. 1.5 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of												
Diseases, Tenth Revision, 1992),												
Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years									
Nontransport accidents (W00-X59,Y86)												
All origins, both sexes	63,959	907	982	445	466	1,431	2,641	2,677	2,834	3,859	5,365	5,651
Male	38,739	523	605	290	332	1,167	2,143	2,105	2,125	2,713	3,661	3,885
Female	25,220	384	377	155	134	264	498	572	709	1,146	1,704	1,766
Hispanic, both sexes	4,398	119	149	42	47	163	343	364	341	447	461	401
Male	3,245	69	91	25	35	135	299	320	290	366	368	322
Female	1,153	50	58	17	12	28	44	44	51	81	93	79
Mexican, both sexes	2,597	93	110	32	31	125	218	240	221	246	261	237
Male	1,989	53	64	17	24	103	197	221	191	211	218	199
Female	608	40	46	15	7	22	21	19	30	35	43	38
Puerto Rican, both sexes	532	4	4	3	7	8	31	44	41	80	60	57
Male	385	2	4	2	5	6	29	35	32	59	47	39
Female	147	2	-	1	2	2	2	9	9	21	13	18
Cuban, both sexes	228	3	2	2	1	3	3	7	5	7	13	13
Male	136	2	2	2	1	2	2	5	5	6	6	9
Female	92	1	-	-	-	1	1	2	-	1	7	4
Central and South American, both sexes.	346	4	16	-	2	13	47	30	34	38	41	20
Male	271	3	9	_	2	11	40	27	30	35	35	18
Female	75	1	7	_	_	2	7	3	4	3	6	2
Other and Unknown Hispanic, both sexes.	695	15	17	5	6	14	44	43	40	76	86	74
Male	464	9	12	4	3	13	31	32	32	55	62	57
Female	231	6	5	1	3	1	13	11	8	21	24	17
Non-Hispanic 1/, both sexes	59,343	787	829	401	418	1,264	2,293	2,304	2,480	3,399	4,886	5,224
Male	35,330	453	512	265	296	1,028	1,840	1,778	1,824	2,336	3,279	3,540
Female	24,013	334	317	136	122	236	453	526	656	1,063	1,607	1,684
Non-Hispanic white, both sexes	51,151	491	595	230	258	1,044	1,991	1,965	2,095	2,821	4,139	4,322
Male	30,062	296	376	148	187	846	1,602	1,516	1,546	1,951	2,784	2,939
Female	21,089	195	219	82	71	198	389	449	549	870	1,355	1,383
Non-Hispanic black, both sexes	6,702	266	188	142	131	163	223	268	302	478	638	794
Male	4,321	139	114	98	87	134	173	204	219	309	421	523
Female	2,381	127	74	44	44	29	50	64	83	169	217	271
Origin not stated 2/, both sexes	218	1	4	2	1	4	5	9	13	13	18	26
Male	164	1	2	-	1	4	4	7	11	11	14	23
Female	54	-	2	2	-	-	1	2	2	2	4	3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for	All	Under	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
non-Hispanic population, and sex	ages	1 year	years	years	years	years	years	years	years	years	years	years
non hispanic populacion, and bex	ages	ı year	<i>y</i> carb	<i>y</i> carb	years	years	years	years	years	yearb	years	years
Falls (W00-W19)												
All origins, both sexes	18,807	23	47	11	26	87	154	151	169	241	418	567
Male	9,856	13	31	5	17	70	136	130	144	191	335	447
Female	8,951	10	16	6	9	17	18	21	25	50	83	120
Hispanic, both sexes	978	6	7	1	1	16	28	47	33	52	60	52
Male	653	4	4	-	1	14	28	42	32	47	49	49
Female	325	2	3	1	-	2	-	5	1	5	11	3
Mexican, both sexes	539	5	5	1	1	13	19	35	19	30	31	33
Male	373	4	2	-	1	12	19	31	19	29	29	32
Female	166	1	3	1	-	1	-	4	-	1	2	1
Puerto Rican, both sexes	102	-	1	-	-	1	2	1	2	10	7	7
Male	66	_	1	-	-	1	2	1	1	7	4	6
Female	36	_	-	-	-	-	-	-	1	3	3	1
Cuban, both sexes	91	_	-	-	-	-	-	1	-	-	1	-
Male	58	_	-	-	-	-	-	1	-	-	_	-
Female	33	_	-	-	-	-	-	-	-	-	1	-
Central and South American, both sexes.	88	_	1	-	-	1	4	8	6	4	11	4
Male	62	_	1	-	-	-	4	8	6	4	9	3
Female	26	_	-	-	-	1	-	-	-	-	2	1
Other and Unknown Hispanic, both sexes.	158	1	-	-	-	1	3	2	6	8	10	8
Male	94	_	-	-	-	1	3	1	6	7	7	8
Female	64	1	-	-	-	-	-	1	-	1	3	-
Non-Hispanic 1/, both sexes	17,780	17	40	9	25	70	126	103	134	189	357	510
Male	9,170	9	27	5	16	55	108	87	111	144	285	393
Female	8,610	8	13	4	9	15	18	16	23	45	72	117
Non-Hispanic white, both sexes	16,379	12	26	8	20	63	110	78	113	149	294	425
Male	8,318	7	19	4	12	50	95	66	95	120	241	329
Female	8,061	5	7	4	8	13	15	12	18	29	53	96
Non-Hispanic black, both sexes	980	3	9	1	3	5	9	15	18	32	49	64
Male	614	_	7	1	2	3	6	13	14	20	36	46
Female	366	3	2	-	1	2	3	2	4	12	13	18
Origin not stated 2/, both sexes	49	_	-	1	-	1	-	1	2	-	1	5
Male	33	_	-	-	-	1	-	1	1	-	1	5
Female	16	_	-	1	-	-	-	-	1	-	_	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Accidental discharge of firearms (W32-W34) All origins, both sexes..... 5.8 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... 2.8 2.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Accidental drowning and submersion (W65-W74) All origins, both sexes..... 3,308 Male.... 2.594 Female..... 2.7 2.4 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 4 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 2,856 2,220 Male.... Female..... Non-Hispanic white, both sexes..... 2,133 Male.... 1.646 Female..... Non-Hispanic black, both sexes..... 3.0 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Accidental exposure to smoke, fire and flames (X00-X09) All origins, both sexes..... 3,229 Male.... 1,902 Female..... 1,327 Hispanic, both sexes..... Male.... Female..... 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 3,038 Non-Hispanic 1/, both sexes..... 1,785 Male.... 1,253 Female..... Non-Hispanic white, both sexes..... 2,151 Male.... 1.265 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years vears vears vears Accidental poisoning and exposure to noxious substances (X40-X49) All origins, both sexes..... 20.950 1,616 1.747 1,894 2,694 3,750 3.696 13.934 1.271 1,328 1,345 1.781 2,395 2,360 Male.... Female..... 7.016 2.0 1,355 1,336 Hispanic, both sexes..... 1.716 Male.... 1.327 Female..... 3.0 2.4 Mexican, both sexes..... Male.... Female..... 1.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 19,153 1,470 1,556 1,703 2,413 3,439 3,443 12,541 1,155 1,565 Male.... 2.4 1,162 1,187 2,153 2.171 6,612 1,286 1,272 Female..... Non-Hispanic white, both sexes..... 16,313 1,354 1,403 1,485 2,052 2,957 2,866 10,684 1,069 1,048 1,856 Male.... 1,051 1,348 1.819 5,629 1,101 1,047 Female..... Non-Hispanic black, both sexes..... 2,460 1,607 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86) All origins, both sexes..... 17,016 Male.... 9.880 Female..... 7.136 2.7 5.8 Hispanic, both sexes..... 1,042 Male.... 2.8 Female..... Mexican, both sexes..... 1.0 Male.... 2.7 Female..... Puerto Rican, both sexes..... Male.... 2.4 Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Ω Male.... Female..... 15,923 Non-Hispanic 1/, both sexes..... 9,092 Male.... 6,831 Female..... Non-Hispanic white, both sexes..... 13,724 7,754 Male.... 5,970 Female..... Non-Hispanic black, both sexes..... 1,825 1,101 Male.... Female..... Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0) All origins, both sexes..... 32,439 2 283 1,700 2,616 2,412 2,662 2.948 3,690 3.743 Male.... 25.566 2 185 1,345 2,251 2,015 2,127 2,292 2,775 2.749 . . . . . . Female..... 6.873 98 355 365 397 535 656 915 994 . . . . . . Hispanic, both sexes..... 2.207 39 214 357 297 248 229 192 176 Male.... 1,840 2.4 170 317 258 158 139 . . . . . . Female..... 367 15 44 40 39 39 36 34 37 . . . . . . Mexican, both sexes..... 1.292 26 140 243 195 156 130 104 93 Male.... 1,094 14 111 217 175 130 114 86 77 . . . . . . Female..... 198 12 29 26 2.0 26 16 18 16 . . . . . . Puerto Rican, both sexes..... 238 15 30 2.8 30 26 22 30 5 . . . . . . Male.... 204 3 11 28 23 28 18 19 27 . . . . . . 34 2 4 2 5 2 3 3 Female..... 8 . . . . . . Cuban, both sexes..... 134 3 7 3 7 10 13 9 . . . Male 110 3 7 3 5 9 12 7 . . . . . . 24 2 1 1 2 Female..... . . . . . . 178 18 18 31 19 Central and South American, both sexes. . . . . . . 14 23 15 Male 133 11 14 2.4 11 20 16 9 . . . . . . Female..... 45 7 4 7 3 6 . . . . . . Other and Unknown Hispanic, both sexes. 365 Ω 38 50 40 41 40 34 29 . . . . . . Male.... 299 7 34 51 33 35 32 25 19 . . . . . . Female..... 66 1 4 8 6 8 9 10 . . . . . . Non-Hispanic 1/, both sexes..... 30,111 2 243 1,477 2,251 2,110 2,406 2,703 3,483 3,550 . . . . . . Male.... 23,633 2 160 1,167 1,928 1,754 1,912 2,087 2,605 2.599 . . . . . . 6,478 83 310 323 356 494 616 878 951 Female..... . . . . . . Non-Hispanic white, both sexes..... 27,001 1 179 1,211 1,834 1,707 2,040 2,404 3,194 3,269 . . . . . . Male..... 1,574 21,214 1 118 961 1,430 1,634 1,864 2,385 2.382 . . . . . . 5,787 61 250 260 277 406 540 809 887 Female..... . . . . . . Non-Hispanic black, both sexes..... 1,970 1 48 150 300 276 234 184 185 167 . . . . . . 1,614 29 120 262 236 184 145 148 128 Male.... . . . 1 . . . Female..... 356 30 3.8 40 50 39 37 39 . . . . . . Origin not stated 2/, both sexes..... 121 1 9 8 5 8 16 15 17 . . . . . . 93 8 6 3 6 12 12 11 Male.... 1 Female..... 2 2 2 3 6 . . . . . .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Intentional self-harm (suicide) by discharge of firearms (X72-X74) All origins, both sexes..... 16.750 59 787 1.317 1,115 1,168 1.233 1,635 1.784 Male.... 14,523 50 688 1,201 987 991 1,039 1,337 1,465 . . . . . . Female..... 2,227 9 99 116 128 177 194 298 319 . . . Hispanic, both sexes..... 888 2 97 169 122 84 69 78 67 . . . Male.... 88 106 75 63 69 61 . . . . . . Female..... 89 9 11 16 9 6 9 6 . . . . . . Mexican, both sexes..... 558 2 67 120 81 58 33 47 39 Male.... 505 2 60 112 71 52 32 40 36 . . . . . . Female..... 53 7 8 1.0 6 1 7 3 . . . . . . Puerto Rican, both sexes..... 76 13 11 8 4 8 6 Male.... 4 13 9 8 5 7 . . . . . . 4 2 1 1 Female..... . . . . . . Cuban, both sexes..... 53 4 2 3 5 7 Male 42 4 2 1 4 6 . . . . . . 11 2 1 1 1 Female..... . . . . . . 51 2 12 7 Central and South American, both sexes. . . . 6 2 8 Male 46 5 2 1.0 2 8 3 . . . . . . Female..... 5 1 2 . . . . . . Other and Unknown Hispanic, both sexes. 30 12 17 150 . . . . . . 2.0 16 10 Male.... 134 19 27 14 12 14 9 . . . . . . Female..... 16 1 3 2 1 3 1 . . . . . . Non-Hispanic 1/, both sexes..... 15,823 57 687 1,143 992 1,083 1,163 1,548 1,714 . . . . . . 13,691 Male.... 48 597 1.039 880 915 975 1,260 1.402 . . . 2,132 9 90 104 112 168 188 288 312 Female..... . . . . . . Non-Hispanic white, both sexes..... 14,507 47 581 928 803 930 1,047 1,437 1,611 . . . . . . 12,534 505 710 781 Male.... 39 844 877 1,165 1.315 . . . . . . 1,973 8 76 84 93 149 170 272 296 Female..... . . . . . . Non-Hispanic black, both sexes..... 995 8 71 180 148 116 81 73 67 . . . . . . 891 7 60 163 136 102 71 66 58 Male.... . . . . . . Female..... 104 11 17 12 14 10 7 9 . . . . . . Origin not stated 2/, both sexes..... 39 3 5 1 1 1 9 3 . . . . . . 33 3 4 1 1 8 2 Male.... 1 Female..... 1 1

. . .

. . .

7

7

7

6

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0) All origins, both sexes..... 15.689 2 224 913 1.299 1,297 1.494 1.715 2,055 1.959 Male.... 11,043 2 135 657 1,050 1,028 1,136 1,253 1,438 1,284 . . . . . . Female..... 4,646 89 256 249 269 358 462 617 675 . . . . . . Hispanic, both sexes..... 1.319 37 117 188 175 164 160 114 109 . . . Male.... 1,041 82 152 134 130 89 78 . . . . . . Female..... 278 15 35 29 23 3.0 30 25 31 . . . . . . Mexican, both sexes..... 734 2.4 73 123 114 98 97 57 54 Male.... 589 12 51 105 104 78 46 41 . . . . . . Female..... 145 12 22 18 1.0 2.0 15 11 13 . . . . . . Puerto Rican, both sexes..... 162 11 17 17 22 2.0 14 23 5 . . . . . . Male.... 132 3 7 15 14 20 13 12 20 . . . . . . 30 2 4 2 3 2 7 2 Female..... . . . . . . Cuban, both sexes..... 81 3 3 1 4 5 6 . . . Male 68 3 3 1 4 5 6 . . . . . . 13 Female..... . . . . . . 127 12 16 19 12 Central and South American, both sexes. . . . . . . 15 12 11 Male 87 6 12 14 9 12 9 . . . . . . Female..... 40 4 5 3 3 . . . . . . Other and Unknown Hispanic, both sexes. 215 Ω 18 29 24 23 25 . . . . . . 28 1 Q Male.... 165 7 15 24 19 23 18 16 10 . . . . . . Female..... 50 1 3 5 5 5 5 9 . . . . . . Non-Hispanic 1/, both sexes..... 14,288 2 186 790 1,108 1,118 1,323 1,540 1,935 1,836 . . . . . . 570 Male.... 9.942 2 112 889 874 997 1,112 1,345 1.197 . . . 4,346 74 220 219 244 326 428 590 639 Female..... . . . . . . Non-Hispanic white, both sexes..... 12,494 1 132 630 906 904 1,110 1,357 1,757 1,658 . . . . . . 8,680 730 720 987 Male.... 1 79 456 853 1,220 1.067 . . . . . . 3,814 53 174 176 184 257 370 537 591 Female..... . . . . . . Non-Hispanic black, both sexes..... 975 1 40 79 120 128 118 103 112 100 . . . . . . 723 22 60 99 100 82 74 82 70 Male.... . . . . . . 1 Female..... 252 18 19 21 28 36 29 30 30 . . . . . . Origin not stated 2/, both sexes..... 82 1 6 3 4 7 15 6 14 . . . . . . 60 1 5 2 2 5 11 4 Male....

. . .

. . .

1

2

2

Female.....

3

2

1

6

5

9

9

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1) All origins, both sexes..... 17.357 1,932 3,153 2,598 1.897 1.530 1,454 1.182 Male.... 13,578 1,631 2.749 2,227 1,537 1,187 1,037 Female..... 3.779 Hispanic, both sexes..... 3.271 2.7 Male.... 2,782 2.8 Female..... 2.7 Mexican, both sexes..... 2.303 2.8 Male.... 2.007 Female..... 2.2 2.8 Puerto Rican, both sexes..... 2.4 Male.... Female..... Cuban, both sexes..... Male Female..... 2.7 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 13,965 1,433 2,397 1,991 1,515 1,252 1,236 1,032 10,699 Male.... 1,181 2,059 1,684 1,214 3,266 Female..... Non-Hispanic white, both sexes..... 5,427 3,561 Male.... 1,866 Female..... Non-Hispanic black, both sexes..... 8,000 1,030 1,741 1,437 6,730 1,588 1,294 Male.... Female..... 1,270 Origin not stated 2/, both sexes..... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Assault (homicide) by discharge of firearms (\*U01.4,X93-X95) All origins, both sexes..... 11.624 1,578 2,549 2,063 1.440 1,025 Male.... 9.921 1,396 2,322 1,876 1,235 Female..... 1.703 Hispanic, both sexes..... 2.241 2.7 Male.... 2,031 Female..... 2.5 3.0 2.8 Mexican, both sexes..... 1.641 Male.... 1.511 Female..... Puerto Rican, both sexes..... 2.7 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. 2.7 Male Female..... Other and Unknown Hispanic, both sexes. 1 0 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 9,323 1,162 1,961 1,611 1,160 7,837 Male.... 1,015 1,769 1,450 1,486 Female..... Non-Hispanic white, both sexes..... 2,921 2,083 Male.... Female..... Non-Hispanic black, both sexes..... 6,119 1,569 1,264 5,518 1,466 1,174 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for A11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1) All origins, both sexes..... 5,733 Male.... 3,657 Female..... 2.076 Hispanic, both sexes..... 1,030 Male.... Female..... 3.0 2.6 2.7 Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... 2.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 4,642 Non-Hispanic 1/, both sexes..... 2,862 Male.... 2.2 1,780 Female..... Non-Hispanic white, both sexes..... 2,506 1,478 Male.... 1,028 Female..... Non-Hispanic black, both sexes..... 1,881 1,212 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Legal intervention (Y35, Y89.0) All origins, both sexes..... 2.7 Male.... 2.8 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... \_ Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... 1.3 1.3 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) All origins, both sexes..... 4,976 Male.... 3,164 Female..... 1,812 Hispanic, both sexes..... Male.... 2.4 Female..... Mexican, both sexes..... Male.... Female..... 5.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 4,625 2,910 Male.... 1,715 Female..... Non-Hispanic white, both sexes..... 3,811 2,365 Male.... 1,446 Female..... Non-Hispanic black, both sexes..... 2.4 Male.... Female..... 2.7 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Discharge of firearms, undetermined intent (Y22-Y24) All origins, both sexes..... 2.2 Male.... Female..... Hispanic, both sexes..... 2.0 Male.... Female..... - 1 Mexican, both sexes..... Male.... Female.... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male..... 1.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9) All origins, both sexes..... 4.741 Male.... 2.985 Female..... 1,756 2.0 Hispanic, both sexes..... 3.8 3.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... 5.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... 2.8 Female..... Non-Hispanic 1/, both sexes..... 4,413 2,753 Male.... 2.1 1,660 Female..... Non-Hispanic white, both sexes..... 3,645 2,243 Male.... 1,402 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 - 45-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years years years years years years years years years vears Operations of war and their sequelae (Y36,Y89.1) All origins, both sexes..... 3 Male.... 4 3 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 2.7 Non-Hispanic 1/, both sexes..... 4 27 Male.... 4 1 3 Female..... Non-Hispanic white, both sexes..... 25 3 2 25 Male.... 3 1 2 Female..... Non-Hispanic black, both sexes..... 2 1 1 Male.... 1 1 Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Complications of medical and surgical care (Y40-Y84,Y88) All origins, both sexes..... 2,883 Male.... 1,364 Female..... 1.519 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 2,708 Non-Hispanic 1/, both sexes..... 1,285 Male.... 1.0 1.3 2.3 1,423 Female..... 2. Non-Hispanic white, both sexes..... 2,219 1,084 Male.... 1,135 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 non-Hispanic population, and sex ages 1 year years Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 3/ All origins, both sexes..... 21,081 1,060 2,299 3.422 Male.... 15,906 1,679 2,482 Female..... 5.175 Hispanic, both sexes..... 2,406 Male.... 2,056 Female..... Mexican, both sexes..... 1.586 2.0 Male.... 1,372 2.8 Female..... Puerto Rican, both sexes..... 2.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. ΩN 1 Q Male.... Female Non-Hispanic 1/, both sexes..... 18,567 1,972 2,971 13,761 Male.... 1,401 2.092 4,806 Female..... Non-Hispanic white, both sexes..... 15,418 1,625 2,471 11,428 Male.... 1,164 1.737 3,990 Female..... Non-Hispanic black, both sexes..... 2.318 1.754 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 ages 1 year years years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 3/

All origins, both sexes	30,711	30	24	12	51	814	2,099	2,378	2,683	3,870	5,358	5,405
Male	19,362	11	16	4	21	599	1,588	1,734	1,820	2,490	3,247	3,281
Female	11,349	19	8	8	30	215	511	644	863	1,380	2,111	2,124
Hispanic, both sexes	2,257	-	1	2	5	65	168	234	248	351	395	320
Male	1,671	-	1	1	2	48	127	188	198	266	290	223
Female	586	-	-	1	3	17	41	46	50	85	105	97
Mexican, both sexes	1,212	-	1	1	3	43	85	139	152	173	219	175
Male	933	-	1	-	2	32	65	121	121	134	165	128
Female	279	-	-	1	1	11	20	18	31	39	54	47
Puerto Rican, both sexes	432	-	-	-	1	5	30	44	42	86	66	64
Male	328	-	-	-	-	3	29	35	32	63	53	41
Female	104	-	-	-	1	2	1	9	10	23	13	23
Cuban, both sexes	77	-	-	-	-	4	1	3	5	8	9	10
Male	46	-	-	-	-	4	-	2	5	6	4	6
Female	31	-	-	-	-	-	1	1	-	2	5	4
Central and South American, both sexes.	124	-	-	-	-	5	19	12	15	15	23	9
Male	92	-	-	-	-	3	12	8	13	14	19	7
Female	32	-	-	-	-	2	7	4	2	1	4	2
Other and Unknown Hispanic, both sexes.	412	-	-	1	1	8	33	36	34	69	78	62
Male	272	-	-	1	-	6	21	22	27	49	49	41
Female	140	-	-	-	1	2	12	14	7	20	29	21
Non-Hispanic 1/, both sexes	28,339	29	22	10	46	745	1,929	2,141	2,427	3,501	4,946	5,061
Male	17,605	11	14	3	19	548	1,460	1,545	1,615	2,209	2,945	3,040
Female	10,734	18	8	7	27	197	469	596	812	1,292	2,001	2,021
Non-Hispanic white, both sexes	24,201	18	13	6	35	695	1,782	1,929	2,097	2,972	4,253	4,207
Male	14,952	7	6	3	17	514	1,357	1,399	1,409	1,893	2,526	2,532
Female	9,249	11	7	3	18	181	425	530	688	1,079	1,727	1,675
Non-Hispanic black, both sexes	3,577	11	9	4	8	31	106	168	261	448	614	774
Male	2,309	4	8	-	2	22	74	117	169	265	368	458
Female	1,268	7	1	4	6	9	32	51	92	183	246	316
Origin not stated 2/, both sexes	115	1	1	-	-	4	2	3	8	18	17	24
Male	86	-	1	-	-	3	1	1	7	15	12	18
Female	29	1	-	-	-	1	1	2	1	3	5	6

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for Δ11 Under 1 – 4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40 - 4445-49 non-Hispanic population, and sex ages 1 year years years years years years years years years vears vears Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 3/ All origins, both sexes..... 29.569 2,494 4.046 3,304 2.717 2.364 2,622 2,499 Male.... 25,498 2,202 3,693 2,979 2,323 2,013 2,117 2,014 Female..... 4.071 Hispanic, both sexes..... 3,278 Male.... 2,973 Female..... 2.0 Mexican, both sexes..... 2.317 Male.... 2.132 2.0 Female..... 2.4 Puerto Rican, both sexes..... 2.4 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. an 5.8 1 2 Male.... Female..... Non-Hispanic 1/, both sexes..... 26,189 1,961 3,243 2,703 2,329 2,099 2,391 2,353 22,436 1,768 Male.... 1,713 2,937 2,421 1,976 1,925 1.886 3,753 Female..... Non-Hispanic white, both sexes..... 18,200 1,359 1,171 1,309 1,430 1,872 1,969 15,283 1,185 1,070 Male.... 1,006 1,181 1,476 1.573 2,917 Female..... Non-Hispanic black, both sexes..... 7.347 1,004 1,799 1,453 6,620 1,675 1,350 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
All causes												
All origins, both sexes	99,770	121,163	143,534	171,984	227,682	310,746	373,484	338,140	227,661	87,446	18,526	346
	62,649	73,400	84,632	98,455	124,436	160,308	173,361	133,667	70,168	19,820	2,995	274
	37,121	47,763	58,902	73,529	103,246	150,438	200,123	204,473	157,493	67,626	15,531	72
	6,725	7,248	7,950	9,231	11,189	13,454	13,425	9,835	6,766	2,703	633	48
Male	4,467	4,438	4,808	5,206	6,073	6,928	6,469	3,988	2,177	749	176	46
Female	2,258	2,810	3,142	4,025	5,116	6,526	6,956	5,847	4,589	1,954	457	
Mexican, both sexes	3,824	4,167	4,415	5,122	6,141	7,306	7,123	4,714	3,133	1,275	316	20
	2,551	2,527	2,662	2,877	3,280	3,734	3,532	2,009	1,124	390	99	19
	1,273	1,640	1,753	2,245	2,861	3,572	3,591	2,705	2,009	885	217	1
Puerto Rican, both sexes	1,034	1,185	1,235	1,289	1,475	1,639	1,592	1,154	751	282	65	10
	690	734	737	729	796	837	718	409	230	77	15	9
	344	451	498	560	679	802	874	745	521	205	50	1
Cuban, both sexes	395 287 108	403 268 135	642 428 214	922 578 344	1,330 843 487	1,831 1,053 778	2,019 986 1,033	1,848 753 1,095	1,369 396 973	574 136 438	111 20 91	-
Central and South American, both sexes. Male	522 333 189	565 328 237	635 356 279	691 354 337	786 371 415	859 367 492	750 294 456	700 256 444	430 100 330	141 30 111	33 8 25	2 2
Female Other and Unknown Hispanic, both sexes. Male	950 606	928 581	1,023 625	1,207 668	1,457 783	1,819 937	1,941 939	1,419 561	1,083	431 116	108 34	16 16
Female  Non-Hispanic 1/, both sexes  Male	344 92,672 57,905	347 113,511 68,678	398 135,213 79,556	539 162,309 92,948	674 215,957 118,031	882 296,698 153,064	1,002 359,434 166,586	858 327,805 129,490	756 220,537 67,875	315 84,605 19,036	74 17,862 2,814	151 112
Female  Non-Hispanic white, both sexes  Male	34,767	44,833	55,657	69,361	97,926	143,634	192,848	198,315	152,662	65,569	15,048	39
	69,774	89,100	109,742	133,916	183,690	259,718	322,736	299,301	201,488	77,094	15,499	113
	44,276	54,247	64,924	77,307	101,335	135,156	150,596	118,869	61,968	17,161	2,359	84
Female Non-Hispanic black, both sexes Male	25,498	34,853	44,818	56,609	82,355	124,562	172,140	180,432	139,520	59,933	13,140	29
	20,036	21,096	21,902	24,048	27,078	30,776	29,972	22,966	15,615	6,267	2,029	34
	11,954	12,498	12,552	13,228	14,000	14,666	12,728	8,181	4,480	1,395	359	25
FemaleOrigin not stated 2/, both sexes	8,082	8,598	9,350	10,820	13,078	16,110	17,244	14,785	11,135	4,872	1,670	9
	373	404	371	444	536	594	625	500	358	138	31	147
Male	277	284	268	301	332	316	306	189	116	35	26	116
Female	96	120	103	143	204	278	319	311	242	103		31

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	
Salmonella infections (A01-A02)												
All origins, both sexes	1	2	2	4	3	4	3	6	_	_	_	_
Male	1	2	2	2	1	4	2	4	-	-	-	-
Female	-	-	-	2	2	-	1	2	-	_	-	-
Hispanic, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mexican, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	_	-	-	-	-	-	_	-	-
Male	-	-	-	_	-	-	-	-	-	_	-	-
Female	-	-	-	_	-	-	-	-	-	_	-	-
Non-Hispanic 1/, both sexes	1	2	2	4	3	3	3	6	-	-	-	_
Male	1	2	2	2	1	3	2	4	-	_	-	-
Female	-	-	-	2	2	-	1	2	-	-	-	_
Non-Hispanic white, both sexes	1	1	1	1	2	2	3	5	-	-	-	_
Male	1	1	1	1	-	2	2	3	-	-	-	_
Female	-	-	-	_	2	-	1	2	-	_	-	-
Non-Hispanic black, both sexes	-	1	-	2	-	-	-	-	-	_	-	-
Male	-	1	-	1	-	-	-	-	-	_	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	_	-	-	-	-	-	_	-	-
Male	-	-	-	_	-	-	-	-	-	_	-	-
Female	-	-	-	_	-	-	-	-	-	_	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Shigellosis and amebiasis (A03,A06)												
All origins, both sexes	_	_	_	_	1	_	_	_	_	_	_	_
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	-	_	_	_
Female	_	_	_	_	_	_	_	_	-	_	_	_
Central and South American, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	_	_	_	_	1	_	_	_	_	_	_	_
Male	_	_	_	_	1	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic white, both sexes					1							
Male					1	_				_	_	
Female					_							_
Non-Hispanic black, both sexes										_	_	_
Male												
Female	_	_	_	_	_	_	_	_	_	_	_	_
Origin not stated 2/, both sexes	_	_	-	_	-	_	_	_	_	_	_	_
Male	_	_	_	_	-	_	_	_	_	_	_	_
	_	_	_	_	_	_	_	_	_	_	_	_
Female	-	_	_	_	_	-	-	-	_	_	_	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Certain other intestinal infections (A04,A07-A09) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... - 1 Male.... Female..... Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 2.5 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Tuberculosis (A16-A19)												
All origins, both sexes	47	35	44	55	82	81	84	68	37	11	_	_
Male	32	26	34	38	55	37	47	38	17	3	-	-
Female	15	9	10	17	27	44	37	30	20	8	-	-
Hispanic, both sexes	7	6	8	8	15	15	11	9	3	1	-	-
Male	3	4	7	4	12	8	9	5	2	-	-	-
Female	4	2	1	4	3	7	2	4	1	1	-	-
Mexican, both sexes	4	5	6	7	9	12	5	6	1	1	-	-
Male	2	4	5	4	6	7	4	4	1	-	-	-
Female	2	1	1	3	3	5	1	2	-	1	-	-
Puerto Rican, both sexes	1	-	1	-	2	_	1	_	_	-	-	-
Male	-	-	1	-	2	_	-	_	_	-	-	-
Female	1	-	-	-	-	_	1	_	_	-	-	-
Cuban, both sexes	-	-	1	-	1	1	3	_	_	-	-	-
Male	-	-	1	-	1	_	3	_	_	-	-	-
Female	-	-	_	-	-	1	-	-	-	-	-	-
Central and South American, both sexes.	-	1	_	1	-	-	1	2	1	-	-	-
Male	_	_	_	_	_	_	1	_	_	_	_	_
Female	_	1	_	1	_	_	_	2	1	_	_	_
Other and Unknown Hispanic, both sexes.	2	_	_	_	3	2	1	1	1	_	_	_
Male	1	_	_	_	3	1	1	1	1	_	_	_
Female	1	_	_	_	_	1	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	40	29	36	46	67	65	73	59	33	8	_	_
Male	29	22	27	33	43	28	38	3.3	1.5	2	_	_
Female	11	7	9	13	24	37	35	26	18	6	_	_
Non-Hispanic white, both sexes	14	13	15	21	34	40	48	38	21	4	_	_
Male	12	11	12	15	22	20	22	22	9	_	_	_
Female	2	2	3	6	12	20	26	16	12	4	_	_
Non-Hispanic black, both sexes	20	11	15	14	26	1.3	17	16		2	_	_
Male	12	9	11	9	16	3	12	8	3	_	_	_
Female	8	2	4	5	1.0	1.0	5	8	3	2	_	_
Origin not stated 2/, both sexes	-	_	_	1		1	_	-	1	2	_	_
Male	_	_	_	1	_	1	_	_	_	1	_	_
Female	_	_	_	_	_	_	_	_	1	1	_	_
									_	_		

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Respiratory tuberculosis (A16) All origins, both sexes..... Male.... 3.0 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female.... Non-Hispanic 1/, both sexes..... Male.... 2.4 2.9 1.5 Female..... Non-Hispanic white, both sexes..... Male..... 1.0 1.3 2.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other tuberculosis (A17-A19) All origins, both sexes..... Male.... Female..... 1.0 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Whooping cough (A37)												
All origins, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	-	_	_	-	-	-	-	-	_	_
Female	_	_	-	_	_	-	-	-	-	-	_	_
Hispanic, both sexes	_	_	-	_	_	-	-	-	-	-	_	_
Male	_	_	-	_	_	-	-	-	-	-	_	_
Female	_	_	-	_	_	-	-	-	-	-	_	_
Mexican, both sexes	-	-	-	-	_	-	-	-	-	_	_	-
Male	-	-	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	_	-	-	-	-	_	_	-
Puerto Rican, both sexes	-	-	-	-	_	-	-	-	-	_	_	-
Male	-	-	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	-	-	-	-	_	-	-	-	-	_	_	-
Male	-	-	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	_	-	-	-	-	_	_	-
Non-Hispanic white, both sexes	-	-	-	-	_	-	-	-	-	_	_	-
Male	-	-	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	_	-	-	-	-	-	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	_	-	-	-	-	-	-	_	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	s Age not s stated
Scarlet fever and erysipelas (A38,A46)												
All origins, both sexes	_	_	_	1	_	_	_	1	_	1	_	_
Male	_	_	_	1	_	_	_	Ξ.	_	_	_	_
Female	_	_	_	_	_	_	_	1	_	1	_	_
Hispanic, both sexes	-	_	_	1	-	-	-	-	-	-	-	-
	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	_	-	-	-	-	-	-	-	_
Mexican, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	_	_	-	_	_	-	-	-	_	-	-
Female	-	_	_	-	_	_	-	-	-	_	-	-
Puerto Rican, both sexes	-	_	_	-	_	_	-	-	-	_	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Non-Hispanic white, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Meningococcal infection (A39)												
All origins, both sexes	8	4	2	3	4	1	2	3	2	2	1	_
Male	7	2	2	2	1	-	1	-	-	-	_	-
Female	1	2	_	1	3	1	1	3	2	2	1	-
Hispanic, both sexes	2	-	_	1	-	-	-	-	1	-	_	-
Male	2	-	_	1	-	-	-	-	-	-	_	-
Female	-	-	_	-	-	-	-	-	1	-	_	-
Mexican, both sexes	1	-	_	1	-	-	-	-	-	-	_	-
Male	1	-	_	1	-	-	-	-	-	-	_	-
Female	-	-	_	-	-	-	-	-	-	-	_	-
Puerto Rican, both sexes	-	-	_	-	-	-	-	-	-	-	_	-
Male	-	-	_	-	-	-	-	-	-	-	_	-
Female	-	-	_	-	-	-	-	-	-	-	_	-
Cuban, both sexes	-	-	_	-	-	-	-	-	-	-	_	-
Male	-	-	_	-	-	-	-	-	-	-	_	-
Female	-	-	_	-	-	-	-	-	-	-	_	-
Central and South American, both sexes.	-	-	_	-	-	-	-	-	-	-	_	-
Male	-	-	_	-	-	-	-	-	-	-	_	-
Female	_	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	1	-	_	-	-	-	-	-	1	-	_	-
Male	1	-	_	-	-	-	-	-	-	-	_	-
Female	-	-	_	-	-	-	-	-	1	-	_	-
Non-Hispanic 1/, both sexes	6	4	2	2	4	1	2	3	1	2	1	_
Male	5	2	2	1	1	_	1	_	_	_	_	_
Female	1	2	_	1	3	1	1	3	1	2	1	_
Non-Hispanic white, both sexes	3	3	2	2	3	1	2	2	1	2	1	_
Male	2	1	2	1	_	_	1	_	_	_	_	_
Female	1	2	_	1	3	1	1	2	1	2	1	_
Non-Hispanic black, both sexes	2	1	_	_	ī	_	_	1	_	_	_	_
Male	2	1	_	_	1	_	_	_	_	_	_	_
Female	Ξ.	_	_	_	_	_	_	1	_	_	_	_
Origin not stated 2/, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Septicemia (A40-A41)												
All origins, both sexes	1,353 720 633	1,700 910 790	2,045 1,024 1,021	2,481 1,247 1,234	3,502 1,770 1,732	4,828	5,758 2,562	4,914 1,904 3,010	2,943 869 2,074	1,045 267 778	173 34 139	3 1 2
Female Hispanic, both sexes Male	83 54	102 51	114 58	137 74	182 80	2,520 214 98	3,196 223 97	145 62	100 29	39 14	6 -	- -
Female Mexican, both sexes Male	29 50 33	51 60 26	56 61 28	63 76 44	102 99 38	116 101 37	126 114 48	83 76 26	71 40 12	25 17 10	6 4 -	- - -
Female Puerto Rican, both sexes Male	17 12 5	34 17 11	33 25 13	32 25 12	61 25 14	64 46 26	66 29 14	50 20 10	28 14 4	7 4 -	4 1 -	- - -
FemaleCuban, both sexesMale	7 4 4	6 8 5	12 5 4	13 10 6	11 23 12	20 27 15	15 27 12	10 18 10	10 14 4	4 13 3	1 -	- - -
Female Central and South American, both sexes. Male	10	3 6 2	1 8 4	4 9	11 12 4	12 16 2	15 22 4	8 17 10	10 13 5	10	-	-
Female Other and Unknown Hispanic, both sexes.	, 3 7 5	4 11 7	4 15 9	4 17	8 23 12	14 24 18	18 31 19	7 14 6	8 19	2 2	1	-
Male  Female  Non-Hispanic 1/, both sexes	1,263	4 1,591	6 1,926	7 10 2,336	11 3,316	6 4,602	12 5,530	8 4,765	4 15 2,836	2 1,004	1 167	- - 2
Male  Female  Non-Hispanic white, both sexes	661 602 849	855 736 1,134	962 964 1,379	1,168 1,168 1,729	1,689 1,627 2,599	2,206 2,396 3,777	2,464 3,066 4,649	1,841 2,924 4,096	837 1,999 2,436	252 752 856	34 133 121	- 2 1
Male  Female  Non-Hispanic black, both sexes	457 392 377	610 524 423	689 690 483	875 854 556	1,338 1,261 647	1,837 1,940 746	2,077 2,572 788	1,611 2,485 588	715 1,721 363	217 639 133	25 96 44	- 1 1
Male Female Origin not stated 2/, both sexes	188 189	229 194	236 247	266 290	316 331 4	336 410 12	339 449	198 390 4	104 259	30 103 2	9 35	- 1 1
MaleFemale	5 2	4 3	4	5 3	1 3	4 8	1 4	1 3	3 4	1	- - -	1 -

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Syphilis (A50-A53) All origins, both sexes..... 3 2 2 3 3 4 8 3 1 Male.... 1 3 4 2 Female..... 2 1 1 2 4 2 6 1 Hispanic, both sexes..... 1 Male.... Female..... Mexican, both sexes..... 1 Male..... 1 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male..... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... 2 Non-Hispanic 1/, both sexes..... 3 4 8 3 Male.... 1 1 3 1 3 2. 2 2. 2 1 Female..... 1 1 2. 1 Non-Hispanic white, both sexes..... Male.... 1 1 1 1 Female..... 3 1 2 2 Non-Hispanic black, both sexes..... 1 2 1 3 3 2 5 2 1 Male..... 2 2 1 3 1 1 Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Acute poliomyelitis (A80)												
All origins, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	-	-	-	-	_	-	-	-	-	-	-	-
Female	-	-	-	_	-	-	-	-	-	-	-	-
Hispanic, both sexes	-	-	-	-	_	-	-	-	-	-	-	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	_	_	_	_	-	-	-	-	-	_	_
Mexican, both sexes	-	_	-	-	_	-	-	-	-	_	_	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	_	_	_	_	-	-	-	-	-	_	_
Puerto Rican, both sexes	-	_	_	_	_	-	-	-	-	-	_	_
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	_	-	-	-	-	-	-	-
Cuban, both sexes	-	_	_	_	_	-	-	-	-	-	_	_
Male	-	_	_	_	_	-	-	-	-	-	_	_
Female	-	_	_	_	_	-	-	-	-	-	_	_
Central and South American, both sexes.	-	_	_	_	_	-	-	-	-	-	_	_
Male	-	-	-	-	-	_	_	-	-	-	-	-
Female	-	-	-	-	-	_	_	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	_	_	-	-	-	-	-
Male	-	-	-	-	-	_	_	-	-	-	-	-
Female	-	-	-	-	-	_	_	-	-	-	-	-
Non-Hispanic 1/, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic white, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic black, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	-	_	_	_	_	_	_	_	_	_
Female	-	-	_	_	-	_	_	-	_	-	-	-
Origin not stated 2/, both sexes	_	_	-	-	-	-	-	-	-	_	_	-
Male	_	_	-	-	-	-	-	-	-	_	_	-
Female	_	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Arthropod-borne viral encephalitis (A83-A84,A85.2) All origins, both sexes..... 1 2 1 Male.... 1 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male..... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female Non-Hispanic 1/, both sexes..... Male.... 1 2 1 Female..... Non-Hispanic white, both sexes..... Male.... 1 2 Female..... Non-Hispanic black, both sexes..... 1 Male..... 1 Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Measles (B05)												
All origins, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	-	_	_	-	-	-	-	-	_	_
Female	_	_	-	_	_	-	-	-	-	-	_	_
Hispanic, both sexes	_	_	-	_	_	-	-	-	-	-	_	_
Male	_	_	-	_	_	-	-	-	-	-	_	_
Female	_	_	-	_	_	-	-	-	-	-	_	_
Mexican, both sexes	-	_	-	-	_	-	-	-	-	_	_	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	_	-	-	_	-	-	-	-	_	_	-
Puerto Rican, both sexes	-	_	-	-	_	-	-	-	-	_	_	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes	-	_	-	-	_	-	-	-	-	_	_	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	_	-	-	_	-	-	-	-	_	_	-
Non-Hispanic white, both sexes	-	_	-	-	_	-	-	-	-	_	_	-
Male	-	_	-	-	_	-	-	-	-	_	_	-
Female	-	_	-	-	_	-	-	-	-	-	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	_	-	-	-	-	-	-	_	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Viral hepatitis (B15-B19) All origins, both sexes..... 1,270 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... 1.8 Female..... 1.0 Puerto Rican, both sexes..... 3.0 1.8 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,122 Male.... Female..... Non-Hispanic white, both sexes..... Male.... 2.8 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Human immunodeficiency virus (HIV) disease (B20-B24) All origins, both sexes..... 1.868 1,038 Male.... 1,416 Female..... Hispanic, both sexes..... Male.... 1.0 Female..... 2.7 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male 2.0 Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,601 1,209 Male.... Female..... Non-Hispanic white, both sexes..... Male.... 2.2 Female..... Non-Hispanic black, both sexes..... 1,040 Male.... Female..... Origin not stated 2/, both sexes...... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Malaria (B50-B54)												
All origins, both sexes	1	_	_	1	_	_	_	_	_	_	_	_
Male	1	_	_	1	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Hispanic, both sexes	_	_	_	_	_	_	_	_	-	_	_	_
Male	_	_	_	_	_	_	_	_	-	_	_	_
Female	_	_	_	_	_	_	_	_	-	_	_	_
Mexican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	-	-	-	-	-	-	_	-	-	-	-	-
Female	-	-	-	-	-	-	_	-	-	-	-	-
Cuban, both sexes	-	-	-	-	-	-	_	-	-	-	-	-
Male	_	_	-	_	_	_	-	-	-	_	_	-
Female	_	_	-	_	_	_	-	-	-	_	_	-
Central and South American, both sexes.	_	_	-	_	_	_	-	-	-	_	_	-
Male	_	_	-	_	_	_	-	-	-	_	_	-
Female	_	_	-	_	_	_	-	-	-	_	_	_
Other and Unknown Hispanic, both sexes.	_	_	-	_	_	_	-	-	-	_	_	_
Male	-	_	-	_	_	-	_	-	-	-	_	_
Female	-	_	-	_	_	-	_	-	-	-	_	_
Non-Hispanic 1/, both sexes	1	-	-	1	_	-	-	-	-	_	-	-
Male	1	_	-	1	_	-	_	-	-	-	_	_
Female	-	_	-	_	_	-	_	-	-	-	_	_
Non-Hispanic white, both sexes	-	_	-	_	_	-	_	-	-	-	_	_
Male	-	_	-	_	_	-	_	-	-	-	_	_
Female	-	_	-	_	_	-	_	-	-	-	-	-
Non-Hispanic black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... 2.8 Female..... 2.8 Mexican, both sexes Male.... Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 3.0 3.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Malignant neoplasms (C00-C97)												
All origins, both sexes	30,330 16,037 14,293 1,787	43,127 23,196 19,931 2,168	53,829 29,542 24,287 2,491	63,503 35,065 28,438 2,847	75,914 41,422 34,492 3,149	86,865 46,582 40,283 3,281	79,220 40,152 39,068 2,689	50,851 23,831 27,020 1,454	22,844 9,325 13,519 700	5,855 2,050 3,805 201	795 221 574 23	8 6 2 3
Male Female	959 828	1,140 1,028	1,388 1,103	1,559 1,288	1,754 1,395	1,828 1,453	1,455 1,234	689 765	289 411	90 111	9 14	3 -
Mexican, both sexes	968 506 462	1,180 588 592	1,304 724 580	1,504 849 655	1,665 922 743	1,727 984 743	1,418 807 611	705 346 359	314 147 167	99 48 51	15 6 9	-
Puerto Rican, both sexes	248 151	335 186	370 213	356 187	382 215	353 197	288 146	140 57	89 32	24 12	2 1	2
FemaleCuban, both sexesMale	97 122 80	149 150 91	157 265 165	169 333 193	167 434 274	156 464 286	142 384 211	83 255 132	57 119 40	12 36 15	1 3 1	-
Female	42 182	59 208	100 232	140 261	160 226	178 272	173 178	123 134	79 53	21 12	2	-
MaleFemale	85 97	108 100	114 118	122 139	100 126	127 145	77 101	61 73	14 39	3 9 30	-	- - 1
Other and Unknown Hispanic, both sexes.  Male  Female	267 137 130	295 167 128	320 172 148	393 208 185	442 243 199	465 234 231	421 214 207	220 93 127	125 56 69	12 18	3 1 2	1 -
Non-Hispanic 1/, both sexes	28,485 15,040	40,858 21,992	51,235 28,086	60,537 33,431	72,628 39,591	83,447 44,685	76,415 38,640	49,322 23,106	22,109 9,020	5,644 1,955	772 212	4 2
Female. Non-Hispanic white, both sexes Male	13,445 22,280 11,793	18,866 33,363 17,937	23,149 43,037 23,531	27,106 51,307 28,376	33,037 62,841 34,234	38,762 73,822 39,534	37,775 68,639 34,644	26,216 44,719 20,976	13,089 19,877 8,017	3,689 5,042 1,713	560 645 172	2 3 1
Female	10,487	15,426 6,267	19,506	22,931 7,698	28,607 8,014	34,288 7,921	33,995	23,743	11,860	3,329	473 115	2
Male  Female  Origin not stated 2/, both sexes	2,770 2,425 58	3,422 2,845 101	3,854 3,028 103	4,230 3,468 119	4,412 3,602 137	4,186 3,735 137	3,236 3,087 116	1,719 2,064 75	792 1,039 35	191 314 10	35 80	1 - 1
MaleFemale	38 20	64 37	68 35	75 44	77 60	69 68	57 59	36 39	16 19	5	- - -	1 -

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Malignant neoplasms of lip, oral cavity and pharynx (COO-C14) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 3.8 3.0 1.0 Male.... Female..... 1.0 Mexican, both sexes..... 2.0 1.0 Male.... Female..... \_ Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.5 Female..... Non-Hispanic white, both sexes..... Male.... 2.3 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Malignant neoplasm of esophagus (C15) All origins, both sexes..... 1,334 1.657 1.739 1,815 1,928 1.539 Male.... 1,130 1.372 1,412 1,405 1,450 1.096 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... 2.0 Female..... Puerto Rican, both sexes..... 1.8 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 a Male.... Female..... Non-Hispanic 1/, both sexes..... 1,289 1,588 1,671 1,747 1,867 1,493 1,093 Male.... 1,313 1,356 1,354 1,405 1,061 2.0 Female..... Non-Hispanic white, both sexes..... 1,063 1,318 1,427 1,507 1,681 1,355 1,172 Male.... 1,114 1,191 1,277 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Malignant neoplasm of stomach (C16) 1,005 1,202 All origins, both sexes..... 1,550 1,733 1,682 1,255 Male.... 1,041 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 3.8 Male.... Female..... 3.0 Puerto Rican, both sexes..... 1.0 1.0 Male.... Female..... Cuban, both sexes..... 1.0 1.0 Male Female..... 2.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 1,063 1,376 1,566 1,539 1,175 Male.... Female..... Non-Hispanic white, both sexes..... 1,010 1,211 1,225 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasms of colon, rectum and anus (C18-C21) All origins, both sexes..... 2,636 3,691 4.517 5,297 6,657 8,245 8,469 6,337 3,474 1,023 Male.... 1,531 2,161 2,624 3,074 3.718 4,307 3,882 2,464 1.092 2.0 Female..... 1,105 1,530 1,893 2,223 2,939 3,938 4,587 3,873 2,382 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... 2.0 Female..... Cuban, both sexes..... Male Female..... 2.4 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 2,457 3,468 4,260 5,015 6,311 7,915 8,173 6,157 3,366 Male.... 1,431 2,032 2,467 2,912 3,530 4,119 3,736 2,388 1,058 2.0 1,026 1,436 1,793 2,103 2,781 3,796 4,437 3,769 2,308 Female..... Non-Hispanic white, both sexes..... 1,879 2,769 3,444 4,106 5,308 6,877 7,243 5,538 3,025 3,004 3,588 Male.... 1,114 1,660 2,023 2,417 3,331 2,172 1,109 1,421 1,689 2,304 3,289 3,912 3,366 2,075 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasms of liver and intrahepatic bile ducts (C22) All origins, both sexes..... 1,476 1.554 1.569 1,654 2,016 2.136 1,923 1,030 Male.... 1,216 1,205 1,113 1,092 1,290 1,202 1,025 Female..... Hispanic, both sexes..... Male.... 1.0 Female..... Mexican, both sexes..... Male.... Female..... 2.0 Puerto Rican, both sexes..... 2.8 2.1 Male.... Female..... Cuban, both sexes..... 2.0 Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. 2.8 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,299 1,371 1,371 1,469 1,803 1,939 1,779 1,099 Male.... 1,070 1,054 1,166 2.4 Female..... Non-Hispanic white, both sexes..... 1,019 1,113 1,414 1,599 1,541 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... 5.8 Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Malignant neoplasm of pancreas (C25)												
All origins, both sexes	1,715	2,509	3,247	3,717	4,461	5,106	4,747	3,056	1,323	317	53	1
Male	1,051	1,490	1,887	2,093	2,337	2,407	2,009	1,090	371	66	11	-
Female	664	1,019	1,360	1,624	2,124	2,699	2,738	1,966	952	251	42	1
Hispanic, both sexes	111	138	183	232	216	229	201	88	45	11	3	-
Male	59	76	108	120	120	101	90	25	15	1	-	-
Female	52	62	75	112	96	128	111	63	30	10	3	-
Mexican, both sexes	60	80	95	139	115	105	105	33	18	4	1	-
Male	28	40	60	69	63	43	48	9	8	1	-	-
Female	32	40	35	70	52	62	57	24	10	3	1	_
Puerto Rican, both sexes	9	15	22	23	26	26	23	7	5	1	1	-
Male	7	11	15	14	15	14	10	2	2	-	-	-
Female	2	4	7	9	11	12	13	5	3	1	1	-
Cuban, both sexes	7	10	18	24	27	27	22	13	12	3	_	_
Male	6	8	10	15	16	16	13	5	2	_	_	_
Female	1	2	8	9	11	11	9	8	10	3	_	_
Central and South American, both sexes.	9	14	19	23	12	23	19	1.3	3	_	_	_
Male	7	9	9	12	3	5	5	3	1	_	_	_
Female	2	5	10	11	9	18	14	10	2	_	_	_
Other and Unknown Hispanic, both sexes.	26	19	29	23	36	48	32	22	7	3	1	_
Male	11	8	14	10	23	23	14	6	2	_	_	_
Female	15	11	15	13	13	25	18	16	5	3	1	_
Non-Hispanic 1/, both sexes	1,602	2,363	3,057	3,477	4,239	4,870	4,535	2,965	1,276	306	50	1
Male	991	1,408	1,772	1,968	2,212	2,302	1,915	1,064	355	65	11	_
Female	611	955	1,285	1,509	2,027	2,568	2,620	1,901	921	241	39	1
Non-Hispanic white, both sexes	1,248	1,905	2,522	2,916	3,580	4,259	4,046	2,658	1,136	271	39	1
Male	787	1,143	1,490	1,667	1,883	2,052	1,742	977	317	56	8	_
Female	461	762	1,032	1,249	1,697	2,032	2,304	1,681	819	215	31	1
Non-Hispanic black, both sexes	293	372	439	461	534	495	371	250	113	213	9	_
Male	162	221	237	248	270	195	125	62	29	7	3	
	131	151	202	213	264	300	246	188	84	21	5 6	_
Female	2	121	202 7	213 8	204 6	300		199		21	б	-
Origin not stated 2/, both sexes	_	Ü		8	-	/	11		2	_	_	_
Male	1	6	7	5	5	4	4	1	1	-	-	_
Female	1	2	-	3	1	3	./	2	1	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Malignant neoplasm of larynx (C32) All origins, both sexes..... 2.2 Male.... Female..... Hispanic, both sexes..... 2.0 3.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.47 Female..... Non-Hispanic white, both sexes..... Male.... 2.44 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasms of trachea, bronchus and lung (C33-C34) All origins, both sexes..... 7.884 12,846 18,240 22,552 26,477 27,608 20,856 10,325 3,418 Male.... 4,751 7,663 10,777 13,243 15,210 15,754 11,238 5,246 1,537 Female..... 3,133 5,183 7,463 9,309 11,267 11,854 9,618 5,079 1,881 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 3 0 aa 1.8 Male.... Female..... Non-Hispanic 1/, both sexes..... 7,634 12,470 17,764 21,915 25,714 26,834 20,330 10,033 3,311 Male.... 4,577 7,414 10,453 12,811 14,716 15,250 10,916 5,086 1,493 3,057 5,056 7,311 10,998 11,584 9,414 4,947 1,818 Female..... 9.104 Non-Hispanic white, both sexes..... 6,056 10,303 15,270 18,980 22,812 24,103 18,412 9,197 2,999 11,035 13,633 1,338 Male.... 3,559 6,040 8,817 12,946 9,832 4,662 2,497 4,263 6,453 7,945 9,866 10,470 8,580 4,535 1,661 Female..... Non-Hispanic black, both sexes..... 1,386 1.892 2,166 2,467 2,343 2,237 1,554 1,210 1,436 1,506 1,434 1,306 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Malignant melanoma of skin (C43) All origins, both sexes..... 1,008 Male.... Female..... Hispanic, both sexes..... 2.0 Male.... Female..... Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years and over stated vears vears vears vears Malignant neoplasm of breast (C50) 4,220 4,792 4,579 All origins, both sexes..... 3,639 4.312 4,162 4,065 3,473 1.823 Male.... 2.0 Female..... 3,619 4,282 4,125 4,022 4,172 4.727 4,535 3,440 1,813 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... - 1 Female..... Puerto Rican, both sexes..... 3.0 Male.... 2.7 Female..... Cuban, both sexes..... Male 2.7 Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 2.7 2.7 Male.... Female..... Non-Hispanic 1/, both sexes..... 3,405 4,062 3,955 3,852 4,045 4,624 4,453 3,372 1,781 Male.... 3,388 4,034 3,919 3,809 4,003 4,559 4,411 3,340 1,772 Female..... Non-Hispanic white, both sexes..... 2,550 3,270 3,253 3,268 3,495 4,115 4,042 3,024 1,628 Male.... 2.5 2,539 3,248 3,225 3,231 3,462 4,054 4,005 2,999 1,620 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Malignant neoplasm of cervix uteri (C53) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... 2.8 2.6 2.0 Mexican, both sexes..... 2.7 Male.... Female..... Puerto Rican, both sexes..... Male.... . . . Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male . . . Female..... Other and Unknown Hispanic, both sexes. Male.... . . . Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55) All origins, both sexes..... 1,025 Male.... Female..... 1,025 Hispanic, both sexes..... 3.0 Male.... . . . Female..... Mexican, both sexes..... 1.0 Male.... Female..... 2.2 Puerto Rican, both sexes..... Male.... . . . . . . Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male . . . . . . . . . . . . Female..... Other and Unknown Hispanic, both sexes. Male.... . . . Female..... Non-Hispanic 1/, both sexes..... Male.... . . . Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Malignant neoplasm of ovary (C56) All origins, both sexes..... 1.007 1,301 1,493 1,775 1,918 2,142 2,023 1,208 510 103 10 Male.... Female..... 1.007 1,301 1,493 1,775 1.918 2,142 2,023 1,208 510 103 10 Hispanic, both sexes..... 54 72 73 80 84 81 62 38 18 3 Male.... Female..... 54 72 73 80 84 81 62 3.8 18 3 Mexican, both sexes..... 31 51 40 38 54 39 36 2.0 6 2 Male.... Female..... 21 51 40 38 54 39 36 20 2 6 Puerto Rican, both sexes..... 5 7 7 12 7 5 2 4 15 1 Male.... 15 7 7 12 7 5 2 1 Female..... Cuban, both sexes..... 1 4 5 8 1.0 8 7 7 5 Male . . . 8 10 7 5 Female..... 5 8 7 1 Central and South American, both sexes. 10 6 9 3 4 5 3 Male . . . . . . Female..... 10 9 5 3 Other and Unknown Hispanic, both sexes. 7 7 6 18 7 5 1.8 10 2 Male.... . . . Female..... 6 18 10 18 7 5 2 Non-Hispanic 1/, both sexes..... 951 1,226 1,418 1,692 1,830 2,060 1,960 1,169 491 100 10 Male.... 951 1,226 1,418 1,692 1,830 2,060 1,960 1,169 491 100 1.0 Female..... Non-Hispanic white, both sexes..... 798 1,053 1,259 1,479 1,624 1,875 1,826 1,078 88 7 Male.... 798 1,053 1,259 1,479 1,624 1,875 1,826 1,078 460 88 Female..... Non-Hispanic black, both sexes..... 110 124 124 176 170 155 117 75 26 9 3 Male.... Female..... 124 124 176 170 155 117 75 26 Origin not stated 2/, both sexes..... 2 3 2 3 4 1 1 1 1 Male.... Female..... 3 3 1

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasm of prostate (C61) All origins, both sexes..... 332 688 1.356 2.196 3,515 5.444 6.549 5,328 2.630 747 87 2 Male.... 332 688 1,356 2,196 3,515 5,444 6,549 5,328 2,630 747 87 2 Female..... Hispanic, both sexes..... 19 39 70 102 182 266 280 158 2.8 86 2 1 Male.... 19 39 70 102 182 266 280 158 86 28 1 Female..... Mexican, both sexes..... 13 19 37 52 103 129 161 80 41 12 1 Male.... 13 19 37 52 103 129 161 80 41 12 1 Female..... . . . . . . Puerto Rican, both sexes..... 25 2.0 6 1.0 15 2.4 19 12 1 Male.... 1 6 10 15 24 25 20 19 12 7 1 Female..... . . . Cuban, both sexes..... 12 50 47 2.8 11 5 3 1.0 2.5 Male 1 3 12 10 25 50 47 28 11 5 Female..... . . . . . . Central and South American, both sexes. 4 4 5 11 13 23 16 14 5 Male 4 4 5 11 13 23 16 14 5 Female..... . . . Other and Unknown Hispanic, both sexes. 39 7 6 1 4 17 36 17 17 4 1 Male.... 7 6 14 17 39 36 17 17 4 1 Female..... Non-Hispanic 1/, both sexes..... 313 647 1,284 2,090 3,328 5,169 6,261 5,162 2,541 719 85 Male.... 313 647 1,284 2,090 3,328 5,169 6,261 5,162 2,541 719 85 Female..... . . . Non-Hispanic white, both sexes..... 204 463 948 1,576 2,571 4,145 5,243 4,459 2,156 602 60 2,156 602 60 Male.... 204 463 948 1,576 2,571 4,145 5,243 4,459 Female..... . . . Non-Hispanic black, both sexes..... 105 175 318 484 703 941 913 625 341 101 23 105 175 318 484 703 941 913 625 341 101 23 Male.... Female..... Origin not stated 2/, both sexes..... 2 2 4 5 9 8 8 3 1 2 2 4 5 9 8 8 3 1 Male.... Female..... .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Malignant neoplasms of kidney and renal pelvis (C64-C65) All origins, both sexes..... 1.114 1,420 1,451 1,624 1,819 1,655 1,003 Male.... 1,054 1,060 Female..... Hispanic, both sexes..... Male.... Female..... 2.7 1.8 2.8 2.0 Mexican, both sexes..... Male.... 2.8 Female..... 2.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1.0 1.0 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1 0 a Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 1,032 1,314 1,351 1,540 1,736 1,574 1,008 Male.... Female..... Non-Hispanic white, both sexes..... 1,163 1,193 1,358 1,553 1,446 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasm of bladder (C67) All origins, both sexes..... 1,143 1,610 2,307 2,540 1,976 1.012 Male.... 1,162 1,646 1,730 1,275 Female..... Hispanic, both sexes..... 3.8 Male.... Female..... 1.0 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... 1.0 1.0 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,097 1,559 2,234 2,462 1,935 Male.... 1,128 1,606 1,685 1,250 1.5 Female..... Non-Hispanic white, both sexes..... 1,004 1,416 2,074 2,301 1,802 1,045 1,512 Male.... 1,604 1,180 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not years and over stated

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

					, , , , , ,	•						
All origins, both sexes	1,070	1,378	1,372	1,392	1,503	1,407	1,136	564	204	34	5	_
Male	675	832	821	741	810	717	520	219	60	11	2	_
Female	395	546	551	651	693	690	616	345	144	23	3	_
Hispanic, both sexes	57	64	67	63	51	50	35	16	11	3	-	_
	37	34	42	32	25	29	14	5	3	1	-	-
Female	20	30	25	31	26	21	21	11	8	2	-	-
Mexican, both sexes	25	33	39	33	23	21	18	9	7	3	-	-
Male	15	15	25	19	10	12	10	4	2	1	-	-
Female	10	18	14	14	13	9	8	5	5	2	-	-
Puerto Rican, both sexes	5	8	8	3	5	6	4	1	-	-	-	-
Male	4	5	5	1	2	4	1	_	-	-	-	-
Female	1	3	3	2	3	2	3	1	-	-	-	-
Cuban, both sexes	5	5	9	11	8	11	8	2	1	-	-	-
Male	4	1	7	6	6	8	2	1	-	-	-	-
Female	1	4	2	5	2	3	6	1	1	-	-	-
Central and South American, both sexes.	12	6	4	8	3	4	2	2	1	-	-	-
Male	9	6	3	2	-	2	-	-	1	-	-	-
Female	3	-	1	6	3	2	2	2	-	-	-	-
Other and Unknown Hispanic, both sexes.	10	12	7	8	12	8	3	2	2	-	-	-
Male	5	7	2	4	7	3	1	-	-	-	-	-
Female	5	5	5	4	5	5	2	2	2	-	-	-
Non-Hispanic 1/, both sexes	1,012	1,309	1,303	1,326	1,451	1,355	1,100	548	193	31	5	-
Male	637	795	778	708	785	688	506	214	57	10	2	-
Female	375	514	525	618	666	667	594	334	136	21	3	-
Non-Hispanic white, both sexes	912	1,215	1,212	1,234	1,364	1,274	1,047	520	180	26	3	-
Male	571	736	726	668	748	643	479	208	54	8	2	-
Female	341	479	486	566	616	631	568	312	126	18	1	-
Non-Hispanic black, both sexes	77	69	72	67	67	58	40	21	11	3	2	-
Male	51	45	42	32	25	31	20	4	3	1	-	-
Female	26	24	30	35	42	27	20	17	8	2	2	-
Origin not stated 2/, both sexes	1	5	2	3	1	2	1	-	-	-	-	-
Male	1	3	1	1	-	-	-	-	-	-	-	-
Female	-	2	1	2	1	2	1	-	-	-	-	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) All origins, both sexes..... 2,264 3,323 4,168 5,423 7,165 9,090 9,042 5,925 2.633 Male.... 1,372 1,993 2,504 3,244 4,124 4,915 4.547 2,661 Female..... 1,330 1,664 2,179 3,041 4,175 4,495 3,264 1,638 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 2.7 Puerto Rican, both sexes..... 2.8 3.0 Male.... Female..... Cuban, both sexes..... 1.0 2.0 Male Female..... 2.0 Central and South American, both sexes. 2.1 Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 2,085 3,104 3,934 5,134 6,825 8,725 8,730 5,779 2,566 1,249 4,726 Male.... 1,876 2,361 3,081 3,939 4,401 2,595 1,228 1,573 2,053 2,886 3,999 4,329 3,184 1,592 Female..... Non-Hispanic white, both sexes..... 1,654 2,574 3,359 4,431 5,963 7,877 8,043 5,375 2,357 1,588 Male.... 2,042 2,674 3,486 4,319 4,087 2,438 1,317 1,757 2,477 3,558 3,956 2,937 1,459 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Hodgkin's disease (C81) All origins, both sexes..... 2.4 Male.... Female..... 3.0 Hispanic, both sexes..... - 1 1.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Non-Hodgkin's lymphoma (C82-C85) All origins, both sexes..... 1.277 1,589 2,100 2,720 3,477 3,603 2,359 1.036 Male.... 1,235 1,503 1,782 1.729 1,012 Female..... 1,217 1,695 1,874 1,347 Hispanic, both sexes..... 2.8 Male.... Female..... 2.8 Mexican, both sexes..... 5.8 1.0 Male.... 2.8 2.0 Female..... 2.8 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1.0 Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. Ω a Male.... Female..... Non-Hispanic 1/, both sexes..... 1,186 1,508 1,989 2,586 3,325 3,468 2,299 1,005 1,672 Male.... 1,174 1,427 1,705 1,159 1,620 1,796 1,317 Female..... Non-Hispanic white, both sexes..... 1,024 1,336 1,789 2,353 3,085 3,294 2,182 1,054 1,307 1,593 1,593 Male.... 1,046 1,492 1,701 1,245 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

Cause of death (Based on the International Classification of												
Diseases, Tenth Revision, 1992),												
Hispanic origin, race for	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years	Age not
non-Hispanic population, and sex	years	and over										
	1	1	2	1	1	2	2	1	2	2		
Leukemia (C91-C95)												
All origins, both sexes	812	1,176	1,512	1,985	2,709	3,377	3,420	2,349	1,165	276	40	_
Male	479	720	943	1,237	1,666	1,985	1,836	1,112	453	95	9	_
Female	333	456	569	748	1,043	1,392	1,584	1,237	712	181	31	_
Hispanic, both sexes	69	66	73	97	102	105	91	45	24	6	1	_
Male	43	38	48	58	56	59	47	21	7	2	1	_
Female	26	28	25	39	46	46	44	24	17	4	_	_
Mexican, both sexes	37	32	35	53	49	51	46	21	11	4	1	_
Male	26	17	25	32	28	30	22	7	3	2	1	_
Female	11	15	10	21	21	21	24	14	8	2	-	-
Puerto Rican, both sexes	8	12	10	9	9	11	8	4	_	1	-	-
Male	4	8	5	7	3	6	5	3	_	-	-	-
Female	4	4	5	2	6	5	3	1	-	1	-	-
Cuban, both sexes	3	3	8	15	17	15	16	10	5	-	-	-
Male	2	1	4	9	11	12	8	4	1	-	-	-
Female	1	2	4	6	6	3	8	6	4	-	-	-
Central and South American, both sexes.	7	7	5	10	9	12	8	5	1	-	-	-
Male	6	5	4	5	5	5	4	3	-	-	-	-
Female	1	2	1	5	4	7	4	2	1	-	-	-
Other and Unknown Hispanic, both sexes.	14	12	15	10	18	16	13	5	7	1	-	-
Male	5	7	10	5	9	6	8	4	3	-	-	_
Female	9	5	5	5	9	10	5	1	4	1	-	_
Non-Hispanic 1/, both sexes	741	1,107	1,436	1,885	2,602	3,270	3,326	2,303	1,138	268	39	-
Male	434	681	893	1,179	1,608	1,924	1,787	1,091	445	92	8	-
Female	307	426	543	706	994	1,346	1,539	1,212	693	176	31	-
Non-Hispanic white, both sexes	606	951	1,243	1,654	2,302	2,991	3,083	2,155	1,053	252	36	-
Male	350	597	786	1,037	1,429	1,778	1,672	1,034	411	85	7	-
Female	256	354	457	617	873	1,213	1,411	1,121	642	167	29	_
Non-Hispanic black, both sexes	115	128	166	184	245	216	189	125	68	15	3	_
Male	70	70	91	114	148	104	81	49	24	6	1	_
Female	45	58	75	70	97	112	108	76	44	9	2	_
Origin not stated 2/, both sexes	2	3	3	3	5	2	3	1	3	2	_	_
Male	2	1	2	-	2	2	2	-	1	1	_	_
Female	-	2	1	3	3	-	1	1	2	1	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Multiple myeloma and immunoproliferative neoplasms (C88,C90) All origins, both sexes..... 1,229 1,611 2,096 1,884 1,130 Male.... 1,083 Female..... 1,013 Hispanic, both sexes..... 2.8 Male.... Female..... 2.6 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,160 1,521 2,000 1,807 1,100 1,036 Male.... Female..... Non-Hispanic white, both sexes..... 1,200 1,679 1,546 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96) All origins, both sexes..... 4 8 4 5 7 9 17 6 5 1 Male.... 3 8 4 1 Female..... 5 2 3 6 9 2 3 Hispanic, both sexes..... 1 Male.... Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1 Male..... 1 Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 8 4 17 Male.... 3 2. 3 3 8 4 2. 1 9 3 Female..... 2. 3 2 Non-Hispanic white, both sexes..... 14 4 Male.... 3 3 2 3 3 7 2 Female..... 3 5 5 2 2 2. 3 Non-Hispanic black, both sexes..... 1 2 1 2 1 1 Male.... 1 Female..... Origin not stated 2/, both sexes..... Male.....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97) All origins, both sexes..... 3,159 4,549 5,448 6,434 7,753 9.366 9.183 6,396 3,047 Male.... 1,792 2,597 3,082 3,649 4,149 4,890 4,391 2,646 1.054 Female..... 1,367 1,952 2,366 2,785 3,604 4,476 4,792 3,750 1,993 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... 2.7 Female..... Cuban, both sexes..... 2.0 Male Female..... Central and South American, both sexes. 1.8 Male Female..... Other and Unknown Hispanic, both sexes. 3.0 Male.... Female..... Non-Hispanic 1/, both sexes..... 2,945 4,305 5,162 6,114 7,376 8,960 8,832 6,209 2,945 Male.... 1,669 2,464 2,929 3,492 3,952 4,682 4,220 2,574 1.019 1,276 1,841 2,233 2,622 3,424 4,278 4,612 3,635 1,926 Female..... Non-Hispanic white, both sexes..... 2,289 3,498 4,320 5,157 6,458 8,002 7,999 5,718 2,697 1,999 3,865 2,383 Male.... 1,291 2,458 2,961 3,481 4,236 1,499 1,862 2,196 2,977 3,766 4,134 3,335 1,762 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... 2.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) All origins, both sexes..... 1,422 2,021 2,623 2,172 1.176 Male.... 1,078 1,300 Female..... 1,323 1,226 Hispanic, both sexes..... 2.0 Male.... 2.7 Female..... 2.6 Mexican, both sexes..... 2.0 2.0 2.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,344 1,941 2,536 2,124 1,141 Male.... 1,036 1,259 1,277 1,201 Female..... 1,978 Non-Hispanic white, both sexes..... 1,188 1,744 2,337 1,070 1,169 Male.... 1,168 1,110 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... 

Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Anemias (D50-D64) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 1.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... 3.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Diabetes mellitus (E10-E14) All origins, both sexes..... 3,350 4.842 5,938 7,102 8,991 11,098 11,847 8,711 4,643 1,337 227 3 Male.... 2,017 2,758 3,395 3,832 4,598 5,335 5,234 3,296 1,408 45 Female..... 1,333 2,084 2,543 3,270 4,393 5,763 6,613 5,415 3,235 1.051 182 2 Hispanic, both sexes..... 300 477 596 734 853 960 809 442 248 67 12 Male.... 341 357 402 433 348 142 71 16 3 Female..... 109 216 255 377 451 527 461 300 177 51 9 Mexican, both sexes..... 194 307 394 486 566 606 530 254 134 38 6 Male.... 125 170 223 232 263 273 236 82 45 11 Female..... 69 137 171 254 303 333 294 172 89 2.7 5 Puerto Rican, both sexes..... 40 67 82 103 94 106 88 63 2.7 7 1 Male.... 37 42 57 44 48 34 15 7 15 30 40 46 50 58 54 48 2.0 1 Female..... Cuban, both sexes..... 14 1.0 24 40 54 82 86 52 32 8 1 Male 11 6 15 21 31 48 37 23 6 3 4 9 19 23 34 49 29 26 1 Female..... 9 29 2.0 29 49 40 16 2.2 1 Central and South American, both sexes. 12 2 Male 5 14 13 11 24 12 6 2 3 1 Female..... 15 7 18 25 28 10 20 9 2 Other and Unknown Hispanic, both sexes. 42 64 76 76 an 126 QQ 51 43 3 12 Male.... 25 34 48 36 40 52 35 20 10 4 1 Female..... 18 30 28 40 50 74 54 31 33 8 2 Non-Hispanic 1/, both sexes..... 3,039 4,346 5,323 6,355 8,116 10,110 11,018 8,248 4,386 1,263 214 3 Male.... 1,819 2,484 3,043 3,468 4,185 4,884 4,879 3,150 1,333 266 42 1 1,220 1,862 2,280 2,887 3,931 5,226 6,139 5,098 3,053 997 172 2 Female..... Non-Hispanic white, both sexes..... 2,073 3,089 3,876 4,704 6,171 8,066 9,154 6,979 3,714 1,055 148 2 4,059 1,147 Male.... 1,309 1,815 2,313 2,659 3,325 4,217 2,751 230 33 764 1,274 1,563 2,045 2,846 4,007 4,937 4,228 2,567 825 115 2 Female..... Non-Hispanic black, both sexes..... 856 1,075 1,229 1,423 1,655 1,733 1,536 1,062 566 176 60 1 446 561 610 692 725 671 536 318 148 2.8 6 1 Male.... Female..... 514 619 731 930 1,062 1,000 744 418 148 54 Origin not stated 2/, both sexes..... 11 19 19 13 22 2.8 2.0 21 9 7 1 7 13 11 7 11 18 7 4 4 Male.... 4 Female..... 6 8 6 11 10 13 17 5

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Nutritional deficiencies (E40-E64)												
All origins, both sexes	77	72	90	120	178	339	558	644	518	248	64	1
Male	34	42	45	60	85	153	204	225	134	48	6	1
Female	43	30	45	60	93	186	354	419	384	200	58	_
Hispanic, both sexes	3	4	4	9	8	14	18	15	11	11	4	-
Male	1	2	1	6	3	4	7	4	4	1	1	-
Female	2	2	3	3	5	10	11	11	7	10	3	_
Mexican, both sexes	3	2	4	8	6	6	12	10	8	8	1	_
Male	1	1	1	5	2	1	5	3	2	1	1	_
Female	2	1	3	3	4	5	7	7	6	7	-	_
Puerto Rican, both sexes	-	2	-	_	1	2	-	1	1	_	1	_
Male	-	1	_	-	-	1	-	-	1	-	-	-
Female	-	1	-	_	1	1	-	1	_	_	1	_
Cuban, both sexes	-	-	_	1	1	1	2	2	-	1	-	-
Male	_	-	-	1	1	-	1	-	-	-	_	-
Female	-	-	_	-	-	1	1	2	-	1	-	-
Central and South American, both sexes.	-	-	_	-	-	3	1	1	-	-	-	-
Male	_	_	_	_	_	1	_	_	_	_	_	_
Female	-	-	_	-	-	2	1	1	-	-	-	-
Other and Unknown Hispanic, both sexes.	_	_	_	_	_	2	3	1	2	2	2	_
Male	_	_	_	_	_	1	1	1	1	_	_	_
Female	_	_	_	_	_	1	2	_	1	2	2	_
Non-Hispanic 1/, both sexes	74	67	86	111	170	323	539	628	507	237	60	_
Male	33	39	44	54	82	149	196	220	130	47	5	_
Female	41	28	42	57	88	174	343	408	377	190	55	_
Non-Hispanic white, both sexes	52	53	63	82	135	264	468	536	444	209	48	_
Male	24	28	32	34	67	121	168	190	112	42	4	_
Female	28	25	31	48	68	143	300	346	332	167	44	_
Non-Hispanic black, both sexes	20	13	21	26	32	48	56	78	52	23	11	_
Male	9	10	11	19	13	23	20	24	15	5	1	_
Female	11	3	10	7	19	25	36	54	37	18	10	_
Origin not stated 2/, both sexes		1		_		2	1	1	-	_		1
Male	_	1	_	_	_	_	1	1	_	_	_	1
Female	_	_	_	_	_	2	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Malnutrition (E40-E46) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... - 1 Female..... 1.0 1.0 Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 8.0 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 2.0 a Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other nutritional deficiencies (E50-E64) All origins, both sexes..... 4 4 6 4 6 11 22 31 47 21 2 Male.... 4 2 5 10 13 14 5 Female..... 2 2 2 2 4 6 12 18 33 16 2 Hispanic, both sexes..... 2 Male.... Female..... 2 Mexican, both sexes..... 1 Male..... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1 Male.... 1 Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 Male.... Female..... 1 Non-Hispanic 1/, both sexes..... 4 6 4 11 20 31 47 21 Male.... 2 4 2. 2 5 1.0 13 14 5 2 10 18 33 Female..... 2 16 Non-Hispanic white, both sexes..... 3 11 18 30 20 Male.... 2 1 1 2 5 9 12 1.3 5 Female..... 2 6 9 18 32 15 1 Non-Hispanic black, both sexes..... 1 2 2 2 1 2 1 Male.... 1 1 1 1 Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Meningitis (G00,G03) All origins, both sexes..... 3.0 Male.... 2.8 2.0 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female.... Non-Hispanic 1/, both sexes..... Male.... 2.5 1.5 Female..... Non-Hispanic white, both sexes..... Male.... 1.3 1.3 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Parkinson's disease (G20-G21)												
All origins, both sexes	48	103	246	640	1,574	3,526	5,226	4,174	1,937	441	46	_
Male	37	71	156	421	1,036	2,189	3,117	2,260	888	160	10	-
Female	11	32	90	219	538	1,337	2,109	1,914	1,049	281	36	-
Hispanic, both sexes	1	9	14	25	62	132	170	113	53	13	-	-
Male	1	5	10	16	38	85	101	62	22	6	-	-
Female	-	4	4	9	24	47	69	51	31	7	-	-
Mexican, both sexes	1	4	10	11	30	68	85	51	26	4	-	-
Male	1	3	7	6	18	41	49	27	11	3	-	-
Female	-	1	3	5	12	27	36	24	15	1	-	-
Puerto Rican, both sexes	-	2	1	3	3	13	13	12	4	_	-	-
Male	-	-	1	2	2	7	6	6	2	_	-	-
Female	-	2	-	1	1	6	7	6	2	_	-	-
Cuban, both sexes	-	-	2	5	11	19	41	22	11	3	-	-
Male	-	-	2	2	8	14	27	11	4	1	-	-
Female	-	-	-	3	3	5	14	11	7	2	-	-
Central and South American, both sexes.	_	1	_	2	5	5	6	5	3	-	-	_
Male	-	1	_	2	2	5	4	3	1	-	-	-
Female	_	_	_	-	3	-	2	2	2	-	-	_
Other and Unknown Hispanic, both sexes.	-	2	1	4	13	27	25	23	9	6	-	-
Male	-	1	_	4	8	18	15	15	4	2	-	-
Female	-	1	1	-	5	9	10	8	5	4	-	-
Non-Hispanic 1/, both sexes	47	93	231	614	1,512	3,385	5,049	4,056	1,881	428	46	_
Male	36	65	145	404	998	2,099	3,010	2,195	865	154	10	_
Female	11	28	86	210	514	1,286	2,039	1,861	1,016	274	36	_
Non-Hispanic white, both sexes	39	87	209	557	1,431	3,187	4,812	3,860	1,783	403	37	_
Male	29	62	128	370	943	1,984	2,880	2,086	822	140	8	_
Female	10	25	81	187	488	1,203	1,932	1,774	961	263	29	_
Non-Hispanic black, both sexes	5	2	19	37	55	132	158	126	63	18	7	_
Male	5	_	15	21	39	75	81	69	24	10	2	_
Female	_	2	4	16	16	57	77	57	39	8	5	_
Origin not stated 2/, both sexes	_	1	1	1		9	7	5	3	_	-	_
Male	_	1	1	1	_	5	6	3	1	_	_	_
Female	_	_	_	_	_	4	1	2	2	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	-
Alzheimer's disease (G30)												
All origins, both sexes	65	152	390	1,004	2,637	7,368	14,514	18,327	14,505	5,871	1,087	_
Male	30	66	187	434	1,124	2,767	5,010	5,282	3,087	883	91	-
Female	35	86	203	570	1,513	4,601	9,504	13,045	11,418	4,988	996	-
Hispanic, both sexes	5	7	15	43	110	259	422	443	377	174	38	-
Male	2	3	7	15	50	100	160	135	78	41	13	-
Female	3	4	8	28	60	159	262	308	299	133	25	-
Mexican, both sexes	2	2	6	17	48	122	206	200	148	68	17	_
Male	-	1	4	5	21	43	82	71	27	21	7	-
Female	2	1	2	12	27	79	124	129	121	47	10	-
Puerto Rican, both sexes	3	2	7	5	24	38	49	43	36	22	1	_
Male	2	1	2	_	11	16	15	11	8	8	_	_
Female	1	1	5	5	13	22	34	32	28	14	1	_
Cuban, both sexes	_	_	_	8	22	52	89	107	101	46	11	_
Male	_	_	_	2	7	24	30	29	27	5	2	_
Female	_	_	_	6	15	28	59	78	74	41	9	_
Central and South American, both sexes.	_	1	_	3	4	12	19	17	17	7	2	_
Male	_	1	_	1	1	3	7	5	2	2	1	_
Female	_	_	_	2	3	9	12	12	15	5	1	_
Other and Unknown Hispanic, both sexes.	_	2	2	10	12	35	59	76	75	31	7	_
Male	_	_	1	7	10	14	26	19	1.4	5	3	_
Female	_	2	1	3	2	21	33	57	61	26	4	_
Non-Hispanic 1/, both sexes	60	145	374	959	2,518	7,098	14,075	17,869	14,107	5,688	1,047	_
Male	28	63	179	418	1,072	2,664	4,843	5,144	3,003	842	78	_
Female	32	82	195	541	1,446	4,434	9,232	12,725	11,104	4,846	969	_
Non-Hispanic white, both sexes	56	134	338	851	2,256	6,458	12,980	16,698	13,147	5,272	923	_
Male	27	60	161	376	959	2,439	4,506	4,821	2,788	780	62	_
Female	29	74	177	475	1,297	4,019	8,474	11,877	10,359	4,492	861	_
Non-Hispanic black, both sexes	4	10	27	98	229	563	947	1,013	837	361	106	_
Male	1	2	13	38	98	196	290	268	170	48	13	_
Female	3	2	14	60	131	367	657	745	667	313	93	_
Origin not stated 2/, both sexes	_	-	1	2	9	11	17	15	21	313	2	_
Male	_	_	1	1	2	3	± /	3	6	_	_	_
Female	_	_	_	1	7	8	10	12	15	9	2	
r-cmarc	_	-	_	Τ.	/	8	Τ0	⊥∠	Τ2	9	2	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Major cardiovascular diseases (I00-I78)												
All origins, both sexes  Male Female.  Hispanic, both sexes.  Male. Female.  Mexican, both sexes.	27,138 18,931 8,207 1,656 1,162 494 955	34,808 23,704 11,104 2,017 1,371 646 1,203	43,176 28,310 14,866 2,477 1,615 862 1,351	53,520 33,114 20,406 3,074 1,880 1,194 1,706	75,296 43,065 32,231 3,984 2,252 1,732 2,174	112,224 58,898 53,326 5,113 2,675 2,438 2,754	151,813 69,592 82,221 5,620 2,716 2,904 2,959	153,212 58,466 94,746 4,535 1,795 2,740 2,115	112,753 33,116 79,637 3,354 1,037 2,317 1,557	45,518 9,712 35,806 1,411 368 1,043 647	10,061 1,514 8,547 336 84 252 161	45 38 7 5 5
Male Female Puerto Rican, both sexes Male Female	667 288 229 161 68	818 385 315 212 103	874 477 418 264 154	1,016 690 455 282 173	1,207 967 559 304 255	1,415 1,339 646 332 314	1,451 1,508 701 316 385	890 1,225 548 188 360	534 1,023 366 102 264	182 465 156 33 123	49 112 43 7 36	2 - 1 1 -
Cuban, both sexes	98 75 23 148 109	104 76 28 161 104	198 143 55 210 134	311 221 90 220 126	507 342 165 285 139	755 448 307 322 137	862 430 432 333 133	931 371 560 329 107	707 202 505 218 45	310 74 236 73 17	52 6 46 20 3	- - - -
Female Other and Unknown Hispanic, both sexes. Male Female Non-Hispanic 1/, both sexes	39 226 150 76 25,380	57 234 161 73	76 300 200 100 40,555	94 382 235 147 50,279	146 459 260 199 71,094	185 636 343 293 106,872	200 765 386 379 145,915	222 612 239 373 148,453	173 506 154 352 109,219	56 225 62 163 44,039	17 60 19 41 9,712	- 2 2 - 33
Male  Female  Non-Hispanic white, both sexes  Male	17,688 7,692 18,175 13,211	32,669 22,245 10,424 24,508 17,171	26,585 13,970 31,633 21,187	31,116 19,163 39,976 25,320	40,671 30,423 58,964 34,478	56,086 50,786 92,057 49,164	66,734 79,181 130,212 60,340	56,593 91,860 135,393 52,117	32,021 77,198 99,906 29,365	9,331 34,708 40,145 8,450	1,427 8,285 8,470 1,204	26 7 19 15
Female. Non-Hispanic black, both sexes Male. Female. Origin not stated 2/, both sexes Male. Female.	4,964 6,460 3,957 2,503 102 81	7,337 7,217 4,450 2,767 122 88 34	10,446 7,856 4,707 3,149 144 110	14,656 8,831 4,926 3,905 167 118 49	24,486 10,401 5,274 5,127 218 142	42,893 12,461 5,702 6,759 239 137	69,872 12,864 5,066 7,798 278 142	83,276 10,426 3,388 7,038 224 78	70,541 7,626 1,977 5,649 180 58 122	31,695 3,243 642 2,601 68 13	7,266 1,068 174 894 13 3	4 13 10 3 7 7

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Diseases of heart (I00-I09,I11,I13,I20-I51) All origins, both sexes..... 22,166 28,596 35,017 42,187 57,812 83,741 111,638 112,382 83,574 34.234 7.734 36 Male.... 15,972 20,188 23,600 26,871 34,129 45,619 53,193 44,648 25,565 7.574 1,215 32 Female..... 6,194 8,408 11,417 15,316 23,683 38,122 58,445 67,734 58,009 26,660 6,519 4 Hispanic, both sexes..... 1,220 1.519 1.941 2,362 3,027 3,839 4.222 3,344 2,536 1,097 269 3 Male.... 1.072 1,304 1,467 1,769 2,083 2,080 1,321 813 294 68 3 Female..... 331 447 637 895 1,258 1,756 2,142 2,023 1.723 803 201 Mexican, both sexes..... 700 901 1,044 1,281 1,629 1.997 2,149 1,470 1.132 480 125 1 Male.... 509 642 698 784 943 1,070 1.072 623 405 144 39 1 Female..... 191 259 346 497 686 927 1,077 847 727 336 86 Puerto Rican, both sexes..... 165 241 333 368 441 514 558 432 294 122 38 1 Male.... 116 164 215 229 243 269 250 147 88 2.7 7 1 49 77 118 139 198 245 308 285 206 95 31 Female..... Cuban, both sexes..... 82 92 165 249 413 608 711 765 583 267 45 Male 64 67 121 171 279 373 357 303 167 63 5 18 25 44 78 134 235 354 462 416 204 40 Female..... 101 169 206 250 253 261 170 63 Central and South American, both sexes. 116 153 15 Male 77 77 97 96 101 107 106 91 37 14 2 Female..... 24 39 56 73 105 143 147 170 133 49 13 Other and Unknown Hispanic, both sexes. 172 169 246 295 338 470 551 416 357 165 46 1 Male.... 123 122 173 187 203 264 295 157 116 46 15 1 Female..... 49 47 73 108 135 206 256 259 241 119 31 27 Non-Hispanic 1/, both sexes..... 20,864 26,976 32,954 39,685 54,603 79,709 107,199 108,868 80,898 33,084 7,455 43,261 7,269 Male.... 15,016 19,041 22,201 25,301 32,238 43,423 50,995 24.704 1,145 23 5,848 7,935 10,753 14,384 22,365 36,286 56,204 65,607 56,194 25,815 6,310 4 Female..... Non-Hispanic white, both sexes..... 15,395 20,700 26,234 31,971 45,681 69,171 96,088 99,626 74,261 30,248 6,534 16 46,330 Male.... 11,527 15,021 18,057 20,843 27,574 38,364 39,954 22,721 6,611 964 13 3,868 5,679 8,177 11,128 18,107 30,807 49,758 59,672 51,540 23,637 5,570 3 Female..... Non-Hispanic black, both sexes..... 4,972 5,601 5,940 6,658 7,714 8,959 9,180 7,400 5,443 2,372 803 10 3.124 3,529 3,620 3,792 4,006 4.211 3,720 2.516 1.483 480 142 9 Male.... Female..... 1,848 2,072 2,320 2,866 3,708 4,748 5,460 4,884 3,960 1.892 661 Origin not stated 2/, both sexes..... 82 101 122 140 182 193 217 170 140 53 1.0 6 67 75 95 103 122 113 118 66 48 11 2 6 Male.... Female..... 15 26 27 37 60 80 99 104 92 42

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 2.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 5.0 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Hypertensive heart disease (I11) 3,676 All origins, both sexes..... 1.760 2,012 1,873 1,867 2,134 2,722 4,053 3,411 1,510 Male.... 1,185 1,363 1,221 1,146 1,157 1,213 1,291 1,167 Female..... 1,509 2,385 2,886 2,665 1.279 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 2.1 Male.... Female..... Cuban, both sexes..... 2.0 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 2.4 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,640 1,867 1,751 1,710 1,980 2,557 3,502 3,901 3,290 1,456 1,070 Male.... 1,110 1,260 1,136 1,059 1,135 1,208 1,114 1,422 2,294 2,787 2,579 1,236 Female..... Non-Hispanic white, both sexes..... 1,124 1,105 1,142 1,366 1,879 2,782 3,306 2,835 1,269 Male.... 1,028 1,823 2,366 2,236 1,086 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes..... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Hypertensive heart and renal disease (I13) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 1.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex  Ischemic heart diseases (I20-I25)	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	_
All origins, both sexes	15,192	20,437	25,565	30,914	42,188	60,283	78,845	76,562	55,554	22,517	5,123	26
Male	11,544	15,004	17,864	20,392	25,881	34,272	39,319	31,859	17,650	5,155	831	23
Female	3,648	5,433	7,701	10,522	16,307	26,011	39,526	44,703	37,904	17,362	4,292	3
Hispanic, both sexes	862	1,135	1,474	1,795	2,352	2,970	3,238	2,533	1,928	823	210	3
Male	664	832	1,020	1,166	1,411	1,670	1,616	1,019	632	217	56	3
Female	198	303	454	629	941	1,300	1,622	1,514	1,296	606	154	-
Mexican, both sexes	493	671	795	963	1,275	1,547	1,637	1,108	837	355	98	1
Male	381	495	538	619	752	876	834	481	319	106	32	1
Female	112	176	257	344	523	671	803	627	518	249	66	-
Puerto Rican, both sexes	118	182	254	299	332	404	435	350	241	101	31	1
Male	85	125	172	194	192	213	197	122	73	22	5	1
Female	33	57	82	105	140	191	238	228	168	79	26	-
Cuban, both sexes	51	68	125	189	332	480	572	578	457	203	38	-
Male	41	53	95	136	221	292	285	225	127	47	5	-
Female	10	15	30	53	111	188	287	353	330	156	33	-
Central and South American, both sexes.	72	88	118	132	157	196	194	195	128	47	12	-
Male	57	60	79	73	85	87	78	73	27	9	2	-
Female	15	28	39	59	72	109	116	122	101	38	10	-
Other and Unknown Hispanic, both sexes.	128	126	182	212	256	343	400	302	265	117	31	1
Male	100	99	136	144	161	202	222	118	86	33	12	1
Female	28	27	46	68	95	141	178	184	179	84	19	-
Non-Hispanic 1/, both sexes	14,278	19,227	23,998	29,010	39,688	57,166	75,436	73,913	53,521	21,654	4,904	18
Male	10,839	14,118	16,773	19,143	24,367	32,511	37,607	30,793	16,980	4,929	773	15
Female	3,439	5,109	7,225	9,867	15,321	24,655	37,829	43,120	36,541	16,725	4,131	3
Non-Hispanic white, both sexes	11,052	15,305	19,572	23,714	33,552	50,014	67,802	67,688	49,153	19,760	4,297	10
Male	8,653	11,522	13,983	16,017	21,045	28,984	34,329	28,525	15,678	4,482	650	8
Female	2,399	3,783	5,589	7,697	12,507	21,030	33,473	39,163	33,475	15,278	3,647	2
Non-Hispanic black, both sexes	2,867	3,433	3,859	4,494	5,223	6,003	6,197	4,900	3,546	1,578	533	7
Male	1,910	2,219	2,396	2,601	2,809	2,886	2,555	1,691	966	329	98	6
Female	957	1,214	1,463	1,893	2,414	3,117	3,642	3,209	2,580	1,249	435	1
Origin not stated 2/, both sexes	52	75	93	109	148	147	171	116	105	40	9	5
Male	41	54	71	83	103	91	96	47	38	9	2	5
Female	11	21	22	26	45	56	75	69	67	31	7	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the												
International Classification of												
Diseases, Tenth Revision, 1992),	50-54	55-59	60 61	65 60	70-74	75-79	00 04	05 00	00 04	05 00	100	7aa nat
Hispanic origin, race for			60-64	65-69			80-84	85-89	90-94	95-99	100 years	_
non-Hispanic population, and sex	years	years	years	years	years	years	years	years	years	years	and over	stated
Acute myocardial infarction (I21-I22)												
All origins, both sexes	6,167	8,346	10,412	12,357	15,970	21,573	26,464	24,512	16,524	5,939	1,277	6
Male	4,643	6,107	7,235	8,017	9,641	11,987	12,906	9,983	5,286	1,393	233	4
Female	1,524	2,239	3,177	4,340	6,329	9,586	13,558	14,529	11,238	4,546	1,044	2
Hispanic, both sexes	313	441	541	677	853	1,038	1,123	863	577	231	46	1
Male	240	314	378	435	493	572	556	348	191	50	11	1
Female	73	127	163	242	360	466	567	515	386	181	35	-
Mexican, both sexes	194	292	327	405	504	573	607	393	262	118	22	-
Male	150	211	226	262	297	320	308	175	108	31	8	-
Female	44	81	101	143	207	253	299	218	154	87	14	-
Puerto Rican, both sexes	45	55	70	90	105	122	123	97	56	20	3	1
Male	35	40	43	55	59	58	53	33	18	2	1	1
Female	10	15	27	35	46	64	70	64	38	18	2	-
Cuban, both sexes	17	26	42	65	119	169	217	217	160	64	10	-
Male	12	19	33	44	69	105	104	77	39	11	1	-
Female	5	7	9	21	50	64	113	140	121	53	9	-
Central and South American, both sexes.	18	26	46	36	39	68	36	59	31	6	2	-
Male	13	10	30	17	19	29	15	23	6	1	-	-
Female	5	16	16	19	20	39	21	36	25	5	2	-
Other and Unknown Hispanic, both sexes.	39	42	56	81	86	106	140	97	68	23	9	-
Male	30	34	46	57	49	60	76	40	20	5	1	-
Female	9	8	10	24	37	46	64	57	48	18	8	-
Non-Hispanic 1/, both sexes	5,841	7,885	9,845	11,643	15,083	20,493	25,293	23,617	15,914	5,697	1,229	5
Male	4,393	5,780	6,837	7,552	9,123	11,389	12,327	9,626	5,080	1,341	222	3
Female	1,448	2,105	3,008	4,091	5,960	9,104	12,966	13,991	10,834	4,356	1,007	2
Non-Hispanic white, both sexes	4,655	6,466	8,282	9,722	12,844	18,084	22,756	21,545	14,529	5,125	1,038	3
Male	3,615	4,858	5,910	6,452	7,936	10,285	11,271	8,881	4,690	1,205	184	2
Female	1,040	1,608	2,372	3,270	4,908	7,799	11,485	12,664	9,839	3,920	854	1
Non-Hispanic black, both sexes	1,053	1,255	1,361	1,612	1,899	2,039	2,084	1,660	1,119	489	176	1
Male	683	789	793	904	990	907	817	583	283	105	33	-
Female	370	466	568	708	909	1,132	1,267	1,077	836	384	143	1
Origin not stated 2/, both sexes	13	20	26	37	34	42	48	32	33	11	2	-
Male	10	13	20	30	25	26	23	9	15	2	-	-
Female	3	7	6	7	9	16	25	23	18	9	2	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Other acute ischemic heart diseases (I24) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.1 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Other forms of chronic ischemic heart disease (I20,I25) All origins, both sexes..... 8,889 11,907 14.978 18,378 25,998 38,402 52,019 51,723 38,784 16.467 3,813 19 Male.... 6,805 8,765 10,510 12,251 16,110 22,131 26,238 21,733 12,288 3.741 18 Female..... 2,084 3,142 4,468 6,127 9,888 16,271 25,781 29,990 26,496 12,726 3.221 1 Hispanic, both sexes..... 547 692 929 1,114 1,497 1,926 2,110 1,669 1,350 592 164 1 Male.... 422 516 640 730 916 1,094 1,058 670 441 167 45 Female..... 125 176 289 384 581 832 1,052 999 909 425 119 Mexican, both sexes..... 299 377 467 556 770 970 1,028 715 575 237 76 Male.... 231 282 311 357 454 552 525 306 211 75 24 Female..... 68 95 156 199 316 418 503 409 364 162 52 Puerto Rican, both sexes..... 72 127 182 209 226 282 310 253 185 81 28 Male.... 85 128 139 132 155 143 89 55 2.0 4 23 42 54 70 94 127 167 164 130 61 24 Female..... Cuban, both sexes..... 33 42 83 124 213 311 355 361 297 139 28 Male 2.8 34 62 92 152 187 181 148 88 36 4 5 8 21 32 61 124 174 213 209 103 24 Female..... 54 62 72 96 118 127 157 136 97 41 Central and South American, both sexes. 1.0 Male 44 50 49 56 66 58 63 50 21 8 2 Female..... 10 12 23 40 52 69 94 86 76 33 Other and Unknown Hispanic, both sexes. QQ Ω4 125 129 170 236 260 204 196 94 22 1 Male.... 70 65 90 86 112 142 146 77 66 2.8 11 1 Female..... 19 19 35 43 58 94 114 127 130 66 11 Non-Hispanic 1/, both sexes..... 8,304 11,160 13,983 17,193 24,387 36,371 49,787 49,972 37,362 15,846 3,642 13 Male.... 6,353 8,208 9,820 11,469 15,116 20,972 25,107 21,026 11,824 3.567 545 12 1,951 2,952 5,724 9,271 15,399 24,680 28,946 25,538 12,279 3,097 Female..... 4.163 1 Non-Hispanic white, both sexes..... 6,308 8,709 11,158 13,858 20,531 31,679 44,738 45,847 34,401 14,531 3,230 7,983 18,576 22,902 3,257 Male.... 4,975 6,570 9,474 13,009 19,519 10,920 461 6 1,333 2,139 3,175 4,384 7,522 13,103 21,836 26,328 23,481 11,274 2,769 Female..... Non-Hispanic black, both sexes..... 1.773 2,129 2,462 2,845 3,291 3,916 4,073 3,217 2,406 1,082 354 6 1,200 1,396 1,578 1,669 1,797 1,953 1.722 1,094 676 223 64 Male.... 6 Female..... 573 733 884 1,176 1,494 1,963 2,351 2,123 1,730 859 290 Origin not stated 2/, both sexes..... 38 55 66 71 114 105 122 82 72 29 7 5 30 41 50 52 78 65 73 37 23 7 2 5 Male.... Female..... 14 16 19 36 40 49 45 49 2.2

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Atherosclerotic cardiovascular disease, so described (I25.0) All origins, both sexes..... 3.555 4.378 4,919 4,957 5,873 7,467 9,263 8,986 6.723 2,989 Male.... 2,821 3,361 3,503 3,389 3,657 4,145 4,461 3,499 1,977 Female..... 1,017 1,416 1,568 2,216 3,322 4,802 5,487 4,746 2,338 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... 2.4 Female..... Cuban, both sexes..... 1.0 2.0 Male Female..... 2.8 2.0 Central and South American, both sexes. 3.0 Male Female..... Other and Unknown Hispanic, both sexes. 4 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 3,282 4,060 4,527 4,596 5,436 7,016 8,820 8,684 6,514 2,895 Male.... 2,587 3,100 3,223 3,145 3,366 3,865 4,218 3,379 1,908 1,304 1,451 2,070 3,151 4,602 5,305 4,606 2,274 Female..... Non-Hispanic white, both sexes..... 2,410 3,028 3,415 3,460 4,279 5,703 7,577 7,712 5,815 2,579 2,425 2,708 3,654 1,703 Male.... 1,973 2,353 2,451 3,171 3,010 1,035 1,571 2,532 3,923 4,702 4,112 2,035 Female..... Non-Hispanic black, both sexes..... 1,043 1,154 1,072 Male.... Female..... Origin not stated 2/, both sexes..... 2.4 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) All origins, both sexes..... 5.334 7,529 10,059 13,421 20,125 30,935 42,756 42,737 32.061 13.478 3,052 5 Male.... 3.984 5,404 7,007 8,862 12,453 17,986 21,777 18,234 10,311 3,090 477 5 Female..... 1,350 2,125 3,052 4,559 7,672 12,949 20,979 24,503 21.750 10,388 2.575 Hispanic, both sexes..... 289 403 566 790 1,112 1,512 1,710 1,389 1,165 503 130 1 Male.... 382 511 664 838 840 562 378 32 Female..... 89 127 184 279 448 674 870 827 787 364 98 Mexican, both sexes..... 153 218 274 389 550 741 821 574 490 194 56 Male.... 104 148 171 244 317 408 414 245 182 64 16 Female..... 49 70 103 145 233 333 407 329 308 130 40 Puerto Rican, both sexes..... 35 86 117 159 177 233 262 217 163 73 24 Male.... 57 82 102 106 126 119 76 48 15 3 15 29 35 57 71 107 143 141 115 58 21 Female..... Cuban, both sexes..... 23 2.7 63 97 183 267 305 330 271 131 26 Male 19 21 47 72 125 161 154 137 78 34 4 6 16 25 58 106 151 193 193 97 22 Female..... 4 29 70 85 101 129 116 84 33 9 Central and South American, both sexes. 32 49 Male 20 23 34 40 43 47 52 42 17 6 Female..... 9 9 15 30 42 54 77 74 67 2.7 7 Other and Unknown Hispanic, both sexes. 4 Q 40 63 75 117 170 102 152 157 72 15 1 Male.... 37 27 48 53 73 96 101 62 53 2.0 7 1 Female..... 12 13 15 22 44 74 92 90 104 52 Non-Hispanic 1/, both sexes..... 5,022 7,100 9,456 12,597 18,951 29,355 40,967 41,288 30,848 12,951 2,918 3 3,766 11,750 Male.... 5,108 6,597 8,324 17,107 20,889 17,647 9,916 2.946 444 3 1,256 1,992 2,859 4,273 7,201 12,248 20,078 23,641 20,932 10,005 2,474 Female..... 3 Non-Hispanic white, both sexes..... 3,898 5,681 7,743 10,398 16,252 25,976 37,161 38,135 28,586 11,952 2,616 7,049 15,405 2,713 Male.... 3,002 4,217 5,532 10,301 19,248 16,509 9,217 381 3 896 1,464 2,211 3,349 5,951 10,571 17,913 21,626 19,369 9,239 2,235 Female..... Non-Hispanic black, both sexes..... 993 1,213 1,485 1,852 2,248 2,762 3,001 2,414 1,810 808 258 661 745 905 1,051 1,200 1,352 1,240 808 509 162 46 Male.... Female..... 332 468 580 801 1,048 1,410 1,761 1,606 1,301 646 212 Origin not stated 2/, both sexes..... 23 26 37 34 62 68 79 60 48 2.4 4 1 18 2.0 28 27 39 41 48 25 17 5 1 1 Male.... Female..... 5 6 9 7 23 27 31 35 31 19 3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Other heart diseases (I26-I51)												
All origins, both sexes	4,987	5,890	7,244	8,989	12,916	19,868	27,985	30,758	23,870	9,971	2,166	7
Male	3,131	3,712	4,359	5,157	6,876	9,799	12,161	11,330	6,977	2,142	325	6
Female	1,856	2,178	2,885	3,832	6,040	10,069	15,824	19,428	16,893	7,829	1,841	1
Hispanic, both sexes	234	231	318	387	506	664	763	646	472	221	47	-
Male	155	141	193	208	265	315	361	241	141	66	10	-
Female	79	90	125	179	241	349	402	405	331	155	37	_
Mexican, both sexes	146	145	169	231	259	335	400	288	226	100	22	-
Male	94	90	110	120	140	137	190	108	63	33	6	_
Female	52	55	59	111	119	198	210	180	163	67	16	_
Puerto Rican, both sexes	32	35	58	50	83	82	94	56	40	19	6	_
Male	23	24	31	26	35	44	41	18	12	3	2	_
Female	9	11	27	24	48	38	53	38	28	16	4	_
Cuban, both sexes	16	11	22	31	63	96	112	162	108	49	6	_
Male	13	6	14	18	46	65	60	67	35	13	_	_
Female	3	5	8	13	17	31	52	95	73	36	6	_
Central and South American, both sexes.	11	12	21	22	34	43	44	48	26	11	2	_
Male	10	7	10	17	12	15	17	13	7	5	_	_
Female	1	5	11	5	22	28	27	35	19	6	2	_
Other and Unknown Hispanic, both sexes.	29	28	48	53	67	108	113	92	72	42	11	_
Male	15	14	28	27	32	54	53	35	24	12	2	_
Female	14	14	20	26	35	54	60	57	48	30	9	_
Non-Hispanic 1/, both sexes	4,739	5,642	6,913	8,586	12,393	19,169	27,185	30,071	23,369	9,742	2,118	7
Male	2,965	3,559	4,154	4,940	6,601	9,465	11,782	11,073	6,828	2,075	315	6
Female	1,774	2,083	2,759	3,646	5,792	9,704	15,403	18,998	16,541	7,667	1,803	1
Non-Hispanic white, both sexes	3,301	4,139	5,371	6,853	10,339	16,598	24,574	27,752	21,643	9,020	1,884	4
Male	2,142	2,662	3,270	3,998	5,615	8,273	10,695	10,237	6,284	1,906	269	3
Female	1,159	1,477	2,101	2,855	4,724	8,325	13,879	17,515	15,359	7,114	1,615	1
Non-Hispanic black, both sexes	1,336	1,379	1,392	1,537	1,825	2,238	2,213	1,926	1,435	603	198	3
Male	761	821	795	830	870	1,026	904	666	406	120	33	3
Female	575	558	597	707	955	1,212	1,309	1,260	1,029	483	165	_
Origin not stated 2/, both sexes	14	17	13	16	17	35	37	41	29	8	1	_
Male	11	12	12	9	10	19	18	16	8	1	_	_
Female	3	5	1	7	7	16	19	25	21	7	1	_
	,	3	_	,	,	10	1.7	23	21	,	_	

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Acute and subacute endocarditis (I33) All origins, both sexes..... Male.... 5.8 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 8.5 1.0 Female..... Non-Hispanic white, both sexes..... Male.... 1.0 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Diseases of pericardium and acute myocarditis (I30-I31,I40) All origins, both sexes..... 5.8 Male.... Female..... 2.4 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female.... Non-Hispanic 1/, both sexes..... Male.... 2.6 2.7 2.1 1.0 Female..... Non-Hispanic white, both sexes..... Male.... 2.2 2.2 1.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Heart failure (I50)												
All origins, both sexes	653	1,003	1,441	2,226	3,725	6,574	10,438	12,876	11,063	5,044	1,203	1
Male	378	599	821	1,223	1,940	3,198	4,475	4,707	3,202	1,049	172	-
Female	275	404	620	1,003	1,785	3,376	5,963	8,169	7,861	3,995	1,031	1
Hispanic, both sexes	31	42	62	98	146	224	275	259	208	107	30	-
Male	18	29	35	52	73	103	132	96	53	35	4	-
Female	13	13	27	46	73	121	143	163	155	72	26	-
Mexican, both sexes	20	28	36	75	81	137	169	138	109	52	17	-
Male	12	20	22	38	45	58	78	50	23	18	3	-
Female	8	8	14	37	36	79	91	88	86	34	14	-
Puerto Rican, both sexes	3	4	10	7	23	28	30	24	21	10	3	_
Male	2	2	6	4	10	13	18	7	4	2	_	_
Female	1	2	4	3	13	15	12	17	17	8	3	_
Cuban, both sexes	_	1	3	4	8	14	22	40	37	15	3	_
Male	_	1	2	2	4	10	13	18	13	5	_	_
Female	_	_	1	2	4	4	9	22	24	10	3	_
Central and South American, both sexes.	2	2	1	3	9	11	8	18	7	3	1	_
Male	2	1	_	2	2	3	2	4	1	_	_	_
Female	_	1	1	1	7	8	6	14	6	3	1	_
Other and Unknown Hispanic, both sexes.	6	7	12	9	25	34	46	39	34	27	6	_
Male	2	5	5	6	12	19	21	17	12	10	1	_
Female	4	2	7	3	13	15	25	22	22	17	5	_
Non-Hispanic 1/, both sexes	619	958	1,376	2,125	3,573	6,338	10,149	12,593	10,843	4,932	1,172	1
Male	358	567	783	1,169	1,863	3,088	4,337	4,598	3,145	1,014	168	_
Female	261	391	593	956	1,710	3,250	5,812	7,995	7,698	3,918	1,004	1
Non-Hispanic white, both sexes	378	668	1,051	1,735	3,032	5,547	9,257	11,689	10,113	4,623	1,051	1
Male	230	399	596	959	1,621	2,721	3,990	4,282	2,937	950	148	_
Female	148	269	455	776	1,411	2,826	5,267	7,407	7,176	3,673	903	1
Non-Hispanic black, both sexes	232	271	290	345	484	705	782	7,407	630	274	104	_
Male	124	157	164	187	210	321	305	267	165	54	16	_
Female	108	114	126	158	274	384	477	508	465	220	88	_
Origin not stated 2/, both sexes	108	3	120	158	6	12	14	24	12	220 5	1	-
Male	3	3	3	2	0	14	6	13	4	5	1	-
	2	3	-		4	, 5	8			_	-	_
Female	1	_	-	1	2	5	8	11	8	5	1	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears All other forms of heart disease (I26-I28.I34-I38.I42-I49.I51) 4,754 All origins, both sexes..... 4.176 5,660 6,598 9,007 13,053 17.311 17,704 12,761 4,910 Male.... 2,653 3,031 3,457 3,831 4,858 6,491 7,566 6.544 3,755 1.086 Female..... 1,523 1,723 2,203 2,767 4,149 6,562 9,745 11,160 9,006 3,824 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 2.7 5.8 Male.... 3.0 Female..... Cuban, both sexes..... 1.0 Male Female..... Central and South American, both sexes. 1.0 3.0 Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 3,972 4,563 5,412 6,307 8,644 12,602 16,813 17,305 12,481 4,793 Male.... 2,516 2,919 3,302 3,673 4,664 6,270 7,333 6,399 3,663 1.054 1,456 2,110 2,634 3,980 6,332 9,480 10,906 8,818 3,739 Female..... 1.644 7,156 Non-Hispanic white, both sexes..... 2,828 3,377 4,224 4,997 10,863 15,116 15,906 11,488 4,384 5,886 3,327 Male.... 1,854 2,205 2,618 2,964 3,929 5,460 6,604 1,172 1,606 2,033 3,227 5,403 8,512 10,020 8,161 3,433 Female..... Non-Hispanic black, both sexes..... 1,055 1,084 1,075 1,163 1,318 1,501 1,414 1,137 Male.... Female..... Origin not stated 2/, both sexes..... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Essential (primary) hypertension and hypertensive renal disease (I10,I12) 1,024 All origins, both sexes..... 1,292 1,860 2,855 3,964 4,283 3,434 1,542 Male.... 1,241 1,541 1,363 Female..... 1,016 1,614 2,423 2,920 2,624 1.255 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 5.0 Male.... Female..... 1 0 5.8 Puerto Rican, both sexes..... 2.0 Male.... 1.0 Female..... Cuban, both sexes..... 1.0 2.0 Male Female..... 1.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 1 0 2.4 Ω Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 1,199 1,734 2,705 3,800 4,116 3,286 1,491 Male.... 1,174 1,469 1,310 2.77 1,531 2,331 2,806 2,516 1,214 Female..... Non-Hispanic white, both sexes..... 1,184 2,016 3,082 3,506 2,781 1,284 Male.... 1,203 1,136 1,144 1,879 2,370 2,112 1,056 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Cerebrovascular diseases (I60-I69)												
All origins, both sexes	3,559	4,368	5,598	7,806	12,095	20,504	29,588	30,162	21,179	7,752	1,452	5
	2,015	2,395	3,128	4,180	6,027	9,333	11,692	9,872	5,346	1,449	192	3
	1,544	1,973	2,470	3,626	6,068	11,171	17,896	20,290	15,833	6,303	1,260	2
	330	375	401	524	708	961	1,045	881	569	221	38	2
Male Female Mexican, both sexes	195	219	235	304	343	437	467	352	145	51	8	2
	135	156	166	220	365	524	578	529	424	170	30	-
	197	243	245	320	406	591	616	490	292	112	20	1
Male  Female  Puerto Rican, both sexes  Male	117 80 46 33	138 105 55 35	144 101 55 32	179 141 61 36	192 214 79 38	267 324 99 48	280 336 95 47	204 286 84 29	82 210 55 12	27 85 27 4	6 14 3 -	1 - -
FemaleCuban, both sexesMale	13	20	23	25	41	51	48	55	43	23	3	-
	11	7	21	43	69	109	109	114	84	33	4	-
	7	5	13	35	42	50	51	49	23	8	-	-
Female. Central and South American, both sexes. Male Female	4 34 19 15	2 31 21 10	8 44 28 16	8 34 19 15	27 61 29 32	59 52 22 30	58 61 21 40	65 44 10 34	61 32 6 26	25 7 2 5	4 3 - 3	- - -
Other and Unknown Hispanic, both sexes. Male Female	42	39	36	66	93	110	164	149	106	42	8	1
	19	20	18	35	42	50	68	60	22	10	2	1
	23	19	18	31	51	60	96	89	84	32	6	-
Non-Hispanic 1/, both sexes  Male	3,216	3,978	5,185	7,265	11,366	19,510	28,499	29,244	20,581	7,522	1,411	2
	1,810	2,168	2,885	3,867	5,674	8,879	11,207	9,510	5,196	1,397	183	-
	1,406	1,810	2,300	3,398	5,692	10,631	17,292	19,734	15,385	6,125	1,228	2
	1,955	2,670	3,651	5,461	9,102	16,445	25,225	26,554	18,757	6,838	1,222	-
Male  Female  Non-Hispanic black, both sexes	1,115	1,435	2,012	2,940	4,565	7,558	9,995	8,699	4,714	1,250	152	-
	840	1,235	1,639	2,521	4,537	8,887	15,230	17,855	14,043	5,588	1,070	-
	1,060	1,108	1,312	1,489	1,873	2,452	2,592	2,101	1,452	554	153	2
Male Female Origin not stated 2/, both sexes Male Female	570 490 13 10	639 469 15 8	742 570 12 8 4	770 719 17 9	915 958 21 10 11	1,026 1,426 33 17	933 1,659 44 18 26	592 1,509 37 10 27	345 1,107 29 5	99 455 9 1 8	24 129 3 1 2	2 1 1
10	,	,	-	U		10	20	27	21	O	2	

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Atherosclerosis (I70)												
All origins, both sexes	93	170	289	409	723	1,294	2,112	2,682	2,382	1,246	349	_
Male	62	98	177	229	390	626	923	996	652	240	45	-
Female	31	72	112	180	333	668	1,189	1,686	1,730	1,006	304	-
Hispanic, both sexes	5	9	13	23	32	60	67	62	57	29	9	-
Male	3	7	4	12	17	31	33	29	23	10	1	-
Female	2	2	9	11	15	29	34	33	34	19	8	-
Mexican, both sexes	4	7	9	17	22	39	33	31	30	20	6	-
Male	2	5	3	8	10	24	17	18	14	6	_	-
Female	2	2	6	9	12	15	16	13	16	14	6	-
Puerto Rican, both sexes	-	1	1	3	5	4	8	4	2	-	_	-
Male	-	1	1	1	5	2	4	1	-	-	_	-
Female	-	-	-	2	_	2	4	3	2	-	_	-
Cuban, both sexes	-	_	-	1	1	7	7	9	14	3	1	-
Male	_	_	-	1	-	2	5	3	4	_	_	_
Female	-	_	-	-	1	5	2	6	10	3	1	-
Central and South American, both sexes.	1	_	-	-	1	2	3	5	1	-	1	-
Male	1	_	_	_	_	1	_	1	_	_	_	_
Female	-	_	-	-	1	1	3	4	1	-	1	-
Other and Unknown Hispanic, both sexes.	_	1	3	2	3	8	16	13	10	6	1	_
Male	_	1	_	2	2	2	7	6	5	4	1	_
Female	_	_	3	_	1	6	9	7	5	2	_	_
Non-Hispanic 1/, both sexes	88	160	275	385	690	1,232	2,043	2,618	2,324	1,214	340	_
Male	59	90	172	217	372	593	889	967	628	230	44	_
Female	29	70	103	168	318	639	1,154	1,651	1,696	984	296	_
Non-Hispanic white, both sexes	68	129	237	320	601	1,111	1,896	2,447	2,177	1,135	305	_
Male	50	74	147	186	332	537	824	918	592	218	41	_
Female	18	55	90	134	269	574	1,072	1,529	1,585	917	264	_
Non-Hispanic black, both sexes	18	27	36	52	74	107	129	136	124	68	32	_
Male	8	12	25	25	35	49	57	35	29	10	3	_
Female	10	15	11	27	39	58	72	101	95	58	29	_
Origin not stated 2/, both sexes		1	1	1	1	2	2	2	1	3		_
Male	_	1	1	_	1	2	1	_	1	_	_	_
Female	_	_	_	1	_	-	1	2	_	3	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other diseases of circulatory system (I71-I78) All origins, both sexes..... 1,248 1,826 2,806 3,830 4,511 3,703 2.184 Male.... 1,143 1,675 2,079 2,243 1,587 Female..... 1,131 1,751 2,268 2,116 1,441 Hispanic, both sexes..... Male.... 5.8 Female..... Mexican, both sexes..... Male.... 2.0 Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... 1.0 Male Female..... 1.0 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 2.4 a Male.... Female..... Non-Hispanic 1/, both sexes..... 1,188 1,745 2,701 3,716 4,374 3,607 2,130 Male.... 1,096 1,600 2,017 2,174 1,545 1,101 1,699 2,200 2,062 1,407 Female..... Non-Hispanic white, both sexes..... 1,468 2,396 3,314 3,921 3,260 1,930 1,453 1,833 Male.... 1,988 1,410 1,481 1,933 1,850 1,261 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Aortic aneurysm and dissection (I71) All origins, both sexes..... 1,137 1,795 2,416 2,715 1,949 Male.... 1,160 1,369 1,490 Female..... 1,047 1,225 1,017 2.0 Hispanic, both sexes..... 3.0 Male.... Female..... Mexican, both sexes..... 2.0 3.0 Male.... 1.0 Female..... 1.0 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,093 1,733 2,357 2,646 1,904 Male.... 1,111 1,336 1,449 1,021 1,197 Female..... 2,418 Non-Hispanic white, both sexes..... 1,574 2,155 1,753 1,024 1,339 Male.... 1,240 1,079 Female..... Non-Hispanic black, both sexes..... 5.8 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other diseases of arteries, arterioles and capillaries (I72-I78) All origins, both sexes..... 1,011 1,414 1,796 1,754 1.234 Male.... Female..... 1,043 1,099 Hispanic, both sexes..... Male.... 2.8 Female..... 1.0 2.4 2.7 Mexican, both sexes..... 2.0 Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... 1,359 1,728 1,703 1,196 Male.... 1,003 1,068 Female..... Non-Hispanic white, both sexes..... 1,159 1,503 1,507 1,047 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other disorders of circulatory system (I80-I99) All origins, both sexes..... Male.... 2.7 Female..... Hispanic, both sexes..... 2.7 Male.... Female..... 1.0 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 9.8 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... 2.8 Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Influenza and pneumonia (J10-J18) All origins, both sexes..... 1,094 1.370 1.784 2,370 4.012 7,017 11,049 12,349 10,192 4,561 1,210 Male.... 1,032 1,380 2,225 3,665 5,348 5,208 3,464 1,272 Female..... 1.787 3,352 5,701 7,141 6,728 3,289 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... 3.0 Puerto Rican, both sexes..... 3.8 Male.... Female..... Cuban, both sexes..... Male Female..... 2.7 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω 1 Q 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,001 1,257 1,660 2,190 3,740 6,596 10,569 11,866 9,843 4,396 1,140 1,278 5,117 5,003 Male.... 2,082 3,451 3,343 1,222 1,658 3,145 5,452 6,863 6,500 3,174 Female..... Non-Hispanic white, both sexes..... 1,322 1,735 3,159 5,705 9,438 10,788 8,997 4,034 1,010 2,970 3,065 1,111 Male.... 1,758 4,579 4,557 1,401 2,735 4,859 6,231 5,932 2,923 Female..... Non-Hispanic black, both sexes..... 3.0 Male.... Female..... Origin not stated 2/, both sexes..... 3.0 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Influenza (J10-J11) All origins, both sexes..... Male.... 2.2 Female..... 2.8 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.2 Female..... Non-Hispanic white, both sexes..... Male..... 1.0 1.3 5.8 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Pneumonia (J12-J18)												
All origins, both sexes	1,083	1,347	1,752	2,339	3,944	6,902	10,873	12,111	9,980	4,477	1,190	_
Male	671	780	1,018	1,362	2,185	3,609	5,279	5,116	3,412	1,250	266	-
Female	412	567	734	977	1,759	3,293	5,594	6,995	6,568	3,227	924	-
Hispanic, both sexes	87	99	115	162	256	401	454	450	335	157	66	-
	57	57	65	87	131	207	216	192	117	46	21	-
Female	30	42	50	75	125	194	238	258	218	111	45	-
Mexican, both sexes	49	53	68	84	142	214	221	230	167	82	33	-
Male	31	32	38	42	76	112	100	107	72	25	10	-
Female	18	21	30	42	66	102	121	123	95	57	23	-
Puerto Rican, both sexes	21	26	26	37	37	67	69	65	42	14	5	-
Male	14	15	12	22	17	29	34	24	13	6	2	-
Female	7	11	14	15	20	38	35	41	29	8	3	-
Cuban, both sexes	4	2	5	7	21	48	68	70	53	33	13	-
Male	3	1	4	4	13	29	37	29	12	9	4	-
Female	1	1	1	3	8	19	31	41	41	24	9	-
Central and South American, both sexes.	5	8	8	12	22	27	36	17	19	10	2	-
Male	2	5	6	8	9	13	14	7	4	1	1	-
Female	3	3	2	4	13	14	22	10	15	9	1	-
Other and Unknown Hispanic, both sexes.	8	10	8	22	34	45	60	68	54	18	13	-
Male	7	4	5	11	16	24	31	25	16	5	4	-
Female	1	6	3	11	18	21	29	43	38	13	9	-
Non-Hispanic 1/, both sexes	991	1,234	1,628	2,160	3,675	6,482	10,395	11,632	9,632	4,313	1,121	-
Male	610	713	945	1,260	2,044	3,396	5,049	4,912	3,291	1,200	244	_
Female	381	521	683	900	1,631	3,086	5,346	6,720	6,341	3,113	877	-
Non-Hispanic white, both sexes	713	951	1,294	1,708	3,098	5,606	9,269	10,561	8,790	3,954	991	-
Male	438	556	749	975	1,723	2,925	4,512	4,469	3,013	1,089	204	-
Female	275	395	545	733	1,375	2,681	4,757	6,092	5,777	2,865	787	-
Non-Hispanic black, both sexes	253	253	280	379	464	672	851	792	594	282	101	-
Male	159	140	170	248	262	360	405	312	184	71	30	-
Female	94	113	110	131	202	312	446	480	410	211	71	-
Origin not stated 2/, both sexes	5	14	9	17	13	19	24	29	13	7	3	-
Male	4	10	8	15	10	6	14	12	4	4	1	-
Female	1	4	1	2	3	13	10	17	9	3	2	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other acute lower respiratory infections (J20-J22) All origins, both sexes..... Male.... Female..... 2.7 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2. β 1.0 Female..... Non-Hispanic white, both sexes..... Male.... 2. 1.0 ρ Female..... Non-Hispanic black, both sexes..... \_ Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Acute bronchitis and bronchiolitis (J20-J21) All origins, both sexes..... 4 6 4 8 1.0 14 18 25 29 13 2 Male.... 4 5 5 4 3 11 12 Female..... 4 3 5 1.0 15 14 17 9 2 Hispanic, both sexes..... 1 1 2 1 1 Male.... Female..... 1 1 1 1 Mexican, both sexes..... 1 1 1 1 Male.... 1 1 Female..... 1 1 Puerto Rican, both sexes..... 1 Male.... Female..... 1 Cuban, both sexes..... 1 Male.... 1 Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 1 Male.... 1 Female..... Non-Hispanic 1/, both sexes..... 7 9 13 18 24 27 13 Male.... 3 2 2 4 3 10 11 4 2 15 Female..... 14 16 Non-Hispanic white, both sexes..... 9 18 21 27 12 Male.... 2. 2 2. 4 5 3 3 8 11 Female..... 3 2 6 13 16 3 1.5 Non-Hispanic black, both sexes..... 1 1 4 2 1 Male.... 1 1 1 Female..... 3 Origin not stated 2/, both sexes...... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Unspecified acute lower respiratory infection (J22) All origins, both sexes..... 4 1 2 15 17 25 10 6 Male.... 1 7 4 6 Female..... 1 3 8 13 19 10 5 Hispanic, both sexes..... 1 Male.... Female..... 1 Mexican, both sexes..... 1 Male.... Female..... 1 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male..... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 4 15 17 24 10 Male.... 1 1 1 7 4 6 1 3 8 13 10 Female..... 1 18 Non-Hispanic white, both sexes..... 1 2 15 17 23 1.0 Male.... 1 1 4 6 1 Female..... 4 2 8 13 17 10 1 Non-Hispanic black, both sexes..... 1 1 Male.... Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Chronic lower respiratory diseases (J40-J47) All origins, both sexes..... 2,279 4,205 7.549 11,323 17,067 23,398 24,170 17,842 8.601 2,459 Male.... 1,193 2,155 3.874 5,843 8,660 11,737 11,518 7,964 3,345 Female..... 1,086 2,050 3,675 5,480 8,407 11,661 12,652 9,878 5,256 1,627 Hispanic, both sexes..... Male.... Female..... 5.8 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... 2.0 Female..... Cuban, both sexes..... 5.8 2.7 Male Female..... Central and South American, both sexes. 1.0 2.4 Male Female..... Other and Unknown Hispanic, both sexes. 6.8 1 4 Male.... Female..... Non-Hispanic 1/, both sexes..... 2,197 4,070 7,385 11,032 16,624 22,849 23,564 17,382 8,315 2,359 1,153 3,779 Male.... 2,088 5,675 8,391 11,431 11,207 7,744 3,225 1,044 1,982 3,606 5,357 8,233 11,418 12,357 9,638 5,090 1,565 Female..... 22,262 Non-Hispanic white, both sexes..... 1,803 3,531 6,659 10,125 15,440 21,351 16,437 7,843 2,201 1,766 7,708 10,600 10,476 7,233 3,003 Male.... 3,360 5,157 1,765 3,299 4,968 7,732 10,751 11,786 9,204 4,840 1,478 Female..... Non-Hispanic black, both sexes..... 1,202 Male.... Female..... Origin not stated 2/, both sexes..... 2.7 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Bronchitis, chronic and unspecified (J40-J42) All origins, both sexes..... 2.0 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female.... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 2.0 Male.... R Female..... Non-Hispanic white, both sexes..... Male..... 1.5 1.3 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Emphysema (J43) All origins, both sexes..... 1,019 1.571 2,222 2,782 2,542 1,629 Male.... 1,200 1,442 1,283 Female..... 1,022 1,340 1,259 Hispanic, both sexes..... 1.0 3.0 3.8 2.0 Male.... 2.7 Female..... 3.0 Mexican, both sexes..... 3.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,006 1,537 2,187 2,725 2,484 1,587 Male.... 1,176 1,415 1,251 1,011 1,310 1,233 Female..... Non-Hispanic white, both sexes..... 1,435 2,056 2,577 2,361 1,505 1,091 1,320 1,179 Male.... 1,257 1,182 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Asthma (J45-J46) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 8.3 8.0 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... 2.4 Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other chronic lower respiratory diseases (J44,J47) All origins, both sexes..... 1.707 3,330 6,254 9,464 14,532 20,221 21,151 15,735 7,639 2,144 Male.... 1,724 3,210 4.876 7.344 10,171 10,095 7,044 2,969 Female..... 1,606 3,044 4,588 7,188 10,050 11,056 8,691 4,670 1,399 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... 2.0 Female..... 2.0 Puerto Rican, both sexes..... Male.... Female..... 2.4 Cuban, both sexes..... 2.4 Male 2.7 Female..... 2.1 2.2 Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. QQ 1 Q Male.... Female..... Non-Hispanic 1/, both sexes..... 1,655 3,231 6,123 9,227 14,151 19,755 20,629 15,339 7,383 2,055 Male.... 1,672 3,129 4,735 7,105 9,900 9,823 6,857 2,865 1,559 2,994 4,492 7,046 9,855 10,806 8,482 4,518 1,347 Female..... Non-Hispanic white, both sexes..... 1,420 2,856 5,577 8,491 13,153 18,483 19,511 14,529 6,971 1,924 1,439 9,194 9,186 Male.... 2,807 4,314 6,531 6,412 2,667 1,417 2,770 4,177 6,622 9,289 10,325 8,117 4,304 1,275 Female..... Non-Hispanic black, both sexes..... 1,022 Male.... Female..... Origin not stated 2/, both sexes..... 1.0 3.8 2.8 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Pneumoconioses and chemical effects (J60-J66,J68) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male..... 1.0 2.3 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Pneumonitis due to solids and liquids (J69) All origins, both sexes..... 1,169 2,107 3,457 3,850 2,808 1.117 Male.... 1,207 1.852 1,898 1,211 Female..... 1,605 1,952 1,597 Hispanic, both sexes..... 2.0 Male.... Female..... Mexican, both sexes..... 2.0 Male.... 2.8 Female..... 2.0 2.8 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 2.0 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,132 2,026 3,349 3,739 2,730 1,086 Male.... 1,167 1,799 1,855 1,180 1,550 1,884 1,550 Female..... 3,034 Non-Hispanic white, both sexes..... 1,777 3,439 2,501 1,025 1,639 1,708 1,086 Male.... 1,395 1,731 1,415 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other diseases of respiratory system (J00-J06, J30-J39, J67, J70-J98) All origins, both sexes..... 1,207 1,562 2,160 3,363 4,551 5,078 3,843 2.120 Male.... 1,233 1,873 2,395 2,503 1,614 Female..... 1,490 2,156 2,575 2,229 1,391 Hispanic, both sexes..... Male.... Female..... 2.0 Mexican, both sexes..... 3.0 Male.... 2.0 2.4 1.0 Female..... 2.7 5.8 Puerto Rican, both sexes..... 2.4 1.0 Male.... Female..... Cuban, both sexes..... 2.8 Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 2.4 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,120 1,470 2,015 3,153 4,293 4,838 3,713 2,044 Male.... 1,148 1,775 2,269 2,391 1,567 2.2 1,378 2,024 2,447 2,146 1,338 Female..... Non-Hispanic white, both sexes..... 1,231 1,739 2,820 3,925 4,464 3,467 1,904 1,003 1,600 2,213 Male.... 2,089 1,458 1,220 1,836 2,251 2,009 1,245 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Peptic ulcer (K25-K28) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 1.0 1.0 2.0 1.0 Male.... Female..... 1.0 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.42 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 2.8 2.1 Male.... Female..... 1.0 2.0 2.0 Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Diseases of appendix (K35-K38) All origins, both sexes..... 2.8 Male.... 1.0 Female..... 2.0 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female.... \_ Non-Hispanic 1/, both sexes..... Male.... 2.4 2.6 Female..... Non-Hispanic white, both sexes..... Male..... 2.2 2.2 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Hernia (K40-K46) All origins, both sexes..... Male.... Female..... 2.0 Hispanic, both sexes..... 1.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1.3 Female..... Non-Hispanic white, both sexes..... Male.... 2.4 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Chronic liver disease and cirrhosis (K70,K73-K74) 2,232 All origins, both sexes..... 4.145 3,580 2,989 2,608 2,511 1,501 2.8 Male.... 3,038 2,581 2,006 1,652 1,470 1,196 Female..... 1,107 1,041 1,036 Hispanic, both sexes..... Male.... Female..... 2.8 Mexican, both sexes..... Male.... Female..... 5.0 Puerto Rican, both sexes..... 5.8 Male.... Female..... Cuban, both sexes..... 1.0 Male Female..... Central and South American, both sexes. 2.0 Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 3,586 3,155 2,624 2,289 2,216 1,968 1,349 2,575 1,756 Male.... 2,260 1,469 1,306 1,062 1,011 Female..... Non-Hispanic white, both sexes..... 2,962 2,609 2,229 1,994 2,009 1,831 1,253 1,883 1,279 1,184 Male.... 2,153 1,504 Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Alcoholic liver disease (K70)												
All origins, both sexes	2,369	1,940	1,453	1,078	763	481	228	86	15	1	_	1
Male	1,772	1,453	1,096	824	568	357	170	56	12	1	_	-
Female	597	487	357	254	195	124	58	30	3	_	-	1
Hispanic, both sexes	312	228	155	117	88	71	29	2	2	_	_	-
Male	277	185	129	91	71	59	20	2	2	-	_	-
Female	35	43	26	26	17	12	9	_	-	_	_	-
Mexican, both sexes	231	149	100	73	63	49	18	2	1	-	_	-
Male	210	121	84	56	50	41	14	2	1	_	-	_
Female	21	28	16	17	13	8	4	_	-	_	_	-
Puerto Rican, both sexes	21	34	16	18	13	11	2	-	_	-	_	-
Male	16	28	12	14	11	9	1	-	-	-	-	-
Female	5	6	4	4	2	2	1	-	-	-	-	-
Cuban, both sexes	4	4	7	4	3	1	2	-	-	-	-	-
Male	4	3	7	3	3	1	2	-	-	-	-	-
Female	_	1	-	1	_	-	-	-	-	_	_	_
Central and South American, both sexes.	9	10	5	7	1	2	_	_	1	_	_	_
Male	7	8	4	4	1	2	_	_	1	_	_	_
Female	2	2	1	3	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	47	31	27	15	8	8	7	_	_	_	_	_
Male	40	25	22	14	6	6	3	_	_	_	_	_
Female	7	6	5	1	2	2	4	_	_	_	_	_
Non-Hispanic 1/, both sexes	2,042	1,704	1,293	957	673	409	199	84	13	1	_	1
Male	1,482	1,261	962	730	495	297	150	54	10	1	_	_
Female	560	443	331	227	178	112	49	30	3	_	_	1
Non-Hispanic white, both sexes	1,727	1,423	1,101	826	603	377	178	79	11	1	_	1
Male	1,274	1,058	825	632	441	274	134	51	9	1	_	_
Female	453	365	276	194	162	103	44	28	2	_	_	1
Non-Hispanic black, both sexes	239	215	145	105	49	16	15	1	2	_	_	_
Male	158	157	103	80	41	9	10	1	1	_	_	_
Female	81	58	42	25	8	7	5	_	1	_	_	_
Origin not stated 2/, both sexes	15	8	5	4	2	í	_	_	_	_	_	_
Male	13	7	5	3	2	1	_	_	_	_	_	_
Female	2	1	_	1	_	_	_	_	_	_	_	_
I CHICATE	2			_								

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Other chronic liver disease and cirrhosis (K73-K74) All origins, both sexes..... 1,776 1.640 1,536 1,530 1.748 1,751 1,273 2.7 Male.... 1,266 1,128 Female..... Hispanic, both sexes..... Male.... Female..... 2.8 Mexican, both sexes..... Male.... 2.7 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. 2.4 2.7 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,544 1,451 1,331 1,332 1,543 1,559 1,150 1,093 Male.... Female..... Non-Hispanic white, both sexes..... 1,235 1,186 1,128 1,168 1,406 1,454 1,075 Male.... Female..... Non-Hispanic black, both sexes..... 2.0 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Cholelithiasis and other disorders of gallbladder (K80-K82) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 1.0 2.7 Male.... Female..... 1.0 1.0 Mexican, both sexes..... 1.0 1.0 2.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... 1.0 Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) All origins, both sexes..... 1,245 1.742 2,221 2,949 4,170 6,329 7.729 7,162 4.733 1,711 Male.... 1.173 1,544 2,133 3,167 3,840 3,200 1,815 Female..... 1,048 1,405 2,037 3,162 3,889 3,962 2,918 1.199 Hispanic, both sexes..... 1.0 Male.... Female..... Mexican, both sexes..... 2.8 Male.... Female..... 2.0 Puerto Rican, both sexes..... Male.... 2.0 Female..... Cuban, both sexes..... 2.0 Male Female..... 2.2 Central and South American, both sexes. 1.8 Male Female..... Other and Unknown Hispanic, both sexes. 2.0 Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 1,118 1,593 2,044 2,696 3,883 6,005 7,387 6,953 4,595 1,663 1,077 Male.... 1,429 1.986 3,016 3,668 3,104 1,756 1,267 1,897 2,989 3,719 3,849 2,839 1,165 Female..... Non-Hispanic white, both sexes..... 1,292 1,838 2,907 4,821 6,285 6,080 4,009 1,456 1,544 2,509 Male.... 3,212 2,763 1,566 1,363 2,312 3,073 3,317 2,443 1,024 Female..... Non-Hispanic black, both sexes..... 1,028 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04) All origins, both sexes..... 2.4 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... \_ Non-Hispanic 1/, both sexes..... Male.... Female..... 2. Non-Hispanic white, both sexes..... Male.... ρ 1.0 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26) All origins, both sexes..... Male.... 2.0 2.0 2.0 Female..... 2.7 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1.3 2.4 Female..... 1.0 Non-Hispanic white, both sexes..... Male.... 1.0 1.0 1.3 2.7 1.3 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Renal failure (N17-N19)												
All origins, both sexes	1,208 710 498	1,705 925 780	2,178 1,147 1,031	2,894 1,518 1,376	4,103 2,098 2,005	6,222 3,113 3,109	7,608 3,782 3,826	7,062 3,157 3,905	4,676 1,793 2,883	1,695 507 1,188	317 83 234	2 1 1
Hispanic, both sexes	119 69	139 80	1,031	1,376 242 109	2,005 272 137	3,109 306 142	3,826 317 161	195 88	2,883 128 54	1,188 44 13	10	-
Female Mexican, both sexes	50 79	59 86	79 100	133 158	135 165	164 181	156 180	107 95	74 64	31	5	-
MaleFemale.	43 36	52 34	54 46	72 86	82 83	80 101	85 95	37 58	27 37	8 20	3	-
Puerto Rican, both sexes	14	26 13	29 15	35 16	43 23	33 14	44 26	28 10	13	3	2	
FemaleCuban, both sexes	5 5	13 3	14 6	19 10	20 22	19 36	18 34	18 30	9 20	3 5	1 -	-
Male Female	5 -	2 1	3	6 4	10 12	23 13	20 14	17 13	12 8	2	-	-
Central and South American, both sexes. Male	5 4	5 2	15 9	13 7	18 10	19 9	16 7	21 12	11 3	-		-
Female Other and Unknown Hispanic, both sexes.	1 16	3 19	6 22	6 26	8 24	10 37	9 43	9 21	8 20	- 8	2	-
Male Female	8	11	12 10	8 18	12 12	16 21	23	12 9	8	5	1	-
Non-Hispanic 1/, both sexes Male	1,085 639	1,562 842	2,003	2,646 1,406	3,820 1,953	5,902 2,964	7,268 3,611	6,860 3,065	4,541 1,736	1,647	307 78	1
Female Non-Hispanic white, both sexes	446 579	720 924	950 1,260	1,240	1,867 2,853	2,938 4,735	3,657 6,181	3,795 5,995	2,805	1,154	229 247	1 1
Male Female Non-Hispanic black, both sexes	340 239 471	527 397 585	679 581 682	978 822 750	1,516 1,337 859	2,466 2,269 1,016	3,161 3,020 952	2,729 3,266 744	1,551 2,415 517	428 1,014 189	62 185 53	1
MaleFemale	280 191	293 292	336 346	375 375	387 472	429 587	388 564	282 462	154 363	58 131	12 41	-
Origin not stated 2/, both sexes Male	4	4 3	3 1	575 6 3	11	14	23	7 4	7	4	-	1 1
Female	2	1	2	3	3	7	13	3	4	3	_	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of												
Diseases, Tenth Revision, 1992),												
Hispanic origin, race for	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years	Nae not
non-Hispanic population, and sex	years	years	years	years	years	years	years	years	years	years	and over	_
non-hispanic population, and sex	years	years	years	years	years	years	years	years	years	years	and over	stateu
Other disorders of kidney (N25,N27)												
All origins, both sexes	_	3	4	4	3	6	5	5	_	_	_	_
Male	_	2	2	1	-	2	2	1	_	_	_	_
Female	_	1	2	3	3	4	3	4	_	_	_	_
Hispanic, both sexes	_	1	_	_	_	_	_	_	_	_	_	_
Male	_	1	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Mexican, both sexes	_	1	_	_	_	_	_	_	_	_	_	_
Male	_	1	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Puerto Rican, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Cuban, both sexes	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Central and South American, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Other and Unknown Hispanic, both sexes.	_	_	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_	_	_
Female	_	_	_	_	_	_	_	_	_	_	_	_
Non-Hispanic 1/, both sexes	_	2	4	4	3	6	5	5	_	_	_	_
Male		1	2	1	_	2	2	1				
Female		1	2	3	3	1	3	1				
Non-Hispanic white, both sexes	_	2	4	4	3		3	5	_	_		_
Male	_	1	2	1	3	2	2	1	_	_	_	_
Female	_	1	2	3	3	3	2	1	_	_	_	_
Non-Hispanic black, both sexes	_		۷	3	3	3 1	∠ 1	4	_	_	_	_
Male			_			_	_	_				
Female	_	_	_	_	_	1	- 1	_	_	_	_	_
Origin not stated 2/, both sexes	-	_	-	_	-	1	1	_	_	_	_	_
Male	-	_	-	_	-	_	-	_	_	_	_	_
Female	-	_	-	_	_	_	-	-	_	_	_	_
rcmarc	_	_	_	_	_	_	_	_	_	_	_	_

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Infections of kidney (N10-N12,N13.6,N15.1) All origins, both sexes..... 2.8 Male.... 1.0 Female..... 1.8 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female.... Non-Hispanic 1/, both sexes..... Male.... 1.0 1.5 2.3 Female..... Non-Hispanic white, both sexes..... Male.... 2.2 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Hyperplasia of prostate (N40) All origins, both sexes..... 3 3 1.0 14 31 58 119 115 109 41 3 Male.... 3 3 1.0 14 31 58 119 115 109 41 3 Female..... . . . . . . . . . . . . Hispanic, both sexes..... 2 1 5 1 3 4 3 Male.... 1 5 1 3 4 3 2 Female..... Mexican, both sexes..... 3 2 2 1 1 1 Male.... 1 3 1 2 1 2 Female..... Puerto Rican, both sexes..... 2 Male.... 2 Female..... . . . . . . Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... . . . . . . . . . . . . Other and Unknown Hispanic, both sexes. 2 1 2 2 Male.... 2 1 2 2 Female..... Non-Hispanic 1/, both sexes..... 3 3 9 14 26 56 116 110 106 39 3 9 Male.... 3 3 14 26 56 116 110 106 39 3 Female..... . . . . . . . . . Non-Hispanic white, both sexes..... 22 49 102 103 97 36 8 49 103 97 36 Male.... 1 3 13 2.2 102 3 Female..... . . . Non-Hispanic black, both sexes..... 1 5 8 4 2 2 5 8 7 Male.... 1 1 4 4 2 Female..... Origin not stated 2/, both sexes...... 1 1 1 1 Male.... Female..... .

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Inflammatory diseases of female pelvic organs (N70-N76) All origins, both sexes..... 3 8 7 7 9 10 11 17 11 6 Male.... Female..... 3 8 7 9 10 11 17 11 6 Hispanic, both sexes..... 1 1 2 1 1 Male.... Female..... 1 1 2 1 1 Mexican, both sexes..... 1 Male.... Female..... 1 Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1 Male Female..... 1 Central and South American, both sexes. 1 Male . . . . . . . . . Female..... 1 Other and Unknown Hispanic, both sexes. \_ 1 1 Male.... . . . Female..... 7 7 Non-Hispanic 1/, both sexes..... 3 6 8 8 10 16 11 6 Male.... . . . Female..... 3 6 1.0 16 11 8 6 Non-Hispanic white, both sexes..... 2 6 7 5 7 7 6 15 10 5 Male.... 7 5 15 10 5 Female..... 2 6 6 Non-Hispanic black, both sexes..... 1 1 1 1 1 4 1 1 1 Male.... Female..... 1 4 Origin not stated 2/, both sexes...... 1 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Pregnancy, childbirth and the puerperium (000-099) All origins, both sexes..... 40 1 1 2 Male.... Female..... 1 1 2 Hispanic, both sexes..... 5 Male.... Female..... Mexican, both sexes..... 4 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male . . . . . . . . . . . . Female..... Other and Unknown Hispanic, both sexes. 1 \_ Male.... Female..... 2 Non-Hispanic 1/, both sexes..... 35 1 1 Male.... . . . . . . . . . . . . . . . Female..... 35 1 1 Non-Hispanic white, both sexes..... 27 1 1 Male.... 27 Female..... 1 1 1 Non-Hispanic black, both sexes..... 7 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Pregnancy with abortive outcome (000-007) All origins, both sexes..... 1 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male . . . . . . . . . . . . . . . Female..... Other and Unknown Hispanic, both sexes. \_ Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... . . . . . . . . . . . . . . . . . . Female..... Non-Hispanic white, both sexes..... Male.... 1 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other complications of pregnancy, childbirth and the puerperium (010-099) All origins, both sexes..... 39 1 1 2 Male.... Female..... 39 1 1 2 Hispanic, both sexes..... 5 Male.... Female..... Mexican, both sexes..... 4 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male . . . . . . . . . . . . Female..... Other and Unknown Hispanic, both sexes. 1 \_ Male.... Female..... 2 Non-Hispanic 1/, both sexes..... 34 1 1 Male.... . . . . . . . . . . . . . . . Female..... 34 1 1 Non-Hispanic white, both sexes..... 26 1 1 Male.... 26 Female..... 1 1 1 Non-Hispanic black, both sexes..... 7 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Certain conditions originating in the perinatal period (P00-P96) All origins, both sexes..... 1 1 2 1 1 1 Male.... 1 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male..... Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male..... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1 2 1 1 1 Female..... Non-Hispanic white, both sexes..... 1 Male.... 1 2. 1 Female..... 1 Non-Hispanic black, both sexes..... Male..... Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) All origins, both sexes..... 2.8 Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 8.0 Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... 1.0 Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) All origins, both sexes..... 1,132 1.047 1,014 1,309 2,075 3,084 3,844 3,806 2,183 Male.... 1.232 1,272 Female..... 1,126 1,852 2,572 2,824 1.793 Hispanic, both sexes..... Male.... 2.6 2.4 Female..... 1.8 3.8 Mexican, both sexes..... 2.8 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... 1.0 Male Female..... Central and South American, both sexes. 1.0 Male Female..... Other and Unknown Hispanic, both sexes. a Ω Male.... Female..... Non-Hispanic 1/, both sexes..... 1,067 1,255 2,015 3,013 3,776 3,722 2,131 Male.... 1,207 1,244 1,092 1,806 2,532 2,765 1,749 Female..... Non-Hispanic white, both sexes..... 1,035 1,759 2,688 3,453 3,396 1,975 1,081 1,139 Male.... 1,607 2,314 2,509 1,628 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... 1.0 1.0 Male.... Female..... 2.0

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13,0-N13,5,N13,7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98) 22,278 All origins, both sexes..... 8.777 9.577 9,909 11,101 15,003 30,845 32,567 25,231 10.926 2,304 12 Male.... 5.335 5,467 5,329 5,702 7,103 9.776 11,646 10,289 6,189 1.892 276 Female..... 3,442 4,110 4,580 5,399 7,900 12,502 19,199 22,278 19,042 9,034 2.028 Hispanic, both sexes..... 563 567 523 570 713 941 953 839 665 275 72 Male.... 336 273 289 331 397 343 302 180 14 Female..... 185 231 250 281 382 544 610 537 485 221 58 Mexican, both sexes..... 324 336 311 342 391 532 487 415 307 137 41 Male.... 214 195 161 175 182 230 180 162 98 29 10 Female..... 110 141 150 167 209 302 307 253 209 108 31 Puerto Rican, both sexes..... 79 93 70 85 90 122 125 93 56 22 4 Male.... 52 56 35 44 39 51 44 35 16 6 1 Female..... 2.7 37 35 41 51 71 81 58 40 16 3 Cuban, both sexes..... 33 2.8 29 53 63 108 140 140 146 55 14 Male 22 20 18 29 31 47 43 40 31 7 2 11 8 11 24 32 61 97 100 115 48 12 Female..... 33 26 34 55 49 44 55 37 5 Central and South American, both sexes. 2.8 13 Male 2.0 10 13 15 26 17 17 16 9 2 Female..... 13 16 15 19 29 32 27 39 28 11 Other and Unknown Hispanic, both sexes. 94 Ω4 25 56 114 130 157 136 119 48 Ω Male.... 70 55 46 26 53 52 59 49 26 10 1 Female..... 24 29 39 30 61 78 98 87 93 38 7 Non-Hispanic 1/, both sexes..... 8,175 8,981 9,359 10,504 14,257 21,301 29,847 31,675 24,520 10,635 2,224 Male.... 4,928 5,112 5,036 5,395 6,758 9,362 11,287 9,969 5,996 1.837 262 3,247 3,869 4,323 5,109 7,499 11,939 18,560 21,706 18,524 8,798 1,962 Female..... Non-Hispanic white, both sexes..... 6,250 7,096 7,627 8,658 12,129 18,685 26,904 29,084 22,478 9,718 1,938 4,141 4,446 10,180 5,489 1,626 218 Male.... 3,838 4,080 5,787 8,187 9,139 2,412 3,016 3,486 4,212 6,342 10,498 16,724 19,945 16,989 8,092 1,720 Female..... Non-Hispanic black, both sexes..... 1,718 1,665 1,520 1,596 1,847 2,257 2,501 2,194 1,732 782 245 1 965 914 782 821 854 1,016 908 679 414 168 35 1 Male.... Female..... 753 751 738 775 993 1,241 1,593 1,515 1,318 614 210 Origin not stated 2/, both sexes..... 39 29 27 27 33 36 45 53 46 16 8 29 19 20 18 14 17 16 18 13 1 Male....

Female.....

10

10

7

9

19

19

29

35

33

15

9

3

2

2

2

2

6

3

3

5

2

3

4

4

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years and over stated vears vears vears Accidents (unintentional injuries) (V01-X59,Y85-Y86) 5,408 All origins, both sexes..... 7.516 4.243 3,819 4.297 5,900 7,557 6,780 4,666 1,709 Male.... 5,386 3,752 2.810 2,448 2,544 3,222 3,808 2,962 1,705 Female..... 2,130 1,656 1,433 1,371 1,753 2,678 3,749 3,818 2,961 1,233 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 2.0 2.4 Male.... Female..... Cuban, both sexes..... 2.8 Male Female..... Central and South American, both sexes. 2.4 1.0 Male Female..... Other and Unknown Hispanic, both sexes. 3.0 Male.... Female..... Non-Hispanic 1/, both sexes..... 6,953 5,013 3,925 3,567 4,052 5,641 7,283 6,604 4,536 1,662 2,277 Male.... 4,956 3,475 2,586 2,400 3,072 3,666 2,880 1,660 1,997 1,538 1,339 1,290 1,652 2,569 3,617 3,724 2,876 1,203 Female..... Non-Hispanic white, both sexes..... 5,620 4,106 3,284 3,027 3,505 5,044 6,730 6,221 4,274 1,585 3,982 2,140 2,083 3,379 Male.... 2,815 1,944 2,761 2,704 1,546 1,638 1,291 1,144 1,083 1,422 2,283 3,351 3,517 2,728 1,149 Female..... Non-Hispanic black, both sexes..... 1,113 Male.... Female..... Origin not stated 2/, both sexes..... 1.0 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Transport accidents (V01-V99,Y85)												
All origins, both sexes	3,178	2,510	2,000	1,689	1,540	1,749	1,561	930	328	58	4	27
Male	2,325	1,773	1,341	1,077	921	1,001	904	501	171	27	2	21
Female	853	737	659	612	619	748	657	429	157	31	2	6
Hispanic, both sexes	247	185	168	114	102	81	69	24	13	5	-	6
Male	185	118	111	75	60	54	43	11	8	1	-	5
Female	62	67	57	39	42	27	26	13	5	4	-	1
Mexican, both sexes	153	126	117	62	51	44	34	12	6	4	-	3
Male	111	86	81	40	30	30	23	6	4	1	-	2
Female	42	40	36	22	21	14	11	6	2	3	_	1
Puerto Rican, both sexes	20	16	9	9	9	4	3	1	4	_	_	-
Male	16	8	9	6	6	3	2	-	1	_	_	-
Female	4	8	_	3	3	1	1	1	3	_	_	-
Cuban, both sexes	15	7	10	17	18	11	17	5	2	_	-	-
Male	11	4	6	13	15	5	10	4	2	_	-	-
Female	4	3	4	4	3	6	7	1	_	_	_	_
Central and South American, both sexes.	29	15	14	8	15	8	1	2	1	1	_	2
Male	24	8	4	6	5	3	_	1	1	_	_	2
Female	5	7	10	2	10	5	1	1	_	1	_	_
Other and Unknown Hispanic, both sexes.	30	21	18	18	9	14	14	4	_	_	_	1
Male	23	12	11	10	4	13	8	_	_	_	_	1
Female	7	9	7	8	5	1	6	4	_	_	_	_
Non-Hispanic 1/, both sexes	2,916	2,319	1,827	1,569	1,435	1,664	1,487	906	314	53	4	9
Male	2,128	1,650	1,226	996	859	943	858	490	162	26	2	7
Female	788	669	601	573	576	721	629	416	152	27	2	2
Non-Hispanic white, both sexes	2,350	1,948	1,550	1,345	1,248	1,467	1,375	857	292	50	4	9
Male	1,718	1,387	1,031	858	745	838	782	465	148	26	2	7
Female	632	561	519	487	503	629	593	392	144	24	2	2
Non-Hispanic black, both sexes	435	286	210	156	120	133	78	32	17	3	_	_
Male	326	202	151	107	78	79	56	17	12	_	_	_
Female	109	84	59	49	42	54	22	15	5	3	_	_
Origin not stated 2/, both sexes	15	6	5	6	3	4	5	-	1	_	_	12
Male	12	5	4	6	2	4	3	_	1	_	_	9
	3	1	1	-	1	4	2	_	_	_	_	3
Female	3	1	T	_	1	_	2	_	_	_	_	3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2) All origins, both sexes..... 2,854 2,293 1,800 1,547 1.427 1,683 1,497 Male.... 2,055 1,596 1.174 2.0 Female..... Hispanic, both sexes..... Male.... Female..... 3.8 Mexican, both sexes..... Male.... Female..... 2.1 Puerto Rican, both sexes..... 2.0 Male.... Female..... Cuban, both sexes..... 1.0 1.0 Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1 Q 1 4 Male.... Female..... Non-Hispanic 1/, both sexes..... 2,614 2,113 1,641 1,434 1,326 1,602 1,423 1,878 1,072 Male.... 1,483 2.5 Female..... Non-Hispanic white, both sexes..... 2,098 1,764 1,391 1,225 1,151 1,417 1,313 1,510 1,239 Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9) All origins, both sexes..... 5.8 Male.... 2.0 Female..... 1.8 Hispanic, both sexes..... 1.0 Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 8.3 Male.... 3.0 1.3 2. Female.... Non-Hispanic white, both sexes..... Male.... 2.8 2.6 Female.... Non-Hispanic black, both sexes..... Male..... Female..... Origin not stated 2/, both sexes..... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85) All origins, both sexes..... 2.2 Male.... Female..... 1.0 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.7 2. ρ Female..... Non-Hispanic white, both sexes..... Male.... 2.0 1.5 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male....

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	_
Nontransport accidents (W00-X59,Y86)												
All origins, both sexes	4,338 3,061 1,277 273 210	2,898 1,979 919 189 142	2,243 1,469 774 136 103	2,130 1,371 759 124 83	2,757 1,623 1,134 133 76	4,151 2,221 1,930 168 89	5,996 2,904 3,092 190 92	5,850 2,461 3,389 143 65	4,338 1,534 2,804 107 34	1,651 449 1,202 41 16	288 65 223 5 3	61 53 8 12 12
Female. Mexican, both sexes Male	63 139 117 22	47 97 71 26	33 72 57 15	41 63 41 22	57 69 41 28	79 92 47 45	98 88 44 44	78 70 32 38	73 58 21 37	25 21 8 13	2 3 1 2	11 11 -
Puerto Rican, both sexes	45 31 14 14	34 27 7 16	14 10 4 11	11 8 3 15	17 11 6 15	21 12 9 26	21 12 9 26	16 6 10 23	10 6 4 16	4 2 2 6	- - - 1	- - -
Male  Female  Central and South American, both sexes.	11 3 20 15	12 4 12	8 3 10	13 2 13	8 7 10	10 16 5	15 11 12 6	8 15 8	5 11 6	3 3 5	1 - -	- - -
Male Female Other and Unknown Hispanic, both sexes. Male	5 55 36	1 30 21	1 29 19	6 22 14	5 22 11	2 24 17	6 43 15	4 26 15	6 17 2	4 5 2	- 1 1	- 1 1
Female. Non-Hispanic 1/, both sexes Male. Female.	19 4,037 2,828 1,209	9 2,694 1,825 869	10 2,098 1,360 738	8 1,998 1,281 717	11 2,617 1,541 1,076	7 3,977 2,129 1,848	28 5,796 2,808 2,988	11 5,698 2,390 3,308	15 4,222 1,498 2,724	3 1,609 433 1,176	283 62 221	- 29 24 5
Non-Hispanic white, both sexes  Male  Female  Non-Hispanic black, both sexes	3,270 2,264 1,006 678	2,158 1,428 730 461	1,734 1,109 625 294	1,682 1,086 596 262	2,257 1,338 919 280	3,577 1,923 1,654 320	5,355 2,597 2,758 326	5,364 2,239 3,125 224	3,982 1,398 2,584 186	1,535 410 1,125 56	260 57 203 21	26 22 4 1
Male	498 180 28 23	345 116 15	203 91 9	166 96 8	169 111 7 6	170 150 6 3	162 164 10	86 138 9	78 108 9	17 39 1	5 16 -	1 - 20 17
Female	5	3	3	1	1	3	6	3	7	1	-	3

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Falls (W00-W19) All origins, both sexes..... 1,370 2,246 3,436 3,447 2,433 Male.... 1,189 1,669 1,465 Female..... 1,057 1,767 1,982 1,542 Hispanic, both sexes..... Male.... Female..... 2.0 Mexican, both sexes..... 3.0 5.0 3.0 Male.... 2.0 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Ω 2.8 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,300 2,146 3,318 3,342 2,364 Male.... 1,138 1,610 1,412 1,008 1,708 1,930 1,494 Female..... 3,145 Non-Hispanic white, both sexes..... 1,169 2,002 3,204 2,269 1,343 Male.... 1,069 1,523 1,622 1,861 1,445 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Accidental discharge of firearms (W32-W34) All origins, both sexes..... 65 25 1.0 2.0 2.0 15 13 5 3 Male.... 2.0 9 15 18 12 11 5 3 Female..... 9 5 1 5 2 3 2 Hispanic, both sexes..... 1 Male.... 1 Female..... 1 Mexican, both sexes..... 3 1 Male.... 1 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 2 Male.... Female..... Non-Hispanic 1/, both sexes..... 58 24 10 20 20 15 13 3 Male.... 50 19 9 15 18 12 11 3 R 5 5 3 2 Female..... 1 2. Non-Hispanic white, both sexes..... 54 22 19 17 13 11 3 Male..... 48 18 14 16 10 9 3 6 4 5 3 2 Female..... 1 1 Non-Hispanic black, both sexes..... 4 2 2 1 3 2 2 1 2 2 1 Male..... 1 2 1 2 Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Accidental drowning and submersion (W65-W74) All origins, both sexes..... Male.... Female..... 2.4 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... \_ Non-Hispanic 1/, both sexes..... Male.... 2.5 Female..... Non-Hispanic white, both sexes..... Male.... 2.1 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Accidental exposure to smoke, fire and flames (X00-X09) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... 1.0 Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Accidental poisoning and exposure to noxious substances (X40-X49) All origins, both sexes..... 2.337 1.111 Male.... 1,550 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 Q Male.... Female..... Non-Hispanic 1/, both sexes..... 2,172 1,018 1,428 Male.... Female..... Non-Hispanic white, both sexes..... 1,747 Male.... 1,118 Female..... Non-Hispanic black, both sexes..... 2.4 3.0 1.0 Male.... Female..... 1.0 Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86) All origins, both sexes..... 1,425 2,080 2,077 1,749 Male.... Female..... 1,103 1,227 1,172 Hispanic, both sexes..... 5.0 5.8 Male.... 1.8 Female..... 1.0 2.0 Mexican, both sexes 2.8 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a 1.0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,380 2,018 2,041 1,709 Male.... 2.5 1,068 1,202 1,143 Female..... Non-Hispanic white, both sexes..... 1,207 1,834 1,890 1,595 Male.... 1,114 1,072 Female..... Non-Hispanic black, both sexes..... Male.... Female..... 5.8 Origin not stated 2/, both sexes..... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Intentional self-harm (suicide) (\*U03.X60-X84.Y87.0) All origins, both sexes..... 3.163 2,349 1,662 1,133 1,146 1,189 Male.... 2,329 1,787 1,301 1,005 Female..... Hispanic, both sexes..... 5.8 1.0 Male.... Female..... Mexican, both sexes..... 2.0 1.0 Male.... Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. a Male.... Female..... Non-Hispanic 1/, both sexes..... 3,026 2,240 1,601 1,090 1,094 1,161 2,219 Male.... 1,705 1,251 2.5 Female..... Non-Hispanic white, both sexes..... 2,832 2,088 1,496 1,022 1,021 1,107 2,083 1,575 1,179 Male.... 2.5 Female..... Non-Hispanic black, both sexes..... 2.4 Male.... Female..... Origin not stated 2/, both sexes..... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Intentional self-harm (suicide) by discharge of firearms (X72-X74) All origins, both sexes..... 1.565 1,283 1,045 Male.... 1,296 1,101 Female..... Hispanic, both sexes..... 2.8 Male.... Female..... Mexican, both sexes..... 1.8 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 1,509 1,247 1,014 1,249 Male.... 1,069 Female..... Non-Hispanic white, both sexes..... 1,442 1,179 1,008 Male.... 1.196 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0) All origins, both sexes..... 1.598 1.066 Male.... 1,033 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 3.0 Male.... 2.7 Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1 0 Male.... Female..... Non-Hispanic 1/, both sexes..... 1,517 Male.... Female..... Non-Hispanic white, both sexes..... 1,390 Male.... Female..... Non-Hispanic black, both sexes..... 2.1 Male.... Female..... Origin not stated 2/, both sexes...... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 1.8 Male.... 2.8 Female..... 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... \_ Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... 2.1 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Assault (homicide) by discharge of firearms (\*U01.4,X93-X95) All origins, both sexes..... 2.2 Male.... 2.0 Female..... 5.8 Hispanic, both sexes..... 2.0 Male.... 2.0 Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 1.0 Female..... 1.0 Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 2.7 1.0 Male.... Female..... 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... 1.0 Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... \_ Non-Hispanic 1/, both sexes..... Male.... Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years	
Legal intervention (Y35,Y89.0)												
All origins, both sexes	27	9	3	4	3	3	2	_	_	_	_	_
Male	27	8	3	3	3	3	2	-	-	-	_	_
Female	_	1	_	1	_	-	_	_	-	-	-	_
Hispanic, both sexes	1	_	-	_	_	-	_	_	-	-	-	_
Male	1	-	-	_	_	-	-	-	-	-	-	-
Female	-	-	-	_	_	-	-	-	-	-	-	-
Mexican, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	_	-	-	_	-	-	-	-	_	_
Female	-	-	_	-	-	_	-	-	-	-	_	_
Puerto Rican, both sexes	-	-	_	-	-	_	-	-	-	-	_	_
Male	-	-	_	-	-	_	-	-	-	-	_	_
Female	-	_	-	-	-	_	-	-	-	-	_	-
Cuban, both sexes	1	_	-	-	-	_	-	-	-	-	_	-
Male	1	_	-	-	-	_	-	-	-	-	_	-
Female	-	_	-	-	-	_	-	-	-	-	_	-
Central and South American, both sexes.	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	_	-	-	-	_	-	-	-	-	_	-
Female	-	_	-	-	-	_	-	-	-	-	_	-
Other and Unknown Hispanic, both sexes.	-	_	-	-	-	_	-	-	-	-	_	-
Male	-	_	-	-	-	_	-	-	-	-	_	-
Female	-	_	-	-	-	_	-	-	-	-	_	-
Non-Hispanic 1/, both sexes	26	9	3	4	3	3	2	-	-	-	_	_
Male	26	8	3	3	3	3	2	-	-	-	_	_
Female	-	1	-	1	-	_	-	-	-	-	_	_
Non-Hispanic white, both sexes	17	5	3	2	3	3	2	-	-	-	_	_
Male	17	5	3	1	3	3	2	-	-	-	_	_
Female	-	-	-	1	-	_	-	-	-	-	_	-
Non-Hispanic black, both sexes	8	4	-	2	-	_	-	-	-	-	_	-
Male	8	3	-	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	_	_	_	_	_	_	-	_	_	-	_	-

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) All origins, both sexes..... Male.... 2.8 Female..... Hispanic, both sexes..... Male.... Female..... 1.0 Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.2 Female..... Non-Hispanic white, both sexes..... Male.... 2.5 1.5 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Discharge of firearms, undetermined intent (Y22-Y24) All origins, both sexes..... 17 11 9 11 5 6 2 3 Male.... 8 6 9 5 3 Female..... 4 3 3 2 1 1 1 1 Hispanic, both sexes..... 1 Male.... Female..... 1 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female..... 1 Central and South American, both sexes. Male Female..... 1 Other and Unknown Hispanic, both sexes. Male.... Female..... 2 Non-Hispanic 1/, both sexes..... 16 10 9 11 6 Male.... 12 8 6 9 1 5 1 4 2 3 2 Female..... 1 1 1 1 Non-Hispanic white, both sexes..... 13 10 6 9 Male.... 8 6 8 1 5 1 4 3 2 1 Female..... 1 1 1 1 Non-Hispanic black, both sexes..... 3 1 1 3 Male.... 1 Female..... Origin not stated 2/, both sexes..... 3 3 Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality,

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9) All origins, both sexes..... 2.4 Male.... Female..... Hispanic, both sexes..... Male.... Female..... 1.0 Mexican, both sexes..... 1.0 Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male.... Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... 2.1 2.1 Female..... Non-Hispanic white, both sexes..... Male.... 2.0 1.5 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... 

Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Operations of war and their sequelae (Y36,Y89.1) All origins, both sexes..... 3 1 1 5 9 Male.... 1 5 9 Female..... Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... 1 Male..... 1 Female..... Puerto Rican, both sexes..... Male.... Female.... Cuban, both sexes..... Male..... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... 3 8 Male.... 3 1 1 5 8 1 Female..... Non-Hispanic white, both sexes..... 3 1 8 Male.... 3 1 1 5 8 1 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes..... Male..... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years years and over stated vears Complications of medical and surgical care (Y40-Y84,Y88) All origins, both sexes..... Male.... Female..... Hispanic, both sexes..... 1.0 Male.... Female..... 1.0 Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female.... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... Female..... Non-Hispanic 1/, both sexes..... Male.... R Female..... Non-Hispanic white, both sexes..... Male.... Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... Male.... Female.....

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years and over stated Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 3/ All origins, both sexes..... 3,793 3,109 2,333 1.690 1.202 Male.... 2,919 2,403 1,821 1,304 Female..... Hispanic, both sexes..... Male.... Female..... 2.0 1.0 Mexican, both sexes..... Male.... Female..... 3.0 2.0 Puerto Rican, both sexes..... Male.... 1.8 Female..... Cuban, both sexes..... Male.... Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. 1.0 2.4 Male.... Female..... Non-Hispanic 1/, both sexes..... 3,365 2,803 2,110 1,525 1,091 2,540 Male.... 2,153 1,631 1,172 2.3 Female..... Non-Hispanic white, both sexes..... 2,796 2,322 1,746 1,261 1,785 Male.... 2,123 1,348 Female..... Non-Hispanic black, both sexes..... Male.... Female..... Origin not stated 2/, both sexes...... 1.0 Male.... Female..... 

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not years and over stated

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 3/

All origins, both sexes	3,616	1,819	832	438	347	349	270	179	98	19	8	12
Male	2,247	1,043	453	222	171	160	118	82	40	5	3	7
Female	1,369	776	379	216	176	189	152	97	58	14	5	5
Hispanic, both sexes	207	138	47	24	15	17	7	7	5	_	_	1
	147	97	35	17	10	9	4	6	1	-	-	1
Female	60	41	12	7	5	8	3	1	4	-	-	-
Mexican, both sexes	99	62	23	14	7	6	2	4	3	-	-	1
Male	77	47	17	10	5	2	1	3	1	-	-	1
Female	22	15	6	4	2	4	1	1	2	-	-	-
Puerto Rican, both sexes	48	33	3	4	2	2	1	-	1	-	-	-
Male	36	27	2	3	2	1	1	-	-	-	-	-
Female	12	6	1	1	-	1	-	-	1	-	-	-
Cuban, both sexes	11	11	5	-	2	4	3	1	-	-	-	-
Male	5	5	4	-	1	2	1	1	-	-	-	-
Female	6	6	1	-	1	2	2	-	-	-	-	-
Central and South American, both sexes.	9	8	4	1	-	3	-	1	-	-	-	-
Male	5	3	4	1	-	2	-	1	-	-	-	-
Female	4	5	-	-	-	1	-	-	-	-	-	-
Other and Unknown Hispanic, both sexes.	40	24	12	5	4	2	1	1	1	-	-	-
Male	24	15	8	3	2	2	1	1	-	-	-	-
Female	16	9	4	2	2	-	-	-	1	-	-	-
Non-Hispanic 1/, both sexes	3,390	1,676	782	413	331	330	262	172	93	19	8	6
Male	2,084	941	417	204	161	150	114	76	39	5	3	2
Female	1,306	735	365	209	170	180	148	96	54	14	5	4
Non-Hispanic white, both sexes	2,735	1,346	647	347	303	304	240	162	82	16	8	4
Male	1,623	701	319	152	144	132	105	72	35	2	3	1
Female	1,112	645	328	195	159	172	135	90	47	14	5	3
Non-Hispanic black, both sexes	592	299	115	57	23	20	17	7	9	3	-	1
Male	419	223	86	45	16	15	8	2	4	3	-	1
Female	173	76	29	12	7	5	9	5	5	-	-	-
Origin not stated 2/, both sexes	19	5	3	1	1	2	1	-	-	-	-	5
Male	16	5	1	1	_	1	-	-	-	-	-	4
Female	3	-	2	-	1	1	1	-	-	-	-	1

Worktable Orig210F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004."

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not non-Hispanic population, and sex years years years years years years years years and over stated vears vears Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 3/ All origins, both sexes..... 2,089 1,603 1.256 2.0 Male.... 1.719 1,349 1.042 Female..... 2.7 Hispanic, both sexes..... Male.... Female..... Mexican, both sexes..... Male.... Female..... Puerto Rican, both sexes..... Male.... Female..... Cuban, both sexes..... Male Female..... Central and South American, both sexes. Male Female..... Other and Unknown Hispanic, both sexes. Male.... 1.8 Female..... Non-Hispanic 1/, both sexes..... 1,978 1,541 1,205 Male.... 1.626 1,296 Female..... Non-Hispanic white, both sexes..... 1,722 1,369 1,111 Male..... 1,411 1.147 Female..... Non-Hispanic black, both sexes..... Male.... Female..... 3.0 Origin not stated 2/, both sexes...... Male.... Female..... 

Source: CDC/NCHS, National Vital Statistics System, Mortality.

<sup>-</sup> Quantity zero.

<sup>...</sup> Category not applicable.

<sup>1/</sup> Includes races other than white and black.

<sup>2/</sup> Includes deaths for which Hispanic origin was not reported on the death certificate.

<sup>3/</sup> Included in selected categories.