

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All origins 3/, both sexes.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7
Male.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8
Female.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3
Hispanic, both sexes.....	296.2	586.9	27.3	12.6	16.3	64.2	87.3	77.1	79.4	112.7
Male.....	321.1	636.5	30.2	13.8	19.8	96.5	130.9	108.8	109.7	151.5
Female.....	269.7	535.1	24.3	11.3	12.7	29.7	33.9	37.7	44.2	68.6
Mexican, both sexes.....	254.4	569.4	28.9	12.4	16.8	68.2	88.5	76.1	76.4	105.9
Male.....	280.2	615.0	31.7	13.6	20.3	103.2	133.4	108.6	105.2	141.9
Female.....	225.7	521.8	25.9	11.1	13.1	29.9	31.7	33.9	42.0	62.4
Puerto Rican, both sexes.....	406.2	666.1	24.3	12.1	14.9	43.7	85.6	91.2	111.8	174.1
Male.....	450.8	*	28.9	*	18.7	62.2	126.5	132.6	169.6	217.3
Female.....	362.7	*	*	*	*	24.2	39.8	51.6	60.3	128.3
Cuban, both sexes.....	762.0	*	*	*	*	51.7	*	*	51.6	93.1
Male.....	761.8	*	*	*	*	*	*	*	*	*
Female.....	762.2	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	293.0	675.4	25.1	13.8	16.2	63.5	85.2	77.9	81.5	110.4
Male.....	314.9	750.9	27.6	15.4	19.6	96.0	124.8	104.5	112.9	155.2
Female.....	271.1	599.9	22.4	12.3	12.4	32.3	38.5	45.6	45.6	65.5
Non-Hispanic 5/, both sexes.....	899.4	705.6	30.5	15.2	19.1	66.2	95.2	100.3	113.8	158.3
Male.....	900.9	778.7	32.9	17.0	22.1	89.6	139.3	142.5	150.4	201.3
Female.....	898.0	628.9	28.1	13.4	15.9	41.7	49.7	58.1	77.5	116.0
Non-Hispanic white, both sexes.....	967.8	564.4	26.8	13.5	17.4	63.4	87.2	89.1	102.2	145.2
Male.....	957.4	625.2	29.0	15.2	20.4	83.4	126.6	126.2	135.2	185.6
Female.....	977.7	500.7	24.4	11.6	14.3	42.4	46.3	51.3	68.7	104.4
Non-Hispanic black, both sexes.....	768.8	1,297.2	45.5	22.3	26.2	83.7	143.3	174.2	198.6	263.3
Male.....	818.7	1,427.6	49.5	23.8	29.7	124.6	216.9	255.3	269.0	334.7
Female.....	723.4	1,161.6	41.3	20.7	22.7	41.7	69.4	98.6	135.2	199.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Whooping cough (A37)

Table with 11 columns representing age groups and 20 rows representing different demographic categories (All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) with asterisks indicating data presence.

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
All origins 3/, both sexes.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5
Male.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5
Female.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4
Hispanic, both sexes.....	4.0	6.1	0.7	*	*	*	*	*	*	1.3
Male.....	3.8	7.8	*	*	*	*	*	*	*	1.3
Female.....	4.2	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	3.3	5.9	0.8	*	*	*	*	*	*	1.3
Male.....	2.9	7.8	*	*	*	*	*	*	*	*
Female.....	3.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	7.2	*	*	*	*	*	*	*	*	*
Male.....	7.4	*	*	*	*	*	*	*	*	*
Female.....	7.0	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	9.8	*	*	*	*	*	*	*	*	*
Male.....	9.7	*	*	*	*	*	*	*	*	*
Female.....	9.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.8	*	*	*	*	*	*	*	*	*
Male.....	3.7	*	*	*	*	*	*	*	*	*
Female.....	3.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	12.5	6.7	0.5	0.2	0.2	0.3	0.5	0.6	1.2	1.5
Male.....	11.5	7.9	0.4	*	*	0.3	0.5	0.6	1.2	1.6
Female.....	13.5	5.4	0.5	*	*	0.2	0.4	0.7	1.1	1.4
Non-Hispanic white, both sexes.....	12.5	4.8	0.4	0.2	*	0.2	0.3	0.5	0.9	1.3
Male.....	11.4	5.7	*	*	*	*	0.3	0.5	0.9	1.4
Female.....	13.6	3.9	0.5	*	*	*	*	0.5	0.9	1.2
Non-Hispanic black, both sexes.....	16.1	14.2	*	*	*	*	1.1	1.3	2.6	2.7
Male.....	15.1	15.8	*	*	*	*	*	*	3.0	2.6
Female.....	17.0	12.6	*	*	*	*	*	*	2.2	2.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Arthropod-borne viral encephalitis (A83-A84,A85.2)

Table with 11 columns representing age groups and 20 rows representing different demographic categories (e.g., All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, Non-Hispanic black). Each cell contains an asterisk (*) indicating data presence.

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
All origins 3/, both sexes.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5
Male.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3
Female.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6
Hispanic, both sexes.....	4.3	*	*	*	*	*	*	1.7	3.9	9.7
Male.....	6.4	*	*	*	*	*	*	2.4	5.9	14.4
Female.....	1.9	*	*	*	*	*	*	*	1.7	4.5
Mexican, both sexes.....	2.3	*	*	*	*	*	*	1.4	3.2	6.1
Male.....	3.9	*	*	*	*	*	*	2.1	5.3	10.4
Female.....	0.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	17.8	*	*	*	*	*	*	*	10.6	36.4
Male.....	23.9	*	*	*	*	*	*	*	*	42.0
Female.....	11.9	*	*	*	*	*	*	*	*	30.5
Cuban, both sexes.....	6.5	*	*	*	*	*	*	*	*	*
Male.....	11.2	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.0	*	*	*	*	*	*	*	4.3	10.5
Male.....	6.2	*	*	*	*	*	*	*	6.4	16.5
Female.....	1.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.4	*	*	*	*	0.1	0.8	2.3	5.3	9.4
Male.....	6.5	*	*	*	*	*	0.9	2.6	6.4	13.1
Female.....	2.4	*	*	*	*	*	0.8	2.0	4.1	5.7
Non-Hispanic white, both sexes.....	1.9	*	*	*	*	*	0.2	0.5	1.8	4.4
Male.....	3.3	*	*	*	*	*	*	0.7	2.9	6.9
Female.....	0.6	*	*	*	*	*	*	*	0.8	1.9
Non-Hispanic black, both sexes.....	19.4	*	*	*	*	0.8	3.9	11.4	24.3	38.7
Male.....	26.7	*	*	*	*	*	4.0	12.6	27.5	51.9
Female.....	12.8	*	*	*	*	*	3.8	10.3	21.4	27.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All origins 3/, both sexes.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5
Male.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1
Female.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9
Hispanic, both sexes.....	59.3	*	2.0	2.6	3.0	4.1	4.5	5.0	8.9	15.9
Male.....	60.9	*	1.6	2.9	3.1	5.5	5.5	4.8	8.2	13.0
Female.....	57.7	*	2.4	2.2	2.8	2.7	3.3	5.3	9.7	19.1
Mexican, both sexes.....	48.3	*	2.0	2.4	3.3	4.2	4.4	4.9	8.0	15.9
Male.....	49.3	*	1.7	2.9	3.5	5.7	5.6	4.5	7.0	12.7
Female.....	47.3	*	2.3	1.8	3.0	2.6	2.9	5.5	9.3	19.8
Puerto Rican, both sexes.....	79.0	*	*	*	*	*	*	*	12.4	13.8
Male.....	85.3	*	*	*	*	*	*	*	*	*
Female.....	72.8	*	*	*	*	*	*	*	*	20.1
Cuban, both sexes.....	167.8	*	*	*	*	*	*	*	*	*
Male.....	188.1	*	*	*	*	*	*	*	*	*
Female.....	146.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	65.2	*	*	*	*	4.6	4.5	4.8	10.0	16.8
Male.....	64.3	*	*	*	*	*	5.3	*	10.2	15.7
Female.....	66.0	*	*	*	*	*	*	*	9.8	18.0
Non-Hispanic 5/, both sexes.....	209.4	1.7	2.6	2.7	2.2	3.4	4.7	6.9	12.0	22.4
Male.....	221.8	1.8	2.8	3.0	2.3	3.8	5.5	7.3	10.8	19.0
Female.....	197.5	1.6	2.4	2.4	2.0	3.0	3.9	6.6	13.2	25.8
Non-Hispanic white, both sexes.....	227.0	1.9	2.7	2.8	2.1	3.4	4.4	6.8	11.6	21.7
Male.....	239.2	2.0	2.8	3.1	2.3	3.8	5.1	7.5	11.0	19.0
Female.....	215.3	1.8	2.5	2.4	1.8	2.9	3.7	6.1	12.3	24.4
Non-Hispanic black, both sexes.....	167.5	*	2.5	2.5	2.7	3.9	6.1	9.2	15.6	29.4
Male.....	182.4	*	2.8	2.5	2.6	4.3	7.1	8.1	12.2	20.6
Female.....	153.9	*	2.2	2.5	2.8	3.5	5.2	10.3	18.6	37.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All origins 3/, both sexes.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0
Male.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1
Female.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0
Hispanic, both sexes.....	5.9	*	*	*	*	*	*	*	0.7	1.6
Male.....	6.2	*	*	*	*	*	*	*	*	1.8
Female.....	5.6	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	4.5	*	*	*	*	*	*	*	*	1.4
Male.....	4.8	*	*	*	*	*	*	*	*	1.7
Female.....	4.1	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	8.4	*	*	*	*	*	*	*	*	*
Male.....	8.9	*	*	*	*	*	*	*	*	*
Female.....	8.0	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	19.3	*	*	*	*	*	*	*	*	*
Male.....	20.0	*	*	*	*	*	*	*	*	*
Female.....	18.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.7	*	*	*	*	*	*	*	*	*
Male.....	7.0	*	*	*	*	*	*	*	*	*
Female.....	6.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.3	*	*	*	*	*	0.2	0.5	1.0	2.1
Male.....	20.7	*	*	*	*	*	0.3	0.5	1.0	2.1
Female.....	19.9	*	*	*	*	*	*	0.4	0.9	2.1
Non-Hispanic white, both sexes.....	21.7	*	*	*	*	*	*	0.4	0.9	2.0
Male.....	22.2	*	*	*	*	*	*	0.5	1.0	2.1
Female.....	21.3	*	*	*	*	*	*	0.4	0.8	1.8
Non-Hispanic black, both sexes.....	17.8	*	*	*	*	*	*	*	1.4	2.9
Male.....	18.0	*	*	*	*	*	*	*	*	2.5
Female.....	17.6	*	*	*	*	*	*	*	1.4	3.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
All origins 3/, both sexes.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8
Hispanic, both sexes.....	4.7	*	*	*	*	*	*	*	1.0	2.8
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	9.6	*	*	*	*	*	*	*	2.1	5.9
Mexican, both sexes.....	3.6	*	*	*	*	*	*	*	1.0	2.8
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	7.5	*	*	*	*	*	*	*	2.1	6.2
Puerto Rican, both sexes.....	6.5	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	12.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	12.5	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	25.7	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.7	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	11.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.6	*	*	*	*	*	*	0.4	1.7	4.1
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	30.2	*	*	*	*	*	*	0.8	3.4	8.1
Non-Hispanic white, both sexes.....	16.4	*	*	*	*	*	*	0.3	1.4	3.5
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	32.0	*	*	*	*	*	*	0.6	2.9	7.0
Non-Hispanic black, both sexes.....	15.0	*	*	*	*	*	*	1.0	3.4	8.3
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	28.4	*	*	*	*	*	*	1.9	6.4	15.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
All origins 3/, both sexes.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1
Male.....
Female.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2
Hispanic, both sexes.....	1.0	*	*	*	*	*	*	*	*	1.2
Male.....
Female.....	2.2	*	*	*	*	*	*	*	*	2.5
Mexican, both sexes.....	1.0	*	*	*	*	*	*	*	*	1.4
Male.....
Female.....	2.2	*	*	*	*	*	*	*	*	3.0
Puerto Rican, both sexes.....	1.6	*	*	*	*	*	*	*	*	*
Male.....
Female.....	3.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.9	*	*	*	*	*	*	*	*	*
Male.....
Female.....	1.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.4	*	*	*	*	*	*	0.4	0.7	1.1
Male.....
Female.....	2.6	*	*	*	*	*	*	0.7	1.5	2.1
Non-Hispanic white, both sexes.....	1.2	*	*	*	*	*	*	0.3	0.7	1.0
Male.....
Female.....	2.4	*	*	*	*	*	*	0.6	1.4	2.0
Non-Hispanic black, both sexes.....	2.1	*	*	*	*	*	*	*	1.0	1.7
Male.....
Female.....	4.0	*	*	*	*	*	*	*	1.8	3.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All origins 3/, both sexes.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7
Male.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2
Female.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2
Hispanic, both sexes.....	6.9	*	0.8	1.2	1.6	2.0	2.0	1.7	2.2	2.5
Male.....	7.6	*	*	1.7	1.9	2.6	2.7	1.8	2.7	2.6
Female.....	6.2	*	*	*	1.4	1.3	1.2	1.5	1.7	2.4
Mexican, both sexes.....	6.0	*	*	1.3	1.9	2.2	2.1	1.7	2.1	2.5
Male.....	6.6	*	*	1.9	2.2	2.8	3.0	1.9	2.4	2.6
Female.....	5.4	*	*	*	1.6	*	*	*	*	2.4
Puerto Rican, both sexes.....	8.0	*	*	*	*	*	*	*	*	*
Male.....	8.7	*	*	*	*	*	*	*	*	*
Female.....	7.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	15.9	*	*	*	*	*	*	*	*	*
Male.....	17.7	*	*	*	*	*	*	*	*	*
Female.....	14.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.7	*	*	*	*	*	*	*	*	3.4
Male.....	8.5	*	*	*	*	*	*	*	*	*
Female.....	7.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.5	*	0.8	0.8	0.7	1.3	1.6	1.8	2.3	2.7
Male.....	22.6	*	0.9	1.0	0.7	1.4	2.0	2.2	2.7	3.3
Female.....	18.5	*	0.6	0.7	0.6	1.1	1.3	1.5	1.9	2.2
Non-Hispanic white, both sexes.....	22.7	*	0.8	0.9	0.6	1.3	1.6	1.7	2.2	2.6
Male.....	25.1	*	0.9	1.0	0.7	1.4	1.8	2.2	2.6	3.2
Female.....	20.4	*	0.6	0.7	0.5	1.1	1.3	1.2	1.8	2.0
Non-Hispanic black, both sexes.....	14.0	*	*	*	0.8	1.4	2.0	3.0	3.3	3.8
Male.....	14.8	*	*	*	*	1.4	2.5	3.1	3.8	4.5
Female.....	13.2	*	*	*	*	1.4	1.4	2.8	2.8	3.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes.....	21.1	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.1
Male.....	21.9	*	0.7	0.8	0.8	1.5	1.8	1.9	2.3	3.6
Female.....	20.3	*	0.9	0.5	0.6	1.0	0.9	1.1	1.7	2.5
Hispanic, both sexes.....	7.2	*	0.6	*	0.6	1.1	1.4	1.3	1.4	1.8
Male.....	7.1	*	*	*	*	1.8	1.9	1.5	1.6	2.0
Female.....	7.3	*	*	*	*	*	*	*	1.2	1.6
Mexican, both sexes.....	6.1	*	*	*	0.8	1.1	1.5	1.2	1.2	1.9
Male.....	6.0	*	*	*	*	*	2.0	1.3	1.5	2.0
Female.....	6.2	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	9.1	*	*	*	*	*	*	*	*	*
Male.....	9.8	*	*	*	*	*	*	*	*	*
Female.....	8.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	19.1	*	*	*	*	*	*	*	*	*
Male.....	19.6	*	*	*	*	*	*	*	*	*
Female.....	18.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.7	*	*	*	*	*	*	*	*	*
Male.....	7.4	*	*	*	*	*	*	*	*	*
Female.....	7.9	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	23.3	0.6	0.8	0.7	0.7	1.3	1.4	1.6	2.1	3.3
Male.....	24.4	*	0.8	0.8	0.8	1.5	1.8	2.0	2.4	3.9
Female.....	22.3	*	0.8	0.6	0.6	1.0	1.0	1.2	1.7	2.7
Non-Hispanic white, both sexes.....	25.4	*	0.8	0.7	0.7	1.2	1.3	1.7	2.1	3.2
Male.....	26.5	*	0.9	0.8	0.8	1.4	1.7	2.1	2.6	3.9
Female.....	24.4	*	0.8	0.6	0.5	1.1	0.9	1.2	1.6	2.6
Non-Hispanic black, both sexes.....	17.8	*	*	0.8	1.1	1.4	1.9	1.5	2.1	4.5
Male.....	19.0	*	*	*	*	1.8	2.2	1.8	2.0	4.9
Female.....	16.7	*	*	*	1.2	*	1.5	*	2.2	4.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
All origins 3/, both sexes.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.4	0.4	0.4
Male.....	1.3	*	*	*	*	0.3	0.4	0.4	0.5	0.5
Female.....	1.8	*	*	*	*	*	0.4	0.5	0.3	0.3
Hispanic, both sexes.....	0.5	*	*	*	*	*	*	*	*	*
Male.....	0.4	*	*	*	*	*	*	*	*	*
Female.....	0.6	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	0.7	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	1.6	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.5	*	*	*	*	*	*	*	*	*
Male.....	0.5	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.7	*	0.2	0.1	0.1	0.3	0.4	0.5	0.5	0.4
Male.....	1.4	*	*	*	*	0.3	0.4	0.5	0.6	0.6
Female.....	2.0	*	*	*	*	*	0.4	0.5	0.3	0.3
Non-Hispanic white, both sexes.....	1.6	*	*	*	*	*	*	*	*	*
Male.....	1.3	*	*	*	*	*	*	*	*	*
Female.....	2.0	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	2.7	*	*	*	*	0.9	2.0	2.3	2.5	2.5
Male.....	2.6	*	*	*	*	1.3	2.0	2.5	3.1	3.2
Female.....	2.8	*	*	*	*	*	2.1	2.1	2.0	1.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
All origins 3/, both sexes.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
Male.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2
Female.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5
Hispanic, both sexes.....	14.5	*	*	*	*	*	*	0.5	1.2	2.0
Male.....	13.4	*	*	*	*	*	*	*	1.3	2.2
Female.....	15.7	*	*	*	*	*	*	*	*	1.8
Mexican, both sexes.....	14.2	*	*	*	*	*	*	*	1.1	2.0
Male.....	13.0	*	*	*	*	*	*	*	*	2.2
Female.....	15.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	19.4	*	*	*	*	*	*	*	*	*
Male.....	18.0	*	*	*	*	*	*	*	*	*
Female.....	20.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	25.8	*	*	*	*	*	*	*	*	*
Male.....	24.5	*	*	*	*	*	*	*	*	*
Female.....	27.2	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	11.4	*	*	*	*	*	*	*	*	*
Male.....	10.4	*	*	*	*	*	*	*	*	*
Female.....	12.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	26.5	*	*	*	0.2	0.3	0.6	1.0	2.2	3.6
Male.....	26.2	*	*	*	*	0.3	0.6	1.2	2.5	4.6
Female.....	26.8	*	*	*	*	*	0.6	0.8	2.0	2.6
Non-Hispanic white, both sexes.....	26.1	*	*	*	*	0.2	0.5	0.8	1.9	3.0
Male.....	26.3	*	*	*	*	*	0.5	0.9	2.1	3.9
Female.....	25.9	*	*	*	*	*	0.5	0.7	1.6	2.2
Non-Hispanic black, both sexes.....	34.4	*	*	*	*	*	1.4	2.1	4.7	7.7
Male.....	31.3	*	*	*	*	*	*	2.6	5.5	9.9
Female.....	37.2	*	*	*	*	*	1.7	1.7	4.0	5.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
All origins 3/, both sexes.....	293.3	13.8	1.5	0.6	1.0	2.2	4.2	7.3	12.6	24.3
Male.....	281.2	14.7	1.5	0.5	1.1	2.8	5.3	9.2	16.3	32.1
Female.....	304.9	12.9	1.5	0.7	1.0	1.6	3.0	5.3	8.9	16.4
Hispanic, both sexes.....	89.3	11.5	1.2	0.5	0.7	1.7	3.2	4.1	7.2	14.2
Male.....	90.6	11.7	1.6	*	*	2.5	4.3	5.7	10.0	19.6
Female.....	87.9	11.3	*	*	*	*	1.9	2.1	3.8	8.0
Mexican, both sexes.....	72.1	10.4	1.4	*	*	2.0	3.2	4.4	6.9	14.2
Male.....	74.1	10.7	1.9	*	*	3.1	4.3	6.2	9.3	19.0
Female.....	69.8	10.0	*	*	*	*	1.9	1.9	4.0	8.4
Puerto Rican, both sexes.....	127.8	*	*	*	*	*	*	*	11.3	19.2
Male.....	132.7	*	*	*	*	*	*	*	*	25.6
Female.....	123.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	308.4	*	*	*	*	*	*	*	*	*
Male.....	299.5	*	*	*	*	*	*	*	*	*
Female.....	317.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	85.7	*	*	*	*	*	3.1	3.4	6.7	12.8
Male.....	86.6	*	*	*	*	*	*	*	10.2	19.0
Female.....	84.7	*	*	*	*	*	*	*	*	6.5
Non-Hispanic 5/, both sexes.....	325.9	14.4	1.5	0.6	1.1	2.3	4.4	8.0	13.8	26.1
Male.....	313.4	15.5	1.4	0.6	1.1	2.8	5.5	10.1	17.8	34.5
Female.....	337.9	13.2	1.6	0.7	1.0	1.7	3.2	6.0	9.9	17.8
Non-Hispanic white, both sexes.....	353.1	12.0	1.2	0.5	0.9	1.8	3.7	6.2	11.3	22.4
Male.....	336.9	13.0	1.1	0.5	1.0	2.3	4.7	8.1	14.8	30.0
Female.....	368.7	10.9	1.2	0.5	0.9	1.4	2.6	4.3	7.7	14.8
Non-Hispanic black, both sexes.....	269.2	22.4	3.0	1.2	1.9	4.3	7.9	17.6	29.9	51.7
Male.....	266.0	24.8	2.9	*	1.9	5.7	10.1	21.2	36.7	68.0
Female.....	272.2	20.0	3.0	1.4	1.9	2.8	5.8	14.3	23.7	37.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All origins 3/, both sexes.....	222.2	10.3	1.2	0.4	0.8	1.8	3.2	5.7	10.0	19.4
Male.....	222.8	10.9	1.1	0.3	0.8	2.3	4.2	7.4	13.6	26.5
Female.....	221.6	9.7	1.2	0.5	0.7	1.2	2.2	3.9	6.4	12.3
Hispanic, both sexes.....	67.2	8.9	1.1	*	*	1.3	2.3	2.8	5.2	9.9
Male.....	70.2	8.9	1.4	*	*	1.9	3.3	4.1	7.7	14.1
Female.....	64.1	9.0	*	*	*	*	*	1.2	2.2	5.2
Mexican, both sexes.....	52.8	7.7	1.2	*	*	1.5	2.3	2.9	4.9	10.1
Male.....	56.3	7.5	1.6	*	*	2.3	3.3	4.3	6.9	14.0
Female.....	48.9	7.9	*	*	*	*	*	*	2.4	5.3
Puerto Rican, both sexes.....	100.6	*	*	*	*	*	*	*	8.5	13.2
Male.....	105.9	*	*	*	*	*	*	*	*	19.0
Female.....	95.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	253.7	*	*	*	*	*	*	*	*	*
Male.....	247.2	*	*	*	*	*	*	*	*	*
Female.....	260.7	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	63.2	*	*	*	*	*	*	2.6	5.1	8.1
Male.....	66.3	*	*	*	*	*	*	*	8.3	11.4
Female.....	60.1	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	247.0	10.7	1.2	0.4	0.8	1.8	3.4	6.4	11.1	21.1
Male.....	248.5	11.5	1.1	0.3	0.9	2.3	4.4	8.2	15.0	28.8
Female.....	245.5	9.8	1.3	0.5	0.8	1.4	2.4	4.5	7.2	13.4
Non-Hispanic white, both sexes.....	269.1	9.1	0.9	0.3	0.7	1.5	2.8	4.9	9.2	18.6
Male.....	269.1	9.8	0.8	*	0.8	1.8	3.7	6.5	12.7	25.8
Female.....	269.1	8.3	1.0	0.4	0.6	1.1	1.9	3.3	5.7	11.3
Non-Hispanic black, both sexes.....	198.7	15.6	2.5	0.9	1.5	3.7	6.5	13.9	23.1	39.6
Male.....	203.2	17.0	2.5	*	1.5	4.8	8.3	17.4	29.6	52.9
Female.....	194.6	14.2	2.4	*	1.6	2.5	4.7	10.7	17.3	27.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All origins 3/, both sexes.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9
Male.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8
Female.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0
Hispanic, both sexes.....	3.6	*	*	*	*	*	*	*	*	1.2
Male.....	3.8	*	*	*	*	*	*	*	*	1.9
Female.....	3.4	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	2.8	*	*	*	*	*	*	*	*	1.1
Male.....	3.1	*	*	*	*	*	*	*	*	1.7
Female.....	2.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	5.4	*	*	*	*	*	*	*	*	*
Male.....	5.7	*	*	*	*	*	*	*	*	*
Female.....	5.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	12.1	*	*	*	*	*	*	*	*	*
Male.....	11.8	*	*	*	*	*	*	*	*	*
Female.....	12.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.8	*	*	*	*	*	*	*	*	*
Male.....	3.7	*	*	*	*	*	*	*	*	*
Female.....	3.9	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.7	*	*	*	*	*	0.1	0.6	1.1	2.0
Male.....	9.8	*	*	*	*	*	*	0.8	1.7	3.0
Female.....	11.6	*	*	*	*	*	*	0.3	0.6	1.1
Non-Hispanic white, both sexes.....	9.8	*	*	*	*	*	*	0.3	0.7	1.3
Male.....	8.6	*	*	*	*	*	*	0.5	1.1	2.0
Female.....	11.0	*	*	*	*	*	*	*	*	0.7
Non-Hispanic black, both sexes.....	18.2	*	*	*	*	*	*	2.2	3.4	6.2
Male.....	19.0	*	*	*	*	*	*	2.6	4.7	9.0
Female.....	17.5	*	*	*	*	*	*	1.9	2.2	3.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All origins 3/, both sexes.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4
Male.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9
Female.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9
Hispanic, both sexes.....	49.6	*	*	*	*	*	*	0.5	2.0	4.2
Male.....	52.7	*	*	*	*	*	*	*	3.3	6.3
Female.....	46.2	*	*	*	*	*	*	*	*	1.8
Mexican, both sexes.....	38.5	*	*	*	*	*	*	*	1.8	4.0
Male.....	42.0	*	*	*	*	*	*	*	2.9	5.8
Female.....	34.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	76.3	*	*	*	*	*	*	*	*	*
Male.....	80.8	*	*	*	*	*	*	*	*	*
Female.....	71.8	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	196.1	*	*	*	*	*	*	*	*	*
Male.....	190.2	*	*	*	*	*	*	*	*	*
Female.....	202.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	45.4	*	*	*	*	*	*	*	*	3.5
Male.....	49.3	*	*	*	*	*	*	*	*	*
Female.....	41.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	170.3	*	*	*	*	0.1	0.5	1.7	4.3	10.3
Male.....	179.9	*	*	*	*	*	0.7	2.4	6.6	15.3
Female.....	161.1	*	*	*	*	*	0.3	1.0	2.1	5.4
Non-Hispanic white, both sexes.....	187.7	*	*	*	*	*	0.3	1.5	4.0	10.3
Male.....	198.0	*	*	*	*	*	0.5	2.2	6.2	15.2
Female.....	177.7	*	*	*	*	*	*	0.8	1.9	5.3
Non-Hispanic black, both sexes.....	124.8	*	*	*	*	*	1.3	3.0	6.8	13.1
Male.....	129.2	*	*	*	*	*	1.9	3.5	10.0	19.0
Female.....	120.8	*	*	*	*	*	*	2.5	3.9	7.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All origins 3/, both sexes.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8
Male.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7
Female.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9
Hispanic, both sexes.....	17.3	*	*	*	*	*	*	*	0.8	1.6
Male.....	18.4	*	*	*	*	*	*	*	1.1	2.2
Female.....	16.0	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	14.6	*	*	*	*	*	*	*	0.8	1.7
Male.....	16.2	*	*	*	*	*	*	*	*	2.3
Female.....	12.8	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	22.2	*	*	*	*	*	*	*	*	*
Male.....	23.2	*	*	*	*	*	*	*	*	*
Female.....	21.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	69.8	*	*	*	*	*	*	*	*	*
Male.....	63.7	*	*	*	*	*	*	*	*	*
Female.....	76.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.7	*	*	*	*	*	*	*	*	*
Male.....	14.8	*	*	*	*	*	*	*	*	*
Female.....	12.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	59.2	*	*	*	*	*	0.2	0.7	1.8	4.2
Male.....	64.0	*	*	*	*	*	0.4	0.9	2.6	6.3
Female.....	54.7	*	*	*	*	*	*	0.5	0.9	2.1
Non-Hispanic white, both sexes.....	65.3	*	*	*	*	*	0.2	0.6	1.8	4.2
Male.....	70.9	*	*	*	*	*	*	0.9	2.8	6.3
Female.....	59.9	*	*	*	*	*	*	0.4	0.8	2.1
Non-Hispanic black, both sexes.....	43.4	*	*	*	*	*	*	1.2	2.4	5.1
Male.....	43.9	*	*	*	*	*	*	*	2.9	8.0
Female.....	43.0	*	*	*	*	*	*	*	1.9	2.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
All origins 3/, both sexes.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5
Male.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1
Female.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0
Hispanic, both sexes.....	32.2	*	*	*	*	*	*	*	1.2	2.7
Male.....	34.2	*	*	*	*	*	*	*	2.2	4.1
Female.....	30.1	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	23.8	*	*	*	*	*	*	*	1.0	2.3
Male.....	25.6	*	*	*	*	*	*	*	1.7	3.4
Female.....	21.8	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	53.9	*	*	*	*	*	*	*	*	*
Male.....	57.3	*	*	*	*	*	*	*	*	*
Female.....	50.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	126.2	*	*	*	*	*	*	*	*	*
Male.....	126.4	*	*	*	*	*	*	*	*	*
Female.....	125.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	31.6	*	*	*	*	*	*	*	*	2.9
Male.....	34.5	*	*	*	*	*	*	*	*	*
Female.....	28.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	110.1	*	*	*	*	*	0.3	1.0	2.5	6.0
Male.....	114.9	*	*	*	*	*	0.3	1.5	3.9	8.8
Female.....	105.6	*	*	*	*	*	*	0.5	1.1	3.3
Non-Hispanic white, both sexes.....	121.4	*	*	*	*	*	0.2	0.9	2.2	5.9
Male.....	126.1	*	*	*	*	*	*	1.4	3.4	8.7
Female.....	116.9	*	*	*	*	*	*	0.4	1.1	3.1
Non-Hispanic black, both sexes.....	80.3	*	*	*	*	*	*	1.7	4.3	7.9
Male.....	83.9	*	*	*	*	*	*	2.1	7.0	10.8
Female.....	76.9	*	*	*	*	*	*	*	1.8	5.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
All origins 3/, both sexes.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8
Male.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4
Female.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1
Hispanic, both sexes.....	13.2	8.5	1.0	*	*	1.2	1.8	2.0	2.6	4.2
Male.....	13.0	8.4	1.3	*	*	1.7	2.6	3.1	3.6	5.6
Female.....	13.5	8.6	*	*	*	*	*	*	1.5	2.7
Mexican, both sexes.....	10.8	7.3	1.1	*	*	1.3	1.7	2.1	2.5	4.6
Male.....	10.7	7.0	*	*	*	2.0	2.5	3.3	3.2	6.0
Female.....	10.9	7.6	*	*	*	*	*	*	1.7	2.8
Puerto Rican, both sexes.....	17.9	*	*	*	*	*	*	*	*	*
Male.....	18.5	*	*	*	*	*	*	*	*	*
Female.....	17.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	43.7	*	*	*	*	*	*	*	*	*
Male.....	42.9	*	*	*	*	*	*	*	*	*
Female.....	44.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.1	*	*	*	*	*	*	*	*	3.1
Male.....	12.4	*	*	*	*	*	*	*	*	*
Female.....	13.7	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	63.6	10.3	1.1	0.4	0.8	1.6	2.7	3.9	5.5	8.4
Male.....	57.0	11.0	1.0	0.3	0.8	2.0	3.4	4.8	6.6	10.1
Female.....	69.9	9.6	1.2	0.5	0.7	1.3	2.0	3.0	4.4	6.7
Non-Hispanic white, both sexes.....	69.3	8.7	0.8	0.3	0.6	1.3	2.3	3.1	4.4	6.9
Male.....	60.9	9.3	0.7	*	0.7	1.5	3.0	3.7	5.4	8.4
Female.....	77.3	8.2	0.9	*	0.5	1.0	1.6	2.4	3.4	5.3
Non-Hispanic black, both sexes.....	52.7	15.3	2.4	0.8	1.4	3.1	4.6	8.4	12.4	18.8
Male.....	52.3	16.4	2.5	*	1.3	4.0	5.7	10.8	14.4	22.6
Female.....	53.1	14.2	2.3	*	1.4	2.1	3.5	6.1	10.6	15.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/ 1 year	Under 1 year 2/ 2 years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All origins 3/, both sexes.....	36.5	8.7	0.9	0.3	0.6	1.4	2.3	3.2	4.5	6.9
Male.....	34.5	9.2	0.9	0.2	0.7	1.8	3.0	4.1	5.5	8.3
Female.....	38.5	8.2	0.8	0.4	0.6	1.0	1.6	2.4	3.5	5.5
Hispanic, both sexes.....	9.1	7.2	0.7	*	*	1.0	1.7	1.8	2.5	3.5
Male.....	9.4	6.9	*	*	*	1.5	2.5	2.8	3.4	4.6
Female.....	8.8	7.4	*	*	*	*	*	*	1.3	2.3
Mexican, both sexes.....	7.1	5.8	0.8	*	*	1.2	1.6	1.9	2.4	3.8
Male.....	7.5	*	*	*	*	1.7	2.5	3.0	3.1	5.0
Female.....	6.7	*	*	*	*	*	*	*	*	2.4
Puerto Rican, both sexes.....	12.7	*	*	*	*	*	*	*	*	*
Male.....	13.8	*	*	*	*	*	*	*	*	*
Female.....	11.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	33.9	*	*	*	*	*	*	*	*	*
Male.....	33.8	*	*	*	*	*	*	*	*	*
Female.....	34.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	9.0	*	*	*	*	*	*	*	*	2.9
Male.....	9.1	*	*	*	*	*	*	*	*	*
Female.....	9.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	40.9	9.1	0.9	0.3	0.7	1.5	2.4	3.6	4.9	7.6
Male.....	38.8	9.7	0.9	*	0.7	1.8	3.1	4.4	6.0	9.0
Female.....	43.0	8.3	0.9	0.4	0.6	1.1	1.8	2.8	3.9	6.1
Non-Hispanic white, both sexes.....	43.7	7.9	0.7	0.3	0.6	1.2	2.1	2.8	4.0	6.2
Male.....	40.6	8.4	0.7	*	0.6	1.4	2.7	3.4	4.9	7.5
Female.....	46.7	7.3	0.7	*	0.5	1.0	1.5	2.2	3.0	5.0
Non-Hispanic black, both sexes.....	37.7	12.8	1.9	*	1.2	2.8	4.2	7.6	11.1	16.4
Male.....	39.0	13.9	1.9	*	*	3.8	5.3	9.8	12.7	20.2
Female.....	36.5	11.6	1.8	*	1.3	1.7	3.2	5.6	9.8	13.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
All origins 3/, both sexes.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
Male.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7
Female.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1
Hispanic, both sexes.....	16.4	2.2	*	*	*	*	0.7	0.9	1.5	3.2
Male.....	15.0	*	*	*	*	*	*	1.2	1.7	4.3
Female.....	17.9	*	*	*	*	*	*	*	1.3	2.0
Mexican, both sexes.....	14.7	*	*	*	*	*	0.8	1.1	1.6	3.2
Male.....	13.5	*	*	*	*	*	*	1.6	1.8	4.0
Female.....	15.9	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	19.4	*	*	*	*	*	*	*	*	*
Male.....	18.9	*	*	*	*	*	*	*	*	*
Female.....	19.9	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	38.8	*	*	*	*	*	*	*	*	*
Male.....	35.9	*	*	*	*	*	*	*	*	*
Female.....	42.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	16.4	*	*	*	*	*	*	*	*	3.3
Male.....	14.2	*	*	*	*	*	*	*	*	*
Female.....	18.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	56.7	3.3	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
Male.....	45.0	3.7	0.3	*	*	0.4	0.7	1.1	1.7	3.6
Female.....	67.8	3.0	*	*	*	0.3	0.7	1.1	2.0	3.3
Non-Hispanic white, both sexes.....	60.6	2.6	0.3	0.2	0.2	0.3	0.6	0.9	1.4	2.8
Male.....	47.0	2.9	*	*	*	0.3	0.6	1.0	1.2	2.7
Female.....	73.7	2.2	*	*	*	*	0.6	0.8	1.6	2.9
Non-Hispanic black, both sexes.....	48.7	6.2	*	*	*	*	1.0	2.3	4.3	7.3
Male.....	43.1	7.1	*	*	*	*	*	1.8	4.3	8.9
Female.....	53.8	*	*	*	*	*	*	2.7	4.3	5.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza and pneumonia (J10-J18)										
All origins 3/, both sexes.....	20.3	6.7	0.7	0.2	0.2	0.3	0.6	0.5	1.0	1.5
Male.....	18.6	8.0	0.6	*	0.3	0.4	0.7	0.7	1.2	1.7
Female.....	22.0	5.4	0.9	*	0.2	0.2	0.4	0.4	0.7	1.2
Hispanic, both sexes.....	7.0	5.8	0.7	*	*	*	0.6	*	0.8	1.1
Male.....	6.6	8.0	*	*	*	*	*	*	1.1	1.1
Female.....	7.6	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.7	5.9	*	*	*	*	*	*	0.8	1.1
Male.....	5.5	8.1	*	*	*	*	*	*	*	*
Female.....	6.0	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	12.0	*	*	*	*	*	*	*	*	*
Male.....	11.6	*	*	*	*	*	*	*	*	*
Female.....	12.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	20.5	*	*	*	*	*	*	*	*	*
Male.....	17.8	*	*	*	*	*	*	*	*	*
Female.....	23.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.6	*	*	*	*	*	*	*	*	*
Male.....	5.9	*	*	*	*	*	*	*	*	*
Female.....	7.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	22.4	6.9	0.8	0.2	0.2	0.3	0.6	0.6	1.0	1.5
Male.....	20.6	8.0	0.6	*	0.3	0.4	0.7	0.8	1.2	1.8
Female.....	24.2	5.9	0.9	*	*	0.2	0.4	0.3	0.8	1.3
Non-Hispanic white, both sexes.....	24.8	4.8	0.6	*	0.2	0.3	0.5	0.5	0.9	1.3
Male.....	22.4	5.5	0.5	*	*	0.4	0.7	0.7	1.1	1.5
Female.....	27.1	4.2	0.7	*	*	*	0.3	0.3	0.8	1.1
Non-Hispanic black, both sexes.....	15.2	14.7	1.4	*	*	*	0.9	0.8	1.6	3.0
Male.....	15.5	17.0	*	*	*	*	*	*	2.0	3.8
Female.....	15.0	12.2	1.8	*	*	*	*	*	*	2.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All origins 3/, both sexes.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4
Male.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7
Female.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2
Hispanic, both sexes.....	7.0	5.2	0.6	*	*	*	0.6	*	0.8	1.1
Male.....	6.5	7.3	*	*	*	*	*	*	1.1	1.1
Female.....	7.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.6	5.0	*	*	*	*	*	*	0.8	1.1
Male.....	5.4	7.2	*	*	*	*	*	*	*	*
Female.....	5.9	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	12.0	*	*	*	*	*	*	*	*	*
Male.....	11.6	*	*	*	*	*	*	*	*	*
Female.....	12.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	20.3	*	*	*	*	*	*	*	*	*
Male.....	17.7	*	*	*	*	*	*	*	*	*
Female.....	23.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.6	*	*	*	*	*	*	*	*	*
Male.....	5.9	*	*	*	*	*	*	*	*	*
Female.....	7.2	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	22.0	6.6	0.6	0.2	0.2	0.3	0.5	0.6	1.0	1.5
Male.....	20.3	7.5	0.5	*	*	0.4	0.7	0.8	1.2	1.8
Female.....	23.7	5.6	0.8	*	*	*	0.4	0.3	0.7	1.2
Non-Hispanic white, both sexes.....	24.3	4.6	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2
Male.....	22.0	5.3	*	*	*	0.4	0.7	0.7	1.1	1.5
Female.....	26.5	3.8	0.6	*	*	*	*	0.3	0.7	1.0
Non-Hispanic black, both sexes.....	15.1	14.0	1.2	*	*	*	0.9	0.8	1.5	3.0
Male.....	15.4	15.8	*	*	*	*	*	*	1.9	3.8
Female.....	14.8	12.2	1.7	*	*	*	*	*	*	2.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
Male.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2
Female.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4
Hispanic, both sexes.....	7.5	*	*	*	*	*	*	*	*	*
Male.....	7.6	*	*	*	*	*	*	*	*	*
Female.....	7.4	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.2	*	*	*	*	*	*	*	*	*
Male.....	5.6	*	*	*	*	*	*	*	*	*
Female.....	4.9	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	12.3	*	*	*	*	*	*	*	*	*
Male.....	11.9	*	*	*	*	*	*	*	*	*
Female.....	12.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	32.9	*	*	*	*	*	*	*	*	*
Male.....	30.8	*	*	*	*	*	*	*	*	*
Female.....	35.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.9	*	*	*	*	*	*	*	*	*
Male.....	8.2	*	*	*	*	*	*	*	*	*
Female.....	7.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	47.0	0.8	0.3	0.3	0.4	0.5	0.5	0.6	0.8	1.5
Male.....	46.2	*	0.4	0.3	0.5	0.6	0.5	0.7	0.9	1.4
Female.....	47.8	*	*	*	0.3	0.3	0.4	0.6	0.6	1.6
Non-Hispanic white, both sexes.....	54.9	*	*	*	0.2	0.3	0.3	0.4	0.6	1.3
Male.....	53.0	*	*	*	*	0.3	0.4	0.3	0.7	1.1
Female.....	56.7	*	*	*	*	*	*	0.5	0.5	1.4
Non-Hispanic black, both sexes.....	19.9	*	0.8	0.9	1.1	1.3	1.2	1.8	2.1	3.1
Male.....	22.8	*	*	1.4	1.3	1.7	1.4	2.5	2.7	3.3
Female.....	17.2	*	*	*	*	*	*	*	1.4	2.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes.....	14.5	4.3	*	*	*	0.1	0.3	0.5	0.7	1.4
Male.....	14.1	5.3	*	*	*	*	0.3	0.5	0.6	1.5
Female.....	14.8	3.2	*	*	*	*	0.2	0.5	0.8	1.3
Hispanic, both sexes.....	5.3	4.1	*	*	*	*	*	*	*	0.8
Male.....	5.2	5.4	*	*	*	*	*	*	*	*
Female.....	5.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	4.8	3.9	*	*	*	*	*	*	*	*
Male.....	4.5	*	*	*	*	*	*	*	*	*
Female.....	5.2	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	8.1	*	*	*	*	*	*	*	*	*
Male.....	8.2	*	*	*	*	*	*	*	*	*
Female.....	8.0	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	11.0	*	*	*	*	*	*	*	*	*
Male.....	12.6	*	*	*	*	*	*	*	*	*
Female.....	9.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.6	*	*	*	*	*	*	*	*	*
Male.....	4.7	*	*	*	*	*	*	*	*	*
Female.....	4.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.9	4.3	*	*	*	0.1	0.3	0.5	0.8	1.5
Male.....	15.6	5.2	*	*	*	*	0.3	0.5	0.8	1.6
Female.....	16.2	3.4	*	*	*	*	0.2	0.5	0.9	1.4
Non-Hispanic white, both sexes.....	15.8	2.6	*	*	*	*	0.2	0.4	0.5	1.0
Male.....	15.6	3.1	*	*	*	*	*	0.4	0.4	1.0
Female.....	16.0	2.1	*	*	*	*	*	*	0.7	1.0
Non-Hispanic black, both sexes.....	21.0	10.4	*	*	*	*	0.8	1.4	2.5	4.3
Male.....	19.9	12.4	*	*	*	*	*	*	3.0	5.4
Female.....	22.0	8.4	*	*	*	*	*	1.6	2.2	3.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
All origins 3/, both sexes.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3
Male.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4
Female.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2
Hispanic, both sexes.....	5.2	3.9	*	*	*	*	*	*	*	0.8
Male.....	5.0	5.2	*	*	*	*	*	*	*	*
Female.....	5.4	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	4.7	3.6	*	*	*	*	*	*	*	*
Male.....	4.3	*	*	*	*	*	*	*	*	*
Female.....	5.1	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	7.9	*	*	*	*	*	*	*	*	*
Male.....	8.0	*	*	*	*	*	*	*	*	*
Female.....	7.9	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	10.8	*	*	*	*	*	*	*	*	*
Male.....	12.3	*	*	*	*	*	*	*	*	*
Female.....	9.2	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.5	*	*	*	*	*	*	*	*	*
Male.....	4.6	*	*	*	*	*	*	*	*	*
Female.....	4.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.7	4.1	*	*	*	0.1	0.3	0.5	0.8	1.4
Male.....	15.3	5.1	*	*	*	*	0.3	0.5	0.7	1.6
Female.....	16.0	3.1	*	*	*	*	*	0.4	0.8	1.3
Non-Hispanic white, both sexes.....	15.5	2.5	*	*	*	*	0.2	0.3	0.4	1.0
Male.....	15.3	3.0	*	*	*	*	*	0.4	0.3	1.0
Female.....	15.7	2.0	*	*	*	*	*	*	0.6	1.0
Non-Hispanic black, both sexes.....	20.7	10.1	*	*	*	*	0.7	1.2	2.4	4.1
Male.....	19.6	12.1	*	*	*	*	*	*	2.9	5.3
Female.....	21.7	8.0	*	*	*	*	*	*	2.0	3.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy, childbirth and the puerperium (000-099)										
All origins 3/, both sexes.....	0.2	*	0.2	0.6	0.7	0.7	0.5
Male.....
Female.....	0.5	*	0.4	1.3	1.5	1.4	0.9
Hispanic, both sexes.....	0.3	*	*	*	0.7	0.6	*
Male.....
Female.....	0.6	*	*	*	1.5	1.3	*
Mexican, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.6	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.7	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	*	0.1	0.7	0.7	0.7	0.4
Male.....
Female.....	0.4	*	0.3	1.3	1.5	1.5	0.9
Non-Hispanic white, both sexes.....	0.2	*	*	0.4	0.4	0.6	0.3
Male.....
Female.....	0.3	*	*	0.7	0.9	1.1	0.7
Non-Hispanic black, both sexes.....	0.7	*	*	2.0	2.3	1.8	1.1
Male.....
Female.....	1.3	*	*	4.1	4.4	3.4	2.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
All origins 3/, both sexes.....	0.0	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*
Hispanic, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.0	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
All origins 3/, both sexes.....	0.2	*	0.2	0.6	0.7	0.7	0.4
Male.....
Female.....	0.4	*	0.4	1.2	1.4	1.4	0.9
Hispanic, both sexes.....	0.3	*	*	*	0.7	0.6	*
Male.....
Female.....	0.6	*	*	*	1.5	1.3	*
Mexican, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.6	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.7	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	*	0.1	0.6	0.7	0.7	0.4
Male.....
Female.....	0.4	*	0.3	1.2	1.4	1.4	0.8
Non-Hispanic white, both sexes.....	0.1	*	*	0.3	0.4	0.5	0.3
Male.....
Female.....	0.3	*	*	0.7	0.9	1.1	0.7
Non-Hispanic black, both sexes.....	0.6	*	*	1.8	2.1	1.7	0.9
Male.....
Female.....	1.2	*	*	3.6	4.1	3.2	1.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All origins 3/, both sexes.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0
Male.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2
Female.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8
Hispanic, both sexes.....	4.3	141.0	4.0	0.9	0.9	0.9	0.8	0.9	*	*
Male.....	4.2	134.3	4.6	1.2	1.3	*	1.1	*	*	*
Female.....	4.3	147.9	3.3	*	*	*	*	*	*	*
Mexican, both sexes.....	4.7	144.7	3.9	1.0	0.9	*	0.8	0.8	*	*
Male.....	4.6	139.5	4.5	*	*	*	*	*	*	*
Female.....	4.9	150.2	3.2	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	2.9	*	*	*	*	*	*	*	*	*
Male.....	3.4	*	*	*	*	*	*	*	*	*
Female.....	2.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.0	158.8	5.0	*	*	*	*	*	*	*
Male.....	4.0	144.9	*	*	*	*	*	*	*	*
Female.....	4.1	172.8	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.4	135.9	3.4	1.1	0.9	1.3	1.1	1.1	1.1	1.1
Male.....	3.6	139.1	3.7	1.0	0.8	1.4	1.3	1.3	1.3	1.3
Female.....	3.2	132.6	3.2	1.1	0.9	1.2	0.9	0.9	1.0	0.8
Non-Hispanic white, both sexes.....	3.3	130.5	2.9	0.9	0.8	1.2	1.0	1.2	1.2	1.1
Male.....	3.4	132.8	3.1	1.0	0.8	1.3	1.1	1.5	1.3	1.2
Female.....	3.2	128.0	2.8	0.9	0.9	1.1	1.0	0.9	1.2	0.9
Non-Hispanic black, both sexes.....	4.6	164.3	5.1	1.5	1.1	1.9	1.7	1.2	1.1	1.3
Male.....	5.2	170.5	5.9	*	*	2.1	2.7	*	1.6	1.8
Female.....	4.1	157.8	4.3	1.8	*	1.7	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6
Male.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6
Female.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7
Hispanic, both sexes.....	3.9	50.4	1.1	*	*	1.2	2.0	2.0	2.2	2.8
Male.....	4.5	60.0	1.2	*	*	1.5	3.0	2.4	3.2	3.8
Female.....	3.3	40.4	*	*	*	*	*	1.5	*	1.5
Mexican, both sexes.....	3.6	47.4	1.0	*	*	1.0	2.0	1.9	1.9	2.4
Male.....	4.2	57.4	*	*	*	*	3.0	2.4	2.6	3.4
Female.....	2.9	36.9	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	6.2	*	*	*	*	*	*	*	*	*
Male.....	8.3	*	*	*	*	*	*	*	*	*
Female.....	4.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	5.5	*	*	*	*	*	*	*	*	*
Male.....	4.8	*	*	*	*	*	*	*	*	*
Female.....	6.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.8	60.8	*	*	*	*	*	*	*	*
Male.....	4.1	*	*	*	*	*	*	*	*	*
Female.....	3.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.2	90.9	1.2	0.3	0.3	1.0	2.2	2.8	2.9	3.8
Male.....	10.2	106.6	1.2	0.3	0.3	1.2	2.9	3.9	3.6	4.8
Female.....	12.1	74.3	1.2	*	0.4	0.7	1.4	1.7	2.1	2.8
Non-Hispanic white, both sexes.....	11.7	71.6	1.0	0.2	0.3	0.9	2.2	2.8	2.8	3.7
Male.....	10.3	83.1	1.0	*	*	1.2	2.9	4.0	3.5	4.7
Female.....	13.0	59.4	1.0	*	0.3	0.7	1.3	1.5	2.0	2.7
Non-Hispanic black, both sexes.....	11.9	174.5	2.0	*	0.7	1.3	2.4	4.0	4.3	5.5
Male.....	12.8	208.3	2.2	*	*	1.6	3.1	4.9	5.6	6.6
Female.....	11.0	139.4	1.7	*	*	*	1.6	3.0	3.1	4.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
All origins 3/, both sexes.....	68.8	29.2	3.8	1.9	2.4	3.8	5.6	7.1	9.0	14.3
Male.....	57.2	33.8	4.0	2.0	2.6	4.4	6.4	8.2	10.1	16.6
Female.....	79.9	24.5	3.6	1.8	2.2	3.1	4.8	6.0	7.8	12.0
Hispanic, both sexes.....	22.3	26.1	3.3	1.5	2.3	3.2	4.1	4.4	6.5	9.8
Male.....	21.3	28.7	3.5	1.6	2.6	3.9	4.9	5.0	7.6	13.4
Female.....	23.4	23.5	3.1	1.5	2.1	2.4	3.1	3.7	5.2	5.7
Mexican, both sexes.....	19.4	24.6	3.7	1.6	2.3	3.1	4.1	4.3	6.8	9.0
Male.....	19.0	25.5	3.8	1.6	2.9	4.0	4.7	5.2	8.0	12.0
Female.....	19.9	23.6	3.5	1.6	1.6	2.3	3.4	3.0	5.5	5.3
Puerto Rican, both sexes.....	29.8	*	*	*	*	*	*	8.2	8.5	15.2
Male.....	30.3	*	*	*	*	*	*	*	*	18.4
Female.....	29.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	54.0	*	*	*	*	*	*	*	*	*
Male.....	40.1	*	*	*	*	*	*	*	*	*
Female.....	68.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	22.2	30.4	*	*	*	3.3	4.3	3.6	5.6	9.9
Male.....	21.4	*	*	*	*	*	5.8	*	6.4	14.9
Female.....	23.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	76.2	30.0	3.9	2.0	2.4	3.9	5.9	7.8	9.5	15.1
Male.....	63.3	35.0	4.1	2.2	2.6	4.5	6.7	9.0	10.6	17.1
Female.....	88.5	24.7	3.7	1.9	2.2	3.2	5.1	6.5	8.4	13.0
Non-Hispanic white, both sexes.....	82.7	24.8	3.5	1.9	2.3	3.6	5.5	6.9	8.1	13.5
Male.....	67.4	29.6	3.7	2.1	2.5	4.3	6.5	8.7	9.2	15.9
Female.....	97.5	19.8	3.3	1.6	2.1	2.9	4.4	5.2	7.0	11.0
Non-Hispanic black, both sexes.....	63.1	50.5	5.4	2.7	3.0	5.2	8.8	13.5	18.4	27.0
Male.....	58.1	56.9	5.8	2.2	3.1	6.0	8.3	13.0	20.6	27.3
Female.....	67.6	43.8	5.0	3.2	2.9	4.4	9.4	13.8	16.5	26.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All origins 3/, both sexes.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8
Male.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7
Female.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7
Hispanic, both sexes.....	25.2	17.5	9.0	4.6	5.6	28.7	37.6	29.7	24.9	28.7
Male.....	36.5	18.8	10.5	5.1	7.2	43.0	57.2	45.1	37.8	43.8
Female.....	13.1	16.2	7.5	4.0	4.0	13.4	13.6	10.6	9.8	11.6
Mexican, both sexes.....	24.8	18.4	10.0	4.7	6.1	30.6	38.2	29.5	24.7	27.7
Male.....	36.2	19.7	11.6	4.9	7.9	45.8	58.6	44.6	37.6	42.0
Female.....	12.2	17.0	8.4	4.6	4.3	14.1	12.4	10.0	9.4	10.5
Puerto Rican, both sexes.....	23.5	*	*	*	*	15.8	27.5	26.7	25.1	40.5
Male.....	35.1	*	*	*	*	23.5	40.7	44.9	41.3	61.7
Female.....	12.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	28.3	*	*	*	*	*	*	*	*	*
Male.....	36.8	*	*	*	*	*	*	*	*	*
Female.....	19.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	26.4	*	7.5	4.6	4.6	28.7	39.3	32.3	26.8	27.8
Male.....	38.0	*	8.7	*	*	43.7	58.1	48.1	39.8	43.9
Female.....	14.8	*	*	*	*	14.4	17.2	13.1	11.9	11.7
Non-Hispanic 5/, both sexes.....	40.1	28.1	10.6	6.0	7.6	33.7	41.8	35.7	31.7	35.8
Male.....	51.9	31.8	12.1	7.2	9.4	44.8	62.8	54.0	46.4	50.7
Female.....	28.8	24.2	9.0	4.8	5.7	21.9	20.2	17.4	17.1	21.1
Non-Hispanic white, both sexes.....	42.8	24.1	10.2	5.3	7.4	37.3	45.7	38.0	33.6	37.4
Male.....	54.6	28.3	11.8	6.3	9.3	49.2	68.9	57.0	48.3	52.6
Female.....	31.4	19.7	8.5	4.3	5.4	24.8	21.5	18.7	18.6	22.0
Non-Hispanic black, both sexes.....	33.7	47.0	12.8	8.8	8.7	22.2	30.6	33.4	31.7	36.0
Male.....	47.9	49.2	14.4	10.8	10.5	31.9	45.5	53.5	50.6	53.1
Female.....	20.8	44.8	11.1	6.8	6.9	12.2	15.6	14.7	14.6	20.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
All origins 3/, both sexes.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4
Male.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0
Female.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8
Hispanic, both sexes.....	14.5	4.4	4.7	3.5	4.4	23.8	28.5	20.4	15.7	15.1
Male.....	21.3	*	5.4	3.9	5.4	35.1	42.8	30.4	23.3	22.8
Female.....	7.3	5.0	4.1	3.1	3.4	11.7	11.0	8.1	6.9	6.3
Mexican, both sexes.....	15.3	4.6	5.7	3.6	4.9	25.0	29.8	20.7	15.9	16.1
Male.....	22.3	*	6.6	3.7	6.1	36.9	45.1	30.2	23.6	23.8
Female.....	7.5	*	4.7	3.4	3.7	12.0	10.6	8.4	6.7	6.9
Puerto Rican, both sexes.....	9.5	*	*	*	*	13.5	17.1	11.7	10.6	13.5
Male.....	14.5	*	*	*	*	20.2	22.3	20.3	*	23.0
Female.....	4.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	14.2	*	*	*	*	*	*	*	*	*
Male.....	20.6	*	*	*	*	*	*	*	*	*
Female.....	7.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	14.4	*	*	3.9	3.4	24.7	27.9	23.4	17.5	12.4
Male.....	21.0	*	*	*	*	36.4	41.7	34.9	25.1	19.5
Female.....	7.8	*	*	*	*	13.5	11.7	9.3	8.7	*
Non-Hispanic 5/, both sexes.....	16.6	3.3	3.9	3.4	5.2	26.4	28.5	21.0	16.9	16.7
Male.....	23.3	3.9	4.1	3.9	6.1	33.3	41.7	31.3	24.6	24.2
Female.....	10.3	2.6	3.8	3.0	4.3	19.1	14.8	10.6	9.3	9.2
Non-Hispanic white, both sexes.....	17.2	2.9	3.7	3.3	5.4	29.4	30.3	21.3	17.0	16.9
Male.....	23.9	3.3	3.8	3.8	6.4	36.6	44.7	31.4	24.1	24.4
Female.....	10.7	2.4	3.6	2.8	4.3	21.7	15.4	11.0	9.8	9.4
Non-Hispanic black, both sexes.....	15.6	5.0	5.2	4.0	4.9	17.0	23.1	23.1	20.2	18.0
Male.....	23.4	6.2	5.4	4.3	5.4	23.5	33.9	37.2	33.0	28.5
Female.....	8.5	*	5.0	3.8	4.3	10.3	12.3	10.0	8.6	8.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All origins 3/, both sexes.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2
Male.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9
Female.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3
Hispanic, both sexes.....	13.9	4.4	4.6	3.5	4.2	23.2	27.8	19.6	14.7	14.2
Male.....	20.2	*	5.3	3.8	5.0	34.3	41.7	29.0	21.6	21.4
Female.....	7.1	5.0	3.9	3.1	3.3	11.3	10.8	7.9	6.6	5.9
Mexican, both sexes.....	14.7	4.6	5.5	3.5	4.7	24.6	29.1	19.9	14.8	15.6
Male.....	21.3	*	6.4	3.6	5.7	36.1	44.0	28.8	21.8	22.9
Female.....	7.4	*	4.6	3.4	3.7	11.9	10.4	8.3	6.6	6.8
Puerto Rican, both sexes.....	8.9	*	*	*	*	13.2	16.8	11.0	9.9	11.8
Male.....	13.5	*	*	*	*	19.6	22.3	*	*	20.4
Female.....	4.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	13.7	*	*	*	*	*	*	*	*	*
Male.....	19.6	*	*	*	*	*	*	*	*	*
Female.....	7.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.5	*	*	3.9	3.1	23.2	27.3	22.2	16.3	10.6
Male.....	19.7	*	*	*	*	34.9	40.5	33.3	23.5	16.8
Female.....	7.4	*	*	*	*	12.0	11.7	8.7	8.1	*
Non-Hispanic 5/, both sexes.....	15.5	3.2	3.8	3.2	4.9	25.5	27.3	19.6	15.7	15.3
Male.....	21.5	3.8	3.9	3.6	5.7	32.1	40.0	29.2	22.6	22.0
Female.....	9.8	2.6	3.7	2.8	4.0	18.7	14.3	10.0	8.9	8.7
Non-Hispanic white, both sexes.....	16.0	2.8	3.5	3.1	5.0	28.5	29.1	19.9	15.7	15.5
Male.....	22.0	3.3	3.6	3.5	6.0	35.4	42.9	29.3	22.0	22.1
Female.....	10.2	2.3	3.5	2.6	4.0	21.2	14.8	10.3	9.3	8.8
Non-Hispanic black, both sexes.....	14.6	4.9	5.0	3.9	4.7	16.2	22.2	21.9	19.2	17.0
Male.....	21.7	*	5.2	4.1	5.3	22.2	32.2	34.8	31.2	26.5
Female.....	8.1	*	4.9	3.6	4.0	10.1	12.2	9.8	8.4	8.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nontransport accidents (W00-X59,Y86)										
All origins 3/, both sexes.....	21.8	22.2	6.1	2.3	2.2	6.9	12.6	13.7	13.8	18.3
Male.....	26.8	25.1	7.4	2.9	3.1	11.0	19.8	21.1	20.5	25.7
Female.....	16.9	19.3	4.8	1.6	1.3	2.6	4.9	6.0	7.0	10.9
Hispanic, both sexes.....	10.6	13.1	4.3	1.1	1.2	4.9	9.1	9.3	9.2	13.7
Male.....	15.2	14.9	5.1	1.3	1.8	7.8	14.4	14.8	14.5	21.0
Female.....	5.8	11.3	3.4	*	*	1.7	2.6	2.5	3.0	5.3
Mexican, both sexes.....	9.5	13.8	4.4	1.2	1.2	5.6	8.4	8.8	8.8	11.6
Male.....	13.9	15.4	5.0	*	1.8	8.9	13.5	14.3	14.0	18.2
Female.....	4.7	12.1	3.7	*	*	2.1	1.8	*	2.6	3.7
Puerto Rican, both sexes.....	14.0	*	*	*	*	*	10.4	15.1	14.5	27.0
Male.....	20.6	*	*	*	*	*	18.4	24.6	24.0	38.7
Female.....	7.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	14.1	*	*	*	*	*	*	*	*	*
Male.....	16.3	*	*	*	*	*	*	*	*	*
Female.....	11.8	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.0	*	5.5	*	*	4.0	11.4	9.0	9.4	15.5
Male.....	17.0	*	*	*	*	7.3	16.4	13.2	14.7	24.4
Female.....	7.0	*	*	*	*	*	*	*	*	6.5
Non-Hispanic 5/, both sexes.....	23.5	24.8	6.6	2.5	2.4	7.3	13.3	14.7	14.8	19.1
Male.....	28.7	27.9	8.0	3.3	3.3	11.5	21.1	22.7	21.9	26.5
Female.....	18.6	21.6	5.2	1.8	1.4	2.8	5.3	6.7	7.8	11.9
Non-Hispanic white, both sexes.....	25.6	21.2	6.5	2.0	2.0	7.9	15.3	16.8	16.5	20.4
Male.....	30.7	25.0	8.0	2.5	2.8	12.5	24.2	25.6	24.2	28.2
Female.....	20.7	17.3	4.9	1.4	1.1	3.1	6.1	7.7	8.7	12.7
Non-Hispanic black, both sexes.....	18.2	42.0	7.6	4.8	3.9	5.2	7.4	10.3	11.5	18.0
Male.....	24.6	43.0	9.1	6.5	5.1	8.4	11.5	16.2	17.6	24.6
Female.....	12.3	40.9	6.1	3.0	2.6	1.9	3.3	4.7	6.0	12.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All origins 3/, both sexes.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7
Male.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9
Female.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4
Hispanic, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.5	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.5	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.5	*	*	*	*	*	*	*	*	*
Male.....	0.6	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.2	0.7	1.6	1.0	0.5	0.4	0.6	0.7	0.6	0.7
Male.....	1.4	*	1.7	1.3	0.5	0.4	0.7	1.0	0.8	1.0
Female.....	1.0	*	1.5	0.7	0.5	0.3	0.6	0.4	0.4	0.5
Non-Hispanic white, both sexes.....	1.1	*	1.3	0.7	0.4	0.4	0.6	0.6	0.6	0.7
Male.....	1.3	*	1.4	0.8	0.4	0.4	0.7	0.9	0.8	0.9
Female.....	0.9	*	1.2	0.5	0.3	*	0.5	0.4	0.4	0.6
Non-Hispanic black, both sexes.....	2.1	*	2.9	2.4	1.0	*	0.8	1.2	0.8	0.9
Male.....	2.7	*	3.0	3.3	*	*	*	1.6	*	*
Female.....	1.7	*	2.9	1.5	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All origins 3/, both sexes.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8
Male.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8
Female.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7
Hispanic, both sexes.....	4.2	*	*	*	*	1.7	3.9	4.8	5.0	8.3
Male.....	6.2	*	*	*	*	2.4	5.6	7.6	7.6	12.0
Female.....	1.9	*	*	*	*	1.8	1.8	1.4	1.9	4.1
Mexican, both sexes.....	3.3	*	*	*	*	1.8	3.0	4.2	4.3	6.6
Male.....	5.1	*	*	*	*	2.4	4.3	6.7	6.6	9.5
Female.....	1.3	*	*	*	*	*	*	*	*	3.0
Puerto Rican, both sexes.....	8.4	*	*	*	*	*	*	11.3	10.6	20.9
Male.....	13.1	*	*	*	*	*	*	*	*	30.2
Female.....	3.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.3	*	*	*	*	*	*	*	*	*
Male.....	3.8	*	*	*	*	*	*	*	*	*
Female.....	2.7	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.1	*	*	*	*	*	5.6	4.5	5.2	9.1
Male.....	7.4	*	*	*	*	*	7.4	6.5	8.1	13.6
Female.....	2.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	7.6	*	*	*	0.2	3.4	8.5	9.9	10.2	13.6
Male.....	10.2	*	*	*	0.3	5.0	13.2	14.8	14.2	17.7
Female.....	5.1	*	*	*	*	1.6	3.7	5.0	6.1	9.5
Non-Hispanic white, both sexes.....	8.2	*	*	*	0.2	4.2	10.4	12.0	11.7	14.9
Male.....	10.9	*	*	*	*	6.3	16.1	17.7	16.4	19.5
Female.....	5.5	*	*	*	*	1.9	4.5	6.1	6.9	10.2
Non-Hispanic black, both sexes.....	6.7	*	*	*	*	0.8	2.9	4.6	6.3	11.5
Male.....	9.1	*	*	*	*	*	4.3	6.8	8.8	14.1
Female.....	4.4	*	*	*	*	*	1.5	2.7	4.1	9.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes.....	5.8	19.2	1.6	0.6	0.6	1.2	1.9	1.9	2.2	2.6
Male.....	6.8	21.7	2.0	0.8	1.0	2.0	3.1	3.1	3.7	4.1
Female.....	4.8	16.5	1.1	0.4	0.3	0.4	0.5	0.6	0.7	1.0
Hispanic, both sexes.....	2.5	9.8	1.2	*	*	1.2	2.1	1.6	2.1	2.4
Male.....	3.5	11.7	1.6	*	*	2.0	3.5	2.5	3.3	4.1
Female.....	1.5	7.9	*	*	*	*	*	*	*	*
Mexican, both sexes.....	2.5	10.4	1.0	*	*	1.5	2.1	1.6	2.3	2.6
Male.....	3.6	11.3	*	*	*	2.3	3.6	2.6	3.7	4.5
Female.....	1.3	9.4	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	2.1	*	*	*	*	*	*	*	*	*
Male.....	2.9	*	*	*	*	*	*	*	*	*
Female.....	1.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.8	*	*	*	*	*	*	*	*	*
Male.....	3.9	*	*	*	*	*	*	*	*	*
Female.....	3.7	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.5	*	*	*	*	*	*	*	*	*
Male.....	3.5	*	*	*	*	*	*	*	*	*
Female.....	1.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.3	21.9	1.7	0.6	0.7	1.2	1.8	1.9	2.2	2.6
Male.....	7.4	24.6	2.1	0.8	1.1	2.0	3.0	3.3	3.8	4.1
Female.....	5.3	19.0	1.2	0.4	0.3	0.4	0.5	0.6	0.7	1.1
Non-Hispanic white, both sexes.....	6.9	18.4	1.6	0.6	0.7	1.4	1.9	2.2	2.3	2.6
Male.....	7.9	22.0	2.2	0.8	1.1	2.2	3.3	3.6	3.9	4.2
Female.....	5.9	14.7	0.9	0.5	*	0.4	0.5	0.7	0.8	1.1
Non-Hispanic black, both sexes.....	4.9	38.0	1.9	*	0.7	0.7	1.3	1.7	2.3	2.9
Male.....	6.3	38.7	1.7	*	*	*	2.1	3.0	4.0	4.7
Female.....	3.7	37.3	2.0	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All origins 3/, both sexes.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0
Male.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7
Female.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3
Hispanic, both sexes.....	5.3	*	1.0	6.4	9.5	7.6	6.7	7.0
Male.....	8.6	*	1.2	9.9	15.3	11.9	10.5	11.1
Female.....	1.8	*	*	2.7	2.4	2.2	2.3	2.4
Mexican, both sexes.....	4.7	*	1.0	6.3	9.3	7.1	6.2	6.1
Male.....	7.6	*	*	9.6	14.9	11.4	9.5	9.8
Female.....	1.5	*	*	2.7	2.3	1.7	2.3	*
Puerto Rican, both sexes.....	6.3	*	*	*	10.1	9.6	10.6	8.8
Male.....	10.9	*	*	*	17.8	*	*	*
Female.....	1.8	*	*	*	*	*	*	*
Cuban, both sexes.....	8.3	*	*	*	*	*	*	*
Male.....	13.2	*	*	*	*	*	*	*
Female.....	3.1	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.3	*	*	8.4	9.6	8.7	7.0	8.6
Male.....	10.0	*	*	13.8	15.0	12.8	10.9	14.1
Female.....	2.6	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.9	*	1.4	8.5	13.1	13.5	14.3	15.2
Male.....	19.2	*	1.8	13.1	22.1	22.4	22.9	23.6
Female.....	5.0	*	1.0	3.7	3.8	4.6	5.9	6.9
Non-Hispanic white, both sexes.....	13.5	*	1.4	9.2	14.1	14.6	16.1	17.4
Male.....	21.6	*	1.8	14.2	23.8	24.1	25.6	26.9
Female.....	5.7	*	1.0	3.9	4.1	4.8	6.5	7.9
Non-Hispanic black, both sexes.....	5.3	*	1.4	4.8	10.0	10.6	8.9	6.9
Male.....	9.2	*	1.7	7.5	17.5	18.8	14.8	11.6
Female.....	1.8	*	*	1.9	2.5	3.0	3.6	2.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All origins 3/, both sexes.....	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9
Male.....	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8
Female.....	1.5	*	*	1.0	1.1	1.3	1.7	1.9
Hispanic, both sexes.....	2.1	*	*	2.9	4.5	3.1	2.3	2.1
Male.....	3.7	*	*	5.1	7.6	4.9	3.8	3.6
Female.....	0.4	*	*	*	*	*	*	*
Mexican, both sexes.....	2.0	*	*	3.0	4.6	3.0	2.3	1.6
Male.....	3.5	*	*	5.2	7.7	4.6	3.8	2.8
Female.....	0.4	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	2.0	*	*	*	*	*	*	*
Male.....	3.8	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.3	*	*	*	*	*	*	*
Male.....	5.0	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.3	*	*	3.9	4.0	3.4	*	3.4
Male.....	4.2	*	*	7.3	6.7	5.4	*	6.0
Female.....	0.5	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.3	*	0.3	3.9	6.6	6.3	6.5	6.5
Male.....	11.1	*	0.5	6.7	11.9	11.2	11.0	11.0
Female.....	1.7	*	*	1.1	1.2	1.4	2.0	2.1
Non-Hispanic white, both sexes.....	7.3	*	0.4	4.4	7.1	6.8	7.3	7.6
Male.....	12.8	*	0.6	7.5	12.8	12.0	12.2	12.7
Female.....	1.9	*	*	1.2	1.3	1.6	2.4	2.5
Non-Hispanic black, both sexes.....	2.7	*	*	2.3	6.0	5.7	4.4	3.0
Male.....	5.1	*	*	3.8	10.9	10.8	8.2	5.7
Female.....	0.5	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes.....	5.3	*	1.1	4.4	6.2	6.6	7.3	8.1
Male.....	7.6	*	1.2	6.2	9.7	10.3	11.0	11.9
Female.....	3.1	*	0.9	2.5	2.4	2.8	3.5	4.4
Hispanic, both sexes.....	3.2	*	1.0	3.5	5.0	4.5	4.4	4.9
Male.....	4.9	*	1.1	4.8	7.7	7.0	6.7	7.5
Female.....	1.4	*	*	2.2	1.7	1.3	1.8	2.0
Mexican, both sexes.....	2.7	*	0.9	3.3	4.7	4.2	3.9	4.6
Male.....	4.1	*	*	4.4	7.2	6.7	5.7	7.1
Female.....	1.1	*	*	2.1	*	*	1.7	*
Puerto Rican, both sexes.....	4.3	*	*	*	*	*	*	*
Male.....	7.0	*	*	*	*	*	*	*
Female.....	1.6	*	*	*	*	*	*	*
Cuban, both sexes.....	5.0	*	*	*	*	*	*	*
Male.....	8.1	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.9	*	*	4.5	5.6	5.3	5.1	5.2
Male.....	5.8	*	*	*	8.3	7.4	7.6	8.1
Female.....	2.1	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.7	*	1.1	4.5	6.4	7.1	7.9	8.7
Male.....	8.1	*	1.3	6.4	10.2	11.2	11.9	12.6
Female.....	3.4	*	0.9	2.6	2.6	3.1	3.9	4.8
Non-Hispanic white, both sexes.....	6.3	*	1.0	4.8	7.0	7.7	8.8	9.8
Male.....	8.9	*	1.2	6.7	11.0	12.2	13.4	14.2
Female.....	3.7	*	0.8	2.7	2.8	3.2	4.1	5.4
Non-Hispanic black, both sexes.....	2.6	*	1.2	2.5	4.0	4.9	4.5	3.9
Male.....	4.1	*	1.3	3.8	6.6	8.0	6.6	5.9
Female.....	1.3	*	*	*	1.4	2.1	2.6	2.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All origins 3/, both sexes.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3
Male.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2
Female.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3
Hispanic, both sexes.....	7.9	8.1	2.3	0.7	1.1	14.8	19.7	15.2	9.9	8.2
Male.....	13.0	9.1	2.5	*	1.4	25.8	32.8	24.5	15.7	13.1
Female.....	2.4	7.0	2.1	*	*	2.9	3.7	3.6	3.3	2.7
Mexican, both sexes.....	8.5	7.1	2.5	*	1.1	16.7	20.9	16.0	10.1	8.6
Male.....	14.0	7.8	2.7	*	*	29.0	34.8	26.0	16.2	13.9
Female.....	2.3	*	2.4	*	*	3.2	3.2	3.0	2.8	2.3
Puerto Rican, both sexes.....	7.5	*	*	*	*	10.9	20.8	19.2	11.7	8.1
Male.....	12.7	*	*	*	*	20.2	32.4	33.7	22.5	*
Female.....	2.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	5.3	*	*	*	*	*	*	*	*	*
Male.....	7.4	*	*	*	*	*	*	*	*	*
Female.....	3.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.9	*	*	*	*	12.1	16.7	11.4	9.5	6.8
Male.....	11.0	*	*	*	*	21.4	27.5	17.2	13.3	9.8
Female.....	2.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.5	7.8	2.4	0.6	1.0	8.2	13.9	12.7	9.0	7.0
Male.....	8.7	7.5	2.5	0.6	1.2	13.2	23.6	21.5	14.5	10.8
Female.....	2.5	8.1	2.3	0.6	0.7	3.0	4.0	3.9	3.6	3.4
Non-Hispanic white, both sexes.....	2.7	5.9	1.4	0.4	0.6	2.6	4.4	4.2	3.8	3.7
Male.....	3.6	6.1	1.6	0.4	0.7	3.3	6.1	5.8	5.1	5.2
Female.....	1.8	5.8	1.3	0.4	0.4	1.8	2.7	2.6	2.4	2.3
Non-Hispanic black, both sexes.....	21.7	15.8	6.2	1.4	2.6	32.8	58.1	55.1	36.8	26.1
Male.....	38.3	13.9	6.3	1.4	3.5	56.9	105.9	102.9	67.0	44.6
Female.....	6.6	17.7	6.1	1.5	1.6	7.9	10.2	10.6	9.5	9.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All origins 3/, both sexes.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9
Male.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3
Female.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4
Hispanic, both sexes.....	5.4	*	*	*	0.7	12.4	15.4	11.3	7.3	5.4
Male.....	9.5	*	*	*	1.1	22.0	26.3	19.3	12.1	9.4
Female.....	1.1	*	*	*	*	2.1	2.0	1.4	1.8	*
Mexican, both sexes.....	6.0	*	*	*	*	14.5	16.6	12.3	7.6	6.1
Male.....	10.6	*	*	*	*	25.6	28.4	21.0	12.6	10.2
Female.....	1.0	*	*	*	*	2.3	1.7	*	*	*
Puerto Rican, both sexes.....	5.0	*	*	*	*	8.3	16.5	12.7	9.6	*
Male.....	8.8	*	*	*	*	15.1	27.3	23.1	*	*
Female.....	1.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.5	*	*	*	*	*	*	*	*	*
Male.....	5.3	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.1	*	*	*	*	8.8	11.5	8.0	6.1	3.4
Male.....	7.2	*	*	*	*	15.9	19.9	12.8	9.5	6.5
Female.....	1.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.7	*	0.2	0.2	0.6	6.7	11.4	10.3	6.9	4.7
Male.....	6.4	*	*	0.2	1.0	11.4	20.3	18.5	11.8	8.0
Female.....	1.2	*	*	*	0.3	1.7	2.3	2.1	2.1	1.5
Non-Hispanic white, both sexes.....	1.5	*	*	*	0.3	1.6	2.7	2.7	2.6	2.2
Male.....	2.1	*	*	*	0.5	2.2	4.0	4.2	3.8	3.5
Female.....	0.8	*	*	*	*	1.0	1.3	1.1	1.3	1.0
Non-Hispanic black, both sexes.....	16.6	*	*	*	1.9	28.8	52.4	48.5	30.4	19.1
Male.....	31.4	*	*	*	3.0	51.8	97.8	93.4	57.3	35.9
Female.....	3.1	*	*	*	*	5.1	6.9	6.7	6.1	4.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
All origins 3/, both sexes.....	2.0	7.8	2.1	0.4	0.3	1.7	2.9	2.7	2.2	2.4
Male.....	2.5	7.9	2.3	0.4	0.3	2.2	4.0	3.5	2.9	2.9
Female.....	1.4	7.7	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9
Hispanic, both sexes.....	2.5	7.7	2.1	*	*	2.4	4.4	3.8	2.6	2.8
Male.....	3.5	8.9	2.3	*	*	3.9	6.5	5.2	3.6	3.7
Female.....	1.4	6.5	1.9	*	*	*	1.8	2.1	1.5	1.8
Mexican, both sexes.....	2.4	6.8	2.2	*	*	2.2	4.2	3.7	2.5	2.5
Male.....	3.5	7.5	2.4	*	*	3.4	6.4	5.1	3.7	3.7
Female.....	1.3	*	2.1	*	*	*	*	1.9	*	*
Puerto Rican, both sexes.....	2.5	*	*	*	*	*	*	*	*	*
Male.....	4.0	*	*	*	*	*	*	*	*	*
Female.....	1.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	1.8	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.8	*	*	*	*	3.3	5.1	3.4	3.4	3.4
Male.....	3.8	*	*	*	*	*	7.6	*	*	*
Female.....	1.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	7.7	2.1	0.4	0.3	1.6	2.5	2.4	2.1	2.3
Male.....	2.3	7.4	2.3	0.4	0.2	1.9	3.3	3.0	2.7	2.7
Female.....	1.4	8.0	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9
Non-Hispanic white, both sexes.....	1.3	5.9	1.4	0.3	0.2	1.0	1.8	1.6	1.2	1.5
Male.....	1.5	6.1	1.5	*	*	1.1	2.1	1.6	1.3	1.7
Female.....	1.0	5.8	1.2	*	*	0.9	1.4	1.5	1.1	1.3
Non-Hispanic black, both sexes.....	5.1	15.3	5.4	0.9	0.7	4.0	5.7	6.6	6.4	7.0
Male.....	6.9	13.6	5.6	*	*	5.1	8.1	9.5	9.7	8.8
Female.....	3.5	17.1	5.3	*	*	2.8	3.3	3.9	3.5	5.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9
Male.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8
Female.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9
Hispanic, both sexes.....	0.8	*	*	*	*	*	0.8	1.0	0.9	1.2
Male.....	1.1	*	*	*	*	*	1.2	1.4	1.3	1.8
Female.....	0.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.7	*	*	*	*	*	*	1.0	*	*
Male.....	1.0	*	*	*	*	*	*	1.4	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	1.7	*	*	*	*	*	*	*	*	*
Male.....	2.5	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.8	*	*	*	*	*	*	*	*	*
Male.....	1.1	*	*	*	*	*	*	*	*	*
Female.....	0.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	2.8	0.3	0.1	0.2	0.8	1.8	2.1	2.4	3.2
Male.....	2.4	3.3	0.3	*	0.3	1.2	2.7	3.0	3.3	4.2
Female.....	1.3	2.3	0.3	*	*	0.4	0.9	1.2	1.6	2.2
Non-Hispanic white, both sexes.....	1.9	2.3	0.3	*	0.2	0.8	2.1	2.4	2.5	3.4
Male.....	2.4	2.7	*	*	*	1.2	3.2	3.5	3.3	4.3
Female.....	1.4	2.0	*	*	*	0.4	1.0	1.4	1.7	2.4
Non-Hispanic black, both sexes.....	1.9	4.9	*	*	*	0.7	1.1	1.0	2.7	3.3
Male.....	2.7	*	*	*	*	*	1.6	*	4.2	4.8
Female.....	1.1	*	*	*	*	*	*	*	1.4	1.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5, F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/										
All origins 3/, both sexes.....	10.5	0.7	0.2	*	0.2	3.9	10.0	12.2	13.1	18.4
Male.....	13.4	*	*	*	0.2	5.6	14.7	17.3	17.6	23.6
Female.....	7.6	*	*	*	0.3	2.1	5.0	6.7	8.5	13.2
Hispanic, both sexes.....	5.5	*	*	*	*	2.0	4.5	6.0	6.7	10.7
Male.....	7.8	*	*	*	*	2.8	6.1	8.7	9.9	15.3
Female.....	2.9	*	*	*	*	*	2.4	2.6	2.9	5.5
Mexican, both sexes.....	4.4	*	*	*	*	1.9	3.3	5.1	6.1	8.2
Male.....	6.5	*	*	*	*	2.8	4.5	7.9	8.8	11.5
Female.....	2.2	*	*	*	*	*	1.7	*	2.7	4.1
Puerto Rican, both sexes.....	11.4	*	*	*	*	*	10.1	15.1	14.9	29.0
Male.....	17.5	*	*	*	*	*	18.4	24.6	24.0	41.4
Female.....	5.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	4.8	*	*	*	*	*	*	*	*	*
Male.....	5.5	*	*	*	*	*	*	*	*	*
Female.....	4.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.2	*	*	*	*	*	6.5	5.9	6.2	11.4
Male.....	8.4	*	*	*	*	*	7.6	6.7	9.5	17.1
Female.....	4.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.2	0.9	0.2	*	0.3	4.3	11.2	13.7	14.5	19.7
Male.....	14.3	*	*	*	*	6.1	16.7	19.7	19.3	25.0
Female.....	8.3	*	*	*	0.3	2.3	5.5	7.6	9.6	14.4
Non-Hispanic white, both sexes.....	12.1	*	*	*	0.3	5.3	13.7	16.4	16.6	21.5
Male.....	15.3	*	*	*	*	7.6	20.5	23.6	22.1	27.3
Female.....	9.1	*	*	*	*	2.8	6.7	9.1	10.9	15.7
Non-Hispanic black, both sexes.....	9.7	*	*	*	*	1.0	3.5	6.4	9.9	16.8
Male.....	13.1	*	*	*	*	1.4	4.9	9.3	13.6	21.1
Female.....	6.6	*	*	*	*	*	2.1	3.8	6.6	13.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All origins 3/, both sexes.....	10.1	*	0.3	0.3	1.1	12.0	19.3	16.9	13.3	11.2
Male.....	17.6	*	0.4	0.3	1.8	20.7	34.2	29.8	22.5	19.0
Female.....	2.7	*	0.3	0.3	0.4	2.9	3.5	3.4	3.9	3.3
Hispanic, both sexes.....	7.9	*	*	*	0.9	15.8	21.0	15.1	10.2	7.9
Male.....	13.9	*	*	*	1.4	28.1	35.9	25.4	17.0	13.7
Female.....	1.5	*	*	*	*	2.7	2.6	2.4	2.3	1.3
Mexican, both sexes.....	8.5	*	*	*	0.9	18.1	22.6	16.1	10.5	8.1
Male.....	14.9	*	*	*	1.5	32.0	38.5	27.1	17.6	13.7
Female.....	1.4	*	*	*	*	3.0	2.4	1.9	2.1	*
Puerto Rican, both sexes.....	7.1	*	*	*	*	9.5	21.2	16.5	12.7	8.1
Male.....	12.9	*	*	*	*	17.4	36.2	29.5	25.5	*
Female.....	1.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	6.8	*	*	*	*	*	*	*	*	*
Male.....	10.4	*	*	*	*	*	*	*	*	*
Female.....	3.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.7	*	*	*	*	13.0	16.2	11.8	8.6	7.1
Male.....	11.8	*	*	*	*	23.9	27.8	18.6	13.8	13.0
Female.....	1.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.4	*	0.4	0.3	1.2	11.3	18.8	17.3	13.9	11.8
Male.....	18.2	*	0.4	0.4	1.9	19.2	33.6	30.9	23.7	20.0
Female.....	2.9	*	*	*	0.4	2.9	3.6	3.6	4.2	3.7
Non-Hispanic white, both sexes.....	9.1	*	*	0.2	0.9	6.6	10.5	10.0	10.3	10.4
Male.....	15.6	*	*	*	1.4	10.8	17.9	17.0	16.8	17.0
Female.....	2.9	*	*	*	0.3	2.3	2.7	2.8	3.8	3.6
Non-Hispanic black, both sexes.....	19.9	*	1.1	0.8	2.3	31.9	60.1	55.7	35.8	22.8
Male.....	37.6	*	*	*	3.7	57.3	111.7	107.4	67.4	42.8
Female.....	3.8	*	*	*	*	5.9	8.3	7.6	7.2	4.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

All origins 3/, both sexes.....	231.7	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
Male.....	289.5	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
Female.....	174.5	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
Hispanic, both sexes.....	168.3	258.4	385.6	557.4	862.6	1,289.9	2,017.4	3,218.2	5,038.4	9,483.8
Male.....	222.3	343.3	518.0	706.7	1,118.2	1,606.8	2,514.1	3,990.1	6,105.0	9,932.8
Female.....	109.8	170.7	256.1	417.9	639.0	1,027.7	1,634.2	2,669.9	4,334.3	9,253.0
Mexican, both sexes.....	166.6	252.4	366.6	554.0	863.3	1,414.2	2,191.3	3,172.1	4,915.8	8,907.9
Male.....	224.3	328.5	487.6	668.7	1,046.8	1,649.4	2,731.4	4,065.8	5,617.8	*
Female.....	103.3	166.3	244.8	438.2	681.8	1,195.7	1,786.4	2,579.4	4,377.8	8,485.7
Puerto Rican, both sexes.....	267.2	415.0	503.1	769.5	1,166.4	1,196.5	2,317.2	*	*	*
Male.....	402.4	609.0	645.1	954.7	*	1,443.0	*	*	*	*
Female.....	162.7	254.4	349.0	584.8	803.9	978.8	*	*	*	*
Cuban, both sexes.....	130.2	228.3	616.0	498.9	749.0	1,030.2	1,858.7	2,612.4	3,795.1	8,007.9
Male.....	162.3	296.6	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	145.9	219.8	341.5	476.4	758.3	1,214.9	1,611.8	3,410.8	*	*
Male.....	179.6	296.7	477.1	681.8	1,085.3	1,783.8	*	*	*	*
Female.....	108.4	151.8	227.6	324.3	527.9	885.5	1,209.9	*	*	*
Non-Hispanic 5/, both sexes.....	239.8	361.6	522.0	747.3	1,158.9	1,756.4	2,715.6	4,243.0	6,789.9	13,997.7
Male.....	298.4	454.3	667.7	931.4	1,428.8	2,157.1	3,321.9	5,232.9	8,272.9	15,260.4
Female.....	182.7	271.9	382.9	573.6	912.5	1,406.4	2,226.0	3,531.1	5,879.4	13,433.1
Non-Hispanic white, both sexes.....	220.9	331.1	477.6	696.6	1,105.0	1,700.5	2,669.8	4,207.6	6,807.8	14,191.1
Male.....	279.0	418.2	612.7	863.8	1,355.2	2,082.3	3,257.1	5,178.6	8,308.7	15,489.2
Female.....	163.0	244.9	345.4	535.3	871.8	1,360.0	2,185.0	3,496.3	5,878.7	13,609.6
Non-Hispanic black, both sexes.....	397.2	621.7	928.8	1,296.4	1,845.3	2,554.0	3,637.4	5,336.6	7,587.7	13,492.2
Male.....	480.0	782.6	1,211.2	1,702.7	2,416.8	3,298.9	4,655.2	6,839.3	9,346.6	14,647.3
Female.....	324.4	481.7	690.6	962.5	1,400.6	2,001.4	2,947.5	4,447.1	6,662.3	13,035.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

All origins 3/, both sexes.....	0.1	0.2	0.3	0.6	1.3	2.2	4.6	9.3	17.5	34.3
Male.....	*	*	0.3	0.6	1.3	2.2	4.3	9.3	17.8	32.0
Female.....	*	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
Hispanic, both sexes.....	*	*	*	*	*	*	*	6.0	9.4	18.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	17.3
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.2	0.3	0.6	1.3	2.2	4.7	9.5	17.9	35.0
Male.....	*	*	0.3	0.6	1.4	2.2	4.4	9.5	18.3	32.4
Female.....	*	*	0.3	0.6	1.2	2.2	5.0	9.5	17.6	36.1
Non-Hispanic white, both sexes.....	*	0.1	0.3	0.6	1.3	2.3	5.0	10.0	18.8	37.3
Male.....	*	*	0.3	0.5	1.5	2.2	4.7	10.0	19.3	34.5
Female.....	*	*	0.3	0.6	1.2	2.4	5.4	10.0	18.5	38.6
Non-Hispanic black, both sexes.....	*	*	*	*	*	2.1	3.4	6.1	11.6	16.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	6.6	12.0	16.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningococcal infection (A39)

Table with 11 columns representing age groups (40-44 years to 85 years and over) and rows representing various demographic categories such as All origins 3/, both sexes; Male; Female; Hispanic, both sexes; Mexican, both sexes; Puerto Rican, both sexes; Cuban, both sexes; Other Hispanic 4/, both sexes; Non-Hispanic 5/, both sexes; Non-Hispanic white, both sexes; Non-Hispanic black, both sexes. Each cell contains an asterisk (*) indicating data presence.

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Septicemia (A40-A41)

All origins 3/, both sexes.....	2.3	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
Male.....	2.4	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
Female.....	2.3	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
Hispanic, both sexes.....	1.6	3.5	4.8	7.8	12.4	19.1	32.8	51.2	83.7	138.0
Male.....	2.0	4.5	6.3	8.1	13.5	22.8	33.1	56.4	91.5	147.1
Female.....	*	2.4	3.3	7.6	11.4	16.1	32.6	47.5	78.5	133.2
Mexican, both sexes.....	1.4	3.3	4.8	8.0	11.9	21.0	35.3	43.9	78.7	129.3
Male.....	*	4.1	6.3	6.9	11.0	25.2	31.6	*	*	*
Female.....	*	*	*	9.1	12.8	17.0	38.1	46.2	80.5	129.9
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	25.2	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.4	4.1	7.1	10.5	16.5	25.3	41.7	65.8	104.5	188.7
Male.....	2.4	4.4	7.6	11.6	17.3	27.1	47.5	75.4	122.4	206.3
Female.....	2.5	3.8	6.6	9.4	15.8	23.7	37.0	58.9	93.5	180.8
Non-Hispanic white, both sexes.....	1.8	3.2	5.8	8.9	13.9	22.0	37.8	61.2	98.1	179.6
Male.....	1.8	3.3	6.3	9.7	14.4	23.6	43.0	70.4	114.6	198.5
Female.....	1.9	3.0	5.3	8.0	13.4	20.5	33.5	54.5	87.8	171.1
Non-Hispanic black, both sexes.....	6.4	10.5	17.5	26.0	40.7	59.0	86.9	129.4	199.5	324.7
Male.....	7.0	12.0	19.0	31.2	45.4	66.3	105.1	156.7	248.9	346.5
Female.....	5.9	9.1	16.1	21.7	37.0	53.6	74.6	113.2	173.5	316.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Measles (B05)

Table with 11 columns representing age groups (40-44 years to 85 years and over) and rows for various demographic categories including All origins 3/, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic 4/, Non-Hispanic 5/, Non-Hispanic white, and Non-Hispanic black, each split by sex. All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

All origins 3/, both sexes.....	2.0	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
Male.....	2.8	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
Female.....	1.3	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
Hispanic, both sexes.....	3.0	5.8	8.2	7.4	8.1	7.3	6.7	7.7	*	*
Male.....	4.6	8.6	12.3	11.3	9.1	9.6	*	*	*	*
Female.....	*	2.8	4.2	3.7	7.3	5.4	7.3	9.8	*	*
Mexican, both sexes.....	3.4	5.0	7.1	6.5	6.6	8.6	*	*	*	*
Male.....	5.1	6.9	11.3	9.5	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	17.1	19.5	19.5	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.9	4.1	6.3	4.6	3.2	3.0	3.4	3.3	3.4	2.6
Male.....	2.4	6.1	9.6	7.0	4.2	3.7	3.6	3.8	3.2	3.0
Female.....	1.3	2.2	3.2	2.4	2.2	2.3	3.2	3.0	3.5	2.4
Non-Hispanic white, both sexes.....	1.9	4.1	5.9	3.9	2.7	2.6	2.9	3.0	2.9	2.4
Male.....	2.5	6.0	9.0	5.9	3.5	3.2	3.2	3.6	2.6	2.7
Female.....	1.3	2.2	2.8	2.0	1.9	2.1	2.6	2.6	3.1	2.3
Non-Hispanic black, both sexes.....	2.0	5.2	10.5	10.4	6.6	4.4	5.0	3.8	6.8	*
Male.....	2.3	8.2	16.4	17.0	10.0	7.0	*	*	*	*
Female.....	1.7	2.6	5.5	4.9	3.9	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes.....	12.3	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
Male.....	17.9	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
Female.....	6.7	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
Hispanic, both sexes.....	12.4	14.3	13.9	9.6	6.6	6.4	*	*	*	*
Male.....	18.1	22.1	21.9	15.6	12.6	9.6	*	*	*	*
Female.....	6.1	6.3	6.0	4.0	*	*	*	*	*	*
Mexican, both sexes.....	7.2	6.4	6.9	4.4	4.3	*	*	*	*	*
Male.....	12.4	11.0	12.2	7.1	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	58.7	72.5	54.5	34.4	*	*	*	*	*	*
Male.....	85.9	111.5	71.1	*	*	*	*	*	*	*
Female.....	37.7	40.3	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	8.8	10.1	8.4	*	*	*	*	*	*	*
Male.....	12.5	16.8	14.7	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	12.2	11.1	9.0	5.9	3.9	2.8	1.4	0.9	0.4	*
Male.....	17.7	16.8	13.9	9.9	6.5	4.6	2.3	1.5	*	*
Female.....	6.7	5.6	4.3	2.1	1.5	1.2	0.6	0.5	*	*
Non-Hispanic white, both sexes.....	5.8	4.9	3.7	2.7	1.6	1.1	0.6	0.4	*	*
Male.....	9.8	8.3	6.4	4.9	2.9	2.1	1.2	0.8	*	*
Female.....	1.8	1.5	1.0	0.5	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	52.8	53.0	48.2	33.6	24.8	17.5	8.6	6.4	*	*
Male.....	71.8	78.3	73.9	56.8	42.4	29.4	15.3	9.8	*	*
Female.....	36.0	31.1	26.6	14.6	11.1	8.7	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/, both sexes.....	1.2	2.5	3.5	4.2	4.4	6.0	8.3	11.2	15.9	22.5
Male.....	1.5	3.3	4.9	5.6	5.2	7.2	9.9	12.9	18.4	29.6
Female.....	0.9	1.6	2.2	2.8	3.6	5.0	7.0	10.0	14.4	19.3
Hispanic, both sexes.....	1.9	3.5	5.3	6.6	5.2	6.1	9.6	14.4	15.0	18.6
Male.....	2.5	5.0	7.4	8.8	7.7	7.7	10.3	16.1	19.8	*
Female.....	*	2.0	3.3	4.6	*	*	8.9	13.1	*	19.4
Mexican, both sexes.....	2.1	4.3	6.0	7.7	6.6	8.0	12.8	16.5	*	*
Male.....	2.9	5.6	8.6	10.3	9.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	2.3	3.4	4.0	4.3	6.0	8.2	11.0	15.9	22.6
Male.....	1.3	3.1	4.7	5.3	5.0	7.1	9.9	12.8	18.2	30.2
Female.....	0.9	1.6	2.1	2.7	3.6	5.0	6.9	9.8	14.5	19.2
Non-Hispanic white, both sexes.....	1.0	2.3	3.3	3.7	4.0	5.7	7.8	10.7	15.6	22.6
Male.....	1.1	3.1	4.5	4.9	4.6	7.0	9.3	12.4	17.9	30.8
Female.....	0.8	1.5	2.1	2.6	3.4	4.5	6.7	9.4	14.2	18.9
Non-Hispanic black, both sexes.....	2.1	3.0	4.0	5.9	6.9	7.4	10.5	15.1	19.2	24.8
Male.....	2.5	4.2	6.0	8.9	9.4	7.5	15.0	19.1	22.0	30.5
Female.....	1.8	1.9	2.3	3.5	4.9	7.4	7.4	12.7	17.8	22.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

All origins 3/, both sexes.....	44.3	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
Male.....	39.2	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
Female.....	49.2	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
Hispanic, both sexes.....	28.6	54.9	102.5	166.7	270.3	397.8	567.8	784.8	1,009.2	1,131.2
Male.....	25.1	52.3	111.2	181.5	322.8	481.2	726.1	1,052.8	1,373.1	1,508.8
Female.....	32.3	57.5	93.9	152.9	224.3	328.9	445.6	594.4	768.9	937.0
Mexican, both sexes.....	27.6	50.8	92.8	156.9	255.0	415.3	594.1	749.8	978.6	1,069.4
Male.....	24.4	45.3	96.7	155.6	284.7	486.7	767.8	1,071.4	1,283.6	*
Female.....	31.0	57.1	88.8	158.2	225.6	348.9	463.9	536.5	744.9	855.0
Puerto Rican, both sexes.....	37.8	73.0	120.7	217.5	349.4	330.4	600.1	*	*	*
Male.....	35.3	88.6	141.2	241.9	*	*	*	*	*	*
Female.....	39.8	60.1	98.4	193.2	253.4	295.4	*	*	*	*
Cuban, both sexes.....	19.8	62.6	190.3	185.7	309.2	372.1	606.5	662.0	721.8	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	29.6	55.7	104.2	160.5	252.5	418.6	480.0	938.7	*	*
Male.....	25.0	52.9	112.8	206.3	316.4	576.0	*	*	*	*
Female.....	34.6	58.1	96.9	126.6	207.4	327.5	361.1	*	*	*
Non-Hispanic 5/, both sexes.....	46.4	90.3	160.5	269.0	439.1	655.1	913.3	1,193.3	1,443.5	1,674.3
Male.....	41.3	89.9	173.4	298.3	504.4	775.9	1,114.3	1,527.7	1,918.9	2,387.3
Female.....	51.4	90.6	148.1	241.4	379.5	549.6	751.0	952.9	1,151.7	1,355.6
Non-Hispanic white, both sexes.....	44.0	85.4	152.5	260.8	433.3	651.5	913.4	1,196.0	1,447.9	1,680.9
Male.....	39.2	84.5	163.2	285.6	491.2	764.3	1,100.3	1,514.8	1,911.4	2,387.1
Female.....	48.7	86.3	142.0	236.9	379.4	550.9	759.0	962.4	1,161.0	1,364.5
Non-Hispanic black, both sexes.....	66.0	132.9	240.8	385.1	579.8	817.5	1,076.5	1,373.5	1,600.7	1,794.3
Male.....	58.0	137.6	280.7	466.2	742.0	1,054.9	1,467.0	1,952.1	2,376.3	2,781.1
Female.....	73.2	128.9	207.2	318.5	453.6	641.5	811.8	1,031.0	1,192.7	1,404.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/, both sexes.....	0.9	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
Male.....	1.2	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
Female.....	0.5	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
Hispanic, both sexes.....	*	1.3	1.9	2.4	5.3	6.1	6.9	7.2	10.9	13.8
Male.....	*	2.1	3.2	4.5	8.1	9.9	12.4	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	4.5	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	2.1	4.0	5.3	7.7	9.6	11.3	13.0	15.1	18.2
Male.....	1.4	3.3	6.4	8.9	12.1	15.2	16.6	19.8	21.9	24.9
Female.....	0.5	1.0	1.6	1.8	3.6	4.8	6.9	8.2	11.0	15.1
Non-Hispanic white, both sexes.....	0.7	1.7	3.5	4.8	7.1	9.2	11.2	13.0	15.2	18.7
Male.....	1.1	2.6	5.6	8.1	11.1	14.2	15.9	19.3	21.8	24.8
Female.....	0.4	0.8	1.5	1.6	3.5	4.8	7.4	8.4	11.1	16.0
Non-Hispanic black, both sexes.....	1.9	4.7	7.9	9.8	13.6	14.1	14.0	15.1	15.7	12.7
Male.....	2.9	7.9	14.1	17.4	24.3	26.7	27.6	26.1	26.4	27.4
Female.....	*	1.9	2.7	3.5	5.4	4.8	4.7	8.6	10.0	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
All origins 3/, both sexes.....	1.0	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
Male.....	1.6	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
Female.....	0.3	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
Hispanic, both sexes.....	*	*	1.4	3.2	6.8	9.1	11.9	14.1	16.5	18.6
Male.....	*	*	2.9	5.4	13.3	16.4	20.7	24.8	31.1	29.4
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	3.1	5.3	8.0	10.3	14.8	*	*
Male.....	*	*	*	*	10.2	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	2.5	5.0	8.5	13.6	18.1	22.0	26.7	28.2	26.2
Male.....	1.8	4.3	8.6	14.8	23.6	31.5	38.1	48.0	52.7	49.4
Female.....	0.3	0.7	1.5	2.5	4.5	6.4	8.9	11.4	13.2	15.8
Non-Hispanic white, both sexes.....	1.1	2.4	4.9	8.3	13.3	18.1	21.9	27.2	28.6	26.5
Male.....	1.9	4.2	8.6	14.6	23.3	31.6	38.3	48.9	54.0	50.1
Female.....	0.3	0.5	1.3	2.2	4.0	6.1	8.4	11.3	12.8	16.0
Non-Hispanic black, both sexes.....	1.2	4.1	6.7	12.3	20.6	23.5	27.4	27.7	26.8	21.9
Male.....	1.7	6.5	10.9	20.3	34.1	40.7	44.2	49.9	44.1	39.6
Female.....	*	2.0	3.1	5.7	10.0	10.7	16.0	14.6	17.8	14.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
All origins 3/, both sexes.....	1.4	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
Male.....	1.7	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
Female.....	1.2	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
Hispanic, both sexes.....	2.6	3.6	6.0	10.1	13.9	19.1	31.2	38.8	52.2	62.8
Male.....	2.4	3.8	6.0	13.2	17.2	26.9	38.9	58.7	80.2	84.1
Female.....	2.9	3.4	6.0	7.1	11.0	12.8	25.2	24.5	33.6	51.9
Mexican, both sexes.....	2.6	3.3	5.9	9.7	15.6	22.1	32.8	41.2	48.3	63.2
Male.....	2.4	3.1	5.9	11.4	17.7	32.7	40.0	67.5	*	*
Female.....	2.9	3.5	6.0	8.0	13.6	*	27.5	23.8	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	4.4	7.4	14.4	*	19.8	34.5	*	*	*
Male.....	*	*	*	24.0	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.3	1.9	3.3	4.9	7.5	11.5	17.3	22.4	29.1	41.2
Male.....	1.5	2.4	4.5	7.0	10.7	16.2	24.0	32.0	40.5	61.7
Female.....	1.0	1.4	2.1	3.0	4.6	7.4	11.9	15.5	22.1	32.1
Non-Hispanic white, both sexes.....	0.9	1.4	2.6	3.9	6.2	9.3	14.7	19.6	25.8	37.6
Male.....	1.1	1.8	3.7	5.7	8.8	13.3	20.0	28.0	35.5	56.9
Female.....	0.6	1.0	1.5	2.2	3.8	5.8	10.3	13.5	19.8	28.9
Non-Hispanic black, both sexes.....	2.8	4.0	7.0	10.8	15.8	27.9	34.4	44.2	60.8	71.4
Male.....	3.2	5.4	9.6	15.7	25.2	41.6	56.9	63.9	94.7	99.6
Female.....	2.4	2.8	4.7	6.8	8.5	17.8	19.2	32.6	42.9	60.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes.....	3.9	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
Male.....	4.2	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
Female.....	3.6	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
Hispanic, both sexes.....	2.2	4.9	10.1	16.8	27.3	37.7	60.2	75.8	105.8	147.5
Male.....	2.7	6.2	11.6	20.1	36.0	48.1	74.5	104.8	129.3	172.3
Female.....	1.7	3.6	8.6	13.8	19.7	29.1	49.2	55.2	90.3	134.7
Mexican, both sexes.....	2.1	4.9	8.4	16.0	24.4	38.4	57.1	67.7	99.4	120.8
Male.....	2.3	6.2	10.3	17.5	27.5	49.9	73.3	106.7	*	*
Female.....	*	3.5	6.5	14.4	21.4	27.7	45.0	41.9	87.8	103.6
Puerto Rican, both sexes.....	*	*	15.6	23.4	*	35.3	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	5.2	10.4	15.6	27.0	38.4	56.8	85.3	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.1	7.7	13.8	22.8	36.5	54.3	79.4	113.2	154.4	228.8
Male.....	4.4	8.6	16.5	27.6	44.3	67.6	99.3	140.8	185.5	258.5
Female.....	3.9	6.8	11.3	18.4	29.4	42.6	63.2	93.3	135.3	215.6
Non-Hispanic white, both sexes.....	3.8	7.0	12.9	21.6	34.7	52.1	77.1	111.4	152.8	228.5
Male.....	4.0	8.0	15.4	26.4	42.2	65.1	96.6	137.5	183.8	259.6
Female.....	3.6	6.1	10.4	17.0	27.6	40.6	61.1	92.3	133.6	214.6
Non-Hispanic black, both sexes.....	6.1	12.4	22.4	36.9	58.0	81.4	115.5	150.3	192.9	262.5
Male.....	6.8	13.3	27.1	43.1	71.0	102.7	149.6	201.9	245.3	277.4
Female.....	5.5	11.6	18.5	31.9	47.8	65.5	92.4	119.8	165.4	256.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes.....	1.2	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
Male.....	1.8	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
Female.....	0.6	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
Hispanic, both sexes.....	1.7	5.1	9.9	13.6	20.8	25.3	37.9	46.2	52.5	50.9
Male.....	2.6	7.2	16.5	23.3	33.7	36.1	50.5	58.7	69.8	71.4
Female.....	*	2.8	3.5	4.6	9.6	16.3	28.1	37.2	41.1	40.3
Mexican, both sexes.....	1.9	5.1	8.8	13.6	20.3	30.6	46.0	51.2	57.3	50.0
Male.....	2.8	6.7	14.0	21.7	28.7	41.9	61.6	70.8	*	*
Female.....	*	3.3	*	*	12.1	20.2	34.3	38.3	*	*
Puerto Rican, both sexes.....	*	*	13.6	20.1	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	4.5	9.7	11.5	19.2	26.9	28.7	*	*	*
Male.....	*	*	16.8	22.5	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	3.8	7.3	9.0	11.8	15.9	22.7	27.7	33.6	31.5
Male.....	1.7	6.0	12.3	14.3	17.3	22.6	32.8	37.6	47.1	40.0
Female.....	0.5	1.6	2.5	4.1	6.7	10.1	14.5	20.7	25.3	27.7
Non-Hispanic white, both sexes.....	0.9	3.0	6.2	7.4	10.3	14.1	20.6	25.9	32.5	30.1
Male.....	1.2	4.7	10.5	11.3	14.8	20.1	29.5	36.0	45.7	38.0
Female.....	0.5	1.3	2.1	3.6	6.0	8.8	13.2	18.5	24.3	26.5
Non-Hispanic black, both sexes.....	1.8	7.2	12.8	18.3	16.9	23.7	29.1	34.0	35.7	36.6
Male.....	3.0	12.0	23.0	32.7	27.3	37.9	45.9	39.6	51.4	47.8
Female.....	*	3.2	4.3	6.5	8.7	13.1	17.8	30.6	27.4	32.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
All origins 3/, both sexes.....	1.7	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
Male.....	2.2	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
Female.....	1.3	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
Hispanic, both sexes.....	1.0	2.4	6.4	10.6	19.9	32.4	38.9	54.8	75.4	69.9
Male.....	1.3	3.2	6.8	12.1	25.1	37.0	49.7	58.2	84.9	57.4
Female.....	*	*	5.9	9.2	15.3	28.6	30.7	52.4	69.2	76.3
Mexican, both sexes.....	*	2.4	5.8	10.6	18.6	38.4	41.0	45.6	72.5	52.9
Male.....	*	2.9	5.4	10.6	23.6	39.6	52.5	46.8	*	*
Female.....	*	*	6.2	10.7	13.6	37.3	32.5	44.8	69.5	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	8.1	10.5	22.0	29.4	34.5	90.4	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	4.4	9.0	15.6	26.2	37.6	53.3	69.6	85.7	98.9
Male.....	2.3	5.7	11.4	19.1	31.8	45.7	62.3	78.7	95.1	104.1
Female.....	1.4	3.1	6.7	12.2	21.1	30.6	46.1	63.1	79.9	96.5
Non-Hispanic white, both sexes.....	1.7	4.2	8.5	14.9	25.4	37.0	52.0	69.0	85.3	98.1
Male.....	2.1	5.6	10.9	18.2	31.1	44.9	60.5	78.6	96.1	105.0
Female.....	1.3	2.9	6.2	11.7	20.1	30.0	45.0	61.9	78.7	95.1
Non-Hispanic black, both sexes.....	2.7	6.1	13.6	22.9	37.0	49.0	71.7	85.8	93.9	115.1
Male.....	3.6	7.7	16.4	30.1	45.6	61.8	89.8	90.9	91.8	102.6
Female.....	1.9	4.7	11.2	16.9	30.3	39.4	59.5	82.8	95.0	120.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/, both sexes.....	8.5	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
Male.....	9.1	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
Female.....	7.9	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
Hispanic, both sexes.....	2.1	6.3	13.5	26.5	48.0	82.4	128.6	173.2	187.3	193.1
Male.....	2.6	7.8	19.0	36.6	69.5	122.5	190.8	274.1	288.8	287.2
Female.....	1.5	4.7	8.1	17.0	29.1	49.3	80.5	101.5	120.3	144.8
Mexican, both sexes.....	1.8	4.7	10.3	21.4	41.6	80.3	127.7	171.9	180.1	202.0
Male.....	*	5.2	15.5	28.3	56.6	115.8	195.7	280.9	265.6	*
Female.....	*	4.3	5.0	14.4	26.8	47.4	76.8	99.7	114.6	140.1
Puerto Rican, both sexes.....	*	*	21.9	38.3	63.3	67.8	150.8	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	60.7	78.2	98.3	171.9	155.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	6.9	11.8	23.9	43.4	88.3	97.7	170.7	*	*
Male.....	*	8.2	*	39.8	67.5	*	*	*	*	*
Female.....	*	*	11.1	*	26.5	58.6	61.1	*	*	*
Non-Hispanic 5/, both sexes.....	9.4	22.2	43.0	82.1	152.2	237.2	323.3	383.7	384.0	301.6
Male.....	10.1	25.3	52.8	100.5	187.7	297.3	414.2	521.4	542.1	474.8
Female.....	8.7	19.2	33.7	64.7	119.9	184.6	250.0	284.8	287.0	224.2
Non-Hispanic white, both sexes.....	9.1	21.5	41.5	80.5	153.8	241.0	331.6	390.5	388.4	306.3
Male.....	9.6	23.6	49.3	96.2	184.0	297.2	416.1	522.4	542.5	480.3
Female.....	8.7	19.3	33.8	65.5	125.5	190.9	261.8	293.9	293.0	228.4
Non-Hispanic black, both sexes.....	13.1	31.8	64.2	116.3	182.5	262.0	314.7	387.9	393.4	281.5
Male.....	15.0	42.1	91.5	164.8	276.5	375.6	476.8	609.0	636.7	488.8
Female.....	11.5	22.9	41.3	76.3	109.4	177.8	204.9	257.0	265.4	199.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

All origins 3/, both sexes.....	7.3	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
Male.....	*	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
Female.....	14.5	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
Hispanic, both sexes.....	4.7	7.3	13.2	18.8	22.0	29.1	29.8	38.8	45.0	72.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	9.7	14.6	25.9	36.1	41.1	53.1	50.8	66.3	73.5	108.8
Mexican, both sexes.....	4.1	5.9	13.6	16.9	20.1	28.2	30.0	29.1	32.4	63.2
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	8.5	12.3	27.3	33.7	39.7	54.3	51.8	48.4	*	97.8
Puerto Rican, both sexes.....	*	*	14.6	26.0	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.7	8.5	12.1	20.7	24.2	34.6	22.3	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	12.1	16.1	22.2	35.5	41.3	54.6	*	*	*	*
Non-Hispanic 5/, both sexes.....	7.7	12.2	19.2	26.7	33.9	41.7	50.9	66.1	84.1	126.3
Male.....	*	*	*	0.4	0.6	1.0	1.2	2.2	2.1	3.3
Female.....	15.1	23.9	37.3	51.6	64.2	77.2	91.0	112.1	134.5	181.3
Non-Hispanic white, both sexes.....	6.7	10.9	17.5	25.6	32.8	41.5	50.8	66.7	85.3	126.8
Male.....	*	*	*	0.4	0.6	1.0	1.1	2.3	2.0	2.9
Female.....	13.3	21.6	34.4	49.9	62.7	77.6	91.9	113.8	136.8	182.4
Non-Hispanic black, both sexes.....	14.0	21.7	32.7	41.0	49.7	52.1	63.9	77.3	91.4	145.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	26.3	40.5	59.7	74.3	87.2	89.7	105.5	122.0	137.5	199.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes.....	1.7	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
Male.....
Female.....	3.4	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
Hispanic, both sexes.....	2.0	2.2	2.6	3.3	5.6	3.9	4.7	4.8	*	*
Male.....
Female.....	4.2	4.5	5.1	6.4	10.6	7.1	8.3	8.2	*	*
Mexican, both sexes.....	2.0	2.5	2.8	3.6	6.6	*	*	*	*	*
Male.....
Female.....	4.2	5.4	5.6	7.2	13.2	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.6	2.0	2.0	2.6	2.7	3.1	3.2	3.2	4.2	5.0
Male.....
Female.....	3.2	4.0	3.9	5.1	5.2	5.9	5.8	5.4	6.8	7.2
Non-Hispanic white, both sexes.....	1.4	1.8	1.7	2.4	2.3	2.6	2.6	2.7	3.6	3.9
Male.....
Female.....	2.9	3.6	3.3	4.7	4.5	4.9	4.8	4.7	5.8	5.7
Non-Hispanic black, both sexes.....	2.9	3.6	4.0	4.8	6.3	7.3	8.2	8.1	10.4	16.4
Male.....
Female.....	5.4	6.8	7.4	8.7	11.2	12.8	13.7	13.0	15.8	22.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins 3/, both sexes.....	0.5	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
Male.....
Female.....	1.0	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
Hispanic, both sexes.....	*	*	*	2.3	5.2	8.9	6.5	11.0	9.8	9.5
Male.....
Female.....	*	*	*	4.5	9.8	16.3	11.5	18.8	16.2	14.4
Mexican, both sexes.....	*	*	*	*	*	8.8	*	10.9	*	*
Male.....
Female.....	*	*	*	*	*	17.0	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.9	1.9	3.5	5.8	8.8	11.8	14.0	18.1	22.8
Male.....
Female.....	1.0	1.8	3.7	6.8	11.1	16.4	21.4	24.0	29.3	33.0
Non-Hispanic white, both sexes.....	0.5	0.9	1.7	3.3	5.1	7.5	10.5	13.1	17.8	23.0
Male.....
Female.....	1.0	1.9	3.3	6.5	9.9	14.3	19.1	22.7	28.8	33.3
Non-Hispanic black, both sexes.....	0.8	1.0	3.1	5.2	12.1	20.6	27.0	27.1	25.8	26.5
Male.....
Female.....	1.5	1.9	5.7	9.5	21.6	35.9	45.3	43.1	39.4	36.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

All origins 3/, both sexes.....	1.4	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
Male.....
Female.....	2.8	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
Hispanic, both sexes.....	1.1	2.5	3.1	5.5	7.9	11.2	15.1	19.4	23.3	28.1
Male.....
Female.....	2.3	5.1	6.1	10.7	14.8	20.4	26.8	33.1	38.6	42.5
Mexican, both sexes.....	*	2.7	3.0	6.8	7.8	10.5	19.3	16.9	24.8	*
Male.....
Female.....	*	5.8	6.0	13.6	15.6	20.2	33.7	28.2	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	3.0	5.4	8.1	12.2	18.3	23.0	29.5	37.0	38.1
Male.....
Female.....	2.9	5.9	10.5	15.7	23.2	34.3	41.6	50.6	59.8	55.1
Non-Hispanic white, both sexes.....	1.5	3.0	5.5	8.2	12.7	18.8	23.6	30.4	38.5	39.1
Male.....
Female.....	3.0	6.0	10.8	16.2	24.5	35.5	43.1	52.6	62.4	56.5
Non-Hispanic black, both sexes.....	1.1	3.1	5.1	7.6	10.4	18.7	22.8	26.9	29.6	32.5
Male.....
Female.....	2.0	5.9	9.4	13.9	18.6	32.6	38.3	42.8	45.2	45.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins 3/, both sexes.....	0.8	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
Male.....	1.2	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
Female.....	0.5	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
Hispanic, both sexes.....	0.9	2.3	4.2	6.1	11.2	13.3	15.0	19.4	29.6	28.1
Male.....	*	3.4	5.9	8.3	15.4	23.8	22.8	29.4	43.4	43.4
Female.....	*	*	2.5	4.0	7.5	*	8.9	12.3	20.6	20.2
Mexican, both sexes.....	*	3.0	3.8	8.1	13.1	17.4	18.2	22.6	35.2	35.9
Male.....	*	4.2	5.5	10.3	17.3	28.1	*	*	*	*
Female.....	*	*	*	5.9	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.8	2.1	4.0	6.8	11.3	14.6	19.4	24.8	29.7	32.9
Male.....	1.2	2.9	5.9	10.0	16.7	20.4	28.1	34.5	44.0	47.1
Female.....	0.5	1.3	2.2	3.8	6.3	9.6	12.3	17.9	21.0	26.5
Non-Hispanic white, both sexes.....	0.8	2.2	4.1	7.1	11.7	15.1	19.7	25.2	30.5	33.8
Male.....	1.1	3.0	6.1	10.3	17.3	20.8	28.6	34.8	45.4	48.6
Female.....	0.5	1.3	2.2	3.9	6.5	10.1	12.4	18.1	21.3	27.2
Non-Hispanic black, both sexes.....	1.2	2.4	4.1	6.4	10.7	13.3	19.5	26.9	25.6	28.8
Male.....	1.7	3.3	6.3	9.5	16.4	19.7	29.9	39.2	36.7	39.6
Female.....	*	1.6	2.3	3.8	6.3	8.5	12.4	19.6	19.7	24.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

All origins 3/, both sexes.....	0.4	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
Male.....	0.5	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
Female.....	0.3	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
Hispanic, both sexes.....	*	*	*	2.6	2.9	6.3	8.7	16.0	28.1	32.3
Male.....	*	*	*	3.8	4.9	9.0	13.7	21.9	39.6	58.8
Female.....	*	*	*	*	*	*	*	11.9	20.6	18.7
Mexican, both sexes.....	*	*	*	*	*	*	8.6	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.4	1.0	1.8	3.7	7.0	11.9	19.6	31.9	46.5	69.4
Male.....	0.5	1.4	2.9	5.8	11.2	19.2	31.7	54.9	83.7	136.5
Female.....	0.3	0.6	0.8	1.7	3.2	5.4	9.8	15.4	23.7	39.4
Non-Hispanic white, both sexes.....	0.4	1.0	1.9	3.9	7.6	12.7	20.6	33.6	48.5	71.1
Male.....	0.5	1.4	3.0	6.2	12.0	21.0	33.6	57.9	88.5	143.1
Female.....	0.3	0.5	0.8	1.7	3.5	5.4	9.8	15.8	23.8	38.9
Non-Hispanic black, both sexes.....	*	1.2	2.1	3.4	5.1	8.1	16.7	23.1	34.2	61.3
Male.....	*	1.6	3.1	4.5	8.1	8.5	23.6	35.4	47.7	83.3
Female.....	*	*	*	2.6	*	7.8	11.9	15.7	27.0	52.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes.....	2.4	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
Male.....	2.9	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
Female.....	1.9	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
Hispanic, both sexes.....	1.3	1.7	3.3	4.9	7.3	8.8	9.2	12.0	13.1	14.3
Male.....	1.5	1.9	4.3	5.4	9.8	9.9	10.3	16.7	*	*
Female.....	*	*	2.3	4.5	5.1	7.9	8.3	8.6	13.1	15.1
Mexican, both sexes.....	1.1	*	2.4	4.4	7.6	9.1	8.2	*	*	*
Male.....	*	*	*	*	9.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	5.1	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.5	4.0	5.7	8.6	11.2	14.3	18.2	19.4	20.8	16.7
Male.....	3.1	5.1	7.3	10.8	14.0	16.4	22.1	23.5	25.1	19.7
Female.....	2.0	3.0	4.1	6.6	8.6	12.5	15.1	16.4	18.1	15.4
Non-Hispanic white, both sexes.....	2.8	4.4	6.2	9.5	12.2	15.7	19.8	20.6	22.1	17.4
Male.....	3.3	5.5	7.9	11.7	15.2	18.0	24.0	24.6	26.4	21.0
Female.....	2.3	3.3	4.6	7.4	9.5	13.6	16.3	17.7	19.4	15.8
Non-Hispanic black, both sexes.....	1.8	2.3	3.6	4.2	6.1	7.1	9.0	10.1	10.1	10.6
Male.....	2.8	3.0	5.2	6.1	8.1	8.0	8.3	14.5	14.7	*
Female.....	*	1.6	2.2	2.7	4.5	6.5	9.5	7.5	7.7	11.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes.....	4.1	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
Male.....	5.0	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
Female.....	3.2	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
Hispanic, both sexes.....	4.0	6.0	10.0	16.5	24.6	39.4	59.7	85.6	110.7	103.2
Male.....	5.8	7.1	13.7	18.2	32.3	49.7	75.8	107.7	132.1	119.1
Female.....	2.0	5.0	6.4	14.9	17.9	30.9	47.3	70.0	96.6	95.1
Mexican, both sexes.....	4.0	5.3	9.1	15.0	23.7	43.1	63.9	81.6	115.3	101.9
Male.....	5.6	5.6	13.0	15.6	30.3	51.0	85.8	107.8	*	*
Female.....	*	4.9	5.2	14.4	17.1	35.7	47.5	64.3	106.1	99.2
Puerto Rican, both sexes.....	*	*	12.2	23.4	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.7	7.4	10.2	16.0	25.2	41.6	58.9	101.9	*	*
Male.....	*	9.3	13.7	22.5	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.1	6.7	11.7	20.4	33.7	55.6	85.8	124.8	164.9	193.0
Male.....	4.9	8.2	14.4	25.4	42.4	71.5	110.9	161.6	218.6	261.1
Female.....	3.3	5.4	9.2	15.7	25.8	41.6	65.6	98.3	132.0	162.6
Non-Hispanic white, both sexes.....	4.0	6.5	11.3	20.1	33.8	56.3	86.7	127.6	169.7	198.8
Male.....	4.8	7.8	13.8	25.3	42.6	72.0	112.0	165.5	225.5	270.9
Female.....	3.2	5.2	8.9	15.1	25.6	42.2	65.7	99.9	135.1	166.5
Non-Hispanic black, both sexes.....	5.4	9.3	17.0	26.9	41.2	61.4	94.4	117.4	136.7	156.3
Male.....	6.2	11.5	21.5	32.2	51.8	81.1	124.0	140.8	171.8	197.1
Female.....	4.7	7.4	13.2	22.5	33.0	46.8	74.4	103.5	118.2	140.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
All origins 3/, both sexes.....	1.5	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
Male.....	2.1	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
Female.....	0.9	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
Hispanic, both sexes.....	1.8	2.7	4.0	6.8	8.5	15.1	23.4	35.6	48.4	43.8
Male.....	2.9	3.8	6.4	7.6	10.5	18.5	31.5	44.3	51.9	50.4
Female.....	*	*	*	6.1	6.7	12.3	17.2	29.5	46.1	40.3
Mexican, both sexes.....	1.5	2.1	4.0	6.6	7.6	16.0	25.7	36.9	53.1	44.4
Male.....	2.4	*	6.1	7.4	*	16.6	36.6	44.6	*	*
Female.....	*	*	*	5.9	*	15.4	17.5	31.8	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	22.3	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	2.8	4.8	7.8	12.9	21.5	32.5	47.5	65.5	76.3
Male.....	2.0	3.7	6.0	10.0	16.1	27.2	40.2	58.3	83.0	97.8
Female.....	1.0	1.9	3.7	5.8	10.1	16.5	26.3	39.8	54.8	66.7
Non-Hispanic white, both sexes.....	1.4	2.7	4.8	8.0	13.5	22.7	34.2	50.0	69.5	80.2
Male.....	1.9	3.5	6.0	10.1	16.6	28.4	42.0	61.0	87.9	103.5
Female.....	0.9	1.8	3.6	6.0	10.5	17.7	27.8	41.9	58.1	69.8
Non-Hispanic black, both sexes.....	1.8	4.0	5.9	7.7	11.0	16.0	21.5	29.3	26.6	39.4
Male.....	2.3	5.6	7.8	10.2	14.4	20.0	28.3	32.2	35.2	42.7
Female.....	*	2.5	4.4	5.7	8.4	13.1	16.9	27.6	22.0	38.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

All origins 3/, both sexes.....	1.8	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
Male.....	2.1	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
Female.....	1.6	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
Hispanic, both sexes.....	1.7	2.2	4.0	5.1	7.9	13.6	18.4	25.1	34.2	36.2
Male.....	2.2	2.1	5.0	6.1	11.2	17.9	23.2	34.0	44.4	43.4
Female.....	*	2.3	2.9	4.2	5.1	10.0	14.7	18.8	27.4	32.4
Mexican, both sexes.....	1.9	2.0	3.5	4.3	6.8	14.6	17.5	22.1	31.7	34.9
Male.....	2.3	*	5.0	*	9.8	18.3	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.9	2.5	4.2	7.3	12.3	20.4	32.7	46.8	62.8	80.6
Male.....	2.1	2.9	5.0	9.2	16.0	27.4	45.3	65.8	88.7	113.9
Female.....	1.7	2.2	3.4	5.5	8.9	14.3	22.6	33.1	46.9	65.7
Non-Hispanic white, both sexes.....	1.8	2.6	4.1	7.4	12.5	21.0	33.5	48.5	65.0	83.6
Male.....	2.0	2.9	4.8	9.5	16.4	27.9	45.9	68.1	92.2	118.8
Female.....	1.7	2.3	3.5	5.4	8.9	14.8	23.2	34.0	48.2	67.8
Non-Hispanic black, both sexes.....	2.3	2.9	5.3	7.9	14.0	19.5	32.9	37.5	47.8	60.7
Male.....	2.4	3.0	7.1	9.5	17.5	28.4	49.2	48.5	59.5	81.3
Female.....	2.1	2.7	3.8	6.5	11.2	12.9	21.9	30.9	41.7	52.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All origins 3/, both sexes.....	0.5	1.0	2.3	4.6	7.7	12.3	18.9	28.3	33.9	33.1
Male.....	0.5	1.1	2.7	5.2	9.3	15.3	23.2	35.0	42.8	44.6
Female.....	0.4	0.9	1.8	4.0	6.2	9.8	15.5	23.5	28.4	28.0
Hispanic, both sexes.....	*	*	1.9	4.2	7.4	9.5	16.2	22.5	26.3	18.6
Male.....	*	*	*	4.1	9.8	11.1	18.2	27.1	34.9	*
Female.....	*	*	*	4.3	5.3	8.2	14.7	19.2	20.6	18.7
Mexican, both sexes.....	*	*	*	4.1	8.4	10.5	19.3	20.4	29.0	*
Male.....	*	*	*	*	11.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	1.0	2.3	4.6	7.7	12.6	19.1	28.6	34.1	33.7
Male.....	0.5	1.2	2.8	5.3	9.2	15.6	23.5	35.4	43.2	45.9
Female.....	0.4	0.9	1.8	3.9	6.3	9.9	15.6	23.7	28.6	28.3
Non-Hispanic white, both sexes.....	0.4	0.9	1.9	3.9	7.0	11.4	17.4	27.2	32.6	32.6
Male.....	0.5	1.0	2.3	4.7	8.5	14.4	22.0	34.1	41.5	44.8
Female.....	0.3	0.8	1.5	3.2	5.7	8.8	13.7	22.1	27.1	27.1
Non-Hispanic black, both sexes.....	1.1	2.1	5.2	10.5	15.5	24.7	39.0	49.4	60.5	54.7
Male.....	*	2.2	6.1	11.6	18.7	30.9	45.2	58.8	75.6	72.1
Female.....	*	2.0	4.4	9.6	13.0	20.2	34.7	43.9	52.5	47.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Table with 11 columns representing age groups (40-44 to 85+ years) and 11 rows representing population groups (All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white/black). All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes.....	5.3	9.6	16.2	27.6	43.3	64.6	91.1	126.4	165.2	213.1
Male.....	5.5	10.6	18.8	32.5	51.4	78.8	109.3	157.8	207.2	260.6
Female.....	5.1	8.6	13.7	23.0	35.9	52.3	76.5	103.8	139.3	191.8
Hispanic, both sexes.....	3.6	6.4	12.0	17.8	29.5	43.6	64.4	93.5	126.9	140.8
Male.....	3.9	6.0	13.8	19.7	33.5	46.9	76.2	115.8	156.7	147.1
Female.....	3.2	6.9	10.3	15.9	26.0	40.9	55.3	77.7	107.2	137.6
Mexican, both sexes.....	3.7	6.5	11.5	16.5	30.3	46.9	64.6	99.0	122.2	125.5
Male.....	3.8	5.7	12.8	16.4	33.0	48.7	75.8	132.8	141.6	*
Female.....	3.5	7.3	10.2	16.6	27.6	45.3	56.2	76.5	107.3	125.5
Puerto Rican, both sexes.....	*	*	12.2	26.6	37.8	36.2	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.7	5.9	11.1	16.6	28.4	42.2	62.5	103.2	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	15.0	25.0	*	50.0	*	*	*
Non-Hispanic 5/, both sexes.....	5.6	9.9	16.6	28.3	44.2	66.2	92.8	128.1	166.8	215.9
Male.....	5.8	11.1	19.2	33.4	52.6	81.0	111.2	160.1	209.6	265.7
Female.....	5.4	8.8	14.1	23.6	36.6	53.2	77.8	105.2	140.6	193.7
Non-Hispanic white, both sexes.....	5.4	9.3	15.7	27.3	43.5	65.5	93.9	129.6	168.7	220.6
Male.....	5.5	10.3	17.9	31.8	51.3	79.8	111.9	162.3	213.2	272.4
Female.....	5.3	8.3	13.5	23.0	36.2	52.8	79.0	105.7	141.2	197.4
Non-Hispanic black, both sexes.....	7.7	15.0	25.6	41.4	58.4	85.4	100.2	134.7	170.1	188.2
Male.....	8.5	17.6	33.5	53.4	75.5	109.0	129.7	172.1	207.1	223.5
Female.....	7.0	12.8	18.9	31.6	45.1	67.9	80.2	112.6	150.7	174.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes.....	0.8	1.4	2.0	3.5	5.8	9.3	16.7	27.3	47.2	77.7
Male.....	0.9	1.4	2.3	4.0	7.1	11.7	20.5	34.8	61.3	98.9
Female.....	0.7	1.3	1.7	3.0	4.6	7.1	13.7	21.9	38.5	68.1
Hispanic, both sexes.....	0.7	1.3	1.4	2.4	5.8	5.0	13.3	17.9	31.9	48.5
Male.....	*	*	*	*	6.3	*	16.6	21.9	37.7	57.4
Female.....	*	*	*	*	5.3	5.4	10.9	15.1	28.0	43.9
Mexican, both sexes.....	*	1.6	*	*	5.3	*	9.6	18.7	28.3	49.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.8	1.4	2.1	3.6	5.8	9.5	16.9	27.8	47.9	78.9
Male.....	0.9	1.5	2.4	4.1	7.2	12.2	20.7	35.4	62.5	101.0
Female.....	0.7	1.4	1.7	3.1	4.6	7.2	13.8	22.2	38.9	69.0
Non-Hispanic white, both sexes.....	0.7	1.4	1.9	3.4	5.7	9.7	17.3	28.3	49.3	81.6
Male.....	0.9	1.4	2.3	3.9	7.0	12.6	21.3	36.1	64.5	105.6
Female.....	0.6	1.3	1.6	2.9	4.5	7.1	14.0	22.5	39.9	70.9
Non-Hispanic black, both sexes.....	1.3	1.7	3.1	5.7	7.2	10.4	16.4	26.5	40.0	57.9
Male.....	*	1.7	3.6	6.8	9.6	12.7	19.6	35.0	47.7	62.0
Female.....	1.4	1.6	2.6	4.7	5.2	8.7	14.2	21.5	35.9	56.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

All origins 3/, both sexes.....	0.5	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
Male.....	0.5	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
Female.....	0.5	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	9.4	28.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	30.3
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.6	0.6	0.9	1.4	1.8	3.6	6.0	12.3	38.5
Male.....	0.5	0.5	0.6	1.0	1.5	2.0	3.8	6.2	13.0	37.1
Female.....	0.6	0.6	0.6	0.7	1.3	1.7	3.4	5.8	12.0	39.1
Non-Hispanic white, both sexes.....	0.2	0.3	0.3	0.5	1.1	1.7	3.3	5.7	12.4	38.1
Male.....	*	0.3	0.3	0.5	1.2	1.9	3.6	5.7	12.8	37.3
Female.....	*	*	*	0.5	1.0	1.5	3.1	5.6	12.1	38.4
Non-Hispanic black, both sexes.....	2.5	2.5	3.1	3.9	4.2	3.6	6.3	10.8	13.4	48.9
Male.....	2.6	2.1	2.9	5.3	4.4	*	*	13.5	14.7	43.7
Female.....	2.5	2.9	3.2	2.8	4.0	*	6.3	9.1	12.7	51.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

All origins 3/, both sexes.....	5.7	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
Male.....	7.1	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
Female.....	4.3	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
Hispanic, both sexes.....	4.2	10.3	17.2	36.7	64.7	102.6	153.8	229.6	303.6	365.8
Male.....	4.9	11.4	22.1	41.6	79.3	110.2	166.4	249.4	328.4	325.0
Female.....	3.5	9.2	12.4	32.1	51.9	96.3	144.1	215.6	287.2	386.8
Mexican, both sexes.....	4.5	12.7	18.6	40.8	77.0	134.2	202.0	263.1	365.8	407.7
Male.....	5.5	13.6	23.9	45.0	87.7	133.0	219.0	297.3	375.4	*
Female.....	3.5	11.7	13.3	36.6	66.5	135.3	189.2	240.5	358.4	427.5
Puerto Rican, both sexes.....	*	*	19.5	43.5	77.4	95.6	147.7	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	117.0	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.9	6.0	12.1	29.7	43.9	67.2	99.9	211.4	*	*
Male.....	*	*	15.2	36.0	67.5	*	*	*	*	*
Female.....	*	*	*	25.0	27.3	58.6	83.3	*	*	*
Non-Hispanic 5/, both sexes.....	5.9	10.0	17.1	28.6	45.6	68.8	102.1	144.6	208.1	303.5
Male.....	7.4	12.5	21.0	33.7	54.7	80.5	117.8	167.0	242.3	333.5
Female.....	4.4	7.5	13.4	23.8	37.4	58.5	89.4	128.5	187.2	290.1
Non-Hispanic white, both sexes.....	5.1	8.6	14.2	24.1	39.0	59.7	89.7	130.7	193.1	284.5
Male.....	6.5	10.9	18.1	28.9	48.3	71.6	106.9	155.5	232.7	321.7
Female.....	3.7	6.4	10.3	19.6	30.4	49.1	75.5	112.5	168.6	267.8
Non-Hispanic black, both sexes.....	11.8	19.8	39.7	66.1	103.5	151.1	222.3	300.5	388.9	536.5
Male.....	14.4	24.9	45.2	76.4	117.4	172.6	241.1	312.9	393.6	508.1
Female.....	9.6	15.3	35.0	57.5	92.7	135.2	209.6	293.2	386.4	547.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

All origins 3/, both sexes.....	0.1	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.0	30.3
Male.....	*	*	0.4	0.5	0.8	1.3	2.2	4.9	9.6	27.4
Female.....	*	0.2	0.4	0.4	0.7	1.1	2.0	4.3	10.3	31.7
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	19.5
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	22.3
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.2	30.8
Male.....	*	*	0.4	0.5	0.8	1.3	2.3	5.1	9.7	28.0
Female.....	*	0.2	0.5	0.4	0.7	1.2	2.0	4.3	10.5	32.1
Non-Hispanic white, both sexes.....	*	0.2	0.4	0.4	0.6	1.0	2.0	4.3	9.9	29.6
Male.....	*	*	0.3	0.4	0.7	0.9	2.2	4.6	9.3	26.9
Female.....	*	*	0.4	0.4	0.6	1.2	1.8	4.0	10.2	30.8
Non-Hispanic black, both sexes.....	*	*	0.9	*	1.8	2.8	4.3	8.3	14.2	47.2
Male.....	*	*	*	*	*	*	*	10.7	14.7	45.7
Female.....	*	*	*	*	*	*	*	6.9	13.9	47.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
All origins 3/, both sexes.....	0.1	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
Male.....	*	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
Female.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	19.5
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	22.3
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	0.2	0.4	0.4	0.7	1.2	2.1	4.5	9.8	28.6
Male.....	*	*	0.4	0.5	0.7	1.2	2.3	4.9	9.2	25.8
Female.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.1	10.2	29.9
Non-Hispanic white, both sexes.....	*	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.5	27.3
Male.....	*	*	0.3	0.4	0.6	0.9	2.1	4.4	8.8	24.6
Female.....	*	*	0.4	0.4	0.6	1.1	1.7	3.8	9.9	28.5
Non-Hispanic black, both sexes.....	*	*	0.9	*	*	2.5	4.3	8.3	13.7	46.3
Male.....	*	*	*	*	*	*	*	10.7	*	43.7
Female.....	*	*	*	*	*	*	*	6.9	13.5	47.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

All origins 3/, both sexes.....	*	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
Male.....	*	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
Female.....	*	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
Hispanic, both sexes.....	*	*	*	*	*	3.5	11.2	31.6	63.8	85.1
Male.....	*	*	*	*	*	*	15.7	49.0	95.3	126.1
Female.....	*	*	*	*	*	*	7.7	19.2	43.0	64.1
Mexican, both sexes.....	*	*	*	*	*	*	10.7	29.5	58.7	76.5
Male.....	*	*	*	*	*	*	*	44.6	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	*	0.3	0.6	2.0	6.6	19.0	48.4	95.4	137.9
Male.....	*	*	0.4	0.9	2.6	9.4	28.1	71.8	149.5	224.4
Female.....	*	*	*	0.4	1.4	4.3	11.7	31.6	62.2	99.2
Non-Hispanic white, both sexes.....	*	*	0.3	0.7	2.1	7.1	20.8	51.6	101.5	145.5
Male.....	*	*	0.4	1.0	2.7	10.0	30.3	76.0	158.9	236.3
Female.....	*	*	*	0.4	1.6	4.5	12.9	33.8	66.0	104.8
Non-Hispanic black, both sexes.....	*	*	*	*	*	3.9	7.4	22.9	40.0	61.6
Male.....	*	*	*	*	*	5.2	13.0	35.0	59.5	106.7
Female.....	*	*	*	*	*	*	*	15.7	29.7	43.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

All origins 3/, both sexes.....	*	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
Male.....	*	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
Female.....	*	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
Hispanic, both sexes.....	*	*	*	*	*	6.0	19.8	62.0	158.4	490.9
Male.....	*	*	*	*	*	*	20.7	57.6	151.0	374.1
Female.....	*	*	*	*	*	7.1	19.2	65.0	163.3	551.0
Mexican, both sexes.....	*	*	*	*	*	*	17.1	53.0	142.2	408.7
Male.....	*	*	*	*	*	*	*	46.8	*	*
Female.....	*	*	*	*	*	*	*	57.0	151.2	447.9
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	0.2	0.3	1.0	3.2	10.4	31.7	101.5	265.9	832.6
Male.....	*	*	0.3	0.9	3.2	9.7	30.2	91.1	240.5	631.2
Female.....	*	0.3	0.4	1.0	3.2	11.0	32.9	109.0	281.5	922.7
Non-Hispanic white, both sexes.....	*	0.2	0.4	1.0	3.4	10.8	32.8	104.6	273.8	861.9
Male.....	*	*	0.4	1.0	3.4	10.1	30.8	93.5	248.6	653.3
Female.....	*	0.3	0.4	1.1	3.4	11.4	34.4	112.8	289.4	955.3
Non-Hispanic black, both sexes.....	*	*	*	*	2.3	10.4	30.8	97.6	239.7	666.9
Male.....	*	*	*	*	*	9.5	32.6	91.4	213.0	507.0
Female.....	*	*	*	*	*	11.1	29.5	101.3	253.8	730.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes.....	47.8	86.0	139.2	211.1	343.0	537.5	885.1	1,514.3	2,730.4	6,616.6
Male.....	64.6	120.2	198.5	296.2	472.0	714.7	1,134.9	1,900.8	3,283.3	6,818.1
Female.....	31.2	52.6	82.4	130.8	225.5	383.3	684.0	1,236.7	2,389.8	6,526.0
Hispanic, both sexes.....	29.1	53.3	95.0	155.1	268.8	429.5	718.3	1,223.0	2,109.2	4,583.7
Male.....	40.4	76.0	134.7	218.3	375.6	580.2	932.3	1,540.6	2,563.2	4,600.7
Female.....	16.8	29.9	56.0	96.1	175.3	304.9	553.3	997.4	1,809.5	4,575.0
Mexican, both sexes.....	28.1	52.6	91.5	159.9	264.2	471.0	775.8	1,195.7	2,042.1	4,228.4
Male.....	39.7	73.3	127.5	216.4	343.7	582.5	1,005.1	1,540.7	2,307.9	*
Female.....	15.4	29.2	55.4	102.9	185.5	367.5	603.8	966.9	1,838.4	4,121.7
Puerto Rican, both sexes.....	41.0	79.3	111.4	204.5	394.8	422.3	878.2	*	*	*
Male.....	66.9	118.4	150.5	275.7	*	*	*	*	*	*
Female.....	20.9	46.9	69.0	133.6	248.6	302.4	*	*	*	*
Cuban, both sexes.....	33.7	58.3	152.8	128.7	231.0	347.5	708.6	1,077.2	1,620.3	*
Male.....	49.7	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	26.3	44.3	86.8	126.0	233.3	385.3	534.6	1,220.1	*	*
Male.....	32.3	66.8	131.6	198.8	369.5	630.1	*	*	*	*
Female.....	19.6	24.3	49.1	72.2	137.2	243.6	383.3	*	*	*
Non-Hispanic 5/, both sexes.....	50.3	89.4	143.0	215.1	347.6	544.1	894.0	1,528.3	2,756.4	6,698.1
Male.....	68.0	125.0	204.0	301.7	477.5	722.1	1,144.7	1,917.5	3,314.1	6,917.7
Female.....	33.0	54.9	84.7	133.4	229.0	388.6	691.5	1,248.5	2,414.0	6,600.0
Non-Hispanic white, both sexes.....	43.7	77.2	124.4	191.6	318.5	507.6	857.0	1,491.4	2,746.7	6,790.0
Male.....	61.8	111.4	182.8	273.4	442.3	682.0	1,108.2	1,883.8	3,329.1	7,045.5
Female.....	25.6	43.4	67.2	112.7	203.2	352.1	649.7	1,204.0	2,386.2	6,675.5
Non-Hispanic black, both sexes.....	100.4	183.9	299.5	443.5	661.9	937.9	1,397.2	2,160.8	3,256.7	6,436.5
Male.....	122.0	240.4	400.9	606.2	906.3	1,228.5	1,753.7	2,659.1	3,720.1	6,280.6
Female.....	81.5	134.7	213.9	309.8	471.7	722.3	1,155.5	1,865.8	3,012.8	6,498.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All origins 3/, both sexes.....	38.3	69.6	113.7	173.4	278.1	423.7	679.6	1,130.0	2,007.8	4,895.9
Male.....	54.2	101.5	167.5	252.3	393.5	580.0	899.4	1,472.2	2,509.6	5,239.3
Female.....	22.7	38.5	62.2	99.1	173.2	287.7	502.6	884.1	1,698.7	4,741.5
Hispanic, both sexes.....	21.1	39.0	70.0	116.8	210.6	330.1	545.8	918.3	1,584.5	3,446.8
Male.....	31.3	58.6	103.1	170.7	303.3	452.8	732.3	1,199.7	1,963.0	3,496.8
Female.....	10.0	18.7	37.5	66.5	129.5	228.5	401.8	718.4	1,334.7	3,421.2
Mexican, both sexes.....	20.1	38.5	67.1	119.8	204.1	353.7	581.3	867.0	1,483.1	3,026.9
Male.....	29.9	56.5	97.3	169.9	274.5	449.5	785.3	1,165.1	1,705.1	*
Female.....	9.4	18.2	36.7	69.2	134.6	264.7	428.3	669.4	1,313.0	2,912.2
Puerto Rican, both sexes.....	32.3	57.7	80.3	156.5	314.5	341.6	692.8	*	*	*
Male.....	58.8	89.6	108.5	213.3	*	*	*	*	*	*
Female.....	*	31.3	49.7	99.9	190.5	242.9	*	*	*	*
Cuban, both sexes.....	29.1	42.3	127.9	113.9	192.5	278.2	577.2	867.5	1,336.5	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	17.8	32.4	63.3	90.9	182.5	297.0	390.9	917.0	*	*
Male.....	23.7	52.9	101.6	149.3	298.7	493.9	*	*	*	*
Female.....	11.2	14.2	31.2	47.8	100.6	183.0	266.6	*	*	*
Non-Hispanic 5/, both sexes.....	40.6	72.8	117.5	177.6	282.4	429.5	686.6	1,139.9	2,025.0	4,953.4
Male.....	57.4	106.2	173.1	258.2	398.7	587.2	907.3	1,484.5	2,532.5	5,317.0
Female.....	24.3	40.5	64.4	101.5	176.3	291.7	508.4	892.1	1,713.5	4,790.8
Non-Hispanic white, both sexes.....	36.4	65.0	105.4	161.8	264.2	406.0	663.9	1,120.6	2,026.9	5,038.3
Male.....	53.5	97.5	159.5	239.2	376.9	561.4	886.3	1,469.9	2,556.1	5,430.9
Female.....	19.3	32.9	52.4	87.2	159.1	267.3	480.4	864.7	1,699.3	4,862.4
Non-Hispanic black, both sexes.....	76.0	138.1	230.5	344.2	500.5	707.1	1,036.2	1,553.5	2,324.0	4,610.3
Male.....	96.2	187.0	316.5	480.8	697.0	945.7	1,332.0	1,963.8	2,731.7	4,695.5
Female.....	58.2	95.5	157.9	232.0	347.5	530.1	835.7	1,310.7	2,109.5	4,576.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins 3/, both sexes.....	0.3	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
Male.....	0.2	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
Female.....	0.3	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
Hispanic, both sexes.....	*	*	*	*	2.2	3.4	3.6	5.0	8.3	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.4	0.6	0.9	1.3	2.5	4.2	7.4	11.1	17.7
Male.....	0.2	0.4	0.5	0.7	0.9	2.0	2.8	5.7	9.6	11.9
Female.....	0.3	0.4	0.7	1.1	1.7	3.0	5.4	8.5	11.9	20.2
Non-Hispanic white, both sexes.....	0.2	0.3	0.6	0.8	1.3	2.5	4.4	7.7	11.7	18.9
Male.....	*	0.3	0.4	0.6	0.9	2.0	3.0	6.1	10.3	12.8
Female.....	*	0.3	0.7	0.9	1.8	2.9	5.6	8.9	12.7	21.6
Non-Hispanic black, both sexes.....	0.8	1.0	1.0	2.0	*	2.7	2.7	4.2	*	6.0
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	5.5	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
All origins 3/, both sexes.....	4.0	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
Male.....	5.2	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
Female.....	2.7	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
Hispanic, both sexes.....	2.3	3.9	6.0	10.5	11.6	20.3	24.7	37.3	62.3	150.3
Male.....	3.5	5.4	7.1	15.1	17.2	24.4	32.3	43.8	74.6	131.7
Female.....	*	2.2	5.0	6.2	6.7	16.9	18.8	32.7	54.2	159.9
Mexican, both sexes.....	1.9	3.5	4.9	10.2	11.1	18.2	27.1	37.3	56.6	134.0
Male.....	3.1	5.2	6.1	14.3	16.5	21.8	35.8	50.1	*	*
Female.....	*	*	*	6.1	*	14.9	20.6	28.9	*	135.7
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	6.0	9.3	*	23.7	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.2	7.1	9.2	12.3	15.0	18.5	24.9	36.6	66.2	194.3
Male.....	5.5	9.6	12.8	17.1	20.4	24.6	30.1	38.8	60.0	145.6
Female.....	2.9	4.7	5.8	7.8	10.1	13.2	20.7	35.0	69.9	216.0
Non-Hispanic white, both sexes.....	2.9	4.9	6.4	8.8	11.1	14.5	19.9	30.4	58.7	184.6
Male.....	4.1	7.1	9.5	12.5	15.1	19.4	24.3	32.6	52.9	135.9
Female.....	1.7	2.7	3.3	5.2	7.4	10.1	16.2	28.9	62.3	206.5
Non-Hispanic black, both sexes.....	12.7	22.7	31.6	42.5	50.5	56.3	76.8	104.9	164.3	318.6
Male.....	15.0	28.5	40.8	60.2	72.8	79.3	98.4	116.6	160.8	271.3
Female.....	10.6	17.6	23.8	27.9	33.1	39.2	62.2	98.0	166.1	337.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

All origins 3/, both sexes.....	0.3	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
Male.....	0.3	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
Female.....	*	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
Hispanic, both sexes.....	*	*	*	*	2.4	*	*	6.7	12.4	20.0
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	18.0
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.5	0.5	0.7	1.2	1.6	2.6	4.3	9.3	24.9
Male.....	0.3	0.6	0.7	0.7	1.6	1.7	2.9	4.9	10.1	24.4
Female.....	*	0.4	0.4	0.7	0.8	1.5	2.4	3.9	8.7	25.2
Non-Hispanic white, both sexes.....	*	*	0.2	0.3	0.5	0.9	1.7	3.3	7.9	23.0
Male.....	*	*	*	*	0.8	0.9	2.1	3.7	8.9	22.8
Female.....	*	*	*	*	*	0.8	1.5	3.0	7.2	23.1
Non-Hispanic black, both sexes.....	1.3	2.8	3.0	4.0	6.5	7.6	9.9	15.4	26.6	49.2
Male.....	2.0	3.4	4.0	4.5	8.9	8.2	9.6	21.0	27.9	42.7
Female.....	*	2.4	2.2	3.6	4.6	7.2	10.1	12.1	25.9	51.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All origins 3/, both sexes.....	22.4	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
Male.....	34.0	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
Female.....	10.8	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
Hispanic, both sexes.....	11.7	24.6	49.4	87.3	159.9	250.8	424.1	710.4	1,215.2	2,613.4
Male.....	18.4	39.0	77.0	132.5	237.2	359.9	584.1	961.8	1,525.1	2,695.4
Female.....	4.5	9.8	22.5	45.1	92.3	160.6	300.6	531.9	1,010.7	2,571.3
Mexican, both sexes.....	11.3	25.2	47.3	89.2	155.4	265.9	455.0	671.7	1,129.8	2,263.3
Male.....	17.8	38.5	72.8	131.0	211.6	354.9	626.2	953.8	1,326.5	*
Female.....	4.2	10.1	21.5	47.0	100.0	183.2	326.6	484.5	978.9	2,130.2
Puerto Rican, both sexes.....	16.9	32.9	57.4	118.2	239.9	277.5	521.6	*	*	*
Male.....	*	54.7	79.5	162.6	*	*	*	*	*	*
Female.....	*	*	*	73.9	*	*	*	*	*	*
Cuban, both sexes.....	*	30.4	*	84.2	145.8	211.2	464.0	684.8	1,075.2	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	9.7	19.1	46.4	68.3	137.2	220.2	296.8	686.5	*	*
Male.....	14.3	32.9	79.8	119.3	237.9	378.7	*	*	*	*
Female.....	*	*	18.4	30.5	66.3	128.4	185.5	*	*	*
Non-Hispanic 5/, both sexes.....	23.8	46.1	80.4	126.6	205.7	313.9	499.1	817.5	1,425.0	3,312.1
Male.....	36.2	71.4	125.0	191.5	301.2	444.3	685.8	1,111.5	1,867.6	3,722.6
Female.....	11.6	21.5	37.9	65.4	118.4	200.1	348.3	606.1	1,153.3	3,128.5
Non-Hispanic white, both sexes.....	23.1	44.1	75.7	119.7	197.1	301.1	487.7	810.3	1,430.2	3,369.7
Male.....	36.1	69.5	119.7	183.5	291.9	431.4	676.4	1,110.5	1,894.0	3,814.0
Female.....	10.1	18.8	32.5	58.1	108.7	184.9	331.8	590.3	1,143.1	3,170.6
Non-Hispanic black, both sexes.....	34.2	68.9	132.9	211.0	325.1	477.3	701.6	1,040.9	1,568.8	3,038.5
Male.....	46.1	98.7	193.5	302.3	461.3	648.7	934.0	1,345.9	1,876.2	3,133.7
Female.....	23.7	43.0	81.8	135.9	219.2	350.2	544.1	860.4	1,407.1	3,000.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
All origins 3/, both sexes.....	8.9	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
Male.....	13.2	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
Female.....	4.5	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
Hispanic, both sexes.....	4.4	9.0	17.9	33.9	58.7	94.6	153.8	248.3	421.5	816.8
Male.....	6.8	14.3	27.8	50.0	87.9	134.3	204.1	329.4	524.7	840.6
Female.....	1.8	3.5	8.3	18.9	33.1	61.8	115.0	190.6	353.3	804.5
Mexican, both sexes.....	4.4	9.8	18.6	38.8	63.9	111.8	179.8	248.8	418.9	750.3
Male.....	6.9	15.2	28.7	55.8	88.9	150.2	247.3	348.4	489.9	*
Female.....	*	3.6	8.5	21.6	39.3	76.2	129.2	182.7	364.5	690.1
Puerto Rican, both sexes.....	*	12.2	21.9	35.7	66.1	83.5	165.0	*	*	*
Male.....	*	*	32.7	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	72.6	166.3	241.1	407.9	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.4	6.0	13.2	21.7	46.7	74.9	89.8	221.6	*	*
Male.....	*	10.4	21.9	33.0	84.1	*	*	*	*	*
Female.....	*	*	*	*	*	43.5	63.3	*	*	*
Non-Hispanic 5/, both sexes.....	9.5	18.7	32.9	51.9	84.4	126.0	189.7	293.1	477.8	999.2
Male.....	14.2	28.8	50.7	78.4	122.8	175.3	256.8	389.4	612.2	1,132.5
Female.....	4.9	8.8	15.9	26.9	49.3	83.0	135.5	223.8	395.3	939.6
Non-Hispanic white, both sexes.....	9.4	18.3	31.9	50.6	83.4	123.5	186.7	293.0	480.0	1,010.1
Male.....	14.5	28.8	50.0	77.4	123.4	173.8	255.1	394.1	621.8	1,156.5
Female.....	4.2	8.0	14.1	24.7	46.1	78.6	130.2	218.9	392.2	944.5
Non-Hispanic black, both sexes.....	12.8	25.8	48.8	77.1	114.7	171.2	255.1	353.6	527.6	991.3
Male.....	15.8	37.3	69.2	107.5	152.7	225.4	329.2	423.0	600.0	1,020.2
Female.....	10.1	15.8	31.6	52.2	85.1	131.0	204.9	312.5	489.5	979.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

All origins 3/, both sexes.....	0.3	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
Male.....	0.4	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
Female.....	*	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.4	0.7	1.2	1.5	1.9	2.7	4.3	6.7	15.3
Male.....	0.4	0.5	1.1	1.8	2.1	2.8	3.6	5.1	8.6	17.0
Female.....	*	0.2	0.4	0.7	0.9	1.1	2.0	3.7	5.6	14.6
Non-Hispanic white, both sexes.....	0.3	0.3	0.6	1.0	1.3	1.7	2.6	4.1	6.5	15.6
Male.....	0.4	0.4	0.9	1.5	1.9	2.5	3.2	4.7	8.6	16.9
Female.....	*	*	0.4	0.6	0.8	1.0	2.0	3.6	5.2	15.0
Non-Hispanic black, both sexes.....	*	1.0	1.9	3.0	3.0	3.9	4.4	8.3	10.1	15.5
Male.....	*	*	2.7	4.6	4.8	7.0	7.3	12.1	*	23.4
Female.....	*	*	*	*	*	*	*	6.1	9.3	12.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes.....	13.2	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
Male.....	20.4	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
Female.....	6.1	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
Hispanic, both sexes.....	7.3	15.6	31.4	53.2	100.8	155.7	269.9	460.7	791.9	1,795.7
Male.....	11.5	24.6	48.9	82.2	148.9	225.3	379.2	630.1	998.5	1,853.5
Female.....	2.7	6.3	14.2	26.2	58.8	98.0	185.6	340.4	655.5	1,766.0
Mexican, both sexes.....	6.8	15.4	28.7	50.1	91.3	153.5	274.8	421.1	709.5	1,513.0
Male.....	10.7	23.2	44.2	74.6	122.3	204.7	378.1	601.0	835.0	*
Female.....	2.5	6.5	13.1	25.4	60.7	106.0	197.3	301.8	613.2	1,440.1
Puerto Rican, both sexes.....	10.2	20.7	35.0	82.5	171.9	194.0	355.0	*	*	*
Male.....	*	*	45.8	110.6	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	96.8	138.5	297.7	443.7	667.3	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.3	12.9	33.2	46.6	90.1	144.0	207.0	462.3	*	*
Male.....	9.4	22.2	57.9	86.3	153.8	247.8	*	*	*	*
Female.....	*	*	12.4	17.2	45.2	83.9	122.2	*	*	*
Non-Hispanic 5/, both sexes.....	14.0	27.0	46.8	73.5	119.8	186.1	306.7	520.1	940.5	2,297.5
Male.....	21.7	42.0	73.3	111.3	176.4	266.2	425.4	717.0	1,246.9	2,573.1
Female.....	6.5	12.5	21.5	37.8	68.2	116.1	210.7	378.6	752.4	2,174.3
Non-Hispanic white, both sexes.....	13.4	25.4	43.2	68.1	112.4	176.0	298.4	513.2	943.7	2,343.9
Male.....	21.2	40.3	68.8	104.6	166.6	255.2	418.1	711.8	1,263.5	2,640.6
Female.....	5.7	10.6	18.1	32.9	61.8	105.3	199.6	367.8	745.7	2,211.1
Non-Hispanic black, both sexes.....	20.9	42.2	82.2	130.8	207.4	302.1	442.1	679.0	1,031.1	2,031.7
Male.....	29.5	60.2	121.6	190.2	303.8	416.2	597.5	910.8	1,264.5	2,090.1
Female.....	13.2	26.5	49.0	82.1	132.4	217.5	336.7	541.9	908.3	2,008.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins 3/, both sexes.....	4.8	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
Male.....	7.7	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
Female.....	2.0	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
Hispanic, both sexes.....	2.8	5.9	14.8	22.2	39.4	45.3	69.4	99.0	150.1	279.7
Male.....	4.7	9.8	25.7	38.2	60.0	67.6	104.3	147.4	205.7	297.0
Female.....	*	1.8	4.1	7.3	21.4	26.8	42.5	64.6	113.4	270.8
Mexican, both sexes.....	2.6	5.8	14.0	21.1	37.7	46.1	78.5	99.4	142.9	272.8
Male.....	4.4	9.6	24.3	35.5	55.1	64.8	114.1	156.8	176.5	*
Female.....	*	*	*	6.7	20.6	28.8	51.8	61.4	117.0	262.6
Puerto Rican, both sexes.....	*	*	18.0	26.6	61.4	46.4	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.0	5.5	15.1	23.6	38.9	51.2	61.8	117.2	*	*
Male.....	*	9.7	29.0	48.8	63.1	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.1	9.9	18.5	26.7	38.8	49.7	68.4	100.3	166.6	404.7
Male.....	8.1	15.9	29.8	42.0	57.9	73.0	94.7	132.1	209.5	418.3
Female.....	2.2	4.1	7.7	12.3	21.4	29.4	47.1	77.5	140.3	398.6
Non-Hispanic white, both sexes.....	4.8	9.0	16.5	23.7	34.4	43.9	62.2	92.4	159.8	399.9
Male.....	7.9	14.9	27.3	37.5	51.2	65.3	87.0	121.5	201.6	412.6
Female.....	1.8	3.1	5.9	10.4	18.8	24.9	41.7	71.1	134.0	394.2
Non-Hispanic black, both sexes.....	7.7	17.8	36.2	56.3	82.3	105.5	140.1	200.1	271.4	509.2
Male.....	10.9	25.5	54.6	88.7	129.6	154.1	198.5	280.3	353.9	540.6
Female.....	4.9	11.1	20.6	29.7	45.5	69.4	100.5	152.7	227.9	496.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes.....	8.4	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
Male.....	12.7	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
Female.....	4.1	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
Hispanic, both sexes.....	4.4	9.7	16.6	31.0	61.4	110.4	200.5	361.7	641.8	1,516.0
Male.....	6.8	14.7	23.2	44.0	88.8	157.7	274.9	482.6	792.7	1,556.5
Female.....	1.9	4.5	10.1	18.9	37.4	71.2	143.1	275.7	542.1	1,495.2
Mexican, both sexes.....	4.2	9.6	14.7	29.0	53.6	107.4	196.3	321.7	566.6	1,240.2
Male.....	6.3	13.6	19.9	39.2	67.2	139.9	264.0	444.3	658.5	*
Female.....	*	5.1	9.4	18.7	40.1	77.2	145.5	240.5	496.2	1,177.4
Puerto Rican, both sexes.....	*	13.5	17.0	55.8	110.5	147.6	278.1	*	*	*
Male.....	*	*	*	74.1	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	73.5	108.4	255.8	380.9	573.3	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.3	7.4	18.1	23.0	51.2	92.8	145.2	345.2	*	*
Male.....	*	12.5	29.0	37.5	90.7	*	*	*	*	*
Female.....	*	*	*	*	*	52.6	95.5	*	*	*
Non-Hispanic 5/, both sexes.....	8.9	17.1	28.3	46.7	81.0	136.3	238.3	419.8	773.9	1,892.8
Male.....	13.6	26.1	43.4	69.3	118.5	193.2	330.7	584.9	1,037.4	2,154.8
Female.....	4.4	8.4	13.8	25.5	46.9	86.6	163.7	301.1	612.1	1,775.7
Non-Hispanic white, both sexes.....	8.6	16.4	26.7	44.4	78.0	132.0	236.2	420.8	783.9	1,944.1
Male.....	13.3	25.4	41.5	67.1	115.5	189.9	331.1	590.3	1,062.0	2,228.0
Female.....	3.9	7.5	12.1	22.5	43.0	80.5	157.9	296.7	611.7	1,816.9
Non-Hispanic black, both sexes.....	13.2	24.4	46.0	74.5	125.1	196.7	302.0	478.9	759.7	1,522.6
Male.....	18.7	34.7	67.0	101.5	174.2	262.1	399.0	630.5	910.6	1,549.6
Female.....	8.3	15.4	28.4	52.4	86.9	148.2	236.2	389.2	680.4	1,511.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
All origins 3/, both sexes.....	11.5	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
Male.....	14.4	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
Female.....	8.6	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
Hispanic, both sexes.....	6.5	9.9	13.4	17.8	34.5	54.1	91.2	158.8	286.4	659.3
Male.....	8.9	13.6	18.0	22.5	44.9	64.2	109.7	181.4	340.7	641.6
Female.....	3.9	6.0	9.0	13.4	25.4	45.7	77.0	142.8	250.5	668.4
Mexican, both sexes.....	6.4	9.6	14.0	19.3	33.0	63.8	92.4	145.4	276.1	600.3
Male.....	8.7	12.7	18.0	23.8	43.3	68.8	116.6	149.2	302.2	*
Female.....	3.9	6.2	10.0	14.7	22.9	59.1	74.3	143.0	256.0	621.5
Puerto Rican, both sexes.....	11.4	14.0	15.6	22.7	54.8	46.4	130.4	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	88.0	137.0	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.9	9.4	9.3	12.8	31.6	48.0	72.6	192.3	*	*
Male.....	6.0	13.9	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	24.2	*	63.3	*	*	*
Non-Hispanic 5/, both sexes.....	12.1	18.8	26.7	37.1	59.2	92.9	155.8	274.1	513.5	1,404.5
Male.....	15.1	24.2	34.2	48.3	74.6	114.6	185.8	323.6	585.1	1,412.5
Female.....	9.2	13.5	19.5	26.7	45.2	73.9	131.7	238.6	469.6	1,400.9
Non-Hispanic white, both sexes.....	10.1	15.7	22.6	32.4	54.1	87.0	150.3	268.9	518.4	1,442.1
Male.....	12.9	20.3	29.6	42.4	68.3	107.7	180.5	317.0	590.1	1,445.4
Female.....	7.3	11.0	15.7	22.7	40.9	68.6	125.3	233.7	474.0	1,440.6
Non-Hispanic black, both sexes.....	27.1	42.6	61.9	84.7	117.3	163.2	245.2	388.1	560.2	1,197.9
Male.....	32.5	55.7	77.1	111.8	153.1	207.0	289.3	478.5	663.8	1,244.7
Female.....	22.3	31.3	49.1	62.5	89.4	130.8	215.2	334.6	505.7	1,179.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
All origins 3/, both sexes.....	0.9	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
Male.....	1.0	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
Female.....	0.7	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
Hispanic, both sexes.....	*	1.0	1.8	3.2	6.7	13.7	26.3	53.6	103.2	287.3
Male.....	*	*	*	4.6	8.1	16.0	30.2	59.3	124.6	263.4
Female.....	*	*	*	*	5.5	11.7	23.3	49.5	89.1	299.6
Mexican, both sexes.....	*	*	1.9	3.7	7.0	20.7	28.9	59.5	116.6	298.3
Male.....	*	*	*	*	*	21.8	37.5	63.2	*	*
Female.....	*	*	*	*	*	19.7	22.5	57.0	110.9	323.9
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	24.4	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	1.8	3.5	6.3	11.8	23.0	44.9	90.6	191.7	635.3
Male.....	1.1	2.3	4.1	7.7	14.1	27.1	52.4	105.6	215.4	621.3
Female.....	0.8	1.4	2.9	5.0	9.7	19.4	38.9	79.9	177.2	641.6
Non-Hispanic white, both sexes.....	0.7	1.4	2.6	5.2	10.6	22.0	44.1	89.9	195.3	657.1
Male.....	0.9	1.6	3.2	6.4	12.4	25.8	52.1	104.3	220.1	643.0
Female.....	0.6	1.1	2.0	4.1	8.9	18.6	37.4	79.3	179.9	663.4
Non-Hispanic black, both sexes.....	2.3	5.2	10.8	16.7	24.4	36.6	65.0	122.2	198.0	513.2
Male.....	2.7	7.1	12.6	21.4	31.6	46.6	69.8	149.7	224.0	510.1
Female.....	1.9	3.5	9.2	12.8	18.9	29.2	61.8	106.0	184.3	514.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes.....	10.2	15.5	21.4	28.8	45.0	66.3	105.9	176.1	311.3	747.7
Male.....	12.9	20.2	27.8	37.9	57.6	82.7	128.0	209.5	357.0	765.2
Female.....	7.6	11.0	15.3	20.3	33.4	52.0	88.0	152.2	283.2	739.9
Hispanic, both sexes.....	5.7	8.2	11.1	13.7	25.8	38.8	63.5	102.4	178.3	369.1
Male.....	8.1	11.2	14.8	16.4	34.0	46.6	77.8	120.4	208.6	374.1
Female.....	3.0	5.2	7.4	11.2	18.7	32.4	52.4	89.6	158.3	366.6
Mexican, both sexes.....	5.5	7.8	11.5	14.4	24.2	41.1	62.4	83.4	155.3	300.1
Male.....	7.6	9.9	14.5	16.4	32.6	44.7	78.3	84.9	171.8	*
Female.....	3.1	5.4	8.5	12.3	15.9	37.8	50.6	82.3	142.6	296.2
Puerto Rican, both sexes.....	9.9	11.3	13.1	20.1	42.5	39.0	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	117.0	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.5	8.4	7.2	9.3	23.3	39.0	46.0	129.9	*	*
Male.....	6.0	13.2	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.8	16.3	22.4	30.0	46.4	68.3	108.7	180.2	317.6	764.1
Male.....	13.6	21.2	29.0	39.6	59.3	85.2	131.3	214.4	364.2	784.1
Female.....	8.2	11.6	16.0	21.0	34.6	53.4	90.5	155.7	289.0	755.1
Non-Hispanic white, both sexes.....	9.1	13.8	19.4	26.4	42.5	63.5	104.0	176.0	318.9	779.9
Male.....	11.7	18.2	25.7	35.1	54.6	79.8	126.3	209.2	364.4	795.1
Female.....	6.5	9.5	13.2	18.0	31.2	48.8	85.6	151.7	290.7	773.1
Non-Hispanic black, both sexes.....	24.0	35.7	48.9	66.6	90.6	123.5	177.0	260.3	358.0	678.7
Male.....	28.6	46.2	61.5	88.6	119.2	155.4	216.5	323.2	434.7	728.6
Female.....	19.9	26.7	38.3	48.6	68.3	99.9	150.3	223.0	317.6	659.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All origins 3/, both sexes.....	1.0	2.2	3.4	4.9	8.1	13.0	21.9	38.5	71.3	198.5
Male.....	1.2	2.6	4.5	5.6	9.8	14.9	22.2	40.0	72.7	166.0
Female.....	0.7	1.7	2.3	4.3	6.6	11.3	21.6	37.4	70.4	213.2
Hispanic, both sexes.....	*	1.4	3.3	4.3	7.1	12.3	21.1	34.9	58.9	172.2
Male.....	*	*	4.6	5.9	7.2	16.0	21.5	38.0	66.1	147.1
Female.....	*	*	*	*	6.9	9.2	20.8	32.7	54.2	185.1
Mexican, both sexes.....	*	*	3.4	4.1	5.7	13.8	24.6	37.3	59.4	184.0
Male.....	*	*	4.8	*	*	*	26.6	*	*	*
Female.....	*	*	*	*	*	*	23.1	37.5	*	207.2
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.0	2.3	3.4	5.0	8.2	13.0	21.8	38.7	71.8	199.4
Male.....	1.3	2.8	4.4	5.5	10.0	14.7	22.1	40.1	73.0	166.7
Female.....	0.8	1.8	2.4	4.5	6.5	11.4	21.5	37.6	71.1	214.0
Non-Hispanic white, both sexes.....	0.6	1.3	2.1	3.1	5.6	9.6	17.2	32.7	65.0	188.3
Male.....	0.8	1.7	2.8	3.6	7.2	11.2	17.8	33.4	66.4	159.6
Female.....	0.4	0.9	1.4	2.6	4.2	8.2	16.7	32.1	64.2	201.2
Non-Hispanic black, both sexes.....	3.7	8.6	12.6	20.0	30.2	42.4	66.0	105.9	152.7	339.6
Male.....	4.4	10.1	17.0	22.9	37.2	50.1	67.5	125.4	160.8	268.3
Female.....	3.1	7.3	8.9	17.7	24.9	36.6	64.9	94.4	148.4	367.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
All origins 3/, both sexes.....	7.1	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
Male.....	7.4	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
Female.....	6.8	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
Hispanic, both sexes.....	6.9	11.4	18.9	28.8	43.5	73.2	127.7	229.9	392.2	813.0
Male.....	7.7	13.7	22.6	34.9	54.7	93.8	142.0	251.7	440.7	778.9
Female.....	6.0	9.0	15.3	23.2	33.8	56.2	116.6	214.4	360.1	830.4
Mexican, both sexes.....	6.7	11.7	18.9	32.3	47.9	88.4	144.9	256.6	425.1	862.7
Male.....	8.0	14.0	22.4	36.5	56.6	102.6	159.9	290.7	445.3	*
Female.....	5.3	9.0	15.4	28.1	39.3	75.1	133.6	234.0	409.6	868.1
Puerto Rican, both sexes.....	*	14.9	22.4	35.7	51.9	56.6	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	48.0	96.4	155.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.4	9.4	17.6	22.3	36.6	64.0	110.7	206.3	*	*
Male.....	7.6	10.7	19.3	30.8	50.9	*	*	*	*	*
Female.....	7.2	8.2	16.2	16.1	26.5	46.5	92.2	*	*	*
Non-Hispanic 5/, both sexes.....	7.1	11.9	18.1	26.2	44.4	78.6	142.9	279.0	538.4	1,263.8
Male.....	7.4	12.6	20.9	29.4	51.8	89.7	159.7	303.6	556.6	1,133.7
Female.....	6.9	11.1	15.5	23.2	37.7	68.9	129.4	261.4	527.2	1,321.9
Non-Hispanic white, both sexes.....	5.4	8.7	13.4	20.9	36.8	69.3	132.3	266.4	532.1	1,276.4
Male.....	5.8	9.1	15.4	22.8	42.0	79.2	146.7	289.6	551.4	1,145.3
Female.....	5.0	8.3	11.4	19.0	31.9	60.6	120.4	249.4	520.1	1,335.1
Non-Hispanic black, both sexes.....	17.8	32.4	49.1	68.1	110.5	158.1	251.6	425.2	656.2	1,226.1
Male.....	17.5	36.4	57.8	87.1	142.9	192.0	304.2	478.5	685.1	1,077.1
Female.....	18.0	28.9	41.9	52.5	85.4	133.0	215.9	393.6	641.0	1,285.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

All origins 3/, both sexes.....	0.1	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
Male.....	*	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
Female.....	*	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
Hispanic, both sexes.....	*	*	*	*	*	3.2	5.8	14.4	25.1	74.7
Male.....	*	*	*	*	*	*	*	17.9	31.1	88.3
Female.....	*	*	*	*	*	*	*	11.9	21.2	67.7
Mexican, both sexes.....	*	*	*	*	*	*	*	16.9	22.8	82.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.3	0.5	1.1	2.4	4.2	8.7	17.6	38.6	139.7
Male.....	*	0.4	0.7	1.2	3.1	5.0	10.5	20.3	44.1	130.1
Female.....	*	*	0.3	0.9	1.7	3.4	7.2	15.7	35.2	144.0
Non-Hispanic white, both sexes.....	0.1	0.3	0.5	1.0	2.4	4.1	8.7	18.0	40.0	145.0
Male.....	*	0.5	0.7	1.2	3.1	5.0	10.7	20.6	45.5	136.8
Female.....	*	*	*	0.8	1.8	3.2	7.1	16.1	36.6	148.7
Non-Hispanic black, both sexes.....	*	*	*	1.7	3.0	5.5	9.9	18.6	32.7	103.6
Male.....	*	*	*	*	4.8	6.2	11.6	22.9	41.9	78.2
Female.....	*	*	*	*	*	5.0	8.8	16.0	27.8	113.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

All origins 3/, both sexes.....	1.2	2.1	3.4	5.2	9.9	18.3	33.0	51.7	81.1	139.3
Male.....	1.7	3.0	4.8	7.2	13.6	24.7	44.1	67.1	105.8	166.5
Female.....	0.8	1.2	2.1	3.3	6.6	12.8	24.0	40.6	65.9	127.0
Hispanic, both sexes.....	0.7	1.6	2.5	4.5	6.2	10.8	18.0	25.6	48.4	77.1
Male.....	*	2.2	4.1	5.7	9.5	13.9	29.4	33.4	62.3	89.7
Female.....	*	*	*	3.3	*	8.2	9.3	20.0	39.3	70.6
Mexican, both sexes.....	*	*	*	2.8	4.7	10.5	17.1	17.8	51.8	72.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.3	2.1	3.5	5.3	10.2	18.9	34.0	53.1	82.6	141.8
Male.....	1.8	3.0	4.8	7.3	13.9	25.4	45.0	69.0	108.0	170.1
Female.....	0.9	1.3	2.1	3.3	6.8	13.2	25.0	41.8	67.1	129.2
Non-Hispanic white, both sexes.....	1.1	1.9	3.1	4.8	9.6	18.6	34.8	53.7	82.7	141.9
Male.....	1.5	2.6	4.4	6.6	13.1	25.2	46.7	70.2	109.7	173.0
Female.....	0.7	1.1	1.8	3.0	6.3	12.8	25.0	41.6	66.0	128.0
Non-Hispanic black, both sexes.....	2.7	4.4	6.4	9.5	17.6	24.7	33.4	57.6	91.1	156.9
Male.....	3.7	6.4	8.8	13.9	24.5	34.4	38.2	68.6	100.6	161.6
Female.....	1.9	2.7	4.4	5.9	12.3	17.6	30.2	51.1	86.2	155.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
All origins 3/, both sexes.....	0.8	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
Male.....	1.3	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
Female.....	0.4	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
Hispanic, both sexes.....	*	1.0	1.7	3.0	3.8	6.0	10.3	13.9	24.4	30.4
Male.....	*	*	3.1	4.3	7.2	8.0	18.6	19.0	35.9	40.6
Female.....	*	*	*	*	*	*	*	10.2	16.8	25.2
Mexican, both sexes.....	*	*	*	*	*	*	8.6	*	20.7	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	1.4	2.2	3.1	6.5	11.8	21.8	33.7	50.0	66.2
Male.....	1.4	2.2	3.4	4.9	9.8	17.6	31.3	45.7	72.0	97.5
Female.....	0.4	0.7	1.1	1.5	3.5	6.8	14.1	25.1	36.5	52.3
Non-Hispanic white, both sexes.....	0.7	1.2	2.0	2.8	6.3	12.2	22.9	34.9	51.0	68.4
Male.....	1.1	1.8	3.0	4.5	9.4	18.1	32.9	47.5	73.9	101.4
Female.....	0.3	0.5	0.9	1.2	3.5	6.9	14.6	25.7	36.8	53.6
Non-Hispanic black, both sexes.....	1.8	3.3	4.0	5.8	10.1	11.8	15.2	26.4	41.3	44.6
Male.....	2.9	5.3	5.9	8.9	16.0	18.5	18.6	29.8	49.2	57.9
Female.....	*	1.6	2.5	3.4	5.5	6.8	12.8	24.3	37.1	39.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes.....	0.4	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
Male.....	0.4	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
Female.....	0.5	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
Hispanic, both sexes.....	*	*	*	*	2.4	4.8	7.8	11.7	24.0	46.6
Male.....	*	*	*	*	*	*	10.8	14.4	26.4	49.0
Female.....	*	*	*	*	*	*	*	9.8	22.4	45.4
Mexican, both sexes.....	*	*	*	*	*	*	8.6	*	31.1	48.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.7	1.3	2.1	3.6	7.1	12.2	19.4	32.6	75.6
Male.....	0.4	0.8	1.5	2.4	4.0	7.8	13.8	23.3	36.0	72.6
Female.....	0.5	0.6	1.0	1.9	3.3	6.4	10.9	16.7	30.6	76.9
Non-Hispanic white, both sexes.....	0.4	0.7	1.1	2.0	3.2	6.5	11.9	18.8	31.7	73.6
Male.....	0.4	0.8	1.3	2.1	3.7	7.2	13.8	22.7	35.8	71.6
Female.....	0.4	0.5	0.9	1.8	2.8	5.8	10.4	15.9	29.2	74.4
Non-Hispanic black, both sexes.....	0.9	1.2	2.4	3.7	7.5	13.0	18.3	31.2	49.9	112.2
Male.....	*	*	2.9	5.0	8.5	16.0	19.6	38.7	51.4	103.6
Female.....	*	*	1.9	2.6	6.7	10.7	17.4	26.8	49.1	115.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

All origins 3/, both sexes.....	0.9	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
Male.....	1.0	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
Female.....	0.8	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
Hispanic, both sexes.....	*	1.3	*	*	*	*	*	*	*	17.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	23.8
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.0	1.3	1.5	1.9	2.5	3.2	4.4	7.3	11.7	23.5
Male.....	1.0	1.4	1.9	2.2	2.9	3.4	4.7	8.1	12.6	20.2
Female.....	0.9	1.1	1.1	1.7	2.1	3.0	4.2	6.7	11.2	25.0
Non-Hispanic white, both sexes.....	0.7	1.1	1.2	1.8	2.3	3.0	4.0	7.0	11.6	23.4
Male.....	0.8	1.3	1.5	2.1	2.7	3.2	4.4	7.8	12.2	20.3
Female.....	0.6	0.9	0.9	1.4	1.9	2.9	3.7	6.3	11.2	24.8
Non-Hispanic black, both sexes.....	2.5	2.5	3.7	3.5	5.1	5.0	9.1	13.2	14.9	29.1
Male.....	2.5	2.2	5.2	3.3	6.4	6.0	9.0	14.5	16.9	25.4
Female.....	2.6	2.7	2.5	3.7	4.2	4.3	9.2	12.4	13.9	30.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

All origins 3/, both sexes.....	2.5	3.6	5.6	8.3	14.2	23.8	47.2	94.7	198.7	582.6
Male.....	2.7	4.3	7.1	9.9	17.2	29.8	58.6	118.3	252.3	677.2
Female.....	2.3	3.0	4.2	6.8	11.4	18.6	37.9	77.7	165.7	540.0
Hispanic, both sexes.....	1.6	2.8	5.0	7.6	12.5	22.8	46.7	96.2	170.8	482.3
Male.....	1.9	3.7	6.6	9.1	15.1	26.9	55.1	119.8	204.8	528.2
Female.....	*	2.0	3.5	6.2	10.2	19.4	40.2	79.4	148.3	458.8
Mexican, both sexes.....	1.3	3.0	4.8	7.0	13.3	23.5	51.0	93.3	152.5	486.1
Male.....	*	4.2	5.9	8.5	14.9	24.1	64.1	123.0	159.1	*
Female.....	*	*	*	*	11.7	22.9	41.2	73.7	147.5	437.7
Puerto Rican, both sexes.....	*	*	*	*	*	34.3	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	21.8	40.2	91.7	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.6	3.7	5.6	8.3	14.2	23.7	47.0	94.3	199.7	586.0
Male.....	2.8	4.3	7.1	9.8	17.2	29.7	58.6	118.0	254.1	683.2
Female.....	2.4	3.1	4.3	6.8	11.5	18.5	37.7	77.3	166.2	542.5
Non-Hispanic white, both sexes.....	2.3	3.2	4.9	7.6	13.3	22.0	45.9	92.4	199.1	593.8
Male.....	2.5	3.7	6.1	9.0	15.9	26.7	56.5	113.8	252.6	691.1
Female.....	2.2	2.8	3.8	6.3	10.9	17.8	37.2	76.8	165.9	550.2
Non-Hispanic black, both sexes.....	5.0	7.2	11.8	15.5	23.9	40.6	62.9	118.6	216.5	511.7
Male.....	5.9	8.8	16.1	19.1	32.9	62.1	88.1	172.1	298.1	607.6
Female.....	4.3	5.9	8.1	12.6	16.9	24.6	45.8	87.0	173.5	473.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
All origins 3/, both sexes.....	2.5	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
Male.....	2.7	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
Female.....	2.3	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
Hispanic, both sexes.....	1.6	2.8	5.0	7.6	12.5	22.6	46.2	95.9	170.4	479.5
Male.....	1.9	3.7	6.6	9.1	15.1	26.9	54.2	119.2	203.8	526.8
Female.....	*	1.8	3.4	6.2	10.2	19.2	39.9	79.4	148.3	455.2
Mexican, both sexes.....	1.3	3.0	4.7	7.0	13.3	23.2	50.7	92.9	152.5	483.2
Male.....	*	4.2	5.9	8.5	14.9	24.1	63.3	122.0	159.1	*
Female.....	*	*	*	*	11.7	22.4	41.2	73.7	147.5	434.8
Puerto Rican, both sexes.....	*	*	*	*	*	34.3	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	21.8	40.2	91.7	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.6	3.6	5.6	8.1	14.0	23.4	46.2	92.7	196.4	574.2
Male.....	2.8	4.3	7.0	9.7	17.0	29.2	57.5	116.1	250.7	671.6
Female.....	2.4	3.0	4.2	6.7	11.2	18.2	37.1	75.9	163.0	530.7
Non-Hispanic white, both sexes.....	2.3	3.2	4.9	7.4	13.0	21.7	45.0	90.8	195.5	581.1
Male.....	2.4	3.7	6.1	8.9	15.6	26.3	55.4	112.1	248.9	678.4
Female.....	2.2	2.7	3.7	6.1	10.6	17.6	36.5	75.3	162.5	537.5
Non-Hispanic black, both sexes.....	5.0	7.2	11.7	15.5	23.6	40.3	62.3	116.5	215.4	509.2
Male.....	5.9	8.7	16.1	19.1	32.7	61.8	87.1	167.9	297.4	606.6
Female.....	4.2	5.9	8.0	12.6	16.5	24.2	45.5	86.1	172.3	470.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes.....	2.6	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
Male.....	2.6	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
Female.....	2.7	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
Hispanic, both sexes.....	1.8	2.1	4.4	9.4	16.3	35.8	72.3	124.4	212.4	391.5
Male.....	2.4	1.7	4.2	9.6	19.8	45.4	99.4	167.0	270.9	518.4
Female.....	*	2.4	4.7	9.2	13.2	27.8	51.4	94.1	173.8	326.3
Mexican, both sexes.....	1.3	1.6	3.5	7.2	12.3	32.6	70.3	111.1	187.7	322.8
Male.....	*	*	*	6.9	16.5	40.1	103.3	150.3	254.5	*
Female.....	*	*	*	7.5	*	25.6	45.6	85.2	136.5	261.2
Puerto Rican, both sexes.....	*	*	14.6	*	34.9	38.1	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	122.7	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	8.6	16.5	39.0	68.3	140.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.8	5.9	12.4	26.8	63.3	119.4	209.0	326.8	445.1	610.3
Male.....	2.6	6.0	13.3	28.3	67.9	131.7	236.2	390.8	556.6	824.6
Female.....	2.9	5.8	11.5	25.4	59.1	108.6	187.1	280.7	376.7	514.5
Non-Hispanic white, both sexes.....	2.5	5.5	12.3	27.6	67.1	128.6	224.4	345.9	469.6	640.0
Male.....	2.4	5.7	13.1	28.1	70.1	138.9	247.7	406.1	578.0	852.5
Female.....	2.7	5.4	11.6	27.1	64.2	119.4	205.1	301.8	402.5	544.8
Non-Hispanic black, both sexes.....	4.9	9.9	16.8	28.6	53.6	83.1	130.2	208.4	251.4	335.9
Male.....	5.0	10.2	19.6	38.6	69.5	110.7	184.9	306.9	398.0	569.0
Female.....	4.8	9.6	14.5	20.4	41.2	62.5	93.1	150.2	174.2	243.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

All origins 3/, both sexes.....	0.2	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
Male.....	0.3	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
Female.....	0.2	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
Hispanic, both sexes.....	*	*	*	*	*	4.2	6.0	13.6	19.9	30.0
Male.....	*	*	*	*	*	*	9.1	15.6	28.3	47.6
Female.....	*	*	*	*	*	*	*	12.3	14.3	20.9
Mexican, both sexes.....	*	*	*	*	*	*	*	13.0	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.6	1.6	3.6	8.6	16.6	27.5	39.0	46.9	50.7
Male.....	0.3	0.9	1.8	4.0	10.1	19.3	33.1	48.4	62.1	75.5
Female.....	0.2	0.4	1.4	3.3	7.3	14.3	23.0	32.2	37.6	39.6
Non-Hispanic white, both sexes.....	0.2	0.7	1.6	3.8	9.2	18.2	29.9	41.7	49.8	53.4
Male.....	0.3	0.8	1.7	4.0	10.5	20.6	35.1	50.6	65.0	78.2
Female.....	*	0.5	1.4	3.7	8.1	16.1	25.6	35.3	40.4	42.3
Non-Hispanic black, both sexes.....	*	*	2.2	3.2	6.7	8.5	14.1	21.2	24.6	27.9
Male.....	*	*	3.1	5.0	10.0	14.0	21.9	35.4	39.7	57.9
Female.....	*	*	*	*	4.0	4.4	8.8	12.7	16.6	16.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
All origins 3/, both sexes.....	0.9	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
Male.....	0.8	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
Female.....	1.1	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
Hispanic, both sexes.....	0.8	0.9	1.5	1.8	*	2.8	4.0	5.0	*	11.4
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.0	1.4	1.4	1.7	1.9	2.5	2.9	4.1	6.6	14.2
Male.....	0.8	1.2	0.9	1.4	1.3	1.9	2.3	2.8	4.5	11.1
Female.....	1.1	1.6	1.8	1.9	2.5	2.9	3.5	5.0	7.8	15.5
Non-Hispanic white, both sexes.....	0.7	0.8	1.0	1.2	1.4	2.0	2.6	3.5	6.2	13.8
Male.....	0.5	0.6	0.7	1.0	0.8	1.5	1.9	2.1	3.9	10.4
Female.....	0.8	1.0	1.3	1.5	2.0	2.5	3.3	4.5	7.6	15.3
Non-Hispanic black, both sexes.....	2.9	4.9	4.3	5.3	6.4	5.8	5.8	9.2	9.6	15.3
Male.....	2.7	4.9	3.0	4.9	5.8	5.0	6.7	*	*	*
Female.....	3.1	5.0	5.3	5.6	6.9	6.5	5.2	9.9	9.3	16.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

All origins 3/, both sexes.....	1.4	3.6	8.8	20.2	49.7	95.1	170.8	272.9	380.4	531.1
Male.....	1.5	3.6	9.8	21.5	53.5	105.2	193.5	328.2	476.3	718.6
Female.....	1.4	3.6	7.8	18.9	46.2	86.2	152.5	233.1	321.3	446.8
Hispanic, both sexes.....	0.9	1.0	2.7	6.6	13.1	28.8	61.8	104.8	183.5	345.4
Male.....	*	*	2.8	7.2	17.2	38.0	87.8	146.9	236.9	455.3
Female.....	*	*	2.6	6.1	9.6	21.2	41.8	74.9	148.3	288.8
Mexican, both sexes.....	*	*	2.5	5.5	10.9	26.0	60.7	93.8	164.3	287.9
Male.....	*	*	*	*	14.5	33.3	90.8	131.8	225.9	*
Female.....	*	*	*	*	*	19.2	38.1	68.6	117.0	237.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	104.2	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	14.6	33.3	58.9	115.9	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	3.9	9.3	21.3	52.5	99.9	177.9	282.5	389.7	538.8
Male.....	1.5	3.9	10.4	22.7	56.2	109.9	200.0	338.5	487.8	730.9
Female.....	1.5	3.8	8.3	19.9	49.1	91.1	160.2	242.3	329.4	453.0
Non-Hispanic white, both sexes.....	1.6	4.0	9.7	22.3	56.2	107.8	191.2	299.4	411.6	566.1
Male.....	1.6	4.1	10.6	22.9	58.6	116.2	209.9	352.3	506.8	756.7
Female.....	1.6	3.9	8.8	21.8	53.9	100.3	175.7	260.7	352.6	480.7
Non-Hispanic black, both sexes.....	1.5	4.2	10.1	19.8	40.1	68.6	109.7	177.2	216.5	288.1
Male.....	1.6	4.0	12.8	28.6	53.1	91.5	155.6	262.5	348.1	495.9
Female.....	1.4	4.4	7.8	12.6	30.0	51.6	78.7	126.7	147.2	206.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

All origins 3/, both sexes.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
Male.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
Female.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
Hispanic, both sexes.....	*	*	*	*	2.2	4.5	6.0	18.2	39.0	101.3
Male.....	*	*	*	*	*	6.5	8.7	21.9	48.1	121.9
Female.....	*	*	*	*	*	*	*	15.5	33.0	90.8
Mexican, both sexes.....	*	*	*	*	*	6.1	*	17.4	36.6	102.9
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.8	1.2	1.9	3.5	6.1	14.2	29.0	63.3	167.4
Male.....	0.6	1.0	1.6	2.4	4.5	8.2	19.3	39.9	89.3	242.7
Female.....	0.4	0.6	0.9	1.6	2.6	4.3	10.2	21.1	47.3	133.8
Non-Hispanic white, both sexes.....	0.4	0.8	1.2	1.9	3.1	6.0	13.9	28.8	64.0	170.4
Male.....	0.6	1.0	1.6	2.2	3.8	7.9	18.5	39.3	90.4	247.4
Female.....	0.3	0.5	0.8	1.6	2.5	4.3	10.1	21.1	47.6	135.9
Non-Hispanic black, both sexes.....	0.9	1.1	1.7	3.1	7.5	9.0	20.7	35.0	66.1	147.4
Male.....	*	*	2.2	4.5	11.4	14.0	31.6	52.2	94.7	207.3
Female.....	*	*	*	*	4.5	5.4	13.3	24.8	51.0	123.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes.....	1.5	2.6	4.0	7.3	12.4	21.7	39.5	61.4	91.3	139.4
Male.....	1.5	2.7	4.5	8.1	14.3	26.6	49.4	77.3	118.1	168.8
Female.....	1.5	2.6	3.4	6.6	10.7	17.4	31.6	50.0	74.8	126.1
Hispanic, both sexes.....	1.0	1.8	3.3	6.3	9.9	19.7	37.0	60.5	87.4	112.7
Male.....	*	1.8	4.3	6.8	12.3	25.3	38.9	70.8	101.0	112.1
Female.....	*	*	2.3	5.8	7.7	15.1	35.5	53.2	78.5	113.1
Mexican, both sexes.....	*	*	2.9	6.2	9.8	19.6	38.5	60.8	95.9	93.4
Male.....	*	*	*	*	11.0	25.2	41.6	76.2	*	*
Female.....	*	*	*	6.9	*	14.4	36.2	50.5	82.9	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	25.6	42.4	73.9	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.6	2.7	4.0	7.4	12.6	21.8	39.6	61.4	91.4	140.5
Male.....	1.6	2.8	4.6	8.2	14.5	26.6	50.0	77.6	118.7	171.6
Female.....	1.6	2.7	3.5	6.6	10.9	17.6	31.3	49.8	74.6	126.6
Non-Hispanic white, both sexes.....	1.4	2.3	3.7	7.1	12.4	22.1	41.0	63.6	94.2	145.4
Male.....	1.4	2.3	4.3	7.9	14.3	27.0	51.4	80.0	122.1	177.3
Female.....	1.4	2.3	3.1	6.4	10.6	17.7	32.4	51.5	76.9	131.0
Non-Hispanic black, both sexes.....	3.1	5.7	7.3	11.2	16.9	23.5	33.4	46.6	64.6	94.4
Male.....	3.4	6.5	7.3	12.9	19.1	29.2	42.2	61.6	83.0	117.9
Female.....	2.9	5.0	7.3	9.9	15.3	19.2	27.5	37.8	54.9	85.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

All origins 3/, both sexes.....	0.3	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
Male.....	0.3	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
Female.....	0.2	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
Hispanic, both sexes.....	*	*	*	*	2.2	*	*	6.0	9.0	16.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	15.8
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.6	0.8	1.0	1.7	2.6	3.9	6.9	11.7	24.6
Male.....	0.3	0.7	1.0	1.2	2.1	2.9	4.6	8.3	13.0	24.8
Female.....	0.2	0.4	0.5	0.7	1.3	2.3	3.4	5.9	10.9	24.6
Non-Hispanic white, both sexes.....	0.2	0.5	0.7	0.9	1.6	2.5	3.8	6.9	11.8	25.4
Male.....	0.3	0.6	1.0	1.2	2.0	2.9	4.4	8.1	12.7	25.7
Female.....	*	0.3	0.5	0.7	1.3	2.2	3.3	6.0	11.2	25.3
Non-Hispanic black, both sexes.....	*	1.3	1.3	1.7	2.7	3.6	5.2	6.8	9.4	16.1
Male.....	*	2.1	*	*	4.0	*	*	10.3	*	*
Female.....	*	*	*	*	*	3.7	4.5	*	7.7	16.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes.....	8.7	15.1	21.3	21.7	23.7	26.2	29.5	30.1	27.0	19.7
Male.....	12.2	21.5	31.9	32.3	33.4	35.7	38.7	38.6	34.0	25.5
Female.....	5.4	9.0	11.1	11.8	14.9	18.0	22.1	24.0	22.7	17.1
Hispanic, both sexes.....	11.4	20.3	30.7	31.7	38.7	43.6	52.1	62.7	56.3	26.2
Male.....	17.8	32.7	51.7	49.4	56.7	54.9	65.4	76.0	62.3	29.4
Female.....	4.5	7.5	10.1	15.2	23.0	34.2	41.8	53.2	52.3	24.5
Mexican, both sexes.....	13.8	24.1	35.3	35.1	44.8	51.6	66.4	72.5	62.8	*
Male.....	21.8	36.6	60.8	52.7	61.3	59.0	81.6	91.5	*	*
Female.....	4.9	10.0	9.6	17.4	28.4	44.7	54.9	59.9	61.0	*
Puerto Rican, both sexes.....	11.4	21.6	28.2	43.5	44.4	42.7	*	*	*	*
Male.....	*	*	39.3	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.8	13.9	22.7	22.7	32.0	40.3	33.8	*	*	*
Male.....	11.5	27.2	39.6	39.8	53.1	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	8.3	14.5	20.2	20.8	22.5	24.8	27.9	28.1	25.5	19.4
Male.....	11.3	20.0	29.7	30.7	31.5	34.1	36.8	36.3	32.5	25.3
Female.....	5.5	9.1	11.1	11.5	14.2	16.6	20.7	22.3	21.2	16.7
Non-Hispanic white, both sexes.....	8.5	14.7	20.3	20.4	22.4	25.3	29.2	29.7	26.4	19.9
Male.....	11.5	20.1	29.8	30.0	31.4	34.4	38.1	38.1	33.7	26.4
Female.....	5.5	9.4	11.0	11.2	14.1	17.2	21.9	23.5	21.9	17.0
Non-Hispanic black, both sexes.....	8.0	14.7	23.5	26.3	25.3	23.3	20.4	13.9	15.2	12.1
Male.....	10.9	21.6	35.1	41.8	37.4	36.2	32.3	19.6	22.0	*
Female.....	5.4	8.6	13.8	13.5	15.9	13.7	12.4	10.5	11.6	11.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

All origins 3/, both sexes.....	5.8	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
Male.....	8.0	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
Female.....	3.6	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
Hispanic, both sexes.....	7.8	13.1	17.9	17.5	16.8	16.3	15.9	17.0	10.9	*
Male.....	12.4	22.6	32.1	29.5	30.0	28.1	29.4	34.0	18.9	*
Female.....	2.8	3.3	4.0	6.4	5.3	6.6	*	*	*	*
Mexican, both sexes.....	9.5	15.1	22.1	19.8	19.6	20.2	22.5	21.3	*	*
Male.....	15.5	25.0	40.1	32.0	33.0	32.1	41.6	44.6	*	*
Female.....	2.9	4.0	4.0	7.5	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	11.3	*	22.1	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.6	11.1	13.0	13.1	14.6	*	*	*	*	*
Male.....	8.1	21.1	23.9	24.8	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.5	8.6	11.5	11.2	11.1	10.4	8.5	5.8	3.8	2.1
Male.....	7.4	11.6	17.1	17.1	17.3	16.9	13.9	10.2	7.4	4.5
Female.....	3.7	5.7	6.2	5.7	5.4	4.6	4.0	2.8	1.5	1.0
Non-Hispanic white, both sexes.....	5.7	8.9	11.8	11.1	11.1	10.5	8.8	6.1	3.8	2.2
Male.....	7.6	11.8	17.6	16.8	17.2	17.0	14.2	10.5	7.4	4.7
Female.....	3.8	6.0	6.1	5.6	5.4	4.7	4.3	2.9	1.5	1.0
Non-Hispanic black, both sexes.....	4.3	7.6	11.1	13.2	12.2	11.2	6.6	*	*	*
Male.....	5.8	10.9	16.0	21.4	19.8	20.0	13.6	*	*	*
Female.....	2.9	4.7	6.9	6.5	6.3	4.6	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/, both sexes.....	2.9	6.0	9.1	9.9	12.2	15.4	20.5	23.6	22.9	17.6
Male.....	4.1	8.6	13.3	14.1	15.2	17.9	23.8	27.1	26.0	21.0
Female.....	1.8	3.5	5.1	6.0	9.5	13.2	18.0	21.1	21.0	16.1
Hispanic, both sexes.....	3.6	7.2	12.8	14.1	21.9	27.2	36.2	45.7	45.4	24.3
Male.....	5.5	10.1	19.6	19.9	26.7	26.9	36.0	42.0	43.4	*
Female.....	1.7	4.2	6.1	8.8	17.7	27.6	36.4	48.3	46.7	24.5
Mexican, both sexes.....	4.3	9.0	13.1	15.3	25.2	31.5	43.9	51.2	50.4	*
Male.....	6.3	11.6	20.6	20.6	28.3	26.9	40.0	46.8	*	*
Female.....	*	6.0	5.6	9.9	22.2	35.7	46.8	54.2	*	*
Puerto Rican, both sexes.....	*	*	18.0	21.4	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	9.7	9.6	17.4	26.2	27.3	*	*	*
Male.....	*	*	15.8	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.8	5.8	8.7	9.6	11.4	14.4	19.4	22.3	21.7	17.3
Male.....	3.9	8.4	12.6	13.5	14.3	17.2	22.8	26.2	25.1	20.7
Female.....	1.8	3.4	5.0	5.8	8.8	12.0	16.6	19.5	19.7	15.7
Non-Hispanic white, both sexes.....	2.7	5.9	8.5	9.3	11.4	14.8	20.4	23.6	22.7	17.7
Male.....	3.8	8.3	12.2	13.1	14.2	17.4	23.9	27.6	26.3	21.7
Female.....	1.7	3.4	4.8	5.5	8.7	12.5	17.6	20.6	20.4	16.0
Non-Hispanic black, both sexes.....	3.7	7.1	12.5	13.1	13.1	12.1	13.8	11.1	11.4	11.2
Male.....	5.1	10.7	19.0	20.4	17.5	16.2	18.6	15.4	14.7	*
Female.....	2.5	3.9	6.9	7.1	9.6	9.1	10.6	8.6	9.7	10.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins 3/, both sexes.....	0.2	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
Male.....	*	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
Female.....	*	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
Hispanic, both sexes.....	*	*	*	*	*	2.9	*	9.3	10.1	27.1
Male.....	*	*	*	*	*	*	*	12.1	*	*
Female.....	*	*	*	*	*	*	*	*	*	31.0
Mexican, both sexes.....	*	*	*	*	*	*	*	10.9	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.3	0.4	0.6	1.1	1.8	3.2	5.6	10.4	23.6
Male.....	*	0.3	0.4	0.7	1.2	2.2	3.9	6.7	12.6	25.8
Female.....	*	0.2	0.4	0.5	1.0	1.4	2.7	4.8	9.0	22.6
Non-Hispanic white, both sexes.....	0.2	0.2	0.3	0.5	1.0	1.7	3.0	5.4	10.3	24.0
Male.....	*	0.2	0.3	0.7	1.2	2.1	3.8	6.6	12.7	27.0
Female.....	*	*	0.3	0.4	0.8	1.3	2.4	4.6	8.8	22.7
Non-Hispanic black, both sexes.....	*	*	1.0	1.3	2.0	2.3	5.2	7.1	12.4	19.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	5.6	6.6	13.1	21.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes.....	2.2	3.7	6.4	10.6	17.6	29.6	49.0	85.4	139.0	286.6
Male.....	2.6	4.4	7.7	11.8	19.6	33.3	56.2	102.2	181.2	372.2
Female.....	1.8	3.1	5.2	9.4	15.9	26.4	43.2	73.3	113.0	248.1
Hispanic, both sexes.....	1.6	2.6	7.1	11.2	18.9	34.5	49.8	74.2	119.7	183.6
Male.....	2.2	3.2	8.1	13.5	22.1	34.6	57.5	82.9	152.9	231.2
Female.....	*	2.0	6.0	8.9	16.1	34.5	43.8	67.9	97.8	159.2
Mexican, both sexes.....	1.8	2.7	7.9	12.0	19.7	44.5	60.3	79.0	124.9	187.8
Male.....	2.6	3.2	8.4	14.6	21.6	42.4	69.9	87.1	135.2	*
Female.....	*	*	7.3	9.4	17.9	46.3	53.1	73.7	117.0	175.1
Puerto Rican, both sexes.....	*	*	*	*	*	32.5	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	8.3	16.9	26.2	30.2	72.6	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.3	3.8	6.3	10.5	17.5	29.2	48.8	85.9	139.5	290.9
Male.....	2.6	4.5	7.6	11.6	19.3	33.2	55.9	103.1	182.2	378.6
Female.....	2.0	3.2	5.1	9.4	15.9	25.7	43.1	73.5	113.4	251.6
Non-Hispanic white, both sexes.....	1.6	2.5	4.1	7.4	13.0	23.3	42.3	78.1	132.6	282.1
Male.....	1.9	2.9	4.9	8.6	14.5	26.8	49.6	96.1	177.2	373.0
Female.....	1.2	2.1	3.4	6.3	11.6	20.2	36.2	64.9	104.9	241.4
Non-Hispanic black, both sexes.....	7.1	12.9	22.1	36.4	58.1	80.8	116.6	178.3	244.3	437.2
Male.....	7.4	15.8	28.6	40.6	65.8	94.5	130.3	202.9	287.9	520.3
Female.....	6.9	10.4	16.6	33.0	52.1	70.7	107.3	163.7	221.4	404.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
All origins 3/, both sexes.....	2.1	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
Male.....	2.5	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
Female.....	1.8	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
Hispanic, both sexes.....	1.6	2.6	6.8	10.7	18.7	33.8	49.0	73.2	119.0	179.3
Male.....	2.2	3.2	8.0	12.7	21.6	33.6	56.7	81.8	151.9	224.2
Female.....	*	2.0	5.7	8.8	16.1	34.0	43.1	67.1	97.2	156.3
Mexican, both sexes.....	1.8	2.7	7.6	11.4	19.6	43.6	58.9	78.6	124.2	182.2
Male.....	2.6	3.2	8.2	13.8	21.2	41.3	68.3	87.1	135.2	*
Female.....	*	*	6.9	9.1	17.9	45.8	51.8	72.9	115.8	172.2
Puerto Rican, both sexes.....	*	*	*	*	*	32.5	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	7.7	16.9	25.0	30.2	71.3	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.2	3.7	6.1	10.3	17.2	28.6	48.0	84.4	137.3	287.2
Male.....	2.5	4.4	7.4	11.4	18.9	32.6	55.0	101.3	179.3	374.0
Female.....	1.9	3.1	4.9	9.2	15.6	25.1	42.4	72.2	111.5	248.5
Non-Hispanic white, both sexes.....	1.5	2.4	4.0	7.2	12.7	22.9	41.5	76.7	130.4	278.6
Male.....	1.9	2.8	4.7	8.4	14.2	26.3	48.7	94.5	174.4	368.8
Female.....	1.1	2.1	3.2	6.1	11.3	19.7	35.5	63.7	103.1	238.2
Non-Hispanic black, both sexes.....	6.8	12.5	21.8	35.9	57.5	79.7	115.4	176.2	241.0	432.6
Male.....	7.1	15.3	28.4	39.9	64.7	93.5	128.7	200.1	284.9	514.2
Female.....	6.6	10.1	16.3	32.7	51.8	69.4	106.4	162.0	217.9	400.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes.....	4.9	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
Male.....	6.5	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
Female.....	3.3	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
Hispanic, both sexes.....	3.5	4.2	2.9	3.0	5.2	4.3	8.7	12.0	23.3	99.4
Male.....	5.2	5.6	3.9	4.3	7.9	*	11.2	12.1	21.7	86.9
Female.....	1.7	2.8	*	*	*	*	6.7	11.9	24.3	105.9
Mexican, both sexes.....	3.1	3.6	2.7	3.1	5.1	*	9.6	10.9	22.1	99.1
Male.....	4.5	5.6	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	106.5
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.0	5.9	6.0	6.6	8.0	10.6	15.8	28.8	56.9	226.7
Male.....	6.6	7.6	7.9	8.5	10.1	13.2	18.2	31.6	59.9	188.0
Female.....	3.5	4.2	4.2	4.7	6.1	8.3	13.8	26.8	55.1	244.0
Non-Hispanic white, both sexes.....	4.8	5.7	5.5	6.0	7.5	10.2	15.0	28.5	56.7	230.0
Male.....	6.3	7.3	7.1	7.7	9.3	12.6	17.6	31.1	59.6	191.0
Female.....	3.3	4.1	4.0	4.4	5.8	8.2	12.9	26.6	54.9	247.5
Non-Hispanic black, both sexes.....	7.7	8.6	11.2	12.4	14.7	16.4	26.5	40.1	72.2	223.3
Male.....	9.6	11.6	16.5	18.1	20.0	22.7	28.9	44.8	82.2	182.9
Female.....	6.1	5.9	6.8	7.7	10.5	11.7	24.8	37.3	66.8	239.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
All origins 3/, both sexes.....	21.9	33.6	45.0	58.1	78.7	111.5	176.4	300.6	554.8	1,461.6
Male.....	26.0	40.6	56.0	68.3	88.8	123.1	187.2	315.5	549.4	1,236.6
Female.....	17.9	26.7	34.6	48.4	69.5	101.4	167.6	289.9	558.0	1,562.8
Hispanic, both sexes.....	16.8	24.3	32.3	43.6	56.7	79.6	128.6	225.1	357.7	880.5
Male.....	21.5	32.8	43.8	53.5	63.5	89.2	137.0	228.6	323.7	770.5
Female.....	11.8	15.5	21.0	34.4	50.8	71.7	122.0	222.6	380.1	937.0
Mexican, both sexes.....	18.1	24.6	31.1	44.7	60.8	94.4	139.5	231.0	336.1	849.4
Male.....	22.8	32.0	40.9	51.6	63.3	100.3	151.6	250.4	286.3	*
Female.....	12.9	16.1	21.2	37.7	58.3	88.9	130.5	218.1	374.3	876.9
Puerto Rican, both sexes.....	22.1	35.1	38.4	60.4	66.1	78.9	141.4	*	*	*
Male.....	34.4	54.7	48.6	72.8	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	59.2	88.0	154.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.3	21.0	29.5	35.1	51.7	57.6	121.4	228.0	*	*
Male.....	17.4	29.0	45.7	48.8	65.3	*	*	*	*	*
Female.....	8.7	13.9	15.8	25.0	42.1	49.5	100.0	*	*	*
Non-Hispanic 5/, both sexes.....	22.6	34.5	46.1	59.1	80.2	113.7	179.3	304.6	563.8	1,485.2
Male.....	26.6	41.3	56.8	69.3	90.4	125.2	190.2	320.1	560.5	1,257.5
Female.....	18.7	27.9	35.8	49.5	70.9	103.6	170.5	293.5	565.8	1,587.0
Non-Hispanic white, both sexes.....	20.9	31.6	42.8	55.5	76.8	109.9	176.3	302.7	567.5	1,511.9
Male.....	25.6	38.8	53.1	65.0	86.4	119.8	186.0	313.7	561.7	1,273.4
Female.....	16.1	24.5	32.7	46.3	67.8	101.2	168.3	294.7	571.1	1,618.7
Non-Hispanic black, both sexes.....	36.5	60.9	79.6	102.3	128.1	169.5	248.1	391.4	633.2	1,425.6
Male.....	38.5	68.3	97.8	124.5	150.6	204.7	284.0	473.8	666.8	1,316.9
Female.....	34.8	54.4	64.3	84.1	110.6	143.4	223.8	342.6	615.5	1,468.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes.....	39.7	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
Male.....	56.0	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
Female.....	23.6	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
Hispanic, both sexes.....	30.6	31.9	29.8	28.8	33.0	33.3	42.4	59.6	97.2	160.8
Male.....	46.0	49.4	45.8	41.4	49.8	48.8	56.3	82.4	127.4	193.3
Female.....	13.8	13.9	14.2	17.0	18.3	20.4	31.6	43.4	77.3	144.0
Mexican, both sexes.....	32.1	33.5	28.0	29.6	37.0	34.5	42.8	59.0	84.2	164.2
Male.....	48.4	52.3	43.6	41.5	54.3	46.4	59.1	83.8	*	*
Female.....	14.2	12.3	12.3	17.6	19.8	23.4	30.6	42.6	67.1	147.4
Puerto Rican, both sexes.....	35.1	38.3	31.6	32.5	*	*	*	*	*	*
Male.....	66.0	60.7	43.9	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	19.2	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	27.8	27.8	31.1	24.9	32.5	39.0	40.2	65.0	*	*
Male.....	40.4	42.5	49.8	39.0	47.6	*	*	*	*	*
Female.....	13.8	14.8	15.4	14.4	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	40.8	43.6	39.2	33.0	33.6	38.6	51.0	80.7	137.6	281.5
Male.....	57.3	61.4	57.1	47.1	46.4	52.8	67.5	105.0	182.1	352.5
Female.....	24.8	26.4	22.0	19.7	22.0	26.2	37.6	63.2	110.3	249.8
Non-Hispanic white, both sexes.....	42.7	44.3	38.5	32.1	33.1	38.4	50.9	81.7	142.0	295.2
Male.....	59.4	61.9	55.1	44.8	44.7	52.4	67.0	105.8	186.4	366.8
Female.....	26.1	26.9	22.2	19.8	22.3	26.0	37.7	64.1	114.4	263.1
Non-Hispanic black, both sexes.....	39.4	50.1	51.6	45.9	42.5	44.4	53.7	78.6	102.3	155.1
Male.....	58.0	73.4	83.5	74.5	68.2	68.1	82.1	116.1	160.1	218.5
Female.....	23.0	29.8	24.7	22.4	22.5	26.8	34.5	56.3	71.9	130.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
All origins 3/, both sexes.....	16.4	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
Male.....	24.1	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
Female.....	8.9	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
Hispanic, both sexes.....	14.6	14.4	14.2	14.2	18.2	15.9	18.4	19.4	25.9	20.0
Male.....	21.5	21.6	21.5	18.8	25.8	23.1	24.8	31.1	40.6	28.0
Female.....	7.1	6.9	7.0	10.0	11.6	10.0	13.4	11.0	16.2	15.8
Mexican, both sexes.....	17.2	16.0	14.7	16.8	22.9	17.1	18.2	19.1	23.5	*
Male.....	24.6	24.5	21.2	22.8	31.9	22.9	*	*	*	*
Female.....	9.1	6.3	8.1	10.7	14.0	*	*	*	*	*
Puerto Rican, both sexes.....	11.4	12.6	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	10.4	12.1	13.7	11.5	14.6	*	*	*	*	*
Male.....	15.1	15.7	23.9	*	*	*	*	*	*	*
Female.....	*	8.8	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	16.6	17.3	16.4	15.3	15.7	17.0	18.0	23.8	28.1	27.5
Male.....	24.4	25.1	24.5	22.4	22.0	23.1	24.2	32.2	42.6	47.3
Female.....	9.1	9.7	8.7	8.6	9.9	11.6	13.1	17.7	19.2	18.6
Non-Hispanic white, both sexes.....	17.0	17.4	16.1	15.2	15.6	17.1	18.1	23.8	29.0	28.8
Male.....	24.7	25.2	23.8	22.1	21.5	23.1	23.9	32.1	43.1	49.6
Female.....	9.2	9.7	8.6	8.6	10.1	11.7	13.3	17.7	20.3	19.5
Non-Hispanic black, both sexes.....	16.8	19.6	20.2	17.6	17.7	16.6	16.1	23.1	19.7	15.0
Male.....	26.2	30.2	33.0	27.5	29.1	26.7	25.9	36.8	41.1	29.5
Female.....	8.6	10.3	9.3	9.4	8.8	9.1	9.5	14.9	8.5	9.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All origins 3/, both sexes.....	15.1	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
Male.....	21.9	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
Female.....	8.3	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
Hispanic, both sexes.....	13.7	13.3	13.1	13.5	16.8	15.0	17.7	18.4	25.9	19.5
Male.....	20.0	19.6	19.6	17.4	23.0	21.3	23.6	28.8	40.6	28.0
Female.....	6.8	6.7	6.8	9.8	11.4	9.7	13.1	11.0	16.2	15.1
Mexican, both sexes.....	16.2	14.8	13.6	16.0	20.9	16.6	17.8	18.7	23.5	*
Male.....	23.0	22.7	19.3	21.4	28.3	21.8	*	*	*	*
Female.....	8.7	6.0	7.9	10.4	13.6	*	*	*	*	*
Puerto Rican, both sexes.....	10.2	11.3	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	9.6	11.1	12.3	10.5	14.2	*	*	*	*	*
Male.....	13.8	13.6	21.3	*	*	*	*	*	*	*
Female.....	*	8.8	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.2	15.7	14.7	13.9	14.1	15.5	16.7	22.9	26.9	26.3
Male.....	22.1	22.5	21.7	20.1	19.3	20.7	21.8	30.8	40.3	45.0
Female.....	8.5	9.1	8.1	8.1	9.3	11.0	12.5	17.3	18.7	18.0
Non-Hispanic white, both sexes.....	15.5	15.8	14.4	13.8	14.0	15.6	16.7	23.0	27.7	27.6
Male.....	22.4	22.5	20.9	19.7	18.7	20.5	21.5	30.7	40.6	47.1
Female.....	8.6	9.1	8.0	8.1	9.6	11.1	12.8	17.3	19.7	18.9
Non-Hispanic black, both sexes.....	15.6	17.8	18.6	16.3	15.8	15.6	15.2	21.0	19.5	15.0
Male.....	23.9	27.3	30.0	25.1	25.6	25.2	23.9	33.6	40.4	29.5
Female.....	8.2	9.5	9.0	9.1	8.1	8.5	9.2	13.5	8.5	9.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
All origins 3/, both sexes.....	23.3	25.5	22.3	17.6	17.8	21.4	32.4	56.0	107.8	249.5
Male.....	31.9	35.6	32.1	24.7	24.5	29.6	42.8	71.7	137.0	299.0
Female.....	14.7	15.8	12.8	10.8	11.7	14.3	24.1	44.8	89.9	227.3
Hispanic, both sexes.....	16.0	17.6	15.7	14.5	14.8	17.3	24.0	40.2	71.3	140.8
Male.....	24.5	27.7	24.4	22.6	24.0	25.6	31.5	51.3	86.8	165.3
Female.....	6.7	7.0	7.1	7.0	6.7	10.5	18.2	32.3	61.1	128.2
Mexican, both sexes.....	14.9	17.6	13.3	12.9	14.1	17.4	24.6	39.9	60.7	143.5
Male.....	23.8	27.8	22.4	18.8	22.4	23.5	34.1	51.2	*	*
Female.....	5.1	6.0	4.2	6.9	*	*	17.5	32.5	*	131.3
Puerto Rican, both sexes.....	23.6	25.7	21.9	22.1	*	*	*	*	*	*
Male.....	42.5	38.8	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	17.4	15.8	17.4	13.4	17.8	22.4	23.0	*	*	*
Male.....	25.3	26.8	25.9	24.0	*	*	*	*	*	*
Female.....	8.7	*	10.2	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	24.2	26.3	22.7	17.7	18.0	21.6	32.9	56.9	109.5	254.1
Male.....	32.9	36.3	32.6	24.8	24.4	29.7	43.4	72.8	139.4	305.1
Female.....	15.7	16.7	13.3	11.1	12.1	14.5	24.5	45.4	91.1	231.2
Non-Hispanic white, both sexes.....	25.7	26.9	22.4	16.9	17.5	21.4	32.8	57.9	113.0	266.4
Male.....	34.7	36.7	31.3	22.7	23.1	29.3	43.0	73.7	143.3	317.3
Female.....	16.8	17.1	13.6	11.2	12.2	14.3	24.4	46.4	94.2	243.7
Non-Hispanic black, both sexes.....	22.6	30.5	31.4	28.3	24.8	27.8	37.6	55.5	82.5	140.2
Male.....	31.8	43.1	50.5	47.0	39.1	41.4	56.2	79.3	119.0	189.0
Female.....	14.4	19.4	15.4	13.0	13.6	17.8	25.0	41.4	63.4	120.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
All origins 3/, both sexes.....	1.8	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
Male.....	2.9	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
Female.....	0.7	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
Hispanic, both sexes.....	2.1	2.3	2.4	4.3	5.8	8.7	12.3	23.7	42.0	87.1
Male.....	3.3	4.2	4.1	7.0	9.8	13.0	15.3	29.4	52.8	106.5
Female.....	*	*	*	*	*	5.1	9.9	19.6	34.9	77.1
Mexican, both sexes.....	1.8	2.4	2.1	4.0	6.6	8.8	12.8	23.9	34.5	83.1
Male.....	3.2	4.5	*	*	10.6	*	*	*	*	*
Female.....	*	*	*	*	*	*	21.7	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.9	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	2.6	3.2	4.0	5.7	8.9	16.3	30.7	62.7	145.5
Male.....	2.9	4.0	4.9	6.1	8.0	12.3	22.0	38.9	80.0	177.2
Female.....	0.7	1.2	1.6	2.0	3.7	5.9	11.8	24.8	52.1	131.3
Non-Hispanic white, both sexes.....	1.8	2.6	3.2	4.0	5.8	9.2	17.0	32.4	66.3	155.1
Male.....	3.0	4.1	4.8	5.9	7.7	12.9	22.5	41.0	84.0	186.9
Female.....	0.7	1.2	1.6	2.1	4.0	5.8	12.4	26.2	55.4	140.9
Non-Hispanic black, both sexes.....	1.7	2.5	3.5	5.0	6.4	8.1	12.8	17.2	26.1	47.5
Male.....	2.7	3.8	6.0	8.9	12.1	10.0	20.6	24.2	38.9	73.2
Female.....	*	*	*	*	*	6.7	7.4	13.0	19.3	37.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes.....	0.7	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
Male.....	1.0	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
Female.....	0.5	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.8	1.2	1.3	1.3	1.7	1.8	2.0	2.9	4.4	5.6
Male.....	1.1	1.6	1.8	1.7	2.2	2.6	2.3	3.5	6.0	7.3
Female.....	0.5	0.9	0.7	1.0	1.3	1.1	1.7	2.5	3.4	4.9
Non-Hispanic white, both sexes.....	0.7	1.0	1.2	1.2	1.5	1.6	1.6	2.4	3.8	4.9
Male.....	0.9	1.3	1.6	1.4	2.0	2.2	1.9	2.9	5.4	6.5
Female.....	0.5	0.8	0.7	0.9	1.1	1.0	1.4	2.1	2.8	4.2
Non-Hispanic black, both sexes.....	1.4	2.7	2.2	2.9	4.3	4.1	6.0	8.0	12.2	15.3
Male.....	2.0	4.0	3.6	4.4	4.8	7.5	8.0	12.1	15.4	*
Female.....	*	1.5	*	*	3.9	*	4.7	5.5	10.4	13.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes.....	16.3	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
Male.....	20.9	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
Female.....	11.7	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
Hispanic, both sexes.....	10.2	10.5	8.7	6.8	3.4	*	*	*	*	*
Male.....	15.3	15.3	12.8	10.4	5.6	*	*	*	*	*
Female.....	4.7	5.5	4.7	3.4	*	*	*	*	*	*
Mexican, both sexes.....	8.6	9.7	6.4	4.4	*	*	*	*	*	*
Male.....	13.2	14.4	10.5	7.1	*	*	*	*	*	*
Female.....	3.6	4.4	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	18.9	20.3	17.0	*	*	*	*	*	*	*
Male.....	36.2	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.0	9.4	9.7	*	*	*	*	*	*	*
Male.....	17.2	14.7	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	17.1	17.4	12.2	6.7	3.7	2.4	2.1	2.3	2.6	3.4
Male.....	21.6	22.2	16.5	8.2	4.1	2.7	2.4	2.5	3.0	4.2
Female.....	12.6	12.6	8.2	5.3	3.3	2.1	1.8	2.2	2.3	3.1
Non-Hispanic white, both sexes.....	18.4	17.8	12.0	6.1	3.5	2.3	2.0	2.3	2.5	3.5
Male.....	23.1	22.7	15.5	7.0	3.4	2.3	2.3	2.3	2.8	4.3
Female.....	13.7	13.0	8.5	5.3	3.5	2.3	1.8	2.3	2.3	3.1
Non-Hispanic black, both sexes.....	15.1	20.0	18.3	12.8	6.2	4.2	3.2	*	*	*
Male.....	19.8	26.1	29.3	20.8	10.4	7.5	*	*	*	*
Female.....	11.0	14.6	9.0	6.2	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes.....	3.2	3.9	4.5	4.5	5.7	7.0	11.0	19.2	37.4	95.5
Male.....	5.0	6.3	7.0	6.7	8.4	9.8	14.3	25.1	46.1	109.8
Female.....	1.4	1.5	2.1	2.3	3.2	4.6	8.4	15.0	32.1	89.1
Hispanic, both sexes.....	2.3	3.6	2.9	2.7	4.6	3.5	7.4	10.0	21.8	44.2
Male.....	3.7	6.2	4.9	4.0	6.7	*	8.7	13.2	24.5	43.4
Female.....	*	*	*	*	*	*	6.4	*	19.9	44.7
Mexican, both sexes.....	2.9	4.1	3.2	3.3	4.3	*	*	10.9	*	51.0
Male.....	4.7	6.7	5.4	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.3	3.9	4.6	4.6	5.8	7.2	11.2	19.7	38.1	97.7
Male.....	5.2	6.2	7.1	7.0	8.5	10.0	14.6	25.7	47.2	113.1
Female.....	1.5	1.6	2.2	2.4	3.2	4.8	8.5	15.4	32.6	90.9
Non-Hispanic white, both sexes.....	3.5	4.0	4.7	4.5	5.7	7.0	11.0	19.6	38.7	101.3
Male.....	5.4	6.4	7.1	6.7	8.5	9.7	14.4	25.4	47.7	116.4
Female.....	1.6	1.7	2.3	2.4	3.1	4.6	8.3	15.2	33.1	94.5
Non-Hispanic black, both sexes.....	3.0	4.1	5.7	6.4	6.9	10.4	13.7	26.5	38.2	71.4
Male.....	5.1	6.8	9.0	10.8	10.0	15.0	19.0	35.9	55.1	89.4
Female.....	*	1.7	2.8	2.8	4.5	7.0	10.1	21.0	29.4	64.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes.....	16.0	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
Male.....	24.2	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
Female.....	7.9	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
Hispanic, both sexes.....	6.6	7.7	7.3	7.5	6.3	5.6	8.5	6.0	13.9	*
Male.....	10.5	12.0	11.9	11.6	10.9	11.4	16.6	13.2	30.2	*
Female.....	2.5	3.3	2.8	3.7	*	*	*	*	*	*
Mexican, both sexes.....	5.9	6.9	6.0	6.2	5.7	*	*	*	*	*
Male.....	9.4	10.8	9.6	9.3	9.4	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	13.5	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.3	7.4	7.2	9.3	*	*	*	*	*	*
Male.....	10.7	10.0	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	17.3	17.9	17.0	14.7	13.7	11.8	13.8	16.6	16.9	16.8
Male.....	26.1	26.6	25.6	23.1	22.5	20.8	25.9	33.5	38.7	46.2
Female.....	8.6	9.4	8.9	6.8	5.7	3.9	4.0	4.4	3.4	3.6
Non-Hispanic white, both sexes.....	19.9	20.3	19.4	16.3	15.1	13.0	14.8	17.9	17.9	17.9
Male.....	29.7	29.8	28.8	25.1	24.6	22.6	27.7	36.0	40.9	49.8
Female.....	10.1	11.0	10.1	7.9	6.2	4.4	4.2	4.7	3.6	3.6
Non-Hispanic black, both sexes.....	6.5	6.4	5.2	6.5	4.9	3.6	5.6	5.7	6.8	*
Male.....	11.2	10.6	8.7	12.7	9.0	8.0	13.0	14.0	17.6	*
Female.....	2.5	2.8	2.2	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes.....	6.3	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
Male.....	9.0	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
Female.....	3.6	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
Hispanic, both sexes.....	7.2	6.1	5.7	3.9	3.6	*	*	*	*	*
Male.....	10.4	9.7	9.4	6.5	6.5	*	*	*	*	*
Female.....	3.7	2.4	*	*	*	*	*	*	*	*
Mexican, both sexes.....	8.3	6.5	5.7	3.9	*	*	*	*	*	*
Male.....	12.8	10.1	9.0	6.4	*	*	*	*	*	*
Female.....	3.3	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.5	5.9	*	*	*	*	*	*	*	*
Male.....	6.0	9.7	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.1	5.2	4.0	3.1	2.8	2.6	2.2	2.2	2.3	2.0
Male.....	8.8	7.5	6.3	4.7	4.0	3.8	3.2	2.6	2.8	2.4
Female.....	3.6	3.0	1.8	1.6	1.7	1.6	1.3	2.0	2.0	1.8
Non-Hispanic white, both sexes.....	3.8	3.4	2.6	2.3	2.1	2.2	1.7	1.8	2.1	1.8
Male.....	4.9	4.6	3.9	3.2	2.9	2.8	2.4	2.0	2.3	1.9
Female.....	2.7	2.3	1.3	1.4	1.4	1.7	1.1	1.6	1.9	1.7
Non-Hispanic black, both sexes.....	20.6	16.8	13.6	9.4	7.9	5.8	6.6	7.1	5.3	*
Male.....	33.6	27.6	24.2	16.6	13.1	12.0	11.6	10.3	*	*
Female.....	9.1	7.4	4.7	3.5	3.9	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes.....	3.8	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
Male.....	5.9	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
Female.....	1.6	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
Hispanic, both sexes.....	4.6	2.9	2.5	1.8	2.2	*	*	*	*	*
Male.....	7.0	4.8	4.2	3.2	*	*	*	*	*	*
Female.....	2.0	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.3	3.3	2.7	*	*	*	*	*	*	*
Male.....	8.4	5.3	4.2	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.0	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.6	2.9	2.1	1.7	1.4	1.2	0.9	0.9	0.7	0.5
Male.....	5.7	4.4	3.4	2.7	2.0	1.7	1.5	1.0	*	*
Female.....	1.6	1.4	0.9	0.8	0.9	0.8	*	0.7	0.8	*
Non-Hispanic white, both sexes.....	2.1	1.9	1.4	1.2	1.2	1.1	0.7	0.8	0.7	*
Male.....	2.9	2.6	2.0	1.8	1.6	1.3	1.1	0.9	*	*
Female.....	1.4	1.1	0.7	0.7	0.8	0.9	*	0.6	0.9	*
Non-Hispanic black, both sexes.....	13.1	9.5	7.5	5.1	4.1	2.7	*	*	*	*
Male.....	24.0	16.6	14.2	9.7	6.9	5.7	*	*	*	*
Female.....	3.5	3.3	1.8	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes.....	2.5	2.5	2.1	1.5	1.3	1.4	1.4	1.6	1.5
Male.....	3.1	3.4	3.2	2.2	2.0	2.0	1.8	2.2	1.7
Female.....	1.9	1.6	1.0	0.8	0.8	0.8	1.0	1.2	1.5
Hispanic, both sexes.....	2.6	3.2	3.2	2.1	*	*	*	*	*
Male.....	3.4	4.8	5.2	3.3	*	*	*	*	*
Female.....	1.7	*	*	*	*	*	*	*	*
Mexican, both sexes.....	3.0	3.1	3.0	*	*	*	*	*	*
Male.....	4.4	4.8	4.8	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.5	2.3	1.9	1.4	1.3	1.4	1.3	1.4	1.5
Male.....	3.1	3.1	2.9	2.0	1.9	2.0	1.7	2.3	1.7
Female.....	2.0	1.6	0.9	0.8	0.8	0.8	1.0	1.2	1.4
Non-Hispanic white, both sexes.....	1.7	1.6	1.2	1.1	1.0	1.1	1.0	1.0	1.3
Male.....	2.0	2.0	1.9	1.5	1.3	1.5	1.4	1.1	*
Female.....	1.3	1.2	0.6	0.6	0.6	0.8	0.7	1.0	1.3
Non-Hispanic black, both sexes.....	7.5	7.3	6.2	4.3	3.8	3.2	4.4	5.2	*
Male.....	9.7	11.0	10.0	6.9	6.2	6.2	*	*	*
Female.....	5.6	4.1	2.9	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes.....	3.4	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
Male.....	4.1	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
Female.....	2.7	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
Hispanic, both sexes.....	1.4	1.4	1.4	*	*	*	*	*	*	*
Male.....	1.6	2.0	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	1.5	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.7	3.8	2.7	1.7	1.0	0.9	0.8	0.7	1.3	1.6
Male.....	4.5	4.8	3.4	2.0	1.1	1.2	0.9	0.8	1.3	1.7
Female.....	2.9	2.8	2.1	1.4	0.9	0.7	0.7	0.6	1.3	1.5
Non-Hispanic white, both sexes.....	3.9	3.9	2.7	1.7	1.0	0.8	0.8	0.6	1.3	1.6
Male.....	4.6	4.8	3.3	2.0	1.0	1.0	0.9	*	1.4	1.9
Female.....	3.2	3.1	2.2	1.5	1.0	0.6	0.7	0.6	1.3	1.5
Non-Hispanic black, both sexes.....	3.6	4.1	3.8	2.0	*	*	*	*	*	*
Male.....	4.8	6.3	5.6	3.3	*	*	*	*	*	*
Female.....	2.5	2.3	2.2	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes.....	3.3	3.5	2.5	1.6	0.9	0.8	0.7	0.6	1.2	1.6
Male.....	3.9	4.5	3.1	1.9	1.0	1.0	0.8	0.7	1.1	1.8
Female.....	2.6	2.6	2.0	1.3	0.8	0.6	0.6	0.6	1.2	1.5
Hispanic, both sexes.....	1.4	1.3	1.4	*	*	*	*	*	*	*
Male.....	1.6	1.8	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	1.5	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.5	3.7	2.6	1.6	0.9	0.8	0.7	0.6	1.2	1.5
Male.....	4.3	4.7	3.3	1.9	1.0	1.0	0.8	0.7	1.0	1.7
Female.....	2.8	2.8	2.0	1.3	0.8	0.6	0.7	0.6	1.2	1.5
Non-Hispanic white, both sexes.....	3.7	3.9	2.6	1.7	0.9	0.7	0.7	0.6	1.2	1.6
Male.....	4.4	4.7	3.1	1.9	0.9	0.8	0.8	*	1.1	1.8
Female.....	3.1	3.1	2.1	1.4	0.9	0.6	0.6	0.6	1.3	1.5
Non-Hispanic black, both sexes.....	3.5	4.1	3.6	1.9	*	*	*	*	*	*
Male.....	4.7	6.3	5.3	3.3	*	*	*	*	*	*
Female.....	2.5	2.3	2.2	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/, both sexes.....	0.4	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
Male.....	0.4	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
Female.....	0.5	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.6	0.8	1.3	1.6	2.6	3.6	5.0	7.6	10.8
Male.....	0.4	0.6	0.9	1.5	1.7	2.5	4.4	6.0	9.5	12.5
Female.....	0.5	0.6	0.8	1.0	1.5	2.8	2.9	4.3	6.5	10.1
Non-Hispanic white, both sexes.....	0.4	0.5	0.8	1.2	1.5	2.5	3.4	4.8	7.4	10.6
Male.....	0.4	0.6	0.8	1.4	1.7	2.5	4.3	6.0	9.4	12.3
Female.....	0.5	0.5	0.7	1.0	1.4	2.5	2.7	4.0	6.2	9.9
Non-Hispanic black, both sexes.....	1.0	1.1	1.5	2.7	2.5	4.7	5.6	7.6	12.2	14.1
Male.....	*	*	*	3.4	*	*	*	*	*	*
Female.....	*	*	1.7	*	*	5.5	5.2	7.5	11.2	13.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

All origins 3/, both sexes.....	10.0	15.5	19.5	18.9	18.5	17.0	14.1	10.7	7.3	4.3
Male.....	14.6	22.7	30.6	30.0	30.4	28.1	24.5	19.4	13.8	9.0
Female.....	5.3	8.4	8.8	8.3	7.8	7.3	5.8	4.4	3.3	2.3
Hispanic, both sexes.....	11.0	18.6	23.2	22.1	23.1	21.8	19.3	20.3	12.0	*
Male.....	18.1	31.9	41.5	37.4	42.1	38.9	36.0	40.9	20.8	*
Female.....	3.4	4.8	5.3	7.9	6.5	7.7	6.4	*	*	*
Mexican, both sexes.....	13.0	20.6	27.0	23.0	25.2	24.8	27.1	23.4	*	*
Male.....	21.7	33.8	48.9	37.8	42.9	40.7	50.8	50.1	*	*
Female.....	3.5	5.5	5.0	8.0	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	16.7	15.1	29.9	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	9.3	17.4	19.5	19.8	22.9	23.0	*	*	*	*
Male.....	14.6	32.2	36.6	36.0	47.6	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	9.8	15.0	19.0	18.5	18.1	16.5	13.7	10.1	7.1	4.4
Male.....	14.1	21.4	29.3	29.2	29.3	27.2	23.6	18.1	13.5	8.9
Female.....	5.6	8.7	9.1	8.3	7.9	7.2	5.8	4.4	3.2	2.3
Non-Hispanic white, both sexes.....	10.1	15.4	19.1	18.2	17.6	16.0	13.5	10.2	7.2	4.5
Male.....	14.5	21.7	29.4	28.4	28.1	25.9	22.7	18.0	13.7	9.3
Female.....	5.7	9.1	9.1	8.2	7.7	7.2	5.9	4.5	3.2	2.4
Non-Hispanic black, both sexes.....	7.8	14.3	21.1	23.7	24.3	22.7	17.3	8.8	6.3	*
Male.....	11.9	22.3	33.9	41.6	43.1	43.4	35.6	17.7	*	*
Female.....	4.2	7.5	10.3	9.1	9.6	7.4	5.0	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All origins 3/, both sexes.....	23.2	24.4	18.5	11.0	6.6	4.4	4.1	4.7	4.9	6.3
Male.....	28.3	30.1	23.6	13.0	7.6	4.8	4.5	5.2	5.6	8.6
Female.....	18.2	19.0	13.7	9.1	5.7	4.1	3.7	4.4	4.4	5.2
Hispanic, both sexes.....	13.7	14.0	11.9	10.6	5.1	3.4	*	*	*	*
Male.....	19.3	19.2	17.0	15.4	8.1	*	*	*	*	*
Female.....	7.6	8.6	6.8	6.1	*	*	*	*	*	*
Mexican, both sexes.....	12.5	13.0	9.5	8.2	4.5	*	*	*	*	*
Male.....	18.0	17.9	14.7	12.4	*	*	*	*	*	*
Female.....	6.5	7.4	4.2	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	26.0	28.8	23.4	21.4	*	*	*	*	*	*
Male.....	47.9	40.8	33.7	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.8	11.9	11.4	10.2	*	*	*	*	*	*
Male.....	17.7	17.2	14.7	*	*	*	*	*	*	*
Female.....	9.5	7.3	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	24.5	25.5	19.1	11.0	6.7	4.5	4.2	4.7	4.9	6.3
Male.....	29.6	31.2	24.0	12.8	7.5	4.7	4.5	5.1	5.7	8.6
Female.....	19.6	20.0	14.4	9.4	6.0	4.2	3.9	4.4	4.5	5.3
Non-Hispanic white, both sexes.....	26.5	26.2	18.7	10.5	6.5	4.4	4.4	4.9	5.1	6.4
Male.....	31.5	31.6	22.5	11.2	6.7	4.1	4.6	5.1	5.8	8.7
Female.....	21.5	20.7	15.1	9.9	6.4	4.7	4.2	4.8	4.6	5.4
Non-Hispanic black, both sexes.....	21.7	29.7	27.4	18.4	9.7	6.1	3.1	3.5	*	*
Male.....	27.8	37.8	42.5	30.4	16.6	11.2	*	*	*	*
Female.....	16.3	22.7	14.8	8.5	4.3	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2004. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All origins 3/, both sexes.....	11.4	11.3	10.7	9.7	10.0	9.5	11.0	13.3	13.4	11.9
Male.....	18.5	18.4	18.0	16.9	17.4	17.8	22.1	28.2	32.0	35.9
Female.....	4.4	4.3	3.7	3.0	3.2	2.3	2.1	2.6	1.9	1.0
Hispanic, both sexes.....	7.5	6.1	6.2	4.4	5.3	3.5	5.6	*	8.6	*
Male.....	11.9	10.6	10.6	8.0	9.5	7.4	10.8	*	19.8	*
Female.....	2.7	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	8.2	6.7	5.8	4.3	4.9	*	*	*	*	*
Male.....	13.2	11.0	9.2	7.7	*	*	*	*	*	*
Female.....	2.7	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.3	5.0	6.0	*	*	*	*	*	*	*
Male.....	8.1	8.9	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.9	11.9	11.1	10.1	10.3	10.0	11.3	13.9	13.6	12.1
Male.....	19.3	19.3	18.7	17.6	17.9	18.6	22.8	29.4	32.7	37.0
Female.....	4.6	4.6	3.9	3.1	3.4	2.5	2.1	2.7	1.9	1.0
Non-Hispanic white, both sexes.....	11.6	12.2	11.8	10.7	11.2	10.8	12.2	14.9	14.6	13.1
Male.....	18.4	19.7	19.5	18.3	19.4	19.7	24.2	31.5	34.9	40.1
Female.....	4.9	4.9	4.2	3.4	3.6	2.9	2.4	2.8	2.1	1.0
Non-Hispanic black, both sexes.....	16.2	12.3	10.2	8.8	7.0	6.2	6.7	7.3	6.1	*
Male.....	29.9	21.8	19.1	16.8	12.3	13.7	16.3	16.3	16.2	*
Female.....	4.1	4.1	2.6	2.4	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes Central and South American and Other and unknown Hispanic.

5/ Includes races other than white and black.

6/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.