

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX:

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Shigellosis and amebiasis (A03,A06)									
All races, both sexes	12	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
Certain other intestinal infections (A04,A07-A09)									
All races, both sexes	1,328	13	10	-	-	-	1	2	1
Male	524	10	6	-	-	-	-	2	1
Female	804	3	4	-	-	-	1	-	-
White, both sexes	1,224	8	7	-	-	-	1	1	1
Male	478	6	5	-	-	-	-	1	1
Female	746	2	2	-	-	-	1	-	-
Black, both sexes	84	4	2	-	-	-	-	1	-
Male	37	3	1	-	-	-	-	1	-
Female	47	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	1	1	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-
Female	8	-	1	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Tuberculosis (A16-A19)									
All races, both sexes	776	1	6	-	1	2	9	7	13
Male	468	-	1	-	-	2	5	3	11
Female	308	1	5	-	1	-	4	4	2
White, both sexes	439	-	4	-	1	1	5	5	6
Male	253	-	-	-	-	1	3	2	5
Female	186	-	4	-	1	-	2	3	1
Black, both sexes	244	-	2	-	-	-	2	2	5
Male	161	-	1	-	-	-	1	1	4
Female	83	-	1	-	-	-	1	1	1
American Indian 1/, both sexes	20	1	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	1	2	-	2
Male	42	-	-	-	-	1	1	-	2
Female	31	-	-	-	-	-	1	-	-
Respiratory tuberculosis (A16)									
All races, both sexes	615	-	3	-	-	2	5	5	6
Male	393	-	1	-	-	2	2	3	5
Female	222	-	2	-	-	-	3	2	1
White, both sexes	351	-	1	-	-	1	3	3	2
Male	216	-	-	-	-	1	2	2	2
Female	135	-	1	-	-	-	1	1	-
Black, both sexes	192	-	2	-	-	-	1	2	4
Male	134	-	1	-	-	-	-	1	3
Female	58	-	1	-	-	-	1	1	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	60	-	-	-	-	1	1	-	-
Male	35	-	-	-	-	1	-	-	-
Female	25	-	-	-	-	-	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Scarlet fever and erysipelas (A38,A46)									
All races, both sexes	2	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Meningococcal infection (A39)									
All races, both sexes	211	22	19	12	6	34	22	9	6
Male	106	14	12	5	2	24	11	3	3
Female	105	8	7	7	4	10	11	6	3
White, both sexes	161	19	14	5	5	29	16	7	3
Male	75	11	8	2	2	20	8	2	-
Female	86	8	6	3	3	9	8	5	3
Black, both sexes	44	3	5	7	1	3	4	1	3
Male	27	3	4	3	-	3	2	-	3
Female	17	-	1	4	1	-	2	1	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	1	2	1	-
Male	4	-	-	-	-	1	1	1	-
Female	1	-	-	-	-	-	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)									
All races, both sexes	5,532	167	71	30	37	44	52	56	85
Male	2,861	94	41	16	19	25	29	29	46
Female	2,671	73	30	14	18	19	23	27	39
White, both sexes	4,550	101	50	24	28	32	39	34	50
Male	2,334	59	27	12	13	16	25	19	29
Female	2,216	42	23	12	15	16	14	15	21
Black, both sexes	826	61	14	5	5	9	13	16	34
Male	432	31	10	3	2	6	4	6	16
Female	394	30	4	2	3	3	9	10	18
American Indian 1/, both sexes	34	2	2	-	-	-	-	1	-
Male	13	1	-	-	-	-	-	-	-
Female	21	1	2	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	122	3	5	1	4	3	-	5	1
Male	82	3	4	1	4	3	-	4	1
Female	40	-	1	-	-	-	-	1	-
Malignant neoplasms (C00-C97)									
All races, both sexes	553,091	92	420	489	525	745	968	1,359	2,557
Male	286,082	50	233	269	300	446	576	692	1,162
Female	267,009	42	187	220	225	299	392	667	1,395
White, both sexes	480,011	75	330	397	421	576	757	1,081	1,991
Male	247,403	42	190	215	241	344	454	559	941
Female	232,608	33	140	182	180	232	303	522	1,050
Black, both sexes	61,945	11	70	72	78	136	165	215	434
Male	32,817	4	34	42	45	82	97	94	168
Female	29,128	7	36	30	33	54	68	121	266
American Indian 1/, both sexes	1,914	2	3	5	3	6	7	9	17
Male	997	2	2	3	1	4	3	5	8
Female	917	-	1	2	2	2	4	4	9
Asian or Pacific Islander 2/, both sexes	9,221	4	17	15	23	27	39	54	115
Male	4,865	2	7	9	13	16	22	34	45
Female	4,356	2	10	6	10	11	17	20	70

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Malignant neoplasm of larynx (C32)									
All races, both sexes	3,861	-	-	-	2	1	-	-	4
Male	3,041	-	-	-	-	-	-	-	2
Female	820	-	-	-	2	1	-	-	2
White, both sexes	3,096	-	-	-	2	1	-	-	4
Male	2,426	-	-	-	-	-	-	-	2
Female	670	-	-	-	2	1	-	-	2
Black, both sexes	719	-	-	-	-	-	-	-	-
Male	578	-	-	-	-	-	-	-	-
Female	141	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
All races, both sexes	155,521	2	2	-	1	3	19	37	160
Male	90,469	2	1	-	-	3	11	28	74
Female	65,052	-	1	-	1	-	8	9	86
White, both sexes	136,912	1	2	-	1	1	13	27	128
Male	78,747	1	1	-	-	1	9	20	62
Female	58,165	-	1	-	1	-	4	7	66
Black, both sexes	16,066	1	-	-	-	2	3	8	24
Male	10,153	1	-	-	-	2	-	6	9
Female	5,913	-	-	-	-	-	3	2	15
American Indian 1/, both sexes	480	-	-	-	-	-	-	-	1
Male	270	-	-	-	-	-	-	-	1
Female	210	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2,063	-	-	-	-	-	3	2	7
Male	1,299	-	-	-	-	-	2	2	2
Female	764	-	-	-	-	-	1	-	5

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Malignant melanoma of skin (C43)									
All races, both sexes	7,420	-	-	-	1	7	34	58	128
Male	4,592	-	-	-	1	4	17	33	77
Female	2,828	-	-	-	-	3	17	25	51
White, both sexes	7,245	-	-	-	1	6	34	55	125
Male	4,514	-	-	-	1	4	17	32	76
Female	2,731	-	-	-	-	2	17	23	49
Black, both sexes	135	-	-	-	-	1	-	2	1
Male	60	-	-	-	-	-	-	1	1
Female	75	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	1	2
Malignant neoplasm of breast (C50)									
All races, both sexes	42,300	-	-	-	-	1	10	83	384
Male	428	-	-	-	-	-	-	2	1
Female	41,872	-	-	-	-	1	10	81	383
White, both sexes	36,127	-	-	-	-	1	4	60	265
Male	360	-	-	-	-	-	-	-	1
Female	35,767	-	-	-	-	1	4	60	264
Black, both sexes	5,424	-	-	-	-	-	5	20	103
Male	63	-	-	-	-	-	-	2	-
Female	5,361	-	-	-	-	-	5	18	103
American Indian 1/, both sexes	131	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-
Female	130	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	618	-	-	-	-	-	1	2	15
Male	4	-	-	-	-	-	-	-	-
Female	614	-	-	-	-	-	1	2	15

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Malignant neoplasm of cervix uteri (C53)									
All races, both sexes	4,200	-	-	-	1	3	14	77	153
Female	4,200	-	-	-	1	3	14	77	153
White, both sexes	3,158	-	-	-	-	3	10	63	111
Female	3,158	-	-	-	-	3	10	63	111
Black, both sexes	869	-	-	-	1	-	4	12	39
Female	869	-	-	-	1	-	4	12	39
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	2
Female	30	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	143	-	-	-	-	-	-	2	1
Female	143	-	-	-	-	-	-	2	1
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)									
All races, both sexes	6,586	-	-	-	-	-	3	2	25
Female	6,586	-	-	-	-	-	3	2	25
White, both sexes	5,397	-	-	-	-	-	1	2	18
Female	5,397	-	-	-	-	-	1	2	18
Black, both sexes	1,061	-	-	-	-	-	1	-	6
Female	1,061	-	-	-	-	-	1	-	6
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	109	-	-	-	-	-	1	-	1
Female	109	-	-	-	-	-	1	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)									
All races, both sexes	12,655	20	104	166	150	105	120	159	247
Male	7,011	10	58	89	87	63	69	94	170
Female	5,644	10	46	77	63	42	51	65	77
White, both sexes	11,672	16	82	135	120	84	99	140	220
Male	6,486	9	46	72	69	52	57	82	145
Female	5,186	7	36	63	51	32	42	58	75
Black, both sexes	772	2	18	26	26	16	19	15	21
Male	414	1	10	15	16	9	12	9	19
Female	358	1	8	11	10	7	7	6	2
American Indian 1/, both sexes	41	-	2	2	2	2	-	1	-
Male	20	-	1	1	1	1	-	-	-
Female	21	-	1	1	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	170	2	2	3	2	3	2	3	6
Male	91	-	1	1	1	1	-	3	6
Female	79	2	1	2	1	2	2	-	-

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races, both sexes	56,518	39	146	168	204	323	395	412	506
Male	29,888	21	76	98	125	202	253	226	313
Female	26,630	18	70	70	79	121	142	186	193
White, both sexes	50,394	36	121	136	170	249	300	315	392
Male	26,766	19	64	77	103	155	196	180	240
Female	23,628	17	57	59	67	94	104	135	152
Black, both sexes	5,071	1	20	21	22	58	74	75	85
Male	2,558	-	9	14	12	36	44	33	58
Female	2,513	1	11	7	10	22	30	42	27
American Indian 1/, both sexes	199	1	1	1	1	3	2	2	7
Male	101	1	1	-	-	2	1	2	4
Female	98	-	-	1	1	1	1	-	3
Asian or Pacific Islander 2/, both sexes	854	1	4	10	11	13	19	20	22
Male	463	1	2	7	10	9	12	11	11
Female	391	-	2	3	1	4	7	9	11

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Hodgkin's disease (C81)									
All races, both sexes	1,287	-	-	3	5	21	52	60	85
Male	686	-	-	1	3	6	29	31	48
Female	601	-	-	2	2	15	23	29	37
White, both sexes	1,156	-	-	2	3	17	40	54	76
Male	616	-	-	1	1	4	26	28	42
Female	540	-	-	1	2	13	14	26	34
Black, both sexes	110	-	-	1	1	4	10	6	7
Male	57	-	-	-	1	2	3	3	5
Female	53	-	-	1	-	2	7	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	1	-	2	-	2
Male	9	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	2	-	1
Non-Hodgkin's lymphoma (C82-C85)									
All races, both sexes	22,729	1	7	23	24	60	120	124	169
Male	11,812	-	5	16	12	46	80	64	108
Female	10,917	1	2	7	12	14	40	60	61
White, both sexes	20,892	-	4	18	22	46	85	89	129
Male	10,838	-	4	12	12	38	58	50	81
Female	10,054	-	-	6	10	8	27	39	48
Black, both sexes	1,369	1	2	4	1	8	27	30	26
Male	728	-	-	3	-	3	16	12	19
Female	641	1	2	1	1	5	11	18	7
American Indian 1/, both sexes	83	-	-	-	-	1	-	-	4
Male	41	-	-	-	-	1	-	-	3
Female	42	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	385	-	1	1	1	5	8	5	10
Male	205	-	1	1	-	4	6	2	5
Female	180	-	-	-	1	1	2	3	5

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)									
All races, both sexes	100	-	2	-	1	-	1	-	-
Male	60	-	2	-	1	-	1	-	-
Female	40	-	-	-	-	-	-	-	-
White, both sexes	89	-	2	-	1	-	-	-	-
Male	51	-	2	-	1	-	-	-	-
Female	38	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, both sexes	63,060	24	129	123	137	243	268	292	409
Male	31,728	13	75	64	76	142	177	189	243
Female	31,332	11	54	59	61	101	91	103	166
White, both sexes	55,066	19	97	105	107	187	223	251	330
Male	27,697	11	62	55	59	110	144	165	206
Female	27,369	8	35	50	48	77	79	86	124
Black, both sexes	6,696	4	25	17	24	48	37	34	54
Male	3,378	1	11	9	15	27	29	20	26
Female	3,318	3	14	8	9	21	8	14	28
American Indian 1/, both sexes	250	-	-	-	-	1	3	3	2
Male	118	-	-	-	-	1	1	2	1
Female	132	-	-	-	-	-	2	1	1
Asian or Pacific Islander 2/, both sexes	1,048	1	7	1	6	7	5	4	23
Male	535	1	2	-	2	4	3	2	10
Female	513	-	5	1	4	3	2	2	13

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)									
All races, both sexes	13,546	60	53	62	37	40	55	68	97
Male	6,559	32	33	32	17	27	35	30	52
Female	6,987	28	20	30	20	13	20	38	45
White, both sexes	12,171	48	44	46	30	29	44	47	82
Male	5,900	25	25	24	15	18	30	24	46
Female	6,271	23	19	22	15	11	14	23	36
Black, both sexes	1,136	9	9	13	6	8	8	18	15
Male	520	4	8	6	2	7	2	5	6
Female	616	5	1	7	4	1	6	13	9
American Indian 1/, both sexes	45	-	-	1	-	2	2	2	-
Male	20	-	-	1	-	1	2	1	-
Female	25	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	194	3	-	2	1	1	1	1	-
Male	119	3	-	1	-	1	1	-	-
Female	75	-	-	1	1	-	-	1	-
Anemias (D50-D64)									
All races, both sexes	4,494	10	35	24	28	27	58	62	91
Male	1,669	6	16	17	12	9	30	29	41
Female	2,825	4	19	7	16	18	28	33	50
White, both sexes	3,450	4	15	9	12	10	11	15	14
Male	1,235	2	8	7	5	2	6	9	6
Female	2,215	2	7	2	7	8	5	6	8
Black, both sexes	967	5	19	15	16	16	46	47	77
Male	395	4	8	10	7	7	23	20	35
Female	572	1	11	5	9	9	23	27	42
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	1	1	-	-	1	1	-	-
Male	30	-	-	-	-	-	1	-	-
Female	27	1	1	-	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Alzheimer's disease (G30)									
All races, both sexes	49,558	-	-	-	-	-	-	1	1
Male	14,438	-	-	-	-	-	-	1	-
Female	35,120	-	-	-	-	-	-	-	1
White, both sexes	46,460	-	-	-	-	-	-	1	1
Male	13,524	-	-	-	-	-	-	1	-
Female	32,936	-	-	-	-	-	-	-	1
Black, both sexes	2,728	-	-	-	-	-	-	-	-
Male	776	-	-	-	-	-	-	-	-
Female	1,952	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	95	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	275	-	-	-	-	-	-	-	-
Male	112	-	-	-	-	-	-	-	-
Female	163	-	-	-	-	-	-	-	-
Major cardiovascular diseases (I00-I78)									
All races, both sexes	936,923	636	234	136	226	493	816	1,295	2,497
Male	435,912	341	128	78	128	295	488	800	1,578
Female	501,011	295	106	58	98	198	328	495	919
White, both sexes	816,219	434	161	95	147	331	527	845	1,660
Male	379,095	233	96	56	90	201	327	540	1,105
Female	437,124	201	65	39	57	130	200	305	555
Black, both sexes	104,280	171	63	34	62	144	255	407	764
Male	47,980	91	27	18	30	80	139	231	426
Female	56,300	80	36	16	32	64	116	176	338
American Indian 1/, both sexes	3,198	7	2	2	5	2	10	12	18
Male	1,670	3	1	1	2	2	6	5	11
Female	1,528	4	1	1	3	-	4	7	7
Asian or Pacific Islander 2/, both sexes	13,226	24	8	5	12	16	24	31	55
Male	7,167	14	4	3	6	12	16	24	36
Female	6,059	10	4	2	6	4	8	7	19

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Ischemic heart diseases (I20-I25)									
All races, both sexes	515,204	30	10	8	8	33	90	236	699
Male	260,574	17	7	5	4	21	60	181	523
Female	254,630	13	3	3	4	12	30	55	176
White, both sexes	455,400	23	10	6	7	28	64	160	516
Male	230,951	14	7	3	4	18	43	124	405
Female	224,449	9	3	3	3	10	21	36	111
Black, both sexes	51,265	4	-	2	-	4	20	66	166
Male	24,625	2	-	2	-	2	12	48	104
Female	26,640	2	-	-	-	2	8	18	62
American Indian 1/, both sexes	1,740	1	-	-	-	-	4	4	7
Male	1,008	-	-	-	-	-	3	3	6
Female	732	1	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	6,799	2	-	-	1	1	2	6	10
Male	3,990	1	-	-	-	1	2	6	8
Female	2,809	1	-	-	1	-	-	-	2
Acute myocardial infarction (I21-I22)									
All races, both sexes	192,898	20	1	5	1	14	38	108	267
Male	100,306	10	1	3	1	8	22	75	194
Female	92,592	10	-	2	-	6	16	33	73
White, both sexes	170,584	16	1	3	1	11	28	71	201
Male	89,383	9	1	1	1	6	18	50	155
Female	81,201	7	-	2	-	5	10	21	46
Black, both sexes	19,112	2	-	2	-	2	9	32	62
Male	9,045	1	-	2	-	1	4	21	35
Female	10,067	1	-	-	-	1	5	11	27
American Indian 1/, both sexes	656	1	-	-	-	-	1	2	3
Male	384	-	-	-	-	-	-	1	3
Female	272	1	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	2,546	1	-	-	-	1	-	3	1
Male	1,494	-	-	-	-	1	-	3	1
Female	1,052	1	-	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other acute ischemic heart diseases (I24)									
All races, both sexes	3,363	1	1	-	-	4	4	4	12
Male	1,749	1	1	-	-	3	3	3	10
Female	1,614	-	-	-	-	1	1	1	2
White, both sexes	2,839	1	1	-	-	4	2	4	8
Male	1,453	1	1	-	-	3	1	3	6
Female	1,386	-	-	-	-	1	1	1	2
Black, both sexes	456	-	-	-	-	-	1	-	4
Male	258	-	-	-	-	-	1	-	4
Female	198	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	43	-	-	-	-	-	1	-	-
Male	26	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-
Other forms of chronic ischemic heart disease (I20,I25)									
All races, both sexes	318,943	9	8	3	7	15	48	124	420
Male	158,519	6	5	2	3	10	35	103	319
Female	160,424	3	3	1	4	5	13	21	101
White, both sexes	281,977	6	8	3	6	13	34	85	307
Male	140,115	4	5	2	3	9	24	71	244
Female	141,862	2	3	1	3	4	10	14	63
Black, both sexes	31,697	2	-	-	-	2	10	34	100
Male	15,322	1	-	-	-	1	7	27	65
Female	16,375	1	-	-	-	1	3	7	35
American Indian 1/, both sexes	1,041	-	-	-	-	-	2	2	4
Male	598	-	-	-	-	-	2	2	3
Female	443	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4,228	1	-	-	1	-	2	3	9
Male	2,484	1	-	-	-	-	2	3	7
Female	1,744	-	-	-	1	-	-	-	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)									
All races, both sexes	69,554	1	-	-	1	-	15	33	111
Male	35,896	-	-	-	1	-	12	26	89
Female	33,658	1	-	-	-	-	3	7	22
White, both sexes	57,794	-	-	-	1	-	13	19	83
Male	29,467	-	-	-	1	-	10	14	69
Female	28,327	-	-	-	-	-	3	5	14
Black, both sexes	10,418	1	-	-	-	-	1	12	24
Male	5,615	-	-	-	-	-	1	10	16
Female	4,803	1	-	-	-	-	-	2	8
American Indian 1/, both sexes	266	-	-	-	-	-	1	1	3
Male	159	-	-	-	-	-	1	1	3
Female	107	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,076	-	-	-	-	-	-	1	1
Male	655	-	-	-	-	-	-	1	1
Female	421	-	-	-	-	-	-	-	-
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
All races, both sexes	249,389	8	8	3	6	15	33	91	309
Male	122,623	6	5	2	2	10	23	77	230
Female	126,766	2	3	1	4	5	10	14	79
White, both sexes	224,183	6	8	3	5	13	21	66	224
Male	110,648	4	5	2	2	9	14	57	175
Female	113,535	2	3	1	3	4	7	9	49
Black, both sexes	21,279	1	-	-	-	2	9	22	76
Male	9,707	1	-	-	-	1	6	17	49
Female	11,572	-	-	-	-	1	3	5	27
American Indian 1/, both sexes	775	-	-	-	-	-	1	1	1
Male	439	-	-	-	-	-	1	1	-
Female	336	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3,152	1	-	-	1	-	2	2	8
Male	1,829	1	-	-	-	-	2	2	6
Female	1,323	-	-	-	1	-	-	-	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other heart diseases (I26-I51)									
All races, both sexes	165,428	457	168	96	152	355	498	687	1,062
Male	72,122	238	98	58	92	218	300	407	642
Female	93,306	219	70	38	60	137	198	280	420
White, both sexes	143,761	315	114	63	95	229	309	433	700
Male	61,933	164	71	38	62	139	194	267	435
Female	81,828	151	43	25	33	90	115	166	265
Black, both sexes	19,298	117	48	28	45	113	167	236	328
Male	8,958	59	24	16	24	68	91	129	186
Female	10,340	58	24	12	21	45	76	107	142
American Indian 1/, both sexes	591	6	1	1	4	2	5	4	9
Male	295	3	-	1	2	2	3	2	5
Female	296	3	1	-	2	-	2	2	4
Asian or Pacific Islander 2/, both sexes	1,778	19	5	4	8	11	17	14	25
Male	936	12	3	3	4	9	12	9	16
Female	842	7	2	1	4	2	5	5	9
Acute and subacute endocarditis (I33)									
All races, both sexes	1,172	3	1	2	1	5	8	13	27
Male	665	2	-	1	-	1	6	7	17
Female	507	1	1	1	1	4	2	6	10
White, both sexes	927	1	1	1	1	1	5	10	17
Male	536	1	-	-	-	-	3	6	13
Female	391	-	1	1	1	1	2	4	4
Black, both sexes	214	2	-	-	-	3	3	3	8
Male	111	1	-	-	-	1	3	1	2
Female	103	1	-	-	-	2	-	2	6
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	1	-	1	-	-	2
Male	11	-	-	1	-	-	-	-	2
Female	7	-	-	-	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
All races, both sexes	767	20	24	9	11	17	21	29	24
Male	404	10	13	4	5	10	11	17	12
Female	363	10	11	5	6	7	10	12	12
White, both sexes	590	11	15	8	2	15	10	21	13
Male	309	6	11	4	1	8	5	11	7
Female	281	5	4	4	1	7	5	10	6
Black, both sexes	142	7	8	1	7	1	7	6	10
Male	72	4	2	-	3	1	2	5	5
Female	70	3	6	1	4	-	5	1	5
American Indian 1/, both sexes	6	1	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	1	-	-
Female	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	1	1	-	2	1	3	2	1
Male	19	-	-	-	1	1	3	1	-
Female	10	1	1	-	1	-	-	1	1
Heart failure (I50)									
All races, both sexes	55,704	31	9	10	11	21	13	27	36
Male	21,175	22	7	4	5	11	9	15	21
Female	34,529	9	2	6	6	10	4	12	15
White, both sexes	50,142	19	5	7	9	12	5	15	23
Male	18,949	14	5	3	4	5	5	9	13
Female	31,193	5	-	4	5	7	-	6	10
Black, both sexes	4,965	12	4	2	2	8	5	12	12
Male	1,951	8	2	1	1	5	3	6	7
Female	3,014	4	2	1	1	3	2	6	5
American Indian 1/, both sexes	175	-	-	-	-	-	1	-	-
Male	79	-	-	-	-	-	1	-	-
Female	96	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	422	-	-	1	-	1	2	-	1
Male	196	-	-	-	-	1	-	-	1
Female	226	-	-	1	-	-	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other diseases of circulatory system (I71-I78)									
All races, both sexes	26,036	16	5	3	8	17	42	49	105
Male	14,084	6	2	1	6	9	24	36	67
Female	11,952	10	3	2	2	8	18	13	38
White, both sexes	22,963	13	4	2	5	14	30	35	76
Male	12,611	4	2	1	5	7	19	24	53
Female	10,352	9	2	1	-	7	11	11	23
Black, both sexes	2,595	3	-	1	3	2	11	11	25
Male	1,207	2	-	-	1	1	5	9	13
Female	1,388	1	-	1	2	1	6	2	12
American Indian 1/, both sexes	87	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	391	-	1	-	-	1	1	3	4
Male	229	-	-	-	-	1	-	3	1
Female	162	-	1	-	-	-	1	-	3
Aortic aneurysm and dissection (I71)									
All races, both sexes	15,810	-	-	-	5	10	26	38	69
Male	9,661	-	-	-	3	7	16	30	48
Female	6,149	-	-	-	2	3	10	8	21
White, both sexes	14,210	-	-	-	2	9	17	27	50
Male	8,797	-	-	-	2	6	12	19	37
Female	5,413	-	-	-	-	3	5	8	13
Black, both sexes	1,277	-	-	-	3	1	8	8	17
Male	671	-	-	-	1	1	4	8	11
Female	606	-	-	-	2	-	4	-	6
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	282	-	-	-	-	-	1	3	2
Male	172	-	-	-	-	-	-	3	-
Female	110	-	-	-	-	-	1	-	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Other diseases of arteries, arterioles and capillaries (I72-I78)									
All races, both sexes	10,226	16	5	3	3	7	16	11	36
Male	4,423	6	2	1	3	2	8	6	19
Female	5,803	10	3	2	-	5	8	5	17
White, both sexes	8,753	13	4	2	3	5	13	8	26
Male	3,814	4	2	1	3	1	7	5	16
Female	4,939	9	2	1	-	4	6	3	10
Black, both sexes	1,318	3	-	1	-	1	3	3	8
Male	536	2	-	-	-	-	1	1	2
Female	782	1	-	1	-	1	2	2	6
American Indian 1/, both sexes	46	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	109	-	1	-	-	1	-	-	2
Male	57	-	-	-	-	1	-	-	1
Female	52	-	1	-	-	-	-	-	1
Other disorders of circulatory system (I80-I99)									
All races, both sexes	4,603	27	4	5	7	10	27	58	75
Male	1,981	16	2	3	3	6	10	18	39
Female	2,622	11	2	2	4	4	17	40	36
White, both sexes	3,787	20	3	4	5	4	18	33	49
Male	1,618	13	1	3	2	3	6	9	25
Female	2,169	7	2	1	3	1	12	24	24
Black, both sexes	761	5	1	1	2	6	8	23	26
Male	335	2	1	-	1	3	4	8	14
Female	426	3	-	1	1	3	4	15	12
American Indian 1/, both sexes	18	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	37	2	-	-	-	-	-	1	-
Male	20	1	-	-	-	-	-	-	-
Female	17	1	-	-	-	-	-	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Pneumonia (J12-J18)									
All races, both sexes	63,548	280	93	44	32	57	119	134	217
Male	27,994	158	50	26	17	32	64	78	121
Female	35,554	122	43	18	15	25	55	56	96
White, both sexes	56,252	165	60	34	27	40	86	91	159
Male	24,385	93	26	21	14	24	50	48	87
Female	31,867	72	34	13	13	16	36	43	72
Black, both sexes	5,909	101	29	7	3	14	29	37	55
Male	2,880	56	21	4	2	6	14	26	32
Female	3,029	45	8	3	1	8	15	11	23
American Indian 1/, both sexes	285	6	-	1	1	1	1	4	1
Male	148	2	-	-	-	1	-	2	1
Female	137	4	-	1	1	-	1	2	-
Asian or Pacific Islander 2/, both sexes	1,102	8	4	2	1	2	3	2	2
Male	581	7	3	1	1	1	-	2	1
Female	521	1	1	1	-	1	3	-	1
Other acute lower respiratory infections (J20-J22)									
All races, both sexes	425	43	11	3	4	4	1	5	3
Male	160	25	4	2	1	4	-	5	1
Female	265	18	7	1	3	-	1	-	2
White, both sexes	372	25	8	2	1	4	1	4	1
Male	134	12	4	2	-	4	-	4	-
Female	238	13	4	-	1	-	1	-	1
Black, both sexes	46	14	3	-	3	-	-	1	1
Male	22	10	-	-	1	-	-	1	1
Female	24	4	3	-	2	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	3	-	1	-	-	-	-	1
Male	3	3	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Chronic lower respiratory diseases (J40-J47)									
All races, both sexes	122,009	36	51	48	91	86	104	125	171
Male	60,004	23	28	26	54	58	62	72	93
Female	62,005	13	23	22	37	28	42	53	78
White, both sexes	112,840	23	28	23	40	40	63	67	99
Male	54,816	16	13	12	21	26	39	36	48
Female	58,024	7	15	11	19	14	24	31	51
Black, both sexes	7,607	11	21	24	46	44	37	51	65
Male	4,238	6	14	14	29	31	20	31	41
Female	3,369	5	7	10	17	13	17	20	24
American Indian 1/, both sexes	429	1	2	-	-	1	2	4	-
Male	228	1	1	-	-	1	2	3	-
Female	201	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1,133	1	-	1	5	1	2	3	7
Male	722	-	-	-	4	-	1	2	4
Female	411	1	-	1	1	1	1	1	3
Bronchitis, chronic and unspecified (J40-J42)									
All races, both sexes	1,167	18	14	2	3	3	2	3	5
Male	503	9	7	2	1	2	2	3	4
Female	664	9	7	-	2	1	-	-	1
White, both sexes	1,073	9	7	2	2	2	2	3	3
Male	458	5	3	2	-	1	2	3	2
Female	615	4	4	-	2	1	-	-	1
Black, both sexes	80	7	6	-	1	1	-	-	1
Male	35	3	3	-	1	1	-	-	1
Female	45	4	3	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	1	-	-	-	-	-	-
Male	2	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	1	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	1
Female	4	1	-	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Emphysema (J43)									
All races, both sexes	16,700	3	2	-	-	1	4	6	5
Male	8,671	3	-	-	-	1	4	1	4
Female	8,029	-	2	-	-	-	-	5	1
White, both sexes	15,607	3	-	-	-	-	2	6	4
Male	7,985	3	-	-	-	-	2	1	3
Female	7,622	-	-	-	-	-	-	5	1
Black, both sexes	901	-	1	-	-	1	2	-	1
Male	559	-	-	-	-	1	2	-	1
Female	342	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	59	-	1	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-
Female	30	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	133	-	-	-	-	-	-	-	-
Male	98	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-
Asthma (J45-J46)									
All races, both sexes	4,487	8	32	43	86	78	89	105	144
Male	1,632	6	20	22	52	52	50	64	78
Female	2,855	2	12	21	34	26	39	41	66
White, both sexes	3,144	5	18	18	36	36	53	49	83
Male	1,057	3	9	8	20	24	31	29	41
Female	2,087	2	9	10	16	12	22	20	42
Black, both sexes	1,158	3	14	24	45	40	34	49	55
Male	481	3	11	14	28	27	17	30	34
Female	677	-	3	10	17	13	17	19	21
American Indian 1/, both sexes	31	-	-	-	-	1	1	4	-
Male	14	-	-	-	-	1	1	3	-
Female	17	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	154	-	-	1	5	1	1	3	6
Male	80	-	-	-	4	-	1	2	3
Female	74	-	-	1	1	1	-	1	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Pneumonitis due to solids and liquids (J69)									
All races, both sexes	16,636	10	11	8	5	12	25	22	40
Male	8,390	4	5	5	4	8	15	11	24
Female	8,246	6	6	3	1	4	10	11	16
White, both sexes	14,869	8	9	8	4	10	17	16	30
Male	7,488	4	4	5	3	7	11	7	15
Female	7,381	4	5	3	1	3	6	9	15
Black, both sexes	1,521	1	2	-	1	2	7	6	9
Male	765	-	1	-	1	1	3	4	8
Female	756	1	1	-	-	1	4	2	1
American Indian 1/, both sexes	49	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	197	1	-	-	-	-	1	-	-
Male	110	-	-	-	-	-	1	-	-
Female	87	1	-	-	-	-	-	-	-
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)									
All races, both sexes	25,535	273	87	36	38	60	71	99	154
Male	12,391	154	45	21	17	30	43	48	79
Female	13,144	119	42	15	21	30	28	51	75
White, both sexes	22,664	166	63	24	29	39	47	72	110
Male	10,979	100	30	14	14	17	29	35	58
Female	11,685	66	33	10	15	22	18	37	52
Black, both sexes	2,361	103	21	10	8	18	22	22	37
Male	1,135	51	14	7	3	11	12	12	18
Female	1,226	52	7	3	5	7	10	10	19
American Indian 1/, both sexes	157	1	2	-	1	1	-	3	4
Male	73	-	1	-	-	1	-	-	2
Female	84	1	1	-	1	-	-	3	2
Asian or Pacific Islander 2/, both sexes	353	3	1	2	-	2	2	2	3
Male	204	3	-	-	-	1	2	1	1
Female	149	-	1	2	-	1	-	1	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Cholelithiasis and other disorders of gallbladder (K80-K82)									
All races, both sexes	2,810	-	-	1	1	2	7	6	13
Male	1,244	-	-	-	1	1	3	2	5
Female	1,566	-	-	1	-	1	4	4	8
White, both sexes	2,457	-	-	-	1	1	6	3	7
Male	1,111	-	-	-	1	1	2	2	3
Female	1,346	-	-	-	-	-	4	1	4
Black, both sexes	280	-	-	1	-	1	1	3	5
Male	96	-	-	-	-	-	1	-	2
Female	184	-	-	1	-	1	-	3	3
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)									
All races, both sexes	37,251	162	14	11	14	32	56	80	164
Male	17,811	103	7	7	10	18	25	44	85
Female	19,440	59	7	4	4	14	31	36	79
White, both sexes	29,598	87	8	8	12	20	43	48	88
Male	14,385	58	5	6	8	11	22	28	45
Female	15,213	29	3	2	4	9	21	20	43
Black, both sexes	6,911	71	5	3	1	12	12	29	71
Male	3,074	42	2	1	1	7	3	13	37
Female	3,837	29	3	2	-	5	9	16	34
American Indian 1/, both sexes	215	1	-	-	-	-	-	2	1
Male	98	-	-	-	-	-	-	2	-
Female	117	1	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	527	3	1	-	1	-	1	1	4
Male	254	3	-	-	1	-	-	1	3
Female	273	-	1	-	-	-	1	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Inflammatory diseases of female pelvic organs (N70-N76)									
All races, both sexes	120	2	-	-	1	-	2	2	4
Female	120	2	-	-	1	-	2	2	4
White, both sexes	94	1	-	-	-	-	2	2	2
Female	94	1	-	-	-	-	2	2	2
Black, both sexes	23	1	-	-	1	-	-	-	1
Female	23	1	-	-	1	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
Pregnancy, childbirth and the puerperium (000-099)									
All races, both sexes	404	-	18	77	87	95
Female	404	-	18	77	87	95
White, both sexes	243	-	11	44	53	55
Female	243	-	11	44	53	55
Black, both sexes	142	-	4	32	31	34
Female	142	-	4	32	31	34
American Indian 1/, both sexes	6	-	1	1	1	3
Female	6	-	1	1	1	3
Asian or Pacific Islander 2/, both sexes	13	-	2	-	2	3
Female	13	-	2	-	2	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Pregnancy with abortive outcome (000-007)									
All races, both sexes	37	-	-	6	12	5
Female	37	-	-	6	12	5
White, both sexes	18	-	-	1	7	1
Female	18	-	-	1	7	1
Black, both sexes	19	-	-	5	5	4
Female	19	-	-	5	5	4
American Indian 1/, both sexes	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Female	-	-	-	-	-	-
Other complications of pregnancy, childbirth and the puerperium (010-099)									
All races, both sexes	367	-	18	71	75	90
Female	367	-	18	71	75	90
White, both sexes	225	-	11	43	46	54
Female	225	-	11	43	46	54
Black, both sexes	123	-	4	27	26	30
Female	123	-	4	27	26	30
American Indian 1/, both sexes	6	-	1	1	1	3
Female	6	-	1	1	1	3
Asian or Pacific Islander 2/, both sexes	13	-	2	-	2	3
Female	13	-	2	-	2	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Certain conditions originating in the perinatal period (P00-P96)									
All races, both sexes	14,069	13,918	79	25	16	10	5	5	2
Male	7,968	7,872	52	13	10	9	2	4	2
Female	6,101	6,046	27	12	6	1	3	1	-
White, both sexes	8,532	8,431	46	18	12	8	5	2	1
Male	4,828	4,762	30	10	10	8	2	1	1
Female	3,704	3,669	16	8	2	-	3	1	-
Black, both sexes	5,021	4,979	27	6	4	1	-	3	1
Male	2,844	2,818	18	3	-	1	-	3	1
Female	2,177	2,161	9	3	4	-	-	-	-
American Indian 1/, both sexes	124	121	2	-	-	1	-	-	-
Male	72	71	1	-	-	-	-	-	-
Female	52	50	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	392	387	4	1	-	-	-	-	-
Male	224	221	3	-	-	-	-	-	-
Female	168	166	1	1	-	-	-	-	-

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes	10,578	5,743	495	198	201	225	216	208	269
Male	5,517	3,028	259	105	102	137	131	120	159
Female	5,061	2,715	236	93	99	88	85	88	110
White, both sexes	8,396	4,371	347	147	145	159	173	163	228
Male	4,368	2,316	180	72	75	93	108	93	136
Female	4,028	2,055	167	75	70	66	65	70	92
Black, both sexes	1,790	1,102	126	44	47	56	35	38	39
Male	944	567	70	29	23	39	19	24	22
Female	846	535	56	15	24	17	16	14	17
American Indian 1/, both sexes	95	68	4	3	-	1	2	-	-
Male	56	42	1	2	-	-	2	-	-
Female	39	26	3	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	297	202	18	4	9	9	6	7	2
Male	149	103	8	2	4	5	2	3	1
Female	148	99	10	2	5	4	4	4	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)									
All races, both sexes	31,876	3,639	248	85	92	317	551	663	891
Male	15,863	2,156	137	48	51	196	385	456	604
Female	16,013	1,483	111	37	41	121	166	207	287
White, both sexes	26,285	2,294	182	57	63	240	416	495	690
Male	12,827	1,363	106	35	33	157	301	349	481
Female	13,458	931	76	22	30	83	115	146	209
Black, both sexes	5,004	1,212	57	21	25	67	119	142	174
Male	2,693	710	29	11	17	32	76	91	107
Female	2,311	502	28	10	8	35	43	51	67
American Indian 1/, both sexes	209	61	3	1	-	4	5	5	12
Male	127	42	1	-	-	3	3	2	5
Female	82	19	2	1	-	1	2	3	7
Asian or Pacific Islander 2/, both sexes	378	72	6	6	4	6	11	21	15
Male	216	41	1	2	1	4	5	14	11
Female	162	31	5	4	3	2	6	7	4

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes	178,378	1,105	609	376	503	732	1,032	1,257	1,840
Male	72,227	629	338	198	276	432	625	737	1,074
Female	106,151	476	271	178	227	300	407	520	766
White, both sexes	154,181	740	422	277	375	558	767	880	1,317
Male	61,390	409	238	152	217	340	482	551	803
Female	92,791	331	184	125	158	218	285	329	514
Black, both sexes	21,272	305	148	79	110	150	233	344	472
Male	9,455	187	78	35	52	80	124	165	239
Female	11,817	118	70	44	58	70	109	179	233
American Indian 1/, both sexes	945	17	17	2	5	7	9	18	28
Male	471	8	7	2	3	4	7	13	24
Female	474	9	10	-	2	3	2	5	4
Asian or Pacific Islander 2/, both sexes	1,980	43	22	18	13	17	23	15	23
Male	911	25	15	9	4	8	12	8	8
Female	1,069	18	7	9	9	9	11	7	15

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
All races, both sexes	97,900	881	1,826	1,391	1,588	6,755	7,358	5,898	5,871
Male	63,817	501	1,098	824	1,037	4,757	5,703	4,606	4,453
Female	34,083	380	728	567	551	1,998	1,655	1,292	1,418
White, both sexes	82,592	569	1,335	996	1,223	5,731	6,092	4,736	4,783
Male	53,329	331	808	587	791	4,007	4,763	3,704	3,665
Female	29,263	238	527	409	432	1,724	1,329	1,032	1,118
Black, both sexes	12,277	274	405	338	293	737	958	892	878
Male	8,531	149	239	209	205	542	719	693	647
Female	3,746	125	166	129	88	195	239	199	231
American Indian 1/, both sexes	1,353	20	39	31	39	149	150	123	116
Male	900	11	25	16	21	102	100	94	78
Female	453	9	14	15	18	47	50	29	38
Asian or Pacific Islander 2/, both sexes	1,678	18	47	26	33	138	158	147	94
Male	1,057	10	26	12	20	106	121	115	63
Female	621	8	21	14	13	32	37	32	31
Transport accidents (V01-V99,Y85)									
All races, both sexes	46,749	174	691	842	1,080	5,475	5,553	3,957	3,443
Male	32,166	94	391	464	674	3,693	4,197	3,016	2,554
Female	14,583	80	300	378	406	1,782	1,356	941	889
White, both sexes	39,088	121	513	646	873	4,713	4,566	3,121	2,769
Male	26,815	68	290	358	538	3,158	3,474	2,377	2,064
Female	12,273	53	223	288	335	1,555	1,092	744	705
Black, both sexes	5,839	40	141	157	160	539	744	634	535
Male	4,199	21	79	87	110	380	548	489	401
Female	1,640	19	62	70	50	159	196	145	134
American Indian 1/, both sexes	829	7	16	19	27	120	121	94	73
Male	537	2	10	9	14	80	82	69	48
Female	292	5	6	10	13	40	39	25	25
Asian or Pacific Islander 2/, both sexes	993	6	21	20	20	103	122	108	66
Male	615	3	12	10	12	75	93	81	41
Female	378	3	9	10	8	28	29	27	25

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
All races, both sexes	43,354	168	651	780	992	5,251	5,309	3,715	3,169
Male	29,451	90	368	429	614	3,527	3,989	2,804	2,331
Female	13,903	78	283	351	378	1,724	1,320	911	838
White, both sexes	36,160	115	484	593	796	4,530	4,351	2,908	2,540
Male	24,470	64	275	330	485	3,026	3,291	2,191	1,876
Female	11,690	51	209	263	311	1,504	1,060	717	664
Black, both sexes	5,499	40	134	149	153	513	721	614	497
Male	3,922	21	74	80	105	358	528	470	372
Female	1,577	19	60	69	48	155	193	144	125
American Indian 1/, both sexes	779	7	14	19	25	113	118	92	71
Male	500	2	9	9	14	75	79	67	47
Female	279	5	5	10	11	38	39	25	24
Asian or Pacific Islander 2/, both sexes	916	6	19	19	18	95	119	101	61
Male	559	3	10	10	10	68	91	76	36
Female	357	3	9	9	8	27	28	25	25

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, both sexes	1,492	2	31	39	59	128	115	94	118
Male	1,125	2	17	21	40	99	95	81	90
Female	367	-	14	18	19	29	20	13	28
White, both sexes	1,221	2	21	34	49	109	94	77	94
Male	924	2	10	18	34	83	78	67	72
Female	297	-	11	16	15	26	16	10	22
Black, both sexes	206	-	7	4	7	14	18	11	21
Male	158	-	5	3	5	11	15	10	15
Female	48	-	2	1	2	3	3	1	6
American Indian 1/, both sexes	28	-	1	-	2	4	1	2	1
Male	19	-	-	-	-	4	1	2	1
Female	9	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	2	1	1	1	2	4	2
Male	24	-	2	-	1	1	1	2	2
Female	13	-	-	1	-	-	1	2	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)									
All races, both sexes	1,903	4	9	23	29	96	129	148	156
Male	1,590	2	6	14	20	67	113	131	133
Female	313	2	3	9	9	29	16	17	23
White, both sexes	1,707	4	8	19	28	74	121	136	135
Male	1,421	2	5	10	19	49	105	119	116
Female	286	2	3	9	9	25	16	17	19
Black, both sexes	134	-	-	4	-	12	5	9	17
Male	119	-	-	4	-	11	5	9	14
Female	15	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	22	-	1	-	-	3	2	-	1
Male	18	-	1	-	-	1	2	-	-
Female	4	-	-	-	-	2	-	-	1
Asian or Pacific Islander 2/, both sexes	40	-	-	-	1	7	1	3	3
Male	32	-	-	-	1	6	1	3	3
Female	8	-	-	-	-	1	-	-	-
Nontransport accidents (W00-X59,Y86)									
All races, both sexes	51,151	707	1,135	549	508	1,280	1,805	1,941	2,428
Male	31,651	407	707	360	363	1,064	1,506	1,590	1,899
Female	19,500	300	428	189	145	216	299	351	529
White, both sexes	43,504	448	822	350	350	1,018	1,526	1,615	2,014
Male	26,514	263	518	229	253	849	1,289	1,327	1,601
Female	16,990	185	304	121	97	169	237	288	413
Black, both sexes	6,438	234	264	181	133	198	214	258	343
Male	4,332	128	160	122	95	162	171	204	246
Female	2,106	106	104	59	38	36	43	54	97
American Indian 1/, both sexes	524	13	23	12	12	29	29	29	43
Male	363	9	15	7	7	22	18	25	30
Female	161	4	8	5	5	7	11	4	13
Asian or Pacific Islander 2/, both sexes	685	12	26	6	13	35	36	39	28
Male	442	7	14	2	8	31	28	34	22
Female	243	5	12	4	5	4	8	5	6

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Falls (W00-W19)									
All races, both sexes	13,322	8	36	16	21	99	138	138	165
Male	7,122	6	23	14	16	88	115	122	150
Female	6,200	2	13	2	5	11	23	16	15
White, both sexes	12,252	5	23	12	18	90	127	125	140
Male	6,449	3	13	10	15	81	106	112	127
Female	5,803	2	10	2	3	9	21	13	13
Black, both sexes	760	2	13	4	3	6	7	5	18
Male	485	2	10	4	1	4	7	3	16
Female	275	-	3	-	2	2	-	2	2
American Indian 1/, both sexes	81	1	-	-	-	1	-	6	3
Male	57	1	-	-	-	1	-	5	3
Female	24	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	229	-	-	-	-	2	4	2	4
Male	131	-	-	-	-	2	2	2	4
Female	98	-	-	-	-	-	2	-	-
Accidental discharge of firearms (W32-W34)									
All races, both sexes	776	1	18	18	49	107	95	74	57
Male	671	-	11	16	46	97	86	65	44
Female	105	1	7	2	3	10	9	9	13
White, both sexes	611	1	11	11	41	76	65	52	42
Male	530	-	6	10	38	70	59	44	31
Female	81	1	5	1	3	6	6	8	11
Black, both sexes	137	-	5	5	7	28	24	18	14
Male	117	-	3	5	7	25	21	17	12
Female	20	-	2	-	-	3	3	1	2
American Indian 1/, both sexes	14	-	2	1	1	3	3	1	-
Male	12	-	2	1	1	2	3	1	-
Female	2	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	1	-	-	3	3	1
Male	12	-	-	-	-	-	3	3	1
Female	2	-	-	1	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Accidental drowning and submersion (W65-W74)									
All races, both sexes	3,482	75	493	201	174	371	275	213	206
Male	2,735	43	316	137	130	339	244	184	180
Female	747	32	177	64	44	32	31	29	26
White, both sexes	2,683	47	415	125	93	253	207	162	160
Male	2,105	28	268	84	67	231	182	141	143
Female	578	19	147	41	26	22	25	21	17
Black, both sexes	593	26	56	67	69	86	48	34	24
Male	478	14	35	48	54	81	45	29	23
Female	115	12	21	19	15	5	3	5	1
American Indian 1/, both sexes	79	2	10	7	4	12	6	2	12
Male	61	1	6	4	4	11	4	2	7
Female	18	1	4	3	-	1	2	-	5
Asian or Pacific Islander 2/, both sexes	127	-	12	2	8	20	14	15	10
Male	91	-	7	1	5	16	13	12	7
Female	36	-	5	1	3	4	1	3	3
Accidental exposure to smoke, fire and flames (X00-X09)									
All races, both sexes	3,377	37	290	182	84	79	113	120	121
Male	2,025	20	174	109	36	44	77	78	79
Female	1,352	17	116	73	48	35	36	42	42
White, both sexes	2,376	23	160	104	54	60	81	83	86
Male	1,435	11	98	61	24	32	58	54	59
Female	941	12	62	43	30	28	23	29	27
Black, both sexes	930	14	121	74	27	16	28	31	34
Male	551	9	71	46	11	10	17	19	20
Female	379	5	50	28	16	6	11	12	14
American Indian 1/, both sexes	37	-	5	2	2	3	1	6	1
Male	23	-	3	1	1	2	1	5	-
Female	14	-	2	1	1	1	-	1	1
Asian or Pacific Islander 2/, both sexes	34	-	4	2	1	-	3	-	-
Male	16	-	2	1	-	-	1	-	-
Female	18	-	2	1	1	-	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Accidental poisoning and exposure to noxious substances (X40-X49)									
All races, both sexes	12,757	14	32	17	28	351	809	980	1,400
Male	9,138	11	21	12	15	273	647	779	1,041
Female	3,619	3	11	5	13	78	162	201	359
White, both sexes	10,508	10	19	10	20	318	732	847	1,184
Male	7,558	8	11	7	11	251	601	674	894
Female	2,950	2	8	3	9	67	131	173	290
Black, both sexes	2,028	4	12	6	3	21	58	115	188
Male	1,433	3	9	5	2	13	37	91	125
Female	595	1	3	1	1	8	21	24	63
American Indian 1/, both sexes	137	-	-	1	3	6	12	11	19
Male	84	-	-	-	-	3	4	9	15
Female	53	-	-	1	3	3	8	2	4
Asian or Pacific Islander 2/, both sexes	84	-	1	-	2	6	7	7	9
Male	63	-	1	-	2	6	5	5	7
Female	21	-	-	-	-	-	2	2	2

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races, both sexes	17,437	572	266	115	152	273	375	416	479
Male	9,960	327	162	72	120	223	337	362	405
Female	7,477	245	104	43	32	50	38	54	74
White, both sexes	15,074	362	194	88	124	221	314	346	402
Male	8,437	213	122	57	98	184	283	302	347
Female	6,637	149	72	31	26	37	31	44	55
Black, both sexes	1,990	188	57	25	24	41	49	55	65
Male	1,268	100	32	14	20	29	44	45	50
Female	722	88	25	11	4	12	5	10	15
American Indian 1/, both sexes	176	10	6	1	2	4	7	3	8
Male	126	7	4	1	1	3	6	3	5
Female	50	3	2	-	1	1	1	-	3
Asian or Pacific Islander 2/, both sexes	197	12	9	1	2	7	5	12	4
Male	129	7	4	-	1	7	4	12	3
Female	68	5	5	1	1	-	1	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Intentional self-harm (suicide) (X60-X84,Y87.0)									
All races, both sexes	29,350	7	300	1,621	2,373	2,341	2,451
Male	23,618	6	238	1,351	2,073	1,956	1,982
Female	5,732	1	62	270	300	385	469
White, both sexes	26,475	6	248	1,354	1,961	1,941	2,143
Male	21,293	5	195	1,130	1,723	1,631	1,732
Female	5,182	1	53	224	238	310	411
Black, both sexes	1,962	-	40	172	308	266	210
Male	1,636	-	35	150	265	224	178
Female	326	-	5	22	43	42	32
American Indian 1/, both sexes	297	1	6	44	45	43	35
Male	237	1	5	34	37	33	26
Female	60	-	1	10	8	10	9
Asian or Pacific Islander 2/, both sexes	616	-	6	51	59	91	63
Male	452	-	3	37	48	68	46
Female	164	-	3	14	11	23	17

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes	16,586	-	110	897	1,370	1,223	1,230
Male	14,454	-	90	793	1,250	1,084	1,039
Female	2,132	-	20	104	120	139	191
White, both sexes	15,174	-	103	741	1,137	1,012	1,086
Male	13,214	-	86	653	1,040	893	922
Female	1,960	-	17	88	97	119	164
Black, both sexes	1,090	-	4	117	191	161	111
Male	965	-	4	109	172	145	95
Female	125	-	-	8	19	16	16
American Indian 1/, both sexes	133	-	1	20	23	14	13
Male	109	-	-	17	20	12	8
Female	24	-	1	3	3	2	5
Asian or Pacific Islander 2/, both sexes	189	-	2	19	19	36	20
Male	166	-	-	14	18	34	14
Female	23	-	2	5	1	2	6

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)									
All races, both sexes	12,764	7	190	724	1,003	1,118	1,221
Male	9,164	6	148	558	823	872	943
Female	3,600	1	42	166	180	246	278
White, both sexes	11,301	6	145	613	824	929	1,057
Male	8,079	5	109	477	683	738	810
Female	3,222	1	36	136	141	191	247
Black, both sexes	872	-	36	55	117	105	99
Male	671	-	31	41	93	79	83
Female	201	-	5	14	24	26	16
American Indian 1/, both sexes	164	1	5	24	22	29	22
Male	128	1	5	17	17	21	18
Female	36	-	-	7	5	8	4
Asian or Pacific Islander 2/, both sexes	427	-	4	32	40	55	43
Male	286	-	3	23	30	34	32
Female	141	-	1	9	10	21	11
Assault (homicide) (X85-Y09,Y87.1)									
All races, both sexes	16,765	349	356	140	231	1,914	3,025	2,312	1,852
Male	12,820	203	198	77	150	1,613	2,590	1,918	1,443
Female	3,945	146	158	63	81	301	435	394	409
White, both sexes	8,339	197	199	98	142	823	1,158	945	869
Male	5,925	125	114	59	89	659	924	721	635
Female	2,414	72	85	39	53	164	234	224	234
Black, both sexes	7,867	140	145	40	84	1,028	1,791	1,301	923
Male	6,482	73	75	16	56	899	1,602	1,146	765
Female	1,385	67	70	24	28	129	189	155	158
American Indian 1/, both sexes	203	5	8	1	-	25	26	28	31
Male	159	3	6	1	-	21	25	22	24
Female	44	2	2	-	-	4	1	6	7
Asian or Pacific Islander 2/, both sexes	356	7	4	1	5	38	50	38	29
Male	254	2	3	1	5	34	39	29	19
Female	102	5	1	-	-	4	11	9	10

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Assault (homicide) by discharge of firearms (X93-X95)									
All races, both sexes	10,801	12	28	50	137	1,549	2,414	1,789	1,285
Male	9,006	9	17	28	110	1,381	2,167	1,574	1,076
Female	1,795	3	11	22	27	168	247	215	209
White, both sexes	4,806	5	17	39	75	617	817	645	547
Male	3,686	2	9	22	59	531	693	528	427
Female	1,120	3	8	17	16	86	124	117	120
Black, both sexes	5,699	7	8	11	58	888	1,546	1,097	715
Male	5,084	7	5	6	47	812	1,431	1,009	628
Female	615	-	3	5	11	76	115	88	87
American Indian 1/, both sexes	86	-	2	-	-	14	18	17	10
Male	70	-	2	-	-	12	17	13	9
Female	16	-	-	-	-	2	1	4	1
Asian or Pacific Islander 2/, both sexes	210	-	1	-	4	30	33	30	13
Male	166	-	1	-	4	26	26	24	12
Female	44	-	-	-	-	4	7	6	1

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races, both sexes	5,964	337	328	90	94	365	611	523	567
Male	3,814	194	181	49	40	232	423	344	367
Female	2,150	143	147	41	54	133	188	179	200
White, both sexes	3,533	192	182	59	67	206	341	300	322
Male	2,239	123	105	37	30	128	231	193	208
Female	1,294	69	77	22	37	78	110	107	114
Black, both sexes	2,168	133	137	29	26	140	245	204	208
Male	1,398	66	70	10	9	87	171	137	137
Female	770	67	67	19	17	53	74	67	71
American Indian 1/, both sexes	117	5	6	1	-	11	8	11	21
Male	89	3	4	1	-	9	8	9	15
Female	28	2	2	-	-	2	-	2	6
Asian or Pacific Islander 2/, both sexes	146	7	3	1	1	8	17	8	16
Male	88	2	2	1	1	8	13	5	7
Female	58	5	1	-	-	-	4	3	9

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Legal intervention (Y35,Y89.0)									
All races, both sexes	359	-	-	-	3	27	46	59	51
Male	348	-	-	-	3	27	44	58	50
Female	11	-	-	-	-	-	2	1	1
White, both sexes	237	-	-	-	1	11	33	36	36
Male	231	-	-	-	1	11	31	35	35
Female	6	-	-	-	-	-	2	1	1
Black, both sexes	111	-	-	-	2	15	13	19	14
Male	108	-	-	-	2	15	13	19	14
Female	3	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	3	1
Male	6	-	-	-	-	-	-	3	1
Female	1	-	-	-	-	-	-	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)									
All races, both sexes	3,819	65	39	18	42	120	250	306	373
Male	2,635	27	26	13	37	93	192	223	266
Female	1,184	38	13	5	5	27	58	83	107
White, both sexes	3,053	41	27	9	29	103	199	252	301
Male	2,078	18	17	6	26	79	149	181	211
Female	975	23	10	3	3	24	50	71	90
Black, both sexes	668	15	11	9	12	13	39	45	60
Male	488	3	8	7	10	13	34	35	45
Female	180	12	3	2	2	-	5	10	15
American Indian 1/, both sexes	46	4	1	-	1	1	4	3	6
Male	36	3	1	-	1	-	3	2	6
Female	10	1	-	-	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	52	5	-	-	-	3	8	6	6
Male	33	3	-	-	-	1	6	5	4
Female	19	2	-	-	-	2	2	1	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Discharge of firearms, undetermined intent (Y22-Y24)									
All races, both sexes	230	-	-	2	8	26	45	32	20
Male	190	-	-	2	7	25	37	26	17
Female	40	-	-	-	1	1	8	6	3
White, both sexes	167	-	-	-	5	20	28	23	11
Male	137	-	-	-	5	19	23	18	10
Female	30	-	-	-	-	1	5	5	1
Black, both sexes	54	-	-	2	3	6	14	8	7
Male	47	-	-	2	2	6	13	7	5
Female	7	-	-	-	1	-	1	1	2
American Indian 1/, both sexes	4	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	1	-	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)									
All races, both sexes	3,589	65	39	16	34	94	205	274	353
Male	2,445	27	26	11	30	68	155	197	249
Female	1,144	38	13	5	4	26	50	77	104
White, both sexes	2,886	41	27	9	24	83	171	229	290
Male	1,941	18	17	6	21	60	126	163	201
Female	945	23	10	3	3	23	45	66	89
Black, both sexes	614	15	11	7	9	7	25	37	53
Male	441	3	8	5	8	7	21	28	40
Female	173	12	3	2	1	-	4	9	13
American Indian 1/, both sexes	42	4	1	-	1	1	2	3	5
Male	33	3	1	-	1	-	2	2	5
Female	9	1	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	47	5	-	-	-	3	7	5	5
Male	30	3	-	-	-	1	6	4	3
Female	17	2	-	-	-	2	1	1	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS
Operations of war and their sequelae (Y36,Y89.1)									
All races, both sexes	16	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Complications of medical and surgical care (Y40-Y84,Y88)									
All races, both sexes	3,059	19	22	18	13	10	22	37	48
Male	1,430	12	11	11	4	5	11	23	24
Female	1,629	7	11	7	9	5	11	14	24
White, both sexes	2,520	14	12	12	11	8	16	22	38
Male	1,210	11	8	7	4	4	9	13	20
Female	1,310	3	4	5	7	4	7	9	18
Black, both sexes	487	5	8	4	1	2	5	13	10
Male	193	1	2	2	-	1	1	9	4
Female	294	4	6	2	1	1	4	4	6
American Indian 1/, both sexes	14	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	-	-
Female	9	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	2	2	-	-	1	2	-
Male	22	-	1	2	-	-	1	1	-
Female	16	-	1	-	-	-	-	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Tuberculosis (A16-A19)

All races, both sexes	17	25	38	53	31	50	70	85	132
Male	11	17	35	41	23	36	46	41	69
Female	6	8	3	12	8	14	24	44	63
White, both sexes	8	5	13	31	8	25	38	49	79
Male	8	3	12	22	6	18	26	27	40
Female	-	2	1	9	2	7	12	22	39
Black, both sexes	8	17	21	21	18	16	25	27	29
Male	2	12	19	18	13	13	17	11	17
Female	6	5	2	3	5	3	8	16	12
American Indian 1/, both sexes	-	-	2	-	3	3	4	1	4
Male	-	-	2	-	2	2	2	1	2
Female	-	-	-	-	1	1	2	-	2
Asian or Pacific Islander 2/, both sexes	1	3	2	1	2	6	3	8	20
Male	1	2	2	1	2	3	1	2	10
Female	-	1	-	-	-	3	2	6	10

Respiratory tuberculosis (A16)

All races, both sexes	15	20	30	42	24	36	52	67	107
Male	11	15	29	35	21	29	38	35	58
Female	4	5	1	7	3	7	14	32	49
White, both sexes	8	3	9	26	7	19	27	39	65
Male	8	3	9	21	5	15	20	23	36
Female	-	-	-	5	2	4	7	16	29
Black, both sexes	6	14	17	15	12	10	21	22	24
Male	2	10	16	13	12	10	16	10	13
Female	4	4	1	2	-	-	5	12	11
American Indian 1/, both sexes	-	-	2	-	3	2	2	-	3
Male	-	-	2	-	2	2	1	-	1
Female	-	-	-	-	1	-	1	-	2
Asian or Pacific Islander 2/, both sexes	1	3	2	1	2	5	2	6	15
Male	1	2	2	1	2	2	1	2	8
Female	-	1	-	-	-	3	1	4	7

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Measles (B05)

All races, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-

Viral hepatitis (B15-B19)

All races, both sexes	219	560	1,044	850	492	412	410	435	401
Male	151	389	787	638	338	239	202	194	198
Female	68	171	257	212	154	173	208	241	203
White, both sexes	180	445	827	635	350	315	320	358	350
Male	123	316	636	487	246	186	156	161	175
Female	57	129	191	148	104	129	164	197	175
Black, both sexes	30	88	172	164	93	50	47	47	32
Male	22	56	121	120	63	25	26	22	13
Female	8	32	51	44	30	25	21	25	19
American Indian 1/, both sexes	3	12	12	13	4	6	4	1	1
Male	2	6	7	9	1	4	1	-	-
Female	1	6	5	4	3	2	3	1	1
Asian or Pacific Islander 2/, both sexes	6	15	33	38	45	41	39	29	18
Male	4	11	23	22	28	24	19	11	10
Female	2	4	10	16	17	17	20	18	8

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79
YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races, both sexes	162	188	277	308	323	384	451	623	739
Male	105	115	164	194	179	204	253	344	367
Female	57	73	113	114	144	180	198	279	372
White, both sexes	104	138	212	232	263	297	380	533	655
Male	68	85	127	147	150	161	211	300	318
Female	36	53	85	85	113	136	169	233	337
Black, both sexes	48	47	62	65	54	65	55	74	65
Male	30	28	35	41	27	35	33	37	36
Female	18	19	27	24	27	30	22	37	29
American Indian 1/, both sexes	5	-	1	4	3	9	1	2	2
Male	2	-	1	4	1	2	-	-	1
Female	3	-	-	-	2	7	1	2	1
Asian or Pacific Islander 2/, both sexes	5	3	2	7	3	13	15	14	17
Male	5	2	1	2	1	6	9	7	12
Female	-	1	1	5	2	7	6	7	5

Malignant neoplasms (C00-C97)

All races, both sexes	5,565	10,955	18,803	29,231	38,930	50,075	65,843	84,288	90,655
Male	2,402	4,942	9,243	14,972	20,912	27,515	36,730	46,461	48,284
Female	3,163	6,013	9,560	14,259	18,018	22,560	29,113	37,827	42,371
White, both sexes	4,316	8,448	14,743	23,735	32,440	42,319	56,673	73,897	80,805
Male	1,907	3,832	7,292	12,061	17,275	23,182	31,634	40,650	42,849
Female	2,409	4,616	7,451	11,674	15,165	19,137	25,039	33,247	37,956
Black, both sexes	1,017	2,091	3,437	4,710	5,499	6,620	7,797	8,839	8,338
Male	383	926	1,664	2,528	3,114	3,715	4,371	4,950	4,584
Female	634	1,165	1,773	2,182	2,385	2,905	3,426	3,889	3,754
American Indian 1/, both sexes	34	71	88	132	190	227	251	288	248
Male	16	33	44	72	106	120	136	155	133
Female	18	38	44	60	84	107	115	133	115
Asian or Pacific Islander 2/, both sexes	198	345	535	654	801	909	1,122	1,264	1,264
Male	96	151	243	311	417	498	589	706	718
Female	102	194	292	343	384	411	533	558	546

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races, both sexes	88	205	408	660	770	872	942	1,017	933
Male	60	155	309	516	618	652	667	659	569
Female	28	50	99	144	152	220	275	358	364
White, both sexes	60	138	291	475	581	674	776	868	832
Male	38	100	222	358	467	499	541	555	495
Female	22	38	69	117	114	175	235	313	337
Black, both sexes	15	46	96	156	164	166	135	121	82
Male	11	36	69	137	133	132	106	87	61
Female	4	10	27	19	31	34	29	34	21
American Indian 1/, both sexes	2	2	-	4	6	4	7	10	3
Male	2	2	-	3	4	1	5	7	2
Female	-	-	-	1	2	3	2	3	1
Asian or Pacific Islander 2/, both sexes	11	19	21	25	19	28	24	18	16
Male	9	17	18	18	14	20	15	10	11
Female	2	2	3	7	5	8	9	8	5

Malignant neoplasm of esophagus (C15)

All races, both sexes	78	219	478	870	1,180	1,430	1,711	1,960	1,817
Male	69	196	415	746	988	1,181	1,349	1,482	1,337
Female	9	23	63	124	192	249	362	478	480
White, both sexes	74	175	377	674	932	1,150	1,441	1,705	1,599
Male	67	162	341	588	791	975	1,158	1,308	1,186
Female	7	13	36	86	141	175	283	397	413
Black, both sexes	3	41	96	184	228	255	244	222	203
Male	1	31	69	150	185	186	172	149	139
Female	2	10	27	34	43	69	72	73	64
American Indian 1/, both sexes	-	2	3	3	3	8	6	7	3
Male	-	2	3	3	1	8	5	7	2
Female	-	-	-	-	2	-	1	-	1
Asian or Pacific Islander 2/, both sexes	1	1	2	9	17	17	20	26	12
Male	1	1	2	5	11	12	14	18	10
Female	-	-	-	4	6	5	6	8	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79
YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Malignant neoplasm of stomach (C16)

All races, both sexes	167	317	468	629	753	1,051	1,317	1,737	2,009
Male	100	184	290	413	506	722	860	1,106	1,201
Female	67	133	178	216	247	329	457	631	808
White, both sexes	113	205	314	463	561	791	988	1,348	1,571
Male	66	115	192	311	379	557	660	874	952
Female	47	90	122	152	182	234	328	474	619
Black, both sexes	37	81	110	130	149	189	243	286	320
Male	26	52	72	83	100	125	148	171	176
Female	11	29	38	47	49	64	95	115	144
American Indian 1/, both sexes	1	3	-	5	7	9	15	10	9
Male	-	3	-	4	5	4	8	5	6
Female	1	-	-	1	2	5	7	5	3
Asian or Pacific Islander 2/, both sexes	16	28	44	31	36	62	71	93	109
Male	8	14	26	15	22	36	44	56	67
Female	8	14	18	16	14	26	27	37	42

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races, both sexes	480	889	1,656	2,581	3,397	4,543	5,913	8,188	9,410
Male	271	488	893	1,466	1,989	2,619	3,407	4,517	4,852
Female	209	401	763	1,115	1,408	1,924	2,506	3,671	4,558
White, both sexes	349	666	1,250	2,050	2,737	3,752	4,991	7,094	8,322
Male	204	363	692	1,182	1,607	2,181	2,906	3,955	4,332
Female	145	303	558	868	1,130	1,571	2,085	3,139	3,990
Black, both sexes	109	184	359	464	538	687	787	945	942
Male	56	105	180	251	312	369	420	479	442
Female	53	79	179	213	226	318	367	466	500
American Indian 1/, both sexes	2	11	7	11	22	20	22	29	27
Male	1	7	4	5	17	13	14	16	15
Female	1	4	3	6	5	7	8	13	12
Asian or Pacific Islander 2/, both sexes	20	28	40	56	100	84	113	120	119
Male	10	13	17	28	53	56	67	67	63
Female	10	15	23	28	47	28	46	53	56

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races, both sexes	113	316	742	944	1,040	1,216	1,526	1,913	2,011
Male	74	223	595	729	761	846	1,039	1,207	1,196
Female	39	93	147	215	279	370	487	706	815
White, both sexes	65	196	510	654	756	924	1,204	1,580	1,717
Male	37	142	401	500	542	641	827	1,006	1,048
Female	28	54	109	154	214	283	377	574	669
Black, both sexes	25	77	166	199	171	173	189	202	162
Male	16	50	138	152	134	122	131	124	77
Female	9	27	28	47	37	51	58	78	85
American Indian 1/, both sexes	1	3	7	13	14	9	11	10	6
Male	1	3	7	10	10	5	8	8	3
Female	-	-	-	3	4	4	3	2	3
Asian or Pacific Islander 2/, both sexes	22	40	59	78	99	110	122	121	126
Male	20	28	49	67	75	78	73	69	68
Female	2	12	10	11	24	32	49	52	58

Malignant neoplasm of pancreas (C25)

All races, both sexes	140	408	834	1,528	2,030	2,678	3,546	4,557	5,223
Male	87	237	531	955	1,206	1,543	1,965	2,224	2,447
Female	53	171	303	573	824	1,135	1,581	2,333	2,776
White, both sexes	110	319	679	1,248	1,675	2,250	2,999	3,945	4,603
Male	69	189	437	787	1,003	1,314	1,709	1,949	2,197
Female	41	130	242	461	672	936	1,290	1,996	2,406
Black, both sexes	24	73	136	246	307	365	455	523	528
Male	15	39	80	151	178	195	214	236	215
Female	9	34	56	95	129	170	241	287	313
American Indian 1/, both sexes	2	6	1	8	16	9	10	14	8
Male	-	5	1	4	10	6	5	7	2
Female	2	1	-	4	6	3	5	7	6
Asian or Pacific Islander 2/, both sexes	4	10	18	26	32	54	82	75	84
Male	3	4	13	13	15	28	37	32	33
Female	1	6	5	13	17	26	45	43	51

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Malignant neoplasm of larynx (C32)									
All races, both sexes	9	55	156	270	384	489	574	645	595
Male	6	43	122	218	319	407	441	504	451
Female	3	12	34	52	65	82	133	141	144
White, both sexes	8	41	108	191	279	364	466	521	501
Male	5	30	86	151	228	302	356	407	383
Female	3	11	22	40	51	62	110	114	118
Black, both sexes	1	14	47	76	101	118	96	115	88
Male	1	13	35	65	87	100	76	90	62
Female	-	1	12	11	14	18	20	25	26
American Indian 1/, both sexes	-	-	-	1	1	5	7	1	1
Male	-	-	-	1	1	3	5	-	1
Female	-	-	-	-	-	2	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	3	2	5	8	5
Male	-	-	1	1	3	2	4	7	5
Female	-	-	-	1	-	-	1	1	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
All races, both sexes	664	2,092	4,220	7,695	12,453	17,268	23,438	28,820	27,305
Male	370	1,172	2,540	4,583	7,546	10,391	14,189	17,193	15,726
Female	294	920	1,680	3,112	4,907	6,877	9,249	11,627	11,579
White, both sexes	499	1,594	3,265	6,272	10,562	14,925	20,532	25,676	24,672
Male	277	859	1,953	3,641	6,274	8,905	12,313	15,221	14,059
Female	222	735	1,312	2,631	4,288	6,020	8,219	10,455	10,613
Black, both sexes	142	442	837	1,295	1,660	2,066	2,534	2,692	2,222
Male	80	282	520	864	1,125	1,312	1,661	1,680	1,390
Female	62	160	317	431	535	754	873	1,012	832
American Indian 1/, both sexes	3	8	17	28	56	70	78	87	68
Male	1	4	9	15	31	45	40	43	44
Female	2	4	8	13	25	25	38	44	24
Asian or Pacific Islander 2/, both sexes	20	48	101	100	175	207	294	365	343
Male	12	27	58	63	116	129	175	249	233
Female	8	21	43	37	59	78	119	116	110

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Malignant melanoma of skin (C43)									
All races, both sexes	234	392	534	665	627	676	753	884	887
Male	118	237	321	448	420	446	495	572	559
Female	116	155	213	217	207	230	258	312	328
White, both sexes	229	377	525	654	611	662	736	862	871
Male	116	229	319	441	411	440	484	563	552
Female	113	148	206	213	200	222	252	299	319
Black, both sexes	4	11	6	7	11	9	11	19	14
Male	2	7	1	5	6	5	7	7	6
Female	2	4	5	2	5	4	4	12	8
American Indian 1/, both sexes	-	-	1	1	2	1	1	-	1
Male	-	-	-	1	1	1	1	-	-
Female	-	-	1	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	4	2	3	3	4	5	3	1
Male	-	1	1	1	2	-	3	2	1
Female	1	3	1	2	1	4	2	1	-
Malignant neoplasm of breast (C50)									
All races, both sexes	982	1,859	2,730	3,655	3,790	3,783	4,135	4,889	5,256
Male	12	7	18	28	44	41	47	68	65
Female	970	1,852	2,712	3,627	3,746	3,742	4,088	4,821	5,191
White, both sexes	717	1,380	2,066	2,934	3,100	3,182	3,584	4,341	4,735
Male	8	3	13	24	34	33	43	60	60
Female	709	1,377	2,053	2,910	3,066	3,149	3,541	4,281	4,675
Black, both sexes	228	417	564	614	592	518	477	492	474
Male	4	4	4	4	8	8	3	8	5
Female	224	413	560	610	584	510	474	484	469
American Indian 1/, both sexes	3	13	11	14	20	19	14	9	11
Male	-	-	-	-	1	-	-	-	-
Female	3	13	11	14	19	19	14	9	11
Asian or Pacific Islander 2/, both sexes	34	49	89	93	78	64	60	47	36
Male	-	-	1	-	1	-	1	-	-
Female	34	49	88	93	77	64	59	47	36

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

All races, both sexes	294	400	448	467	416	332	342	344	320
Female	294	400	448	467	416	332	342	344	320
White, both sexes	223	288	334	352	314	243	243	255	253
Female	223	288	334	352	314	243	243	255	253
Black, both sexes	62	88	88	92	83	70	85	73	61
Female	62	88	88	92	83	70	85	73	61
American Indian 1/, both sexes	2	5	3	6	2	3	1	2	2
Female	2	5	3	6	2	3	1	2	2
Asian or Pacific Islander 2/, both sexes	7	19	23	17	17	16	13	14	4
Female	7	19	23	17	17	16	13	14	4

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races, both sexes	41	97	190	317	478	644	800	981	1,039
Female	41	97	190	317	478	644	800	981	1,039
White, both sexes	33	73	158	262	394	496	602	771	866
Female	33	73	158	262	394	496	602	771	866
Black, both sexes	6	16	24	44	71	131	177	198	156
Female	6	16	24	44	71	131	177	198	156
American Indian 1/, both sexes	-	1	1	-	-	2	5	3	4
Female	-	1	1	-	-	2	5	3	4
Asian or Pacific Islander 2/, both sexes	2	7	7	11	13	15	16	9	13
Female	2	7	7	11	13	15	16	9	13

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of ovary (C56)

All races, both sexes	145	327	617	966	1,123	1,321	1,669	1,932	2,295
Female	145	327	617	966	1,123	1,321	1,669	1,932	2,295
White, both sexes	120	277	541	838	1,008	1,179	1,489	1,752	2,130
Female	120	277	541	838	1,008	1,179	1,489	1,752	2,130
Black, both sexes	17	43	51	89	86	119	152	147	147
Female	17	43	51	89	86	119	152	147	147
American Indian 1/, both sexes	1	-	2	1	3	4	2	3	6
Female	1	-	2	1	3	4	2	3	6
Asian or Pacific Islander 2/, both sexes	7	7	23	38	26	19	26	30	12
Female	7	7	23	38	26	19	26	30	12

Malignant neoplasm of prostate (C61)

All races, both sexes	6	19	108	263	623	1,306	2,466	4,342	6,314
Male	6	19	108	263	623	1,306	2,466	4,342	6,314
White, both sexes	4	12	70	192	444	960	1,851	3,380	5,167
Male	4	12	70	192	444	960	1,851	3,380	5,167
Black, both sexes	2	7	35	67	174	323	586	905	1,070
Male	2	7	35	67	174	323	586	905	1,070
American Indian 1/, both sexes	-	-	2	2	1	5	12	18	18
Male	-	-	2	2	1	5	12	18	18
Asian or Pacific Islander 2/, both sexes	-	-	1	2	4	18	17	39	59
Male	-	-	1	2	4	18	17	39	59

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races, both sexes	83	251	450	789	1,026	1,169	1,432	1,714	1,819
Male	53	175	309	571	727	803	942	1,068	1,054
Female	30	76	141	218	299	366	490	646	765
White, both sexes	59	211	381	685	903	1,031	1,271	1,542	1,657
Male	42	150	266	495	637	716	829	966	965
Female	17	61	115	190	266	315	442	576	692
Black, both sexes	21	31	58	87	100	114	134	139	132
Male	9	18	35	61	69	74	96	84	68
Female	12	13	23	26	31	40	38	55	64
American Indian 1/, both sexes	1	4	6	6	5	5	7	12	9
Male	1	3	3	5	5	3	5	6	7
Female	-	1	3	1	-	2	2	6	2
Asian or Pacific Islander 2/, both sexes	2	5	5	11	18	19	20	21	21
Male	1	4	5	10	16	10	12	12	14
Female	1	1	-	1	2	9	8	9	7

Malignant neoplasm of bladder (C67)

All races, both sexes	40	71	160	280	486	737	1,119	1,736	2,155
Male	21	41	110	212	369	547	838	1,256	1,542
Female	19	30	50	68	117	190	281	480	613
White, both sexes	36	63	137	254	439	670	1,020	1,585	1,977
Male	20	37	96	195	335	508	779	1,170	1,447
Female	16	26	41	59	104	162	241	415	530
Black, both sexes	3	7	20	24	41	59	86	134	160
Male	-	4	11	16	28	35	48	72	87
Female	3	3	9	8	13	24	38	62	73
American Indian 1/, both sexes	1	-	-	1	1	1	2	1	5
Male	1	-	-	-	1	1	2	1	1
Female	-	-	-	1	-	-	-	-	4
Asian or Pacific Islander 2/, both sexes	-	1	3	1	5	7	11	16	13
Male	-	-	3	1	5	3	9	13	7
Female	-	1	-	-	-	4	2	3	6

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, both sexes	454	642	815	1,109	1,110	1,227	1,409	1,623	1,437
Male	293	411	481	667	660	719	789	889	705
Female	161	231	334	442	450	508	620	734	732
White, both sexes	405	564	745	1,026	1,019	1,142	1,321	1,543	1,362
Male	264	366	449	609	611	673	747	846	669
Female	141	198	296	417	408	469	574	697	693
Black, both sexes	34	57	52	70	73	66	68	66	58
Male	19	35	24	49	40	37	30	37	25
Female	15	22	28	21	33	29	38	29	33
American Indian 1/, both sexes	7	4	4	1	2	1	3	1	6
Male	4	1	1	1	1	-	2	-	4
Female	3	3	3	-	1	1	1	1	2
Asian or Pacific Islander 2/, both sexes	8	17	14	12	16	18	17	13	11
Male	6	9	7	8	8	9	10	6	7
Female	2	8	7	4	8	9	7	7	4

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races, both sexes	776	1,077	1,675	2,350	3,156	4,128	5,932	8,109	9,743
Male	438	637	1,015	1,384	1,850	2,355	3,445	4,593	5,168
Female	338	440	660	966	1,306	1,773	2,487	3,516	4,575
White, both sexes	617	868	1,340	1,957	2,693	3,562	5,251	7,307	8,944
Male	354	536	827	1,150	1,591	2,048	3,103	4,146	4,784
Female	263	332	513	807	1,102	1,514	2,148	3,161	4,160
Black, both sexes	130	176	295	328	393	482	563	678	643
Male	66	87	163	188	223	262	281	375	313
Female	64	89	132	140	170	220	282	303	330
American Indian 1/, both sexes	4	5	9	11	15	19	22	24	25
Male	3	1	6	8	9	8	10	16	10
Female	1	4	3	3	6	11	12	8	15
Asian or Pacific Islander 2/, both sexes	25	28	31	54	55	65	96	100	131
Male	15	13	19	38	27	37	51	56	61
Female	10	15	12	16	28	28	45	44	70

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Hodgkin's disease (C81)									
All races, both sexes	78	73	73	88	64	94	92	140	134
Male	42	40	46	53	30	57	58	76	71
Female	36	33	27	35	34	37	34	64	63
White, both sexes	66	61	66	75	56	82	86	125	134
Male	36	34	42	45	26	48	55	67	71
Female	30	27	24	30	30	34	31	58	63
Black, both sexes	12	12	7	10	8	9	4	11	-
Male	6	6	4	5	4	7	2	5	-
Female	6	6	3	5	4	2	2	6	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	3	-
Male	-	-	-	-	-	-	1	3	-
Female	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	2	1	1	-
Male	-	-	-	3	-	2	-	1	-
Female	-	-	-	-	-	-	1	-	-
Non-Hodgkin's lymphoma (C82-C85)									
All races, both sexes	290	461	730	1,015	1,434	1,736	2,386	3,212	3,984
Male	192	290	466	646	865	984	1,342	1,763	2,018
Female	98	171	264	369	569	752	1,044	1,449	1,966
White, both sexes	226	384	603	879	1,290	1,564	2,179	2,982	3,761
Male	150	251	388	556	771	887	1,244	1,631	1,917
Female	76	133	215	323	519	677	935	1,351	1,844
Black, both sexes	56	65	107	103	119	138	150	170	146
Male	37	34	67	68	82	76	74	96	65
Female	19	31	40	35	37	62	76	74	81
American Indian 1/, both sexes	1	3	5	7	4	6	11	10	11
Male	1	-	3	5	2	3	3	5	6
Female	-	3	2	2	2	3	8	5	5
Asian or Pacific Islander 2/, both sexes	7	9	15	26	21	28	46	50	66
Male	4	5	8	17	10	18	21	31	30
Female	3	4	7	9	11	10	25	19	36

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Leukemia (C91-C95)									
All races, both sexes	368	433	622	789	1,041	1,373	2,073	2,937	3,515
Male	185	238	356	435	629	823	1,287	1,803	1,982
Female	183	195	266	354	412	550	786	1,134	1,533
White, both sexes	300	348	500	669	900	1,188	1,870	2,690	3,272
Male	154	199	292	366	550	717	1,176	1,651	1,855
Female	146	149	208	303	350	471	694	1,039	1,417
Black, both sexes	49	68	107	97	107	154	162	213	195
Male	20	33	54	53	61	90	86	130	105
Female	29	35	53	44	46	64	76	83	90
American Indian 1/, both sexes	3	1	2	2	8	7	9	7	5
Male	2	-	1	2	4	3	5	6	1
Female	1	1	1	-	4	4	4	1	4
Asian or Pacific Islander 2/, both sexes	16	16	13	21	26	24	32	27	43
Male	9	6	9	14	14	13	20	16	21
Female	7	10	4	7	12	11	12	11	22
Multiple myeloma and immunoproliferative neoplasms (C88,C90)									
All races, both sexes	38	108	245	455	611	919	1,374	1,805	2,097
Male	18	67	143	248	321	487	751	946	1,087
Female	20	41	102	207	290	432	623	859	1,010
White, both sexes	24	73	168	331	442	722	1,109	1,496	1,766
Male	14	50	103	181	240	392	621	793	932
Female	10	23	65	150	202	330	488	703	834
Black, both sexes	13	31	72	118	158	181	247	284	300
Male	3	14	36	62	75	89	119	144	142
Female	10	17	36	56	83	92	128	140	158
American Indian 1/, both sexes	-	1	2	2	3	5	1	3	9
Male	-	1	2	1	3	2	1	1	3
Female	-	-	-	1	-	3	-	2	6
Asian or Pacific Islander 2/, both sexes	1	3	3	4	8	11	17	22	22
Male	1	2	2	4	3	4	10	8	10
Female	-	1	1	-	5	7	7	14	12

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races, both sexes	2	2	5	3	6	6	7	15	13
Male	1	2	4	2	5	4	7	5	10
Female	1	-	1	1	1	2	-	10	3
White, both sexes	1	2	3	3	5	6	7	14	11
Male	-	2	2	2	4	4	7	4	9
Female	1	-	1	1	1	2	-	10	2
Black, both sexes	-	-	2	-	1	-	-	-	2
Male	-	-	2	-	1	-	-	-	1
Female	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, both sexes	771	1,319	2,114	3,193	4,088	5,205	6,819	8,897	10,087
Male	424	717	1,186	1,773	2,286	2,937	3,791	4,781	5,098
Female	347	602	928	1,420	1,802	2,268	3,028	4,116	4,989
White, both sexes	595	1,001	1,652	2,554	3,432	4,362	5,908	7,822	9,026
Male	336	539	928	1,437	1,921	2,430	3,328	4,244	4,553
Female	259	462	724	1,117	1,511	1,932	2,580	3,578	4,473
Black, both sexes	154	280	397	538	557	710	775	882	876
Male	75	156	228	285	312	430	392	446	448
Female	79	124	169	253	245	280	383	436	428
American Indian 1/, both sexes	4	4	14	16	14	33	26	47	36
Male	2	2	8	10	8	17	14	21	18
Female	2	2	6	6	6	16	12	26	18
Asian or Pacific Islander 2/, both sexes	18	34	51	85	85	100	110	146	149
Male	11	20	22	41	45	60	57	70	79
Female	7	14	29	44	40	40	53	76	70

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, both sexes	155	213	293	397	513	711	1,007	1,539	2,142
Male	86	102	160	225	270	387	587	828	1,120
Female	69	111	133	172	243	324	420	711	1,022
White, both sexes	124	168	225	333	447	611	888	1,351	1,962
Male	68	79	127	192	240	324	517	730	1,032
Female	56	89	98	141	207	287	371	621	930
Black, both sexes	24	39	56	58	57	78	96	152	150
Male	13	19	24	31	27	49	60	71	73
Female	11	20	32	27	30	29	36	81	77
American Indian 1/, both sexes	3	1	1	1	2	4	4	6	5
Male	1	-	-	-	-	3	1	5	1
Female	2	1	1	1	2	1	3	1	4
Asian or Pacific Islander 2/, both sexes	4	5	11	5	7	18	19	30	25
Male	4	4	9	2	3	11	9	22	14
Female	-	1	2	3	4	7	10	8	11

Anemias (D50-D64)

All races, both sexes	85	105	110	117	127	133	210	305	482
Male	34	44	52	56	66	64	95	141	198
Female	51	61	58	61	61	69	115	164	284
White, both sexes	17	26	38	56	84	85	160	250	414
Male	7	14	16	27	45	42	81	118	169
Female	10	12	22	29	39	43	79	132	245
Black, both sexes	65	76	69	59	40	43	43	46	58
Male	26	29	35	29	18	19	13	18	23
Female	39	47	34	30	22	24	30	28	35
American Indian 1/, both sexes	-	2	1	-	1	-	2	2	6
Male	-	1	-	-	1	-	1	-	5
Female	-	1	1	-	-	-	1	2	1
Asian or Pacific Islander 2/, both sexes	3	1	2	2	2	5	5	7	4
Male	1	-	1	-	2	3	-	5	1
Female	2	1	1	2	-	2	5	2	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Diabetes mellitus (E10-E14)

All races, both sexes	684	1,242	1,917	3,037	3,992	5,194	7,054	9,620	11,607
Male	396	755	1,124	1,697	2,173	2,770	3,556	4,670	5,288
Female	288	487	793	1,340	1,819	2,424	3,498	4,950	6,319
White, both sexes	493	926	1,344	2,170	2,967	3,912	5,393	7,682	9,561
Male	285	581	808	1,269	1,662	2,161	2,849	3,890	4,508
Female	208	345	536	901	1,305	1,751	2,544	3,792	5,053
Black, both sexes	177	285	517	762	894	1,121	1,441	1,677	1,781
Male	99	155	280	374	438	524	593	671	659
Female	78	130	237	388	456	597	848	1,006	1,122
American Indian 1/, both sexes	7	16	25	53	59	75	77	109	81
Male	6	10	18	23	26	42	33	37	37
Female	1	6	7	30	33	33	44	72	44
Asian or Pacific Islander 2/, both sexes	7	15	31	52	72	86	143	152	184
Male	6	9	18	31	47	43	81	72	84
Female	1	6	13	21	25	43	62	80	100

Nutritional deficiencies (E40-E64)

All races, both sexes	30	26	46	53	70	103	192	268	505
Male	9	7	25	21	45	52	87	134	207
Female	21	19	21	32	25	51	105	134	298
White, both sexes	21	19	37	36	52	78	159	211	425
Male	7	4	20	13	30	35	70	102	171
Female	14	15	17	23	22	43	89	109	254
Black, both sexes	9	7	8	16	18	23	29	49	68
Male	2	3	5	8	15	15	15	29	32
Female	7	4	3	8	3	8	14	20	36
American Indian 1/, both sexes	-	-	-	1	-	-	2	4	5
Male	-	-	-	-	-	-	1	1	2
Female	-	-	-	1	-	-	1	3	3
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	2	4	7
Male	-	-	-	-	-	2	1	2	2
Female	-	-	1	-	-	-	1	2	5

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Meningitis (G00,G03)									
All races, both sexes	35	37	45	55	43	47	48	60	68
Male	22	24	25	31	26	28	26	34	41
Female	13	13	20	24	17	19	22	26	27
White, both sexes	24	20	35	36	29	36	32	47	56
Male	15	12	21	22	20	22	19	29	33
Female	9	8	14	14	9	14	13	18	23
Black, both sexes	10	17	9	17	13	9	15	12	8
Male	6	12	3	8	6	5	6	5	5
Female	4	5	6	9	7	4	9	7	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	2	1	1	2
Male	-	-	1	1	-	1	1	-	2
Female	-	-	-	1	1	1	-	1	-
Parkinson's disease (G20-G21)									
All races, both sexes	4	3	16	33	82	188	626	1,482	3,322
Male	3	2	10	19	55	115	398	940	2,052
Female	1	1	6	14	27	73	228	542	1,270
White, both sexes	3	1	12	29	75	176	585	1,397	3,163
Male	2	1	8	16	49	109	372	890	1,943
Female	1	-	4	13	26	67	213	507	1,220
Black, both sexes	1	2	2	4	5	9	28	66	118
Male	1	1	1	3	4	5	19	41	80
Female	-	1	1	1	1	4	9	25	38
American Indian 1/, both sexes	-	-	-	-	1	1	3	6	9
Male	-	-	-	-	1	-	1	3	5
Female	-	-	-	-	-	1	2	3	4
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	2	10	13	32
Male	-	-	1	-	1	1	6	6	24
Female	-	-	1	-	-	1	4	7	8

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79
YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Alzheimer's disease (G30)

All races, both sexes	2	4	17	47	142	349	868	2,563	6,319
Male	1	3	10	24	60	150	420	1,134	2,433
Female	1	1	7	23	82	199	448	1,429	3,886
White, both sexes	2	4	14	41	132	318	796	2,343	5,864
Male	1	3	8	21	55	138	389	1,043	2,271
Female	1	1	6	20	77	180	407	1,300	3,593
Black, both sexes	-	-	1	4	7	23	67	194	400
Male	-	-	1	2	5	11	30	83	137
Female	-	-	-	2	2	12	37	111	263
American Indian 1/, both sexes	-	-	-	-	1	2	2	12	21
Male	-	-	-	-	-	-	1	3	6
Female	-	-	-	-	1	2	1	9	15
Asian or Pacific Islander 2/, both sexes	-	-	2	2	2	6	3	14	34
Male	-	-	1	1	-	1	-	5	19
Female	-	-	1	1	2	5	3	9	15

Major cardiovascular diseases (I00-I78)

All races, both sexes	5,491	11,133	17,785	25,798	33,454	44,028	61,769	94,418	135,296
Male	3,538	7,572	12,544	17,991	22,817	28,606	37,721	54,789	70,196
Female	1,953	3,561	5,241	7,807	10,637	15,422	24,048	39,629	65,100
White, both sexes	3,812	7,867	12,794	19,428	26,035	34,836	50,692	80,400	118,684
Male	2,573	5,616	9,522	14,011	18,204	23,256	31,747	47,612	62,471
Female	1,239	2,251	3,272	5,417	7,831	11,580	18,945	32,788	56,213
Black, both sexes	1,510	2,946	4,531	5,675	6,597	8,068	9,612	12,167	14,047
Male	854	1,736	2,710	3,503	4,074	4,613	5,104	6,104	6,348
Female	656	1,210	1,821	2,172	2,523	3,455	4,508	6,063	7,699
American Indian 1/, both sexes	53	97	130	179	224	274	367	351	436
Male	30	66	90	118	142	165	215	209	220
Female	23	31	40	61	82	109	152	142	216
Asian or Pacific Islander 2/, both sexes	116	223	330	516	598	850	1,098	1,500	2,129
Male	81	154	222	359	397	572	655	864	1,157
Female	35	69	108	157	201	278	443	636	972

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)									
All races, both sexes	4,333	8,848	14,313	21,167	27,676	35,723	49,178	73,227	101,764
Male	2,945	6,341	10,596	15,331	19,456	23,829	30,890	43,699	54,421
Female	1,388	2,507	3,717	5,836	8,220	11,894	18,288	29,528	47,343
White, both sexes	3,083	6,449	10,645	16,346	21,927	28,654	40,751	62,698	89,644
Male	2,185	4,835	8,267	12,213	15,794	19,628	26,246	38,218	48,669
Female	898	1,614	2,378	4,133	6,133	9,026	14,505	24,480	40,975
Black, both sexes	1,127	2,178	3,356	4,324	5,133	6,252	7,367	9,223	10,333
Male	674	1,335	2,091	2,746	3,237	3,645	3,973	4,699	4,752
Female	453	843	1,265	1,578	1,896	2,607	3,394	4,524	5,581
American Indian 1/, both sexes	44	78	103	146	180	214	291	257	326
Male	26	57	77	102	118	134	175	161	180
Female	18	21	26	44	62	80	116	96	146
Asian or Pacific Islander 2/, both sexes	79	143	209	351	436	603	769	1,049	1,461
Male	60	114	161	270	307	422	496	621	820
Female	19	29	48	81	129	181	273	428	641
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)									
All races, both sexes	49	92	119	133	183	237	327	492	601
Male	25	36	49	55	72	94	98	139	177
Female	24	56	70	78	111	143	229	353	424
White, both sexes	34	62	90	107	148	202	293	441	575
Male	19	27	39	48	62	85	88	122	169
Female	15	35	51	59	86	117	205	319	406
Black, both sexes	11	24	23	20	26	24	31	27	21
Male	5	7	8	5	7	7	8	8	7
Female	6	17	15	15	19	17	23	19	14
American Indian 1/, both sexes	1	1	1	-	3	3	1	3	1
Male	-	-	-	-	1	-	-	1	-
Female	1	1	1	-	2	3	1	2	1
Asian or Pacific Islander 2/, both sexes	3	5	5	6	6	8	2	21	4
Male	1	2	2	2	2	2	2	8	1
Female	2	3	3	4	4	6	-	13	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Hypertensive heart disease (I11)

All races, both sexes	323	658	952	1,152	1,249	1,423	1,570	2,035	2,694
Male	210	433	625	811	789	895	909	1,062	1,135
Female	113	225	327	341	460	528	661	973	1,559
White, both sexes	174	368	521	647	721	867	1,032	1,379	2,012
Male	126	259	383	480	485	564	616	749	855
Female	48	109	138	167	236	303	416	630	1,157
Black, both sexes	145	279	420	475	507	524	494	612	625
Male	81	167	235	308	291	308	269	287	248
Female	64	112	185	167	216	216	225	325	377
American Indian 1/, both sexes	-	-	1	10	5	6	6	5	5
Male	-	-	1	7	2	3	4	4	3
Female	-	-	-	3	3	3	2	1	2
Asian or Pacific Islander 2/, both sexes	4	11	10	20	16	26	38	39	52
Male	3	7	6	16	11	20	20	22	29
Female	1	4	4	4	5	6	18	17	23

Hypertensive heart and renal disease (I13)

All races, both sexes	22	61	68	79	103	105	149	268	376
Male	12	37	43	44	57	51	74	132	152
Female	10	24	25	35	46	54	75	136	224
White, both sexes	9	28	21	33	46	44	72	173	263
Male	5	17	12	19	25	21	38	88	120
Female	4	11	9	14	21	23	34	85	143
Black, both sexes	13	33	45	46	55	58	75	89	99
Male	7	20	30	25	31	27	36	41	29
Female	6	13	15	21	24	31	39	48	70
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	1
Male	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	3	1	6	13
Male	-	-	1	-	-	3	-	3	3
Female	-	-	-	-	1	-	1	3	10

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Ischemic heart diseases (I20-I25)

All races, both sexes	2,233	5,286	9,638	15,252	20,724	27,013	37,257	55,628	76,569
Male	1,665	4,122	7,601	11,548	15,154	18,641	24,225	34,329	42,452
Female	568	1,164	2,037	3,704	5,570	8,372	13,032	21,299	34,117
White, both sexes	1,745	4,106	7,570	12,265	16,873	22,082	31,304	48,138	67,860
Male	1,336	3,326	6,194	9,559	12,586	15,635	20,885	30,317	38,199
Female	409	780	1,376	2,706	4,287	6,447	10,419	17,821	29,661
Black, both sexes	432	1,041	1,845	2,624	3,383	4,302	5,124	6,495	7,298
Male	281	682	1,223	1,701	2,227	2,561	2,796	3,409	3,451
Female	151	359	622	923	1,156	1,741	2,328	3,086	3,847
American Indian 1/, both sexes	18	55	75	101	135	167	229	185	246
Male	14	42	61	73	91	110	143	114	138
Female	4	13	14	28	44	57	86	71	108
Asian or Pacific Islander 2/, both sexes	38	84	148	262	333	462	600	810	1,165
Male	34	72	123	215	250	335	401	489	664
Female	4	12	25	47	83	127	199	321	501

Acute myocardial infarction (I21-I22)

All races, both sexes	904	2,228	4,230	6,647	9,030	11,983	15,898	22,561	30,059
Male	657	1,730	3,319	4,946	6,490	8,148	10,177	13,646	16,295
Female	247	498	911	1,701	2,540	3,835	5,721	8,915	13,764
White, both sexes	744	1,752	3,442	5,458	7,574	10,085	13,575	19,710	26,733
Male	556	1,413	2,797	4,206	5,558	7,028	8,918	12,171	14,738
Female	188	339	645	1,252	2,016	3,057	4,657	7,539	11,995
Black, both sexes	146	423	699	1,038	1,277	1,658	1,982	2,465	2,761
Male	90	272	449	626	798	950	1,041	1,254	1,254
Female	56	151	250	412	479	708	941	1,211	1,507
American Indian 1/, both sexes	7	23	30	42	54	59	97	66	96
Male	5	18	25	30	38	35	64	37	54
Female	2	5	5	12	16	24	33	29	42
Asian or Pacific Islander 2/, both sexes	7	30	59	109	125	181	244	320	469
Male	6	27	48	84	96	135	154	184	249
Female	1	3	11	25	29	46	90	136	220

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Other acute ischemic heart diseases (I24)									
All races, both sexes	30	62	93	189	219	265	294	350	440
Male	24	47	76	141	163	178	194	204	232
Female	6	15	17	48	56	87	100	146	208
White, both sexes	23	45	69	145	160	207	236	294	373
Male	19	35	61	107	124	142	154	171	199
Female	4	10	8	38	36	65	82	123	174
Black, both sexes	7	15	23	40	51	52	47	46	56
Male	5	11	15	31	35	32	32	27	28
Female	2	4	8	9	16	20	15	19	28
American Indian 1/, both sexes	-	2	1	4	4	5	8	7	4
Male	-	1	-	3	2	3	6	4	3
Female	-	1	1	1	2	2	2	3	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	4	1	3	3	7
Male	-	-	-	-	2	1	2	2	2
Female	-	-	-	-	2	-	1	1	5
Other forms of chronic ischemic heart disease (I20,I25)									
All races, both sexes	1,299	2,996	5,315	8,416	11,475	14,765	21,065	32,717	46,070
Male	984	2,345	4,206	6,461	8,501	10,315	13,854	20,479	25,925
Female	315	651	1,109	1,955	2,974	4,450	7,211	12,238	20,145
White, both sexes	978	2,309	4,059	6,662	9,139	11,790	17,493	28,134	40,754
Male	761	1,878	3,336	5,246	6,904	8,465	11,813	17,975	23,262
Female	217	431	723	1,416	2,235	3,325	5,680	10,159	17,492
Black, both sexes	279	603	1,123	1,546	2,055	2,592	3,095	3,984	4,481
Male	186	399	759	1,044	1,394	1,579	1,723	2,128	2,169
Female	93	204	364	502	661	1,013	1,372	1,856	2,312
American Indian 1/, both sexes	11	30	44	55	77	103	124	112	146
Male	9	23	36	40	51	72	73	73	81
Female	2	7	8	15	26	31	51	39	65
Asian or Pacific Islander 2/, both sexes	31	54	89	153	204	280	353	487	689
Male	28	45	75	131	152	199	245	303	413
Female	3	9	14	22	52	81	108	184	276

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races, both sexes	391	985	1,894	3,086	3,841	4,401	5,309	7,258	9,271
Male	312	787	1,533	2,444	2,978	3,147	3,557	4,512	5,086
Female	79	198	361	642	863	1,254	1,752	2,746	4,185
White, both sexes	291	743	1,393	2,339	2,858	3,255	4,093	5,765	7,717
Male	237	615	1,164	1,903	2,252	2,408	2,811	3,636	4,249
Female	54	128	229	436	606	847	1,282	2,129	3,468
Black, both sexes	87	208	459	681	882	1,029	1,083	1,325	1,349
Male	65	145	332	489	646	662	648	769	716
Female	22	63	127	192	236	367	435	556	633
American Indian 1/, both sexes	3	9	13	20	27	25	31	24	33
Male	2	6	11	14	21	18	18	15	18
Female	1	3	2	6	6	7	13	9	15
Asian or Pacific Islander 2/, both sexes	10	25	29	46	74	92	102	144	172
Male	8	21	26	38	59	59	80	92	103
Female	2	4	3	8	15	33	22	52	69

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, both sexes	908	2,011	3,421	5,330	7,634	10,364	15,756	25,459	36,799
Male	672	1,558	2,673	4,017	5,523	7,168	10,297	15,967	20,839
Female	236	453	748	1,313	2,111	3,196	5,459	9,492	15,960
White, both sexes	687	1,566	2,666	4,323	6,281	8,535	13,400	22,369	33,037
Male	524	1,263	2,172	3,343	4,652	6,057	9,002	14,339	19,013
Female	163	303	494	980	1,629	2,478	4,398	8,030	14,024
Black, both sexes	192	395	664	865	1,173	1,563	2,012	2,659	3,132
Male	121	254	427	555	748	917	1,075	1,359	1,453
Female	71	141	237	310	425	646	937	1,300	1,679
American Indian 1/, both sexes	8	21	31	35	50	78	93	88	113
Male	7	17	25	26	30	54	55	58	63
Female	1	4	6	9	20	24	38	30	50
Asian or Pacific Islander 2/, both sexes	21	29	60	107	130	188	251	343	517
Male	20	24	49	93	93	140	165	211	310
Female	1	5	11	14	37	48	86	132	207

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other heart diseases (I26-I51)

All races, both sexes	1,706	2,751	3,536	4,551	5,417	6,945	9,875	14,804	21,524
Male	1,033	1,713	2,278	2,873	3,384	4,148	5,584	8,037	10,505
Female	673	1,038	1,258	1,678	2,033	2,797	4,291	6,767	11,019
White, both sexes	1,121	1,885	2,443	3,294	4,139	5,459	8,050	12,567	18,934
Male	699	1,206	1,639	2,107	2,636	3,323	4,619	6,942	9,326
Female	422	679	804	1,187	1,503	2,136	3,431	5,625	9,608
Black, both sexes	526	801	1,023	1,159	1,162	1,344	1,643	2,000	2,290
Male	300	459	595	707	681	742	864	954	1,017
Female	226	342	428	452	481	602	779	1,046	1,273
American Indian 1/, both sexes	25	22	25	35	36	38	54	64	73
Male	12	15	15	22	23	21	28	42	39
Female	13	7	10	13	13	17	26	22	34
Asian or Pacific Islander 2/, both sexes	34	43	45	63	80	104	128	173	227
Male	22	33	29	37	44	62	73	99	123
Female	12	10	16	26	36	42	55	74	104

Acute and subacute endocarditis (I33)

All races, both sexes	41	75	89	97	88	77	105	119	166
Male	24	49	59	61	52	44	59	63	95
Female	17	26	30	36	36	33	46	56	71
White, both sexes	27	53	57	65	59	63	89	102	148
Male	14	36	39	44	38	38	47	57	87
Female	13	17	18	21	21	25	42	45	61
Black, both sexes	11	20	30	27	26	13	15	16	16
Male	9	12	20	15	12	6	11	5	6
Female	2	8	10	12	14	7	4	11	10
American Indian 1/, both sexes	1	2	2	3	1	1	-	-	1
Male	1	1	-	2	1	-	-	-	1
Female	-	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	2	2	-	1	1	1
Male	-	-	-	-	1	-	1	1	1
Female	2	-	-	2	1	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races, both sexes	45	41	45	40	47	42	52	79	82
Male	30	28	28	18	31	21	24	43	42
Female	15	13	17	22	16	21	28	36	40
White, both sexes	27	31	26	33	39	33	41	67	69
Male	18	21	17	15	28	17	15	37	34
Female	9	10	9	18	11	16	26	30	35
Black, both sexes	14	10	14	7	7	8	10	8	8
Male	9	7	7	3	2	4	8	3	5
Female	5	3	7	4	5	4	2	5	3
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	2
Male	-	-	1	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	4	-	1	1	1	3	3
Male	3	-	3	-	1	-	1	2	2
Female	1	-	1	-	-	1	-	1	1

Heart failure (I50)

All races, both sexes	77	185	283	528	828	1,383	2,284	4,119	6,936
Male	40	119	191	305	479	785	1,243	2,171	3,286
Female	37	66	92	223	349	598	1,041	1,948	3,650
White, both sexes	44	120	180	378	627	1,088	1,870	3,514	6,178
Male	25	84	127	224	383	624	1,049	1,892	2,959
Female	19	36	53	154	244	464	821	1,622	3,219
Black, both sexes	32	62	98	143	189	266	373	544	674
Male	14	33	60	78	90	143	175	248	285
Female	18	29	38	65	99	123	198	296	389
American Indian 1/, both sexes	1	-	2	5	4	10	17	25	29
Male	1	-	1	3	1	5	8	15	16
Female	-	-	1	2	3	5	9	10	13
Asian or Pacific Islander 2/, both sexes	-	3	3	2	8	19	24	36	55
Male	-	2	3	-	5	13	11	16	26
Female	-	1	-	2	3	6	13	20	29

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races, both sexes	1,543	2,450	3,119	3,886	4,454	5,443	7,434	10,487	14,340
Male	939	1,517	2,000	2,489	2,822	3,298	4,258	5,760	7,082
Female	604	933	1,119	1,397	1,632	2,145	3,176	4,727	7,258
White, both sexes	1,023	1,681	2,180	2,818	3,414	4,275	6,050	8,884	12,539
Male	642	1,065	1,456	1,824	2,187	2,644	3,508	4,956	6,246
Female	381	616	724	994	1,227	1,631	2,542	3,928	6,293
Black, both sexes	469	709	881	982	940	1,057	1,245	1,432	1,592
Male	268	407	508	611	577	589	670	698	721
Female	201	302	373	371	363	468	575	734	871
American Indian 1/, both sexes	23	20	20	27	31	27	37	38	41
Male	10	14	13	17	21	16	20	26	21
Female	13	6	7	10	10	11	17	12	20
Asian or Pacific Islander 2/, both sexes	28	40	38	59	69	84	102	133	168
Male	19	31	23	37	37	49	60	80	94
Female	9	9	15	22	32	35	42	53	74

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races, both sexes	105	243	352	526	597	843	1,094	1,690	2,507
Male	61	155	224	331	352	466	504	804	1,049
Female	44	88	128	195	245	377	590	886	1,458
White, both sexes	41	117	169	269	345	506	706	1,200	1,925
Male	30	79	117	184	212	291	340	586	823
Female	11	38	52	85	133	215	366	614	1,102
Black, both sexes	62	117	171	238	242	317	365	441	520
Male	30	70	99	134	137	166	154	191	197
Female	32	47	72	104	105	151	211	250	323
American Indian 1/, both sexes	1	2	2	3	4	10	4	9	9
Male	-	2	1	2	1	4	1	2	3
Female	1	-	1	1	3	6	3	7	6
Asian or Pacific Islander 2/, both sexes	1	7	10	16	6	10	19	40	53
Male	1	4	7	11	2	5	9	25	26
Female	-	3	3	5	4	5	10	15	27

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Cerebrovascular diseases (I60-I69)

All races, both sexes	880	1,719	2,625	3,386	4,145	5,811	8,671	14,978	24,678
Male	430	867	1,401	1,834	2,294	3,202	4,513	7,529	11,192
Female	450	852	1,224	1,552	1,851	2,609	4,158	7,449	13,486
White, both sexes	558	1,087	1,627	2,287	2,930	4,309	6,775	12,449	21,385
Male	282	560	894	1,244	1,607	2,391	3,540	6,288	9,746
Female	276	527	733	1,043	1,323	1,918	3,235	6,161	11,639
Black, both sexes	283	553	878	958	1,043	1,248	1,582	2,107	2,674
Male	127	270	449	520	594	664	818	1,017	1,153
Female	156	283	429	438	449	584	764	1,090	1,521
American Indian 1/, both sexes	7	15	22	25	32	47	44	64	86
Male	4	7	10	12	17	26	27	38	30
Female	3	8	12	13	15	21	17	26	56
Asian or Pacific Islander 2/, both sexes	32	64	98	116	140	207	270	358	533
Male	17	30	48	58	76	121	128	186	263
Female	15	34	50	58	64	86	142	172	270

Atherosclerosis (I70)

All races, both sexes	16	32	71	105	170	281	495	928	1,685
Male	10	25	48	71	108	186	290	510	838
Female	6	7	23	34	62	95	205	418	847
White, both sexes	12	24	50	73	140	226	431	815	1,535
Male	9	19	36	50	93	154	252	461	775
Female	3	5	14	23	47	72	179	354	760
Black, both sexes	4	8	19	28	28	51	51	97	132
Male	1	6	11	17	14	29	33	42	54
Female	3	2	8	11	14	22	18	55	78
American Indian 1/, both sexes	-	-	-	-	2	1	8	8	3
Male	-	-	-	-	1	1	3	3	1
Female	-	-	-	-	1	-	5	5	2
Asian or Pacific Islander 2/, both sexes	-	-	2	4	-	3	5	8	15
Male	-	-	1	4	-	2	2	4	8
Female	-	-	1	-	-	1	3	4	7

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Other diseases of circulatory system (I71-I78)									
All races, both sexes	157	291	424	614	866	1,370	2,331	3,595	4,662
Male	92	184	275	424	607	923	1,524	2,247	2,696
Female	65	107	149	190	259	447	807	1,348	1,966
White, both sexes	118	190	303	453	693	1,141	2,029	3,238	4,195
Male	67	123	208	320	498	792	1,369	2,059	2,458
Female	51	67	95	133	195	349	660	1,179	1,737
Black, both sexes	34	90	107	127	151	200	247	299	388
Male	22	55	60	86	92	109	126	155	192
Female	12	35	47	41	59	91	121	144	196
American Indian 1/, both sexes	1	2	3	5	6	2	20	13	12
Male	-	-	2	2	5	-	9	5	6
Female	1	2	1	3	1	2	11	8	6
Asian or Pacific Islander 2/, both sexes	4	9	11	29	16	27	35	45	67
Male	3	6	5	16	12	22	20	28	40
Female	1	3	6	13	4	5	15	17	27
Aortic aneurysm and dissection (I71)									
All races, both sexes	103	200	268	394	585	908	1,552	2,460	3,043
Male	72	147	191	301	438	673	1,097	1,633	1,885
Female	31	53	77	93	147	235	455	827	1,158
White, both sexes	75	129	193	303	475	782	1,381	2,270	2,795
Male	52	97	147	237	364	592	1,006	1,527	1,747
Female	23	32	46	66	111	190	375	743	1,048
Black, both sexes	25	65	68	69	98	102	129	152	190
Male	18	46	39	52	63	62	68	83	101
Female	7	19	29	17	35	40	61	69	89
American Indian 1/, both sexes	-	-	1	-	1	2	13	8	9
Male	-	-	1	-	1	-	7	4	5
Female	-	-	-	-	-	2	6	4	4
Asian or Pacific Islander 2/, both sexes	3	6	6	22	11	22	29	30	49
Male	2	4	4	12	10	19	16	19	32
Female	1	2	2	10	1	3	13	11	17

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races, both sexes	54	91	156	220	281	462	779	1,135	1,619
Male	20	37	84	123	169	250	427	614	811
Female	34	54	72	97	112	212	352	521	808
White, both sexes	43	61	110	150	218	359	648	968	1,400
Male	15	26	61	83	134	200	363	532	711
Female	28	35	49	67	84	159	285	436	689
Black, both sexes	9	25	39	58	53	98	118	147	198
Male	4	9	21	34	29	47	58	72	91
Female	5	16	18	24	24	51	60	75	107
American Indian 1/, both sexes	1	2	2	5	5	-	7	5	3
Male	-	-	1	2	4	-	2	1	1
Female	1	2	1	3	1	-	5	4	2
Asian or Pacific Islander 2/, both sexes	1	3	5	7	5	5	6	15	18
Male	1	2	1	4	2	3	4	9	8
Female	-	1	4	3	3	2	2	6	10

Other disorders of circulatory system (I80-I99)

All races, both sexes	125	194	249	266	231	285	326	469	573
Male	70	120	145	150	126	138	170	220	245
Female	55	74	104	116	105	147	156	249	328
White, both sexes	87	143	189	203	175	222	242	395	488
Male	47	94	111	117	94	107	131	192	214
Female	40	49	78	86	81	115	111	203	274
Black, both sexes	36	51	58	60	53	56	79	68	82
Male	22	26	34	32	29	27	36	26	28
Female	14	25	24	28	24	29	43	42	54
American Indian 1/, both sexes	-	-	1	1	-	2	3	4	1
Male	-	-	-	-	-	1	2	1	1
Female	-	-	1	1	-	1	1	3	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	3	5	2	2	2
Male	1	-	-	1	3	3	1	1	2
Female	1	-	1	1	-	2	1	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races, both sexes	310	620	1,174	2,077	3,985	6,754	11,833	19,324	24,309
Male	145	292	617	1,044	2,072	3,492	6,149	10,061	12,305
Female	165	328	557	1,033	1,913	3,262	5,684	9,263	12,004
White, both sexes	194	430	882	1,702	3,469	6,063	10,793	17,972	22,832
Male	85	207	472	849	1,791	3,102	5,523	9,263	11,418
Female	109	223	410	853	1,678	2,961	5,270	8,709	11,414
Black, both sexes	104	171	259	340	445	628	881	1,133	1,191
Male	52	78	125	169	240	355	532	677	691
Female	52	93	134	171	205	273	349	456	500
American Indian 1/, both sexes	4	6	10	7	36	35	58	66	66
Male	1	3	7	5	19	22	24	26	39
Female	3	3	3	2	17	13	34	40	27
Asian or Pacific Islander 2/, both sexes	8	13	23	28	35	28	101	153	220
Male	7	4	13	21	22	13	70	95	157
Female	1	9	10	7	13	15	31	58	63

Bronchitis, chronic and unspecified (J40-J42)

All races, both sexes	7	13	19	19	20	47	67	91	156
Male	4	10	8	8	9	23	41	38	84
Female	3	3	11	11	11	24	26	53	72
White, both sexes	7	7	17	15	17	44	58	82	144
Male	4	4	8	7	8	20	37	34	78
Female	3	3	9	8	9	24	21	48	66
Black, both sexes	-	6	2	4	1	3	7	9	10
Male	-	6	-	1	-	3	2	4	4
Female	-	-	2	3	1	-	5	5	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	2	-	2
Male	-	-	-	-	1	-	2	-	2
Female	-	-	-	-	1	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Emphysema (J43)

All races, both sexes	23	59	162	312	680	1,127	1,976	3,144	3,522
Male	15	31	101	179	366	577	1,077	1,661	1,839
Female	8	28	61	133	314	550	899	1,483	1,683
White, both sexes	18	45	121	262	609	1,036	1,819	2,964	3,342
Male	11	23	72	146	325	522	966	1,553	1,735
Female	7	22	49	116	284	514	853	1,411	1,607
Black, both sexes	4	13	38	47	63	84	135	146	138
Male	3	8	26	30	36	52	97	90	74
Female	1	5	12	17	27	32	38	56	64
American Indian 1/, both sexes	1	-	1	1	6	5	8	11	13
Male	1	-	1	1	3	2	3	3	7
Female	-	-	-	-	3	3	5	8	6
Asian or Pacific Islander 2/, both sexes	-	1	2	2	2	2	14	23	29
Male	-	-	2	2	2	1	11	15	23
Female	-	1	-	-	-	1	3	8	6

Asthma (J45-J46)

All races, both sexes	183	275	318	296	270	270	328	406	420
Male	81	115	128	110	106	94	121	132	120
Female	102	160	190	186	164	176	207	274	300
White, both sexes	95	161	178	182	162	185	231	316	348
Male	37	68	79	71	57	62	79	106	102
Female	58	93	99	111	105	123	152	210	246
Black, both sexes	79	100	128	100	89	79	80	73	60
Male	38	41	44	30	38	29	33	21	10
Female	41	59	84	70	51	50	47	52	50
American Indian 1/, both sexes	2	5	1	-	6	1	3	3	1
Male	-	2	-	-	3	-	1	-	1
Female	2	3	1	-	3	1	2	3	-
Asian or Pacific Islander 2/, both sexes	7	9	11	14	13	5	14	14	11
Male	6	4	5	9	8	3	8	5	7
Female	1	5	6	5	5	2	6	9	4

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79
YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Pneumonitis due to solids and liquids (J69)

All races, both sexes	65	116	157	209	257	341	642	1,257	2,258
Male	45	70	99	126	153	201	405	790	1,291
Female	20	46	58	83	104	140	237	467	967
White, both sexes	43	99	115	166	214	271	530	1,050	2,006
Male	32	61	72	99	131	160	329	653	1,139
Female	11	38	43	67	83	111	201	397	867
Black, both sexes	21	17	39	41	39	63	102	188	219
Male	12	9	26	27	20	35	67	123	132
Female	9	8	13	14	19	28	35	65	87
American Indian 1/, both sexes	-	-	2	1	4	3	4	5	6
Male	-	-	-	-	2	2	4	3	5
Female	-	-	2	1	2	1	-	2	1
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	4	6	14	27
Male	1	-	1	-	-	4	5	11	15
Female	-	-	-	1	-	-	1	3	12

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races, both sexes	231	384	527	805	1,044	1,367	2,184	3,264	4,487
Male	132	203	281	431	572	752	1,179	1,782	2,353
Female	99	181	246	374	472	615	1,005	1,482	2,134
White, both sexes	152	291	374	635	867	1,150	1,888	2,948	4,123
Male	87	154	199	341	479	639	1,035	1,615	2,167
Female	65	137	175	294	388	511	853	1,333	1,956
Black, both sexes	70	83	142	146	157	170	254	250	280
Male	41	44	79	74	81	85	125	121	143
Female	29	39	63	72	76	85	129	129	137
American Indian 1/, both sexes	4	4	6	11	6	18	15	24	17
Male	3	2	1	9	3	11	6	9	10
Female	1	2	5	2	3	7	9	15	7
Asian or Pacific Islander 2/, both sexes	5	6	5	13	14	29	27	42	67
Male	1	3	2	7	9	17	13	37	33
Female	4	3	3	6	5	12	14	5	34

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Hernia (K40-K46)									
All races, both sexes	13	18	26	58	51	62	84	126	224
Male	8	5	13	32	20	29	39	53	91
Female	5	13	13	26	31	33	45	73	133
White, both sexes	9	13	20	47	45	54	68	117	198
Male	6	4	9	25	18	25	31	51	85
Female	3	9	11	22	27	29	37	66	113
Black, both sexes	3	5	5	8	5	8	14	7	22
Male	1	1	4	5	1	4	7	2	4
Female	2	4	1	3	4	4	7	5	18
American Indian 1/, both sexes	1	-	-	2	1	-	2	1	1
Male	1	-	-	1	1	-	1	-	-
Female	-	-	-	1	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	3
Male	-	-	-	1	-	-	-	-	2
Female	-	-	1	-	-	-	-	1	1
Chronic liver disease and cirrhosis (K70,K73-K74)									
All races, both sexes	1,083	2,288	3,374	3,280	2,947	2,827	2,751	2,731	2,367
Male	716	1,598	2,515	2,460	2,066	1,939	1,729	1,588	1,240
Female	367	690	859	820	881	888	1,022	1,143	1,127
White, both sexes	872	1,873	2,817	2,698	2,497	2,451	2,434	2,491	2,223
Male	588	1,333	2,160	2,045	1,775	1,700	1,553	1,453	1,162
Female	284	540	657	653	722	751	881	1,038	1,061
Black, both sexes	144	307	453	484	349	284	249	185	103
Male	92	192	289	350	234	180	146	112	58
Female	52	115	164	134	115	104	103	73	45
American Indian 1/, both sexes	56	81	72	75	69	54	39	20	19
Male	27	51	42	49	39	32	15	6	11
Female	29	30	30	26	30	22	24	14	8
Asian or Pacific Islander 2/, both sexes	11	27	32	23	32	38	29	35	22
Male	9	22	24	16	18	27	15	17	9
Female	2	5	8	7	14	11	14	18	13

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcoholic liver disease (K70)

All races, both sexes	741	1,485	2,000	1,888	1,575	1,394	1,085	806	533
Male	486	1,047	1,510	1,458	1,172	1,044	821	612	419
Female	255	438	490	430	403	350	264	194	114
White, both sexes	596	1,214	1,670	1,576	1,329	1,204	961	728	487
Male	397	867	1,301	1,237	1,013	917	735	553	387
Female	199	347	369	339	316	287	226	175	100
Black, both sexes	94	193	259	249	187	138	107	63	35
Male	61	126	164	177	124	87	74	50	24
Female	33	67	95	72	63	51	33	13	11
American Indian 1/, both sexes	42	57	52	50	44	35	15	8	8
Male	20	36	30	33	25	26	11	3	5
Female	22	21	22	17	19	9	4	5	3
Asian or Pacific Islander 2/, both sexes	9	21	19	13	15	17	2	7	3
Male	8	18	15	11	10	14	1	6	3
Female	1	3	4	2	5	3	1	1	-

Other chronic liver disease and cirrhosis (K73-K74)

All races, both sexes	342	803	1,374	1,392	1,372	1,433	1,666	1,925	1,834
Male	230	551	1,005	1,002	894	895	908	976	821
Female	112	252	369	390	478	538	758	949	1,013
White, both sexes	276	659	1,147	1,122	1,168	1,247	1,473	1,763	1,736
Male	191	466	859	808	762	783	818	900	775
Female	85	193	288	314	406	464	655	863	961
Black, both sexes	50	114	194	235	162	146	142	122	68
Male	31	66	125	173	110	93	72	62	34
Female	19	48	69	62	52	53	70	60	34
American Indian 1/, both sexes	14	24	20	25	25	19	24	12	11
Male	7	15	12	16	14	6	4	3	6
Female	7	9	8	9	11	13	20	9	5
Asian or Pacific Islander 2/, both sexes	2	6	13	10	17	21	27	28	19
Male	1	4	9	5	8	13	14	11	6
Female	1	2	4	5	9	8	13	17	13

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, both sexes	22	31	43	66	80	109	184	266	419
Male	12	15	23	30	48	62	100	142	224
Female	10	16	20	36	32	47	84	124	195
White, both sexes	17	21	26	49	64	87	151	215	373
Male	10	12	16	23	41	51	87	119	203
Female	7	9	10	26	23	36	64	96	170
Black, both sexes	4	10	14	14	13	16	28	37	37
Male	1	3	5	6	5	9	10	15	15
Female	3	7	9	8	8	7	18	22	22
American Indian 1/, both sexes	-	-	1	1	1	1	-	2	5
Male	-	-	1	-	1	-	-	1	2
Female	-	-	-	1	-	1	-	1	3
Asian or Pacific Islander 2/, both sexes	1	-	2	2	2	5	5	12	4
Male	1	-	1	1	1	2	3	7	4
Female	-	-	1	1	1	3	2	5	-

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races, both sexes	278	462	687	964	1,269	1,831	2,811	4,179	5,893
Male	152	277	433	549	697	954	1,444	2,120	3,039
Female	126	185	254	415	572	877	1,367	2,059	2,854
White, both sexes	144	268	376	560	789	1,220	1,966	3,178	4,857
Male	80	163	237	321	444	662	1,071	1,698	2,567
Female	64	105	139	239	345	558	895	1,480	2,290
Black, both sexes	130	176	300	369	436	555	756	897	931
Male	71	104	191	207	227	262	334	382	421
Female	59	72	109	162	209	293	422	515	510
American Indian 1/, both sexes	1	11	5	17	17	22	36	29	22
Male	-	7	2	9	9	12	18	10	7
Female	1	4	3	8	8	10	18	19	15
Asian or Pacific Islander 2/, both sexes	3	7	6	18	27	34	53	75	83
Male	1	3	3	12	17	18	21	30	44
Female	2	4	3	6	10	16	32	45	39

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79
YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races, both sexes	1	2	5	11	2	8	13	16	16
Male	-	1	2	9	2	6	7	5	11
Female	1	1	3	2	-	2	6	11	5
White, both sexes	-	1	2	4	2	8	8	14	12
Male	-	1	2	3	2	6	4	5	9
Female	-	-	-	1	-	2	4	9	3
Black, both sexes	1	1	3	7	-	-	4	2	3
Male	-	-	-	6	-	-	2	-	1
Female	1	1	3	1	-	-	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1
Male	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07,N26)

All races, both sexes	14	20	18	21	31	28	43	60	78
Male	8	12	13	14	16	15	24	30	45
Female	6	8	5	7	15	13	19	30	33
White, both sexes	7	13	15	11	24	19	32	53	67
Male	5	7	11	7	13	10	18	27	39
Female	2	6	4	4	11	9	14	26	28
Black, both sexes	7	3	3	9	5	7	9	5	9
Male	3	2	2	6	2	5	6	2	4
Female	4	1	1	3	3	2	3	3	5
American Indian 1/, both sexes	-	2	-	1	1	-	1	-	1
Male	-	2	-	1	1	-	-	-	1
Female	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	2	1	2	1
Male	-	1	-	-	-	-	-	1	1
Female	-	1	-	-	1	2	1	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races, both sexes	9	6	6	3	4	4	11	10	17
Female	9	6	6	3	4	4	11	10	17
White, both sexes	5	2	4	3	2	4	9	8	15
Female	5	2	4	3	2	4	9	8	15
Black, both sexes	3	4	2	-	2	-	2	2	2
Female	3	4	2	-	2	-	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-

Pregnancy, childbirth and the puerperium (000-099)

All races, both sexes	86	35	5	1
Female	86	35	5	1
White, both sexes	56	21	2	1
Female	56	21	2	1
Black, both sexes	26	12	3	-
Female	26	12	3	-
American Indian 1/, both sexes	-	-	-	-
Female	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	-	-
Female	4	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Pregnancy with abortive outcome (000-007)									
All races, both sexes	10	4	-	-
Female	10	4	-	-
White, both sexes	6	3	-	-
Female	6	3	-	-
Black, both sexes	4	1	-	-
Female	4	1	-	-
American Indian 1/, both sexes	-	-	-	-
Female	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-
Female	-	-	-	-
Other complications of pregnancy, childbirth and the puerperium (010-099)									
All races, both sexes	76	31	5	1
Female	76	31	5	1
White, both sexes	50	18	2	1
Female	50	18	2	1
Black, both sexes	22	11	3	-
Female	22	11	3	-
American Indian 1/, both sexes	-	-	-	-
Female	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	-	-
Female	4	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

All races, both sexes	2	2	1	1	1	1	-	-	-
Male	-	2	1	1	-	-	-	-	-
Female	2	-	-	-	1	1	-	-	-
White, both sexes	2	2	1	1	1	1	-	-	-
Male	-	2	1	1	-	-	-	-	-
Female	2	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes	277	292	287	325	296	273	248	216	242
Male	154	162	157	190	130	134	120	102	99
Female	123	130	130	135	166	139	128	114	143
White, both sexes	223	236	245	294	255	241	219	191	225
Male	127	132	132	168	114	123	99	89	94
Female	96	104	113	126	141	118	120	102	131
Black, both sexes	48	45	39	28	32	27	25	18	14
Male	25	26	22	19	11	7	18	8	5
Female	23	19	17	9	21	20	7	10	9
American Indian 1/, both sexes	2	2	-	1	2	3	2	3	1
Male	-	1	-	1	1	3	1	2	-
Female	2	1	-	-	1	-	1	1	1
Asian or Pacific Islander 2/, both sexes	4	9	3	2	7	2	2	4	2
Male	2	3	3	2	4	1	2	3	-
Female	2	6	-	-	3	1	-	1	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, both sexes	1,284	1,703	1,727	1,415	1,152	1,101	1,353	1,687	2,235
Male	857	1,137	1,157	966	743	686	832	876	1,100
Female	427	566	570	449	409	415	521	811	1,135
White, both sexes	1,035	1,360	1,364	1,087	907	876	1,072	1,446	1,965
Male	714	935	929	744	574	551	658	750	974
Female	321	425	435	343	333	325	414	696	991
Black, both sexes	214	314	325	298	216	208	256	223	249
Male	115	178	200	203	152	123	159	114	116
Female	99	136	125	95	64	85	97	109	133
American Indian 1/, both sexes	12	9	13	13	13	6	7	4	7
Male	7	7	12	9	8	5	4	3	4
Female	5	2	1	4	5	1	3	1	3
Asian or Pacific Islander 2/, both sexes	23	20	25	17	16	11	18	14	14
Male	21	17	16	10	9	7	11	9	6
Female	2	3	9	7	7	4	7	5	8

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes	3,161	4,952	6,227	7,099	7,152	8,004	10,108	14,899	21,425
Male	1,873	2,928	3,811	4,100	3,967	4,269	5,174	7,147	9,237
Female	1,288	2,024	2,416	2,999	3,185	3,735	4,934	7,752	12,188
White, both sexes	2,269	3,647	4,572	5,526	5,694	6,488	8,481	12,879	19,016
Male	1,398	2,220	2,913	3,245	3,198	3,443	4,349	6,184	8,195
Female	871	1,427	1,659	2,281	2,496	3,045	4,132	6,695	10,821
Black, both sexes	793	1,176	1,517	1,413	1,287	1,338	1,423	1,753	2,096
Male	415	632	819	767	695	725	724	837	907
Female	378	544	698	646	592	613	699	916	1,189
American Indian 1/, both sexes	58	74	69	82	74	60	77	71	74
Male	35	49	38	44	31	30	37	34	27
Female	23	25	31	38	43	30	40	37	47
Asian or Pacific Islander 2/, both sexes	41	55	69	78	97	118	127	196	239
Male	25	27	41	44	43	71	64	92	108
Female	16	28	28	34	54	47	63	104	131

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races, both sexes	7,471	7,942	6,870	5,408	4,021	3,484	3,375	4,323	5,596
Male	5,537	5,907	5,103	3,960	2,804	2,329	2,117	2,522	3,039
Female	1,934	2,035	1,767	1,448	1,217	1,155	1,258	1,801	2,557
White, both sexes	6,171	6,492	5,554	4,347	3,334	2,941	2,855	3,762	5,039
Male	4,608	4,854	4,143	3,179	2,297	1,945	1,770	2,207	2,712
Female	1,563	1,638	1,411	1,168	1,037	996	1,085	1,555	2,327
Black, both sexes	1,058	1,229	1,109	879	566	431	417	451	424
Male	772	906	817	658	434	316	285	258	253
Female	286	323	292	221	132	115	132	193	171
American Indian 1/, both sexes	129	127	108	89	55	39	31	32	29
Male	81	85	80	70	34	22	17	19	20
Female	48	42	28	19	21	17	14	13	9
Asian or Pacific Islander 2/, both sexes	113	94	99	93	66	73	72	78	104
Male	76	62	63	53	39	46	45	38	54
Female	37	32	36	40	27	27	27	40	50

Transport accidents (V01-V99,Y85)

All races, both sexes	3,889	3,772	3,248	2,717	2,115	1,768	1,599	1,716	1,860
Male	2,802	2,758	2,338	1,922	1,456	1,161	996	1,025	1,050
Female	1,087	1,014	910	795	659	607	603	691	810
White, both sexes	3,208	3,110	2,679	2,245	1,796	1,523	1,375	1,503	1,683
Male	2,312	2,277	1,945	1,598	1,226	987	845	899	942
Female	896	833	734	647	570	536	530	604	741
Black, both sexes	527	542	457	361	244	180	165	160	109
Male	396	408	321	258	188	133	120	100	73
Female	131	134	136	103	56	47	45	60	36
American Indian 1/, both sexes	80	66	53	43	26	24	19	15	14
Male	50	42	38	32	12	13	9	10	9
Female	30	24	15	11	14	11	10	5	5
Asian or Pacific Islander 2/, both sexes	74	54	59	68	49	41	40	38	54
Male	44	31	34	34	30	28	22	16	26
Female	30	23	25	34	19	13	18	22	28

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races, both sexes	3,545	3,382	2,936	2,425	1,917	1,589	1,436	1,602	1,764
Male	2,507	2,433	2,091	1,689	1,293	1,013	870	932	980
Female	1,038	949	845	736	624	576	566	670	784
White, both sexes	2,918	2,772	2,421	1,989	1,622	1,362	1,234	1,402	1,598
Male	2,061	1,996	1,739	1,392	1,085	852	736	816	879
Female	857	776	682	597	537	510	498	586	719
Black, both sexes	485	502	415	342	225	169	150	151	102
Male	361	374	288	242	171	124	107	94	68
Female	124	128	127	100	54	45	43	57	34
American Indian 1/, both sexes	74	61	45	37	24	23	17	14	13
Male	45	38	32	26	10	13	9	9	8
Female	29	23	13	11	14	10	8	5	5
Asian or Pacific Islander 2/, both sexes	68	47	55	57	46	35	35	35	51
Male	40	25	32	29	27	24	18	13	25
Female	28	22	23	28	19	11	17	22	26

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, both sexes	149	168	125	109	64	58	52	48	51
Male	121	137	92	74	48	50	38	37	33
Female	28	31	33	35	16	8	14	11	18
White, both sexes	118	132	94	92	50	51	43	42	44
Male	98	108	70	62	36	46	30	32	30
Female	20	24	24	30	14	5	13	10	14
Black, both sexes	26	28	25	10	11	4	7	3	4
Male	21	22	20	7	9	3	7	2	2
Female	5	6	5	3	2	1	-	1	2
American Indian 1/, both sexes	3	5	3	3	1	1	1	-	-
Male	2	4	1	3	1	-	-	-	-
Female	1	1	2	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	3	4	2	2	1	3	3
Male	-	3	1	2	2	1	1	3	1
Female	2	-	2	2	-	1	-	-	2

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races, both sexes	195	222	187	183	134	121	111	66	45
Male	174	188	155	159	115	98	88	56	37
Female	21	34	32	24	19	23	23	10	8
White, both sexes	172	206	164	164	124	110	98	59	41
Male	153	173	136	144	105	89	79	51	33
Female	19	33	28	20	19	21	19	8	8
Black, both sexes	16	12	17	9	8	7	8	6	3
Male	14	12	13	9	8	6	6	4	3
Female	2	-	4	-	-	1	2	2	-
American Indian 1/, both sexes	3	-	5	3	1	-	1	1	1
Male	3	-	5	3	1	-	-	1	1
Female	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	4	1	7	1	4	4	-	-
Male	4	3	1	3	1	3	3	-	-
Female	-	1	-	4	-	1	1	-	-

Nontransport accidents (W00-X59,Y86)

All races, both sexes	3,582	4,170	3,622	2,691	1,906	1,716	1,776	2,607	3,736
Male	2,735	3,149	2,765	2,038	1,348	1,168	1,121	1,497	1,989
Female	847	1,021	857	653	558	548	655	1,110	1,747
White, both sexes	2,963	3,382	2,875	2,102	1,538	1,418	1,480	2,259	3,356
Male	2,296	2,577	2,198	1,581	1,071	958	925	1,308	1,770
Female	667	805	677	521	467	460	555	951	1,586
Black, both sexes	531	687	652	518	322	251	252	291	315
Male	376	498	496	400	246	183	165	158	180
Female	155	189	156	118	76	68	87	133	135
American Indian 1/, both sexes	49	61	55	46	29	15	12	17	15
Male	31	43	42	38	22	9	8	9	11
Female	18	18	13	8	7	6	4	8	4
Asian or Pacific Islander 2/, both sexes	39	40	40	25	17	32	32	40	50
Male	32	31	29	19	9	18	23	22	28
Female	7	9	11	6	8	14	9	18	22

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Falls (W00-W19)									
All races, both sexes	238	370	409	462	448	501	595	1,065	1,641
Male	199	297	332	358	326	354	378	613	895
Female	39	73	77	104	122	147	217	452	746
White, both sexes	194	310	352	377	374	428	514	966	1,539
Male	163	253	290	289	269	305	326	563	833
Female	31	57	62	88	105	123	188	403	706
Black, both sexes	29	45	47	67	60	56	64	73	65
Male	23	34	37	54	47	42	41	37	39
Female	6	11	10	13	13	14	23	36	26
American Indian 1/, both sexes	7	6	5	11	7	5	-	6	9
Male	5	5	1	10	6	1	-	2	8
Female	2	1	4	1	1	4	-	4	1
Asian or Pacific Islander 2/, both sexes	8	9	5	7	7	12	17	20	28
Male	8	5	4	5	4	6	11	11	15
Female	-	4	1	2	3	6	6	9	13
Accidental discharge of firearms (W32-W34)									
All races, both sexes	75	78	38	40	36	14	24	17	17
Male	60	65	31	38	30	14	20	14	16
Female	15	13	7	2	6	-	4	3	1
White, both sexes	60	68	30	37	34	13	22	15	16
Male	49	58	26	35	28	13	19	12	15
Female	11	10	4	2	6	-	3	3	1
Black, both sexes	13	6	7	3	2	1	1	2	-
Male	9	5	4	3	2	1	-	2	-
Female	4	1	3	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	-	-	-	-	-	-	1
Male	2	2	-	-	-	-	-	-	1
Female	-	1	-	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
Accidental drowning and submersion (W65-W74)									
All races, both sexes	250	230	182	172	120	97	80	99	71
Male	212	194	156	143	102	77	63	70	38
Female	38	36	26	29	18	20	17	29	33
White, both sexes	205	180	147	130	105	82	70	86	58
Male	172	151	125	107	89	70	55	59	31
Female	33	29	22	23	16	12	15	27	27
Black, both sexes	36	43	23	29	9	8	7	8	9
Male	32	37	21	24	8	5	6	7	4
Female	4	6	2	5	1	3	1	1	5
American Indian 1/, both sexes	4	3	5	6	3	1	1	1	-
Male	4	3	4	5	3	1	1	1	-
Female	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	7	7	3	6	2	4	4
Male	4	3	6	7	2	1	1	3	3
Female	1	1	1	-	1	5	1	1	1
Accidental exposure to smoke, fire and flames (X00-X09)									
All races, both sexes	188	214	214	225	187	182	188	213	257
Male	133	135	145	158	126	115	107	125	136
Female	55	79	69	67	61	67	81	88	121
White, both sexes	134	157	155	175	129	131	139	158	189
Male	97	98	106	122	89	80	76	93	100
Female	37	59	49	53	40	51	63	65	89
Black, both sexes	44	55	52	48	53	49	45	53	64
Male	30	35	35	34	35	34	29	31	35
Female	14	20	17	14	18	15	16	22	29
American Indian 1/, both sexes	5	2	1	1	3	2	1	1	1
Male	2	2	1	1	1	1	-	1	1
Female	3	-	-	-	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	6	1	2	-	3	1	3
Male	4	-	3	1	1	-	2	-	-
Female	1	-	3	-	1	-	1	1	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races, both sexes	2,159	2,504	1,992	1,069	447	241	146	132	118
Male	1,554	1,830	1,455	768	260	141	79	68	49
Female	605	674	537	301	187	100	67	64	69
White, both sexes	1,811	2,031	1,568	790	349	194	122	111	111
Male	1,330	1,498	1,143	558	187	105	60	55	46
Female	481	533	425	232	162	89	62	56	65
Black, both sexes	317	429	389	266	92	43	24	19	7
Male	206	301	288	199	70	34	19	12	3
Female	111	128	101	67	22	9	5	7	4
American Indian 1/, both sexes	21	28	23	8	3	-	-	1	-
Male	10	17	16	8	2	-	-	-	-
Female	11	11	7	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	16	12	5	3	4	-	1	-
Male	8	14	8	3	1	2	-	1	-
Female	2	2	4	2	2	2	-	-	-

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races, both sexes	672	774	787	723	668	681	743	1,081	1,632
Male	577	628	646	573	504	467	474	607	855
Female	95	146	141	150	164	214	269	474	777
White, both sexes	559	636	623	593	547	570	613	923	1,443
Male	485	519	508	470	409	385	389	526	745
Female	74	117	115	123	138	185	224	397	698
Black, both sexes	92	109	134	105	106	94	111	136	170
Male	76	86	111	86	84	67	70	69	99
Female	16	23	23	19	22	27	41	67	71
American Indian 1/, both sexes	12	21	20	20	13	7	9	8	5
Male	10	16	19	14	10	6	6	5	2
Female	2	5	1	6	3	1	3	3	3
Asian or Pacific Islander 2/, both sexes	9	8	10	5	2	10	10	14	14
Male	6	7	8	3	1	9	9	7	9
Female	3	1	2	2	1	1	1	7	5

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (X60-X84,Y87.0)

All races, both sexes	3,112	3,450	2,987	2,450	1,708	1,237	1,073	1,219	1,227
Male	2,457	2,657	2,307	1,842	1,306	959	870	1,018	1,062
Female	655	793	680	608	402	278	203	201	165
White, both sexes	2,796	3,146	2,762	2,273	1,604	1,171	1,017	1,151	1,174
Male	2,197	2,423	2,128	1,706	1,225	908	823	961	1,020
Female	599	723	634	567	379	263	194	190	154
Black, both sexes	214	224	156	119	62	41	39	46	30
Male	180	176	129	90	51	31	36	40	25
Female	34	48	27	29	11	10	3	6	5
American Indian 1/, both sexes	37	30	16	15	13	3	3	5	1
Male	31	22	11	13	12	3	3	5	1
Female	6	8	5	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	50	53	43	29	22	14	17	22
Male	49	36	39	33	18	17	8	12	16
Female	16	14	14	10	11	5	6	5	6

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes	1,521	1,670	1,491	1,367	1,001	832	791	909	917
Male	1,255	1,402	1,263	1,146	854	714	698	830	856
Female	266	268	228	221	147	118	93	79	61
White, both sexes	1,384	1,549	1,390	1,290	941	791	759	867	894
Male	1,133	1,298	1,178	1,083	800	679	667	790	836
Female	251	251	212	207	141	112	92	77	58
Black, both sexes	101	93	77	61	39	33	28	34	19
Male	89	80	66	48	35	27	27	33	17
Female	12	13	11	13	4	6	1	1	2
American Indian 1/, both sexes	12	13	8	10	8	3	3	4	1
Male	10	10	4	9	8	3	3	4	1
Female	2	3	4	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	15	16	6	13	5	1	4	3
Male	23	14	15	6	11	5	1	3	2
Female	1	1	1	-	2	-	-	1	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races, both sexes	1,591	1,780	1,496	1,083	707	405	282	310	310
Male	1,202	1,255	1,044	696	452	245	172	188	206
Female	389	525	452	387	255	160	110	122	104
White, both sexes	1,412	1,597	1,372	983	663	380	258	284	280
Male	1,064	1,125	950	623	425	229	156	171	184
Female	348	472	422	360	238	151	102	113	96
Black, both sexes	113	131	79	58	23	8	11	12	11
Male	91	96	63	42	16	4	9	7	8
Female	22	35	16	16	7	4	2	5	3
American Indian 1/, both sexes	25	17	8	5	5	-	-	1	-
Male	21	12	7	4	4	-	-	1	-
Female	4	5	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	35	37	37	16	17	13	13	19
Male	26	22	24	27	7	12	7	9	14
Female	15	13	13	10	9	5	6	4	5

Assault (homicide) (X85-Y09,Y87.1)

All races, both sexes	1,732	1,487	1,059	696	455	283	233	210	172
Male	1,235	1,084	773	511	338	199	147	132	85
Female	497	403	286	185	117	84	86	78	87
White, both sexes	909	816	616	432	309	188	150	153	138
Male	614	588	423	295	224	126	86	90	67
Female	295	228	193	137	85	62	64	63	71
Black, both sexes	764	612	397	231	123	81	74	51	30
Male	590	450	315	193	97	63	58	37	16
Female	174	162	82	38	26	18	16	14	14
American Indian 1/, both sexes	24	22	9	9	2	4	2	2	2
Male	16	16	7	7	1	3	1	2	1
Female	8	6	2	2	1	1	1	-	1
Asian or Pacific Islander 2/, both sexes	35	37	37	24	21	10	7	4	2
Male	15	30	28	16	16	7	2	3	1
Female	20	7	9	8	5	3	5	1	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (X93-X95)

All races, both sexes	1,064	842	582	377	235	132	99	85	54
Male	842	645	445	285	172	94	58	50	23
Female	222	197	137	92	63	38	41	35	31
White, both sexes	525	460	338	229	165	95	70	64	47
Male	384	340	246	160	117	60	37	33	20
Female	141	120	92	69	48	35	33	31	27
Black, both sexes	509	354	223	131	56	29	25	20	7
Male	439	284	183	113	43	26	20	17	3
Female	70	70	40	18	13	3	5	3	4
American Indian 1/, both sexes	9	8	2	2	-	2	1	-	-
Male	7	4	2	1	-	2	-	-	-
Female	2	4	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	21	20	19	15	14	6	3	1	-
Male	12	17	14	11	12	6	1	-	-
Female	9	3	5	4	2	-	2	1	-

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races, both sexes	668	645	477	319	220	151	134	125	118
Male	393	439	328	226	166	105	89	82	62
Female	275	206	149	93	54	46	45	43	56
White, both sexes	384	356	278	203	144	93	80	89	91
Male	230	248	177	135	107	66	49	57	47
Female	154	108	101	68	37	27	31	32	44
Black, both sexes	255	258	174	100	67	52	49	31	23
Male	151	166	132	80	54	37	38	20	13
Female	104	92	42	20	13	15	11	11	10
American Indian 1/, both sexes	15	14	7	7	2	2	1	2	2
Male	9	12	5	6	1	1	1	2	1
Female	6	2	2	1	1	1	-	-	1
Asian or Pacific Islander 2/, both sexes	14	17	18	9	7	4	4	3	2
Male	3	13	14	5	4	1	1	3	1
Female	11	4	4	4	3	3	3	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Legal intervention (Y35,Y89.0)

All races, both sexes	60	41	38	17	6	4	2	3	1
Male	58	38	37	17	6	3	2	3	1
Female	2	3	1	-	-	1	-	-	-
White, both sexes	38	25	32	11	5	4	1	2	1
Male	37	25	32	11	5	3	1	2	1
Female	1	-	-	-	-	1	-	-	-
Black, both sexes	20	15	6	5	-	-	1	1	-
Male	20	13	5	5	-	-	1	1	-
Female	-	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races, both sexes	599	687	543	298	154	74	55	49	43
Male	415	470	374	196	105	47	38	32	29
Female	184	217	169	102	49	27	17	17	14
White, both sexes	492	539	432	240	116	61	45	39	38
Male	342	368	292	154	74	38	29	24	27
Female	150	171	140	86	42	23	16	15	11
Black, both sexes	98	136	103	53	31	12	8	7	4
Male	67	95	77	38	24	9	7	6	2
Female	31	41	26	15	7	3	1	1	2
American Indian 1/, both sexes	6	6	4	3	6	-	-	1	-
Male	5	4	3	2	6	-	-	-	-
Female	1	2	1	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	6	4	2	1	1	2	2	1
Male	1	3	2	2	1	-	2	2	-
Female	2	3	2	-	-	1	-	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races, both sexes	18	21	9	14	12	3	7	4	5
Male	12	18	6	12	10	1	5	4	4
Female	6	3	3	2	2	2	2	-	1
White, both sexes	13	18	8	12	8	3	6	3	5
Male	8	17	5	10	6	1	4	3	4
Female	5	1	3	2	2	2	2	-	1
Black, both sexes	4	3	1	1	3	-	1	1	-
Male	4	1	1	1	3	-	1	1	-
Female	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races, both sexes	581	666	534	284	142	71	48	45	38
Male	403	452	368	184	95	46	33	28	25
Female	178	214	166	100	47	25	15	17	13
White, both sexes	479	521	424	228	108	58	39	36	33
Male	334	351	287	144	68	37	25	21	23
Female	145	170	137	84	40	21	14	15	10
Black, both sexes	94	133	102	52	28	12	7	6	4
Male	63	94	76	37	21	9	6	5	2
Female	31	39	26	15	7	3	1	1	2
American Indian 1/, both sexes	6	6	4	3	5	-	-	1	-
Male	5	4	3	2	5	-	-	-	-
Female	1	2	1	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	6	4	1	1	1	2	2	1
Male	1	3	2	1	1	-	2	2	-
Female	1	3	2	-	-	1	-	-	1

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	35-39 YEARS	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All races, both sexes	-	-	-	2	-	1	1	2	4
Male	-	-	-	2	-	1	1	2	4
Female	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	2	4
Male	-	-	-	1	-	1	1	2	4
Female	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-

Complications of medical and surgical care (Y40-Y84,Y88)

All races, both sexes	65	84	128	167	157	219	263	384	427
Male	33	43	59	84	77	104	123	208	212
Female	32	41	69	83	80	115	140	176	215
White, both sexes	52	63	97	134	120	165	215	320	361
Male	27	37	45	70	64	80	98	175	186
Female	25	26	52	64	56	85	117	145	175
Black, both sexes	13	18	27	31	35	51	42	59	59
Male	6	6	12	13	12	22	24	28	22
Female	7	12	15	18	23	29	18	31	37
American Indian 1/, both sexes	-	1	-	1	-	-	4	1	1
Male	-	-	-	1	-	-	-	1	1
Female	-	1	-	-	-	-	4	-	-
Asian or Pacific Islander 2/, both sexes	-	2	4	1	2	3	2	4	6
Male	-	-	2	-	1	2	1	4	3
Female	-	2	2	1	1	1	1	-	3

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
All Causes						
All races, both sexes	362,745	335,813	219,283	84,641	18,434	356
Male	166,892	128,877	64,439	18,552	2,874	289
Female	195,853	206,936	154,844	66,089	15,560	67
White, both sexes	328,472	307,495	201,942	77,167	16,157	275
Male	151,919	118,321	59,009	16,624	2,455	231
Female	176,553	189,174	142,933	60,543	13,702	44
Black, both sexes	28,903	23,919	14,680	6,407	2,032	72
Male	12,344	8,428	4,249	1,532	351	50
Female	16,559	15,491	10,431	4,875	1,681	22
American Indian 1/, both sexes	932	728	442	169	43	3
Male	422	286	155	50	10	3
Female	510	442	287	119	33	-
Asian or Pacific Islander 2/, both sexes	4,438	3,671	2,219	898	202	6
Male	2,207	1,842	1,026	346	58	5
Female	2,231	1,829	1,193	552	144	1

Salmonella infections (A01-A02)

All races, both sexes	2	1	2	-	-	-
Male	-	-	1	-	-	-
Female	2	1	1	-	-	-
White, both sexes	2	1	2	-	-	-
Male	-	-	1	-	-	-
Female	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Shigellosis and amebiasis (A03,A06)						
All races, both sexes	5	-	-	-	1	-
Male	2	-	-	-	-	-
Female	3	-	-	-	1	-
White, both sexes	5	-	-	-	1	-
Male	2	-	-	-	-	-
Female	3	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Certain other intestinal infections (A04,A07-A09)						
All races, both sexes	260	267	168	53	11	-
Male	107	81	48	9	2	-
Female	153	186	120	44	9	-
White, both sexes	251	247	162	49	10	-
Male	101	76	46	8	2	-
Female	150	171	116	41	8	-
Black, both sexes	7	17	5	4	-	-
Male	5	4	2	1	-	-
Female	2	13	3	3	-	-
American Indian 1/, both sexes	-	1	-	-	-	-
Male	-	-	-	-	-	-
Female	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	-	1	-
Male	1	1	-	-	-	-
Female	1	1	1	-	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Tuberculosis (A16-A19)						
All races, both sexes	120	74	31	10	-	1
Male	70	38	13	5	-	1
Female	50	36	18	5	-	-
White, both sexes	76	53	26	6	-	-
Male	38	27	12	3	-	-
Female	38	26	14	3	-	-
Black, both sexes	31	13	2	4	-	1
Male	23	6	-	2	-	1
Female	8	7	2	2	-	-
American Indian 1/, both sexes	1	-	1	-	-	-
Male	1	-	-	-	-	-
Female	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	12	8	2	-	-	-
Male	8	5	1	-	-	-
Female	4	3	1	-	-	-
Respiratory tuberculosis (A16)						
All races, both sexes	102	64	26	8	-	1
Male	58	33	13	4	-	1
Female	44	31	13	4	-	-
White, both sexes	66	45	22	5	-	-
Male	31	24	12	2	-	-
Female	35	21	10	3	-	-
Black, both sexes	24	12	2	3	-	1
Male	19	5	-	2	-	1
Female	5	7	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	7	2	-	-	-
Male	8	4	1	-	-	-
Female	4	3	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other tuberculosis (A17-A19)						
All races, both sexes	18	10	5	2	-	-
Male	12	5	-	1	-	-
Female	6	5	5	1	-	-
White, both sexes	10	8	4	1	-	-
Male	7	3	-	1	-	-
Female	3	5	4	-	-	-
Black, both sexes	7	1	-	1	-	-
Male	4	1	-	-	-	-
Female	3	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	-	-	-
Male	1	-	-	-	-	-
Female	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Female	-	-	-	-	-	-
Whooping cough (A37)						
All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
----------------	----------------	----------------	----------------	-----------------------	-------------------

Scarlet fever and erysipelas (A38,A46)

All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

Meningococcal infection (A39)

All races, both sexes	8	2	4	1	-	-
Male	3	-	-	1	-	-
Female	5	2	4	-	-	-
White, both sexes	8	2	4	1	-	-
Male	3	-	-	1	-	-
Female	5	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Septicemia (A40-A41)						
All races, both sexes	5,193	4,900	2,949	1,079	216	1
Male	2,233	1,690	812	199	35	1
Female	2,960	3,210	2,137	880	181	-
White, both sexes	4,364	4,206	2,546	875	163	1
Male	1,904	1,448	692	161	25	1
Female	2,460	2,758	1,854	714	138	-
Black, both sexes	766	635	380	187	53	-
Male	297	218	109	34	10	-
Female	469	417	271	153	43	-
American Indian 1/, both sexes	15	12	4	6	-	-
Male	8	2	2	1	-	-
Female	7	10	2	5	-	-
Asian or Pacific Islander 2/, both sexes	48	47	19	11	-	-
Male	24	22	9	3	-	-
Female	24	25	10	8	-	-
Syphilis (A50-A53)						
All races, both sexes	6	6	3	1	-	-
Male	4	3	2	-	-	-
Female	2	3	1	1	-	-
White, both sexes	4	4	3	1	-	-
Male	2	1	2	-	-	-
Female	2	3	1	1	-	-
Black, both sexes	2	2	-	-	-	-
Male	2	2	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Acute poliomyelitis (A80)						
All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)						
All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Measles (B05)						
All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Viral hepatitis (B15-B19)						
All races, both sexes	259	123	35	10	-	1
Male	113	55	11	4	-	1
Female	146	68	24	6	-	-
White, both sexes	222	110	30	9	-	1
Male	98	49	11	3	-	1
Female	124	61	19	6	-	-
Black, both sexes	25	9	2	-	-	-
Male	12	4	-	-	-	-
Female	13	5	2	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-
Male	1	-	-	-	-	-
Female	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	10	4	2	1	-	-
Male	2	2	-	1	-	-
Female	8	2	2	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX:

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Human immunodeficiency virus (HIV) disease (B20-B24)						
All races, both sexes	16	4	3	2	-	1
Male	11	1	1	-	-	1
Female	5	3	2	2	-	-
White, both sexes	7	2	1	-	-	1
Male	5	-	-	-	-	1
Female	2	2	1	-	-	-
Black, both sexes	9	2	2	2	-	-
Male	6	1	1	-	-	-
Female	3	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Malaria (B50-B54)						
All races, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
----------------	----------------	----------------	----------------	-----------------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races, both sexes	636	513	265	93	28	-
Male	298	216	97	22	4	-
Female	338	297	168	71	24	-
White, both sexes	571	470	235	77	25	-
Male	269	197	81	17	3	-
Female	302	273	154	60	22	-
Black, both sexes	57	33	26	15	3	-
Male	24	10	12	5	1	-
Female	33	23	14	10	2	-
American Indian 1/, both sexes	1	1	-	-	-	-
Male	-	1	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	9	4	1	-	-
Male	5	8	4	-	-	-
Female	2	1	-	1	-	-

Malignant neoplasms (C00-C97)

All races, both sexes	74,444	49,155	21,564	5,653	764	11
Male	37,623	22,714	8,506	1,850	190	10
Female	36,821	26,441	13,058	3,803	574	1
White, both sexes	67,321	44,566	19,488	4,995	630	7
Male	33,865	20,503	7,611	1,600	149	7
Female	33,456	24,063	11,877	3,395	481	-
Black, both sexes	6,023	3,897	1,784	591	118	3
Male	3,183	1,832	742	219	38	2
Female	2,840	2,065	1,042	372	80	1
American Indian 1/, both sexes	164	103	46	16	4	-
Male	80	44	20	9	1	-
Female	84	59	26	7	3	-
Asian or Pacific Islander 2/, both sexes	936	589	246	51	12	1
Male	495	335	133	22	2	1
Female	441	254	113	29	10	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)						
All races, both sexes	693	488	249	74	13	-
Male	370	217	96	19	-	-
Female	323	271	153	55	13	-
White, both sexes	629	449	229	67	11	-
Male	332	202	86	18	-	-
Female	297	247	143	49	11	-
Black, both sexes	56	26	15	6	2	-
Male	34	8	8	-	-	-
Female	22	18	7	6	2	-
American Indian 1/, both sexes	-	3	-	-	-	-
Male	-	1	-	-	-	-
Female	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	10	5	1	-	-
Male	4	6	2	1	-	-
Female	4	4	3	-	-	-
Malignant neoplasm of esophagus (C15)						
All races, both sexes	1,346	764	274	55	7	-
Male	896	437	122	23	4	-
Female	450	327	152	32	3	-
White, both sexes	1,218	709	253	51	7	-
Male	830	412	112	19	4	-
Female	388	297	141	32	3	-
Black, both sexes	106	48	19	4	-	-
Male	54	21	8	4	-	-
Female	52	27	11	-	-	-
American Indian 1/, both sexes	6	1	1	-	-	-
Male	4	-	1	-	-	-
Female	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	6	1	-	-	-
Male	8	4	1	-	-	-
Female	8	2	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of stomach (C16)						
All races, both sexes	1,824	1,379	641	188	26	-
Male	943	617	247	56	8	-
Female	881	762	394	132	18	-
White, both sexes	1,465	1,129	525	149	13	-
Male	770	510	205	45	4	-
Female	695	619	320	104	9	-
Black, both sexes	266	197	86	30	12	-
Male	125	80	27	8	3	-
Female	141	117	59	22	9	-
American Indian 1/, both sexes	6	6	2	-	-	-
Male	2	3	1	-	-	-
Female	4	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	87	47	28	9	1	-
Male	46	24	14	3	1	-
Female	41	23	14	6	-	-
Malignant neoplasms of colon, rectum and anus (C18-C21)						
All races, both sexes	8,643	6,735	3,518	1,061	148	-
Male	3,908	2,571	1,012	260	30	-
Female	4,735	4,164	2,506	801	118	-
White, both sexes	7,710	6,090	3,180	933	129	-
Male	3,485	2,327	905	218	23	-
Female	4,225	3,763	2,275	715	106	-
Black, both sexes	801	561	295	117	18	-
Male	354	199	82	39	7	-
Female	447	362	213	78	11	-
American Indian 1/, both sexes	16	13	8	3	-	-
Male	7	6	3	1	-	-
Female	9	7	5	2	-	-
Asian or Pacific Islander 2/, both sexes	116	71	35	8	1	-
Male	62	39	22	2	-	-
Female	54	32	13	6	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms of liver and intrahepatic bile ducts (C22)						
All races, both sexes	1,547	943	359	81	4	-
Male	856	407	114	26	1	-
Female	691	536	245	55	3	-
White, both sexes	1,376	830	325	70	2	-
Male	766	352	105	21	-	-
Female	610	478	220	49	2	-
Black, both sexes	101	71	21	7	2	-
Male	54	34	4	3	1	-
Female	47	37	17	4	1	-
American Indian 1/, both sexes	11	8	1	1	-	-
Male	5	2	-	1	-	-
Female	6	6	1	-	-	-
Asian or Pacific Islander 2/, both sexes	59	34	12	3	-	-
Male	31	19	5	1	-	-
Female	28	15	7	2	-	-
Malignant neoplasm of pancreas (C25)						
All races, both sexes	4,236	2,571	1,196	286	34	1
Male	1,742	891	310	59	2	1
Female	2,494	1,680	886	227	32	-
White, both sexes	3,786	2,297	1,059	250	27	1
Male	1,570	812	272	49	2	1
Female	2,216	1,485	787	201	25	-
Black, both sexes	369	228	118	32	5	-
Male	138	60	27	8	-	-
Female	231	168	91	24	5	-
American Indian 1/, both sexes	10	3	2	1	1	-
Male	4	-	1	1	-	-
Female	6	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	71	43	17	3	1	-
Male	30	19	10	1	-	-
Female	41	24	7	2	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of larynx (C32)						
All races, both sexes	401	191	73	11	1	-
Male	314	155	51	8	-	-
Female	87	36	22	3	1	-
White, both sexes	358	171	70	10	1	-
Male	281	140	48	7	-	-
Female	77	31	22	3	1	-
Black, both sexes	41	19	2	1	-	-
Male	32	14	2	1	-	-
Female	9	5	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	-	-
Male	1	1	1	-	-	-
Female	1	-	-	-	-	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)						
All races, both sexes	18,531	9,302	2,902	542	60	5
Male	10,240	4,840	1,335	204	16	5
Female	8,291	4,462	1,567	338	44	-
White, both sexes	17,054	8,508	2,649	479	49	3
Male	9,356	4,398	1,206	177	11	3
Female	7,698	4,110	1,443	302	38	-
Black, both sexes	1,216	651	208	53	9	1
Male	732	355	105	23	5	1
Female	484	296	103	30	4	-
American Indian 1/, both sexes	38	21	3	2	-	-
Male	24	9	3	1	-	-
Female	14	12	-	1	-	-
Asian or Pacific Islander 2/, both sexes	223	122	42	8	2	1
Male	128	78	21	3	-	1
Female	95	44	21	5	2	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant melanoma of skin (C43)						
All races, both sexes	737	495	252	52	4	-
Male	454	277	100	13	-	-
Female	283	218	152	39	4	-
White, both sexes	717	483	243	50	4	-
Male	444	276	96	13	-	-
Female	273	207	147	37	4	-
Black, both sexes	18	12	7	2	-	-
Male	9	1	2	-	-	-
Female	9	11	5	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-
Male	1	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-
Male	-	-	2	-	-	-
Female	1	-	-	-	-	-
Malignant neoplasm of breast (C50)						
All races, both sexes	4,494	3,596	1,880	658	115	-
Male	44	36	13	1	1	-
Female	4,450	3,560	1,867	657	114	-
White, both sexes	4,107	3,267	1,696	586	102	-
Male	38	29	12	1	1	-
Female	4,069	3,238	1,684	585	101	-
Black, both sexes	356	307	173	71	13	-
Male	5	7	1	-	-	-
Female	351	300	172	71	13	-
American Indian 1/, both sexes	9	2	4	-	-	-
Male	-	-	-	-	-	-
Female	9	2	4	-	-	-
Asian or Pacific Islander 2/, both sexes	22	20	7	1	-	-
Male	1	-	-	-	-	-
Female	21	20	7	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of cervix uteri (C53)						
All races, both sexes	279	180	92	35	3	-
Female	279	180	92	35	3	-
White, both sexes	225	147	67	25	2	-
Female	225	147	67	25	2	-
Black, both sexes	48	30	22	10	1	-
Female	48	30	22	10	1	-
American Indian 1/, both sexes	-	1	1	-	-	-
Female	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	2	2	-	-	-
Female	6	2	2	-	-	-
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)						
All races, both sexes	919	617	333	84	15	1
Female	919	617	333	84	15	1
White, both sexes	785	557	294	74	11	-
Female	785	557	294	74	11	-
Black, both sexes	124	56	36	10	4	1
Female	124	56	36	10	4	1
American Indian 1/, both sexes	2	-	1	-	-	-
Female	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	8	4	2	-	-	-
Female	8	4	2	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of ovary (C56)						
All races, both sexes	1,803	1,176	453	98	9	-
Female	1,803	1,176	453	98	9	-
White, both sexes	1,684	1,099	418	86	8	-
Female	1,684	1,099	418	86	8	-
Black, both sexes	97	66	31	12	1	-
Female	97	66	31	12	1	-
American Indian 1/, both sexes	3	2	-	-	-	-
Female	3	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	9	4	-	-	-
Female	19	9	4	-	-	-
Malignant neoplasm of prostate (C61)						
All races, both sexes	6,757	5,532	2,638	628	75	-
Male	6,757	5,532	2,638	628	75	-
White, both sexes	5,637	4,785	2,257	524	56	-
Male	5,637	4,785	2,257	524	56	-
Black, both sexes	1,037	678	348	96	18	-
Male	1,037	678	348	96	18	-
American Indian 1/, both sexes	15	10	6	2	1	-
Male	15	10	6	2	1	-
Asian or Pacific Islander 2/, both sexes	68	59	27	6	-	-
Male	68	59	27	6	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms of kidney and renal pelvis (C64-C65)						
All races, both sexes	1,463	945	397	83	7	-
Male	811	477	165	31	3	-
Female	652	468	232	52	4	-
White, both sexes	1,344	880	361	76	6	-
Male	744	444	152	29	2	-
Female	600	436	209	47	4	-
Black, both sexes	103	57	30	7	1	-
Male	58	30	10	2	1	-
Female	45	27	20	5	-	-
American Indian 1/, both sexes	1	3	1	-	-	-
Male	-	1	-	-	-	-
Female	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	15	5	5	-	-	-
Male	9	2	3	-	-	-
Female	6	3	2	-	-	-
Malignant neoplasm of bladder (C67)						
All races, both sexes	2,252	1,790	875	256	31	-
Male	1,512	1,110	478	110	12	-
Female	740	680	397	146	19	-
White, both sexes	2,079	1,665	830	242	27	-
Male	1,414	1,037	460	106	12	-
Female	665	628	370	136	15	-
Black, both sexes	158	103	41	12	3	-
Male	87	59	15	3	-	-
Female	71	44	26	9	3	-
American Indian 1/, both sexes	2	1	1	-	-	-
Male	2	1	1	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	21	3	2	1	-
Male	9	13	2	1	-	-
Female	4	8	1	1	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)						
All races, both sexes	1,041	521	166	28	2	-
Male	482	212	53	9	1	-
Female	559	309	113	19	1	-
White, both sexes	978	486	157	27	1	-
Male	459	201	51	9	-	-
Female	519	285	106	18	1	-
Black, both sexes	51	26	6	1	1	-
Male	18	6	2	-	1	-
Female	33	20	4	1	-	-
American Indian 1/, both sexes	2	1	-	-	-	-
Male	1	1	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	8	3	-	-	-
Male	4	4	-	-	-	-
Female	6	4	3	-	-	-

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races, both sexes	8,705	5,716	2,329	559	70	-
Male	4,231	2,475	783	185	15	-
Female	4,474	3,241	1,546	374	55	-
White, both sexes	8,129	5,292	2,145	511	59	-
Male	3,976	2,309	725	169	14	-
Female	4,153	2,983	1,420	342	45	-
Black, both sexes	466	356	159	38	8	-
Male	199	132	49	13	1	-
Female	267	224	110	25	7	-
American Indian 1/, both sexes	23	13	7	4	-	-
Male	11	5	1	2	-	-
Female	12	8	6	2	-	-
Asian or Pacific Islander 2/, both sexes	87	55	18	6	3	-
Male	45	29	8	1	-	-
Female	42	26	10	5	3	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Hodgkin's disease (C81)						
All races, both sexes	121	70	24	10	-	-
Male	50	30	10	5	-	-
Female	71	40	14	5	-	-
White, both sexes	114	68	22	9	-	-
Male	47	29	9	5	-	-
Female	67	39	13	4	-	-
Black, both sexes	5	1	2	-	-	-
Male	3	-	1	-	-	-
Female	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	-	-
Male	-	1	-	-	-	-
Female	2	-	-	1	-	-
Non-Hodgkin's lymphoma (C82-C85)						
All races, both sexes	3,551	2,330	856	193	23	-
Male	1,616	959	274	62	4	-
Female	1,935	1,371	582	131	19	-
White, both sexes	3,396	2,212	817	185	21	-
Male	1,548	917	260	59	4	-
Female	1,848	1,295	557	126	17	-
Black, both sexes	100	80	30	6	-	-
Male	38	23	12	3	-	-
Female	62	57	18	3	-	-
American Indian 1/, both sexes	15	4	1	-	-	-
Male	8	1	-	-	-	-
Female	7	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	40	34	8	2	2	-
Male	22	18	2	-	-	-
Female	18	16	6	2	2	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Leukemia (C91-C95)						
All races, both sexes	3,218	2,194	1,041	271	37	-
Male	1,713	1,018	359	85	8	-
Female	1,505	1,176	682	186	29	-
White, both sexes	3,026	2,053	967	250	32	-
Male	1,616	949	338	76	7	-
Female	1,410	1,104	629	174	25	-
Black, both sexes	155	124	64	18	5	-
Male	76	61	16	7	1	-
Female	79	63	48	11	4	-
American Indian 1/, both sexes	5	6	3	1	-	-
Male	3	2	1	1	-	-
Female	2	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes	32	11	7	2	-	-
Male	18	6	4	1	-	-
Female	14	5	3	1	-	-
Multiple myeloma and immunoproliferative neoplasms (C88,C90)						
All races, both sexes	1,799	1,109	401	84	10	-
Male	845	462	137	33	3	-
Female	954	647	264	51	7	-
White, both sexes	1,578	947	333	66	6	-
Male	758	409	116	29	3	-
Female	820	538	217	37	3	-
Black, both sexes	205	150	62	14	3	-
Male	82	47	19	3	-	-
Female	123	103	43	11	3	-
American Indian 1/, both sexes	3	3	3	3	-	-
Male	-	2	-	1	-	-
Female	3	1	3	2	-	-
Asian or Pacific Islander 2/, both sexes	13	9	3	1	1	-
Male	5	4	2	-	-	-
Female	8	5	1	1	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)						
All races, both sexes	16	13	7	1	-	-
Male	7	6	3	-	-	-
Female	9	7	4	1	-	-
White, both sexes	15	12	6	1	-	-
Male	7	5	2	-	-	-
Female	8	7	4	1	-	-
Black, both sexes	1	1	1	-	-	-
Male	-	1	1	-	-	-
Female	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, both sexes	8,773	6,214	2,937	874	140	4
Male	4,063	2,460	989	218	22	4
Female	4,710	3,754	1,948	656	118	-
White, both sexes	8,040	5,722	2,730	785	115	3
Male	3,763	2,269	919	195	20	3
Female	4,277	3,453	1,811	590	95	-
Black, both sexes	609	405	167	82	20	1
Male	247	148	52	19	1	1
Female	362	257	115	63	19	-
American Indian 1/, both sexes	19	15	8	3	2	-
Male	4	5	3	1	-	-
Female	15	10	5	2	2	-
Asian or Pacific Islander 2/, both sexes	105	72	32	4	3	-
Male	49	38	15	3	1	-
Female	56	34	17	1	2	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
----------------	----------------	----------------	----------------	-----------------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, both sexes	2,503	2,103	1,143	314	40	1
Male	1,199	881	362	86	7	1
Female	1,304	1,222	781	228	33	-
White, both sexes	2,340	1,960	1,066	291	35	-
Male	1,136	818	341	82	7	-
Female	1,204	1,142	725	209	28	-
Black, both sexes	134	118	63	20	4	1
Male	46	45	18	3	-	1
Female	88	73	45	17	4	-
American Indian 1/, both sexes	2	6	1	1	1	-
Male	2	2	-	-	-	-
Female	-	4	1	1	1	-
Asian or Pacific Islander 2/, both sexes	27	19	13	2	-	-
Male	15	16	3	1	-	-
Female	12	3	10	1	-	-

Anemias (D50-D64)

All races, both sexes	689	790	638	290	78	-
Male	267	237	183	58	14	-
Female	422	553	455	232	64	-
White, both sexes	617	718	566	261	68	-
Male	238	212	156	52	13	-
Female	379	506	410	209	55	-
Black, both sexes	64	64	61	28	10	-
Male	23	20	22	5	1	-
Female	41	44	39	23	9	-
American Indian 1/, both sexes	2	1	3	-	-	-
Male	1	-	-	-	-	-
Female	1	1	3	-	-	-
Asian or Pacific Islander 2/, both sexes	6	7	8	1	-	-
Male	5	5	5	1	-	-
Female	1	2	3	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Diabetes mellitus (E10-E14)						
All races, both sexes	10,577	8,002	4,104	1,250	200	-
Male	4,450	2,789	1,165	260	20	-
Female	6,127	5,213	2,939	990	180	-
White, both sexes	8,917	6,905	3,525	1,067	148	-
Male	3,911	2,473	1,023	228	16	-
Female	5,006	4,432	2,502	839	132	-
Black, both sexes	1,434	962	507	160	49	-
Male	454	260	109	21	2	-
Female	980	702	398	139	47	-
American Indian 1/, both sexes	52	36	13	2	2	-
Male	13	15	5	1	1	-
Female	39	21	8	1	1	-
Asian or Pacific Islander 2/, both sexes	174	99	59	21	1	-
Male	72	41	28	10	1	-
Female	102	58	31	11	-	-
Nutritional deficiencies (E40-E64)						
All races, both sexes	780	947	781	302	90	-
Male	264	289	182	52	13	-
Female	516	658	599	250	77	-
White, both sexes	687	836	712	255	71	-
Male	233	256	167	41	11	-
Female	454	580	545	214	60	-
Black, both sexes	80	96	61	41	16	-
Male	27	31	12	11	1	-
Female	53	65	49	30	15	-
American Indian 1/, both sexes	3	8	3	1	3	-
Male	1	-	1	-	1	-
Female	2	8	2	1	2	-
Asian or Pacific Islander 2/, both sexes	10	7	5	5	-	-
Male	3	2	2	-	-	-
Female	7	5	3	5	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Malnutrition (E40-E46)						
All races, both sexes	721	888	712	280	84	-
Male	240	275	165	46	13	-
Female	481	613	547	234	71	-
White, both sexes	634	781	650	235	67	-
Male	209	242	152	35	11	-
Female	425	539	498	200	56	-
Black, both sexes	74	94	57	39	14	-
Male	27	31	10	11	1	-
Female	47	63	47	28	13	-
American Indian 1/, both sexes	3	6	2	1	3	-
Male	1	-	1	-	1	-
Female	2	6	1	1	2	-
Asian or Pacific Islander 2/, both sexes	10	7	3	5	-	-
Male	3	2	2	-	-	-
Female	7	5	1	5	-	-
Other nutritional deficiencies (E50-E64)						
All races, both sexes	59	59	69	22	6	-
Male	24	14	17	6	-	-
Female	35	45	52	16	6	-
White, both sexes	53	55	62	20	4	-
Male	24	14	15	6	-	-
Female	29	41	47	14	4	-
Black, both sexes	6	2	4	2	2	-
Male	-	-	2	-	-	-
Female	6	2	2	2	2	-
American Indian 1/, both sexes	-	2	1	-	-	-
Male	-	-	-	-	-	-
Female	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-
Male	-	-	-	-	-	-
Female	-	-	2	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Meningitis (G00,G03)						
All races, both sexes	55	32	16	2	-	-
Male	21	13	5	1	-	-
Female	34	19	11	1	-	-
White, both sexes	48	24	14	2	-	-
Male	19	9	4	1	-	-
Female	29	15	10	1	-	-
Black, both sexes	5	7	2	-	-	-
Male	2	4	1	-	-	-
Female	3	3	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-
Male	-	-	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-
Male	-	-	-	-	-	-
Female	1	1	-	-	-	-
Parkinson's disease (G20-G21)						
All races, both sexes	4,330	3,567	1,627	369	27	1
Male	2,478	1,879	726	115	8	1
Female	1,852	1,688	901	254	19	-
White, both sexes	4,143	3,418	1,560	350	26	1
Male	2,364	1,804	689	109	7	1
Female	1,779	1,614	871	241	19	-
Black, both sexes	133	102	47	15	1	-
Male	78	44	24	6	1	-
Female	55	58	23	9	-	-
American Indian 1/, both sexes	11	6	5	2	-	-
Male	7	4	4	-	-	-
Female	4	2	1	2	-	-
Asian or Pacific Islander 2/, both sexes	43	41	15	2	-	-
Male	29	27	9	-	-	-
Female	14	14	6	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Alzheimer's disease (G30)						
All races, both sexes	10,934	13,337	10,257	4,014	701	2
Male	3,795	3,687	2,101	564	54	1
Female	7,139	9,650	8,156	3,450	647	1
White, both sexes	10,260	12,562	9,723	3,765	633	1
Male	3,583	3,466	1,974	525	45	1
Female	6,677	9,096	7,749	3,240	588	-
Black, both sexes	608	682	464	219	58	1
Male	185	181	99	36	6	-
Female	423	501	365	183	52	1
American Indian 1/, both sexes	18	19	15	5	-	-
Male	7	4	4	1	-	-
Female	11	15	11	4	-	-
Asian or Pacific Islander 2/, both sexes	48	74	55	25	10	-
Male	20	36	24	2	3	-
Female	28	38	31	23	7	-
Major cardiovascular diseases (I00-I78)						
All races, both sexes	160,906	165,313	117,092	47,427	10,618	62
Male	72,601	60,379	32,104	9,636	1,539	43
Female	88,305	104,934	84,988	37,791	9,079	19
White, both sexes	145,286	151,317	108,139	43,327	9,357	45
Male	66,225	55,631	29,586	8,639	1,320	34
Female	79,061	95,686	78,553	34,688	8,037	11
Black, both sexes	13,174	11,860	7,551	3,492	1,133	17
Male	5,217	3,778	1,942	758	188	9
Female	7,957	8,082	5,609	2,734	945	8
American Indian 1/, both sexes	388	322	211	90	18	-
Male	176	123	60	23	2	-
Female	212	199	151	67	16	-
Asian or Pacific Islander 2/, both sexes	2,058	1,814	1,191	518	110	-
Male	983	847	516	216	29	-
Female	1,075	967	675	302	81	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Diseases of heart (I00-I09,I11,I13,I20-I51)						
All races, both sexes	118,296	121,003	86,424	35,605	8,210	59
Male	55,272	45,756	24,475	7,430	1,235	41
Female	63,024	75,247	61,949	28,175	6,975	18
White, both sexes	107,189	111,070	80,035	32,666	7,279	44
Male	50,626	42,311	22,657	6,708	1,059	33
Female	56,563	68,759	57,378	25,958	6,220	11
Black, both sexes	9,469	8,478	5,414	2,503	834	15
Male	3,827	2,776	1,408	548	153	8
Female	5,642	5,702	4,006	1,955	681	7
American Indian 1/, both sexes	271	231	150	65	13	-
Male	126	91	46	18	-	-
Female	145	140	104	47	13	-
Asian or Pacific Islander 2/, both sexes	1,367	1,224	825	371	84	-
Male	693	578	364	156	23	-
Female	674	646	461	215	61	-

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races, both sexes	575	416	231	54	11	-
Male	144	96	40	13	1	-
Female	431	320	191	41	10	-
White, both sexes	554	400	216	50	11	-
Male	140	93	36	11	1	-
Female	414	307	180	39	10	-
Black, both sexes	15	12	11	4	-	-
Male	1	3	3	2	-	-
Female	14	9	8	2	-	-
American Indian 1/, both sexes	2	2	-	-	-	-
Male	1	-	-	-	-	-
Female	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	4	-	-	-
Male	2	-	1	-	-	-
Female	2	2	3	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Hypertensive heart disease (I11)						
All races, both sexes	3,268	3,684	2,886	1,330	286	3
Male	1,134	960	545	181	21	2
Female	2,134	2,724	2,341	1,149	265	1
White, both sexes	2,614	3,122	2,487	1,132	236	1
Male	893	813	456	147	18	1
Female	1,721	2,309	2,031	985	218	-
Black, both sexes	584	509	360	172	46	2
Male	212	134	77	24	2	1
Female	372	375	283	148	44	1
American Indian 1/, both sexes	11	6	2	2	-	-
Male	5	1	1	1	-	-
Female	6	5	1	1	-	-
Asian or Pacific Islander 2/, both sexes	59	47	37	24	4	-
Male	24	12	11	9	1	-
Female	35	35	26	15	3	-
Hypertensive heart and renal disease (I13)						
All races, both sexes	487	494	361	153	24	-
Male	198	188	109	37	7	-
Female	289	306	252	116	17	-
White, both sexes	361	399	305	129	12	-
Male	153	150	90	28	2	-
Female	208	249	215	101	10	-
Black, both sexes	110	82	50	19	10	-
Male	40	29	16	6	4	-
Female	70	53	34	13	6	-
American Indian 1/, both sexes	1	4	-	-	-	-
Male	1	1	-	-	-	-
Female	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	9	6	5	2	-
Male	4	8	3	3	1	-
Female	11	1	3	2	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Ischemic heart diseases (I20-I25)						
All races, both sexes	87,123	86,890	60,269	24,477	5,683	48
Male	42,295	34,024	17,603	5,216	847	34
Female	44,828	52,866	42,666	19,261	4,836	14
White, both sexes	79,215	79,909	55,931	22,506	5,046	36
Male	38,901	31,567	16,346	4,726	729	27
Female	40,314	48,342	39,585	17,780	4,317	9
Black, both sexes	6,668	5,891	3,644	1,676	568	12
Male	2,748	1,939	957	369	104	7
Female	3,920	3,952	2,687	1,307	464	5
American Indian 1/, both sexes	192	156	113	45	7	-
Male	93	64	38	15	-	-
Female	99	92	75	30	7	-
Asian or Pacific Islander 2/, both sexes	1,048	934	581	250	62	-
Male	553	454	262	106	14	-
Female	495	480	319	144	48	-
Acute myocardial infarction (I21-I22)						
All races, both sexes	32,151	29,922	18,701	6,755	1,367	8
Male	15,336	11,765	5,701	1,539	238	5
Female	16,815	18,157	13,000	5,216	1,129	3
White, both sexes	29,233	27,383	17,270	6,112	1,174	7
Male	14,094	10,840	5,246	1,363	210	4
Female	15,139	16,543	12,024	4,749	964	3
Black, both sexes	2,475	2,151	1,207	542	178	1
Male	1,016	727	348	128	27	1
Female	1,459	1,424	859	414	151	-
American Indian 1/, both sexes	70	50	38	15	2	-
Male	33	22	12	7	-	-
Female	37	28	26	8	2	-
Asian or Pacific Islander 2/, both sexes	373	338	186	86	13	-
Male	193	176	95	41	1	-
Female	180	162	91	45	12	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other acute ischemic heart diseases (I24)						
All races, both sexes	466	439	318	136	35	1
Male	214	153	68	28	5	1
Female	252	286	250	108	30	-
White, both sexes	417	400	288	130	32	-
Male	193	142	60	27	4	-
Female	224	258	228	103	28	-
Black, both sexes	43	35	26	6	3	1
Male	18	9	7	1	1	1
Female	25	26	19	5	2	-
American Indian 1/, both sexes	4	1	2	-	-	-
Male	1	1	1	-	-	-
Female	3	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	2	-	-	-
Male	2	1	-	-	-	-
Female	-	2	2	-	-	-
Other forms of chronic ischemic heart disease (I20,I25)						
All races, both sexes	54,506	56,529	41,250	17,586	4,281	39
Male	26,745	22,106	11,834	3,649	604	28
Female	27,761	34,423	29,416	13,937	3,677	11
White, both sexes	49,565	52,126	38,373	16,264	3,840	29
Male	24,614	20,585	11,040	3,336	515	23
Female	24,951	31,541	27,333	12,928	3,325	6
Black, both sexes	4,150	3,705	2,411	1,128	387	10
Male	1,714	1,203	602	240	76	5
Female	2,436	2,502	1,809	888	311	5
American Indian 1/, both sexes	118	105	73	30	5	-
Male	59	41	25	8	-	-
Female	59	64	48	22	5	-
Asian or Pacific Islander 2/, both sexes	673	593	393	164	49	-
Male	358	277	167	65	13	-
Female	315	316	226	99	36	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Atherosclerotic cardiovascular disease, so described (I25.0)						
All races, both sexes	10,274	10,397	7,953	3,429	885	19
Male	4,639	3,740	2,205	691	122	15
Female	5,635	6,657	5,748	2,738	763	4
White, both sexes	8,892	9,245	7,202	3,112	760	13
Male	4,037	3,348	1,997	611	93	12
Female	4,855	5,897	5,205	2,501	667	1
Black, both sexes	1,204	1,010	663	280	114	6
Male	505	340	175	67	26	3
Female	699	670	488	213	88	3
American Indian 1/, both sexes	25	30	15	5	1	-
Male	15	9	6	1	-	-
Female	10	21	9	4	1	-
Asian or Pacific Islander 2/, both sexes	153	112	73	32	10	-
Male	82	43	27	12	3	-
Female	71	69	46	20	7	-

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, both sexes	44,232	46,132	33,297	14,157	3,396	20
Male	22,106	18,366	9,629	2,958	482	13
Female	22,126	27,766	23,668	11,199	2,914	7
White, both sexes	40,673	42,881	31,171	13,152	3,080	16
Male	20,577	17,237	9,043	2,725	422	11
Female	20,096	25,644	22,128	10,427	2,658	5
Black, both sexes	2,946	2,695	1,748	848	273	4
Male	1,209	863	427	173	50	2
Female	1,737	1,832	1,321	675	223	2
American Indian 1/, both sexes	93	75	58	25	4	-
Male	44	32	19	7	-	-
Female	49	43	39	18	4	-
Asian or Pacific Islander 2/, both sexes	520	481	320	132	39	-
Male	276	234	140	53	10	-
Female	244	247	180	79	29	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other heart diseases (I26-I51)						
All races, both sexes	26,843	29,519	22,677	9,591	2,206	8
Male	11,501	10,488	6,178	1,983	359	5
Female	15,342	19,031	16,499	7,608	1,847	3
White, both sexes	24,445	27,240	21,096	8,849	1,974	7
Male	10,539	9,688	5,729	1,796	309	5
Female	13,906	17,552	15,367	7,053	1,665	2
Black, both sexes	2,092	1,984	1,349	632	210	1
Male	826	671	355	147	43	-
Female	1,266	1,313	994	485	167	1
American Indian 1/, both sexes	65	63	35	18	6	-
Male	26	25	7	2	-	-
Female	39	38	28	16	6	-
Asian or Pacific Islander 2/, both sexes	241	232	197	92	16	-
Male	110	104	87	38	7	-
Female	131	128	110	54	9	-
Acute and subacute endocarditis (I33)						
All races, both sexes	136	83	30	5	1	-
Male	74	40	9	2	-	-
Female	62	43	21	3	1	-
White, both sexes	122	74	26	5	-	-
Male	66	37	8	2	-	-
Female	56	37	18	3	-	-
Black, both sexes	10	6	4	-	1	-
Male	5	1	1	-	-	-
Female	5	5	3	-	1	-
American Indian 1/, both sexes	1	1	-	-	-	-
Male	1	-	-	-	-	-
Female	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	-	-	-
Male	2	2	-	-	-	-
Female	1	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Diseases of pericardium and acute myocarditis (I30-I31,I40)						
All races, both sexes	63	43	24	8	1	-
Male	31	17	7	1	1	-
Female	32	26	17	7	-	-
White, both sexes	61	38	21	8	1	-
Male	31	15	6	1	1	-
Female	30	23	15	7	-	-
Black, both sexes	2	5	2	-	-	-
Male	-	2	-	-	-	-
Female	2	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-
Male	-	-	1	-	-	-
Female	-	-	-	-	-	-
Heart failure (I50)						
All races, both sexes	9,884	12,315	10,543	4,969	1,208	4
Male	4,212	4,256	2,819	982	191	2
Female	5,672	8,059	7,724	3,987	1,017	2
White, both sexes	9,038	11,433	9,847	4,628	1,098	4
Male	3,872	3,961	2,618	905	166	2
Female	5,166	7,472	7,229	3,723	932	2
Black, both sexes	742	783	608	292	102	-
Male	288	255	167	60	22	-
Female	454	528	441	232	80	-
American Indian 1/, both sexes	29	28	14	8	2	-
Male	13	12	2	1	-	-
Female	16	16	12	7	2	-
Asian or Pacific Islander 2/, both sexes	75	71	74	41	6	-
Male	39	28	32	16	3	-
Female	36	43	42	25	3	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)						
All races, both sexes	16,760	17,078	12,080	4,609	996	4
Male	7,184	6,175	3,343	998	167	3
Female	9,576	10,903	8,737	3,611	829	1
White, both sexes	15,224	15,695	11,202	4,208	875	3
Male	6,570	5,675	3,097	888	142	3
Female	8,654	10,020	8,105	3,320	733	-
Black, both sexes	1,338	1,190	735	340	107	1
Male	533	413	187	87	21	-
Female	805	777	548	253	86	1
American Indian 1/, both sexes	35	34	21	10	4	-
Male	12	13	5	1	-	-
Female	23	21	16	9	4	-
Asian or Pacific Islander 2/, both sexes	163	159	122	51	10	-
Male	69	74	54	22	4	-
Female	94	85	68	29	6	-

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races, both sexes	3,116	3,273	2,375	1,017	242	-
Male	1,155	991	569	155	32	-
Female	1,961	2,282	1,806	862	210	-
White, both sexes	2,573	2,772	2,056	856	196	-
Male	960	839	483	123	24	-
Female	1,613	1,933	1,573	733	172	-
Black, both sexes	462	440	283	142	41	-
Male	163	127	75	27	5	-
Female	299	313	208	115	36	-
American Indian 1/, both sexes	7	13	4	2	2	-
Male	2	3	1	1	1	-
Female	5	10	3	1	1	-
Asian or Pacific Islander 2/, both sexes	74	48	32	17	3	-
Male	30	22	10	4	2	-
Female	44	26	22	13	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Cerebrovascular diseases (I60-I69)						
All races, both sexes	32,342	34,030	23,205	8,530	1,611	2
Male	12,754	10,864	5,588	1,564	196	1
Female	19,588	23,166	17,617	6,966	1,415	1
White, both sexes	28,982	30,998	21,356	7,711	1,397	1
Male	11,464	9,892	5,095	1,367	172	1
Female	17,518	21,106	16,261	6,344	1,225	-
Black, both sexes	2,725	2,492	1,514	688	197	1
Male	1,024	730	369	145	21	-
Female	1,701	1,762	1,145	543	176	1
American Indian 1/, both sexes	92	66	47	16	-	-
Male	39	24	12	3	-	-
Female	53	42	35	13	-	-
Asian or Pacific Islander 2/, both sexes	543	474	288	115	17	-
Male	227	218	112	49	3	-
Female	316	256	176	66	14	-
Atherosclerosis (I70)						
All races, both sexes	2,592	3,188	2,870	1,518	429	-
Male	1,097	1,090	738	283	55	-
Female	1,495	2,098	2,132	1,235	374	-
White, both sexes	2,375	2,984	2,693	1,431	383	-
Male	1,008	1,015	687	261	48	-
Female	1,367	1,969	2,006	1,170	335	-
Black, both sexes	191	173	153	76	40	-
Male	79	63	41	19	5	-
Female	112	110	112	57	35	-
American Indian 1/, both sexes	10	8	3	4	2	-
Male	5	3	-	-	1	-
Female	5	5	3	4	1	-
Asian or Pacific Islander 2/, both sexes	16	23	21	7	4	-
Male	5	9	10	3	1	-
Female	11	14	11	4	3	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other diseases of circulatory system (I71-I78)						
All races, both sexes	4,560	3,819	2,218	757	126	1
Male	2,323	1,678	734	204	21	1
Female	2,237	2,141	1,484	553	105	-
White, both sexes	4,167	3,493	1,999	663	102	-
Male	2,167	1,574	664	180	17	-
Female	2,000	1,919	1,335	483	85	-
Black, both sexes	327	277	187	83	21	1
Male	124	82	49	19	4	1
Female	203	195	138	64	17	-
American Indian 1/, both sexes	8	4	7	3	1	-
Male	4	2	1	1	-	-
Female	4	2	6	2	1	-
Asian or Pacific Islander 2/, both sexes	58	45	25	8	2	-
Male	28	20	20	4	-	-
Female	30	25	5	4	2	-
Aortic aneurysm and dissection (I71)						
All races, both sexes	2,792	2,134	985	213	25	-
Male	1,558	1,063	425	68	6	-
Female	1,234	1,071	560	145	19	-
White, both sexes	2,603	1,988	900	188	23	-
Male	1,480	1,015	389	62	6	-
Female	1,123	973	511	126	17	-
Black, both sexes	143	114	64	20	1	-
Male	54	35	21	4	-	-
Female	89	79	43	16	1	-
American Indian 1/, both sexes	3	-	3	1	-	-
Male	2	-	1	-	-	-
Female	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	43	32	18	4	1	-
Male	22	13	14	2	-	-
Female	21	19	4	2	1	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other diseases of arteries, arterioles and capillaries (I72-I78)						
All races, both sexes	1,768	1,685	1,233	544	101	1
Male	765	615	309	136	15	1
Female	1,003	1,070	924	408	86	-
White, both sexes	1,564	1,505	1,099	475	79	-
Male	687	559	275	118	11	-
Female	877	946	824	357	68	-
Black, both sexes	184	163	123	63	20	1
Male	70	47	28	15	4	1
Female	114	116	95	48	16	-
American Indian 1/, both sexes	5	4	4	2	1	-
Male	2	2	-	1	-	-
Female	3	2	4	1	1	-
Asian or Pacific Islander 2/, both sexes	15	13	7	4	1	-
Male	6	7	6	2	-	-
Female	9	6	1	2	1	-
Other disorders of circulatory system (I80-I99)						
All races, both sexes	588	571	343	149	21	-
Male	227	175	72	26	-	-
Female	361	396	271	123	21	-
White, both sexes	529	516	309	132	21	-
Male	205	158	64	22	-	-
Female	324	358	245	110	21	-
Black, both sexes	50	51	30	15	-	-
Male	16	14	8	4	-	-
Female	34	37	22	11	-	-
American Indian 1/, both sexes	3	-	1	-	-	-
Male	2	-	-	-	-	-
Female	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	3	2	-	-
Male	4	3	-	-	-	-
Female	2	1	3	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Influenza and pneumonia (J10-J18)						
All races, both sexes	11,909	13,752	11,318	5,116	1,361	3
Male	5,667	5,575	3,724	1,373	273	2
Female	6,242	8,177	7,594	3,743	1,088	1
White, both sexes	10,783	12,677	10,480	4,728	1,216	1
Male	5,119	5,128	3,415	1,261	241	1
Female	5,664	7,549	7,065	3,467	975	-
Black, both sexes	869	827	657	315	122	2
Male	408	325	219	86	25	1
Female	461	502	438	229	97	1
American Indian 1/, both sexes	34	36	36	9	3	-
Male	15	11	16	4	-	-
Female	19	25	20	5	3	-
Asian or Pacific Islander 2/, both sexes	223	212	145	64	20	-
Male	125	111	74	22	7	-
Female	98	101	71	42	13	-
Influenza (J10-J11)						
All races, both sexes	303	372	305	130	48	-
Male	115	127	86	35	6	-
Female	188	245	219	95	42	-
White, both sexes	289	353	295	125	47	-
Male	112	119	83	32	6	-
Female	177	234	212	93	41	-
Black, both sexes	8	17	8	4	1	-
Male	1	7	1	3	-	-
Female	7	10	7	1	1	-
American Indian 1/, both sexes	-	1	1	-	-	-
Male	-	-	1	-	-	-
Female	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	1	1	-	-
Male	2	1	1	-	-	-
Female	4	-	-	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Pneumonia (J12-J18)						
All races, both sexes	11,606	13,380	11,013	4,986	1,313	3
Male	5,552	5,448	3,638	1,338	267	2
Female	6,054	7,932	7,375	3,648	1,046	1
White, both sexes	10,494	12,324	10,185	4,603	1,169	1
Male	5,007	5,009	3,332	1,229	235	1
Female	5,487	7,315	6,853	3,374	934	-
Black, both sexes	861	810	649	311	121	2
Male	407	318	218	83	25	1
Female	454	492	431	228	96	1
American Indian 1/, both sexes	34	35	35	9	3	-
Male	15	11	15	4	-	-
Female	19	24	20	5	3	-
Asian or Pacific Islander 2/, both sexes	217	211	144	63	20	-
Male	123	110	73	22	7	-
Female	94	101	71	41	13	-
Other acute lower respiratory infections (J20-J22)						
All races, both sexes	41	80	77	23	12	-
Male	14	26	15	6	-	-
Female	27	54	62	17	12	-
White, both sexes	39	75	72	22	12	-
Male	12	24	12	6	-	-
Female	27	51	60	16	12	-
Black, both sexes	2	5	4	1	-	-
Male	2	2	2	-	-	-
Female	-	3	2	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-
Male	-	-	1	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Acute bronchitis and bronchiolitis (J20-J21)						
All races, both sexes	23	45	44	15	7	-
Male	9	14	10	5	-	-
Female	14	31	34	10	7	-
White, both sexes	22	42	42	14	7	-
Male	8	14	8	5	-	-
Female	14	28	34	9	7	-
Black, both sexes	1	3	2	1	-	-
Male	1	-	2	-	-	-
Female	-	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Unspecified acute lower respiratory infection (J22)						
All races, both sexes	18	35	33	8	5	-
Male	5	12	5	1	-	-
Female	13	23	28	7	5	-
White, both sexes	17	33	30	8	5	-
Male	4	10	4	1	-	-
Female	13	23	26	7	5	-
Black, both sexes	1	2	2	-	-	-
Male	1	2	-	-	-	-
Female	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-
Male	-	-	1	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Chronic lower respiratory diseases (J40-J47)						
All races, both sexes	23,413	17,085	7,823	2,238	350	2
Male	11,533	7,843	3,165	776	92	2
Female	11,880	9,242	4,658	1,462	258	-
White, both sexes	22,193	16,186	7,360	2,070	310	1
Male	10,836	7,340	2,938	702	78	1
Female	11,357	8,846	4,422	1,368	232	-
Black, both sexes	961	671	360	133	31	-
Male	532	360	173	60	8	-
Female	429	311	187	73	23	-
American Indian 1/, both sexes	60	49	15	3	4	-
Male	38	24	8	1	3	-
Female	22	25	7	2	1	-
Asian or Pacific Islander 2/, both sexes	199	179	88	32	5	1
Male	127	119	46	13	3	1
Female	72	60	42	19	2	-
Bronchitis, chronic and unspecified (J40-J42)						
All races, both sexes	177	236	161	80	24	-
Male	90	92	46	19	1	-
Female	87	144	115	61	23	-
White, both sexes	166	229	157	77	23	-
Male	87	88	46	18	1	-
Female	79	141	111	59	22	-
Black, both sexes	10	5	3	3	1	-
Male	3	2	-	1	-	-
Female	7	3	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	-	-	-
Male	-	2	-	-	-	-
Female	1	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Emphysema (J43)						
All races, both sexes	2,989	1,832	685	150	18	-
Male	1,534	898	317	58	5	-
Female	1,455	934	368	92	13	-
White, both sexes	2,851	1,731	644	134	16	-
Male	1,453	827	293	47	3	-
Female	1,398	904	351	87	13	-
Black, both sexes	109	74	31	14	-	-
Male	63	47	20	9	-	-
Female	46	27	11	5	-	-
American Indian 1/, both sexes	5	5	1	-	1	-
Male	2	4	1	-	1	-
Female	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	22	9	2	1	-
Male	16	20	3	2	1	-
Female	8	2	6	-	-	-
Asthma (J45-J46)						
All races, both sexes	429	388	227	81	11	-
Male	123	104	39	15	-	-
Female	306	284	188	66	11	-
White, both sexes	378	338	195	69	8	-
Male	105	87	30	9	-	-
Female	273	251	165	60	8	-
Black, both sexes	35	32	27	9	3	-
Male	11	9	8	5	-	-
Female	24	23	19	4	3	-
American Indian 1/, both sexes	1	2	-	-	-	-
Male	1	1	-	-	-	-
Female	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	16	5	3	-	-
Male	6	7	1	1	-	-
Female	9	9	4	2	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Other chronic lower respiratory diseases (J44,J47)						
All races, both sexes	19,818	14,629	6,750	1,927	297	2
Male	9,786	6,749	2,763	684	86	2
Female	10,032	7,880	3,987	1,243	211	-
White, both sexes	18,798	13,888	6,364	1,790	263	1
Male	9,191	6,338	2,569	628	74	1
Female	9,607	7,550	3,795	1,162	189	-
Black, both sexes	807	560	299	107	27	-
Male	455	302	145	45	8	-
Female	352	258	154	62	19	-
American Indian 1/, both sexes	54	42	14	3	3	-
Male	35	19	7	1	2	-
Female	19	23	7	2	1	-
Asian or Pacific Islander 2/, both sexes	159	139	73	27	4	1
Male	105	90	42	10	2	1
Female	54	49	31	17	2	-
Pneumoconioses and chemical effects (J60-J66,J68)						
All races, both sexes	235	217	80	20	-	-
Male	226	208	79	19	-	-
Female	9	9	1	1	-	-
White, both sexes	222	210	76	18	-	-
Male	214	201	75	17	-	-
Female	8	9	1	1	-	-
Black, both sexes	10	7	3	1	-	-
Male	9	7	3	1	-	-
Female	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-
Male	-	-	1	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	1	-	-
Male	3	-	-	1	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Pneumonitis due to solids and liquids (J69)						
All races, both sexes	3,247	3,825	2,779	1,111	239	-
Male	1,747	1,839	1,116	367	65	-
Female	1,500	1,986	1,663	744	174	-
White, both sexes	2,942	3,532	2,562	1,026	211	-
Male	1,607	1,720	1,028	341	60	-
Female	1,335	1,812	1,534	685	151	-
Black, both sexes	253	236	180	74	21	-
Male	114	90	67	21	4	-
Female	139	146	113	53	17	-
American Indian 1/, both sexes	8	8	5	1	1	-
Male	3	4	2	1	-	-
Female	5	4	3	-	1	-
Asian or Pacific Islander 2/, both sexes	44	49	32	10	6	-
Male	23	25	19	4	1	-
Female	21	24	13	6	5	-
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)						
All races, both sexes	4,413	3,394	1,850	629	135	3
Male	2,062	1,409	612	156	28	2
Female	2,351	1,985	1,238	473	107	1
White, both sexes	4,085	3,193	1,711	576	120	1
Male	1,921	1,328	557	138	21	1
Female	2,164	1,865	1,154	438	99	-
Black, both sexes	252	148	111	42	13	2
Male	98	54	41	15	5	1
Female	154	94	70	27	8	1
American Indian 1/, both sexes	20	11	8	1	-	-
Male	6	5	3	1	-	-
Female	14	6	5	-	-	-
Asian or Pacific Islander 2/, both sexes	56	42	20	10	2	-
Male	37	22	11	2	2	-
Female	19	20	9	8	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Peptic ulcer (K25-K28)						
All races, both sexes	773	801	498	177	33	-
Male	326	291	150	49	9	-
Female	447	510	348	128	24	-
White, both sexes	717	731	468	164	29	-
Male	298	263	138	42	8	-
Female	419	468	330	122	21	-
Black, both sexes	39	50	23	8	4	-
Male	19	18	5	3	1	-
Female	20	32	18	5	3	-
American Indian 1/, both sexes	3	1	-	-	-	-
Male	2	-	-	-	-	-
Female	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	19	7	5	-	-
Male	7	10	7	4	-	-
Female	7	9	-	1	-	-
Diseases of appendix (K35-K38)						
All races, both sexes	68	39	24	4	1	-
Male	30	15	9	-	-	-
Female	38	24	15	4	1	-
White, both sexes	58	38	22	4	1	-
Male	25	14	8	-	-	-
Female	33	24	14	4	1	-
Black, both sexes	7	1	2	-	-	-
Male	3	1	1	-	-	-
Female	4	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-
Male	2	-	-	-	-	-
Female	1	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Hernia (K40-K46)						
All races, both sexes	242	294	188	63	10	-
Male	104	96	42	12	4	-
Female	138	198	146	51	6	-
White, both sexes	223	270	176	57	9	-
Male	95	88	39	10	4	-
Female	128	182	137	47	5	-
Black, both sexes	16	24	12	3	1	-
Male	8	8	3	-	-	-
Female	8	16	9	3	1	-
American Indian 1/, both sexes	1	-	-	-	-	-
Male	-	-	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	3	-	-
Male	1	-	-	2	-	-
Female	1	-	-	1	-	-
Chronic liver disease and cirrhosis (K70,K73-K74)						
All races, both sexes	1,462	742	203	30	4	3
Male	634	325	83	12	1	3
Female	828	417	120	18	3	-
White, both sexes	1,368	697	188	29	4	2
Male	600	305	76	12	1	2
Female	768	392	112	17	3	-
Black, both sexes	62	30	13	1	-	1
Male	23	12	5	-	-	1
Female	39	18	8	1	-	-
American Indian 1/, both sexes	6	2	-	-	-	-
Male	1	1	-	-	-	-
Female	5	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	13	2	-	-	-
Male	10	7	2	-	-	-
Female	16	6	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Alcoholic liver disease (K70)						
All races, both sexes	218	76	17	2	-	2
Male	142	52	12	1	-	2
Female	76	24	5	1	-	-
White, both sexes	194	70	16	2	-	1
Male	130	48	11	1	-	1
Female	64	22	5	1	-	-
Black, both sexes	15	4	1	-	-	1
Male	9	3	1	-	-	1
Female	6	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-
Male	1	-	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	2	-	-	-	-
Male	2	1	-	-	-	-
Female	5	1	-	-	-	-
Other chronic liver disease and cirrhosis (K73-K74)						
All races, both sexes	1,244	666	186	28	4	1
Male	492	273	71	11	1	1
Female	752	393	115	17	3	-
White, both sexes	1,174	627	172	27	4	1
Male	470	257	65	11	1	1
Female	704	370	107	16	3	-
Black, both sexes	47	26	12	1	-	-
Male	14	9	4	-	-	-
Female	33	17	8	1	-	-
American Indian 1/, both sexes	4	2	-	-	-	-
Male	-	1	-	-	-	-
Female	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	11	2	-	-	-
Male	8	6	2	-	-	-
Female	11	5	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Cholelithiasis and other disorders of gallbladder (K80-K82)						
All races, both sexes	507	528	358	143	24	-
Male	237	204	105	27	3	-
Female	270	324	253	116	21	-
White, both sexes	454	489	335	135	23	-
Male	220	193	99	26	2	-
Female	234	296	236	109	21	-
Black, both sexes	39	33	18	6	-	-
Male	12	9	3	-	-	-
Female	27	24	15	6	-	-
American Indian 1/, both sexes	3	3	1	1	-	-
Male	1	-	1	-	-	-
Female	2	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	11	3	4	1	1	-
Male	4	2	2	1	1	-
Female	7	1	2	-	-	-
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)						
All races, both sexes	6,565	6,150	3,915	1,443	269	2
Male	3,174	2,672	1,473	453	73	2
Female	3,391	3,478	2,442	990	196	-
White, both sexes	5,613	5,390	3,447	1,258	216	2
Male	2,786	2,398	1,320	387	66	2
Female	2,827	2,992	2,127	871	150	-
Black, both sexes	858	671	413	171	44	-
Male	344	234	126	60	5	-
Female	514	437	287	111	39	-
American Indian 1/, both sexes	24	11	13	2	1	-
Male	9	5	6	2	-	-
Female	15	6	7	-	1	-
Asian or Pacific Islander 2/, both sexes	70	78	42	12	8	-
Male	35	35	21	4	2	-
Female	35	43	21	8	6	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)						
All races, both sexes	36	27	13	8	1	-
Male	15	13	5	4	-	-
Female	21	14	8	4	1	-
White, both sexes	34	23	12	8	1	-
Male	14	10	4	4	-	-
Female	20	13	8	4	1	-
Black, both sexes	2	2	-	-	-	-
Male	1	1	-	-	-	-
Female	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-
Male	-	1	1	-	-	-
Female	-	-	-	-	-	-
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07,N26)						
All races, both sexes	94	78	23	15	2	-
Male	38	40	9	6	-	-
Female	56	38	14	9	2	-
White, both sexes	83	64	17	12	1	-
Male	36	34	6	4	-	-
Female	47	30	11	8	1	-
Black, both sexes	10	9	4	2	-	-
Male	2	4	2	1	-	-
Female	8	5	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-
Male	-	-	-	-	-	-
Female	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	5	2	1	-	-
Male	-	2	1	1	-	-
Female	1	3	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Renal failure (N17-N19)						
All races, both sexes	6,429	6,040	3,878	1,420	265	2
Male	3,121	2,617	1,458	443	73	2
Female	3,308	3,423	2,420	977	192	-
White, both sexes	5,491	5,299	3,417	1,238	214	2
Male	2,736	2,352	1,309	379	66	2
Female	2,755	2,947	2,108	859	148	-
Black, both sexes	846	659	409	169	43	-
Male	341	229	124	59	5	-
Female	505	430	285	110	38	-
American Indian 1/, both sexes	24	10	13	2	-	-
Male	9	4	6	2	-	-
Female	15	6	7	-	-	-
Asian or Pacific Islander 2/, both sexes	68	72	39	11	8	-
Male	35	32	19	3	2	-
Female	33	40	20	8	6	-
Other disorders of kidney (N25,N27)						
All races, both sexes	6	5	1	-	1	-
Male	-	2	1	-	-	-
Female	6	3	-	-	1	-
White, both sexes	5	4	1	-	-	-
Male	-	2	1	-	-	-
Female	5	2	-	-	-	-
Black, both sexes	-	1	-	-	1	-
Male	-	-	-	-	-	-
Female	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-
Male	-	-	-	-	-	-
Female	1	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Infections of kidney (N10-N12,N13.6,N15.1)						
All races, both sexes	134	145	114	44	17	-
Male	40	32	30	5	2	-
Female	94	113	84	39	15	-
White, both sexes	114	134	104	38	15	-
Male	35	32	27	4	2	-
Female	79	102	77	34	13	-
Black, both sexes	12	10	8	6	1	-
Male	3	-	3	1	-	-
Female	9	10	5	5	1	-
American Indian 1/, both sexes	1	-	1	-	1	-
Male	-	-	-	-	-	-
Female	1	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	7	1	1	-	-	-
Male	2	-	-	-	-	-
Female	5	1	1	-	-	-
Hyperplasia of prostate (N40)						
All races, both sexes	95	106	83	29	4	1
Male	95	106	83	29	4	1
White, both sexes	88	96	74	27	4	-
Male	88	96	74	27	4	-
Black, both sexes	7	7	7	1	-	1
Male	7	7	7	1	-	1
American Indian 1/, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	1	-	-
Male	-	2	2	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Inflammatory diseases of female pelvic organs (N70-N76)						
All races, both sexes	12	11	12	3	1	-
Female	12	11	12	3	1	-
White, both sexes	11	9	12	2	1	-
Female	11	9	12	2	1	-
Black, both sexes	1	2	-	-	-	-
Female	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-
Female	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Female	-	-	-	-	-	-

Pregnancy, childbirth and the puerperium (000-099)

All races, both sexes	-
Female	-
White, both sexes	-
Female	-
Black, both sexes	-
Female	-
American Indian 1/, both sexes	-
Female	-
Asian or Pacific Islander 2/, both sexes	-
Female	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Pregnancy with abortive outcome (000-007)						
All races, both sexes	-
Female	-
White, both sexes	-
Female	-
Black, both sexes	-
Female	-
American Indian 1/, both sexes	-
Female	-
Asian or Pacific Islander 2/, both sexes	-
Female	-
Other complications of pregnancy, childbirth and the puerperium (010-099)						
All races, both sexes	-
Female	-
White, both sexes	-
Female	-
Black, both sexes	-
Female	-
American Indian 1/, both sexes	-
Female	-
Asian or Pacific Islander 2/, both sexes	-
Female	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
----------------	----------------	----------------	----------------	-----------------------	-------------------

Certain conditions originating in the perinatal period (P00-P96)

All races, both sexes	-	-	-	-	-	1
Male	-	-	-	-	-	-
Female	-	-	-	-	-	1
White, both sexes	-	-	-	-	-	1
Male	-	-	-	-	-	-
Female	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes	246	178	97	39	6	1
Male	98	74	41	13	2	-
Female	148	104	56	26	4	1
White, both sexes	232	168	94	34	5	1
Male	94	70	40	12	1	-
Female	138	98	54	22	4	1
Black, both sexes	11	7	3	5	1	-
Male	4	3	1	1	1	-
Female	7	4	2	4	-	-
American Indian 1/, both sexes	1	-	-	-	-	-
Male	-	-	-	-	-	-
Female	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	-	-	-	-
Male	-	1	-	-	-	-
Female	2	2	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX80-84 85-89 90-94 95-99 100 YEARS AGE NOT
YEARS YEARS YEARS YEARS AND OVER STATED

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, both sexes	2,794	3,286	3,014	1,798	726	115
Male	1,144	1,078	733	331	100	90
Female	1,650	2,208	2,281	1,467	626	25
White, both sexes	2,509	3,015	2,802	1,680	647	83
Male	1,036	988	687	308	88	66
Female	1,473	2,027	2,115	1,372	559	17
Black, both sexes	260	239	184	95	78	28
Male	99	76	39	14	11	21
Female	161	163	145	81	67	7
American Indian 1/, both sexes	6	11	8	8	-	1
Male	2	4	2	3	-	1
Female	4	7	6	5	-	-
Asian or Pacific Islander 2/, both sexes	19	21	20	15	1	3
Male	7	10	5	6	1	2
Female	12	11	15	9	-	1

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes	26,594	28,517	21,457	9,192	2,126	11
Male	9,865	8,712	4,966	1,593	265	11
Female	16,729	19,805	16,491	7,599	1,861	-
White, both sexes	24,061	26,146	19,763	8,434	1,861	8
Male	8,912	7,958	4,509	1,441	225	8
Female	15,149	18,188	15,254	6,993	1,636	-
Black, both sexes	2,192	2,067	1,474	659	240	3
Male	822	622	362	130	35	3
Female	1,370	1,445	1,112	529	205	-
American Indian 1/, both sexes	78	62	43	17	3	-
Male	33	24	17	2	2	-
Female	45	38	26	15	1	-
Asian or Pacific Islander 2/, both sexes	263	242	177	82	22	-
Male	98	108	78	20	3	-
Female	165	134	99	62	19	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Accidents (unintentional injuries) (V01-X59,Y85-Y86)						
All races, both sexes	6,162	5,933	3,939	1,419	304	85
Male	3,024	2,563	1,391	397	62	83
Female	3,138	3,370	2,548	1,022	242	2
White, both sexes	5,701	5,570	3,693	1,317	272	79
Male	2,781	2,400	1,290	357	53	77
Female	2,920	3,170	2,403	960	219	2
Black, both sexes	354	280	191	80	28	5
Male	185	121	79	32	7	5
Female	169	159	112	48	21	-
American Indian 1/, both sexes	22	13	6	3	2	1
Male	12	9	2	1	-	1
Female	10	4	4	2	2	-
Asian or Pacific Islander 2/, both sexes	85	70	49	19	2	-
Male	46	33	20	7	2	-
Female	39	37	29	12	-	-

Transport accidents (V01-V99,Y85)

All races, both sexes	1,482	985	298	49	3	33
Male	808	539	167	27	1	33
Female	674	446	131	22	2	-
White, both sexes	1,377	914	275	44	3	31
Male	743	503	154	25	1	31
Female	634	411	121	19	2	-
Black, both sexes	74	50	16	3	-	1
Male	47	27	10	2	-	1
Female	27	23	6	1	-	-
American Indian 1/, both sexes	5	3	2	1	-	1
Male	4	2	1	-	-	1
Female	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	26	18	5	1	-	-
Male	14	7	2	-	-	-
Female	12	11	3	1	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)						
All races, both sexes	1,409	953	285	47	3	26
Male	759	517	161	27	1	26
Female	650	436	124	20	2	-
White, both sexes	1,308	885	262	43	3	24
Male	696	482	148	25	1	24
Female	612	403	114	18	2	-
Black, both sexes	71	47	16	2	-	1
Male	46	26	10	2	-	1
Female	25	21	6	-	-	-
American Indian 1/, both sexes	5	3	2	1	-	1
Male	4	2	1	-	-	1
Female	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	25	18	5	1	-	-
Male	13	7	2	-	-	-
Female	12	11	3	1	-	-
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)						
All races, both sexes	40	22	12	1	-	7
Male	24	14	5	-	-	7
Female	16	8	7	1	-	-
White, both sexes	36	20	12	-	-	7
Male	22	14	5	-	-	7
Female	14	6	7	-	-	-
Black, both sexes	3	2	-	1	-	-
Male	1	-	-	-	-	-
Female	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-
Male	1	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX80-84 85-89 90-94 95-99 100 YEARS AGE NOT
YEARS YEARS YEARS YEARS AND OVER STATED

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races, both sexes	33	10	1	1	-	-
Male	25	8	1	-	-	-
Female	8	2	-	1	-	-
White, both sexes	33	9	1	1	-	-
Male	25	7	1	-	-	-
Female	8	2	-	1	-	-
Black, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

Nontransport accidents (W00-X59,Y86)

All races, both sexes	4,680	4,948	3,641	1,370	301	52
Male	2,216	2,024	1,224	370	61	50
Female	2,464	2,924	2,417	1,000	240	2
White, both sexes	4,324	4,656	3,418	1,273	269	48
Male	2,038	1,897	1,136	332	52	46
Female	2,286	2,759	2,282	941	217	2
Black, both sexes	280	230	175	77	28	4
Male	138	94	69	30	7	4
Female	142	136	106	47	21	-
American Indian 1/, both sexes	17	10	4	2	2	-
Male	8	7	1	1	-	-
Female	9	3	3	1	2	-
Asian or Pacific Islander 2/, both sexes	59	52	44	18	2	-
Male	32	26	18	7	2	-
Female	27	26	26	11	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Falls (W00-W19)						
All races, both sexes	2,200	2,352	1,682	618	120	-
Male	1,052	992	592	173	27	-
Female	1,148	1,360	1,090	445	93	-
White, both sexes	2,088	2,262	1,613	584	111	-
Male	997	951	559	159	25	-
Female	1,091	1,311	1,054	425	86	-
Black, both sexes	72	50	46	21	7	-
Male	31	20	23	9	1	-
Female	41	30	23	12	6	-
American Indian 1/, both sexes	3	6	2	2	1	-
Male	2	5	1	1	-	-
Female	1	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	37	34	21	11	1	-
Male	22	16	9	4	1	-
Female	15	18	12	7	-	-
Accidental discharge of firearms (W32-W34)						
All races, both sexes	7	7	4	-	-	-
Male	7	7	4	-	-	-
Female	-	-	-	-	-	-
White, both sexes	6	7	4	-	-	-
Male	6	7	4	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-
Male	1	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Accidental drowning and submersion (W65-W74)						
All races, both sexes	85	37	19	8	-	24
Male	50	22	7	5	-	23
Female	35	15	12	3	-	1
White, both sexes	79	34	16	6	-	23
Male	50	20	7	3	-	22
Female	29	14	9	3	-	1
Black, both sexes	3	3	2	2	-	1
Male	-	2	-	2	-	1
Female	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-
Male	-	-	-	-	-	-
Female	3	-	1	-	-	-
Accidental exposure to smoke, fire and flames (X00-X09)						
All races, both sexes	215	158	78	24	3	5
Male	108	69	34	11	1	5
Female	107	89	44	13	2	-
White, both sexes	156	129	51	17	-	5
Male	81	61	22	8	-	5
Female	75	68	29	9	-	-
Black, both sexes	57	29	26	7	3	-
Male	26	8	12	3	1	-
Female	31	21	14	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-
Male	1	-	-	-	-	-
Female	1	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Accidental poisoning and exposure to noxious substances (X40-X49)						
All races, both sexes	127	109	60	14	1	7
Male	53	51	23	2	-	6
Female	74	58	37	12	1	1
White, both sexes	118	92	54	11	-	6
Male	49	43	21	1	-	5
Female	69	49	33	10	-	1
Black, both sexes	8	16	6	3	1	1
Male	4	8	2	1	-	1
Female	4	8	4	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-
Male	-	-	-	-	-	-
Female	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-
Male	-	-	-	-	-	-
Female	1	-	-	-	-	-

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races, both sexes	2,046	2,285	1,798	706	177	16
Male	946	883	564	179	33	16
Female	1,100	1,402	1,234	527	144	-
White, both sexes	1,877	2,132	1,680	655	158	14
Male	855	815	523	161	27	14
Female	1,022	1,317	1,157	494	131	-
Black, both sexes	139	132	95	44	17	2
Male	76	56	32	15	5	2
Female	63	76	63	29	12	-
American Indian 1/, both sexes	14	3	2	-	1	-
Male	6	2	-	-	-	-
Female	8	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	16	18	21	7	1	-
Male	9	10	9	3	1	-
Female	7	8	12	4	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Intentional self-harm (suicide) (X60-X84,Y87.0)						
All races, both sexes	954	590	205	34	4	7
Male	822	507	168	26	4	7
Female	132	83	37	8	-	-
White, both sexes	916	574	198	32	2	6
Male	797	495	162	24	2	6
Female	119	79	36	8	-	-
Black, both sexes	23	9	2	-	1	-
Male	15	8	2	-	1	-
Female	8	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	7	5	2	1	1
Male	10	4	4	2	1	1
Female	5	3	1	-	-	-

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes	694	410	129	18	1	5
Male	649	388	119	18	1	5
Female	45	22	10	-	-	-
White, both sexes	679	403	126	17	1	4
Male	637	381	116	17	1	4
Female	42	22	10	-	-	-
Black, both sexes	13	6	2	-	-	-
Male	10	6	2	-	-	-
Female	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	-	1
Male	2	1	1	1	-	1
Female	-	-	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

80-84 YEARS 85-89 YEARS 90-94 YEARS 95-99 YEARS 100 YEARS AND OVER AGE NOT STATED

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races, both sexes	260	180	76	16	3	2
Male	173	119	49	8	3	2
Female	87	61	27	8	-	-
White, both sexes	237	171	72	15	1	2
Male	160	114	46	7	1	2
Female	77	57	26	8	-	-
Black, both sexes	10	3	-	-	1	-
Male	5	2	-	-	1	-
Female	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	6	4	1	1	-
Male	8	3	3	1	1	-
Female	5	3	1	-	-	-

Assault (homicide) (X85-Y09,Y87.1)

All races, both sexes	129	68	23	8	1	30
Male	69	25	10	5	-	15
Female	60	43	13	3	1	15
White, both sexes	89	57	18	7	1	25
Male	42	20	7	4	-	13
Female	47	37	11	3	1	12
Black, both sexes	34	8	5	1	-	4
Male	23	3	3	1	-	1
Female	11	5	2	-	-	3
American Indian 1/, both sexes	-	2	-	-	-	1
Male	-	2	-	-	-	1
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	-	-	-	-
Male	4	-	-	-	-	-
Female	2	1	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Assault (homicide) by discharge of firearms (X93-X95)						
All races, both sexes	27	23	4	1	-	12
Male	14	6	2	1	-	7
Female	13	17	2	-	-	5
White, both sexes	17	19	4	1	-	10
Male	5	5	2	1	-	5
Female	12	14	2	-	-	5
Black, both sexes	10	4	-	-	-	1
Male	9	1	-	-	-	1
Female	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1
Male	-	-	-	-	-	1
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races, both sexes	102	45	19	7	1	18
Male	55	19	8	4	-	8
Female	47	26	11	3	1	10
White, both sexes	72	38	14	6	1	15
Male	37	15	5	3	-	8
Female	35	23	9	3	1	7
Black, both sexes	24	4	5	1	-	3
Male	14	2	3	1	-	-
Female	10	2	2	-	-	3
American Indian 1/, both sexes	-	2	-	-	-	-
Male	-	2	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	-	-	-	-
Male	4	-	-	-	-	-
Female	2	1	-	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Legal intervention (Y35,Y89.0)						
All races, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-
Male	-	1	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)						
All races, both sexes	36	26	25	5	1	11
Male	20	14	5	2	-	11
Female	16	12	20	3	1	-
White, both sexes	34	22	21	5	-	8
Male	18	11	4	2	-	8
Female	16	11	17	3	-	-
Black, both sexes	2	3	3	-	1	3
Male	2	2	1	-	-	3
Female	-	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-
Male	-	1	-	-	-	-
Female	-	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Discharge of firearms, undetermined intent (Y22-Y24)						
All races, both sexes	1	2	-	-	-	1
Male	1	2	-	-	-	1
Female	-	-	-	-	-	-
White, both sexes	1	2	-	-	-	1
Male	1	2	-	-	-	1
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)						
All races, both sexes	35	24	25	5	1	10
Male	19	12	5	2	-	10
Female	16	12	20	3	1	-
White, both sexes	33	20	21	5	-	7
Male	17	9	4	2	-	7
Female	16	11	17	3	-	-
Black, both sexes	2	3	3	-	1	3
Male	2	2	1	-	-	3
Female	-	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-
Male	-	1	-	-	-	-
Female	-	-	1	-	-	-

TABLE 210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(DATA FOR SPECIFIED RACES OTHER THAN WHITE AND BLACK SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING RACE ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	80-84 YEARS	85-89 YEARS	90-94 YEARS	95-99 YEARS	100 YEARS AND OVER	AGE NOT STATED
Operations of war and their sequelae (Y36,Y89.1)						
All races, both sexes	5	1	-	-	-	-
Male	5	1	-	-	-	-
Female	-	-	-	-	-	-
White, both sexes	5	1	-	-	-	-
Male	5	1	-	-	-	-
Female	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-
Male	-	-	-	-	-	-
Female	-	-	-	-	-	-

Complications of medical and surgical care (Y40-Y84,Y88)

All races, both sexes	398	337	176	54	11	-
Male	190	134	48	13	1	-
Female	208	203	128	41	10	-
White, both sexes	357	297	155	41	10	-
Male	177	119	44	11	1	-
Female	180	178	111	30	9	-
Black, both sexes	37	34	20	12	1	-
Male	12	12	3	1	-	-
Female	25	22	17	11	1	-
American Indian 1/, both sexes	2	3	-	-	-	-
Male	1	1	-	-	-	-
Female	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	1	-	-
Male	-	2	1	1	-	-
Female	2	1	-	-	-	-

- QUANTITY ZERO.

... CATEGORY NOT APPLICABLE.

1/ INCLUDES ALEUTS AND ESKIMOS.

2/ INCLUDES CHINESE, FILIPINO, HAWAIIAN, JAPANESE, AND OTHER ASIAN OR PACIFIC ISLANDER.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.