

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
All races 3/, both sexes	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7
Male	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9
Female	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5
White, both sexes	2.0	3.4	0.4	0.2	0.2	0.2	0.3	0.2	0.3	0.6
Male	2.1	3.9	0.4	*	*	*	0.3	*	0.3	0.7
Female	1.9	2.9	0.4	*	*	*	*	*	0.3	0.4
Black, both sexes	2.3	9.9	*	*	*	*	*	*	1.2	1.6
Male	2.5	9.9	*	*	*	*	*	*	*	2.2
Female	2.1	9.9	*	*	*	*	*	*	*	*
Malignant neoplasms (C00-C97)										
All races 3/, both sexes	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
Male	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
Female	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
White, both sexes	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
Male	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
Female	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
Black, both sexes	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
Male	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
Female	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes	4.5	*	*	*	*	*	*	0.2	0.4	0.7
Male	5.3	*	*	*	*	*	*	0.2	0.5	0.9
Female	3.7	*	*	*	*	*	*	*	0.4	0.6
White, both sexes	4.2	*	*	*	*	*	*	0.1	0.3	0.6
Male	5.0	*	*	*	*	*	*	*	0.4	0.7
Female	3.5	*	*	*	*	*	*	*	0.3	0.5
Black, both sexes	5.9	*	*	*	*	*	*	*	0.9	1.3
Male	7.0	*	*	*	*	*	*	*	*	1.9
Female	5.0	*	*	*	*	*	*	*	*	*
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All races 3/, both sexes	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
Male	20.6	*	*	*	*	*	*	0.4	1.1	2.4
Female	20.2	*	*	*	*	*	*	0.4	0.8	1.8
White, both sexes	21.5	*	*	*	*	*	*	0.4	0.9	1.9
Male	21.6	*	*	*	*	*	*	0.4	1.0	2.2
Female	21.4	*	*	*	*	*	*	0.3	0.7	1.6
Black, both sexes	18.8	*	*	*	*	*	*	*	1.5	3.7
Male	19.1	*	*	*	*	*	*	*	1.7	4.0
Female	18.5	*	*	*	*	*	*	*	1.4	3.4

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Malignant melanoma of skin (C43)										
All races 3/, both sexes	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
Male	3.3	*	*	*	*	*	*	0.3	0.7	1.0
Female	2.0	*	*	*	*	*	*	0.3	0.5	1.0
White, both sexes	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
Male	4.0	*	*	*	*	*	*	0.4	0.9	1.2
Female	2.3	*	*	*	*	*	*	0.3	0.6	1.2
Black, both sexes	0.4	*	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*	*
Malignant neoplasm of breast (C50)										
All races 3/, both sexes	15.0	*	*	*	*	*	*	0.4	1.9	4.3
Male	0.3	*	*	*	*	*	*	*	*	*
Female	29.2	*	*	*	*	*	*	0.8	3.8	8.5
White, both sexes	15.7	*	*	*	*	*	*	0.4	1.6	3.9
Male	0.3	*	*	*	*	*	*	*	*	*
Female	30.7	*	*	*	*	*	*	0.8	3.3	7.7
Black, both sexes	14.8	*	*	*	*	*	*	0.7	3.7	7.7
Male	0.4	*	*	*	*	*	*	*	*	*
Female	27.9	*	*	*	*	*	*	*	7.1	14.3

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
Male	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
Female	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
White, both sexes	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
Male	5.7	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
Female	4.4	*	0.6	0.8	0.7	0.4	0.6	0.8	0.9	1.5
Black, both sexes	2.1	*	*	0.7	0.8	*	*	*	0.8	1.2
Male	2.4	*	*	*	*	*	*	*	*	*
Female	1.9	*	*	*	*	*	*	*	*	*

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
Male	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
Female	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
White, both sexes	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
Male	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
Female	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
Black, both sexes	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
Male	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
Female	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
All races 3/, both sexes	0.0	*	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4
Male	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7
Female	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0
White, both sexes	23.9	*	0.8	0.7	0.7	1.2	1.5	1.6	2.0	3.2
Male	24.4	*	1.0	0.7	0.7	1.3	1.9	2.1	2.5	3.6
Female	23.5	*	0.6	0.6	0.6	1.0	1.1	1.1	1.5	2.8
Black, both sexes	18.3	*	1.0	*	0.7	1.5	1.3	1.3	2.0	5.2
Male	19.4	*	*	*	*	1.7	2.1	1.6	2.0	5.4
Female	17.3	*	*	*	*	1.4	*	*	1.9	5.1

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All races 3/, both sexes	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7
Male	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8
Female	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6
White, both sexes	5.3	1.6	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7
Male	5.2	1.6	0.4	0.3	*	*	0.4	0.3	0.5	0.7
Female	5.4	1.6	*	0.3	*	*	*	0.3	0.4	0.6
Black, both sexes	3.1	*	*	*	*	*	*	*	*	0.8
Male	3.0	*	*	*	*	*	*	*	*	*
Female	3.2	*	*	*	*	*	*	*	*	*
Anemias (D50-D64)										
All races 3/, both sexes	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4
Male	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
Female	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4
White, both sexes	1.5	*	*	*	*	*	*	*	*	*
Male	1.1	*	*	*	*	*	*	*	*	*
Female	1.9	*	*	*	*	*	*	*	*	*
Black, both sexes	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
Male	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
Female	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Alzheimer's disease (G30)										
All races 3/, both sexes	17.6	*	*	*	*	*	*	*	*	*
Male	10.5	*	*	*	*	*	*	*	*	*
Female	24.5	*	*	*	*	*	*	*	*	*
White, both sexes	20.2	*	*	*	*	*	*	*	*	*
Male	11.9	*	*	*	*	*	*	*	*	*
Female	28.2	*	*	*	*	*	*	*	*	*
Black, both sexes	7.5	*	*	*	*	*	*	*	*	*
Male	4.5	*	*	*	*	*	*	*	*	*
Female	10.2	*	*	*	*	*	*	*	*	*
Major cardiovascular diseases (I00-I78)										
All races 3/, both sexes	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2
Male	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3
Female	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1
White, both sexes	354.7	14.6	1.3	0.6	0.9	2.1	3.5	5.5	10.1	20.6
Male	334.2	15.3	1.6	0.7	1.1	2.4	4.2	6.9	13.2	27.6
Female	374.8	13.9	1.1	0.5	0.7	1.7	2.8	4.1	6.9	13.5
Black, both sexes	285.0	27.8	2.5	1.0	1.9	4.6	9.1	15.1	27.7	51.2
Male	275.6	29.1	2.1	*	1.8	5.1	10.2	18.0	32.7	61.6
Female	293.4	26.5	2.9	*	2.0	4.2	8.1	12.4	23.3	42.0

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
Male	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
Female	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
White, both sexes	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
Male	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
Female	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
Black, both sexes	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
Male	141.5	*	*	*	*	*	*	3.7	8.0	20.3
Female	138.8	*	*	*	*	*	*	*	4.3	9.7
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
Male	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
Female	64.6	*	*	*	*	*	*	0.3	0.7	2.2
White, both sexes	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
Male	78.8	*	*	*	*	*	*	0.6	1.8	6.0
Female	69.6	*	*	*	*	*	*	0.3	0.6	2.0
Black, both sexes	52.2	*	*	*	*	*	*	1.2	2.3	4.9
Male	52.0	*	*	*	*	*	*	1.6	2.7	6.5
Female	52.5	*	*	*	*	*	*	*	1.9	3.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes	1.2	*	*	*	*	*	*	*	*	0.1
Male	1.3	*	*	*	*	*	*	*	*	0.2
Female	1.1	*	*	*	*	*	*	*	*	*
White, both sexes	1.2	*	*	*	*	*	*	*	*	0.1
Male	1.3	*	*	*	*	*	*	*	*	*
Female	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes	1.2	*	*	*	*	*	*	*	*	*
Male	1.5	*	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	*	*	*	*
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
Male	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
Female	111.9	*	*	*	*	*	*	0.2	1.0	2.8
White, both sexes	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
Male	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
Female	121.6	*	*	*	*	*	*	*	0.8	2.4
Black, both sexes	86.6	*	*	*	*	*	*	1.3	3.6	9.5
Male	88.0	*	*	*	*	*	*	2.1	5.0	13.4
Female	85.3	*	*	*	*	*	*	*	2.4	5.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes	24.7	*	*	*	*	*	*	0.2	0.5	1.7
Male	26.0	*	*	*	*	*	*	0.3	0.9	2.8
Female	23.5	*	*	*	*	*	*	*	0.2	0.7
White, both sexes	25.1	*	*	*	*	*	*	*	0.5	1.6
Male	26.0	*	*	*	*	*	*	*	0.8	2.5
Female	24.3	*	*	*	*	*	*	*	*	0.6
Black, both sexes	28.5	*	*	*	*	*	*	*	0.9	2.9
Male	32.3	*	*	*	*	*	*	*	*	4.7
Female	25.0	*	*	*	*	*	*	*	*	1.4

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
Male	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
Female	88.4	*	*	*	*	*	*	*	0.8	2.1
White, both sexes	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
Male	97.5	*	*	*	*	*	*	0.7	2.1	5.6
Female	97.3	*	*	*	*	*	*	*	0.6	1.8
Black, both sexes	58.1	*	*	*	*	*	*	0.8	2.8	6.5
Male	55.8	*	*	*	*	*	*	*	3.8	8.7
Female	60.3	*	*	*	*	*	*	*	1.9	4.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Emphysema (J43)										
All races 3/, both sexes	5.9	*	*	*	*	*	*	*	*	0.1
Male	6.3	*	*	*	*	*	*	*	*	*
Female	5.6	*	*	*	*	*	*	*	*	*
White, both sexes	6.8	*	*	*	*	*	*	*	*	*
Male	7.0	*	*	*	*	*	*	*	*	*
Female	6.5	*	*	*	*	*	*	*	*	*
Black, both sexes	2.5	*	*	*	*	*	*	*	*	*
Male	3.2	*	*	*	*	*	*	*	*	*
Female	1.8	*	*	*	*	*	*	*	*	*
Asthma (J45-J46)										
All races 3/, both sexes	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
Male	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
Female	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
White, both sexes	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
Male	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
Female	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
Black, both sexes	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
Male	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
Female	3.5	*	*	*	*	*	*	*	1.4	2.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Hernia (K40-K46)										
All races 3/, both sexes	0.5	1.2	*	*	*	*	*	*	*	*
Male	0.4	1.4	*	*	*	*	*	*	*	*
Female	0.7	*	*	*	*	*	*	*	*	*
White, both sexes	0.6	1.2	*	*	*	*	*	*	*	*
Male	0.5	1.5	*	*	*	*	*	*	*	*
Female	0.7	*	*	*	*	*	*	*	*	*
Black, both sexes	0.4	*	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*	*	*

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8
Male	12.5	*	*	*	*	*	*	0.6	2.1	6.3
Female	6.5	*	*	*	*	*	*	0.3	1.1	3.2
White, both sexes	10.0	*	*	*	*	*	*	0.4	1.5	4.7
Male	13.2	*	*	*	*	*	*	0.6	2.1	6.3
Female	6.8	*	*	*	*	*	*	*	1.0	3.1
Black, both sexes	7.5	*	*	*	*	*	*	0.8	1.3	4.9
Male	9.9	*	*	*	*	*	*	*	*	6.6
Female	5.2	*	*	*	*	*	*	*	1.4	3.3

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes	1.0	*	*	*	*	*	*	*	*	0.1
Male	0.9	*	*	*	*	*	*	*	*	*
Female	1.1	*	*	*	*	*	*	*	*	*
White, both sexes	1.1	*	*	*	*	*	*	*	*	*
Male	1.0	*	*	*	*	*	*	*	*	*
Female	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes	0.8	*	*	*	*	*	*	*	*	*
Male	0.6	*	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	*	*	*	*
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2
Male	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3
Female	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1
White, both sexes	12.9	2.9	*	*	*	0.1	0.3	0.3	0.5	0.8
Male	12.7	3.8	*	*	*	*	0.3	0.4	0.5	0.9
Female	13.0	2.0	*	*	*	*	0.3	0.3	0.5	0.7
Black, both sexes	18.9	11.5	*	*	*	*	*	1.1	2.6	4.4
Male	17.7	13.4	*	*	*	*	*	*	2.8	5.1
Female	20.0	9.6	*	*	*	*	*	*	2.3	3.8

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Inflammatory diseases of female pelvic organs (N70-N76)										
All races 3/, both sexes	0.0	*	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*	*
Pregnancy, childbirth and the puerperium (000-099)										
All races 3/, both sexes	0.1	*	*	0.4	0.4	0.5	0.4
Female	0.3	*	*	0.8	0.9	0.9	0.8
White, both sexes	0.1	*	*	0.3	0.3	0.3	0.3
Female	0.2	*	*	0.6	0.7	0.7	0.6
Black, both sexes	0.4	*	*	1.1	1.1	1.2	0.9
Female	0.7	*	*	2.2	2.2	2.3	1.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Pregnancy with abortive outcome (000-007)										
All races 3/, both sexes	0.0	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (010-099)										
All races 3/, both sexes	0.1	*	*	0.4	0.4	0.4	0.3
Female	0.3	*	*	0.8	0.8	0.9	0.7
White, both sexes	0.1	*	*	0.3	0.3	0.3	0.3
Female	0.2	*	*	0.6	0.6	0.7	0.5
Black, both sexes	0.3	*	*	1.0	1.0	1.1	0.7
Female	0.6	*	*	1.9	1.8	2.1	1.4

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Certain conditions originating in the perinatal period (P00-P96)										
All races 3/, both sexes	5.0	365.7	0.5	0.1	*	*	*	*	*	*
Male	5.8	403.9	0.7	*	*	*	*	*	*	*
Female	4.3	325.6	0.4	*	*	*	*	*	*	*
White, both sexes	3.7	283.7	0.4	*	*	*	*	*	*	*
Male	4.3	312.4	0.5	*	*	*	*	*	*	*
Female	3.2	253.5	*	*	*	*	*	*	*	*
Black, both sexes	13.7	809.5	1.1	*	*	*	*	*	*	*
Male	16.3	901.3	*	*	*	*	*	*	*	*
Female	11.3	714.6	*	*	*	*	*	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
Male	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
Female	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
White, both sexes	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
Male	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
Female	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
Black, both sexes	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
Male	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
Female	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All races 3/, both sexes	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
Male	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
Female	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
White, both sexes	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
Male	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
Female	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
Black, both sexes	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
Male	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
Female	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All races 3/, both sexes	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9
Male	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5
Female	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3
White, both sexes	67.0	24.9	3.5	1.7	2.3	3.5	5.1	5.7	8.0	12.3
Male	54.1	26.8	3.9	1.9	2.6	4.1	6.2	7.0	9.6	15.0
Female	79.6	22.9	3.2	1.6	2.0	2.8	3.9	4.4	6.4	9.5
Black, both sexes	58.1	49.6	5.9	2.3	3.3	4.8	8.3	12.8	17.1	26.9
Male	54.3	59.8	6.1	2.0	3.1	5.1	9.1	12.9	18.3	29.9
Female	61.6	39.0	5.7	2.6	3.5	4.6	7.6	12.6	16.1	24.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
Male	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
Female	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
White, both sexes	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
Male	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
Female	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
Black, both sexes	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
Male	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
Female	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
Male	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
Female	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
White, both sexes	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
Male	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
Female	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
Black, both sexes	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
Male	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
Female	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All races 3/, both sexes	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9
Male	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5
Female	0.2	*	*	*	*	0.3	*	*	0.2	0.2
White, both sexes	0.7	*	*	*	0.2	0.5	0.8	0.9	0.8	0.9
Male	1.3	*	*	*	*	0.6	1.4	1.5	1.4	1.6
Female	0.2	*	*	*	*	0.3	*	*	*	*
Black, both sexes	0.4	*	*	*	*	*	*	*	*	*
Male	0.7	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8
Male	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2
Female	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4
White, both sexes	18.9	15.1	6.9	2.2	2.2	6.4	10.2	10.5	12.2	16.0
Male	23.4	17.3	8.4	2.8	3.1	10.3	16.7	16.9	19.1	24.6
Female	14.6	12.8	5.2	1.6	1.2	2.2	3.3	3.8	5.1	7.3
Black, both sexes	17.6	38.0	10.6	5.2	4.0	6.4	7.7	9.6	12.5	18.0
Male	24.9	40.9	12.6	6.9	5.6	10.3	12.6	15.9	18.9	27.1
Female	11.0	35.1	8.5	3.5	2.3	2.3	3.0	3.8	6.7	9.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
Male	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
Female	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
White, both sexes	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
Male	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
Female	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
Black, both sexes	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
Male	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
Female	0.6	*	1.7	*	*	*	*	*	*	*

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
Male	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
Female	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
White, both sexes	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
Male	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
Female	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
Black, both sexes	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
Male	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
Female	2.0	*	4.1	1.6	*	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
Male	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
Female	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
White, both sexes	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
Male	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
Female	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
Black, both sexes	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
Male	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
Female	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0
Male	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1
Female	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8
White, both sexes	6.6	12.2	1.6	0.6	0.8	1.4	2.1	2.3	2.4	3.0
Male	7.4	14.0	2.0	0.7	1.2	2.2	3.7	3.8	4.1	5.2
Female	5.7	10.3	1.2	0.4	0.3	0.5	0.4	0.6	0.7	0.8
Black, both sexes	5.4	30.6	2.3	0.7	0.7	1.3	1.8	2.0	2.4	3.1
Male	7.3	32.0	2.5	*	1.2	1.8	3.2	3.5	3.8	5.5
Female	3.8	29.1	2.0	*	*	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Intentional self-harm (suicide) (X60-X84,Y87.0)										
All races 3/, both sexes	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
Male	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
Female	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
White, both sexes	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
Male	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
Female	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
Black, both sexes	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
Male	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
Female	1.7	*	*	1.4	3.0	3.0	2.2	2.2

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
Male	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
Female	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
White, both sexes	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
Male	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
Female	1.7	*	*	1.1	1.3	1.6	2.0	2.7
Black, both sexes	3.0	*	*	3.8	6.8	6.0	4.0	3.4
Male	5.5	*	*	6.9	12.6	11.3	7.3	6.4
Female	0.7	*	*	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)										
All races 3/, both sexes	4.5	*	0.9	3.6	5.3	5.8	6.0	7.0
Male	6.6	*	1.4	5.4	8.5	8.9	9.1	10.6
Female	2.5	*	0.4	1.7	1.9	2.6	2.7	3.4
White, both sexes	4.9	*	0.9	3.8	5.5	6.1	6.4	7.6
Male	7.1	*	1.3	5.8	8.8	9.4	9.7	11.4
Female	2.8	*	0.5	1.8	1.9	2.5	3.1	3.8
Black, both sexes	2.4	*	1.1	1.8	4.2	3.9	3.6	3.8
Male	3.9	*	1.8	2.6	6.8	6.2	6.4	6.6
Female	1.0	*	*	*	1.7	1.8	*	1.4

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
Male	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
Female	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
White, both sexes	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
Male	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
Female	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
Black, both sexes	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
Male	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
Female	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS
Assault (homicide) by discharge of firearms (X93-X95)										
All races 3/, both sexes	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
Male	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
Female	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
White, both sexes	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
Male	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
Female	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
Black, both sexes	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
Male	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
Female	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9
Male	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5
Female	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4
White, both sexes	1.5	6.5	1.5	0.4	0.4	1.3	2.3	2.0	2.0	2.1
Male	2.0	8.1	1.7	0.5	0.4	1.6	3.0	2.5	2.5	2.5
Female	1.1	4.8	1.3	0.3	0.5	1.0	1.5	1.4	1.4	1.7
Black, both sexes	5.9	21.6	5.5	0.8	0.8	4.5	8.8	7.6	7.6	8.6
Male	8.0	21.1	5.5	*	*	5.5	12.6	10.7	10.5	10.9
Female	4.0	22.2	5.5	*	*	3.4	5.2	4.7	4.9	6.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Measles (B05)										
All races 3/, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)										
All races 3/, both sexes	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
Male	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
Female	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
White, both sexes	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
Male	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
Female	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
Black, both sexes	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
Male	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
Female	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
All races 3/, both sexes	0.8	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
Male	1.0	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
Female	0.6	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
White, both sexes	0.7	1.3	1.6	2.3	3.2	4.6	6.8	9.8	12.7	21.0
Male	0.9	1.5	2.0	2.7	3.6	5.5	8.6	11.6	16.1	26.9
Female	0.6	1.0	1.1	1.9	2.8	3.8	5.4	8.6	10.8	18.7
Black, both sexes	1.7	2.6	3.5	4.0	5.9	6.1	9.9	11.6	16.1	24.1
Male	2.1	3.2	4.8	4.4	7.2	8.6	12.4	17.0	20.2	32.3
Female	*	2.1	2.4	3.6	4.9	4.2	8.2	8.3	14.0	21.0
Malignant neoplasms (C00-C97)										
All races 3/, both sexes	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
Male	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
Female	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
White, both sexes	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
Male	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
Female	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
Black, both sexes	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
Male	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
Female	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
Male	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
Female	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
White, both sexes	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
Male	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
Female	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
Black, both sexes	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
Male	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
Female	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
Malignant neoplasm of esophagus (C15)										
All races 3/, both sexes	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
Male	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
Female	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
White, both sexes	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
Male	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
Female	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
Black, both sexes	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
Male	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
Female	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
Male	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
Female	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
White, both sexes	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
Male	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
Female	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
Black, both sexes	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
Male	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
Female	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All races 3/, both sexes	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
Male	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
Female	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
White, both sexes	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
Male	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
Female	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
Black, both sexes	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
Male	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
Female	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
Male	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
Female	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
White, both sexes	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
Male	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
Female	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
Black, both sexes	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
Male	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
Female	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
Male	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
Female	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
White, both sexes	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
Male	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
Female	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
Black, both sexes	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
Male	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
Female	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of larynx (C32)										
All races 3/, both sexes	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
Male	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
Female	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
White, both sexes	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
Male	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
Female	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
Black, both sexes	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
Male	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
Female	*	*	*	*	*	3.8	5.5	7.4	*	*
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
Male	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
Female	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
White, both sexes	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
Male	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
Female	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
Black, both sexes	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
Male	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
Female	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant melanoma of skin (C43)										
All races 3/, both sexes	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
Male	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
Female	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
White, both sexes	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
Male	2.5	3.8	6.0	7.3	9.9	12.5	16.2	20.1	26.6	34.7
Female	1.6	2.5	2.8	3.4	4.6	5.7	6.9	8.2	9.7	14.5
Black, both sexes	*	*	*	*	*	*	*	*	*	6.6
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Malignant neoplasm of breast (C50)										
All races 3/, both sexes	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
Male	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
Female	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
White, both sexes	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
Male	*	*	0.3	0.6	0.7	1.1	1.7	2.2	2.3	3.9
Female	15.0	24.4	38.7	51.8	65.4	80.2	98.6	119.7	144.9	205.5
Black, both sexes	14.8	23.9	32.8	43.9	47.3	52.7	65.6	84.2	100.5	176.4
Male	*	*	*	*	*	*	*	*	*	*
Female	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of cervix uteri (C53)										
All races 3/, both sexes	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
Female	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
White, both sexes	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
Female	3.1	4.0	4.7	5.3	5.0	5.5	5.9	6.5	8.0	8.8
Black, both sexes	3.1	3.7	4.9	6.2	6.4	9.4	9.7	10.8	13.6	19.7
Female	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
All races 3/, both sexes	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
Female	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
White, both sexes	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
Female	0.8	1.9	3.5	6.7	10.3	13.6	17.8	22.2	28.0	34.3
Black, both sexes	*	1.0	2.4	5.3	12.0	19.6	26.4	27.7	35.0	33.2
Female	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of ovary (C56)										
All races 3/, both sexes	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
Female	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
White, both sexes	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
Female	3.0	6.4	11.1	17.0	24.5	33.7	40.3	54.5	60.0	59.0
Black, both sexes	1.5	2.2	4.8	6.4	10.9	16.8	19.6	26.1	27.4	34.4
Female	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
Malignant neoplasm of prostate (C61)										
All races 3/, both sexes	*	0.5	1.5	4.6	12.1	25.9	49.0	85.1	136.6	209.3
Male	*	1.1	3.1	9.6	25.4	56.0	111.3	207.4	368.2	723.1
White, both sexes	*	0.4	1.3	3.8	10.4	22.4	43.2	77.6	125.8	198.6
Male	*	0.8	2.6	7.9	21.6	48.0	97.0	188.0	337.4	687.5
Black, both sexes	*	1.5	3.6	12.9	29.5	64.7	120.7	190.1	292.8	356.6
Male	*	3.2	7.8	28.7	66.9	152.5	302.6	505.6	874.2	1,314.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
Male	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
Female	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
White, both sexes	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
Male	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
Female	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
Black, both sexes	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
Male	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
Female	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
Malignant neoplasm of bladder (C67)										
All races 3/, both sexes	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
Male	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
Female	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
White, both sexes	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
Male	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
Female	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
Black, both sexes	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
Male	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
Female	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
Male	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
Female	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
White, both sexes	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
Male	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
Female	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
Black, both sexes	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
Male	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
Female	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
Male	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
Female	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
White, both sexes	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
Male	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
Female	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
Black, both sexes	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
Male	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
Female	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Hodgkin's disease (C81)										
All races 3/, both sexes	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
Male	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
Female	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
White, both sexes	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
Male	0.4	0.5	0.6	0.5	1.1	1.4	1.9	2.6	2.8	3.9
Female	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.6	2.4	2.1
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
Male	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
Female	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
White, both sexes	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
Male	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
Female	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
Black, both sexes	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
Male	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
Female	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Leukemia (C91-C95)										
All races 3/, both sexes	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
Male	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
Female	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
White, both sexes	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
Male	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
Female	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
Black, both sexes	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
Male	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
Female	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All races 3/, both sexes	0.5	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
Male	0.6	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
Female	0.4	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
White, both sexes	0.4	1.0	2.2	3.8	7.8	13.4	19.1	26.5	35.2	35.2
Male	0.5	1.2	2.5	4.3	8.8	16.1	22.8	33.9	45.4	50.2
Female	0.2	0.8	2.0	3.4	6.8	11.1	16.2	21.3	29.2	29.1
Black, both sexes	1.1	3.1	6.3	11.7	16.5	27.3	37.9	53.3	57.9	71.6
Male	*	3.3	7.2	12.3	18.4	31.0	48.1	67.1	69.1	79.6
Female	*	2.8	5.5	11.2	15.0	24.6	31.1	45.0	52.2	68.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

40-44 YEARS 45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes	*	*	*	*	*	*	*	*	*	0.5
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	5.9	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
Male	6.4	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
Female	5.3	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
White, both sexes	5.4	9.9	17.2	29.7	47.1	71.4	99.9	135.6	179.5	243.7
Male	5.8	11.2	19.5	34.1	54.7	86.2	121.8	165.7	225.2	307.0
Female	5.0	8.6	14.8	25.6	40.1	58.4	82.4	114.5	152.3	218.0
Black, both sexes	10.0	16.8	28.8	41.3	64.8	85.6	117.7	155.7	172.0	210.8
Male	11.8	20.8	33.1	51.4	89.0	102.0	149.1	211.7	208.2	253.7
Female	8.3	13.4	25.1	33.1	45.8	73.5	96.8	121.9	153.7	194.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All races 3/, both sexes	0.9	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
Male	0.9	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
Female	1.0	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
White, both sexes	0.9	1.3	2.2	3.9	6.6	10.7	17.3	29.5	52.2	87.3
Male	0.9	1.5	2.6	4.3	7.3	13.4	21.0	37.6	68.0	112.6
Female	1.0	1.2	1.9	3.5	6.0	8.4	14.3	23.8	42.9	77.1
Black, both sexes	1.4	2.4	3.1	4.2	7.1	10.6	20.3	26.7	37.8	64.1
Male	*	2.2	3.6	4.4	10.1	15.6	23.7	34.5	38.8	76.1
Female	1.3	2.5	2.7	4.0	4.7	6.9	18.0	21.9	37.4	59.7
Anemias (D50-D64)										
All races 3/, both sexes	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
Male	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
Female	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
White, both sexes	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
Male	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
Female	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
Black, both sexes	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
Male	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
Female	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Diabetes mellitus (E10-E14)										
All races 3/, both sexes	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
Male	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
Female	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
White, both sexes	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
Male	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
Female	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
Black, both sexes	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
Male	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
Female	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
Nutritional deficiencies (E40-E64)										
All races 3/, both sexes	0.1	0.2	0.3	0.5	1.0	2.0	3.0	6.8	15.8	50.0
Male	*	0.3	0.2	0.7	1.0	2.0	3.4	6.8	14.4	43.7
Female	*	0.2	0.4	0.4	0.9	2.0	2.7	6.8	16.6	52.6
White, both sexes	*	0.2	0.2	0.5	0.8	1.9	2.7	6.4	15.3	48.8
Male	*	0.2	*	0.5	0.8	1.8	2.9	6.2	13.9	42.8
Female	*	*	0.3	0.4	0.9	2.0	2.5	6.5	16.2	51.3
Black, both sexes	*	*	*	*	2.1	3.2	6.5	12.1	22.6	66.9
Male	*	*	*	*	*	*	9.7	15.1	22.8	63.4
Female	*	*	*	*	*	*	4.4	10.3	22.5	68.3

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Meningitis (G00,G03)										
All races 3/, both sexes	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
Male	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
Female	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
White, both sexes	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
Male	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
Female	*	*	*	*	*	*	*	0.6	1.0	1.0
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Parkinson's disease (G20-G21)										
All races 3/, both sexes	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
Male	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
Female	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
White, both sexes	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
Male	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
Female	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
Black, both sexes	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
Male	*	*	*	*	*	*	13.7	37.8	65.8	86.5
Female	*	*	*	*	*	*	5.5	10.8	23.4	38.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Alzheimer's disease (G30)										
All races 3/, both sexes	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
Male	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
Female	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
White, both sexes	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
Male	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
Female	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
Black, both sexes	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
Male	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
Female	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
Major cardiovascular diseases (I00-I78)										
All races 3/, both sexes	49.6	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
Male	68.0	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
Female	31.5	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
White, both sexes	42.6	76.5	130.6	225.6	376.3	612.7	1,027.2	1,783.3	3,243.6	8,133.9
Male	60.7	114.5	190.5	323.5	523.7	822.7	1,366.4	2,273.2	3,963.6	8,585.0
Female	24.4	39.0	72.0	132.4	240.4	429.2	755.0	1,438.7	2,815.1	7,950.6
Black, both sexes	104.9	192.0	303.5	489.3	736.9	1,061.9	1,623.2	2,496.2	3,719.8	7,518.9
Male	131.7	247.2	407.0	670.8	955.1	1,328.0	2,040.7	2,999.5	4,397.8	7,687.4
Female	81.2	144.0	215.2	340.5	564.6	865.5	1,346.0	2,192.8	3,378.3	7,456.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All races 3/, both sexes	39.4	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
Male	57.0	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
Female	22.2	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
White, both sexes	34.9	63.7	109.9	190.0	309.5	492.6	801.0	1,347.0	2,393.0	6,020.8
Male	52.3	99.4	166.1	280.7	442.0	680.1	1,096.8	1,771.0	3,030.0	6,560.8
Female	17.5	28.3	54.9	103.7	187.4	328.6	563.7	1,048.7	2,014.1	5,801.4
Black, both sexes	77.5	142.2	231.2	380.7	571.0	813.9	1,230.4	1,836.2	2,673.7	5,389.5
Male	101.3	190.8	319.1	533.0	754.7	1,033.7	1,570.9	2,245.4	3,226.1	5,633.5
Female	56.5	100.1	156.3	255.9	426.1	651.7	1,004.3	1,589.5	2,395.4	5,298.7

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
Male	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
Female	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
White, both sexes	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
Male	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.1	8.4	12.7
Female	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.4	14.7	19.6
Black, both sexes	0.9	1.0	1.1	1.9	2.2	3.4	3.6	3.7	*	8.4
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	4.4	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Hypertensive heart disease (I11)										
All races 3/, both sexes	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
Male	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
Female	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
White, both sexes	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
Male	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
Female	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
Black, both sexes	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
Male	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
Female	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
Male	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
Female	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
White, both sexes	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
Male	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
Female	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
Black, both sexes	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
Male	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
Female	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
Male	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
Female	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
White, both sexes	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
Male	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
Female	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
Black, both sexes	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
Male	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
Female	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
Male	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
Female	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
White, both sexes	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
Male	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
Female	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
Black, both sexes	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
Male	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
Female	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
Male	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
Female	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
White, both sexes	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
Male	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
Female	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
Black, both sexes	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
Male	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
Female	*	*	*	*	3.3	*	*	8.0	10.6	22.3
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
Male	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
Female	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
White, both sexes	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
Male	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
Female	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
Black, both sexes	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
Male	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
Female	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
Male	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
Female	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
White, both sexes	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
Male	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
Female	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
Black, both sexes	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
Male	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
Female	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
Male	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
Female	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
White, both sexes	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
Male	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
Female	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
Black, both sexes	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
Male	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
Female	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
Male	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
Female	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
White, both sexes	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
Male	0.2	*	*	0.5	*	*	1.1	1.2	1.9	2.1
Female	*	*	*	*	*	0.6	0.7	0.9	1.1	1.6
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Heart failure (I50)										
All races 3/, both sexes	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
Male	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
Female	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
White, both sexes	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
Male	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
Female	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
Black, both sexes	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
Male	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
Female	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes	10.9	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
Male	13.6	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
Female	8.2	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
White, both sexes	9.1	13.0	18.9	29.6	46.2	73.1	113.5	188.4	339.9	833.4
Male	11.5	17.5	24.8	38.9	59.5	90.9	142.2	227.3	393.2	884.2
Female	6.7	8.6	13.2	20.7	33.9	57.6	90.4	161.1	308.1	812.7
Black, both sexes	25.2	37.3	52.5	69.7	96.5	137.5	191.0	282.9	377.8	742.0
Male	30.9	46.3	71.0	95.0	122.0	174.3	233.4	340.7	449.3	816.5
Female	20.3	29.5	36.8	49.0	76.5	110.4	162.9	248.1	341.8	714.3

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes	1.1	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
Male	1.4	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
Female	0.8	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
White, both sexes	0.6	1.0	1.8	3.0	5.5	8.5	15.3	28.9	57.4	153.2
Male	0.9	1.4	2.5	3.8	6.6	8.8	16.8	29.9	57.5	132.5
Female	0.4	0.6	1.1	2.2	4.5	8.3	14.1	28.2	57.4	161.6
Black, both sexes	4.2	7.2	12.7	17.9	29.0	40.3	58.8	92.4	130.4	283.4
Male	5.3	9.0	15.6	22.6	34.4	40.1	63.9	93.1	137.4	269.9
Female	3.2	5.7	10.3	14.2	24.7	40.5	55.5	92.0	126.9	288.5

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
Male	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
Female	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
White, both sexes	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
Male	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
Female	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
Black, both sexes	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
Male	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
Female	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
Atherosclerosis (I70)										
All races 3/, both sexes	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
Male	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
Female	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
White, both sexes	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
Male	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
Female	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
Black, both sexes	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
Male	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
Female	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Other diseases of circulatory system (I71-I78)										
All races 3/, both sexes	1.3	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
Male	1.7	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
Female	0.9	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
White, both sexes	1.0	1.8	3.0	6.0	12.3	24.5	41.4	63.0	93.0	163.0
Male	1.3	2.5	4.4	8.9	17.8	35.5	59.1	89.4	129.7	219.6
Female	0.7	1.1	1.8	3.3	7.2	15.0	27.1	44.5	71.2	140.1
Black, both sexes	3.2	4.5	6.8	11.2	18.3	27.3	39.9	68.9	92.3	177.7
Male	4.2	5.5	10.0	15.1	22.6	32.8	51.8	90.7	104.5	177.6
Female	2.3	3.7	4.1	8.0	14.9	23.2	32.0	55.8	86.2	177.7
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
Male	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
Female	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
White, both sexes	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
Male	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
Female	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
Black, both sexes	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
Male	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
Female	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
Male	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
Female	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
White, both sexes	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
Male	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
Female	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
Black, both sexes	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
Male	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
Female	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
Male	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
Female	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
White, both sexes	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
Male	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
Female	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
Black, both sexes	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
Male	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
Female	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Emphysema (J43)										
All races 3/, both sexes	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
Male	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
Female	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
White, both sexes	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
Male	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
Female	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
Black, both sexes	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
Male	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
Female	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
Asthma (J45-J46)										
All races 3/, both sexes	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
Male	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
Female	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
White, both sexes	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
Male	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
Female	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
Black, both sexes	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
Male	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
Female	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
Male	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
Female	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
White, both sexes	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
Male	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
Female	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
Black, both sexes	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
Male	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
Female	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
All races 3/, both sexes	1.7	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
Male	1.8	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
Female	1.6	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
White, both sexes	1.6	2.2	4.3	7.5	12.4	22.8	37.7	62.0	91.2	145.9
Male	1.7	2.4	4.6	8.5	14.4	26.8	46.3	78.9	115.0	184.4
Female	1.5	2.1	3.9	6.6	10.6	19.3	30.7	50.1	77.1	130.3
Black, both sexes	3.0	6.0	7.8	11.6	15.5	28.1	33.4	49.8	71.2	98.2
Male	3.3	7.2	8.6	13.3	17.6	32.5	40.5	67.6	82.6	132.6
Female	2.6	5.0	7.1	10.3	13.9	24.8	28.6	39.0	65.4	85.4

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Hernia (K40-K46)										
All races 3/, both sexes	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
Male	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
Female	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
White, both sexes	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
Male	*	*	0.3	*	0.6	0.8	1.5	3.1	5.7	12.7
Female	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
Black, both sexes	*	*	*	*	*	*	*	3.9	*	12.5
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	12.4

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes	10.2	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
Male	14.4	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
Female	6.1	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
White, both sexes	10.1	16.9	18.1	21.6	26.5	29.4	31.8	33.4	30.5	23.9
Male	14.4	26.0	27.8	31.5	38.3	40.2	41.7	42.3	35.9	35.5
Female	5.9	7.8	8.7	12.2	15.6	20.0	23.9	27.2	27.3	19.2
Black, both sexes	10.9	19.2	25.9	25.9	25.9	27.5	24.7	18.3	17.5	13.8
Male	14.6	26.4	40.7	38.5	37.3	38.0	37.4	27.4	19.4	*
Female	7.7	13.0	13.3	15.5	17.0	19.8	16.2	12.8	16.6	11.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Alcoholic liver disease (K70)										
All races 3/, both sexes	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
Male	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
Female	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
White, both sexes	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
Male	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
Female	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
Black, both sexes	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
Male	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
Female	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes	3.6	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
Male	5.0	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
Female	2.2	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
White, both sexes	3.6	6.9	7.5	10.1	13.5	17.8	22.5	26.1	26.2	21.6
Male	5.0	10.3	11.0	13.5	17.6	21.2	25.8	28.2	28.1	30.1
Female	2.1	3.4	4.2	6.9	9.6	14.8	19.9	24.6	25.1	18.2
Black, both sexes	4.1	8.2	12.6	12.0	13.3	15.7	16.3	12.1	13.3	12.2
Male	5.0	11.4	20.1	18.1	19.3	18.7	20.7	16.1	*	*
Female	3.2	5.5	6.1	7.0	8.7	13.4	13.3	9.7	14.0	11.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
Male	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
Female	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
White, both sexes	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
Male	*	*	0.3	0.7	1.1	2.3	3.4	7.4	13.2	28.9
Female	*	*	0.3	0.4	0.7	1.4	2.2	4.4	8.3	24.3
Black, both sexes	*	*	*	*	*	3.1	4.9	6.6	11.0	17.8
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	4.9	6.3	11.5	19.3
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes	2.1	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
Male	2.5	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
Female	1.6	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
White, both sexes	1.5	2.2	3.8	6.8	13.2	23.8	40.6	73.0	125.3	268.7
Male	1.8	2.9	4.4	7.9	14.9	27.8	48.7	93.4	166.7	376.2
Female	1.1	1.7	3.2	5.8	11.6	20.3	34.1	58.6	100.7	225.0
Black, both sexes	6.3	12.7	19.7	32.3	50.7	83.5	119.7	165.4	242.3	406.4
Male	7.9	17.4	24.1	37.4	54.2	86.9	127.7	198.9	290.0	490.1
Female	4.8	8.6	16.1	28.2	47.9	81.0	114.3	145.3	218.2	375.2

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX40-44 YEARS 45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS
AND OVER

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
Male	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
Female	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
White, both sexes	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
Male	1.4	1.6	2.3	2.0	2.8	2.6	2.6	3.4	5.6	11.1
Female	1.1	1.3	1.7	2.4	2.4	2.7	2.3	3.4	4.9	6.5
Black, both sexes	1.6	1.7	1.5	2.4	2.5	2.8	*	*	*	*
Male	2.0	2.0	*	*	*	*	*	*	*	*
Female	*	*	*	2.8	3.3	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

40-44 YEARS 45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
Male	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
Female	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
White, both sexes	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
Male	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
Female	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
Black, both sexes	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
Male	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
Female	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes	22.1	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
Male	26.3	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
Female	17.9	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
White, both sexes	19.8	27.4	37.1	49.3	70.1	102.5	164.5	285.7	537.2	1,464.6
Male	24.0	35.0	44.1	56.8	77.5	112.7	177.5	298.2	533.4	1,274.8
Female	15.5	19.8	30.3	42.2	63.2	93.6	154.2	277.0	539.4	1,541.7
Black, both sexes	41.9	64.3	75.6	95.5	122.2	157.2	233.9	372.5	618.9	1,388.9
Male	48.0	74.7	89.1	114.4	150.1	188.4	279.8	428.6	692.9	1,325.1
Female	36.5	55.2	64.0	79.9	100.2	134.2	203.3	338.6	581.7	1,412.7

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
Male	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
Female	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
White, both sexes	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
Male	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
Female	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
Black, both sexes	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
Male	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
Female	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
Male	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
Female	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
White, both sexes	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
Male	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
Female	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
Black, both sexes	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
Male	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
Female	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All races 3/, both sexes	1.0	0.9	1.0	1.0	1.1	1.2	0.7	0.6	0.7	*
Male	1.7	1.6	1.8	1.8	1.9	2.0	1.4	1.2	1.4	*
Female	0.3	0.3	0.3	*	0.4	0.4	*	*	*	*
White, both sexes	1.1	1.0	1.1	1.1	1.2	1.2	0.8	0.6	0.7	*
Male	1.9	1.6	2.0	1.9	2.0	2.0	1.5	1.2	1.5	*
Female	0.4	0.3	0.3	*	0.4	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes	18.6	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
Male	28.3	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
Female	9.0	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
White, both sexes	18.3	17.2	14.1	13.3	15.3	17.9	28.9	50.4	96.5	250.6
Male	27.9	26.4	21.5	19.0	21.6	24.0	37.5	64.4	122.0	308.2
Female	8.7	8.1	6.9	7.9	9.5	12.6	21.9	40.6	81.4	227.2
Black, both sexes	24.5	27.6	27.7	23.9	22.9	27.8	38.8	56.0	79.1	159.5
Male	37.8	45.3	46.5	40.5	37.9	42.9	52.8	85.1	116.3	230.6
Female	12.7	12.3	11.7	10.3	11.1	16.7	29.5	38.4	60.3	133.1

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
Male	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
Female	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
White, both sexes	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
Male	1.6	1.5	1.5	1.6	1.6	1.4	1.7	1.1	3.0	2.7
Female	0.3	0.3	0.3	*	*	*	0.6	0.7	1.0	1.0
Black, both sexes	1.5	1.0	1.6	*	*	*	*	*	*	*
Male	2.8	1.9	2.8	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
Male	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
Female	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
White, both sexes	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
Male	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
Female	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
Black, both sexes	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
Male	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
Female	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
Male	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
Female	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
White, both sexes	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
Male	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
Female	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
Black, both sexes	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
Male	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
Female	8.6	8.0	6.6	3.0	*	*	*	*	*	*

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes	3.4	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
Male	5.6	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
Female	1.3	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
White, both sexes	3.4	3.7	4.0	4.7	6.2	7.4	11.8	21.7	41.9	120.5
Male	5.6	6.1	6.4	7.3	8.7	10.1	15.1	27.1	51.2	137.6
Female	1.3	1.4	1.6	2.3	3.8	5.1	9.1	17.9	36.4	113.6
Black, both sexes	3.9	5.7	5.6	7.9	8.6	12.3	18.1	30.2	39.2	90.1
Male	6.5	10.1	10.0	13.8	13.9	18.2	23.1	46.8	64.1	124.5
Female	1.5	1.8	*	3.0	4.4	7.9	14.9	20.2	26.7	77.3

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)										
All races 3/, both sexes	7.9	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
Male	11.3	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
Female	4.6	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
White, both sexes	8.7	8.2	6.6	5.7	4.1	3.1	3.6	4.2	5.3	6.7
Male	12.2	11.4	8.5	7.6	5.2	4.0	4.9	6.7	9.6	15.2
Female	5.1	5.0	4.8	4.0	3.1	2.3	2.6	2.5	2.7	3.3
Black, both sexes	4.7	3.3	3.1	1.7	*	*	*	*	*	*
Male	7.3	5.7	4.9	*	*	*	*	*	*	*
Female	2.3	*	*	*	*	*	*	*	*	*

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
Male	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
Female	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
White, both sexes	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
Male	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
Female	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
Black, both sexes	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
Male	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
Female	10.9	6.5	3.8	3.5	*	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Assault (homicide) by discharge of firearms (X93-X95)										
All races 3/, both sexes	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
Male	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
Female	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
White, both sexes	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
Male	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
Female	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
Black, both sexes	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
Male	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
Female	4.7	3.2	*	*	*	*	*	*	*	*

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	2.9	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
Male	3.9	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
Female	1.8	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
White, both sexes	1.9	1.7	1.4	1.2	1.0	1.0	1.1	1.4	1.6	1.5
Male	2.7	2.1	1.8	1.9	1.5	1.3	1.6	1.7	2.2	2.1
Female	1.2	1.2	0.9	0.6	0.6	0.7	0.7	1.1	1.2	1.3
Black, both sexes	9.2	7.4	5.3	5.0	4.7	5.4	4.1	4.1	6.8	*
Male	12.6	12.0	9.3	8.9	7.7	9.9	6.7	*	*	*
Female	6.2	3.3	2.0	*	*	*	*	*	*	*

TABLE 210. DEATH RATES FOR 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	40-44 YEARS	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Operations of war and their sequelae (Y36,Y89.1)										
All races 3/, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
Male	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
Female	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
White, both sexes	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
Male	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
Female	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
Black, both sexes	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
Male	*	*	*	*	4.6	6.2	9.4	10.4	*	*
Female	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.