

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Measles (B05)									
All races 3/, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)									
All races 3/, both sexes	1.9	*	*	*	*	0.2	1.7	5.0	3.7
Male	2.4	*	*	*	*	0.2	2.4	7.7	5.0
Female	1.4	*	*	*	*	0.2	1.1	2.4	2.6
White, both sexes	1.8	*	*	*	*	0.2	1.7	4.6	3.2
Male	2.4	*	*	*	*	0.2	2.4	7.2	4.3
Female	1.3	*	*	*	*	0.1	1.0	2.1	2.2
Black, both sexes	2.2	*	*	*	*	0.4	2.0	7.9	5.9
Male	2.9	*	*	*	*	*	2.9	12.3	8.1
Female	1.5	*	*	*	*	*	1.3	4.2	4.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)									
All races 3/, both sexes	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9
Male	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3
Female	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6
White, both sexes	2.0	3.4	0.4	0.2	0.2	0.3	0.7	1.4	2.7
Male	2.1	3.9	0.4	0.2	0.3	0.3	0.8	1.7	3.1
Female	1.9	2.9	0.4	0.2	0.2	0.2	0.5	1.1	2.3
Black, both sexes	2.3	9.9	*	*	0.4	0.9	1.6	3.0	4.9
Male	2.5	9.9	*	*	*	0.9	2.1	3.9	5.7
Female	2.1	9.9	*	*	*	1.0	1.2	2.2	4.2
Malignant neoplasms (C00-C97)									
All races 3/, both sexes	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7
Male	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8
Female	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3
White, both sexes	208.6	2.5	2.8	2.6	4.3	9.7	34.5	121.8	359.4
Male	218.1	2.8	3.1	2.8	5.0	9.2	30.9	123.5	401.9
Female	199.4	2.3	2.4	2.3	3.6	10.1	38.2	120.1	319.7
Black, both sexes	169.3	*	2.8	2.2	5.1	11.9	54.0	192.6	496.0
Male	188.5	*	2.7	2.5	6.1	10.1	48.4	214.2	626.4
Female	151.8	*	2.9	1.9	4.1	13.5	58.9	173.9	391.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
All races 3/, both sexes	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8
Male	3.6	*	*	*	*	0.1	1.0	4.5	10.9
Female	1.8	*	*	*	*	0.1	0.3	1.3	2.9
White, both sexes	2.7	*	*	*	*	0.1	0.5	2.4	6.0
Male	3.5	*	*	*	*	0.1	0.7	3.7	9.6
Female	1.9	*	*	*	*	*	0.3	1.2	2.7
Black, both sexes	3.0	*	*	*	*	*	1.1	6.0	13.5
Male	4.8	*	*	*	*	*	1.7	10.5	24.3
Female	1.4	*	*	*	*	*	*	2.0	4.8
Malignant neoplasm of esophagus (C15)									
All races 3/, both sexes	4.3	*	*	*	*	0.1	0.7	3.6	10.8
Male	6.7	*	*	*	*	0.2	1.2	6.3	18.6
Female	2.1	*	*	*	*	*	0.1	1.0	3.5
White, both sexes	4.5	*	*	*	*	0.1	0.7	3.3	10.0
Male	7.0	*	*	*	*	0.2	1.2	5.9	17.5
Female	2.1	*	*	*	*	*	0.1	0.8	2.9
Black, both sexes	4.5	*	*	*	*	*	0.8	6.6	19.8
Male	6.7	*	*	*	*	*	1.2	11.2	34.0
Female	2.5	*	*	*	*	*	*	2.7	8.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasm of stomach (C16)									
All races 3/, both sexes	4.5	*	*	*	*	0.3	1.1	2.9	7.4
Male	5.3	*	*	*	*	0.3	1.3	3.8	10.5
Female	3.7	*	*	*	*	0.3	0.9	2.1	4.6
White, both sexes	4.2	*	*	*	*	0.2	0.9	2.5	6.5
Male	5.0	*	*	*	*	0.3	1.0	3.2	9.3
Female	3.5	*	*	*	*	0.2	0.7	1.7	3.9
Black, both sexes	5.9	*	*	*	*	0.7	2.0	5.7	13.8
Male	7.0	*	*	*	*	0.8	2.9	7.9	20.6
Female	5.0	*	*	*	*	*	1.3	3.7	8.4
Malignant neoplasms of colon, rectum and anus (C18-C21)									
All races 3/, both sexes	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7
Male	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6
Female	20.2	*	*	*	*	0.6	2.7	9.8	26.4
White, both sexes	21.5	*	*	*	0.1	0.6	2.7	10.4	31.2
Male	21.6	*	*	*	*	0.7	3.1	12.0	37.6
Female	21.4	*	*	*	*	0.5	2.4	9.0	25.2
Black, both sexes	18.8	*	*	*	*	1.0	5.1	19.5	50.1
Male	19.1	*	*	*	*	1.0	6.0	22.0	62.5
Female	18.5	*	*	*	*	1.0	4.3	17.2	40.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasms of liver and intrahepatic bile ducts (C22)									
All races 3/, both sexes	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3
Male	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8
Female	3.3	*	*	*	*	0.1	0.6	1.9	5.1
White, both sexes	4.5	*	*	*	0.1	0.1	0.7	3.7	8.1
Male	5.7	*	*	*	*	0.2	1.0	5.7	11.8
Female	3.3	*	*	*	*	*	0.4	1.7	4.6
Black, both sexes	4.4	*	*	*	*	0.4	1.8	8.6	14.1
Male	6.1	*	*	*	*	*	2.4	14.8	23.5
Female	2.8	*	*	*	*	*	1.2	3.3	6.5
Malignant neoplasm of pancreas (C25)									
All races 3/, both sexes	10.4	*	*	*	*	0.1	1.2	6.3	19.4
Male	10.3	*	*	*	*	0.2	1.4	8.0	23.6
Female	10.5	*	*	*	*	0.1	1.0	4.6	15.5
White, both sexes	11.0	*	*	*	*	0.1	1.2	6.1	18.9
Male	10.9	*	*	*	*	0.2	1.4	7.8	23.0
Female	11.1	*	*	*	*	*	0.9	4.4	15.0
Black, both sexes	9.3	*	*	*	*	*	1.7	9.0	27.5
Male	9.0	*	*	*	*	*	2.0	11.8	34.2
Female	9.7	*	*	*	*	*	1.4	6.6	22.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasm of larynx (C32)									
All races 3/, both sexes	1.4	*	*	*	*	*	0.1	1.1	3.6
Male	2.2	*	*	*	*	*	0.2	1.8	6.2
Female	0.6	*	*	*	*	*	*	0.4	1.2
White, both sexes	1.3	*	*	*	*	*	0.1	0.9	3.1
Male	2.1	*	*	*	*	*	0.2	1.5	5.3
Female	0.6	*	*	*	*	*	*	0.4	1.1
Black, both sexes	2.0	*	*	*	*	*	*	2.9	9.0
Male	3.3	*	*	*	*	*	*	5.1	17.2
Female	0.7	*	*	*	*	*	*	1.0	2.4
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
All races 3/, both sexes	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4
Male	65.5	*	*	*	*	0.5	6.9	38.5	154.0
Female	45.4	*	*	*	*	0.5	5.3	25.0	93.3
White, both sexes	59.5	*	*	*	*	0.5	5.7	30.2	122.5
Male	69.4	*	*	*	*	0.5	6.1	35.7	150.8
Female	49.9	*	*	*	*	0.5	5.2	24.8	96.1
Black, both sexes	43.9	*	*	*	*	0.6	10.1	50.4	152.5
Male	58.3	*	*	*	*	*	13.4	70.7	223.5
Female	30.8	*	*	*	*	*	7.3	32.9	95.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant melanoma of skin (C43)									
All races 3/, both sexes	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4
Male	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4
Female	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5
White, both sexes	3.1	*	*	*	0.1	0.6	1.6	3.7	6.1
Male	4.0	*	*	*	0.1	0.7	1.9	4.8	8.5
Female	2.3	*	*	*	*	0.5	1.4	2.6	3.9
Black, both sexes	0.4	*	*	*	*	*	*	*	0.8
Male	0.3	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*
Malignant neoplasm of breast (C50)									
All races 3/, both sexes	15.0	*	*	*	*	1.2	6.3	16.9	31.2
Male	0.3	*	*	*	*	*	*	0.2	0.7
Female	29.2	*	*	*	*	2.3	12.4	33.0	59.3
White, both sexes	15.7	*	*	*	*	1.0	5.7	15.8	30.2
Male	0.3	*	*	*	*	*	*	0.2	0.7
Female	30.7	*	*	*	*	2.1	11.3	31.2	57.9
Black, both sexes	14.8	*	*	*	*	2.3	11.2	27.8	45.4
Male	0.4	*	*	*	*	*	*	*	*
Female	27.9	*	*	*	*	4.2	20.9	51.5	80.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX:

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasm of cervix uteri (C53)									
All races 3/, both sexes	1.5	*	*	*	*	0.6	1.5	2.4	3.1
Female	2.9	*	*	*	*	1.2	3.1	4.8	5.9
White, both sexes	1.4	*	*	*	*	0.5	1.4	2.2	2.7
Female	2.7	*	*	*	*	1.1	2.8	4.3	5.2
Black, both sexes	2.4	*	*	*	*	0.9	2.6	4.3	6.3
Female	4.5	*	*	*	*	1.8	4.9	7.9	11.3
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)									
All races 3/, both sexes	2.3	*	*	*	*	0.1	0.3	1.3	4.6
Female	4.6	*	*	*	*	0.1	0.6	2.6	8.9
White, both sexes	2.3	*	*	*	*	0.1	0.3	1.3	4.3
Female	4.6	*	*	*	*	0.1	0.6	2.6	8.3
Black, both sexes	2.9	*	*	*	*	*	0.4	1.6	8.3
Female	5.5	*	*	*	*	*	0.7	3.0	14.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasm of ovary (C56)									
All races 3/, both sexes	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1
Female	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4
White, both sexes	5.5	*	*	*	*	0.3	1.1	4.4	10.5
Female	10.9	*	*	*	*	0.5	2.2	8.7	20.4
Black, both sexes	2.9	*	*	*	*	*	1.0	3.3	8.4
Female	5.6	*	*	*	*	*	2.0	6.2	15.2
Malignant neoplasm of prostate (C61)									
All races 3/, both sexes	11.0	*	*	*	*	*	0.1	1.0	7.9
Male	22.5	*	*	*	*	*	0.1	2.0	16.6
White, both sexes	11.0	*	*	*	*	*	*	0.8	6.8
Male	22.3	*	*	*	*	*	*	1.7	13.9
Black, both sexes	14.6	*	*	*	*	*	*	2.4	20.3
Male	30.7	*	*	*	*	*	*	5.2	45.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)									
All races 3/, both sexes	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6
Male	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8
Female	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6
White, both sexes	5.1	*	0.7	0.8	0.6	1.1	2.6	5.6	10.4
Male	5.7	*	0.7	0.9	0.7	1.4	3.4	6.8	12.8
Female	4.4	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2
Black, both sexes	2.1	*	*	0.8	0.6	0.7	1.6	2.9	5.7
Male	2.4	*	*	0.9	0.7	1.1	2.0	3.7	7.1
Female	1.9	*	*	0.6	*	*	1.2	2.2	4.6
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
All races 3/, both sexes	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0
Male	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1
Female	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4
White, both sexes	21.9	1.2	1.0	1.0	1.8	2.2	4.0	10.4	30.1
Male	23.6	*	1.0	1.1	2.2	2.6	4.8	12.6	36.1
Female	20.3	*	1.0	0.8	1.3	1.8	3.2	8.3	24.4
Black, both sexes	13.9	*	0.8	0.6	2.2	2.9	5.3	14.7	35.8
Male	14.7	*	*	0.8	2.7	3.5	5.7	17.9	44.5
Female	13.1	*	*	*	1.8	2.4	5.0	12.0	28.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Hodgkin's disease (C81)									
All races 3/, both sexes	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7
Male	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7
Female	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6
White, both sexes	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7
Male	0.5	*	*	*	0.2	0.4	0.4	0.6	0.7
Female	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6
Black, both sexes	0.3	*	*	*	*	*	0.4	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	0.3	*	*	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)									
All races 3/, both sexes	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1
Male	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9
Female	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5
White, both sexes	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.7
Male	9.6	*	*	0.1	0.6	0.8	2.2	6.0	16.5
Female	8.6	*	*	*	0.2	0.6	1.1	3.4	11.1
Black, both sexes	3.7	*	*	*	0.6	1.0	2.1	5.0	10.5
Male	4.2	*	*	*	*	1.2	2.6	6.9	14.5
Female	3.3	*	*	*	*	0.9	1.6	3.3	7.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Leukemia (C91-C95)									
All races 3/, both sexes	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9
Male	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5
Female	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6
White, both sexes	8.3	1.2	1.0	0.8	1.2	1.1	1.8	3.7	10.0
Male	9.4	*	0.9	0.9	1.4	1.3	1.9	4.2	12.6
Female	7.4	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7
Black, both sexes	4.8	*	*	0.5	1.4	1.6	2.0	4.8	10.7
Male	5.3	*	*	0.6	1.9	1.9	2.0	5.5	13.8
Female	4.2	*	*	*	0.9	1.3	2.1	4.3	8.1
Multiple myeloma and immunoproliferative neoplasms (C88,C90)									
All races 3/, both sexes	3.9	*	*	*	*	*	0.3	1.9	6.3
Male	4.0	*	*	*	*	*	0.4	2.1	6.9
Female	3.8	*	*	*	*	*	0.3	1.6	5.7
White, both sexes	3.9	*	*	*	*	*	0.3	1.6	5.6
Male	4.1	*	*	*	*	*	0.3	1.8	6.3
Female	3.8	*	*	*	*	*	0.2	1.4	5.0
Black, both sexes	5.0	*	*	*	*	*	0.8	4.5	13.9
Male	4.8	*	*	*	*	*	*	5.0	15.0
Female	5.2	*	*	*	*	*	0.9	4.0	12.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)									
All races 3/, both sexes	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3
Male	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9
Female	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2
White, both sexes	23.9	*	0.8	0.7	1.3	1.8	4.3	13.3	37.5
Male	24.4	*	1.0	0.7	1.6	2.3	4.7	15.1	43.2
Female	23.5	*	0.6	0.6	1.0	1.3	3.9	11.6	32.1
Black, both sexes	18.3	*	1.0	0.6	1.4	1.6	7.5	22.1	51.9
Male	19.4	*	*	0.7	1.9	1.8	8.5	26.2	68.1
Female	17.3	*	*	*	1.0	1.5	6.6	18.6	38.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)									
All races 3/, both sexes	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0
Male	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6
Female	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5
White, both sexes	5.3	1.6	0.4	0.2	0.2	0.4	0.8	1.8	5.1
Male	5.2	1.6	0.4	0.2	0.3	0.4	0.8	2.0	5.6
Female	5.4	1.6	*	0.2	0.2	0.4	0.8	1.5	4.6
Black, both sexes	3.1	*	*	*	*	0.6	1.1	2.7	5.5
Male	3.0	*	*	*	*	*	1.2	2.8	7.0
Female	3.2	*	*	*	*	0.8	1.0	2.6	4.4
Anemias (D50-D64)									
All races 3/, both sexes	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1
Male	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1
Female	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0
White, both sexes	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8
Male	1.1	*	*	*	*	*	0.1	0.3	0.9
Female	1.9	*	*	*	*	*	0.1	0.3	0.8
Black, both sexes	2.6	*	*	0.5	1.1	2.3	2.4	3.0	3.4
Male	2.3	*	*	*	1.0	2.1	2.0	3.3	3.4
Female	3.0	*	*	*	1.1	2.4	2.8	2.8	3.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Alzheimer's disease (G30)									
All races 3/, both sexes	17.6	*	*	*	*	*	*	0.2	2.0
Male	10.5	*	*	*	*	*	*	0.2	1.8
Female	24.5	*	*	*	*	*	*	0.2	2.2
White, both sexes	20.2	*	*	*	*	*	*	0.2	2.2
Male	11.9	*	*	*	*	*	*	0.2	1.9
Female	28.2	*	*	*	*	*	*	0.2	2.4
Black, both sexes	7.5	*	*	*	*	*	*	*	1.2
Male	4.5	*	*	*	*	*	*	*	*
Female	10.2	*	*	*	*	*	*	*	*
Major cardiovascular diseases (I00-I78)									
All races 3/, both sexes	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2
Male	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6
Female	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3
White, both sexes	354.7	14.6	1.3	0.8	2.8	7.9	31.6	102.0	292.7
Male	334.2	15.3	1.6	0.9	3.3	10.1	44.1	150.2	411.8
Female	374.8	13.9	1.1	0.6	2.2	5.5	19.0	54.6	180.9
Black, both sexes	285.0	27.8	2.5	1.4	6.8	21.5	77.4	241.3	600.3
Male	275.6	29.1	2.1	1.4	7.5	25.4	95.8	317.5	796.8
Female	293.4	26.5	2.9	1.4	6.1	17.9	61.1	175.6	441.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)									
All races 3/, both sexes	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2
Male	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7
Female	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3
White, both sexes	270.2	11.5	1.1	0.6	2.1	6.1	25.8	85.4	243.2
Male	265.8	11.9	1.3	0.7	2.6	8.1	37.8	130.7	351.8
Female	274.5	11.1	0.8	0.4	1.6	4.0	13.7	40.9	141.3
Black, both sexes	211.8	19.8	2.0	1.1	5.6	17.2	57.4	181.5	466.0
Male	211.1	19.8	1.9	1.2	6.6	21.2	74.3	247.2	631.2
Female	212.6	19.8	2.0	1.0	4.7	13.6	42.4	125.0	332.8

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes	1.3	*	*	*	*	0.1	0.3	0.7	1.7
Male	0.8	*	*	*	*	*	0.3	0.6	1.4
Female	1.8	*	*	*	*	0.1	0.4	0.8	2.0
White, both sexes	1.4	*	*	*	*	0.1	0.3	0.6	1.7
Male	0.9	*	*	*	*	*	0.2	0.6	1.5
Female	1.9	*	*	*	*	*	0.3	0.7	1.9
Black, both sexes	0.7	*	*	*	*	*	0.6	1.0	2.0
Male	0.5	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	0.8	1.3	2.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Hypertensive heart disease (I11)									
All races 3/, both sexes	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0
Male	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5
Female	9.7	*	*	*	*	0.4	1.5	3.5	7.8
White, both sexes	7.6	*	*	*	*	0.3	1.5	3.7	7.6
Male	6.1	*	*	*	*	0.4	2.1	5.5	10.4
Female	9.0	*	*	*	*	0.2	0.9	1.9	5.0
Black, both sexes	16.1	*	*	*	*	2.1	7.4	21.2	42.2
Male	15.6	*	*	*	*	2.6	9.2	27.8	54.9
Female	16.5	*	*	*	*	1.7	5.8	15.5	31.9
Hypertensive heart and renal disease (I13)									
All races 3/, both sexes	1.0	*	*	*	*	0.1	0.2	0.4	0.9
Male	0.8	*	*	*	*	*	0.2	0.5	0.9
Female	1.1	*	*	*	*	*	0.1	0.3	0.8
White, both sexes	0.8	*	*	*	*	*	0.1	0.2	0.4
Male	0.7	*	*	*	*	*	0.1	0.2	0.5
Female	1.0	*	*	*	*	*	*	0.1	0.4
Black, both sexes	2.2	*	*	*	*	*	0.8	2.2	4.6
Male	2.0	*	*	*	*	*	1.0	2.8	5.3
Female	2.4	*	*	*	*	*	*	1.6	4.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Ischemic heart diseases (I20-I25)									
All races 3/, both sexes	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7
Male	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2
Female	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4
White, both sexes	197.9	0.8	*	*	0.3	2.1	15.8	62.8	187.3
Male	203.6	*	*	*	0.4	3.3	25.1	100.5	280.3
Female	192.4	*	*	*	0.2	0.9	6.5	25.6	100.0
Black, both sexes	140.1	*	*	*	0.4	4.3	25.6	105.6	314.6
Male	141.5	*	*	*	*	5.9	35.6	149.4	439.2
Female	138.8	*	*	*	*	2.8	16.7	68.0	214.1
Acute myocardial infarction (I21-I22)									
All races 3/, both sexes	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6
Male	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7
Female	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5
White, both sexes	74.1	*	*	*	0.1	0.9	6.8	28.2	84.9
Male	78.8	*	*	*	0.2	1.3	10.6	44.7	125.0
Female	69.6	*	*	*	*	0.4	2.9	11.9	47.3
Black, both sexes	52.2	*	*	*	*	1.7	9.9	41.1	120.1
Male	52.0	*	*	*	*	2.2	13.4	54.9	160.3
Female	52.5	*	*	*	*	1.3	6.8	29.1	87.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other acute ischemic heart diseases (I24)									
All races 3/, both sexes	1.2	*	*	*	*	*	0.2	0.7	2.0
Male	1.3	*	*	*	*	*	0.3	1.2	2.9
Female	1.1	*	*	*	*	*	0.1	0.3	1.1
White, both sexes	1.2	*	*	*	*	*	0.2	0.7	1.8
Male	1.3	*	*	*	*	*	0.3	1.1	2.6
Female	1.2	*	*	*	*	*	*	0.3	0.9
Black, both sexes	1.2	*	*	*	*	*	0.4	1.5	4.2
Male	1.5	*	*	*	*	*	*	2.4	6.1
Female	1.0	*	*	*	*	*	*	*	2.7
Other forms of chronic ischemic heart disease (I20,I25)									
All races 3/, both sexes	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1
Male	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6
Female	111.9	*	*	*	*	0.6	4.3	16.0	58.8
White, both sexes	122.6	*	*	*	0.2	1.2	8.9	33.9	100.6
Male	123.5	*	*	*	0.2	1.9	14.2	54.8	152.7
Female	121.6	*	*	*	*	0.5	3.5	13.4	51.8
Black, both sexes	86.6	*	*	*	*	2.5	15.3	63.1	190.2
Male	88.0	*	*	*	*	3.6	21.6	92.1	272.7
Female	85.3	*	*	*	*	1.5	9.7	38.1	123.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)									
All races 3/, both sexes	24.7	*	*	*	*	0.4	3.0	13.2	34.0
Male	26.0	*	*	*	*	0.6	4.9	21.5	52.6
Female	23.5	*	*	*	*	0.1	1.2	5.2	16.8
White, both sexes	25.1	*	*	*	*	0.3	2.8	11.8	29.4
Male	26.0	*	*	*	*	0.5	4.6	19.6	46.3
Female	24.3	*	*	*	*	*	1.0	4.2	13.5
Black, both sexes	28.5	*	*	*	*	0.7	5.1	26.9	78.2
Male	32.3	*	*	*	*	1.0	7.8	42.0	120.0
Female	25.0	*	*	*	*	*	2.8	14.0	44.6
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
All races 3/, both sexes	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1
Male	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0
Female	88.4	*	*	*	*	0.5	3.0	10.7	42.0
White, both sexes	97.4	*	*	*	0.1	0.9	6.1	22.1	71.2
Male	97.5	*	*	*	0.1	1.4	9.6	35.2	106.4
Female	97.3	*	*	*	*	0.4	2.5	9.3	38.3
Black, both sexes	58.1	*	*	*	*	1.8	10.2	36.1	112.0
Male	55.8	*	*	*	*	2.6	13.9	50.2	152.7
Female	60.3	*	*	*	*	1.1	6.9	24.1	79.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other heart diseases (I26-I51)									
All races 3/, both sexes	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9
Male	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7
Female	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2
White, both sexes	62.5	10.6	1.0	0.5	1.7	3.6	8.1	18.2	46.1
Male	54.6	10.8	1.2	0.6	2.1	4.3	10.3	23.9	59.2
Female	70.2	10.4	0.7	0.4	1.4	2.8	6.0	12.5	33.9
Black, both sexes	52.7	19.0	1.9	1.1	4.7	10.3	23.0	51.6	102.6
Male	51.5	18.9	1.9	1.2	5.4	12.2	28.1	66.5	130.5
Female	53.9	19.2	2.0	1.0	4.1	8.7	18.6	38.7	80.1
Acute and subacute endocarditis (I33)									
All races 3/, both sexes	0.4	*	*	*	*	0.1	0.3	0.5	0.7
Male	0.5	*	*	*	*	0.1	0.3	0.6	0.8
Female	0.4	*	*	*	*	*	0.2	0.3	0.5
White, both sexes	0.4	*	*	*	*	0.1	0.2	0.4	0.6
Male	0.5	*	*	*	*	*	0.3	0.5	0.8
Female	0.3	*	*	*	*	*	0.2	0.2	0.4
Black, both sexes	0.6	*	*	*	*	*	0.5	1.3	1.6
Male	0.6	*	*	*	*	*	0.8	1.8	*
Female	0.5	*	*	*	*	*	*	1.0	1.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
All races 3/, both sexes	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4
Male	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4
Female	0.3	*	*	*	*	0.1	0.1	0.2	0.3
White, both sexes	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3
Male	0.3	*	*	*	*	*	0.2	0.2	0.4
Female	0.2	*	*	*	*	*	*	0.2	0.3
Black, both sexes	0.4	*	*	*	*	*	0.4	0.5	*
Male	0.4	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*
Heart failure (I50)									
All races 3/, both sexes	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1
Male	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9
Female	24.1	*	*	*	*	0.1	0.5	1.6	7.5
White, both sexes	21.8	*	*	*	*	0.1	0.4	1.8	8.2
Male	16.7	*	*	*	*	0.1	0.6	2.2	10.0
Female	26.7	*	*	*	*	*	0.3	1.3	6.6
Black, both sexes	13.6	*	*	*	*	0.4	1.6	5.7	18.6
Male	11.2	*	*	*	*	*	1.7	7.1	21.4
Female	15.7	*	*	*	*	*	1.5	4.5	16.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)									
All races 3/, both sexes	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8
Male	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6
Female	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9
White, both sexes	40.0	9.6	0.8	0.4	1.6	3.3	7.3	15.8	37.0
Male	37.1	9.4	0.9	0.5	1.9	4.0	9.2	20.9	48.0
Female	42.8	9.7	0.7	0.3	1.2	2.5	5.4	10.8	26.6
Black, both sexes	38.2	15.6	1.4	0.9	4.3	9.4	20.5	44.0	81.7
Male	39.2	14.7	1.6	1.0	4.9	11.2	25.0	57.2	106.9
Female	37.3	16.5	*	0.8	3.7	7.8	16.5	32.7	61.4
Essential (primary) hypertension and hypertensive renal disease (I10,I12)									
All races 3/, both sexes	6.4	*	*	*	*	0.2	0.8	2.3	5.9
Male	5.0	*	*	*	*	0.2	1.0	3.0	7.0
Female	7.8	*	*	*	*	0.2	0.6	1.7	4.9
White, both sexes	6.0	*	*	*	*	0.1	0.4	1.4	4.1
Male	4.5	*	*	*	*	0.1	0.6	1.9	5.0
Female	7.4	*	*	*	*	*	0.3	0.9	3.2
Black, both sexes	10.6	*	*	*	*	0.6	3.1	9.7	22.9
Male	9.1	*	*	*	*	*	3.7	11.9	27.8
Female	12.0	*	*	*	*	*	2.6	7.7	18.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other diseases of circulatory system (I71-I78)									
All races 3/, both sexes	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2
Male	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1
Female	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6
White, both sexes	10.0	*	*	*	0.1	0.3	0.8	2.4	8.8
Male	11.1	*	*	*	0.2	0.5	1.0	3.4	12.8
Female	8.9	*	*	*	*	0.2	0.6	1.4	5.1
Black, both sexes	7.1	*	*	*	*	0.7	2.2	5.5	14.4
Male	6.9	*	*	*	*	0.9	2.8	7.5	18.4
Female	7.2	*	*	*	*	*	1.5	3.9	11.1
Aortic aneurysm and dissection (I71)									
All races 3/, both sexes	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2
Male	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5
Female	4.3	*	*	*	*	0.1	0.4	0.9	3.0
White, both sexes	6.2	*	*	*	0.1	0.2	0.6	1.6	6.0
Male	7.8	*	*	*	*	0.3	0.8	2.5	9.5
Female	4.6	*	*	*	*	0.1	0.3	0.7	2.8
Black, both sexes	3.5	*	*	*	*	0.5	1.6	3.2	8.2
Male	3.9	*	*	*	*	*	2.4	4.7	11.5
Female	3.2	*	*	*	*	*	0.9	2.0	5.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Other diseases of arteries, arterioles and capillaries (I72-I78)									
All races 3/, both sexes	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1
Male	3.2	*	*	*	*	0.1	0.3	1.1	3.6
Female	4.0	*	*	*	*	0.1	0.4	0.9	2.6
White, both sexes	3.8	*	*	*	*	0.1	0.3	0.8	2.8
Male	3.4	*	*	*	*	0.1	0.2	0.9	3.3
Female	4.2	*	*	*	*	*	0.3	0.7	2.3
Black, both sexes	3.6	*	*	*	*	*	0.6	2.3	6.2
Male	3.1	*	*	*	*	*	*	2.8	7.0
Female	4.1	*	*	*	*	*	0.7	1.8	5.5
Other disorders of circulatory system (I80-I99)									
All races 3/, both sexes	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1
Male	1.4	*	*	*	*	0.3	0.8	1.6	2.3
Female	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0
White, both sexes	1.6	0.7	*	*	0.1	0.3	0.6	1.2	1.9
Male	1.4	*	*	*	*	0.2	0.8	1.5	2.0
Female	1.9	*	*	*	*	0.3	0.5	1.0	1.8
Black, both sexes	2.1	*	*	*	*	0.9	1.5	2.8	4.5
Male	1.9	*	*	*	*	0.9	1.8	3.4	5.1
Female	2.2	*	*	*	*	0.9	1.3	2.3	3.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Emphysema (J43)									
All races 3/, both sexes	5.9	*	*	*	*	*	0.2	1.3	7.4
Male	6.3	*	*	*	*	*	0.2	1.5	8.1
Female	5.6	*	*	*	*	*	0.2	1.0	6.8
White, both sexes	6.8	*	*	*	*	*	0.2	1.2	7.9
Male	7.0	*	*	*	*	*	0.2	1.4	8.4
Female	6.5	*	*	*	*	*	0.2	1.0	7.4
Black, both sexes	2.5	*	*	*	*	*	*	2.0	6.0
Male	3.2	*	*	*	*	*	*	2.9	8.1
Female	1.8	*	*	*	*	*	*	1.3	4.4
Asthma (J45-J46)									
All races 3/, both sexes	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2
Male	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7
Female	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7
White, both sexes	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7
Male	0.9	*	*	0.2	0.3	0.4	0.6	1.0	1.2
Female	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1
Black, both sexes	3.2	*	*	1.0	1.3	1.9	3.1	5.4	6.9
Male	2.8	*	*	1.2	1.5	2.5	2.9	3.8	6.1
Female	3.5	*	*	0.8	1.0	1.4	3.3	6.8	7.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Pneumonitis due to solids and liquids (J69)									
All races 3/, both sexes	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5
Male	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0
Female	5.8	*	*	*	*	0.1	0.3	0.7	1.9
White, both sexes	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3
Male	6.6	*	*	*	*	0.1	0.5	1.1	2.9
Female	6.3	*	*	*	*	0.2	0.3	0.7	1.8
Black, both sexes	4.2	*	*	*	*	*	0.7	1.9	4.2
Male	4.4	*	*	*	*	*	0.8	2.7	5.0
Female	3.9	*	*	*	*	*	*	1.2	3.5
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)									
All races 3/, both sexes	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9
Male	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4
Female	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6
White, both sexes	9.9	5.6	0.5	0.2	0.3	0.6	1.2	3.2	9.7
Male	9.7	6.6	0.5	0.2	0.3	0.6	1.3	3.4	11.1
Female	10.0	4.6	0.6	0.2	0.3	0.6	1.1	2.9	8.4
Black, both sexes	6.5	16.7	0.8	*	0.7	1.1	2.7	6.8	13.4
Male	6.5	16.3	*	*	0.8	1.2	3.1	7.8	15.2
Female	6.4	17.2	*	*	*	1.0	2.2	5.9	11.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Hernia (K40-K46)									
All races 3/, both sexes	0.5	1.2	*	*	*	*	0.1	0.2	0.5
Male	0.4	1.4	*	*	*	*	*	0.2	0.4
Female	0.7	*	*	*	*	*	*	0.2	0.5
White, both sexes	0.6	1.2	*	*	*	*	0.1	0.2	0.5
Male	0.5	1.5	*	*	*	*	*	0.2	0.4
Female	0.7	*	*	*	*	*	*	0.2	0.5
Black, both sexes	0.4	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*	*
Chronic liver disease and cirrhosis (K70,K73-K74)									
All races 3/, both sexes	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8
Male	12.5	*	*	*	*	1.4	10.3	26.9	34.4
Female	6.5	*	*	*	*	0.7	4.7	8.8	14.0
White, both sexes	10.0	*	*	*	0.1	1.0	7.4	17.5	23.8
Male	13.2	*	*	*	*	1.4	10.3	26.8	34.5
Female	6.8	*	*	*	*	0.6	4.5	8.2	13.7
Black, both sexes	7.5	*	*	*	*	1.0	7.8	22.1	25.9
Male	9.9	*	*	*	*	1.0	10.5	32.7	38.0
Female	5.2	*	*	*	*	1.0	5.5	13.1	16.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Alcoholic liver disease (K70)									
All races 3/, both sexes	4.3	*	*	*	*	0.7	4.9	10.3	12.2
Male	6.5	*	*	*	*	1.0	6.8	16.0	19.0
Female	2.2	*	*	*	*	0.4	3.1	4.8	6.0
White, both sexes	4.5	*	*	*	*	0.7	4.9	10.3	12.2
Male	6.8	*	*	*	*	1.0	6.8	16.2	19.2
Female	2.1	*	*	*	*	0.3	3.0	4.4	5.6
Black, both sexes	3.8	*	*	*	*	0.6	5.0	12.0	13.3
Male	5.3	*	*	*	*	*	6.9	17.4	19.4
Female	2.4	*	*	*	*	*	3.3	7.3	8.4
Other chronic liver disease and cirrhosis (K73-K74)									
All races 3/, both sexes	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6
Male	6.0	*	*	*	*	0.4	3.5	10.9	15.4
Female	4.3	*	*	*	*	0.3	1.6	4.0	8.0
White, both sexes	5.5	*	*	*	*	0.3	2.5	7.2	11.6
Male	6.4	*	*	*	*	0.3	3.5	10.6	15.3
Female	4.7	*	*	*	*	0.2	1.5	3.8	8.1
Black, both sexes	3.7	*	*	*	*	0.4	2.8	10.1	12.6
Male	4.7	*	*	*	*	*	3.6	15.2	18.6
Female	2.8	*	*	*	*	*	2.2	5.8	7.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Cholelithiasis and other disorders of gallbladder (K80-K82)									
All races 3/, both sexes	1.0	*	*	*	*	*	0.1	0.3	0.8
Male	0.9	*	*	*	*	*	0.1	0.3	0.9
Female	1.1	*	*	*	*	*	0.1	0.3	0.6
White, both sexes	1.1	*	*	*	*	*	0.1	0.2	0.7
Male	1.0	*	*	*	*	*	0.1	0.2	0.9
Female	1.2	*	*	*	*	*	*	0.2	0.5
Black, both sexes	0.8	*	*	*	*	*	*	0.7	1.2
Male	0.6	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	*	*	*
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)									
All races 3/, both sexes	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8
Male	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2
Female	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5
White, both sexes	12.9	2.9	*	0.1	0.2	0.4	1.1	3.0	9.7
Male	12.7	3.8	*	*	0.2	0.4	1.3	3.6	11.0
Female	13.0	2.0	*	*	0.2	0.4	0.9	2.4	8.4
Black, both sexes	18.9	11.5	*	*	0.4	1.8	5.3	15.8	40.6
Male	17.7	13.4	*	*	*	1.9	6.5	20.3	44.9
Female	20.0	9.6	*	*	*	1.7	4.3	11.9	37.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Inflammatory diseases of female pelvic organs (N70-N76)									
All races 3/, both sexes	0.0	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*
Pregnancy, childbirth and the puerperium (000-099)									
All races 3/, both sexes	0.1	*	0.2	0.5	0.3	*	...
Female	0.3	*	0.5	0.9	0.5	*	...
White, both sexes	0.1	*	0.2	0.3	0.2	*	...
Female	0.2	*	0.4	0.7	0.4	*	...
Black, both sexes	0.4	*	0.6	1.2	0.7	*	...
Female	0.7	*	1.2	2.3	1.2	*	...

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Pregnancy with abortive outcome (000-007)									
All races 3/, both sexes	0.0	*	*	*	*	*	...
Female	0.0	*	*	*	*	*	...
White, both sexes	*	*	*	*	*	*	...
Female	*	*	*	*	*	*	...
Black, both sexes	*	*	*	*	*	*	...
Female	*	*	*	*	*	*	...
Other complications of pregnancy, childbirth and the puerperium (010-099)									
All races 3/, both sexes	0.1	*	0.2	0.4	0.2	*	...
Female	0.3	*	0.5	0.8	0.5	*	...
White, both sexes	0.1	*	0.2	0.3	0.2	*	...
Female	0.2	*	0.4	0.6	0.4	*	...
Black, both sexes	0.3	*	0.5	1.0	0.6	*	...
Female	0.6	*	1.0	2.0	1.1	*	...

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes	5.0	365.7	0.5	0.1	*	*	*	*	*
Male	5.8	403.9	0.7	0.1	*	*	*	*	*
Female	4.3	325.6	0.4	*	*	*	*	*	*
White, both sexes	3.7	283.7	0.4	0.1	*	*	*	*	*
Male	4.3	312.4	0.5	0.1	*	*	*	*	*
Female	3.2	253.5	*	*	*	*	*	*	*
Black, both sexes	13.7	809.5	1.1	*	*	*	*	*	*
Male	16.3	901.3	*	*	*	*	*	*	*
Female	11.3	714.6	*	*	*	*	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3
Male	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3
Female	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4
White, both sexes	3.6	147.1	2.9	0.9	1.1	1.2	1.2	1.7	2.4
Male	3.9	151.9	2.9	0.9	1.3	1.4	1.4	1.9	2.4
Female	3.5	142.0	2.9	0.9	0.9	1.0	1.1	1.5	2.4
Black, both sexes	4.9	179.2	5.0	1.3	1.5	1.4	1.6	1.6	2.4
Male	5.4	181.4	5.5	1.5	2.0	1.8	1.9	2.1	*
Female	4.4	176.9	4.6	1.2	1.1	1.1	1.4	1.1	3.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3
Male	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3
Female	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5
White, both sexes	11.4	77.2	1.5	0.4	2.1	3.7	6.5	7.8	8.6
Male	11.3	89.4	1.7	0.4	2.9	5.1	8.9	10.7	11.2
Female	11.5	64.3	1.3	0.3	1.3	2.3	4.1	4.9	6.1
Black, both sexes	13.7	197.1	2.3	0.7	3.2	5.8	9.2	14.7	17.4
Male	15.5	227.1	2.3	0.8	3.7	7.7	10.8	20.6	25.2
Female	12.0	166.0	2.3	*	2.6	4.1	7.7	9.7	11.0

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4
Male	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7
Female	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8
White, both sexes	67.0	24.9	3.5	2.0	4.3	6.9	16.0	32.0	58.6
Male	54.1	26.8	3.9	2.2	5.2	8.3	19.5	39.3	66.0
Female	79.6	22.9	3.2	1.8	3.4	5.4	12.5	24.7	51.6
Black, both sexes	58.1	49.6	5.9	2.8	6.5	15.0	34.2	69.3	107.4
Male	54.3	59.8	6.1	2.5	7.0	15.6	38.7	81.1	130.2
Female	61.6	39.0	5.7	3.0	6.0	14.4	30.2	59.1	89.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
All races 3/, both sexes	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9
Male	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1
Female	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8
White, both sexes	35.9	19.1	11.1	6.9	38.3	29.9	34.3	31.3	30.2
Male	47.0	21.7	13.2	8.4	55.0	45.4	51.0	46.7	42.1
Female	25.1	16.4	9.0	5.4	20.4	13.8	17.4	16.2	18.9
Black, both sexes	33.5	44.5	16.2	9.3	28.7	32.5	39.7	47.0	40.8
Male	49.0	47.7	18.8	12.0	43.0	51.8	62.0	75.4	68.8
Female	19.5	41.3	13.5	6.5	14.6	15.0	19.9	22.6	18.3
Transport accidents (V01-V99,Y85)									
All races 3/, both sexes	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0
Male	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5
Female	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0
White, both sexes	17.0	4.1	4.3	4.7	30.0	18.5	17.1	15.6	16.0
Male	23.6	4.5	4.7	5.5	41.6	27.4	24.7	22.6	22.0
Female	10.5	3.7	3.8	4.0	17.7	9.3	9.4	8.7	10.3
Black, both sexes	16.0	6.5	5.6	4.7	21.7	21.4	18.6	19.3	17.4
Male	24.1	6.7	6.2	5.7	31.6	34.4	29.7	29.6	29.4
Female	8.5	*	5.0	3.6	11.9	9.7	8.7	10.5	7.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)									
All races 3/, both sexes	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1
Male	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8
Female	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3
White, both sexes	0.7	*	*	0.1	0.6	0.9	1.0	1.0	1.1
Male	1.3	*	*	0.2	1.0	1.4	1.8	1.8	1.9
Female	0.2	*	*	*	0.3	0.2	0.3	0.3	0.4
Black, both sexes	0.4	*	*	*	*	0.5	0.5	0.6	*
Male	0.7	*	*	*	*	0.9	1.0	1.1	*
Female	*	*	*	*	*	*	*	*	*
Nontransport accidents (W00-X59,Y86)									
All races 3/, both sexes	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9
Male	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6
Female	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8
White, both sexes	18.9	15.1	6.9	2.2	8.2	11.4	17.2	15.8	14.2
Male	23.4	17.3	8.4	2.9	13.4	18.0	26.2	24.1	20.2
Female	14.6	12.8	5.2	1.4	2.7	4.5	8.0	7.5	8.6
Black, both sexes	17.6	38.0	10.6	4.6	7.0	11.0	21.1	27.7	23.5
Male	24.9	40.9	12.6	6.3	11.4	17.4	32.3	45.8	39.3
Female	11.0	35.1	8.5	2.9	2.7	5.3	11.3	12.1	10.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Accidental drowning and submersion (W65-W74)									
All races 3/, both sexes	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9
Male	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5
Female	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3
White, both sexes	1.2	1.6	3.5	0.7	1.5	1.0	1.0	0.9	0.9
Male	1.9	1.8	4.4	0.9	2.6	1.7	1.7	1.5	1.6
Female	0.5	*	2.5	0.4	0.3	0.2	0.3	0.3	0.3
Black, both sexes	1.6	4.2	2.2	2.0	2.3	1.1	1.4	1.2	*
Male	2.7	*	2.8	3.0	4.3	2.0	2.6	2.3	*
Female	0.6	*	1.7	1.0	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)									
All races 3/, both sexes	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5
Male	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1
Female	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0
White, both sexes	1.0	0.8	1.3	0.5	0.5	0.5	0.8	1.0	1.3
Male	1.3	*	1.6	0.5	0.6	0.7	1.1	1.5	1.7
Female	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.8
Black, both sexes	2.5	*	4.8	1.5	0.7	1.2	1.7	2.4	4.2
Male	3.2	*	5.6	1.7	0.9	1.5	2.4	3.5	6.3
Female	2.0	*	4.1	1.3	*	0.9	1.1	1.4	2.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Accidental poisoning and exposure to noxious substances (X40-X49)									
All races 3/, both sexes	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8
Male	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4
Female	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3
White, both sexes	4.6	*	*	0.1	3.4	6.4	10.4	7.5	2.6
Male	6.7	*	*	*	5.3	9.7	15.2	10.9	2.9
Female	2.5	*	*	*	1.3	3.0	5.5	4.1	2.3
Black, both sexes	5.5	*	*	*	1.3	5.6	13.0	15.5	5.5
Male	8.2	*	*	*	1.7	8.4	18.7	24.9	9.5
Female	3.1	*	*	*	1.0	3.0	7.8	7.4	2.3
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)									
All races 3/, both sexes	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6
Male	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3
Female	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0
White, both sexes	6.6	12.2	1.6	0.7	1.7	2.4	3.2	3.8	5.4
Male	7.4	14.0	2.0	0.9	2.9	4.0	5.4	6.2	7.9
Female	5.7	10.3	1.2	0.4	0.5	0.6	1.0	1.5	3.0
Black, both sexes	5.4	30.6	2.3	0.7	1.5	2.2	3.5	5.6	8.2
Male	7.3	32.0	2.5	1.0	2.5	3.7	6.0	10.1	13.8
Female	3.8	29.1	2.0	*	*	0.9	1.3	1.8	3.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Intentional self-harm (suicide) (X60-X84,Y87.0)									
All races 3/, both sexes	10.4	0.7	10.2	12.0	14.5	14.4	12.1
Male	17.1	1.2	17.1	19.6	22.8	22.4	19.4
Female	4.0	0.3	3.0	4.3	6.4	6.7	5.4
White, both sexes	11.5	0.8	10.7	12.8	16.1	15.9	13.3
Male	18.8	1.2	17.9	20.7	24.9	24.5	21.2
Female	4.4	0.3	3.1	4.6	7.2	7.5	6.0
Black, both sexes	5.4	0.6	8.1	8.7	7.6	6.5	4.2
Male	9.4	1.0	14.2	15.5	13.2	11.2	7.5
Female	1.7	*	2.2	2.6	2.7	2.5	1.6
Intentional self-harm (suicide) by discharge of firearms (X72-X74)									
All races 3/, both sexes	5.9	0.3	5.8	6.1	7.1	7.6	7.6
Male	10.5	0.4	10.2	10.6	11.8	13.0	13.5
Female	1.5	0.1	1.2	1.7	2.4	2.3	2.1
White, both sexes	6.6	0.3	6.1	6.6	7.9	8.5	8.3
Male	11.6	0.5	10.6	11.2	13.1	14.4	14.7
Female	1.7	*	1.2	1.8	2.7	2.6	2.4
Black, both sexes	3.0	*	5.2	5.0	3.4	3.3	2.9
Male	5.5	*	9.6	9.3	6.2	5.8	5.7
Female	0.7	*	0.9	1.1	0.8	1.1	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races 3/, both sexes	4.5	0.5	4.4	5.9	7.5	6.8	4.6
Male	6.6	0.7	6.9	9.0	10.9	9.4	6.0
Female	2.5	0.2	1.8	2.7	4.0	4.4	3.3
White, both sexes	4.9	0.5	4.6	6.2	8.1	7.5	5.0
Male	7.1	0.7	7.3	9.5	11.8	10.0	6.5
Female	2.8	0.2	1.9	2.8	4.5	4.9	3.6
Black, both sexes	2.4	0.5	2.9	3.7	4.2	3.2	1.3
Male	3.9	0.9	4.6	6.3	6.9	5.4	1.8
Female	1.0	*	1.3	1.5	1.9	1.4	*

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0
Male	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6
Female	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6
White, both sexes	3.6	6.6	1.7	0.7	6.4	5.7	4.7	3.3	2.4
Male	5.2	8.2	1.9	0.9	9.9	8.4	6.5	4.6	3.5
Female	2.1	5.0	1.5	0.6	2.7	2.9	2.8	2.1	1.4
Black, both sexes	21.5	22.8	5.8	1.8	47.8	40.8	23.9	14.8	8.3
Male	37.2	23.3	5.9	2.1	85.3	73.9	38.5	26.0	14.7
Female	7.2	22.2	5.7	1.6	10.7	10.9	11.0	5.3	3.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Assault (homicide) by discharge of firearms (X93-X95)									
All races 3/, both sexes	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5
Male	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3
Female	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8
White, both sexes	2.1	*	*	0.4	4.6	3.7	2.7	1.8	1.3
Male	3.2	*	*	0.5	7.7	5.9	3.9	2.6	1.8
Female	1.0	*	*	0.2	1.4	1.5	1.4	1.0	0.8
Black, both sexes	15.6	*	*	1.0	41.2	33.2	15.0	8.4	3.5
Male	29.2	*	*	1.5	76.5	63.3	26.7	15.1	6.3
Female	3.2	*	*	*	6.4	6.1	4.6	2.6	*

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5
Male	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3
Female	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8
White, both sexes	1.5	6.5	1.5	0.4	1.8	2.0	2.0	1.5	1.1
Male	2.0	8.1	1.7	0.4	2.3	2.5	2.6	2.0	1.7
Female	1.1	4.8	1.3	0.4	1.3	1.4	1.4	1.1	0.6
Black, both sexes	5.9	21.6	5.5	0.8	6.5	7.6	8.9	6.5	4.9
Male	8.0	21.1	5.5	*	8.8	10.6	11.7	10.8	8.3
Female	4.0	22.2	5.5	1.1	4.3	4.8	6.4	2.7	2.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Legal intervention (Y35,Y89.0)									
All races 3/, both sexes	0.1	*	*	*	0.2	0.3	0.2	0.1	*
Male	0.3	*	*	*	0.4	0.5	0.4	0.3	*
Female	*	*	*	*	*	*	*	*	*
White, both sexes	0.1	*	*	*	0.1	0.2	0.2	0.1	*
Male	0.2	*	*	*	0.3	0.4	0.3	0.3	*
Female	*	*	*	*	*	*	*	*	*
Black, both sexes	0.3	*	*	*	0.5	0.6	0.6	*	*
Male	0.6	*	*	*	1.0	1.3	1.2	*	*
Female	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)									
All races 3/, both sexes	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9
Male	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3
Female	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6
White, both sexes	1.3	1.4	0.2	0.1	1.0	1.7	2.8	2.1	0.9
Male	1.8	*	*	0.2	1.4	2.4	3.8	2.8	1.1
Female	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6
Black, both sexes	1.8	*	*	0.3	0.9	1.9	4.1	3.7	1.8
Male	2.8	*	*	*	1.6	3.1	6.0	5.9	3.0
Female	0.9	*	*	*	*	0.9	2.4	1.8	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Discharge of firearms, undetermined intent (Y22-Y24)									
All races 3/, both sexes	0.1	*	*	*	0.2	0.1	0.1	0.1	*
Male	0.1	*	*	*	0.3	0.2	0.1	*	*
Female	0.0	*	*	*	*	*	*	*	*
White, both sexes	0.1	*	*	*	0.2	0.1	0.1	0.1	*
Male	0.1	*	*	*	0.3	0.2	0.1	*	*
Female	0.0	*	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	0.3	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9
Male	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2
Female	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6
White, both sexes	1.3	1.4	0.2	0.1	0.8	1.6	2.7	2.1	0.8
Male	1.7	*	*	0.2	1.2	2.2	3.7	2.8	1.0
Female	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6
Black, both sexes	1.7	*	*	*	0.5	1.7	3.9	3.6	1.6
Male	2.5	*	*	*	1.0	2.6	5.8	5.8	2.8
Female	0.9	*	*	*	*	0.8	2.3	1.8	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Operations of war and their sequelae (Y36,Y89.1)									
All races 3/, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)									
All races 3/, both sexes	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5
Male	1.0	*	*	*	*	0.2	0.3	0.8	1.6
Female	1.1	*	*	*	*	0.2	0.3	0.8	1.5
White, both sexes	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4
Male	1.1	*	*	*	*	0.2	0.3	0.7	1.4
Female	1.1	*	*	*	*	0.2	0.3	0.7	1.3
Black, both sexes	1.3	*	*	*	*	0.4	0.5	1.4	3.5
Male	1.1	*	*	*	*	*	*	1.3	3.1
Female	1.5	*	*	*	*	*	*	1.5	3.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

All Causes

All races 3/, both sexes	2,399.1	5,666.5	15,524.4
Male	2,979.6	6,972.6	17,501.4
Female	1,921.2	4,814.7	14,719.2
White, both sexes	2,341.5	5,637.7	15,707.0
Male	2,905.7	6,933.1	17,716.4
Female	1,868.3	4,785.3	14,890.7
Black, both sexes	3,335.6	6,701.3	14,714.3
Male	4,253.5	8,486.0	16,791.0
Female	2,689.6	5,696.5	13,941.3

Salmonella infections (A01-A02)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84
YEARS 85 YEARS
AND OVER

Shigellosis and amebiasis (A03,A06)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes	1.2	3.7	11.8
Male	1.3	4.0	11.4
Female	1.2	3.6	11.9
White, both sexes	1.3	4.0	12.2
Male	1.3	4.2	11.9
Female	1.2	3.8	12.3
Black, both sexes	*	*	8.1
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84 85 YEARS
YEARS YEARS AND OVER

Tuberculosis (A16-A19)

All races 3/, both sexes	0.8	2.0	2.7
Male	1.0	2.8	4.6
Female	0.7	1.5	2.0
White, both sexes	0.5	1.4	2.2
Male	0.7	1.8	3.8
Female	0.4	1.1	1.6
Black, both sexes	3.1	6.5	*
Male	4.1	12.1	*
Female	2.5	3.4	*

Respiratory tuberculosis (A16)

All races 3/, both sexes	0.6	1.7	2.3
Male	0.9	2.4	4.1
Female	0.5	1.2	1.6
White, both sexes	0.4	1.2	1.9
Male	0.6	1.5	3.4
Female	0.3	1.0	1.2
Black, both sexes	2.6	5.2	*
Male	3.8	9.7	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84 85 YEARS
YEARS YEARS AND OVER

Other tuberculosis (A17-A19)

All races 3/, both sexes	0.2	0.3	*
Male	*	0.5	*
Female	0.2	0.3	*
White, both sexes	0.1	0.2	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Whooping cough (A37)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Scarlet fever and erysipelas (A38,A46)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Meningococcal infection (A39)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Septicemia (A40-A41)

All races 3/, both sexes	31.0	80.4	215.7
Male	34.5	91.9	223.0
Female	28.1	72.9	212.7
White, both sexes	27.4	73.9	203.0
Male	30.8	84.6	209.8
Female	24.6	66.8	200.2
Black, both sexes	71.9	173.4	392.6
Male	81.2	207.1	427.8
Female	65.3	154.4	379.5

Syphilis (A50-A53)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Acute poliomyelitis (A80)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Arthropod-borne viral encephalitis (A83-A84,A85.2)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Measles (B05)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Viral hepatitis (B15-B19)

All races 3/, both sexes	4.6	5.3	4.0
Male	4.8	6.4	5.7
Female	4.5	4.7	3.3
White, both sexes	4.2	5.1	3.9
Male	4.3	6.2	5.7
Female	4.1	4.5	3.2
Black, both sexes	5.7	6.2	*
Male	7.0	7.6	*
Female	4.7	5.5	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS

75-84
YEARS

85 YEARS
AND OVER

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes	2.2	0.7	*
Male	3.8	1.3	*
Female	0.8	0.3	*
White, both sexes	1.0	0.4	*
Male	1.9	0.7	*
Female	0.3	*	*
Black, both sexes	13.5	4.8	*
Male	24.6	9.4	*
Female	5.8	*	*

Malaria (B50-B54)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes	5.8	11.1	21.2
Male	7.2	13.6	27.6
Female	4.7	9.5	18.6
White, both sexes	5.7	11.0	21.0
Male	7.0	13.3	26.9
Female	4.6	9.5	18.7
Black, both sexes	7.8	13.3	24.1
Male	10.2	18.2	32.3
Female	6.1	10.6	21.0

Malignant neoplasms (C00-C97)

All races 3/, both sexes	816.3	1,335.6	1,819.4
Male	1,001.9	1,760.6	2,710.7
Female	663.6	1,058.5	1,456.4
White, both sexes	811.0	1,330.3	1,815.7
Male	984.3	1,736.0	2,693.7
Female	665.6	1,063.4	1,459.1
Black, both sexes	1,005.3	1,566.3	1,998.9
Male	1,363.8	2,351.8	3,264.8
Female	753.1	1,124.0	1,527.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes	10.7	13.2	19.4
Male	16.0	19.2	27.1
Female	6.3	9.2	16.3
White, both sexes	10.2	13.1	19.7
Male	14.9	18.7	27.6
Female	6.3	9.4	16.5
Black, both sexes	15.5	15.1	15.3
Male	28.2	28.8	*
Female	6.5	7.3	14.2

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes	20.0	25.6	25.9
Male	34.1	45.8	47.8
Female	8.3	12.4	17.1
White, both sexes	19.5	25.3	26.6
Male	33.6	45.6	49.3
Female	7.8	11.9	17.3
Black, both sexes	28.2	33.7	22.2
Male	47.0	58.4	38.1
Female	14.9	19.8	16.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes	16.6	31.0	52.7
Male	23.7	43.9	75.6
Female	10.8	22.6	43.4
White, both sexes	14.5	27.3	47.3
Male	20.9	39.0	68.9
Female	9.2	19.6	38.6
Black, both sexes	32.0	63.9	101.7
Male	46.7	91.1	136.1
Female	21.6	48.6	88.9

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes	76.7	146.0	270.4
Male	95.4	179.5	315.6
Female	61.2	124.2	251.9
White, both sexes	75.1	144.0	269.2
Male	93.4	176.9	313.3
Female	59.7	122.3	251.3
Black, both sexes	104.7	190.1	310.0
Male	131.5	241.0	377.1
Female	85.8	161.4	285.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX

65-74 YEARS 75-84 YEARS 85 YEARS AND OVER

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes	18.7	28.8	32.7
Male	27.0	42.1	44.7
Female	11.8	20.1	27.8
White, both sexes	17.3	27.8	32.0
Male	25.0	41.1	43.1
Female	10.9	19.0	27.4
Black, both sexes	23.6	28.7	31.6
Male	37.3	39.7	48.4
Female	14.0	22.5	25.3

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes	44.1	76.5	96.4
Male	50.4	85.9	102.9
Female	38.8	70.4	93.8
White, both sexes	43.1	75.3	94.7
Male	49.8	85.2	102.4
Female	37.5	68.8	91.5
Black, both sexes	59.1	97.8	119.8
Male	65.8	106.9	109.6
Female	54.4	92.7	123.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Malignant neoplasm of larynx (C32)

All races 3/, both sexes	6.6	8.1	6.5
Male	11.4	15.7	17.4
Female	2.7	3.1	2.1
White, both sexes	6.1	7.7	6.6
Male	10.4	15.0	17.6
Female	2.6	2.9	2.1
Black, both sexes	12.8	14.1	6.9
Male	24.3	28.5	*
Female	4.6	6.0	*

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes	284.2	370.8	302.1
Male	377.9	532.2	521.2
Female	206.9	265.6	212.8
White, both sexes	287.0	374.7	304.5
Male	374.9	529.9	522.4
Female	213.2	272.7	215.9
Black, both sexes	315.8	375.0	288.1
Male	488.8	642.5	562.8
Female	194.1	224.3	185.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Malignant melanoma of skin (C43)

All races 3/, both sexes	8.9	13.1	18.9
Male	12.9	20.8	31.8
Female	5.7	8.2	13.7
White, both sexes	9.9	14.3	20.3
Male	14.3	22.5	34.7
Female	6.3	8.8	14.5
Black, both sexes	1.8	3.5	6.6
Male	*	*	*
Female	*	*	*

Malignant neoplasm of breast (C50)

All races 3/, both sexes	49.1	78.9	147.4
Male	1.4	2.2	4.2
Female	88.3	128.9	205.7
White, both sexes	49.2	79.4	147.3
Male	1.4	2.2	3.9
Female	89.3	130.2	205.5
Black, both sexes	58.6	90.5	176.4
Male	*	*	*
Female	98.6	139.8	238.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of cervix uteri (C53)			
All races 3/, both sexes	3.7	4.8	7.3
Female	6.8	8.0	10.3
White, both sexes	3.1	4.3	6.3
Female	5.7	7.1	8.8
Black, both sexes	9.5	11.9	19.7
Female	16.3	18.6	27.0
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)			
All races 3/, both sexes	9.7	15.8	24.7
Female	17.7	26.2	34.8
White, both sexes	8.5	14.8	24.4
Female	15.7	24.6	34.3
Black, both sexes	22.7	30.5	33.2
Female	38.6	47.7	45.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of ovary (C56)			
All races 3/, both sexes	19.6	33.2	40.9
Female	35.7	54.8	57.6
White, both sexes	20.1	34.3	42.0
Female	37.0	56.8	59.0
Black, both sexes	18.1	26.6	34.4
Female	30.8	41.6	47.2
Malignant neoplasm of prostate (C61)			
All races 3/, both sexes	37.0	105.7	209.3
Male	82.0	267.9	723.1
White, both sexes	32.5	97.0	198.6
Male	71.2	244.5	687.5
Black, both sexes	90.1	229.8	356.6
Male	218.2	638.0	1,314.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races 3/, both sexes	17.1	26.6	33.8
Male	24.2	38.2	55.1
Female	11.3	18.9	25.1
White, both sexes	17.5	27.0	34.5
Male	24.4	38.7	56.6
Female	11.6	19.2	25.5
Black, both sexes	16.5	25.6	29.7
Male	26.3	38.2	49.6
Female	9.6	18.6	22.3

Malignant neoplasm of bladder (C67)

All races 3/, both sexes	15.5	35.7	69.6
Male	25.2	62.6	139.4
Female	7.5	18.1	41.2
White, both sexes	16.2	36.4	72.0
Male	26.5	64.7	145.7
Female	7.5	17.8	42.1
Black, both sexes	13.3	34.7	49.7
Male	17.6	52.7	88.8
Female	10.3	24.5	35.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes	16.5	20.0	16.9
Male	20.2	24.3	22.4
Female	13.4	17.3	14.7
White, both sexes	17.8	21.0	17.5
Male	21.7	25.5	23.5
Female	14.5	18.0	15.0
Black, both sexes	8.1	11.9	10.6
Male	9.8	13.0	*
Female	6.9	11.3	10.7

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes	76.3	149.2	204.6
Male	96.8	192.6	281.8
Female	59.5	120.9	173.1
White, both sexes	78.0	153.3	208.7
Male	98.7	198.2	290.2
Female	60.6	123.8	175.5
Black, both sexes	75.0	121.0	175.5
Male	96.0	155.0	224.9
Female	60.2	101.8	157.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Hodgkin's disease (C81)			
All races 3/, both sexes	1.3	2.1	2.5
Male	1.6	2.5	3.7
Female	1.0	1.8	2.0
White, both sexes	1.3	2.2	2.6
Male	1.7	2.7	3.9
Female	1.0	1.9	2.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Non-Hodgkin's lymphoma (C82-C85)			
All races 3/, both sexes	30.4	61.0	80.2
Male	37.4	74.5	105.9
Female	24.7	52.1	69.8
White, both sexes	32.1	64.3	84.3
Male	39.2	78.4	111.8
Female	26.1	55.0	73.1
Black, both sexes	19.3	26.8	36.3
Male	24.9	31.2	43.8
Female	15.4	24.4	33.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Leukemia (C91-C95)

All races 3/, both sexes	27.2	54.5	83.6
Male	37.2	75.7	119.8
Female	19.0	40.6	68.8
White, both sexes	28.3	56.6	86.0
Male	38.5	78.5	123.6
Female	19.8	42.1	70.8
Black, both sexes	22.7	38.2	66.0
Male	31.6	54.8	98.0
Female	16.4	28.8	54.1

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes	17.3	31.5	37.8
Male	20.4	39.6	51.8
Female	14.7	26.3	32.2
White, both sexes	16.2	30.0	35.2
Male	19.3	38.2	50.2
Female	13.6	24.6	29.1
Black, both sexes	32.1	55.1	71.6
Male	38.5	67.8	79.6
Female	27.6	47.9	68.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

65-74 YEARS 75-84 YEARS 85 YEARS AND OVER

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes	0.1	0.2	0.5
Male	*	*	*
Female	*	*	*
White, both sexes	0.1	0.2	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	85.5	152.6	239.8
Male	103.2	187.8	300.7
Female	70.8	129.6	215.0
White, both sexes	85.3	153.3	243.7
Male	103.1	188.2	307.0
Female	70.3	130.3	218.0
Black, both sexes	100.1	162.0	210.8
Male	122.6	210.4	253.7
Female	84.3	134.7	194.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX65-74 75-84 85 YEARS
YEARS YEARS AND OVER

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes	13.8	37.6	84.9
Male	17.0	47.5	108.9
Female	11.2	31.1	75.2
White, both sexes	13.9	38.6	87.3
Male	17.0	49.1	112.6
Female	11.3	31.8	77.1
Black, both sexes	15.0	31.0	64.1
Male	19.2	36.0	76.1
Female	12.0	28.1	59.7

Anemias (D50-D64)

All races 3/, both sexes	2.8	9.5	42.4
Male	2.8	9.5	40.1
Female	2.8	9.4	43.3
White, both sexes	2.5	9.3	42.0
Male	2.7	9.2	39.1
Female	2.4	9.3	43.2
Black, both sexes	5.4	13.3	51.0
Male	4.5	13.9	55.4
Female	6.0	13.0	49.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Diabetes mellitus (E10-E14)			
All races 3/, both sexes	90.7	179.5	319.7
Male	99.1	199.6	345.1
Female	83.7	166.3	309.4
White, both sexes	81.2	166.0	303.5
Male	91.8	190.5	337.4
Female	72.4	149.8	289.7
Black, both sexes	188.4	350.6	524.9
Male	184.9	337.0	452.1
Female	190.9	358.3	552.0
Nutritional deficiencies (E40-E64)			
All races 3/, both sexes	2.5	10.4	50.0
Male	2.7	9.7	43.7
Female	2.4	10.9	52.6
White, both sexes	2.3	10.0	48.8
Male	2.3	9.1	42.8
Female	2.3	10.5	51.3
Black, both sexes	4.7	16.1	66.9
Male	6.4	17.9	63.4
Female	3.5	15.2	68.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Malnutrition (E40-E46)			
All races 3/, both sexes	2.3	9.7	46.3
Male	2.5	8.9	40.7
Female	2.1	10.2	48.6
White, both sexes	2.1	9.3	45.2
Male	2.2	8.4	39.7
Female	2.0	9.9	47.4
Black, both sexes	4.7	14.9	63.8
Male	6.3	17.3	61.1
Female	3.5	13.6	64.8
Other nutritional deficiencies (E50-E64)			
All races 3/, both sexes	0.2	0.7	3.7
Male	*	0.8	3.0
Female	0.2	0.7	4.0
White, both sexes	0.2	0.7	3.7
Male	*	0.7	3.2
Female	0.3	0.6	3.9
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Meningitis (G00,G03)			
All races 3/, both sexes	0.6	1.0	1.2
Male	0.7	1.3	*
Female	0.5	0.8	1.0
White, both sexes	0.5	0.9	1.0
Male	0.7	1.2	*
Female	0.4	0.8	1.0
Black, both sexes	1.6	*	*
Male	*	*	*
Female	*	*	*
Parkinson's disease (G20-G21)			
All races 3/, both sexes	11.5	61.9	131.9
Male	16.1	92.8	222.3
Female	7.6	41.7	95.0
White, both sexes	12.3	65.6	139.5
Male	17.2	97.5	235.3
Female	8.2	44.7	100.6
Black, both sexes	5.7	27.4	51.6
Male	8.8	47.8	86.5
Female	3.5	15.9	38.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Alzheimer's disease (G30)			
All races 3/, both sexes	18.7	139.6	667.7
Male	18.7	127.6	522.1
Female	18.6	147.4	727.0
White, both sexes	19.5	144.8	695.3
Male	19.5	132.5	542.1
Female	19.5	152.9	757.6
Black, both sexes	15.8	109.9	445.1
Male	16.5	97.5	371.3
Female	15.2	116.9	472.6
Major cardiovascular diseases (I00-I78)			
All races 3/, both sexes	849.3	2,396.2	8,030.3
Male	1,114.1	2,926.6	8,448.1
Female	631.2	2,050.4	7,860.1
White, both sexes	814.2	2,370.8	8,133.9
Male	1,080.7	2,912.4	8,585.0
Female	590.8	2,014.4	7,950.6
Black, both sexes	1,316.2	2,968.8	7,518.9
Male	1,639.9	3,501.8	7,687.4
Female	1,088.3	2,668.7	7,456.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION

OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS

75-84
YEARS

85 YEARS
AND OVER

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes	665.6	1,780.3	5,926.1
Male	898.3	2,248.1	6,430.0
Female	474.0	1,475.1	5,720.9
White, both sexes	642.5	1,767.8	6,020.8
Male	877.8	2,247.0	6,560.8
Female	445.2	1,452.4	5,801.4
Black, both sexes	1,002.6	2,159.7	5,389.5
Male	1,268.8	2,597.6	5,633.5
Female	815.2	1,913.1	5,298.7

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes	4.5	9.5	16.8
Male	2.9	6.6	12.2
Female	5.8	11.4	18.7
White, both sexes	4.6	10.1	17.6
Male	2.9	7.0	12.7
Female	6.0	12.2	19.6
Black, both sexes	3.5	3.9	8.4
Male	*	*	*
Female	4.3	4.8	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS

75-84
YEARS

85 YEARS
AND OVER

Hypertensive heart disease (I11)

All races 3/, both sexes	19.6	48.2	193.1
Male	23.7	46.5	139.1
Female	16.2	49.4	215.1
White, both sexes	15.0	41.5	181.8
Male	18.6	39.6	129.3
Female	11.9	42.9	203.1
Black, both sexes	66.8	131.9	340.0
Male	81.4	139.3	273.3
Female	56.6	127.7	364.9

Hypertensive heart and renal disease (I13)

All races 3/, both sexes	2.3	7.0	24.3
Male	2.5	7.2	27.8
Female	2.1	6.9	22.9
White, both sexes	1.5	5.6	22.0
Male	1.7	6.2	24.4
Female	1.4	5.2	21.1
Black, both sexes	9.9	22.8	50.4
Male	11.3	20.9	63.4
Female	9.0	23.9	45.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes	505.1	1,324.2	4,182.5
Male	705.2	1,736.8	4,701.7
Female	340.3	1,055.2	3,971.0
White, both sexes	493.4	1,320.9	4,257.7
Male	697.3	1,744.8	4,813.9
Female	322.5	1,042.0	4,031.8
Black, both sexes	702.2	1,523.2	3,684.7
Male	907.9	1,877.0	3,885.2
Female	557.4	1,324.0	3,610.0

Acute myocardial infarction (I21-I22)

All races 3/, both sexes	209.1	503.3	1,338.5
Male	286.9	648.3	1,568.3
Female	145.1	408.7	1,244.8
White, both sexes	206.7	502.6	1,353.5
Male	287.2	652.5	1,592.9
Female	139.3	404.1	1,256.2
Black, both sexes	268.7	571.1	1,275.7
Male	335.8	687.3	1,418.5
Female	221.6	505.6	1,222.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other acute ischemic heart diseases (I24)

All races 3/, both sexes	3.5	7.3	21.9
Male	4.8	9.1	20.7
Female	2.4	6.1	22.4
White, both sexes	3.3	7.1	22.1
Male	4.4	8.9	21.0
Female	2.3	5.9	22.6
Black, both sexes	5.6	10.8	21.9
Male	8.6	13.9	*
Female	3.5	9.0	22.3

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes	292.4	813.6	2,822.1
Male	413.5	1,079.4	3,112.7
Female	192.8	640.3	2,703.8
White, both sexes	283.4	811.2	2,882.1
Male	405.6	1,083.4	3,200.0
Female	180.9	632.0	2,753.0
Black, both sexes	427.8	941.3	2,387.1
Male	563.5	1,175.7	2,446.0
Female	332.3	809.4	2,365.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes	68.3	158.1	534.6
Male	97.2	199.3	550.8
Female	44.6	131.3	528.0
White, both sexes	61.2	149.2	529.5
Male	87.8	187.5	545.6
Female	39.0	123.9	522.9
Black, both sexes	145.5	278.4	646.6
Male	207.3	369.7	701.2
Female	102.0	227.1	626.3

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes	224.1	655.5	2,287.5
Male	316.3	880.1	2,561.9
Female	148.2	509.0	2,175.8
White, both sexes	222.2	662.0	2,352.7
Male	317.8	895.9	2,654.4
Female	141.9	508.1	2,230.1
Black, both sexes	282.3	662.9	1,740.5
Male	356.1	806.0	1,744.8
Female	230.3	582.3	1,738.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION

OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Other heart diseases (I26-I51)

All races 3/, both sexes	134.2	391.3	1,509.4
Male	164.0	451.0	1,549.1
Female	109.6	352.3	1,493.2
White, both sexes	128.1	389.6	1,541.6
Male	157.4	449.5	1,580.5
Female	103.4	350.1	1,525.8
Black, both sexes	220.2	477.9	1,306.0
Male	266.0	558.0	1,402.3
Female	187.9	432.8	1,270.2

Acute and subacute endocarditis (I33)

All races 3/, both sexes	1.2	2.4	2.8
Male	1.5	3.5	4.2
Female	1.0	1.8	2.3
White, both sexes	1.2	2.4	2.7
Male	1.4	3.5	4.2
Female	1.0	1.7	2.1
Black, both sexes	1.9	2.8	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION

OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes	0.7	1.2	1.8
Male	0.8	1.5	2.1
Female	0.6	1.0	1.7
White, both sexes	0.7	1.2	1.8
Male	0.7	1.5	2.1
Female	0.6	1.0	1.6
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Heart failure (I50)

All races 3/, both sexes	34.8	136.1	684.9
Male	41.1	153.7	672.2
Female	29.6	124.6	690.0
White, both sexes	33.4	136.7	703.7
Male	40.0	154.6	690.0
Female	27.9	124.9	709.3
Black, both sexes	55.4	154.4	558.4
Male	61.9	173.5	581.2
Female	50.9	143.7	549.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION

OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes	97.4	251.6	820.0
Male	120.7	292.4	870.7
Female	78.3	225.0	799.3
White, both sexes	92.8	249.3	833.4
Male	115.3	290.0	884.2
Female	73.9	222.6	812.7
Black, both sexes	161.8	319.6	742.0
Male	200.2	379.7	816.5
Female	134.8	285.7	714.3

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes	15.1	45.5	162.9
Male	15.8	45.2	142.4
Female	14.6	45.7	171.3
White, both sexes	11.8	40.4	153.2
Male	12.6	40.3	132.5
Female	11.2	40.4	161.6
Black, both sexes	48.7	107.1	283.4
Male	50.5	109.0	269.9
Female	47.5	106.0	288.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Cerebrovascular diseases (I60-I69)			
All races 3/, both sexes	128.6	461.3	1,589.2
Male	145.0	490.8	1,484.3
Female	115.1	442.1	1,632.0
White, both sexes	119.4	452.4	1,601.6
Male	133.8	480.0	1,490.7
Female	107.3	434.2	1,646.7
Black, both sexes	222.9	588.8	1,530.0
Male	268.5	659.2	1,458.8
Female	190.9	549.2	1,556.5
Atherosclerosis (I70)			
All races 3/, both sexes	7.7	34.6	188.8
Male	9.6	39.7	176.5
Female	6.2	31.3	193.8
White, both sexes	7.7	35.1	195.2
Male	9.7	40.3	181.4
Female	6.1	31.7	200.8
Black, both sexes	8.9	35.2	138.3
Male	11.0	40.3	147.6
Female	7.5	32.4	134.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS

75-84
YEARS

85 YEARS
AND OVER

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes	32.2	74.6	163.2
Male	45.4	102.9	214.9
Female	21.4	56.2	142.2
White, both sexes	32.7	75.1	163.0
Male	46.7	104.7	219.6
Female	21.0	55.6	140.1
Black, both sexes	33.0	78.0	177.7
Male	41.1	95.7	177.6
Female	27.3	68.0	177.7

Aortic aneurysm and dissection (I71)

All races 3/, both sexes	21.8	47.2	79.2
Male	32.9	70.6	127.3
Female	12.7	32.0	59.6
White, both sexes	22.7	48.5	80.8
Male	34.5	73.0	132.8
Female	12.8	32.3	59.6
Black, both sexes	17.0	36.3	62.3
Male	22.1	46.9	69.2
Female	13.4	30.3	59.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84 85 YEARS
YEARS YEARS AND OVER

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes	10.4	27.4	84.0
Male	12.5	32.3	87.6
Female	8.7	24.2	82.6
White, both sexes	10.0	26.6	82.3
Male	12.2	31.6	86.9
Female	8.2	23.3	80.4
Black, both sexes	16.0	41.7	115.4
Male	19.0	48.7	108.4
Female	13.9	37.7	118.0

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes	4.3	9.4	25.6
Male	4.7	9.7	22.2
Female	4.0	9.2	26.9
White, both sexes	4.0	9.1	25.5
Male	4.4	9.5	22.0
Female	3.6	8.9	26.9
Black, both sexes	8.9	14.4	30.0
Male	9.1	13.3	30.0
Female	8.8	15.0	30.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes	39.1	160.3	744.1
Male	48.3	199.8	892.0
Female	31.5	134.6	683.9
White, both sexes	37.1	160.0	758.3
Male	45.1	197.8	906.1
Female	30.3	135.0	698.3
Black, both sexes	60.3	171.1	600.9
Male	83.5	234.1	755.4
Female	43.9	135.7	543.4

Influenza (J10-J11)

All races 3/, both sexes	1.1	4.0	20.2
Male	1.3	4.0	20.7
Female	0.9	4.0	19.9
White, both sexes	1.1	4.2	21.4
Male	1.3	4.2	21.6
Female	0.9	4.2	21.3
Black, both sexes	*	*	9.4
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Pneumonia (J12-J18)			
All races 3/, both sexes	38.0	156.4	723.9
Male	47.1	195.8	871.3
Female	30.6	130.6	663.9
White, both sexes	36.0	155.8	737.0
Male	43.8	193.6	884.4
Female	29.4	130.8	677.1
Black, both sexes	59.3	169.7	591.5
Male	82.8	232.5	742.7
Female	42.7	134.3	535.3
Other acute lower respiratory infections (J20-J22)			
All races 3/, both sexes	0.2	0.7	4.5
Male	0.3	0.7	3.8
Female	0.2	0.6	4.8
White, both sexes	0.3	0.7	4.7
Male	0.3	0.7	3.8
Female	0.2	0.7	5.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84
YEARS 85 YEARS
AND OVER

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes	0.2	0.4	2.6
Male	*	0.6	2.4
Female	*	0.3	2.7
White, both sexes	0.2	0.4	2.7
Male	*	0.6	2.4
Female	*	0.3	2.9
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Unspecified acute lower respiratory infection (J22)

All races 3/, both sexes	*	0.2	1.9
Male	*	*	*
Female	*	0.3	2.1
White, both sexes	*	0.3	2.0
Male	*	*	*
Female	*	0.3	2.2
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes	169.4	386.1	648.6
Male	195.2	488.5	967.9
Female	148.2	319.2	518.5
White, both sexes	178.7	404.4	675.6
Male	201.4	503.6	997.4
Female	159.6	339.1	544.8
Black, both sexes	121.7	234.7	373.8
Male	176.9	370.3	693.1
Female	82.9	158.4	255.0

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes	0.9	2.7	11.8
Male	1.0	3.6	12.9
Female	0.8	2.1	11.4
White, both sexes	0.9	2.8	12.7
Male	1.0	3.7	13.8
Female	0.8	2.2	12.2
Black, both sexes	*	2.2	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Emphysema (J43)			
All races 3/, both sexes	27.8	52.7	63.3
Male	33.0	69.1	104.2
Female	23.6	41.9	46.7
White, both sexes	29.7	55.6	65.8
Male	34.3	72.1	105.5
Female	25.9	44.7	49.7
Black, both sexes	17.0	26.9	37.2
Male	27.4	41.5	87.6
Female	9.7	18.8	18.5
Asthma (J45-J46)			
All races 3/, both sexes	4.0	6.9	16.7
Male	3.0	5.0	12.9
Female	4.8	8.1	18.2
White, both sexes	3.4	6.5	15.9
Male	2.5	4.7	11.4
Female	4.1	7.7	17.7
Black, both sexes	9.2	10.4	22.2
Male	7.9	6.4	25.4
Female	10.2	12.6	21.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Other chronic lower respiratory diseases (J44,J47)

All races 3/, both sexes	136.7	323.8	556.7
Male	158.3	410.9	838.0
Female	119.0	267.1	442.2
White, both sexes	144.7	339.5	581.2
Male	163.6	423.0	866.7
Female	128.9	284.4	465.2
Black, both sexes	94.5	195.2	310.6
Male	140.8	320.4	576.6
Female	62.0	124.8	211.6

Pneumoconioses and chemical effects (J60-J66,J68)

All races 3/, both sexes	1.5	3.9	7.5
Male	3.2	9.6	24.9
Female	*	*	*
White, both sexes	1.7	4.1	7.9
Male	3.4	10.0	26.4
Female	*	*	*
Black, both sexes	*	2.5	*
Male	*	6.4	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS 75-84 YEARS 85 YEARS AND OVER

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes	10.3	44.5	187.6
Male	14.4	62.3	276.0
Female	7.0	33.0	151.6
White, both sexes	9.8	44.4	191.0
Male	13.4	62.1	284.0
Female	6.8	32.8	153.2
Black, both sexes	17.5	51.5	159.8
Male	27.8	74.5	209.9
Female	10.3	38.5	141.2

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes	29.6	72.0	141.7
Male	35.7	90.5	179.7
Female	24.7	59.9	126.2
White, both sexes	30.0	73.7	145.9
Male	36.1	92.5	184.4
Female	25.0	61.4	130.3
Black, both sexes	30.5	58.0	98.2
Male	36.0	73.0	132.6
Female	26.6	49.6	85.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Peptic ulcer (K25-K28)

All races 3/, both sexes	4.1	11.4	35.6
Male	5.3	13.7	40.7
Female	3.0	10.0	33.5
White, both sexes	4.0	11.5	36.3
Male	5.2	13.5	40.7
Female	3.0	10.2	34.5
Black, both sexes	4.8	10.0	26.6
Male	7.0	13.9	31.1
Female	3.3	7.8	24.9

Diseases of appendix (K35-K38)

All races 3/, both sexes	0.4	1.0	1.6
Male	0.5	1.2	2.0
Female	0.3	0.8	1.5
White, both sexes	0.3	0.9	1.7
Male	0.4	1.2	2.0
Female	0.3	0.7	1.6
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Hernia (K40-K46)

All races 3/, both sexes	1.1	3.8	13.1
Male	1.1	4.0	12.6
Female	1.2	3.6	13.3
White, both sexes	1.1	3.8	13.3
Male	1.1	4.1	12.7
Female	1.2	3.6	13.6
Black, both sexes	1.3	4.1	12.5
Male	*	*	*
Female	*	4.4	12.4

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes	29.8	31.0	23.1
Male	39.9	38.4	34.3
Female	21.5	26.1	18.5
White, both sexes	30.6	32.3	23.9
Male	40.9	39.9	35.5
Female	21.9	27.2	19.2
Black, both sexes	26.2	18.0	13.8
Male	37.7	24.5	*
Female	18.1	14.3	11.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Alcoholic liver disease (K70)

All races 3/, both sexes	10.3	6.1	2.2
Male	17.3	11.5	5.3
Female	4.5	2.5	1.0
White, both sexes	10.5	6.1	2.3
Male	17.5	11.7	5.4
Female	4.6	2.4	1.0
Black, both sexes	10.3	5.5	*
Male	18.1	10.0	*
Female	4.7	*	*

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes	19.5	24.9	20.9
Male	22.7	26.9	29.0
Female	16.9	23.6	17.5
White, both sexes	20.1	26.1	21.6
Male	23.4	28.2	30.1
Female	17.3	24.8	18.2
Black, both sexes	16.0	12.5	12.2
Male	19.6	14.5	*
Female	13.4	11.4	11.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes	2.4	7.5	24.8
Male	2.9	9.4	27.6
Female	2.1	6.2	23.7
White, both sexes	2.3	7.4	25.6
Male	2.8	9.6	28.9
Female	1.8	6.0	24.3
Black, both sexes	3.9	8.3	17.8
Male	3.7	8.2	*
Female	4.1	8.4	19.3

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes	38.0	100.8	277.8
Male	42.9	127.3	380.7
Female	34.0	83.5	235.9
White, both sexes	31.9	94.0	268.7
Male	37.7	121.1	376.2
Female	27.1	76.2	225.0
Black, both sexes	99.9	195.1	406.4
Male	104.8	231.6	490.1
Female	96.5	174.6	375.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes	0.2	0.4	1.2
Male	*	0.5	1.8
Female	*	0.3	0.9
White, both sexes	0.1	0.4	1.1
Male	*	0.5	*
Female	*	0.3	1.0
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07,N26)

All races 3/, both sexes	0.6	1.4	2.8
Male	0.7	1.7	4.5
Female	0.5	1.2	2.1
White, both sexes	0.5	1.3	2.4
Male	0.6	1.7	4.0
Female	0.5	1.1	1.8
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Renal failure (N17-N19)

All races 3/, both sexes	37.2	98.9	273.7
Male	42.1	125.1	374.2
Female	33.3	81.8	232.8
White, both sexes	31.2	92.2	265.0
Male	36.9	118.9	370.4
Female	26.5	74.6	222.1
Black, both sexes	98.6	192.4	400.4
Male	103.3	229.2	480.9
Female	95.3	171.7	370.4

Other disorders of kidney (N25,N27)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes	0.6	2.0	7.5
Male	0.3	1.5	5.6
Female	0.8	2.3	8.3
White, both sexes	0.5	1.9	7.6
Male	*	1.5	5.9
Female	0.7	2.2	8.3
Black, both sexes	*	3.1	7.8
Male	*	*	*
Female	*	3.6	9.0

Hyperplasia of prostate (N40)

All races 3/, both sexes	0.2	1.3	5.2
Male	0.5	3.3	18.1
White, both sexes	0.2	1.3	5.2
Male	0.5	3.3	18.1
Black, both sexes	*	*	*
Male	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
---	----------------	----------------	----------------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races 3/, both sexes	0.1	0.2	0.6
Female	0.2	0.4	0.9
White, both sexes	*	0.2	0.6
Female	*	0.4	0.9
Black, both sexes	*	*	*
Female	*	*	*

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
Pregnancy with abortive outcome (000-007)			
All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female
Other complications of pregnancy, childbirth and the puerperium (010-099)			
All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	2.5	3.9	7.5
Male	2.7	4.0	10.6
Female	2.4	3.9	6.3
White, both sexes	2.5	4.1	7.8
Male	2.6	4.3	11.1
Female	2.5	4.0	6.5
Black, both sexes	2.6	2.7	*
Male	3.8	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes	16.5	40.7	208.1
Male	20.6	46.0	182.7
Female	13.2	37.2	218.5
White, both sexes	15.6	40.2	212.2
Male	19.2	45.5	186.8
Female	12.7	36.7	222.5
Black, both sexes	28.9	55.5	186.4
Male	39.9	65.1	161.5
Female	21.2	50.1	195.7

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes	136.0	388.5	1,445.7
Male	148.4	391.5	1,266.2
Female	125.8	386.5	1,518.8
White, both sexes	132.7	386.9	1,464.6
Male	143.4	387.1	1,274.8
Female	123.6	386.7	1,541.7
Black, both sexes	191.9	467.7	1,388.9
Male	228.4	523.5	1,325.1
Female	166.3	436.2	1,412.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes	41.9	95.1	273.5
Male	55.9	124.3	359.7
Female	30.3	76.1	238.4
White, both sexes	41.1	96.5	282.8
Male	54.2	124.3	369.8
Female	30.1	78.1	247.4
Black, both sexes	52.5	84.9	181.1
Male	79.4	132.6	275.6
Female	33.5	58.0	145.9

Transport accidents (V01-V99,Y85)

All races 3/, both sexes	18.0	27.0	31.5
Male	24.3	38.1	59.8
Female	12.8	19.8	19.9
White, both sexes	17.9	27.5	32.2
Male	23.7	38.1	61.6
Female	12.9	20.5	20.3
Black, both sexes	19.6	20.0	21.6
Male	32.2	36.3	45.0
Female	10.8	10.7	12.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes	16.5	25.7	30.4
Male	21.7	35.6	57.5
Female	12.3	19.2	19.3
White, both sexes	16.4	26.1	31.1
Male	21.1	35.6	59.2
Female	12.4	19.8	19.7
Black, both sexes	18.2	18.9	20.3
Male	29.4	34.5	43.8
Female	10.3	10.1	11.6

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes	0.5	0.7	0.8
Male	0.9	1.2	*
Female	0.2	0.5	*
White, both sexes	0.5	0.7	0.8
Male	0.8	1.2	*
Female	0.3	0.4	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes	1.0	0.6	*
Male	1.7	1.3	*
Female	0.3	*	*
White, both sexes	1.0	0.7	*
Male	1.8	1.3	*
Female	0.3	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes	23.8	68.1	242.0
Male	31.5	86.2	299.8
Female	17.5	56.3	218.4
White, both sexes	23.2	69.0	250.6
Male	30.4	86.2	308.2
Female	17.2	57.7	227.2
Black, both sexes	32.8	64.9	159.5
Male	47.3	96.3	230.6
Female	22.7	47.2	133.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Falls (W00-W19)

All races 3/, both sexes	9.0	31.1	112.6
Male	11.9	39.9	145.4
Female	6.6	25.3	99.2
White, both sexes	9.2	32.6	119.1
Male	12.1	41.4	152.8
Female	6.7	26.8	105.4
Black, both sexes	8.3	14.9	38.8
Male	11.4	21.2	61.1
Female	6.1	11.4	30.5

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes	0.2	0.2	*
Male	0.4	0.5	*
Female	*	*	*
White, both sexes	0.2	0.2	*
Male	0.4	0.5	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74
YEARS 75-84
YEARS 85 YEARS
AND OVER

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes	1.0	1.3	1.5
Male	1.6	1.8	2.8
Female	0.5	0.9	1.0
White, both sexes	1.0	1.2	1.5
Male	1.6	1.8	2.7
Female	0.5	0.8	1.0
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes	2.2	3.8	6.2
Male	2.8	5.0	9.4
Female	1.7	3.0	4.9
White, both sexes	1.8	3.1	5.1
Male	2.3	4.1	8.2
Female	1.5	2.4	3.9
Black, both sexes	5.9	13.2	20.3
Male	8.8	18.5	27.7
Female	3.9	10.2	17.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

65-74 YEARS 75-84 YEARS 85 YEARS AND OVER

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes	1.5	2.0	4.3
Male	1.8	2.1	6.2
Female	1.3	1.9	3.6
White, both sexes	1.4	2.1	4.1
Male	1.6	2.1	5.9
Female	1.3	2.0	3.4
Black, both sexes	2.6	*	8.1
Male	4.5	*	*
Female	*	*	*

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes	9.9	29.8	117.1
Male	13.0	36.9	135.2
Female	7.4	25.1	109.8
White, both sexes	9.5	29.8	120.5
Male	12.5	36.2	137.6
Female	7.1	25.6	113.6
Black, both sexes	14.9	33.7	90.1
Male	20.3	53.0	124.5
Female	11.1	22.8	77.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Intentional self-harm (suicide) (X60-X84,Y87.0)

All races 3/, both sexes	12.5	17.6	19.6
Male	22.7	38.6	57.5
Female	4.0	4.0	4.2
White, both sexes	13.5	18.8	21.0
Male	24.3	41.1	61.6
Female	4.4	4.1	4.5
Black, both sexes	5.1	5.8	*
Male	11.1	12.1	*
Female	*	*	*

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes	9.2	13.0	13.2
Male	18.4	30.8	42.9
Female	1.7	1.4	1.1
White, both sexes	10.1	14.1	14.3
Male	19.8	33.3	46.5
Female	1.9	1.5	1.2
Black, both sexes	3.7	3.5	*
Male	8.8	8.2	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, AND SEX

65-74 YEARS 75-84 YEARS 85 YEARS AND OVER

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races 3/, both sexes	3.2	4.6	6.5
Male	4.3	7.8	14.6
Female	2.3	2.6	3.2
White, both sexes	3.4	4.6	6.7
Male	4.5	7.8	15.2
Female	2.5	2.6	3.3
Black, both sexes	1.4	2.3	*
Male	*	*	*
Female	*	*	*

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	2.4	2.4	2.4
Male	3.4	3.2	3.3
Female	1.6	2.0	2.0
White, both sexes	1.9	2.0	2.2
Male	2.4	2.5	2.8
Female	1.5	1.8	1.9
Black, both sexes	7.6	7.0	*
Male	13.9	11.8	*
Female	3.1	4.3	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (X93-X95)

All races 3/, both sexes	1.0	0.7	0.7
Male	1.3	0.8	*
Female	0.8	0.6	*
White, both sexes	0.8	0.6	0.6
Male	1.0	0.6	*
Female	0.7	0.6	*
Black, both sexes	2.7	*	*
Male	5.4	*	*
Female	*	*	*

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	1.4	1.8	1.7
Male	2.1	2.4	2.5
Female	0.9	1.4	1.4
White, both sexes	1.0	1.5	1.5
Male	1.4	1.9	2.1
Female	0.7	1.2	1.3
Black, both sexes	4.8	5.1	*
Male	8.5	8.2	*
Female	2.3	3.4	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Legal intervention (Y35,Y89.0)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes	0.6	0.6	1.3
Male	0.8	1.0	1.7
Female	0.3	0.4	1.2
White, both sexes	0.5	0.6	1.3
Male	0.7	1.0	*
Female	0.4	0.4	1.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes	0.5	0.6	1.3
Male	0.7	0.9	*
Female	0.3	0.4	1.2
White, both sexes	0.5	0.6	1.2
Male	0.6	0.9	*
Female	0.3	0.4	1.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

RACE, AND SEX

	65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------------

Operations of war and their sequelae (Y36,Y89.1)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes	3.5	6.7	13.6
Male	4.0	8.2	16.0
Female	3.1	5.7	12.7
White, both sexes	3.3	6.4	13.1
Male	3.7	8.2	15.8
Female	3.0	5.3	12.0
Black, both sexes	6.1	10.5	21.0
Male	7.6	10.3	*
Female	5.0	10.6	21.9

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.