

Provisional COVID-19 Deaths Among Ages 0-18 Years

Data as of 7/13/2020

Data are reported for deaths that occurred from the week ending February 1, 2020 through July 11, 2020. Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. COVID-19 deaths are those with confirmed or presumed COVID-19, coded to ICD-10 code U07.1.

Table 1. Provisional COVID-19 deaths by age group: United States, Week ending 2/1/2020 to 7/11/2020. (n = 120,373)

Age Category	Number of COVID-19 deaths	Percentage of COVID-19 deaths
0-18 years	52	0.0%
19-44 years	3,134	2.6%
45-64 years	20,507	17.0%
65-74 years	25,040	20.8%
> 75 years	71,639	59.5%

Table 2. Provisional COVID-19 deaths among ages 0-18 years, by sex: United States, Week ending 2/1/2020 to 7/11/2020. (n=52)

Sex	Number of COVID-19 deaths
Male	27
Female	25

Table 3. Provisional COVID-19 deaths among ages 0-18 years, by race and Hispanic origin: United States, Week ending 2/1/2020 to 7/11/2020. (n=52)

Race/Hispanic Origin	Number of COVID-19 deaths	Percentage of COVID-19 deaths
Hispanic or Latino	15	29%
Non-Hispanic American Indian or Alaska Native	2	4%
Non-Hispanic Asian	1	2%
Non-Hispanic Black	17	33%
Non-Hispanic More than one race	1	2%
Non-Hispanic Native Hawaiian or Other Pacific Islander	1	2%
Non-Hispanic White	13	25%
Unknown	2	4%

Source: NCHS. National Vital Statistics System. Based on provisional data.

Table 4. Provisional COVID-19 deaths mentioning specific conditions among ages 0-18 years, by condition: United States, Week ending 2/1/2020 to 7/11/2020.

Condition mentioned on death certificate	Number of COVID-19 deaths mentioning condition	Percentage of COVID-19 deaths
Chronic lower respiratory diseases	4	7.7%
Diabetes	4	7.7%
Heart failure	2	3.8%
Hypertensive diseases	2	3.8%
Influenza and pneumonia	11	21.2%
Malignant neoplasms	6	11.5%
Obesity	7	13.5%
Renal failure	2	3.8%
Respiratory distress syndrome	9	17.3%
Respiratory failure	9	17.3%
Sepsis	4	7.7%

Source: NCHS. National Vital Statistics System. Based on provisional data.