

Table 1. Number of pregnancies for teenagers 15-19 years by outcome of pregnancy, by age: United States, 1990, 1995, 2000, and 2002
 [Numbers in thousands]

Year and pregnancy outcome	15-19 years			15-17 years			18-19 years					
	All races ¹	Non-Hispanic white	Non-Hispanic black	Hispanic ²	All races ¹	Non-Hispanic white	Non-Hispanic black	Hispanic ²	All races ¹	Non-Hispanic white	Non-Hispanic black	Hispanic ²
All pregnancies ³												
2002	757	311	207	207	265	96	80	78	492	215	127	129
2000	834	357	234	211	297	113	91	81	537	245	142	129
1995	903	421	257	195	365	156	116	82	538	265	142	112
1990	1017	527	301	164	377	177	126	64	641	350	174	100
Live births ⁴												
2002	425	181	102	128	139	50	37	47	287	131	65	81
2000	469	207	116	129	157	60	44	48	312	148	73	81
1995	500	234	132	118	193	80	58	49	307	154	74	70
1990	522	259	150	98	183	79	62	37	338	180	88	61
Induced abortions ⁵												
2002	215	81	77	44	78	28	30	15	137	53	47	28
2000	235	94	84	45	85	32	33	16	151	62	52	29
1995	263	122	88	43	106	49	37	17	157	74	51	26
1990	351	198	108	38	130	71	42	14	221	126	65	24
Fetal losses ⁶												
2002	117	49	28	36	48	17	13	16	69	31	15	19
2000	129	56	33	36	55	21	15	17	75	35	17	19
1995	140	65	38	34	67	28	20	17	74	37	18	17
1990	145	70	43	27	64	27	22	13	81	43	21	15

¹Includes races other than white and black and origin not stated

²Includes all persons of Hispanic origin of any race

³Pregnancy estimates are sums of live births, induced abortions, and fetal losses

⁴Live births are compiled from birth data collected from all birth certificates through the National Vital Statistics System; see references 5 and 6

⁵Induced abortion estimates are based on the age distribution of abortions collected by the Centers for Disease Control and Prevention's National Center for Chronic Disease Prevention and Health Promotion, which are adjusted to national totals compiled by The Guttmacher Institute from their surveys of all known abortion providers; see references 7-9.

⁶Spontaneous fetal losses from recognized pregnancies of all gestational periods as estimated from reports from women in the 1982, 1988, 1995, and 2002 National Surveys of Family Growth conducted by the National Center for Health Statistics, CDC. The rate of fetal loss depends on the degree to which losses at very early gestations are detected.

NOTES: Pregnancy and fetal loss estimates for 1990-2000 have been revised and may differ from figures previously published.

Due to sample size limitations in any given year, fetal loss proportions for teenagers for 1990-2002 are based on cycles 3 through 6 of the NSFG (conducted 1982, 1988, 1995, and 2002). Due to rounding, figures may not add to totals.

Table 2. Pregnancy, live birth, induced abortion, and fetal loss rates for teenagers 15-19 years, by age, race, and Hispanic origin: United States, 1990-2002

Year and pregnancy outcome	15- 19 years				15- 17 years				18- 19 years			
	All races ¹	Non-Hispanic White	Non-Hispanic Black	Hispanic ²	All races ¹	Non-Hispanic White	Non-Hispanic Black	Hispanic ²	All races ¹	Non-Hispanic White	Non-Hispanic Black	Hispanic ²
All pregnancies ³												
2002	76.4	49.0	138.9	135.2	44.4	25.1	88.4	85.1	125.0	85.3	217.0	210.9
2001	80.6	52.3	147.7	139.4	46.9	26.9	94.2	88.3	130.9	90.5	228.8	214.1
2000	84.8	56.3	158.8	142.1	50.8	29.8	104.8	93.3	134.5	95.4	237.8	211.6
1999	86.9	59.0	161.9	143.2	53.1	32.4	106.4	97.0	136.6	98.2	242.9	209.6
1998	90.1	61.8	168.4	146.3	56.7	35.4	114.5	100.3	140.3	101.8	247.8	214.5
1997	92.7	64.5	174.7	147.7	59.5	37.9	121.3	102.6	144.3	106.1	257.3	215.5
1996	97.0	67.2	182.2	157.1	63.4	40.3	129.7	109.4	149.0	109.1	264.0	228.4
1995	101.1	70.6	189.6	163.3	67.4	43.0	139.4	115.5	153.4	113.7	268.4	234.2
1994	106.1	73.4	207.3	169.0	71.1	44.7	152.9	120.3	159.6	117.4	292.2	240.7
1993	109.4	75.9	219.0	170.3	72.7	46.0	161.9	117.8	164.1	120.8	305.1	247.9
1992	112.3	78.3	226.0	174.0	73.5	46.6	165.2	119.7	169.3	125.3	314.8	253.9
1991	116.4	83.8	231.8	173.7	76.1	50.1	170.5	117.4	172.1	130.4	316.0	254.3
1990	116.8	86.8	232.7	167.4	77.1	52.5	172.0	113.0	167.7	129.8	312.6	242.4
Live births ⁴												
2002	43.0	28.5	68.3	83.4	23.2	13.1	41.0	50.7	72.8	51.9	110.3	133.0
2001	45.3	30.3	73.5	86.4	24.7	14.0	44.9	52.8	76.1	54.8	116.7	135.5
2000	47.7	32.6	79.2	87.3	26.9	15.8	50.1	55.5	78.1	57.5	121.9	132.6
1999	48.8	34.1	81.0	86.8	28.2	17.1	51.7	56.9	79.1	59.4	123.9	129.5
1998	50.3	35.3	85.7	87.9	29.9	18.3	56.8	58.5	80.9	60.9	128.2	131.5
1997	51.3	36.0	88.3	89.6	31.4	19.3	60.7	61.1	82.1	62.1	131.0	132.4
1996	53.5	37.6	91.9	94.6	33.3	20.6	64.8	64.2	84.7	64.0	134.1	140.0
1995	56.0	39.3	97.2	99.3	35.5	22.0	70.4	68.3	87.7	66.2	139.2	145.4
1994	58.2	40.4	105.7	101.3	37.2	22.7	77.0	69.9	90.2	67.6	150.4	147.5
1993	59.0	40.7	110.5	101.8	37.5	22.7	81.1	68.5	91.1	67.7	154.6	151.1
1992	60.3	41.6	114.7	103.3	37.6	22.7	82.9	68.9	93.6	69.7	161.0	153.8
1991	61.8	43.3	118.2	104.6	38.6	23.6	86.1	69.2	94.0	70.5	162.2	155.4
1990	59.9	42.6	116.2	100.2	37.5	23.3	84.9	65.8	88.6	66.9	157.5	147.6
Induced abortions ⁵												
2002	21.7	12.8	51.6	28.5	13.1	7.4	33.1	16.8	34.8	21.0	80.2	46.2
2001	22.8	13.9	53.7	28.9	13.6	8.0	33.7	17.1	36.6	22.6	84.1	46.2
2000	24.0	14.8	57.4	30.3	14.5	8.5	37.3	18.4	37.7	24.1	86.8	47.3
1999	24.7	15.5	58.1	32.1	15.2	9.4	36.7	20.2	38.6	24.6	89.4	49.1
1998	25.8	16.9	58.6	33.7	16.4	10.7	38.0	21.5	40.0	26.3	89.0	51.6
1997	27.1	18.6	61.3	32.7	17.2	11.8	39.5	20.3	42.6	29.1	95.0	51.4
1996	28.6	19.3	64.1	35.7	18.6	12.6	42.4	22.9	44.0	29.8	97.9	54.9
1995	29.4	20.5	64.6	35.8	19.5	13.4	44.6	23.5	44.8	31.6	96.0	54.0
1994	31.6	21.8	71.2	39.0	21.0	14.1	49.1	26.1	47.8	33.7	105.8	58.0
1993	33.9	24.0	76.9	39.7	22.2	15.4	52.6	25.5	51.2	36.9	113.5	60.7
1992	35.2	25.3	78.6	41.6	22.9	16.0	53.5	26.8	53.3	38.9	115.2	63.3
1991	37.4	28.7	80.0	39.6	24.2	18.3	54.5	24.2	55.7	43.0	115.0	61.7
1990	40.3	32.5	83.5	39.1	26.5	21.1	57.7	24.3	57.9	46.8	117.4	59.5
Fetal losses ⁶												
2002	11.8	7.7	19.0	23.3	8.1	4.6	14.2	17.6	17.4	12.4	26.4	31.8
2001	12.4	8.2	20.5	24.1	8.6	4.9	15.6	18.3	18.2	13.1	27.9	32.4
2000	13.1	8.8	22.2	24.4	9.4	5.5	17.4	19.3	18.7	13.8	29.1	31.7
1999	13.5	9.3	22.7	24.4	9.8	5.9	18.0	19.8	18.9	14.2	29.6	31.0
1998	14.0	9.6	24.1	24.8	10.4	6.4	19.7	20.3	19.4	14.6	30.7	31.4
1997	14.3	9.9	25.1	25.4	10.9	6.7	21.1	21.2	19.6	14.9	31.3	31.7
1996	15.0	10.3	26.2	26.8	11.6	7.2	22.5	22.3	20.3	15.3	32.1	33.5
1995	15.7	10.8	27.9	28.2	12.3	7.6	24.4	23.7	21.0	15.8	33.3	34.8
1994	16.3	11.2	30.4	28.7	12.9	7.9	26.8	24.3	21.6	16.2	36.0	35.3
1993	16.5	11.2	31.7	28.8	13.0	7.9	28.2	23.8	21.8	16.2	37.0	36.1
1992	16.8	11.4	32.7	29.1	13.0	7.9	28.8	23.9	22.4	16.7	38.5	36.8
1991	17.2	11.8	33.7	29.4	13.4	8.2	29.9	24.0	22.5	16.9	38.8	37.2
1990	16.6	11.6	33.0	28.1	13.0	8.1	29.5	22.9	21.2	16.0	37.7	35.3

¹ Includes races other than white and black and origin not stated.² Includes all persons of Hispanic origin of any race.³ Pregnancy rates are estimated by summing rates of live births, induced abortions, and fetal losses.⁴ Live birth rates are computed from birth data collected from all birth certificates through the National Vital Statistics System; see references 5 and 6.⁵ Induced abortion rates are based on the age distribution of abortions collected by CDC's National Center for Chronic Disease Prevention and Health Promotion, which are adjusted to national totals compiled by The Guttmacher Institute from their surveys of all known abortion providers; see references 7-9.⁶ Spontaneous fetal losses from recognized pregnancies of all gestational periods as estimated from reports from women in the 1982, 1988, 1995 and 2002 National Surveys of Family Growth conducted by the National Center for Health Statistics. The rate of fetal loss depends on the degree to which losses at very early gestations are detected; see references 1, 9, and 10.

NOTES: Pregnancy and fetal loss rates for 1990-2000 have been revised and may differ from rates previously published. Due to sample size limitations in any given year, fetal loss proportions for teenagers for 1990-2002 are based on cycles 3 through 6 of the NSFG (conducted 1982, 1988, 1995, and 2002).