


Diabetes-related death rates, 1986-96


[Objective 17.9]



NOTE: Data include deaths from diabetes as an underlying or contributing cause.

SOURCE: CDC/NCHS, National Vital Statistics System, 1986-96


Proportion of persons 18 years and over who are diagnosed with diabetes, 1997 [Objective 17.11]



SOURCE: CDC/NCCDPHP, Behavioral Risk Factor Surveillance System, 1997


Diabetes prevalence, 1988-96

[Objective 17.11]



NOTE: Data for Total and Black are 3-year moving averages plotted on the last year.
SOURCE: CDC/NCHS, National Health Interview Survey, 1988-96. For American Indian/Alaska Native data: Ambulatory Utilization Data, Indian Health Service.


Diabetes prevalence: 1994-96



¹Hispanic origin can be of any race.

SOURCE: CDC/NCHS, National Health Interview Survey, 1994-96.

Diagnosed and undiagnosed diabetes prevalence, 1976-80 and 1988-94




NOTE: Data are for persons 40-74 years.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 1976-80 and 1988-94

Overweight (measured) prevalence, 1976-80 and 1988-94⁶

[Objective 17.12]




NOTE: All data are for persons 20-74 years, except for adolescents (12-19 years). Overweight is defined as a Body Mass Index (weight in kilograms divided by the square of height in meters) of at least 27.8 for men and 27.3 for women.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 1976-80 and 1988-94


Lower extremity amputations due to diabetes, 1987-96

[Objective 17.10]




SOURCE: CDC/NCHS, National Health Interview Survey; National Hospital Discharge Survey, 1987-96

ESRD treatment by primary cause, 1987-96



NOTE: Of those alive on December 31 of each year.
SOURCE: United States Renal Data System, 1987-96


Proportion of people with diabetes 18 years and over who had a dilated eye exam in the past year, 1995-97 [Objective 17.23]



NOTE: Data shown represent two or three year averages, depending on the State. Data are not available for States that have less than 2 years of data. SOURCE: CDC/NCCDPHP, Behavioral Risk Factor Surveillance System, 1995-97

Diabetes/asthma education, 1983-93


[Objective 17.14]



SOURCE: CDC/NCHS, National Health Interview Survey, 1983-93

Emergency room visits for asthma, 1992-97


[HP2010 objective 24.3]



NOTE: Rate is based on the number of visits that had asthma as the first-listed diagnosis.
SOURCE: CDC/NCHS, National Hospital Ambulatory Medical Care Survey, 1992-97


Asthma hospitalizations, 1987-96

[Objective 11.1]



SOURCE: CDC/NCHS, National Hospital Discharge Survey, 1987-96

Limitation of activity due to asthma, 1988-96 [Objective 17.4]



NOTE: Data are 3-year moving averages plotted on the last year.

*Data are unreliable due to small number of events.

SOURCE: CDC/NCHS, National Health Interview Survey, 1988-96