

Suicide rates [Objective 6.1]




NOTE: Rates for all persons and American Indians/Alaska Native males are age-adjusted. SOURCE: CDC/NCHS, National Vital Statistics System, 1987-97

Suicide among persons 65 years and over




¹ Hispanic origin can be of any race. SOURCE: CDC/NCHS, National Vital Statistics Surveillance System, 1996

Adolescent injurious suicide attempts (Objective 6.2)




SOURCE: CDC/DASH, Youth Risk Behavior Survey, 1990-97

Suicide attempts and suicides among adolescents




SOURCE: CDC/DASH, Youth Risk Behavior Survey, 1991, 1993, 1995, 1997; CDC/NCHS, National Vital Statistics System, 1991, 1993, 1995, 1997

Prevalence of mental disorders (Objective 6.4)




SOURCE: NIH/NIMH, Epidemiologic Catchment Area Study, 1981-85; NIH/NIMH, National Comorbidity Survey, 1990-92

Persons with adverse effects from stress (Objective 6.5)




SOURCE: CDC/NCHS, National Health Interview Survey, 1985, 1990, 1993. and 1995

Use of community support by persons with mental illness (Objective 6.6)




SOURCE: NIH/NIMH, Community Support Program Followup, 1986; CDC/NCHS, National Health Interview Survey, 1994

Treatment for depression (Objective 6.7)




SOURCE: NIH/NIMH, Epidemiologic Catchment Area Study, 1981-85; NIH/NIMH, National Comorbidity Survey, 1990-92

Persons seeking help for emotional problems (Objective 6.8)



SOURCE: CDC/NCHS, National Health Interview Survey, 1985, 1990, 1993. and 1995

Prevalence of depression (Objective 6.15)



SOURCE: NIH/NIMH, Epidemiologic Catchment Area Study, 1981-85; NIH/NIMH, National Comorbidity Survey, 1990-92