| | The state of the s | Baseline | | | | | | | | | | Target |
|-------|--|----------|----------|---------|-------|----------|------------|----------|-------------|----------|-------------|--------|
| | Objective | year | Baseline | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 2000 |
| 10.1 | Work-related injury deaths (per 100,000 full-time workers) | 1983-87 | 6.0 | 4.3 | 4.3 | 5.0 | 5.0 | 5.0 | 4.9 | 4.8 | 1/4.5 | 4.0 |
| | a. Mine workers | 1983-87 | 30.3 | 2/17.3 | 15.6 | 27.0 | 26.0 | 27.0 | 25.0 | 26.8 | 1/23.6 | 21 |
| | b. Construction workers | 1983-87 | 25.0 | 20.6 | 16.6 | 14.0 | 14.0 | 15.0 | 14.6 | 13.9 | 1/14.6 | 17 |
| | c. Transportation workers | 1983-87 | 15.2 | 10.0 | 8.1 | 13.0 | 13.0 | 14.0 | 13.1 | 13.1 | 1/11.8 | 10 |
| | d. Farm workers | 1983-87 | 14.0 | 23.8 | | 24.0 | 26.0 | 24.0 | 22.5 | 22.8 | 1/24.1 | 9.5 |
| | u. Tailii workers | 1303-07 | 14.0 | 25.0 | | 24.0 | 20.0 | 24.0 | 22.0 | 22.0 | 1/24.1 | 3.5 |
| 10.2 | Nonfatal work-related injuries (per 100 full-time workers)/3 | 1983-87 | 7.7 | 8.3 | 7.9 | 8.3 | 7.9 | 7.7 | 7.5 | 6.9 | 6.6 | 6 |
| | a. Construction workers/3 | 1983-87 | 14.9 | 14.1 | 12.8 | 12.9 | 12.0 | 11.5 | 10.4 | 9.7 | 9.3 | 10 |
| | b. Nursing and personal care workers/3 | 1983-87 | 12.7 | 15.4 | 15.0 | 18.2 | 16.9 | 16.5 | 17.8 | 16.2 | 15.9 | 9 |
| | c. Farm workers/3 | 1983-87 | 12.4 | 12.3 | 11.1 | 11.5 | 10.9 | 9.7 | 9.9 | 8.9 | 8.7 | 8 |
| | d. Transportation workers/3 | 1983-87 | 8.3 | 9.4 | 9.1 | 8.8 | 9.1 | 9.0 | 8.7 | 8.4 | 7.9 | 6 |
| | e. Mine workers/3 | 1983-87 | 8.3 | 8.1 | 7.1 | 7.0 | 6.5 | 6.0 | 6.0 | 5.3 | 5.7 | 6 |
| | f. Adolescent workers (15-17 years) | 1992 | 4/5.8 | | | | | | | 5/4.8 | | 3.8 |
| | in Audiococia monicio (10 11 years) | .002 | ., 0.0 | | | | | | | 0, | | 0.0 |
| 10.3 | Cumulative trauma disorders (per 100,000 full-time workers) | 1987 | 100 | 241 | 297 | 368 | 383 | 411 | 378 | 335 | 320 | 60 |
| | Manufacturing industry workers | 1987 | 355 | 867 | 1,046 | 1,241 | 1,267 | 1,362 | 1,258 | 1,104 | 1,061 | 150 |
| | b. Meat product workers | 1987 | 3,920 | 8,245 | 8,802 | 8,475 | 8,532 | 8,750 | 7,720 | 6,116 | 6,860 | 2,000 |
| 10.4 | Occupational skin disorders (per 100,000 full-time workers) | 1983-87 | 64 | 79 | 77 | 82 | 76 | 81 | 79 | 69 | 67 | 55 |
| 10.5* | Hepatitis B infections among occupationally exposed workers (number of cases) | 1987 | 3,090 | 1,258 | 2,576 | 1,923 | 727 | 506 | 407 | 391 | | 623 |
| 10.6 | Worksite occupant protection system mandates | 1992 | 82.4% | | | | | | 85% | | 6/47% | 95% |
| 10.7 | Occupational noise exposure/7 (average noise levels exceeding 85 db) | 1989 | 16% | 20.5% | 23.8% | 21.5% | 19.9% | | | | | 15% |
| 10.8 | Occupational lead exposure (blood concentration greater than 25 ug/dL) | 1988 | 8/4,804 | 9/4,531 | 1 | 10/8,886 | 11/9,57112 | 2/11,068 | 12/11,660 1 | 3/12,706 | 1,14/10,501 | 0 |
| 10.9* | Hepatitis B immunizations among occupationally exposed workers/15 | 1989 | 37% | | | 50% | | 67% | | | | 90% |
| 10.10 | Number of States with occupational health and safety plans | 1989 | 10 | | | 23 | | | | | 6/23 | 50 |
| 10.11 | Number of States with occupational lung disease exposure standards | 1969 | 16/50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 6/50 | 50 |
| 10.12 | Worksite health and safety programs | 1992 | 63.8% | | | | | | | | | 70% |
| 10.13 | Worksite back injury prevention and rehabilitation programs | 1985 | 28.6% | | | 32.5% | | | | | 6/53% | 50% |
| | Back injury classes, workshops or lectures | | | | | 24% | | | 26% | | | |
| 10.14 | Number of States with programs for small business safety and health | 1991 | 26 | | | | 50 | 50 | 50 | 50 | 50 | 50 |
| 10.15 | Clinician assessment of occupational health exposures Percent of clinicians routinely providing service to 81-100% of patients | | | | | | | | | | | 75% |
| | Inquiry about work-related health risks (16 years and over) Pediatricians | 1992 | 7% | | | | | | | | 17,18/ | 75% |

| | · · · · · · · · · · · · · · · · · · · | Baseline | | | | | | | | | | Target |
|--------|--|----------|----------|-------|-------|-------|-------|-------|-------|-------|---------|--------|
| | Objective | year | Baseline | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 2000 |
| | Nurse practitioners | 1992 | 14% | | | | | | | | 17/14% | 75% |
| | Obstetricians/gynecologists | 1992 | 6% | | | | | | | | 17,18/ | 75% |
| | Internists | 1992 | 14% | | | | | | | | 17,18/ | 75% |
| | Family physicians | 1992 | 7% | | | | | | | | 17,18/ | 75% |
| | Counseling about work-related health risks | | | | | | | | | | | |
| | Pediatricians | 1992 | 8% | | | | | | | | 17,18/ | 75% |
| | Nurse practitioners | 1992 | 10% | | | | | | | | 17/12% | 75% |
| | Obstetricians/gynecologists | 1992 | 10% | | | | | | | | 17,18/ | 75% |
| | Internists | 1992 | 9% | | | | | | | | 17,18/ | 75% |
| | Family physicians | 1992 | 8% | | | | | | | | 17,18/ | 75% |
| 10.16 | Work-related homicides (per 100,000 full-time workers) | 1980-89 | 0.7 | | | 0.9 | 0.8 | 0.9 | 0.8 | 0.7 | 1/0.5 | 0.5 |
| 10.17 | Occupational lung disease deaths (age adjusted per million/19) | 1990 | 9.6 | | | | | | | | | 7.7 |
| | Number of pneumoconiosis deaths among people 15 years and over/20 | | | 3,644 | 3,486 | 3,230 | 3,238 | 3,216 | 3,151 | 3,114 | | |
| 10.18* | Worksites with smoking policies | | | | | | | | | | | |
| | Policy that bans smoking or limits it to separately ventilated areas | | | | | | | | | | | |
| | 50 or more employees | 1985 | 27% | | | 59% | | | | | 6/79% | 100% |
| | Any smoking policy | | | | | | | | | | | |
| | Medium and large companies | 1987 | 54% | | 85% | | | | | | | 100% |
| | 50 or more employees | | | | | 86% | | | 87% | | | 100% |
| 10.19* | Number of States with comprehensive laws for clean indoor air/21 | | | | | | | | | | | |
| | Private workplaces | 1995 | a/1 | | | | | | | 1 | 1,15/1 | 51 |
| | Public workplaces | 1995 | a/9 | | | | | | | 9 | 1,15/13 | 51 |
| | Restaurants | 1995 | 2 | | | | | | | 3 | 1,15/3 | 51 |
| | Public transportation | 1995 | a/17 | | | | | | | 17 | 1,15/16 | 51 |
| | Hospitals | 1995 | a/8 | | | | | | | 8 | 8 | 51 |
| | Day care centers | 1995 | 21 | | | | | | | 21 | 1,15/22 | 51 |
| | Grocery stores | 1995 | a/4 | | | | | | | 4 | 4 | 51 |
| 10.20* | Preemptive clean indoor air laws | | | | | | | | | | | |
| | States with laws | 1995 | 17 | | | | | | | 19 | 1/17 | 0 |

^{---/}Data not available.

^{.../}Category not applicable.

a/Baseline has been revised.

^{1/1998} data

^{2/1989} data.

^{3/}Data include work-related injuries and illnesses.

^{4/}Data are for adolescents age 15-17 who sought medical treatment in an emergency room and are reported by fiscal year.

^{5/}Data are for the fiscal year 1996.

^{6/1999} data

^{7/}Data represent a cross-section of civilian and military employees.

^{8/}Data are from 7 States.

^{9/}Data are from 13 States.

^{10/}Data are from 18 States.

^{11/}Data are from 17 States.

^{12/}Data are from 22 States.

^{13/}Data are from 24 States.

^{14/}Data are from 25 States.

| | Baseline | | | | | | | | | Target | |
|-----------|----------|----------|------|------|------|------|------|------|------|--------|------|
| Objective | year | Baseline | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 2000 |

15/Health care workers only.

16/Pursuant to the enactment of the Federal Coal Mine Health and Safety Act of 1969 (PL91-173, amended by PL95-164) and the Occupational Safety and Health Act of 1970 (PL91-596), Federal Standards have been established for occupational exposure to airborne asbestos fibers, cotton dust, coal mine dust and silica dust. These exposure limits apply in all 50 States and U.S. Territories.

17/1997-98 data.

18/Response rate for this group was too low to produce reliable estimates.

19/ICD-9 codes: 500-502, 504. 20/ICD-9 codes: 500-505.

21/Includes the District of Columbia.

NOTE: Data include revisions and, therefore, may differ from data previously published in these reports and other publications.

Data Sources:

| Objective number | Data source |
|------------------|---|
| 10.1, 10.1a-d | Baseline and 1990-91 updates: Annual Survey of Occupational Injuries and Illnesses, DOL, BLS. |
| | 1992-98 updates: Census of Fatal Occupational Injuries, DOL, BLS. |
| 10.2, 10.2a-e | Annual Survey of Occupational Injuries and Illnesses, DOL, BLS. |
| 10.2f | National Electronic Injury Surveillance System, CPSC. |
| 10.3, 10.3a-b | Annual Survey of Occupational Injuries and Illnesses, DOL, BLS. |
| 10.4 | Annual Survey of Occupational Injuries and Illnesses, DOL, BLS. |
| 10.5* | National Notifiable Disease Surveillance System, CDC, EPO. |
| | Sentinel Counties Surveillance of Acute Viral Hepatitis, Viral Hepatitis Surveillance Program, CDC, NCID. |
| 10.6 | Baseline: National Survey of Worksite Health Promotion Activities, OPHS, ODPHP. |
| | 1995 Update: Business Responds to AIDS Benchmark Survey, CDC, NCHSTP; 1999 Update: National Worksite Health Promotion Survey, |
| | OPHS, ODPHP. |
| 10.7 | U.S. Air Force Hearing Conservation Database, DoD. |
| 10.8 | Adult Elevated Blood Lead Level Registries, CDC, NIOSH. |
| 10.9* | Baseline: Regulatory Impact Analysis of OSHA Final Rule on Occupational Exposure |
| | to Bloodborne Pathogens, DOL, OSHA, ORA. |
| | Updates: CDC, NCID. |
| 10.10 | Baseline: Association of State and Territorial Health Officials Reporting System: |
| | Unintentional Injuries Survey, PHF. |
| | Updates: DOL, OSHA. |
| 10.11 | CDC. NIOSH. |
| 10.12 | National Survey of Worksite Health Promotion Activities, OPHS, ODPHP. |
| 10.13 | Baseline and 1992 updates: National Survey of Worksite Health Promotion Activities, OPHS, ODPHP. |
| | 1995 update: Business Responds to AIDS Benchmark Survey, CDC, NCHSTP; 1999 Update: National Worksite Health Promotion Survey, |
| | OPHS, ODPHP. |
| 10.14 | CDC, NIOSH. |
| | |

| . • | E | 3aselin ₆ | | | | | | | | | | Target |
|-----------|---|----------------------|--------------|-----------|----------|-----------|------------|----------|-----------|-------------|------------|---------|
| Objective | У | /ear | Baseline | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 2000 |
| 10.15 | Baseline: Primary Care Provider Survey | s, OPHS | S, ODPHP. | | | | | | | | | |
| | Update: Prevention in Primary Care Study, American College of Preventive Medicine. | | | | | | | | | | | |
| 10.16 | Baseline: National Traumatic Occupatio | nal Fata | lities, CDC, | NIOSH. | | | | | | | | |
| | Updates: Census of Fatal Occupational Injuries, DOL, BLS. | | | | | | | | | | | |
| 0.17 | National Vital Statistics System, CDC, N | NCHS. | | | | | | | | | | |
| 0.18* | Baseline, 1991, and 1992 updates for worksites with 50 or more employees: National Survey of Worksite Health Promotion Activities, OPHS, ODPI | | | | | | | | | | | |
| | 1995 data: Business Responds to AIDS | Benchm | nark Survey, | CDC, NO | CHSTP; | 1999 upd | ate: Natio | nal Work | site Heal | th Promotic | on Survey, | OPHS, O |
| | For medium and large companies: Nation | onwide S | Survey on Sr | noking in | the Work | kplace, C | DC, OSH | l; | | | | |
| | Bureau of National Affairs; American S | Society f | or Personne | l Adminis | tration. | | | | | | | |
| 10.19* | Office of Smoking and Health Legislative | e Trackii | ng System, (| CDC, NC | CDPHP. | | | | | | | |
| 10.20* | Office of Smoking and Health Legislative | e Trackii | na Svstem. (| CDC. NC | CDPHP. | | | | | | | |