nens Munthly VITAL STATISTICS REPURT

Final Data From the National Center for Health Statistics

Vol. 32, No. 10, Supplement (2) ● February 17, 1984

Multiple Causes of Death in the United States

Introduction

Presented here are summary data on multiple causes of death in the United States for 1978. These statistics reflect all medical information reported on the death certificates for the 1,927,788 resident deaths occurring during the year. They complement the 1978 statistics on underlying cause of death.

For many years national cause-of-death statistics have been based solely on the underlying cause of death. The underlying cause is defined as the disease or injury that initiated the train of events leading directly to death or as the circumstances of the accident or violence that produced the fatal injury. It is selected from the conditions entered by the physician in the cause-of-death section of the death certificate. When more than one cause or condition is entered by the physician, the underlying cause is determined by the sequence of conditions on the certificate, provisions of the International Classification of Diseases, and associated classification rules. This is the cause by which deaths have been routinely tabulated and cross-classified by demographic variables such as age, race, sex, and geographic area of occurrence and residence. When only information related to the underlying cause of death is made available, a great deal of information appearing on the death certificates is lost to the data user, including the conditions recorded as the immediate cause of death, any diseases that intervene between the underlying and immediate cause, and any contributory conditions outside the sequence of conditions leading to death.

As an example of the magnitude of this loss, in 1978 nearly 75 percent of the death certificates had more than one condition listed. One-third of the certificates had two conditions, one-fourth had three, and 15 percent had four or more conditions listed. When so many conditions are reported and only one is tabulated for mortality statistics, a substantial amount of diagnostic information remains unused.

To capture this information, the National Center for Health Statistics (NCHS) has developed and implemented a multiple-cause-of-death statistical program. Its objective is to disseminate data on all diseases, conditions, and injuries entered on the death certificates by the certifying physicians. Multiple causes of death are now available on public use tapes for deaths occurring during each year of the period 1968-80. The multiple-cause-of-death program is now a part of the basic data program of NCHS, and multiple cause statistics will be released annually. This first report presents selected summary data from the file of resident deaths for 1978, the last year of the decade for which cause-of-death data were classified according to the Eighth Revision International Classification of Diseases, Adapted for Use in the United States (ICDA-8).

Multiple cause data are not a replacement for the traditional underlying cause-of-death data. They give additional information, information on diseases that are a factor in death whether or not they are the underlying cause of death; on associations among diseases; and on injuries leading to death. They also give information that will aid in evaluating and strengthening the certification process, thereby improving the quality of the medical information entered on the death certificate. The information presented here should help users understand and interpret them.

Underlying causes versus reported causes

Some causes are selected as the underlying cause of death much more often than others, relative to the number of deaths for which they are reported on the death certificate. This relationship can be expressed by a ratio obtained by dividing the total number of deaths for which a given cause

¹National Center for Health Statistics: Eighth Revision International Classification of Diseases, Adapted for Use in the United States. PHS Pub. No. 1693. Public Health Service. Washington. U.S. Government Printing Office, 1967.

appears on the death certificate by the number of those deaths for which it is selected as the underlying cause. The ratio is close to 1.0 if the cause is usually selected as the underlying cause when it is on the death certificate; it is larger for causes less often selected as the underlying cause relative to the total number of deaths for which they are reported.

Table I shows for a list of 69 selected causes the number of deaths for which they were reported on the death certificate, the number of those deaths for which they were selected as the underlying cause, and the number for which they were not selected as the underlying cause. Also shown for each of the 69-cause categories is the ratio relating the total number of deaths for which the cause was reported to the number of deaths for which it was selected as the underlying cause.

Disease categories with large ratios are those that appear much more frequently as reported causes than as underlying causes. They therefore include diseases that in certain cases would not be fatal by themselves but, combined with another serious disease, contribute to the death. Also in this group are diseases that were part of a sequence of conditions that they did not initiate and diseases that occurred either as the immediate or precipitating cause or as a complication where another cause initiated the sequence of events leading to death.

The ratios for several of the causes in the List of 69 Selected Causes of Death developed by the National Center for Health Statistics show that the number of death certificates upon which the causes appeared was more than 10 times the number from which they were selected as the underlying

cause: Angina pectoris (16.2), Symptoms and ill-defined conditions (12.9), and Avitaminoses and other nutritional deficiencies (11.1). Acute poliomyelitis also had a high ratio (10.1), but this was due to certification practices that resulted in a number of cases of late effects of poliomyelitis being assigned to the category for Acute poliomyelitis. Causes that appeared on 5 to 8 times as many death certificates as the number from which they were selected as the underlying cause were Anemias (8.2), All other forms of heart disease (a residual category with a large number of cases of congestive heart failure and cardiac arrest, not otherwise specified) (8.2), Septicemia (6.6), Arteriosclerosis (6.5), Acute nephritis and nephrotic syndrome (5.7), and Hyperplasia of prostate (5.2). These and other causes with large ratios are shown in table A.

The ratios indicate that the following causes from the List of 69 Selected Causes were usually selected as the underlying cause if they were on the death certificate: Motor vehicle accidents (1.0); Suicide (1.0); Homicide (1.0); Other external causes (1.1); Meningococcal infections (1.1); Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues (1.1); Acute myocardial infarction (1.1); Ischemic heart disease (1.2); Complications of pregnancy, childbirth, and the puerperium (1.2); Hypertensive heart and renal disease (1.3); and Certain causes of mortality in early infancy (1.3). The ratios for malignant neoplasms by specified site and for other causes with ratios close to 1 are shown in table B.

The number of deaths and the death rates for the 69 selected causes as reported causes are shown by race and sex in tables 2 and 3. The relationship between these numbers

Table A. Reported causes of death not frequently selected as underlying causes and ratios of reported to underlying causes: United States, 1978 [Based on NCHS List of 69 Selected Causes of Death]

Cause of death and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Total reported	Selected as underlying cause	Ratio of reported to underlying cause
Tuberculosis of respiratory system	8.576	2.256	3.8
Septicemia	51,265	7,800	6.6
Acute poliomyelitis	131	13	10.1
Syphilis and its sequelae	524	169	3.1
Other infective and parasitic diseases Remainder of 000–136	15,619	4,813	3.2
Malignant neoplasms of all other and unspecified sites 170-173, 190-199	201,218	48,943	4.1
Diabetes mellitus	130,672	33,841	3.9
Avitaminoses and other nutritional deficiencies	31,700	2,855	11.1
nemias	27,156	3,300	8.2
Other acute and subacute forms of ischemic heart disease 411	16,115	4,090	3.9
Ingina pectoris	4,108	253	16.2
Chronic disease of endocardium and other myocardial insufficiency 424, 428	14,389	4,072	3.5
All other forms of heart disease	488,000	59,430	8.2
Cerebral embolism	3,555	760	4.4
urteriosclerosis	187,126	28,940	6.5
cute bronchitis and bronchiolitis	2,752	756	3.6
neumonia	180,771	54,267	3.3
sthma	6,278	1,872	3.4
lernia and intestinal obstruction	19,798	5,568	3.6
cute nephritis and nephrotic syndrome	11,905	2,073	5.7
yperplasia of prostate	4,469	854	5.2
ymptoms and ill-defined conditions	402,871	31,324	12.9
If other diseases	509,290	137,396	3.7

Table B. Reported causes of death usually selected as underlying causes and ratios of reported to underlying causes: United States, 1978 [Based on NCHS List of 69 Selected Causes of Death]

Cause of death and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Total : reported	Selected as underlying cause	Ratio of reported to underlying cause
leningococcal infections	439	403	1.1
lalignant neoplasms of buccal cavity and pharynx	10.079	8,341	1.2
lalignant neoplasms of digestive organs and peritoneum 150–159	119,053	105,427	1.1
lalignant neoplasms of respiratory system	109,550	99,898	1.1
lalignant neoplasm of breast	40,602	34,609	1.2
falignant neoplasms of genital organs	57,528	45,131	1.3
falignant neoplasms of urinary organs	21,995	17,574	1.3
eukemia	19,633	15,391	1.3
Other neoplasms of lymphatic and hematopoietic tissues 200–203, 208, 209	27,293	21,678	1.3
lypertensive heart and renal disease	4,101	3,249	1.3
schemic heart disease	763,810	642,270	1.2
cute myocardial infarction	326,643	302,664	1.1
complications of pregnancy, childbirth, and the puerperium 630-678	372	321	1.2
ertain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778	28,962	22,033	1.3
lotor vehicle accidents	53,559	52,411	1.0
uicide	27,459	27,294	1.0
omicide	20,634	20,432	1.0
Other external causes	4,433	4,163	1.1

and rates and those published earlier for the 69 selected causes as underlying causes can be better summarized through use of the ratios discussed above, ratios relating the number of deaths for which a cause is reported to the number of deaths for which it was selected as the underlying cause. The ratios by race and sex are shown in table 4.

Although many of the ratios for males and females are similar, ratios tend to be greater for males than for females. Among the most frequently reported causes, ratios were greater for males than for females for Arteriosclerosis, Diabetes mellitus, and Influenza and pneumonia, indicating that when these causes appeared on the death certificate they were relatively less often selected as the underlying cause for males than for females. The reverse was true for one frequently reported cause, All other accidents (accidents other than motor vehicle accidents).

By race, many of the ratios are similar for the two racial groups shown, but they tend to be greater for the white population than for the population of all other races. Among the frequently reported causes for which ratios were greater for the white population were Diabetes mellitus, Septicemia and All other accidents.

Leading causes of death

Since multiple cause data provide a means for looking at diseases in terms of the number of deaths for which they were mentioned on the death certificates (reported causes) as well as of the number for which they were selected as the underlying cause, rankings of causes of death by the two criteria can be compared. This comparison is made in tables 5 and 6, and in the text figure for the 10 leading causes of death.

Diseases of heart, Malignant neoplasms, and Cerebrovascular diseases were the three most frequent causes in the same order whether ranked as underlying cause or reported cause; but a number of causes changed in relative importance when ranked by numbers of deaths for which they were reported. The greatest change in ranking was for Arteriosclerosis, which ranked 8th as an underlying cause but ranked 4th as a reported cause. Cirrhosis of liver changed rank from 7th to 10th. Suicide and Certain causes of mortality in early infancy, ranked 9th and 10th as underlying causes of death, were replaced in the 10 leading causes by Bronchitis, emphysema, and asthma and by Septicemia, which ranked 8th and 9th as reported causes.

It is of interest that while the number of deaths for which Diabetes mellitus was reported (130,672) was far greater than the number for which it was selected as the underlying cause (33,841), its relative ranking among causes changed little, from 7th to 6th. Similarly, the category Influenza and pneumonia was reported for three times the number of deaths for which it was selected as the underlying cause, yet retained the same rank. Also, Cirrhosis of liver was reported for nearly 50 percent more deaths than the number for which it was selected as underlying cause, yet its ranking dropped from 7th among underlying causes to 10th as a reported cause.

These differences in ranking of a given cause ranked by the two criteria (the number of deaths for which selected as the underlying cause and the number of deaths for which reported) occur because, as discussed earlier, some causes are selected more often than others as the underlying cause of death relative to the number of deaths for which they are reported. If all reported causes were selected as underlying causes with the same relative frequency, each of the causes would be ranked the same in both groups.

The ranking of reported causes, as of underlying causes, is different for males than for females. Among reported causes, Arteriosclerosis, Diabetes mellitus, Septicemia, and




Figure. Percent distribution and rank order of 10 leading causes of death for underlying causes and for reported causes: United States, 1978

Avitaminoses and other nutritional deficiencies rank higher for females than for males. Influenza and pneumonia; Accidents; Bronchitis, emphysema, and asthma; and Cirrhosis of liver rank lower for females.

Rankings of the leading causes of death as underlying causes and as reported causes are shown by age in table 6. At certain ages, several causes of death ranked higher as reported causes than as underlying causes: Diseases of heart at 1-4 and 5-14 years of age; Anemias at 15-24 years of age; Arteriosclerosis at 45-64; Avitaminoses and other

nutritional deficiencies at 65 years of age and over; and Septicemia at all ages. Suicide and Homicide declined in rank at 45-64 years of age, dropping from the leading causes. Cirrhosis of liver and Nephritis and nephrosis also dropped from the top 10 causes for 65 years of age and over.

Nature of injury

National cause-of-death statistics that show only the underlying cause, the cause that initiated the train of events leading to death, show the circumstances of the accident or violence that produced the fatal injuries, but not the nature of the injuries. Since, in selecting one cause to represent a death, a choice has to be made between the two, it is considered more important to know the number of deaths resulting from such external causes as motor vehicle accidents, falls, homicide, and suicide than from causes categorized by the nature of the injury, such as fractures, gunshot wounds, and lacerations.

For example, in underlying cause statistics a death resulting from a skull fracture sustained in an automobile accident would be classified as a motor vehicle accident rather than as a skull fracture. Consequently, underlying cause statistics provide us with information on deaths resulting from automobile accidents but not on deaths involving skull fractures. Multiple cause statistics yield information on both. We now have information on the nature of injuries incurred in accidents, including fractures (by site), intracranial injuries, internal injuries, lacerations, burns, and many other injuries, as well as information on the external causes, such as motor vehicle accidents, falls, fires, poisonings, drownings, suicide, and homicide.

Injuries were reported for 210,942 deaths in 1978, 11 percent of all deaths in that year (table 7). By broad category,

the injuries reported for the most deaths were those classified to Adverse effects of chemical substances and other adverse effects (N960-N999), a category that includes injuries of specified, unspecified, and multiple sites of the body, and complications of those injuries; adverse effects of medical and nonmedical agents; complications of surgical procedures and other medical care; and effects of a variety of other external causes, including drowning, asphyxiation and strangulation, radiation, extreme temperature, and lightning. A total of 95,619 deaths, nearly half (45 percent) of the deaths with a reported injury, were in this broad category. The category Intracranial injury, excluding those with skull fracture (N850-N854), was reported for 49,797 deaths; and Fractures, dislocations, and sprains and strains of joints and adjacent muscles (N800-N848) was reported for 49,420 deaths.

Injuries were reported for more deaths for males (139,891) than for females (71,051), 13 percent of all deaths for males and 8 percent of all deaths for females. Deaths with mention of Intracranial injury (excluding those with skull fracture) were more frequent among the injury deaths for males than among those for females. The reverse was true for Fractures, dislocations, and sprains and strains of joints and adjacent muscles. However, that category assumed greater prominence in the upper age groups for both men and women.

Table 1. Deaths with 69 selected causes as reported causes, by whether selected as underlying causes, and ratios of reported to underlying cause: United States, 1978

Classification of Diseases, Adapted, 1965 cillary dysentery and amebiasis	Total reported 54 3.757	underlying cause	underlying cause	underlyin cause
teritis and other diarrheal diseases				Cause
teritis and other diarrheal diseases		2/	20	1.6
berculosis, all forms 010–019 Tuberculosis of respiratory system 010–012 Tuberculosis, other forms 013–019	3,757	34 1,874	1,883	2.0
Tuberculosis of respiratory system	10,029	2,914	7,115	3.4
Tuberculosis, other forms	8,576	2,256	6,320	3.8
	1,541	2,256 658	883	2.3
	1,541	6	6	2.0
nooping cough	12	5	7	2.4
iningococcal infections	439	403	36	1.1
oticemia	51,265	7,800	43,465	6.6
ute poliomyelitis	131	13	118	10.1
asles	27	11	16	2,5
philis and its sequelae	524	169	355	3.1
ner infective and parasitic diseases	15,619	4,813	10,806	3.2
lignant neoplasms, including neoplasms of lymphatic	10,010	1,010	10,000	
and hematopoietic tissues	451,612	396,992	54,620	1.1
Malignant neoplasms of buccal cavity and pharynx	10,079	8,341	1,738	1.2
Malignant neoplasms of digestive organs and peritoneum 150–159	119,053	105,427	13,626	1.1
Malignant neoplasms of respiratory system	109,550	99,898	9,652	1.1
Malignant neoplasm of breast	40,602	34,609	5,993	1.2
Malignant neoplasms of genital organs	57,528	45,131	12,397	1.3
Malignant neoplasms of urinary organs	21,995	17,574	4,421	1.3
Malignant neoplasms of all other and unspecified sites 170-173,190-199	201,218	48,943	152,275	4.1
Leukemia	19,633	15,391	4,242	1.3
Other neoplasms of lymphatic and hematopoietic tissues 200–203,208,209	27,293	21,678	5,615	1.3
nign neoplasms and neoplasms of unspecified nature	9,179	4,963	4,216	1.8
abetes mellitus	130,672	33,841	96,831	3.9
itaminoses and other nutritional deficiencies	31,700	2,855	28,845	11.1
emias	27,156	3,300	23,856	8.2
ningitis	3,200	1,560	1,640	2.1
jor cardiovascular diseases	1,240,359	966,120	274,239	1.3
Diseases of heart	1,014,863	729,510	285,353	1.4
Active rheumatic fever and chronic rheumatic heart disease 390-398	24,632	13,402	11,280	1.8
Hypertensive heart disease	13,654	7,087	6,567	1.9
Hypertensive heart and renal disease	4,101	3,249	852	1.3
Ischemic heart disease	763,810	642,270	121,540	1.2
Acute myocardial infarction	326,643	302,664	23,979	1.1
Other acute and subacute forms of ischemic heart disease 411	16,115	4,090	12,025	3.9
Chronic ischemic heart disease	604,945	335,263	269,682	1.8
Angina pectoris	4,108	253	3,855	16.2
Chronic disease of endocardium and other myocardial insufficiency 424,428	14,389	4,072	10,317	3.5
All other forms of heart disease	488,000	59,430	428,570	8.2
Hypertension	13,217	5,490	7,727	2.4
Cerebrovascular diseases	289,246	175,629	113,617	1.6
Cerebral hemorrhage	32,743	22,633	10,110	1.4
Cerebral thrombosis	55,876	38,710	17,166	1.4
Cerebral embolism	3,355	760	2,595	4.4
All other cerebrovascular diseases	225,720	113,526	112,194	2.0
Arteriosclerosis	187,126	28,940	158,186	6.5
Other diseases of arteries, arterioles, and capillaries	53,318	26,551	26,767	2.0
ute bronchitis and bronchiolitis	2,752	756	1,996	3.6 3.2
luenza and pneumonia	186,512	58,319	128,193	1.4
Influenza	5,754	4,052	1,702	3.3
Pneumonia	180,771	54,267	126,504	2.5
onchitis, emphysema, and asthma	54,508 10,425	21,875 4,376	32,633 6,049	2.5
Chronic and unqualified bronchitis	10,425 42,364	4,376 15,627	26,737	2.4
Emphysema	42,364 6,278	1,872	4,406	3.4
Asthma				2.8
otic ulcer	15,393	5,545 735	9,848 485	1.7
pendicitis	1,220	735 5,568	14,230	3.6
ernia and intestinal obstruction	19,798	•	14,023	1.5
rrhosis of liver	44,089 g 151	30,066 2,941	5,210	2.8
nolelithiasis, cholecystitis, and cholangitis	8,151	•	19,830	3.2
ephritis and nephrosis	28,698	8,868 2,073	·	5.2 5.7
Acute nephritis and nephrotic syndrome	11,905	2,073	9,832	2.6
Chronic and unqualified nephritis and renal sclerosis	17,846	6,795 3,096	11,051 5.727	2.8
rections of kidney	8,823 4,469	3,096 854	5,727 3,615	5.2

Table 1. Deaths with 69 selected causes as reported causes, by whether selected as underlying causes, and ratios of reported to underlying cause: United States, 1978—Con.

Cause of death and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Total reported	Selected as underlying cause	Not selected as underlying cause	Ratio of reported to underlying cause
Complications of pregnancy, childbirth, and the puerperium 630–678	372	321	51	1.2
Abortions	22	16	6	1.4
and the puerperium	353	305	- 48	1.2
Congenital anomalies	17,861	12,968	4,893	1.4
Certain causes of mortality in early infancy 760–769.2,769.4–772,774–778 Birth injury, difficult labor, and other anoxic	28,962	22,033	6,929	1.3
and hypoxic conditions	17,236	11,710	5,526	1.5
Other causes of mortality in early infancy Remainder of 760–778	26,023	10,323	15,700	2.5
Symptoms and ill-defined conditions	402,871	31,324	371,547	12.9
All other diseases	509,290	137,396	371,894	3.7
Accidents	161,561	105,561	56,000	1.5
Motor vehicle accidents	53,559	52,411	1,148	1.0
All other accidents	108,516	53,150	55,366	2.0
Suicide	27,459	27,294	165	1.0
Homicide	20,634	20,432	202	1.0
Other external causes	4,433	4,163	270	1.1

Table 2. Deaths with 69 selected causes as reported causes, by race and sex: United States, 1978

Cause of death and category number		All races			White		All other		
of the Eighth Revision International Classification	Both			Both			Both		
of Diseases, Adapted, 1965	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Bacillary dysentery and amebiasis	54	34	20	40	25	15	14	9	5
Enteritis and other diarrheal diseases	3,757	1,707	2,050	2,983	1,305	1,678	774	402	372
Tuberculosis, all forms	10,029	6,054	3,975	7,881	4,603	3,278	2,148	1,451	697
Tuberculosis of respiratory system	8,576	5,184	3,392	6,786	3,960	2,826	1,790	1,224	566
Tuberculosis, other forms	1,541	926	615	1,145	680	465	396	246	150
Whooping cough	12	4	8	6	2	4	6	2	4
Streptococcal sore throat and scarlet fever	12	6	6	12	6	6	-	-	-
Meningococcal infections	439	233	206	353	192	161	86	41	45
Septicemia	51,265	26,694	24,571	41,775	21,690	20,085	9,490	5,004	4,486
Acute poliomyelitis	131	78	53	127	75	52	4	3	1
Measles	27	12	15	23	11	12	4	1	3
Syphilis and its sequelae	524	311	213	309	192	117	215	119	96
Other infective and parasitic diseases Remainder of 000–136	15,619	8,061	7,558	12,979	6,682	6,297	2,640	1,379	1,261
Malignant neoplasms, including neoplasms of lymphatic and									
hematopoietic tissues	451,612	248,425	203,187	401,787	219,485	182,302	49,825	28,940	20,885
Malignant neoplasms of buccal cavity and pharynx 140-149	10,079	7,109	2,970	8,712	6,084	2,628	1,367	1,025	342
Malignant neoplasms of digestive organs and									
peritoneum	119,053	62,674	56,379	105,306	54,931	50,375	13,747	7,743	6,004
Malignant neoplasms of respiratory system	109,550	82,204	27,346	97,577	72,758	24,819	11,973	9,446	2,527
Malignant neoplasm of breast	40,602	363	40,239	36,840	316	36,524	3,762	47	3,715
Malignant neoplasms of genital organs 180-187	57,528	31,897	25,631	49,635	27,342	22,293	7,893	4,555	3,338
Malignant neoplasms of urinary organs	21,995	14,817	7,178	20,262	13,733	6,529	1,733	1,084	649
Malignant neoplasms of all other and unspecified									
sites	201,218	104,431	96,787	180,230	92,918	87,312	20,988	11,513	9,475
Leukemia	19,633	11,203	8,430	18,025	10,279	7,746	1,608	924	684
Other neoplasms of lymphatic and hematopoietic									
tissues	27,293	14,605	12,688	24,749	13,206	11,543	2,544	1,399	1,145
Benign neoplasms and neoplasms of unspecified nature 210-239	9,179	4,312	4,867	8,098	3,841	4,257	1,081	471	610
Diabetes mellitus	130,672	57,322	73,350	113,738	50,754	62,984	16,934	6,568	10,366
Avitaminoses and other nutritional deficiencies	31,700	15,209	16,491	27,742	13,026	14,716	3,958	2,183	1,775
Anemias	27,156	12,393	14,763	23,020	10,460	12,560	4,136	1,933	2,203
Meningitis	3,200	1,820	1,380	2,408	1,360	1,048	792	460	332
Major cardiovascular diseases	1,240,359	649,901	590,458	1,105,391	579,033	526,358	134,968	70,868	64,100
Diseases of heart	1,014,863	548,644	466,219	906,652	490,601	416,051	108,211	58,043	50,168
Active rheumatic fever and chronic rheumatic heart									
disease	24,682	11,184	13,498	22,701	10,247	12,454	1,981	937	1,044
Hypertensive heart disease	13,654	5,721	7,933	10,389	4,250	6,139	3,265	1,471	1,794
Hypertensive heart and renal disease	4,101	1,851	2,250	3,431	1,529	1,902	670	322	348
Ischemic heart disease	763,810	416,338	347,472	694,673	379,379	315,294	69,137	36,959	32,178
Acute myocardial infarction 410	326,643	197,873	128,770	301,305	183,556	117,749	25,338	14,317	11,021
Other acute and subacute forms of ischemic									
heart disease	16,115	9,954	6,161	14,378	8,874	5,504	1,737	1,080	657
Chronic ischemic heart disease	604,945	318,489	286,456	550,284	289,839	260,445	54,661	28,650	26,011
Angina pectoris	4,108	2,390	1,718	3,855	2,257	1,598	253	133	120
Chronic disease of endocardium and other myocardial	•	•	, -	• • •	•	•			
insufficiency	14,389	7,708	6,681	12,882	6,834	6,048	1,507	874	633
All other forms of heart	•	•		•	• • •	•	•		
disease	488,000	260,396	227,604	433,820	231,238	202,582	54,180	29,158	25,022
Hypertension	13,217	6,110	7,107	10,704	4,910	5,794	2,513	1,200	1,313
21							•		•

Cerebrovascular diseases	289,246	125,215	164,031	254,255	108,795	145,460	34,991	16,420	18,571
Cerebral hemorrhage	32,743	14,990	17,753	27,484	12,321	15,163	5,259	2,669	2,590
Cerebral thrombosis	55,876	23,842	32,034	49,939	21,049	28,890	5,937	2,793	3,144
Cerebral embolism	3,355	1,435	1,920	3,110	1,312	1,798	245	123	122
All other cerebrovascular diseases 430, 432, 435–438	225,720	96,611	129,109	199,421	84,492	114,929	26,299	12,119	14,180
Arteriosclerosis	187,126	83,854	103,272	173,078	77,274	95,804	14,048	6,580	7,468
Other diseases of arteries, arterioles, and capillaries 441–448	53,318	29,958	23,360	48,314	27,415	20,899	5,004	2,543	2,461
Acute bronchitis and bronchiolitis	2,752	1,542	1,210	2,530	1,410	1,120	222	132	90
Influenza and pneumonia	186,512	103,202	83,310	166,601	90,911	75,690	19,911	12,291	7,620
Influenza	5,754	2,365	3,389	5,405	2,199	3,206	349	166	183
Pneumonia	180,771	100,841	79,930	161,209	88,716	72,493	19,562	12,125	7,437
Bronchitis, emphysema, and asthma	54,508	38,753	15,755	51,016	36,311	14,705	3,492	2,442	1,050
Chronic and unqualified bronchitis	10,425	6,381	4,044	9,830	6,018	3,812	595	363	232
Emphysema	42,364	32,727	9,637	40,166	30,930	9,236	2,198	1,797	401
Asthma	6,278	2,781	3,497	5,345	2,329	3,016	933	452	481
Peptic ulcer	15,393	9,271	6,122	13,906	8,317	5,589	1,487	954	533
Appendicitis	1,220	675	545	1,017	561	456	203	114	89
Hernia and intestinal obstruction	19,798	8,188	11,610	17,596	7,093	10,503	2,202	1,095	1,107
Cirrhosis of liver	44,089	29,361	14,728	36,733	24,586	12,147	7,356	4,775	2,581
Cholelithiasis, cholecystitis, and cholangitis 574, 575	8,151	3,446	4,705	7,480	3,184	4,296	671	262	409
Nephritis and nephrosis	28,698	15,687	13,011	22,407	12,562	9,845	6,291	3,125	3,166
Acute nephritis and nephrotic syndrome 580, 581	11,905	6,608	5,297	9,993	5,622	4,371	1,912	986	926
Chronic and unqualified nephritis and renal sclerosis 582–584	17,846	9,656	8,190	13,249	7,418	5,831	4,597	2,238	2,359
Infections of kidney	8,823	3,865	4,958	7,707	3,313	4,394	1,116	552	564
Hyperplasia of prostate	4,469	4,469	• • •	4,006	4,006	• • • •	463	463	• • •
Complications of pregnancy, childbirth, and the									470
puerperium	372		372	202	• • •	202	170	* * *	170
Abortions	22	• • • •	22	8		8	14	• • •	14
Other complications of pregnancy, childbirth,			0.50	405		405	450		450
and the puerperium	353		353	195		195	158		158
Congenital anomalies	17,861	9,616	8,245	14,891	8,028	6,863	2,970	1,588	1,382
Certain causes of mortality in early	28.962	16 600	10.054	00.007	11,728	8.559	0.675	4,880	3,795
infancy	28,902	16,608	12,354	20,287	11,726	6,559	8,675	4,000	3,793
Birth injury, difficult labor, and other anoxic	17.000	10 107	7.000	10.404	7.075	5.049	4.812	2.792	0.000
and hypoxic conditions	17,236	10,167	7,069	12,424	7,375	5,049	4,012	2,792	2,020
Other causes of mortality in early	26.023	14.866	11,157	18,034	10,406	7.628	7,989	4,460	3,529
infancy	402.871	214,000	188.844	350,426	185,319	165,107	7,969 52,445	28,708	23,737
All other diseases	509,290	274,027	235,277	449,024	240,671	208,353	60,266	33,342	26,924
Accidents	161,561	102,770	235,277 58,791	140,558	88,468	52,090	21,003	14,302	6,701
Motor vehicle accidents	53,559	38,990	14,569	46,987	34,090	12.897	6.572	4,900	1.672
All other accidents	108,516	64,181	44,335	94,024	54,729	39,295	14,492	9,452	5,040
Suicide	27,459	20,287	7,172	25,402	18,707	6,695	2.057	1,580	477
Homicide	20,634	15,979	4,655	11,334	8,519	2,815	9,300	7,460	1,840
Other external causes	4,433	3,050	1,383	3,431	2,340	1,091	1,002	710	292
Offici Catomai Gauses		0,000	1,000	0,401	2,040	1,001	1,002	, 10	2.52

Table 3. Death rates for 69 selected causes as reported causes, by race and sex: United States, 1978

[Rates per 100,000 population in specified group. Refers only to resident deaths occurring within the United States. Excludes fetal deaths]

Cause of death and category number	_	All races			White			All other	
of the Eighth Revision International Classification	Both			Both			Both		
of Diseases, Adapted, 1965	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Bacillary dysentery and amebiasis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Enteritis and other diarrheal diseases	1.7	1.6	1.8	1.6	1.4	1.7	2.6	2.8	2.4
Tuberculosis, all forms	4.5	5.6	3.5	4.1	4.9	3.3	7.1	10.1	4.4
Tuberculosis of respiratory system	3.9	4.8	3.0	3.5	4.2	2.9	5.9	8.5	3.6
Tuberculosis, other forms	0.7	0.9	0.5	0.6	0.7	0.5	1.3	1.7	0.9
Whooping cough	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Streptococcal sore throat and scarlet fever	0.0	0.0	0.0	0.0	0.0	0.0	-	-	-
Meningococcal infections	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
Septicemia	23.1	24.7	21.5	21.8	23.2	20.4	31.5	34.9	28.4
Acute poliomyelitis	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0
Measles	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Syphilis and its sequelae	0.2	0.3	0.2	0.2	0.2	0.1	0.7	8.0	0.6
Other infective and parasitic diseases Remainder of 000–136	7.0	7.5	6.6	6.8	7,1	6.4	8.8	9.6	8.0
Malignant neoplasms, including neoplasms of lymphatic and									
hematopoietic tissues	203.3	230.1	178.0	209.3	234.4	185.4	165.3	202.0	132.1
Malignant neoplasms of buccal cavity and pharynx 140-149	4.5	6.6	2.6	4.5	6.5	2.7	4.5	7.2	2.2
Malignant neoplasms of digestive organs and									
peritoneum	53.6	58.1	49.4	54.9	58.7	51.2	45.6	54.0	38.0
Malignant neoplasms of respiratory system	49.3	76.1	24.0	50.8	77.7	25.2	39.7	65.9	16.0
Malignant neoplasm of breast	18.3	0.3	35.3	19.2	0.3	37.1	12.5	0.3	23.5
Malignant neoplasms of genital organs	25.9	29.5	22.5	25.9	29.2	22.7	26.2	31.8	21.1
Malignant neoplasms of urinary organs	9.9	13.7	6.3	10.6	14.7	6.6	5.8	7.6	4.1
Malignant neoplasms of all other and unspecified	0.0			, 0,0	• • • • • • • • • • • • • • • • • • • •		0.0		
sites	90.6	96.7	84.8	93.9	99.2	88.8	69.6	80.3	59.9
Leukemia	8.8	10.4	7.4	9.4	11.0	7.9	5.3	6.4	4.3
Other neoplasms of lymphatic and hematopoietic	0.0			U. .		7.0	0.0	0	
tissues	12.3	13.5	11.1	12.9	14,1	11.7	8.4	9.8	7.2
Benign neoplasms and neoplasms of unspecified nature	4.1	4.0	4.3	4.2	4.1	4.3	3.6	3.3	3.9
Diabetes mellitus	58.8	53.1	64.3	59.3	54.2	64.1	56.2	45.8	65.6
Avitaminoses and other nutritional deficiencies	14.3	14.1	14.4	14.5	13.9	15.0	13.1	15.2	11,2
Anemias	12.2	11.5	12.9	12.0	11.2	12.8	13.7	13.5	13.9
Meningitis	1.4	1.7	1.2	1.3	1.5	1.1	2.6	3.2	2.1
Major cardiovascular diseases	558.5	602.0	517.3	575.8	618.4	535.3	447.9	494.5	405.5
Diseases of heart	456.9	508.2	408.5	472.3	524.0	423.1	359.1	405.0	317.4
Active rheumatic fever and chronic rheumatic heart	450.5	300.2	400.5	472.0	324.0	720.1	000.1	403.0	017.4
disease	11.1	10.4	11.8	11.8	10.9	12.7	6.6	6.5	6.6
	6.1	5.3	7.0	5.4	4.5	6.2	10.8	10.3	11.4
Hypertensive heart disease	1.8	1.7	2.0	1.8	1.6	1.9	2.2	2.2	2.2
Ischemic heart disease	343.9	385.6	304.4	361.9	405.2	320.7	229.4	257.9	203.6
	147.1	183.3	112.8	157.0	196.0	119.8	84.1	99.9	69.7
Acute myocardial infarction	147.1	103.3	112.0	157.0	196.0	119.6	04.1	99.9	69.7
Other acute and subacute forms of ischemic	7.0	0.0	E 4	7.5	0.5	F. C		7.5	4.2
heart disease	7.3	9.2 295.0	5.4 251.0	7.5 286.7	9.5 309.5	5.6 264.9	5.8	7.5	
Chronic ischemic heart disease	272.4				309.5 2.4		181.4	199.9	164.6
Angina pectoris	1.8	2.2	1.5	2.0	2.4	1.6	8.0	0.9	8.0
Chronic disease of endocardium and other myocardial	0.5	→ 4	۲0	^ 7	7.0	2.0		o 4	4.0
insufficiency	6.5	7.1	5.9	6.7	7.3	6.2	5.0	6.1	4.0
All other forms of heart	040.7	044.0	100.1	000.0	047.0	000.0	470.0	000.5	450.0
disease	219.7	241.2	199.4	226.0	247.0	206.0	179.8	203.5	158.3
Hypertension	6.0	5.7	6.2	5.6	5.2	5.9	8.3	8.4	8.3

Cerebrovascular diseases	130.2	116.0	143.7	132.5	116.2	147.9	116.1	114.6	117.5
Cerebral hemorrhage	14.7	13.9	15.6	14.3	13.2	15.4	17.5	18.6	16.4
Cerebral thrombosis	25.2	22.1	28.1	26.0	22.5	29.4	19.7	19.5	19.9
Cerebral embolism	1.5	1.3	1.7	1.6	1.4	1.8	0.8	0.9	0.8
All other cerebrovascular diseases	101.6	89.5	113.1	103.9	90.2	116.9	87.3	84.6	89.7
Arteriosclerosis	84.3	77.7	90.5	90.2	82.5	97.4	46.6	45.9	47.2
Other diseases of arteries, arterioles, and capillaries	24.0	27.7	20.5	25.2	29.3	21.3	16.6	17.7	15.6
Acute bronchitis and bronchiolitis	1.2	1.4	1.1	1.3	1.5	1.1	0.7	0.9	0.6
Influenza and pneumonia	84.0	95.6	73.0	86.8	97.1	77.0	66.1	85.8	48.2
Influenza	2.6	2.2	3.0	2.8	2.3	3.3	1.2	1.2	1.2
Pneumonia	81.4	93.4	70.0	84.0	94.7	73.7	64.9	84.6	47.1
Bronchitis, emphysema, and asthma	24.5	35.9	13.8	26.6	38.8	15.0	11.6	17.0	6.6
Chronic and unqualified bronchitis	4.7	5.9	3.5	5.1	6.4	3.9	2.0	2.5	1.5
Emphysema	19.1	30.3	8.4	20.9	33.0	9.4	7.3	12.5	2.5
Asthma	2.8	2.6	3.1	2.8	2.5	3.1	3.1	3.2	3.0
Peptic ulcer	6.9	8.6	5.4	7.2	8.9	5.7	4.9	6.7	3.4
Appendicitis	0.5	0.6	0.5	0.5	0.6	0.5	0.7	0.8	0.6
Hernia and intestinal obstruction	8.9	7.6	10.2	9.2	7.6	10.7	7.3	7.6	7.0
Cirrhosis of liver	19.9	27.2	12.9	19.1	26.3	12.4	24.4	33.3	16.3
Cholelithiasis, cholecystitis, and cholangitis	3.7	3.2	4.1	3.9	3.4	4.4	2.2	1.8	2.6
Nephritis and nephrosis	12.9	14.5	11.4	11.7	13.4	10.0	20.9	21.8	20.0
Acute nephritis and nephrotic syndrome	5.4	6.1	4.6	5.2	6.0	4.4	6.3	6.9	5.9
Chronic and unqualified nephritis and renal sclerosis 582–584	8.0	8.9	7.2	5.2 6.9	7.9	5.9	15.3	15.6	14.9
Infections of kidney	4.0	3.6	4.3	4.0	7.9 3.5	4.5	3.7	3.9	3.6
	2.0	3.0 4.1	4.3	2.1	3.5 4.3	4.5	3.7 1.5	3.9	
Hyperplasia of prostate	2.0	4.1	• • • •	2.1	4.3	• • •	1.5	3.2	• • •
	0.2		0.3	0.1		0.2	0.6		1.1
puerperium								• • •	
Abortions	0.0	• • • •	0.0	0.0		0.0	0.0		0.1
Other complications of pregnancy, childbirth,	0.0			0.4			0.5		4.0
and the puerperium	0.2		0.3	0.1		0.2	0.5		1.0
Congenital anomalies	8.0	8.9	7.2	7.8	8.6	7.0	9.9	11.1	8.7
Certain causes of mortality in early	40.0	45.4	40.0	40.0	40.5				24.0
infancy	13.0	15.4	10.8	10.6	12.5	8.7	28.8	34.1	24.0
Birth injury, difficult labor, and other anoxic									
and hypoxic conditions	7.8	9.4	6.2	6.5	7.9	5.1	16.0	19.5	12.8
Other causes of mortality in early									
infancy Remainder of 760–778	11.7	13.8	9.8	9.4	11.1	7.8	26.5	31.1	22.3
Symptoms and ill-defined conditions	181.4	198.2	165.5	182.6	197.9	167.9	174.0	200.3	150.2
All other diseases	229.3	253.8	206.1	233.9	257.0	211.9	200.0	232.7	170.3
Accidents	72.7	95.2	51.5	73.2	94.5	53.0	69.7	99.8	42.4
Motor vehicle accidents	24.1	36.1	12.8	24.5	36.4	13.1	21.8	34.2	10.6
All other accidents	48.9	59.4	38.8	49.0	58.5	40.0	48.1	66.0	31.9
Suicide	12.4	18.8	6.3	13.2	20.0	6.8	6.8	11.0	3.0
Homicide	9.3	14.8	4.1	5.9	9.1	2.9	30.9	52.1	11.6
Other external causes	2.0	2.8	1.2	1.8	2.5	1.1	3.3	5.0	1.8

Table 4. Ratios of reported causes of death to underlying causes for 69 selected causes, by race and sex: United States, 1978

Cause of death and category number		All races			White		All other		
of the Eighth Revision International Classification	Both			Both			Both		
of Diseases, Adapted, 1965	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Bacillary dysentery and amebiasis	1.6	1.6	1.5	1.8	1.9	1.7	1.2	1.1	1.3
Enteritis and other diarrheal diseases	2.0	1.9	2.1	2.2	2.1	2.3	1.5	1.5	1.5
Tuberculosis, all forms	3.4	3.1	4.0	3.9	3.5	4.6	2.4	2.4	2.6
Tuberculosis of respiratory system	3.8	3.3	4.8	4.3	3.8	5.4	2.6	2.4	3.0
Tuberculosis, other forms	2.3	2.4	2.2	2.4	2.5	2.3	2.1	2.3	1.9
Whooping cough	2.0	1.3	2.7	2.0	2.0	2.0	2.0	1.0	4.0
Streptococcal sore throat and scarlet fever	2.4	1.5	6.0	2.4	1.5	6.0	-	-	_
Meningococcal infections	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Septicemia	6.6	6.6	6.6	6.9	6.9	6.9	5.5	5.6	5.4
Acute poliomyelitis	10.1	8.7	13.3	9.8	8.3	13.0	-	-	
Measles	2.5	3.0	2.1	2.6	3.7	2.0	2.0	1.0	3.0
Syphilis and its sequelae	3.1	3.5	2.6	3.2	3.6	2.6	3.0	3.4	2.7
Other infective and parasitic diseases Remainder of 000–136	3.1	3.3	3.2	3.4	3.4	3.3	2.8	2.9	2.7
·	3.2	3.3	3.2	3.4	3.4	5.5	2.0	2.3	2.1
Malignant neoplasms, including neoplasms of lymphatic and	4.4	1.0	4.4		1.0	1.1	1.1	1.1	1.1
hematopoietic tissues	1.1	1.2	1.1	1.1	1.2	1.2	1.1	1.2	1.1
Malignant neoplasms of buccal cavity and pharynx 140–149	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
Malignant neoplasms of digestive organs and									
peritoneum	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Malignant neoplasms of respiratory system 160-163	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Malignant neoplasm of breast	1.2	1.3	1.2	1.2	1.3	1.2	1.1	1.1	1.1
Malignant neoplasms of genital organs	1.3	1.4	1.1	1.3	1.4	1.1	1.2	1.3	1.1
Malignant neoplasms of urinary organs 188, 189	1.3	1.3	1.2	1.3	1.3	1.2	1.2	1.2	1.2
Malignant neoplasms of all other and unspecified									
sites	4.1	4.1	4.2	4.1	4.1	4.2	3.8	3.9	3.7
Leukemia	1.3	1.3	1.3	1.3	1.3	1.3	1.2	1.2	1.2
Other neoplasms of lymphatic and hematopoietic									
tissues	1.3	1.3	1.2	1.3	1.3	1.2	1.2	1.2	1.2
Benign neoplasms and neoplasms of unspecified nature 210-239	1.8	1.8	1.9	1.9	1.9	1.9	1.8	1.7	1.9
Diabetes mellitus	3.9	4.1	3.7	4.0	4.3	3.8	3.1	3.2	3.0
Avitaminoses and other nutritional deficiencies 260–269	11.1	13.1	9.7	11.5	13.7	10.0	9.1	10.4	7.8
Anemias	8.2	8.3	8.1	8.7	8.9	8.6	6.2	6.1	6.3
Meningitis	2.1	2.1	2.0	2.1	2.2	2.1	1.8	2.0	1.7
Major cardiovascular diseases	1.3	1.3	1.3	1.3	1.3	1.3	1.4	1.4	1.3
Diseases of heart	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.5
Active rheumatic fever and chronic rheumatic heart		***	***	***					
disease	1.8	1.9	1.8	1.9	1.9	1.8	1.7	1.8	1.6
Hypertensive heart disease	1.9	2.0	1.9	1.9	1.9	1.9	2.1	2.0	2.1
Hypertensive heart and renal disease	1.3	1.3	1.3	1.3	1.3	1.2	1.3	1.3	1.3
Ischemic heart disease	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.3
	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Acute myocardial infarction	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Other acute and subacute forms of ischemic	3.9	3.8	4.2	4.1	3.9	4.3	3.1	3.0	3.3
heart disease									
Chronic ischemic heart disease	1.8	1.9	1.7	1.8	1.9	1.7	1.6	1.6	1.6
Angina pectoris	16.2	17.4	14.8	17.3	19.0	15.4	8.4	7.4	10.0
Chronic disease of endocardium and other myocardial									
insufficiency	3.5	3.8	3.2	3.6	4.0	3.3	2.8	3.0	2.6
disease	8.2	8.0	8.4	8.7	8.5	9.0	5.6	5.5	5.6

Hypertension	2.4	2.4	2.4	2.4	2.4	2.5	2.3	2.3	2.4
Cerebrovascular diseases	1.6	1.7	1.6	1.7	1.7	1.6	1.6	1.6	1.6
Cerebral hemorrhage	1.4	1.5	1.4	1.4	1.5	1.4	1.4	1.4	1.5
Cerebral thrombosis	1.4	1.5	1.4	1.4	1.5	1.4	1.5	1.5	1.5
Cerebral embolism	4.4	4.3	4.5	4.4	4.3	4.5	4.3	4.0	4.7
All other cerebrovascular diseases 430, 432, 435-438	2.0	2.0	1.9	2.0	2.1	2.0	1.9	1.9	1.8
Arteriosclerosis	6.5	7.3	5.9	6.5	7.3	5.9	6.6	6.9	6.4
Other diseases of arteries, arterioles, and capillaries 441–448	2.0	1.9	2.2	2.0	1.9	2.2	2.1	2.1	2.2
Acute bronchitis and bronchiolitis	3.6	3.7	3.6	3.8	4.0	3.6	2.3	2.0	3.0
Influenza and pneumonia	3.2	3.4	3.0	3.2	3.4	3.0	2.9	2.9	2.9
Influenza	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Pneumonia	3.3	3.5	3.2	3.4	3.5	3.2	2.9	3.0	2.9
Bronchitis, emphysema, and asthma	2.5	2.5	2.4	2.5	2.6	2.4	2.4	2.4	2.3
Chronic and unqualified bronchitis	2.4	2.3	2.5	2.4	2.3	2.5	2.3	2.3	2.5
Emphysema	2.7	2.8	2.5	2.7	2.8	2.5	2.6	2.6	2.7
Asthma	3.4	3.8	3.1	3.6	4.2	3.2	2.4	2.5	2.4
Peptic ulcer	2.8	2.9	2.6	2.8	2.9	2.6	2.7	2.7	2.8
Appendicitis	1.7	1.6	1.7	1.7	1.7	1.8	1.5	1.6	1.4
Hernia and intestinal obstruction	3.6	3.5	3.6	3.6	3.6	3.7	3.1	2.9	3.2
Cirrhosis of liver	1.5	1.5	1.4	1.5	1.5	1.4	1,4	1.5	1.4
Cholelithiasis, cholecystitis, and cholangitis	2.8	2.6	2.9	2.8	2.6	2.9	2.8	2.6	2.9
Nephritis and nephrosis	3.2	3.3	3.2	3.4	3.5	3.3	2.7	2.7	2.5
Acute nephritis and nephrotic syndrome	5.7	5.8	5.6	5.4 5.9	6.1	5.6	5.2	4.7	5.8
Chronic and unqualified nephritis and renal sclerosis									
·	2.6	2.7	2.6	2.7	2.8	2.7	2.4	2.4	2.4
Infections of kidney	2.8	3.1	2.7	2.9	3.2	2.7	2.6	2.7	2.4
Hyperplasia of prostate	5.2	5.2	• • • •	5.5	5.5	• • •	3.8	3.8	
Complications of pregnancy, childbirth, and the									
puerperium	1.2	• • • •	1.2	1.2	• • •	1.2	1.1	• • •	1.1
Abortions	1.4	• • •	1.4	2.0	• • •	2.0	1.2	• • •	1.2
and the puerperium 630–639, 650–678	1.2		1.2	1.2		1.2	1.1	• • •	1.1
Congenital anomalies	1.4	1.4	1.4	1.4	1.4	1.4	1.3	1.3	1.3
Certain causes of mortality in early									
infancy	1.3	1.3	1.3	1.3	1.3	1.4	1.2	1.2	1.2
Birth injury, difficult labor, and other anoxic									
and hypoxic conditions	1.5	1.5	1.5	1.5	1.5	1.5	1.4	1.4	1.4
Other causes of mortality in early									
infancy Remainder of 760–778	2.5	2.6	2.4	2.7	2.8	2.6	2.2	2.3	2.1
Symptoms and ill-defined conditions	12.9	11.9	14.2	15.3	14.1	17.0	6.2	6.0	6.5
All other diseases	3.7	3.5	3.9	3.8	3.6	4.0	3.2	3.1	3.4
Accidents	1.5	1.4	1.9	1.6	1.4	1.9	1.4	1.3	1.6
Motor vehicle accidents	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
All other accidents	2.0	1.8	2.5	2.1	1.9	2.6	1.7	1.5	2.0
Suicide	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Homicide	1.0	1.0	1.0	1.0	1,0	1.0	1.0	1.0	1.0
Other external causes	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.0
	***	•••	•••	1.1	1.1				1.0

Table 5. 10 leading causes of death according to underlying causes and reported causes, number of deaths, percent of total, and rank order, by race and sex: United States, 1978

Race. sex. cause of death, and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965 All races, both sexes All other accidents E800–8949 All other accidents E800–E949 All other accident	Number of deaths 1,927,788 729,510 396,992 175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	Percent of total 100.0 37.8 20.6 9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	Rank order 1 2 3 4 5 6 7 8 9	Number of deaths 1,927,788 1,014,863 451,612 289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459 28,962	Percent of total 100.0 52.6 23.4 15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7 1.4	## Rank order
Ill causes	729,510 396,992 175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	37.8 20.6 9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	1 2 3 4 5 6 7 8 9	1,014,863 451,612 289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	52.6 23.4 15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7 1.4	1 2 3 6 5 7 10 4
Second S	729,510 396,992 175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	37.8 20.6 9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	1 2 3 4 5 6 7 8 9	1,014,863 451,612 289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	52.6 23.4 15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7 1.4	1 2 3 6 5 7 10 4
iseases of heart	729,510 396,992 175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	37.8 20.6 9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	2 3 4 5 6 7 8 9	1,014,863 451,612 289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	52.6 23.4 15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7 1.4	2 3 6 5 7 10 4
alignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues 140–209 erebrovascular diseases 430–438 ccidents E800–E949 Motor vehicle accidents E800–E807,E825–E949 fluenza and pneumonia 470–474,480–486 abetes mellitus 250 rrhosis of liver 571 rteriosclerosis 440 uicide E950–E959 ertain causes of mortality in early infancy 760–769.2,769.4–772,774–778 ronchitis, emphysema, and asthma 490–493 expticemia 038 White, male	396,992 175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	20.6 9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	2 3 4 5 6 7 8 9	451,612 289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	23.4 15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7 1.4	2 3 6 5 7 10 4
of lymphatic and hematopoietic tissues 140–209 erebrovascular diseases 430–438 cidents E800–E949 Motor vehicle accidents E800–E807,E825–E949 filtenza and pneumonia 470–474,480–486 dabetes mellitus 250 rrhosis of liver 571 teriosclerosis 440 uicide E950–E959 ertain causes of mortality in early infancy 760–769.2,769.4–772,774–778 onchitis, emphysema, and asthma 490–493 expticemia 038 White, male causes seases of heart 390–398,402,404,410–429	175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	3 4 5 6 7 8 9	289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7	3 6 5 7 10 4
### Property of the color of th	175,629 105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	9.1 5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	3 4 5 6 7 8 9	289,246 161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	15.0 8.4 2.8 5.6 9.7 6.8 2.3 9.7	3 6 5 7 10 4
Motor vehicle accidents E800–E949	105,561 52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	5.5 2.7 2.8 3.0 1.8 1.6 1.5 1.4	4 5 6 7 8 9	161,561 53,559 108,516 186,512 130,672 44,089 187,126 27,459	8.4 2.8 5.6 9.7 6.8 2.3 9.7	6 5 7 10 4
Motor vehicle accidents	52,411 53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	2.7 2.8 3.0 1.8 1.6 1.5 1.4	 5 6 7 8 9	53,559 108,516 186,512 130,672 44,089 187,126 27,459	2.8 5.6 9.7 6.8 2.3 9.7 1.4	 5 7 10 4
All other accidents	53,150 58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	2.8 3.0 1.8 1.6 1.5 1.4	 5 6 7 8 9	108,516 186,512 130,672 44,089 187,126 27,459	5.6 9.7 6.8 2.3 9.7 1.4	 5 7 10 4
fluenza and pneumonia 470–474,480–486 abetes mellitus 250 rrhosis of liver 571 teriosclerosis 440 uicide E950–E959 ertain causes of mortality in early infancy 760–769.2,769.4–772,774–778 onchitis, emphysema, and asthma 490–493 epticemia 038 White, male causes seases of heart 390–398,402,404,410–429	58,319 33,841 30,066 28,940 27,294 22,033 21,875 7,800	3.0 1.8 1.6 1.5 1.4 1.1	5 6 7 8 9	186,512 130,672 44,089 187,126 27,459	9.7 6.8 2.3 9.7 1.4	5 7 10 4
abetes mellitus	33,841 30,066 28,940 27,294 22,033 21,875 7,800	1.8 1.6 1.5 1.4 1.1	6 7 8 9	130,672 44,089 187,126 27,459	6.8 2.3 9.7 1.4	7 10 4
rrhosis of liver	30,066 28,940 27,294 22,033 21,875 7,800	1.6 1.5 1.4 1.1	7 8 9	44,089 187,126 27,459	2.3 9.7 1.4	10 4
teriosclerosis	28,940 27,294 22,033 21,875 7,800	1.5 1.4 1.1 1.1	8 9 10	187,126 27,459	9.7 1.4	4
white, male causes of heart	27,294 22,033 21,875 7,800	1.4 1.1 1.1	9 10	27,459	1.4	
ertain causes of mortality in early infancy	22,033 21,875 7,800	1.1 1.1	10			
early infancy	21,875 7,800	1.1		28.962		
white, male causes 390–398,402,404,410–429	21,875 7,800	1.1		28.962		
white, male causes 390–398,402,404,410–429	7,800				1.5	
White, male causes		0.4		54,508	2.8	8
causes	000 : 55			51,265	2.7	9
causes	000 : 00					
seases of heart	000 100	100.0		920.123	100.0	
	920,123					
alignant neoplasms, including neoplasms	359,700	39.1	1	490,601	53.3	1
			_			
of lymphatic and hematopoletic tissues	189,956	20.6	2	219,485	23.9	2
rebrovascular diseases	63,439	6.9	3	108,795	11.8	3
cidents	62,868	6.8	4	88,468	9.6	5
Motor vehicle accidents	33,341	-3.6		34,090	3.7	
All other accidents E800-E807,E825-E949	29.527	3.2		54,729	5.9	
luenza and pneumonia 470–474,480–486	26,578	2.9	5	90,911	9.9	4
iicide	18,619	2.0	6	18,707	2.0	
rrhosis of liver	16,413	1.8	7	24,586	2.7	9
onchitis, emphysema, and asthma	14,228	1.5	8	36,311	3.9	8
abetes mellitus	•	1.3	9	50,754	5.5	7
	11,846					
teriosclerosis	10,583	1.2	10	77,274	8.4	6
pticemia	3,150	0.3	• • •	21,690	2.4	10
White, female						
causes	769,599	100.0		769,599	100.0	
seases of heart	298,002	38.7	1	416,051	54.1	1
alignant neoplasms, including neoplasms		*				
of lymphatic and hematopoietic tissues	162,083	21,1	2	182,302	23.7	2
rebrovascular diseases	90,361	11,7	3	145,460	18.9	3
cidents	27,524	3.6	4	52,090	6.8	7
Motor vehicle accidents	12,633	1.6		12,897	1.7	
				39,295	5.1	
All other accidents	14,891	1.9				
luenza and pneumonia	24,854	3.2	5	75,690	9.8	5
abetes mellitus	16,453	2.1	6	62,984	8.2	6
teriosclerosis	16,230	2.1	7	95,804	12.4	4
rrhosis of liver	8,464	1.1	8	12,147	1.6	
icide E950–E959	6,631	0.9	9	6,695	0.9	
ertain causes of mortality in						
early infancy 760-769.2,769.4-772,774-778	6,297	8.0	10	8,559	1.1	
epticemia	2,921	0.4		20,085	2.6	8
vitaminoses and other nutritional deficiencies	1,466	0.2		14,716	1.9	9
ronchitis, emphysema, and asthma 490–493	6,173	0.8		14,705	1.9	10

Table 5. 10 leading causes of death according to underlying causes and reported causes, number of deaths, percent of total, and rank order, by race and sex: United States, 1978—Con.

Race, sex, cause of death, and	Uni	derlying cause		Reported cause			
category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Number of deaths	Percent of total	Rank order	Number of deaths	Percent of total	Rank	
All other, male							
All causes	135,167	100.0		135,167	100.0		
Diseases of heart	38,567	28.5	1	58,043	42.9	1	
Malignant neoplasms, including neoplasms							
of lymphatic and hematopoietic tissues 140-209	26,041	19.3	2	28,940	21.4	2	
Accidents	11,013	8.1	3	14,302	10.6	4	
Motor vehicle accidents	4,798	3.5		4,900	3.6		
All other accidents E800-E807,E825-E949	6,215	4.6		9,452	7.0		
Cerebrovascular diseases	10,209	7.6	4	16,420	12.1	3	
Homicide	7,409	5.5	5	7,460	5.5	6	
Influenza and pneumonia 470-474,480-486	4,218	3.1	6	12.291	9.1	5	
Dertain causes of mortality in	.,	-	-	1-,		•	
early infancy	3,960	2.9	7	4,880	3.6	10	
Cirrhosis of liver	3,280	2.4	8	4,775	3.5		
Diabetes mellitus	2,064	1.5	9	6,568	4.9	8	
Suicide	1,569	1.2	10	1,580	1.2		
Arteriosclerosis	958	0.7		6.580	4.9	7	
Septicemia	900	0.7		5.004	3.7	9	
Ali other, female				-,		-	
All causes	102.899	100.0		102,899	100.0		
Diseases of heart	33,241	32.3	1	50,168	48.8	1	
Malignant neoplasms, including neoplasms	00,241	02.0	•	00,100	₹0.0	•	
of lymphatic and hematopoietic tissues 140–209	18.912	18.4	2	20.885	20.3	2	
Cerebrovascular diseases	11,620	11.3	3	18.571	18.0	3	
Accidents	4,156	4.0	4	6.701	6.5	7	
Motor vehicle accidents	1,639	1.6		1,672	1.6		
All other accidents	2.517	2.4		5.040	4.9		
Diabetes mellitus	3,478	3.4	5	10,366	10.1	4	
Certain causes of mortality in	0,470	0.4	3	70.000	10.1	7	
early infancy	3.043	3.0	6	3.795	3.7	9	
nfluenza and pneumonia	2.669	2.6	7	7.620	7.4	5	
Cirrhosis of liver	1,909	1.9	8	2.581	2.5		
Homicide	1.823	1.8	9	1.840	2.5 1.8		
Arteriosclerosis	1,169	1.1	10	7,468	7.3	6	
Septicemia	829	0.8		4,486	7.3 4.4	8	
Nephritis and nephrosis	1,152	1,1		3,166	3.1	10	
Topining and hepittosis 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1,102	1, 1		9,100	J. J	10	

Table 6. 10 leading causes of death according to underlying causes and reported causes, number of deaths, percent of total, and rank order, by age: United States, 1978

Age, cause of death, and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Underlying cause				ported cause	
	Number of deaths	Percent of total	Rank order	Number of deaths	Percent of total	Ran orde
1–4 years						
I causes	8,429	100.0		8,429	100.0	
ocidents	3,504	41.6	1	3,988	47.3	1
Motor vehicle accidents	1,287	15.3		1,298	15.4	
All other accidents E800-E807,E825-E949	2,217	26.3		2,698	32.0	
ongenital anomalies	1,027	12.2	2	1,245	14.8	3
alignant neoplasms, including neoplasms of	,			.,		_
lymphatic and hematopoietic tissues	599	71	3	645	7.7	5
luenza and pneumonia	354	4.2	4	839	10.0	4
omicide	313	3.7	5	317	3.8	8
seases of heart	279	3.3	6	1,647	19.5	2
eningitis	220	2.6	7	284	3.4	9
eningococcal infections	112	1.3	8	115	1.4	
erebrovascular diseases	100	1.2	9	432	5.1	6
nemias	78	0.9	10	166	2.0	
epticemia	65	0.3		332	2.0 3.9	10 7
puterna	00	0.5		332	3.9	,
5-14 years						
causes	12,030	100.0	• • •	12,030	100.0	
cidents ,	6,118	50.9	1	6,534	54.3	1
Motor vehicle accidents	3,130	26.0		3,158	26.3	• • •
All other accidents E800-E807,E825-E949	2,988	24.8		3,395	28.2	• • •
alignant neoplasms, including neoplasms of						
lymphatic and hematopoietic tissues	1,500	12.5	2	1,584	13.2	3
ongenital anomalies	650	5.4	3	858	7.1	5
omicide	454	3.8	4	455	3.8	7
seases of heart	357	3.0	5	1,960	16.3	2
fluenza and pneumonia 470–474,480–486	304	2.5	6	928	7.7	4
erebrovascular diseases	199	1.7	7	608	5.1	6
uicide	153	1.3	8	154	1,3	10
nemias	87	0.7	9	214		9
·			-	-	1.8	-
enign neoplasms and neoplasms of unspecified nature 210–239 epticemia	83 42	0.7 0.3	10	105 376	0.9 3.1	8
15-24 years						
l causes	48,500	100.0		48,500	100.0	
ccidents	26,622	54.9	1	27,613	56.9	1
Motor vehicle accidents	19,164	39.5		•	39.9	
All other accidents	•			19,341		
	7,458	15.4	• • • •	8,453	17.4	
pmicide	5,443	11.2	5	5,466	11.3	2
ulcide	5,115	10.5	3	5,144	10.6	3
alignant neoplasms, including neoplasms						
of lymphatic and hematopoietic tissues	2,588	5.3	4	2,740	5.6	5
seases of heart	1,098	2.3	5	4,096	8.4	4
ongenital anomalies	648	1.3	6	925	1.9	8
fluenza and pneumonia 470-474,480-486	553	1.1	7	1,958	4.0	6
erebrovascular diseases	473	1.0	8	1,386	2.9	7
abetes mellitus	140	0.3	9	239	0.5	• •
enign neoplasms and neoplasms of unspecified nature 210-239	136	0.3	10	176	0.4	
epticemia	89	0.2		734	1.5	9
nemias	122	0.3		301	0.6	10
25-44 years						
causes	103,991	100.0		103,991	100.0	
cidents	25,024	24.1	1	27,879	26.8	1
Motor vehicle accidents	14,574	14.0		14,741	14.2	
All other accidents E800-E807,E825-E949	10,450	10.0		13,282	12.8	
alignant neoplasms, including neoplasms of	,			,	-	
lymphatic and hematopoietic tissues	16,866	16.2	2	17,543	16.9	3
seases of heart	14,167	13.6	3	23,925	23.0	2
uicide	9,503	9.1	4	9,562	9.2	4
omicide				•		
	9,127	8.8 4.5	5 6	9,178 5,996	8.8 5.8	5
irrhosis of liver	4,686					

Table 6. 10 leading causes of death according to underlying causes and reported causes, number of deaths, percent of total, and rank order, by age: United States, 1978—Con.

Age, cause of death, and category number of the Eighth Revision International Classification of Diseases, Adapted, 1965	Underlying cause			Reported cause		
	Number of deaths	Percent of total	Rank order	Number of deaths	Percent of total	Rank order
25–44 years—Con.						
Cerebrovascular diseases	3,286	3.2	7	5,367	5.2	8
Influenza and pneumonia 470–474,480–486	1,804	1.7	8	5,802	5.6	7
Diabetes mellitus	1,484	1.4	9	2,909	2.8	9
Congenital anomalies	709	0.7	10	1,193	1.1	
Septicemia	344	0.3	• • •	2,720	2.6	10
45-64 years						
All causes	434,246	100.0		434,246	100.0	
Diseases of heart	151,564	34.9	1	208,868	48.1	1
Malignant neoplasms, including neoplasms of						
lymphatic and hematopoietic tissues	134,115	30.9	2	143,281	33.0	2
Cerebrovascular diseases	21,670	5.0	3	35,468	8.2	3
Accidents	18,774	4.3	4.	30,696	7.1	4
Motor vehicle accidents	8,048	1.9		8,332	1.9	
All other accidents E800-E807,E825-E949	10,726	2.5		22,448	5.2	
Dirrhosis of liver	16,449	3.8	5	22,815	5.3	7
Diabetes mellitus	7,790	1.8	6	27,905	6.4	6
Suicide	7,724	1.8	7	7,766	1.8	
Influenza and pneumonia 470-474,480-486	7,269	1.7	8	29,287	6.7	5
Bronchitis, emphysema, and asthma	5,243	1.2	9	11,774	2.7	10
Homicide	3,696	0.9	10	3,737	0.9	
Arteriosclerosis	1,212	0.3		15,115	3.5	8
Septicemia	1,470	0.3		11,875	2.7	9
65 years and over						
All causes	1,273,882	100.0		1,273,882	100.0	
Diseases of heart	560,983	44.0	1	770,208	60.5	1
Malignant neoplasms, including neoplasms of						
lymphatic and hematopoietic tissues 140–209	241,142	18.9	2	285,621	22.4	2
Cerebrovascular diseases	149,695	11.8	3	245,430	19.3	3
nfluenza and pneumonia 470-474,480-486	46,487	3.6	4	145,050	11.4	5
Arteriosclerosis	27,670	2.2	5	171,279	13.4	4
Diabetes mellitus	24,372	1.9	6	99,518	7.8	6
Accidents	24,135	1.9	7	62,766	4.9	7
Motor vehicle accidents	5,905	0.5		6,383	0.5	
All other accidents E800-E807,E825-E949	18,230	1.4		56,459	4.4	
Bronchitis, emphysema, and asthma 490–493	15,912	1.2	8	41,462	3.3	8
Cirrhosis of liver	8,738	0.7	9	14,910	1.2	
Nephritis and nephrosis	6,158	0.5	10	20,021	1.6	
Septicemia	4,687	0.4		32,709	2.6	9
Avitaminoses and other nutritional deficiencies 260–269	2,366	0.2		24,011	1.9	10

Table 7. Nature of injuries reported for deaths from accidents, poisonings, and violence, by age and sex: United States, 1978 [Refers only to resident deaths occurring within the United States. Excludes fetal deaths]

Both sexes Accidents, poisonings, and violence (Nature of Injury)	•	No state
Accidents, poisonings, and violence (Nature of Injury)	18,035	
Nature of Injury) Nature of Injury) Nature of Injury Nature of	18,035	
Fractures, dislocations, and sprains and strains of joints and adjacent muscles	18,035	
and adjacent muscles N800-N848 49,420 201 600 1,448 7,933 4,452 2,575 2,604 3,081 4,835 10,089 Intracranial injury (excluding those with skull fracture) N850-N854 49,797 396 1,123 2,261 13,733 8,743 5,321 5,029 4,756 4,076 3,049 Internal injury of chest, abdomen, and pelvis N860-N869 31,385 85 482 1,165 9,400 6,473 3,784 3,302 2,813 2,138 1,338 Laceration and open wound N870-N907 31,370 22 147 713 7,841 7,767 4,790 3,776 2,939 1,979 1,020 Superficial injury and contusion N910-N929 2,145 20 64 106 430 265 164 168 175 217 318 Effects of foreign body, entering through orifice N930-N939 12,172 494 322 275 696 643 746 1,200 1,772 2,280 2,360 Burn N940-N949 6,550 137 510 398 901 808 614 699 732 766 661 Injury to nerves and spinal cord N950-N959 2,700 5 33 115 647 500 281 269 312 261 201 Adverse effects of chemical substances and other adverse effects of chemical substances and other adverse effects of chemical substances and other adverse effects of injury) N800-N999 95,619 1,170 2,461 3,672 16,368 12,719 8,453 9,820 12,439 13,683 10,286 Fractures, dislocations, and sprains and strains of joints		2
Intracranial injury (excluding those with skull fracture)		
Internal injury of chest, abdomen, and pelvis N860–N869 31,385 85 482 1,165 9,400 6,473 3,784 3,302 2,813 2,138 1,338 Laceration and open wound N870–N907 31,370 22 147 713 7,841 7,767 4,790 3,776 2,939 1,979 1,020 Superficial injury and contusion N910–N929 2,145 20 64 106 430 265 164 168 175 217 318 Effects of foreign body, entering through orifice N930–N939 12,172 494 322 275 696 643 746 1,200 1,772 2,280 2,360 Burn N940–N949 6,550 137 510 398 901 808 614 699 732 766 661 njury to nerves and spinal cord N950–N959 2,700 5 33 115 647 500 281 269 312 261 201 Adverse effects of chemical substances and other adverse effects of chemical substances and other adverse effects N960–N999 95,619 1,170 2,461 3,672 16,368 12,719 8,453 9,820 12,439 13,683 10,286 Male Accidents, poisonings, and violence (Nature of Injury) N800–N999 139,891 1,207 2,579 4,816 30,156 22,487 13,811 13,961 15,583 15,747 12,615 Fractures, dislocations, and sprains and strains of joints	11,577	
Laceration and open wound	1,246	
Superficial injury and contusion . N910-N929 2,145 20 64 106 430 265 164 168 175 217 318 Effects of foreign body, entering through orifice . N930-N939 12,172 494 322 275 696 643 746 1,200 1,772 2,280 2,360 Burn	379	
Effects of foreign body, entering through orifice . N930–N939 12,172 494 322 275 696 643 746 1,200 1,772 2,280 2,360 Burn	344	
Burn	217	
Injury to nerves and spinal cord	1,381	
Adverse effects of chemical substances and other adverse effects of chemical substances and other adverse effects	314	
Male Accidents, poisonings, and violence (Nature of Injury) N800–N999 139,891 1,207 2,579 4,816 30,156 22,487 13,811 13,961 15,583 15,747 12,615 Fractures, dislocations, and sprains and strains of joints	73	
Accidents, poisonings, and violence (Nature of Injury)	4,407	1
(Nature of Injury)		
Fractures, dislocations, and sprains and strains of joints	0.700	
	6,730	1
and adjacent muscles N800-N848 26,997 99 346 1,001 6,053 3,506 1,981 1,929 2,049 2,604 3,877 Intracranial injury	3,532	
(excluding those with skull fracture) N850-N854 37,654 215 657 1,506 10,811 7,039 4,108 3,916 3,653 2,981 1,999	714	
Internal injury of chest, abdomen, and pelvis N860–N869 24,005 45 286 786 7,473 5,229 2,979 2,543 2,085 1,470 854	233	
Laceration and open wound N870-N907 25,434 13 94 513 6,496 6,436 3,783 3,040 2,383 1,609 803	239	
Superficial injury and contusion N910-N929 1,344 12 33 63 321 191 117 118 126 119 154	89	
Effects of foreign body, entering through orifice N930–N939 7,473 281 192 154 497 452 498 808 1,195 1,450 1,279	665	
Burn	154	
Injury to nerves and spinal cord N950–N959 2,042 3 17 81 491 398 217 213 249 197 126 Adverse effects of chemical substances and other	47	
adverse effects	1,990	1
Female		
Accidents, poisonings, and violence		
(Nature of Injury)	11,305	
Fractures, dislocations, and sprains and strains of joints and adjacent muscles	8,045	
Intracranial injury	•	
(excluding those with skull fracture) N850-N854 12,143 181 466 755 2,922 1,704 1,213 1,113 1,103 1,095 1,050	532	
Internal injury of chest, abdomen, and pelvis N860-N869 7,380 40 196 379 1,927 1,244 805 759 728 668 484	146	
Laceration and open wound	105	
Superficial injury and contusion	128	
Effects of foreign body, entering through orifice N930–N939 4,699 213 130 121 199 191 248 392 577 830 1,081	716	
Burn	160	
Injury to nerves and spinal cord N950–N959 658 2 16 34 156 102 64 56 63 64 75 Adverse effects of chemical substances and other	160	
adverse effects	26 2,417	

Technical notes

Nature and sources of data

Data shown in this publication are based on information from all death certificates from the 50 States and the District of Columbia. Mortality statistics are based on information obtained directly from copies of the original certificates received from the State registration offices and from precoded data provided to the National Center for Health Statistics (NCHS) by States through the Vital Statistics Cooperative Program.

Thirty-six States participated in the cooperative program in 1978. Seven of the participating States submitted all mortality data, including cause of death, precoded on computer tape. The remaining 29 States in the program submitted computer tapes with precoded demographic data, along with copies of the original certificates from which NCHS coded the cause of death. For the balance of States, which were not participants in the cooperative system in 1978, NCHS coded both the demographic and the medical data directly from copies of the original death certificates.

Data for the United States as a whole refer to events occurring to residents within the United States.

Cause-of-death classification

The mortality statistics presented here are compiled in accordance with the World Health Organization regulations, which specify that member nations classify causes of death in accordance with the current revision of the International Statistical Classification of Diseases, Injuries, and Causes of Death. An adaptation of the Eighth Revision of the classification-Eighth Revision International Classification of Diseases, Adapted for Use in the United States-was used for the years 1968-78. Besides specifying the classification, the World Health Organization regulations outline the form of medical certification and the coding procedures to be used. Underlying cause-of-death data and reported cause-of-death data presented in this publication were coded by procedures outlined in Vital Statistics Instruction Manuals issued by NCHS, Part 2a, "Classifying the Underlying Cause of Death" and Part 2b, "Medical Classification of Death Records for Automated Classification of Medical Entities."

National coding of more than a single cause of death has been undertaken previously by NCHS and its predecessor agencies. The underlying cause and one associated cause were coded in 1917, 1925, 1936, and 1940. In 1955 up to five additional causes were coded. The multiple cause-of-death statistics in this publication are the first national multiple cause data to be published by the United States since publication of the 1955 data. These data are now being prepared routinely, using automated procedures, and are being made available annually.

Multiple cause-of-death data allow for the maximum use of medical information reported on the death certificate. The medical information reported on the death certificate depends on the certifying physician's evaluation of the part each condition played in the death and on the physician's desire and ability to report that evaluation on the death certificate. It is reasonable to assume that not all conditions present at death are reported on the death certificate and that the variation in reporting depends on a number of factors, including the nature of the conditions that brought about death. For example, if a person with diabetes dies in a motor vehicle accident, the certifying physician may not report Diabetes mellitus on the death certificate.

Population bases for computing rates

The population for computing rates shown in this publication represents the population residing in the specified area, as published by the U.S. Bureau of the Census in *Current Population Reports*, "Population Estimates and Projections," Series P-25, No. 917. These populations are revised estimates based on the 1980 Census enumeration.

Cause-of-death ranking

The cause-of-death rankings shown in this report are based on the List of 69 Selected Causes of Death. This list is adapted from one of the special lists for mortality tabulations recommended by the World Health Organization for use with the Eighth Revision International Classification of Diseases.

The cause-of-death categories introduced by NCHS during the period of the Eighth Revision are not included in the List of 69 Selected Causes and therefore are not ranked among the leading causes of death. These categories include Chronic obstructive lung disease without mention of asthma, bronchitis, or emphysema (ICDA No. *519.3²).

The group titles Major cardiovascular diseases and Symptoms and ill-defined conditions are not ranked from the List of 69 Selected Causes. Also, category titles that begin with the words "Other" or "All other" are not ranked to determine the leading causes of death. When one of the titles is ranked (for example, Tuberculosis, all forms), its components (in this case, Tuberculosis of respiratory system and Tuberculosis, other forms) are not ranked.

Nature of injury titles are not included in the List of 69 Selected Causes and are therefore not included in the leading causes based on reported causes.

²The asterisk before category number 519.3 indicates that the title to which this number corresponds was not included in ICDA but was introduced by the National Center for Health Statistics.

Symbols

- - Data not available
- . . . Category not applicable
- Quantity zero
- 0.0 Quantity more than zero but less than 0.05
- Z Quantity more than zero but less than 500 where numbers are rounded to thousands
- * Figure does not meet standards of reliability or precision

Suggested Citation

National Center for Health Statistics: Multiple causes of death in the United States. *Monthly Vital Statistics Report.* Vol. 32-No. 10, Supp. (2) DHHS Pub No. (PHS) 84-1120. Public Health Service, Hyattsville, Md., Feb. 17, 1984.

Copyright Information

This report may be reprinted without further permission.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES Public Health Service National Center for Health Statistics 3700 East-West Highway Hyattsville, Maryland 20782

OFFICIAL BUSINESS PENALTY FOR PRIVATE USE, \$300

To receive this publication regularly, contact the National Center for Health Statistics by calling 301 436–NCHS.

FIRST CLASS MAIL POSTAGE & FEES PAID PHS PERMIT No. G29