

2016 National Study of Long-Term Care Providers

Web Tables of State Estimates on Residential Care Community Residents

Supplement to: *Caffrey C, Sengupta M. Variation in residential care community resident characteristics, by size of community: United States, 2016. NCHS Data Brief, no 299. Hyattsville, MD: National Center for Health Statistics. 2018. Available from: <https://www.cdc.gov/nchs/data/databriefs/db299.htm>.*

About the 2016 State Estimate Web Tables Supplement to NCHS Data Brief No. 299

The Long-Term Care Statistics Branch is pleased to release state estimate Web tables on residential care community residents, using data from the third wave of the National Study of Long-Term Care Providers (NSLTCP) conducted in 2016 by the National Center for Health Statistics (NCHS). **These state Web tables on residential care community residents complement Data Brief 299.** Table 1 provides state estimates of the characteristics presented in Data Brief 299, and Table 2 provides the standard errors for those estimates.

Estimates are presented as percentages. Data are presented in the state Web tables only if they meet NCHS confidentiality and reliability standards. Footnotes are included in the state Web tables to indicate estimates that do not meet these confidentiality or reliability standards. The data sources and methodology used to develop the state Web tables are the same as those used for Data Brief 299. For more information on the data source and methodology, see the “Data source and methods” section in Data Brief 299.

About NSLTCP

NSLTCP provides reliable, accurate, relevant, and timely statistical information to support and inform long-term care services policy, research, and practice. The main goals of NSLTCP are to: (1) estimate the supply, provision, and use of paid, regulated long-term care services in five major sectors; (2) estimate key policy-relevant characteristics of providers and service users, and provider practices; (3) produce national and state estimates, where feasible; (4) compare estimates among sectors; and (5) monitor trends over time.

Information on NSLTCP and additional products is available from:

<https://www.cdc.gov/nchs/nsltcp.htm>.

NSLTCP comprises two components: (1) survey data collected by NCHS on assisted living and similar residential care communities and residents, and adult day services centers and participants; and (2) administrative data on nursing homes and residents, home health agencies and patients, and hospices and patients obtained from the Centers for Medicare & Medicaid Services. The restricted 2016 survey data files for the residential care community and adult day services center sectors are available through the NCHS Research Data Center (https://www.cdc.gov/nchs/nsltcp/nsltcp_rdc.htm).

The residential care community survey was conducted between August 2016 and February 2017. NSLTCP uses a multimode survey protocol with mail, Web, and computer-assisted telephone follow-up for nonresponse.

The questionnaires included survey items on provider characteristics such as ownership, size, services offered, selected practices, and staffing, in addition to aggregate service-user characteristics such as age, sex, race, and the number of residents needing assistance with activities of daily living. Two versions of the questionnaire were used, each containing: (1) survey items included on both questionnaires that were asked of all respondents, which can be used to produce national and state estimates, and (2) survey items unique to each questionnaire that can be used to produce national estimates only. The current document includes variables that can be estimated at the state level.

RTI International collected the survey data under contract to NCHS. The 2016 residential care community mail questionnaire and survey documentation are available from:

https://www.cdc.gov/nchs/nsltcp/nsltcp_questionnaires.htm.

The residential care community sampling frame was constructed from lists of licensed residential care communities obtained from the licensing agencies in each of the 50 states and the District of Columbia. A sample of 11,688 residential care communities was selected from the sampling frame of 42,149 communities. Of the 8,626 in-scope and presumed in-scope residential care communities, 4,578 completed the survey questionnaire, for a weighted response rate (for differential probabilities of selection) of 50.7% (calculated using the American Association for Public Opinion Research [AAPOR] Response Rate 4).

For further information, contact the Long-Term Care Statistics Branch at ltcsbfeedback@cdc.gov or (301) 458-4747.

Table 1. State estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Under age 65	Aged 65–74	Aged 75–84	Aged 85 and over	Non-Hispanic white	Female	Some or all services paid by Medicaid in last 30 days
	Percent						
All RCCs ¹	7.0	11.0	30.0	52.0	81.0	71.0	17.0
Alabama	1.3	7.8	40.5	50.4	93.7	78.4	3.8
Alaska	11.0	13.3	31.4	44.7	73.7	63.2	40.3
Arizona	3.6	11.1	31.3	54.0	83.0	69.1	15.0
Arkansas	9.4	14.0	31.2	45.3	92.9	72.3	25.2
California	3.0	10.4	31.9	54.8	67.9	71.2	*3.0
Colorado	9.6	14.1	31.4	44.9	74.2	71.7	25.2
Connecticut	37.3	20.0	20.8	21.9	80.9	56.6	71.3
Delaware	3.8	11.7	24.9	59.6	91.3	67.0	9.8
District of Columbia	*	*	*	*	*	*	*
Florida	8.6	11.8	32.3	47.3	79.3	67.2	20.2
Georgia	6.6	13.7	35.6	44.1	85.4	72.4	8.5
Hawaii	3.9	7.7	20.6	67.8	16.7	66.9	6.8
Idaho	8.0	13.0	30.9	48.4	86.6	72.3	32.5
Illinois	1.3	10.0	33.6	55.1	90.5	70.6	14.4
Indiana	5.4	10.0	28.2	54.3	89.8	69.8	14.3
Iowa	23.8	17.3	18.2	40.6	94.6	58.6	21.7
Kansas	3.2	8.7	25.0	63.1	88.8	71.9	28.4
Kentucky	11.5	13.3	29.7	45.5	87.6	70.5	11.5
Louisiana	2.1	7.3	35.9	54.7	94.8	75.0	0.0
Maine	8.4	13.0	31.3	47.3	92.5	69.5	56.4
Maryland	5.9	15.7	29.0	49.3	79.1	68.1	5.2
Massachusetts	4.7	9.5	27.5	58.3	86.7	74.0	10.7
Michigan	9.6	9.6	34.5	46.3	87.7	69.6	17.7
Minnesota	4.8	11.1	33.2	51.0	80.1	72.4	20.4
Mississippi	7.1	11.5	38.7	42.7	88.4	77.8	11.6
Missouri	17.6	12.9	29.3	40.3	87.1	66.8	23.7
Montana	4.8	6.8	23.4	63.0	51.9	71.7	16.2
Nebraska	4.5	10.5	30.4	54.7	85.4	71.9	21.8
Nevada	7.1	11.0	33.1	48.9	78.0	69.5	9.3
New Hampshire	4.0	8.8	25.7	61.4	84.7	72.3	12.2
New Jersey	1.6	6.0	25.7	65.9	83.9	76.5	16.9
New Mexico	5.4	10.4	36.0	48.1	80.9	69.2	8.2
New York	6.7	12.8	29.2	51.4	73.1	69.3	23.4
North Carolina	18.3	15.5	25.9	40.3	76.8	68.8	44.1
North Dakota	4.8	7.5	29.8	57.9	93.5	71.7	22.4
Ohio	2.7	9.8	30.8	56.6	90.6	74.0	13.5
Oklahoma	2.7	7.7	30.9	58.2	86.4	75.7	5.8
Oregon	8.5	15.0	26.1	50.5	74.4	68.2	44.6
Pennsylvania	5.0	7.8	23.3	63.9	96.7	69.9	*4.0
Rhode Island	6.9	10.3	21.2	61.6	66.2	72.3	15.3
South Carolina	10.0	8.1	29.5	52.6	89.5	71.4	15.1
South Dakota	6.4	8.4	26.6	58.6	89.4	68.4	22.5
Tennessee	3.0	10.4	32.8	53.8	94.0	73.8	5.1
Texas	13.2	13.2	27.1	46.4	86.1	66.5	15.4
Utah	5.7	9.4	30.0	54.9	94.0	73.8	18.6
Vermont	4.5	9.5	24.7	61.3	95.3	74.6	35.7
Virginia	7.4	11.2	32.0	49.5	76.8	71.2	6.9
Washington	5.2	8.5	36.6	49.7	64.7	73.0	27.6
West Virginia	3.7	13.1	25.5	57.7	82.6	76.1	1.0
Wisconsin	9.9	11.2	25.1	53.7	91.8	68.3	20.0
Wyoming	6.7	14.6	36.5	42.2	86.7	70.0	18.8

See footnotes at end of table.

Table 1. State estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Diagnosed with diabetes ²	Diagnosed with		
		Alzheimer disease or other dementias	Depression ²	Heart disease ^{2,3}
Percent				
All RCCs ¹	18.0	42.0	31.0	34.0
Alabama	16.0	35.4	27.5	33.4
Alaska	25.9	60.0	39.1	34.8
Arizona	16.3	56.7	35.9	33.4
Arkansas	17.4	41.9	27.3	27.8
California	15.2	35.3	23.4	31.1
Colorado	17.5	47.8	35.9	28.5
Connecticut	21.9	11.8	36.9	25.6
Delaware	19.1	37.1	27.5	42.7
District of Columbia	*	*	*	*
Florida	16.1	45.6	31.5	33.8
Georgia	18.3	47.9	29.7	27.7
Hawaii	19.5	56.9	23.4	27.1
Idaho	18.3	48.4	38.4	28.2
Illinois	17.7	43.4	28.7	37.5
Indiana	20.8	33.1	34.4	37.2
Iowa	21.8	25.3	32.6	36.5
Kansas	23.2	43.7	45.2	42.6
Kentucky	20.8	30.6	29.4	27.2
Louisiana	13.9	32.5	24.8	29.1
Maine	25.6	44.8	40.4	37.2
Maryland	17.1	53.7	35.5	40.6
Massachusetts	14.5	37.4	26.6	35.8
Michigan	20.0	43.3	29.1	30.7
Minnesota	15.1	33.2	25.0	24.2
Mississippi	14.2	36.5	27.3	32.1
Missouri	20.5	34.8	36.2	35.7
Montana	14.0	45.0	37.2	28.4
Nebraska	18.1	35.0	28.7	36.9
Nevada	12.3	38.6	26.9	31.5
New Hampshire	14.5	48.2	37.4	39.5
New Jersey	18.4	42.4	29.3	41.0
New Mexico	15.7	50.2	31.8	33.5
New York	23.1	31.8	25.0	38.9
North Carolina	24.5	40.4	31.4	32.9
North Dakota	20.2	29.3	30.0	32.8
Ohio	20.5	43.7	27.8	36.4
Oklahoma	17.2	43.2	33.7	35.9
Oregon	20.6	58.4	39.1	39.4
Pennsylvania	20.9	40.3	35.8	45.0
Rhode Island	16.1	33.0	30.5	31.9
South Carolina	15.8	42.8	29.7	34.2
South Dakota	17.3	28.8	28.7	37.6
Tennessee	18.2	48.9	28.9	36.6
Texas	19.0	44.5	38.8	30.8
Utah	17.0	33.3	27.5	27.4
Vermont	15.1	41.9	37.1	39.1
Virginia	18.3	41.2	28.9	32.0
Washington	15.1	54.5	35.3	36.3
West Virginia	22.4	47.5	30.9	49.4
Wisconsin	20.5	38.4	33.0	35.6
Wyoming	16.3	28.9	21.9	26.2

See footnotes at end of table.

Table 1. State estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Needed assistance with bathing	Needed assistance with walking or locomotion	Needed assistance with dressing	Needed assistance with toileting	Needed assistance with transferring in or out of bed or chair	Needed assistance with eating
All RCCs ¹	64.0	56.5	48.0	40.0	29.0	19.0
	Percent					
Alabama	59.4	66.7	42.7	28.1	22.1	12.2
Alaska	75.6	73.3	69.2	63.3	53.5	52.0
Arizona	84.0	61.6	67.9	60.6	43.3	32.1
Arkansas	60.9	50.0	39.6	22.3	14.3	10.8
California	56.5	55.5	47.7	44.1	30.9	19.5
Colorado	54.2	51.2	40.4	33.8	21.3	17.7
Connecticut	32.8	31.6	6.9	3.1	1.1	1.7
Delaware	62.0	61.9	43.0	33.3	26.0	13.2
District of Columbia	*	*	*	*	*	*
Florida	67.2	57.3	52.0	45.0	31.4	17.6
Georgia	64.1	56.2	46.9	37.8	21.2	14.3
Hawaii	71.1	58.3	63.3	51.9	36.1	28.0
Idaho	64.3	62.3	46.1	45.7	37.9	28.6
Illinois	60.6	62.4	47.5	40.6	30.8	18.9
Indiana	48.9	52.2	35.4	26.6	23.5	11.6
Iowa	53.0	37.9	19.6	16.0	7.2	9.7
Kansas	75.9	66.1	60.2	49.4	45.8	23.8
Kentucky	53.7	47.4	36.6	28.0	23.1	14.2
Louisiana	39.3	54.2	34.8	25.6	18.0	10.7
Maine	70.6	55.6	53.7	42.4	33.4	26.5
Maryland	81.1	65.8	66.9	60.1	45.5	33.2
Massachusetts	57.1	44.9	43.2	23.6	14.5	9.6
Michigan	77.0	65.8	57.3	55.5	43.8	31.4
Minnesota	52.5	47.6	37.4	30.8	22.6	17.2
Mississippi	59.3	56.6	43.7	36.4	17.7	13.3
Missouri	54.7	32.0	23.5	16.6	6.1	8.8
Montana	64.1	51.4	40.1	33.6	27.3	16.5
Nebraska	68.6	55.4	34.1	24.3	17.6	11.7
Nevada	57.4	58.3	50.2	40.2	33.2	16.5
New Hampshire	64.0	56.9	49.7	41.3	27.4	18.6
New Jersey	59.9	61.8	50.8	40.9	30.8	15.7
New Mexico	68.7	59.0	57.8	46.4	41.7	23.5
New York	67.5	52.5	41.6	24.4	12.3	9.9
North Carolina	81.3	58.2	66.9	57.4	34.6	24.6
North Dakota	57.4	58.1	32.3	20.4	9.8	10.8
Ohio	66.8	56.2	51.8	42.0	34.8	20.9
Oklahoma	49.0	53.6	32.9	25.9	23.9	12.5
Oregon	76.4	63.9	60.0	52.3	40.9	36.4
Pennsylvania	73.0	59.5	51.8	41.5	33.5	18.4
Rhode Island	63.1	54.0	47.9	25.2	15.9	8.3
South Carolina	58.0	42.3	44.3	32.2	16.9	9.9
South Dakota	60.9	49.2	29.1	20.0	17.8	8.3
Tennessee	60.7	54.5	46.9	37.0	20.5	17.2
Texas	57.6	66.0	45.6	36.6	30.3	18.1
Utah	49.4	52.5	41.7	35.3	32.7	14.4
Vermont	68.7	65.2	45.3	36.9	19.5	18.4
Virginia	56.5	55.4	42.8	33.1	19.6	14.8
Washington	55.6	50.3	46.5	39.0	28.6	23.3
West Virginia	73.9	59.2	51.4	47.2	31.0	16.7
Wisconsin	68.4	56.4	45.4	37.2	30.1	20.9
Wyoming	49.2	63.2	28.1	13.6	8.5	10.3

See footnotes at end of table.

Table 1. State estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Emergency department visit in last 90 days	Percent	Discharge from overnight hospital stay in last 90 days
All RCCs ¹	14.0		8.0
Alabama	14.6		9.7
Alaska	16.3		8.6
Arizona	10.7		6.6
Arkansas	12.8		6.8
California	14.8		8.3
Colorado	14.2		7.4
Connecticut	17.1		8.1
Delaware	19.7		11.1
District of Columbia	*		*
Florida	16.5		8.1
Georgia	13.8		7.0
Hawaii	*13		7.4
Idaho	13.6		7.3
Illinois	13.4		8.8
Indiana	11.2		6.8
Iowa	7.4		7.8
Kansas	10.1		9.1
Kentucky	12.3		7.2
Louisiana	11.2		7.2
Maine	12.3		7.3
Maryland	13.6		6.4
Massachusetts	18.5		11.4
Michigan	18.0		11.9
Minnesota	9.7		7.0
Mississippi	10.9		6.3
Missouri	13.7		10.2
Montana	11.1		5.9
Nebraska	10.5		6.3
Nevada	13.4		6.2
New Hampshire	13.1		8.2
New Jersey	15.8		9.1
New Mexico	13.8		7.4
New York	20.0		11.6
North Carolina	16.4		9.2
North Dakota	9.5		6.6
Ohio	12.0		8.0
Oklahoma	12.4		7.0
Oregon	12.5		5.8
Pennsylvania	17.0		10.5
Rhode Island	18.3		10.7
South Carolina	13.4		8.4
South Dakota	12.9		8.0
Tennessee	16.2		11.9
Texas	14.2		10.1
Utah	11.0		5.7
Vermont	12.8		6.9
Virginia	14.8		6.9
Washington	13.2		7.5
West Virginia	15.4		9.1
Wisconsin	11.7		5.8
Wyoming	14.6		9.5

* Estimate is not presented because it does not meet confidentiality or reliability standards. Estimates with an asterisk have a sample size greater than 30 but a relative standard error greater than 30%.

¹ Residential care communities.

² Item nonresponse (weighted) was higher than 10%: 13% of RCCs did not report number of residents with depression, diabetes, and heart disease.

³ Changes in question wording may have contributed to a difference in the estimates from earlier National Study of Long-Term Care Providers surveys.

NOTES: Cases with missing data were excluded on a variable-by-variable basis. Variables are identified in a footnote if cases with missing data were more than 10% of the weighted total.

SOURCE: NCHS, National Study of Long-Term Care Providers, 2016.

Table 2. Standard errors for state estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Under age 65	Aged 65–74	Aged 75–84	Aged 85 and over	Non-Hispanic white	Female	Some or all services paid by Medicaid in last 30 days
All RCCs ¹	0.3	0.3	0.5	0.7	0.5	0.3	0.6
Alabama	0.2	0.9	1.9	2.5	1.1	0.7	2.8
Alaska	1.5	1.0	2.6	3.3	3.7	2.6	2.5
Arizona	1.1	1.8	3.4	4.3	4.1	1.9	4.0
Arkansas	1.6	1.3	1.2	1.9	1.3	1.0	2.3
California	0.6	1.2	1.8	2.5	3.9	0.9	*1.3
Colorado	2.5	2.8	2.7	3.6	5.5	1.7	6.0
Connecticut	3.4	1.5	1.9	2.8	3.3	2.4	4.0
Delaware	1.4	3.9	2.3	4.2	1.7	5.0	2.9
District of Columbia	*	*	*	*	*	*	*
Florida	2.4	1.6	2.5	3.1	2.6	1.5	2.6
Georgia	0.8	1.6	2.6	3.4	3.4	2.4	1.6
Hawaii	1.1	1.6	4.0	4.7	2.0	3.8	1.9
Idaho	1.0	1.2	1.4	2.4	3.6	0.8	2.4
Illinois	0.4	1.8	2.4	3.3	3.1	1.6	3.8
Indiana	1.5	1.0	1.7	2.4	1.8	1.3	1.8
Iowa	3.6	3.5	1.5	6.1	1.4	5.2	2.9
Kansas	0.8	1.1	2.9	3.8	2.9	1.2	4.1
Kentucky	1.4	1.3	1.4	2.4	2.8	1.0	1.6
Louisiana	0.4	1.3	2.0	2.5	1.4	0.8	0.0
Maine	0.9	0.9	1.4	2.0	1.8	1.3	3.4
Maryland	1.3	2.8	2.6	4.7	4.1	2.3	1.5
Massachusetts	0.8	0.8	1.3	1.8	2.2	0.7	1.6
Michigan	1.2	2.0	3.1	3.4	3.2	1.4	4.0
Minnesota	1.1	2.5	1.9	3.7	5.2	1.8	2.6
Mississippi	1.5	1.5	2.5	2.5	4.1	1.1	2.9
Missouri	4.1	2.0	3.1	4.2	3.8	2.6	4.8
Montana	0.6	0.9	1.9	2.8	4.3	0.7	2.5
Nebraska	0.7	1.2	1.5	2.0	3.0	0.8	2.1
Nevada	2.2	1.6	3.3	4.1	4.0	1.8	0.9
New Hampshire	0.6	1.3	1.8	2.8	4.0	1.1	1.4
New Jersey	0.2	0.5	1.3	1.5	2.2	0.5	1.3
New Mexico	2.0	1.6	2.3	3.5	1.7	2.5	2.2
New York	1.5	1.9	1.7	3.0	5.2	1.2	4.6
North Carolina	3.2	1.7	1.7	3.8	3.3	2.4	4.4
North Dakota	0.5	0.5	1.7	2.1	1.6	0.9	2.0
Ohio	0.4	1.3	2.0	2.5	2.6	0.6	1.9
Oklahoma	0.5	0.8	1.7	2.2	1.8	0.7	2.2
Oregon	1.1	1.8	2.0	3.5	6.1	2.4	3.8
Pennsylvania	1.0	1.3	2.1	2.8	1.3	1.4	*2.2
Rhode Island	1.7	1.1	2.3	3.3	8.1	1.7	3.0
South Carolina	1.9	0.8	2.4	3.1	1.7	1.7	2.7
South Dakota	1.8	1.3	1.1	2.6	2.4	1.4	4.1
Tennessee	0.3	0.5	1.0	1.2	0.8	0.5	0.7
Texas	2.9	2.9	3.3	4.5	2.1	2.6	4.5
Utah	0.8	0.7	1.8	2.3	1.0	0.8	1.3
Vermont	0.7	1.3	2.1	2.4	1.7	1.1	4.0
Virginia	1.7	1.8	2.0	3.4	6.3	1.5	1.9
Washington	1.0	1.0	3.1	3.6	6.3	2.0	4.6
West Virginia	1.0	4.4	2.9	5.0	6.8	1.4	0.5
Wisconsin	1.2	1.8	1.6	2.6	3.2	1.3	2.9
Wyoming	1.8	2.6	2.9	3.1	1.6	0.9	1.8

See footnotes at end of table.

Table 2. Standard errors for state estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Diagnosed with diabetes ²	Diagnosed with Alzheimer disease or other dementias	Diagnosed with depression ²	Diagnosed with heart disease ^{2,3}
Standard error				
All RCCs ¹	0.4	0.7	0.6	0.6
Alabama	0.8	2.9	1.7	1.9
Alaska	1.6	4.3	2.9	3.3
Arizona	1.7	3.5	4.7	3.1
Arkansas	0.9	2.0	1.5	1.9
California	1.4	2.9	2.2	2.5
Colorado	2.4	5.0	3.5	3.0
Connecticut	0.9	1.9	3.3	2.8
Delaware	3.7	4.0	4.9	3.3
District of Columbia	*	*	*	*
Florida	1.1	3.5	3.2	2.7
Georgia	1.3	3.4	5.0	3.0
Hawaii	5.2	7.0	5.4	4.2
Idaho	1.1	2.3	1.9	1.5
Illinois	1.5	3.8	3.6	3.0
Indiana	1.2	1.9	2.1	2.1
Iowa	2.6	2.6	4.1	5.6
Kansas	4.1	3.7	4.0	3.7
Kentucky	1.7	1.9	2.1	2.1
Louisiana	1.1	3.8	2.5	3.2
Maine	1.3	2.0	1.6	1.4
Maryland	1.2	3.8	2.7	5.4
Massachusetts	0.9	2.0	2.1	2.0
Michigan	1.9	3.1	2.2	4.1
Minnesota	1.4	3.8	3.3	2.2
Mississippi	0.9	2.9	2.9	3.5
Missouri	1.9	3.9	4.0	4.2
Montana	1.2	3.9	5.0	2.5
Nebraska	1.0	2.3	1.9	1.6
Nevada	1.4	4.0	3.2	5.4
New Hampshire	1.1	3.7	2.3	3.0
New Jersey	0.9	1.9	2.0	2.0
New Mexico	1.2	3.3	3.0	3.4
New York	2.1	3.3	3.5	3.6
North Carolina	3.0	3.0	4.1	3.5
North Dakota	0.8	2.1	1.5	1.8
Ohio	1.2	2.2	2.3	2.1
Oklahoma	1.2	2.6	3.0	2.5
Oregon	1.6	5.8	4.1	3.8
Pennsylvania	2.0	2.4	2.7	3.4
Rhode Island	1.9	3.2	3.6	4.8
South Carolina	1.0	3.0	3.4	3.7
South Dakota	2.4	1.6	5.4	1.4
Tennessee	1.0	1.7	1.5	1.3
Texas	1.6	4.5	3.2	2.6
Utah	1.1	2.5	2.2	2.2
Vermont	1.0	2.3	2.5	3.2
Virginia	1.8	3.8	3.2	3.8
Washington	1.5	5.0	4.1	2.9
West Virginia	2.2	3.6	2.4	3.2
Wisconsin	2.1	2.9	2.6	2.7
Wyoming	1.7	4.7	4.3	2.8

See footnotes at end of table.

Table 2. Standard errors for state estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Needed assistance with bathing	Needed assistance with walking or locomotion	Needed assistance with dressing	Needed assistance with toileting	Needed assistance with transferring in or out of bed or chair	Needed assistance with eating
	Standard error					
All RCCs ¹	0.8	0.8	0.7	0.7	0.7	0.5
Alabama	2.5	2.1	2.2	2.6	2.4	2.1
Alaska	4.4	2.8	4.6	4.7	4.2	4.5
Arizona	2.5	4.2	2.8	3.8	3.2	3.0
Arkansas	1.9	2.5	2.0	1.6	1.2	1.3
California	3.4	3.1	3.1	3.0	2.5	1.8
Colorado	3.5	5.9	3.6	3.8	2.7	3.5
Connecticut	3.5	3.2	1.1	0.7	0.4	0.3
Delaware	3.0	4.3	5.0	2.9	5.6	1.3
District of Columbia	*	*	*	*	*	*
Florida	3.0	3.4	3.2	3.0	2.9	1.8
Georgia	3.1	4.4	2.7	2.6	3.1	1.9
Hawaii	5.2	9.1	6.8	4.5	5.6	3.8
Idaho	2.5	2.0	1.6	1.8	2.1	1.9
Illinois	3.6	3.7	3.4	4.0	3.9	2.9
Indiana	2.0	2.4	2.1	2.1	2.1	1.4
Iowa	6.9	7.6	2.5	2.3	1.1	1.0
Kansas	4.1	4.1	4.7	4.8	5.2	3.5
Kentucky	2.2	2.9	1.9	1.9	1.9	1.6
Louisiana	2.3	4.4	2.2	2.3	1.8	1.4
Maine	2.4	2.3	2.3	2.0	2.1	1.8
Maryland	3.5	4.4	3.6	3.2	4.2	3.9
Massachusetts	1.9	2.7	2.0	1.9	1.6	1.0
Michigan	3.9	5.3	3.4	4.3	4.2	2.7
Minnesota	4.3	3.2	3.7	3.6	3.0	2.5
Mississippi	4.2	4.2	4.7	3.9	2.7	2.2
Missouri	5.1	5.3	4.0	4.1	1.9	2.3
Montana	2.5	3.8	2.1	3.0	2.9	2.0
Nebraska	2.1	2.2	1.9	1.7	1.6	1.4
Nevada	3.9	3.2	4.1	3.9	3.1	1.3
New Hampshire	3.3	3.3	3.4	3.0	3.2	1.9
New Jersey	1.9	2.2	1.6	1.8	1.7	1.0
New Mexico	4.2	4.6	3.5	3.9	4.1	2.8
New York	4.0	5.0	3.8	3.0	2.9	1.5
North Carolina	2.5	3.9	2.8	3.0	3.6	3.0
North Dakota	2.7	2.2	1.7	1.9	1.4	1.2
Ohio	2.2	2.6	2.1	2.7	2.4	2.3
Oklahoma	2.4	2.8	2.4	2.5	2.4	1.8
Oregon	3.5	4.1	4.2	4.5	3.9	4.5
Pennsylvania	2.8	3.8	3.1	3.6	4.0	2.6
Rhode Island	3.8	3.9	5.7	4.3	3.8	2.0
South Carolina	3.5	4.1	3.5	2.7	1.9	1.3
South Dakota	2.6	2.5	2.0	2.0	2.5	1.0
Tennessee	1.4	1.7	1.4	1.6	1.5	1.2
Texas	4.2	4.0	4.6	3.8	3.6	2.9
Utah	2.2	2.8	2.1	2.0	2.5	1.2
Vermont	2.5	3.4	3.8	3.4	2.0	2.2
Virginia	5.4	5.1	4.4	3.9	2.9	2.6
Washington	4.9	5.8	3.4	4.5	3.7	3.2
West Virginia	3.4	4.3	3.3	4.2	3.3	2.2
Wisconsin	2.9	3.5	2.6	2.8	2.6	2.1
Wyoming	2.7	3.8	2.8	2.7	2.0	1.9

See footnotes at end of table.

Table 2. Standard errors for state estimates for NCHS Data Brief No. 299, Variation in Residential Care Community Resident Characteristics, by Size of Community: United States, 2016

Area	Emergency department visit in last 90 days	Discharge from overnight hospital stay in last 90 days
	Standard error	
All RCCs ¹	0.3	0.2
Alabama	0.9	1.0
Alaska	1.2	1.0
Arizona	1.0	0.9
Arkansas	0.5	0.4
California	1.3	0.9
Colorado	1.7	1.0
Connecticut	1.9	0.9
Delaware	3.9	2.1
District of Columbia	*	*
Florida	1.3	0.7
Georgia	1.2	0.8
Hawaii	*4.5	1.4
Idaho	2.2	1.1
Illinois	1.1	0.9
Indiana	0.6	0.5
Iowa	1.3	1.7
Kansas	1.1	2.0
Kentucky	0.8	0.7
Louisiana	0.8	0.9
Maine	0.6	0.5
Maryland	1.9	1.0
Massachusetts	1.1	1.0
Michigan	3.1	2.5
Minnesota	1.3	0.8
Mississippi	1.0	0.6
Missouri	1.2	1.7
Montana	0.9	0.4
Nebraska	0.6	0.3
Nevada	1.3	0.8
New Hampshire	0.6	0.6
New Jersey	0.9	0.6
New Mexico	1.7	0.8
New York	2.3	1.9
North Carolina	1.8	1.1
North Dakota	0.6	0.4
Ohio	0.9	0.7
Oklahoma	0.9	0.7
Oregon	1.0	0.7
Pennsylvania	1.4	0.9
Rhode Island	1.7	1.4
South Carolina	1.1	1.0
South Dakota	1.5	0.8
Tennessee	0.6	1.1
Texas	1.6	1.5
Utah	0.6	0.4
Vermont	1.4	0.8
Virginia	2.2	1.1
Washington	1.2	0.9
West Virginia	1.4	1.0
Wisconsin	0.8	0.7
Wyoming	1.5	0.9

* Estimate is not presented because it does not meet confidentiality or reliability standards. Estimates with an asterisk have a sample size greater than 30 but a relative standard error greater than 30%.

¹ Residential care communities.

² Item nonresponse (weighted) was higher than 10%: 13% of RCCs did not report number of residents with depression, diabetes, and heart disease.

³ Changes in question wording may have contributed to a difference in the estimates from earlier National Study of Long-Term Care Providers surveys.

NOTES: Cases with missing data were excluded on a variable-by-variable basis. Variables are identified in a footnote if cases with missing data were more than 10% of the weighted total.

SOURCE: NCHS, National Study of Long-Term Care Providers, 2016.