

VITAL and HEALTH STATISTICS

DATA FROM THE NATIONAL HEALTH SURVEY

PROPERTY OF THE
PUBLICATIONS BRANCH
EDITORIAL LIBRARY

Acute Conditions

incidence and associated disability

United States - July 1965 - June 1966

Statistics on the incidence of acute conditions and the associated days of restricted activity, bed disability, and time lost from work and school, by age, sex, calendar quarter, residence, and geographic region. Based on data collected in household interviews during the period July 1965-June 1966.

Washington, D. C.

June 1967

U.S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
John W. Gardner
Secretary

Public Health Service
William H. Stewart
Surgeon General



Public Health Service Publication No. 1000-Series 10- No. 38

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C., 20402 - Price 45 cents

NATIONAL CENTER FOR HEALTH STATISTICS

FORREST E. LINDER, PH.D., *Director*

THEODORE D. WOOLSEY, *Deputy Director*

OSWALD K. SAGEN, PH.D., *Assistant Director for State Relations*

WALT R. SIMMONS, M.A., *Statistical Advisor*

PHILIP S. LAWRENCE, Sc.D., *Planning Officer*

ALICE M. WATERHOUSE, M.D., *Medical Consultant*

JAMES E. KELLY, D.D.S., *Dental Advisor*

LOUIS R. STOLCIS, M.A., *Executive Officer*

DONALD GREEN, *Information Officer*

DIVISION OF HEALTH INTERVIEW STATISTICS

ELIJAH L. WHITE, AM., *Director*

ROBERT R. FUCHSBERG, *Chief Survey Methods Branch*

GERALDINE A. GLEESON, *Chief. Analysis and Reports Branch*

COOPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Health Survey, the Bureau of the Census, under a contractual arrangement, participates in most aspects of survey planning, selects the sample, collects the data, and carries out certain parts of the statistical processing.

Public Health Service Publication No. 1000-Series 10-No. 38

Library of Congress Catalog Card Number 65-60041

CONTENTS

	Page
Selected Findings-----	1
Source of Data-----	2
Condition Group, Sex, and Age-----	3
Geographic Distribution-----	7
Seasonal Variation: Calendar Quarter-----	9
Detailed Tables-----	10
Appendix I. Technical Notes on Methods-----	46
Background of This Report-----	46
Statistical Design of the Health Interview Survey-----	46
General Qualifications-----	47
Reliability of Estimates-----	47
Appendix II. Definitions of Certain Terms Used in This Report-----	56
Terms Relating to Conditions-----	56
Terms Relating to Disability-----	57
Demographic Terms-----	58
Appendix III. Condition Pages From Questionnaire-----	60

IN THIS REPORT statistics are presented on the annual incidence of acute illnesses and injuries, involving medical attention and/or reduced daily activity, which occurred during the period July 1965-June 1966 among the civilian, noninstitutional population of the United States. Estimates of disability days of restricted activity, bed-stay, and time lost from work and school associated with these acute conditions are also presented. These statistics, issued annually, are based on data collected in household interviews for the Health Interview Survey.

The information presented in this report is distributed by age, sex, residence, geographic region, and calendar quarter.

During July 1965-June 1966 an estimated 404.4 million acute conditions occurred in the population, an incidence rate of 212.0 conditions per 100 persons per year. This rate was slightly below that for the year ending in June 1965—212.7 per 100 persons. An epidemic of influenza-like illnesses occurred during the current year, totaling some 83.3 million conditions. The incidence rate for these conditions was 43.7 per 100 persons compared with a rate of 33.9 for the previous year. This increase was offset to some extent by a decline in reporting of injuries and of measles. A decline in the rate of injury from 29.9 to 25.4 per 100 persons for the current year was unusual and may be an artifact. The incidence rate of measles dropped from 3.6 per 100 persons to 1.5 per 100 persons per year.

SYMBOLS

Data not available-----	---
Category not applicable-----	...
Quantity zero-----	-
Quantity more than 0 but less than 0.05----	0.0
Figure does not meet standards of reliability or precision-----	*

ACUTE CONDITIONS

INCIDENCE AND ASSOCIATED DISABILITY

Charles S. Wilder, *Division of Health Interview Statistics*

SELECTED FINDINGS

During the year ending in June 1966, the incidence of acute illness and injury involving medical attention or reduced daily activity was an estimated 404.4 million conditions, or an annual rate of 212.0 conditions per 100 persons in the civilian, noninstitutional population of the United States. This incidence rate was slightly below that of 212.7 conditions per 100 persons during July 1964-June 1965.

The incidence rate for respiratory illness was 125.9 per 100 persons during the 12 months ending in June 1966, compared with a rate of 116.4 during the previous year. Contributing to the increased incidence of respiratory illness was an epidemic of influenza-like illnesses totaling 83.3 million conditions. The incidence rate for conditions reported as influenza was 43.7 per 100 persons, about 30 percent higher than the rate of 33.9 per 100 persons during the year ending in June 1965.

During the current year there was a reduction in reporting of injuries involving activity restriction or medical attention. An estimated 48.4 million injuries occurred, or 25.4 injuries per 100 persons. This rate represents a drop of 15 percent from the rate of injury of 29.9 for the preceding year. This decline in rate may be an artifact due to change in questionnaire format, variations in reporting, or unusual sampling variability (see explanation on pages 3 and 5).

An estimated 2.9 million cases of measles occurred during the current year, or 1.5 cases per 100 persons. The incidence of measles has dropped sharply from the rate of 3.6 per 100 persons during July 1964-June 1965. Undoubtedly this decline is related to the success of measles vaccine programs.

During July 1965-June 1966, acute illnesses or injuries caused an average of 8.2 days of restricted activity per person in the population, including 3.7 days spent in bed. Acute conditions were associated with an average of 4.6 days lost from school among children aged 6-16 years and 3.7 days lost from work among the currently employed population. (A currently employed person is one who had a job or business or worked at any time during the 2-week period prior to the week of interview.)

Respiratory illness accounted for about half of all acute conditions. These illnesses were the leading cause of disability days, causing 48 percent of all days of restricted activity, about 54 percent of days spent in bed, about 62 percent of days lost from school, and 44 percent of days lost from work.

Persons living in metropolitan areas had an incidence rate of 216.0 acute illnesses or injuries per 100 persons during July 1965-June 1966. This rate was slightly below the rate of 217.9 reported for the previous year. Residents of nonfarm areas outside of metropolitan areas reported 209.0 conditions per 100 persons, the same rate as that

for the preceding year. Farm residents in non-metropolitan areas experienced about 183.5 conditions per 100 persons, somewhat higher than the incidence rate of 175.6 for the year ending in June 1965.

The annual incidence rate for the current year was highest in the West Region (235.9 per 100 persons) and lowest in the Northeast Region (202.7 per 100 persons). Persons living in the North Central and South Regions experienced similar rates, 208.8 and 209.8, respectively. The comparable rates for July 1964-June 1965 were as follows: Northeast—210.2, North Central—212.8, South—201.4, and West—237.6.

SOURCE OF DATA

The information in this publication is derived from household interviews conducted by the Health Interview Survey in cooperation with the U.S. Bureau of the Census in a probability sample of the civilian, noninstitutional population of the United States. The sample is designed so that interviews are conducted during every week of the

year. During the 52-week period from July 1965-June 1966, the sample was composed of approximately 42,000 households containing about 134,000 persons living at the time of the interview.

A description of the design of the survey, of the methods used in estimation, and of general qualifications of the data obtained from surveys is presented in Appendix I. Since estimates shown in this report are based on a sample of the population rather than on the entire population, they are subject to sampling error. Therefore, particular attention should be paid to the section entitled "Reliability of Estimates." Sampling errors for most of the estimates are of relatively low magnitude. However, where an estimated number or the numerator or the denominator of a rate or percentage is small, the sampling error may be high.

Certain terms used in this report are defined in Appendix II. Some of the terms have specialized meanings for the purpose of the survey. For example, estimates of the incidence of acute conditions include, with certain exceptions, those conditions which had started within the 2 weeks prior to the week of interview and which had

<p>This survey covers all kinds of illnesses. These first questions refer to LAST WEEK AND THE WEEK BEFORE, that is, the 2-week period outlined in red on this calendar. (<i>Hand calendar</i>)</p> <p>6. a. Was -- sick at any time LAST WEEK OR THE WEEK BEFORE? (the 2 weeks shown on that calendar?) b. What was the matter? c. Did -- have anything else during that 2-week period?</p>	<input type="checkbox"/> Yes 01 <input type="checkbox"/> No
<p>7. a. LAST WEEK OR THE WEEK BEFORE, did -- take any medicine or treatment for any condition (besides . . . which you told me about)? b. For what condition? c. Did -- take any medicine for any other condition?</p>	<input type="checkbox"/> Yes <input type="checkbox"/> No
<p>8. a. LAST WEEK OR THE WEEK BEFORE, did -- have any accidents or injuries? b. What were they? c. Did -- have any other accidents or injuries during that 2-week period?</p>	<input type="checkbox"/> Yes <input type="checkbox"/> No
<p>9. a. Did -- EVER have an (any other) accident or injury that still bothers him or affects him in anyway? b. In what way does it bother him? (<i>Record present effects</i>)</p>	<input type="checkbox"/> Yes <input type="checkbox"/> No
<p>Open your Flashcard Booklet to Card A.</p> <p>10. Read both sides of Card A, condition by condition; record in his column any conditions mentioned for the person.</p>	<input type="checkbox"/> Yes 01 <input type="checkbox"/> No
<p>Turn to Card B.</p> <p>11. Read both sides of Card B, condition by condition; record in his column any conditions mentioned for the person.</p>	<input type="checkbox"/> Yes <input type="checkbox"/> No
<p>12. a. Does -- have any other ailments, conditions, or problems with his health? b. What is the condition? (<i>Record condition itself if still present; otherwise record present effects.</i>) c. Any other problems with his health?</p>	<input type="checkbox"/> Yes <input type="checkbox"/> No

Figure 1. Illness-recall questions, July 1965-June 1966.

involved either medical attention or restricted activity. The exceptions, listed in Appendix II, are certain conditions, such as heart trouble and diabetes, which are always considered chronic, regardless of duration or onset.

Annual estimates of the number of disability days associated with each acute condition are derived from the number of such days experienced during the 2-week period prior to the week of interview and include all those reported, even if the illness causing the disability had its onset prior to the 2-week period. Since any particular day of disability may be associated with more than one condition, the sum of days for all conditions adds to more than the total number of person-days of disability.

The questionnaire used by the Health Interview Survey during July 1965-June 1966 is illustrated in the publication, "Current Estimates From the Health Interview Survey, United States, July 1965-June 1966" (*Vital and Health Statistics*, Series 10, No. 37). Each condition reported in response to the "illness-recall" questions 6-12, shown in figure 1, was entered on a separate condition page (see Appendix III). Further questions were asked about the specific condition to obtain a better description and to obtain information about the actions taken by the person, such as seeking medical attention or restricting usual activities.

By comparing the questionnaires used for July 1964-June 1965 and July 1965-June 1966, as presented in the Current Estimates reports for these years (*Vital and Health Statistics*, Series 10, Nos. 25 and 37), it can be seen that the format of the questionnaires was markedly different. In the current year's questionnaire a new format was used to permit the use of an electronic device called FOSDIC (film optical sensing device for input to computers) in the processing of the data. On the condition pages of the questionnaire the interviewer coded all items except those shown in the locations marked "FOR WASHINGTON USE." These items were coded at the Bureau of the Census by trained medical coders. After coding was completed the pages of the questionnaire were microfilmed and processed for conversion to magnetic tape.

A series of tests have been made on the material collected in the current year to determine

its comparability with that collected for the same items in prior years. There is no tangible evidence that the change in the processing method has affected the estimates. However, changes in the levels of estimates from those of previous years may reflect, to some extent, response variation due to the revised format and administration of the questionnaire.

The incidence rates of acute conditions per quarter for the eight quarters from July 1959-June 1961, shown in figure 2, are presented in *Health Statistics*, Series B, No. 33. The rates for the period July 1961-June 1965 are presented in *Vital and Health Statistics*, Series 10, Nos. 1, 10, 15, and 26.

CONDITION GROUP, SEX, AND AGE

During the 12 months ending in June 1966 the civilian population, not residing in institutions, experienced an estimated 404.4 million acute illnesses or injuries requiring either medical attention or reduction of usual daily activities (table 1). The incidence rate per 100 persons was 212.0 during the current year compared with 212.7 conditions per 100 persons during July 1964-June 1965, as presented in the Acute Conditions report (*Vital and Health Statistics*, Series 10, No. 26).

Respiratory conditions comprised 59.4 percent of all acute conditions during July 1965-June 1966. There was an estimated incidence of 117.5 million common colds and 83.3 million conditions reported as influenza during the year. About four out of five of all respiratory conditions were classified in these categories.

The incidence rate per 100 persons per year for respiratory illnesses was 125.9 for the current year. This rate was about 8 percent higher than the rate of 116.4 for the previous year. The incidence rate for influenza-like illnesses was about 29 percent higher than that for the preceding year, 43.7 per 100 persons compared with an annual rate of 33.9 during the earlier period. The incidence rate of conditions reported as influenza during the current year was almost as high as that reported during the most recent influenza epidemic year, July 1962-June 1963 (shown in




Figure 2. Incidence of acute conditions per 100 persons per quarter, by condition group.

NOTE: Figures 2, 3, and 4 are plotted on a semilogarithmic scale so that visual comparisons of rates of change within and between curves may be made.

figures 2 and 3), during which the rate was 45.6 per 100 persons, about 4 percent higher than the current rate.

Figure 3 shows that the pattern of the influenza epidemic during the current year was different from that during the 1962-1963 season. The influenza incidence rate began rising in October-December 1965 and reached a peak in January-March 1966 somewhat below the level of January-March 1963, but the rate for the April-June quarter was higher than in the comparable quarter of 1963. The Communicable Disease Center's Influenza Surveillance report¹ indicates that the influenza season lasted from December 1965 through May 1966 and was caused by Type A and/or Type B influenza virus. Figure 3 also




Figure 3. Incidence of all acute conditions, upper respiratory conditions, and influenza per 100 persons per quarter.

¹Communicable Disease Center: *Influenza-Respiratory Disease Surveillance*. Report No. 82. Public Health Service. Atlanta, Ga., June 30, 1966.

shows a reversal in the usual pattern of the incidence of upper respiratory illnesses. During the current season the peak incidence was attained in the October-December interval rather than during the January-March interval, the period during which these conditions usually reach a maximum rate. This is the first time this early peak has occurred since the Health Interview Survey began in 1957. While it is possible that sampling variability may have contributed to this change of pattern, it is also possible that the severity of the respiratory illnesses caused increased reporting of influenza with a compensating decrease in such conditions as the common cold.

During July 1965-June 1966 the incidence rate of infective and parasitic diseases was lower than in the previous year, 25.1 per 100 persons compared with a rate of 27.5. A reduction in reporting of cases of measles was primarily responsible for the decline. The incidence of measles has fallen rapidly since the epidemic year of 1964 as shown below.

*Measles incidence
rate per 100 persons
per year*

July 1963-June 1964-----	5.4
July 1964-June 1965-----	3.6
July 1965-June 1966-----	1.5

It is probable that immunization campaigns have contributed substantially to the reduced incidence rate of measles.

The incidence of digestive conditions was somewhat lower during July 1965-June 1966 than in the preceding year. As shown in figure 2, much of the reduction in rate occurred during the last quarter of the year, April-June 1966.

The incidence of injuries during July 1965-June 1966 was 25.4 injuries per 100 persons per year, compared with a rate of 29.9 during July 1964-June 1965. Comparison of the rates by type of injury shown in table 1 with those for the previous year (*Vital and Health Statistics*, Series 10, No. 26) shows that much of the decline occurred in the categories "open wounds and lacerations," "contusions and superficial injuries," and "other

current injuries." In the last group the largest drop was noted for injuries of "other or unspecified nature." As mentioned earlier a new questionnaire form was employed, which obtained more information about the injury. It is quite possible that injuries which would have previously been classified to this category were in the current year sufficiently described to permit more accurate classification to some category other than injury.

Reduction in the incidence in some of the other classes of injury may have resulted from the application of severity criteria to the collected data. In the survey an injury (or other acute condition) which is not medically attended or activity restricting is excluded from the data. Perhaps, in some instances an injury sustained in the current year was not medically attended while formerly, because of medical practice, costs, or availability of services, the same type of injury would have been seen by a physician. In this instance the injury incidence figures might be reduced for the current year even though the actual reporting levels remained about the same.

The use of a 2-week reference period in estimating incidence of acute conditions makes it necessary to assign a substantial weight to each event in the derivation of annual incidence estimates. For example, a change in reporting level of 29 events means that the estimate based on data collected in a 2-week-reference period is lowered or increased by approximately 1 million. Similarly, a change from a medically attended or activity restricting event to one with neither criterion affects the estimate in the same degree.

Females experienced a higher rate of acute conditions than did males, 220.1 conditions per 100 females compared with 203.4 per 100 males. This sex difference was particularly noticeable for respiratory conditions and "all other acute conditions" in the latter group (primarily because of the inclusion of genitourinary disorders, and of deliveries and disorders of pregnancy and the puerperium). In only one condition group, namely, injuries, was the rate for males higher than that for females.

Acute conditions caused an estimated 8.2 days of restricted activity per person in the population, including 3.7 days spent in bed (tables 2 and 3). Respiratory conditions caused about half




Figure 4. Incidence of acute conditions per 100 persons per year, by sex and age.

of the days of restricted activity and bed-stay. Injury was the second leading cause of reduced activity, while the category "infective and parasitic diseases" was the second leading cause of bed disability.

Table 4 shows the distribution of acute conditions by measures of impact, that is, the action taken by the person sustaining the condition. An estimated 27.8 percent of the acute conditions were medically attended only, while 34.9 percent required activity restriction only. The remaining 37.3 percent required both medical attention and activity restriction.

As age increased, the incidence rates of acute conditions declined (table 5). Figure 4 shows that the sex difference in rates, noted above, began about age 15 and continued through the age span. Among the condition groups, the age-specific rates for males declined as age advanced for most of the groups, with the exception of injuries where the rate was highest in the age group 17-44 years. Among females the only exception to the pattern shown in figure 4 was "all other acute conditions" (including the sex-specific conditions of the childbearing years) in which the highest rate was noted in the 17-44-year age group.

Comparison between table 5 in this report and the same table for the previous year (*Vital and Health Statistics*, Series 10, No. 26) shows that the reduction in incidence of injuries was confined primarily to persons under 17 years of age, as shown below:

	Incidence of injuries per 100 persons per year	
	July 1964- June 1965	July 1965- June 1966
All ages-----	29.9	25.4
Under 6 years -----	37.5	26.2
6-16 years -----	36.0	27.0
17-44 years -----	29.5	29.4
45 years and over----	22.5	18.9

As previously indicated the change in questionnaire format, the effect of the severity criteria (medical attention or activity restriction), and the sampling variability may have contributed to this reduction in injuries among children.

For all acute conditions the rate of restricted-activity days and bed-days was highest in the youngest age group, under 6 years (tables 6 and 7). Infective and parasitic diseases and respiratory conditions were primarily responsible for the high rate of disability among children. The rate of disability was highest in the group 45 years and older for digestive conditions and injuries. Disability-day rates were highest for persons aged 17-44 years for "all other acute conditions," due primarily to disability days among women in the childbearing years.

Children aged 6-16 years are classified as the school-age population. Days lost from school are counted solely for this population. The incidence of acute conditions was 251.4 conditions per 100 persons among this population group. Acute illness or injury caused an average of 8.4 days of reduced daily activity, 4.0 days spent in bed, and 4.6 days lost from school (table 8). Respiratory disease was the leading cause of both illness and disability among children 6-16 years.

The incidence rate of acute illness or injury among the currently employed population aged 17 years and older was 165.0 per 100 workers (table

9). (A currently employed person is defined for the purposes of the Health Interview Survey as a person who had a job or business or worked at any time during the 2-week period prior to the week of interview.) Respiratory conditions accounted for about 3 of each 5 acute conditions in this population group. Female workers experienced a higher rate of acute conditions than did male workers. The incidence rate was higher in the age group 17-44 years than for persons aged 45 years and older.

Acute conditions caused an average of 3.7 days lost from work per currently employed person (table 10). Males had a higher rate of time lost from work than did females, 3.8 days per male compared with 3.6 days per female in the currently employed population. Respiratory conditions and injuries were the leading causes of absenteeism.

Contrary to the pattern noted for the past 4 years, estimates of time lost from work associated with acute conditions during the current year are higher for males than for females, as shown below:

	<i>Work-loss days per currently employed person</i>	
	<i>Male</i>	<i>Female</i>
July 1961-June 1962 -----	3.3	3.7
July 1962-June 1963 -----	3.6	4.4
July 1963-June 1964 -----	3.2	3.3
July 1964-June 1965 -----	3.2	3.8
July 1965-June 1966 -----	3.8	3.6

The increase in rate of time lost from work among male workers over that for the prior year occurred as a result of influenza (an increase of .2 of 1 day) and of injuries (an increase of .3 of 1 day).

GEOGRAPHIC DISTRIBUTION

During July 1965-June 1966 residents of the 212 standard metropolitan statistical areas established for the 1960 Decennial Census experienced an incidence of 216.0 acute illnesses or injuries per 100 persons per year (table 11). Persons living in nonfarm areas outside of the metropolitan areas had an incidence rate of 209.0.




Figure 5. Incidence of acute conditions per 100 persons per year, by residence and age.

Persons living on farms outside of SMSA's had a rate of 183.5 conditions per 100 persons. During July 1964-June 1965 the incidence rates for persons living in metropolitan and nonfarm areas were closely comparable to those shown above, 217.9 for SMSA's and 209.3 for nonfarm areas. The current year's rate for farm residents was about 4 percent higher than the rate of 175.6 reported for the period July 1964-June 1965.

Figure 5 shows that the excess in rate for metropolitan areas occurred among persons under 25 years of age. Among persons 45 years and older the incidence rate was lowest for this residence group.

The rates of restricted-activity and bed-disability days associated with acute conditions were lower among farm residents than in the other areas (tables 12 and 13). The low rate of restricted activity for farm residents occurred primarily among young persons under 25 years of age. Among persons aged 25-44 years the rate of restricted activity was markedly higher in farm areas than in other areas of residence. However, for those 45 years and older the rate of activity restriction among farm residents was similar to that for nonfarm residents living in nonmetropolitan areas. The rate of bed disability associated with acute conditions was lowest for farm residents in five of the six age groups shown in table 13.

Comparison of incidence rates for the current year and the preceding year by condition group, shown in table 14, indicates that the increase in rate of respiratory illnesses was inversely related to the density of the population. The percentage increase in rate for respiratory conditions was as follows: metropolitan areas, 6.4 percent; outside SMSA-nonfarm, 8.6 percent; and outside SMSA-farm, 28.0 percent. The increase in rate in the metropolitan and nonfarm areas was caused by the higher incidence of influenza-like illnesses, while the rise in rate for farm residents occurred in both the upper respiratory and the influenza groups. The reduction in rate of infective and parasitic diseases occurred in each residence group, but to a greater extent in the farm population than elsewhere. The reduction in incidence of injuries was constant among the places of residence, averaging about 15 percent in metropolitan and nonfarm areas and about 11 percent in farm areas.

The rate of activity restriction due to acute illness among residents of SMSA's was about the same during both years (table 15). Similarly, the rate for farm residents was relatively unchanged. Among nonfarm residents, however, the rate fell slightly (about 4 percent). Rates of bed-stay rose somewhat in the metropolitan areas, remained the same in nonfarm areas, and fell by 13 percent in farm areas (table 16). The decline in bed-day rate for farm residents occurred in most of the condition groups with the exception of influenza and the category "all other acute conditions."

The incidence of acute conditions in the four geographic regions followed the usual pattern of




Figure 6. Incidence of acute conditions per 100 persons per year, by geographic region and age.

distribution established during the preceding years of the survey, with the rate for the West Region markedly higher than that in each of the other regions (table 17). The rates of disability days were highest in the West and lowest in the North Central Region (tables 18 and 19). Figure 6 shows that the age distribution of the incidence rates was consistent for the regions. However, the rates in the West Region for persons aged 15-64 years were somewhat higher than comparable rates in the other regions. The rise in the incidence rate for persons aged 65 and over who live in the South Region may be due to sampling variability of the data, since this pattern was not exhibited in data for previous years.

The increase in incidence of respiratory illnesses during the current year occurred in each region (table 20). However, the rise in rate of conditions reported as influenza was not constant in each region, increasing by about 15 percent in the Northeast, 25 percent in the North Central Region, 30 percent in the South, and 44 percent in the West Region. The Communicable Disease Center bulletin referred to earlier shows that Type A influenza was not active in the Northeast during the influenza season. Type B influenza was widespread throughout New England, but spread to a lesser extent through the remainder of the Northeast Region.

The reduction in rate of injuries occurred in each region, but to a somewhat greater extent in the West Region where the rate fell by one-third from 33.8 per 100 persons to 23.8.

The pattern of disability-day rates by region was not consistent (tables 21 and 22). The rate of restricted activity fell in the Northeast and North Central Regions, remained constant in the South, and rose in the West Region. The number of bed-days per 100 persons rose in all regions.

SEASONAL VARIATION: CALENDAR QUARTER

Figures 2 and 3 and tables 23-30 show the seasonal variation in incidence and associated disability days for acute conditions. Variation in rates for respiratory conditions has been discussed earlier in this report. The quarterly incidence rates for injuries were somewhat below the rates shown for the other years in figure 2.

The rate for digestive system disorders for the April-June quarter of 1966 was lower than in any preceding quarter shown in the figure and may be due to sampling variability.

Rates of restricted activity per quarter during the current year were quite similar to those for the preceding year (tables 25 and 26). The rate for the October-December 1965 quarter was about 7 percent higher than that for the same period in 1964, while the rate for April-June 1966 was about 11 percent lower than the rate for April-June 1965.

The number of bed-days per 100 persons per quarter was higher during October 1965-March 1966 than during these 2 quarters in the prior year (tables 27 and 28). Increased bed-stay for respiratory conditions was primarily responsible for the rise in the rate.

Table 30 shows the quarterly incidence rate of injuries and indicates that the decline in rate occurred to a greater extent in the latter half of the year than in the earlier portion. The rates for each quarter for the 2 years with the corresponding percent of decrease during the current year are shown below:

	<i>Incidence of injuries per 100 persons per quarter</i>		
	<i>July 1964- June 1965</i>	<i>July 1965- June 1966</i>	<i>Percent decrease</i>
July-Sept ---	8.2	7.7	-6.1
Oct.-Dec ---	7.5	6.2	-17.3
Jan.-Mar ---	6.3	5.5	-12.7
Apr.-June---	7.9	6.0	-24.1



DETAILED TABLES

Page

EXTENDED LIST OF CONDITION GROUPS, BY SEX

Table	1.	Incidence of acute conditions, percent distribution, and number of acute conditions per 100 persons per year, by sex and condition group: United States, July 1965-June 1966-----	13
	2.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by sex and condition group: United States, July 1965-June 1966-----	14
	3.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by sex and condition group: United States, July 1965-June 1966-----	15

MEASURES OF THE IMPACT OF ILLNESS

	4.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by measures of the impact of illness, sex, and condition group: United States, July 1965-June 1966-----	16
--	----	--	----

SEX AND AGE

	5.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by age, sex, and condition group: United States, July 1965-June 1966-----	17
	6.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by age, sex, and condition group: United States, July 1965-June 1966-----	18
	7.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by age, sex, and condition group: United States, July 1965-June 1966-----	19

TIME LOST FROM SCHOOL AND WORK

	8.	Days lost from school associated with acute conditions and days lost from school per 100 children (6-16) per year, by sex and condition group: United States, July 1965-June 1966-----	20
	9.	Incidence of acute conditions among currently employed persons and number of acute conditions per 100 currently employed persons per year, by age, sex, and condition group: United States, July 1965-June 1966-----	21
	10.	Days lost from work associated with acute conditions and days lost from work per 100 currently employed persons per year, by age, sex, and condition group: United States, July 1965-June 1966-----	22

DETAILED TABLES--Continued

Page

RESIDENCE

Table 11.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by residence, sex, and age: United States, July 1965-June 1966-----	23
12.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by residence, sex, and age: United States, July 1965-June 1966-----	24
13.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by residence, sex, and age: United States, July 1965-June 1966-----	25
14.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by residence, sex, and condition group: United States, July 1965-June 1966-----	26
15.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by residence, sex, and condition group: United States, July 1965-June 1966-----	27
16.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by residence, sex, and condition group: United States, July 1965-June 1966-----	28

REGION

17.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by region, sex, and age: United States, July 1965-June 1966-----	29
18.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by region, sex, and age: United States, July 1965-June 1966-----	30
19.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by region, sex, and age: United States, July 1965-June 1966-----	31
20.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by region, sex, and condition group: United States, July 1965-June 1966---	32
21.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by region, sex, and condition group: United States, July 1965-June 1966-----	33
22.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by region, sex, and condition group: United States, July 1965-June 1966-----	34

CALENDAR QUARTER

23.	Incidence of acute conditions per quarter and number of acute conditions per 100 persons per quarter, by sex and age: United States, July 1965-June 1966-----	35
24.	Incidence of acute conditions per quarter and number of acute conditions per 100 persons per quarter, by sex and condition group: United States, July 1965-June 1966-----	36
25.	Days of restricted activity associated with acute conditions per quarter and days of restricted activity per 100 persons per quarter, by sex and age: United States, July 1965-June 1966-----	37
26.	Days of restricted activity associated with acute conditions per quarter and days of restricted activity per 100 persons per quarter, by sex and condition group: United States, July 1965-June 1966-----	38
27.	Days of bed disability associated with acute conditions per quarter and days of bed disability per 100 persons per quarter, by sex and age: United States, July 1965-June 1966-----	39

DETAILED TABLES--Continued

Page

CALENDAR QUARTER--Continued

Table 28.	Days of bed disability associated with acute conditions per quarter and days of bed disability per 100 persons per quarter, by sex and condition group: United States, July 1965-June 1966-----	40
29.	Incidence of acute respiratory conditions and associated disability days per quarter and number of acute respiratory conditions per 100 persons per quarter, by age: United States, July 1965-June 1966-----	41
30.	Incidence of injuries and associated disability days per quarter and number of injuries per 100 persons per quarter, by age: United States, July 1965-June 1966-----	42

POPULATION

31.	Population used in obtaining rates shown in this publication, by residence, region, sex, and age: United States, July 1965-June 1966-----	43
32.	Population of currently employed persons, by sex and age: United States, July 1965-June 1966-----	44
33.	Population used in obtaining rates shown in this publication, by quarter, sex, and age: United States, July 1965-June 1966-----	45

TABLE 1. INCIDENCE OF ACUTE CONDITIONS, PERCENT DISTRIBUTION, AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

CONDITION GROUP	BOTH SEXES			PERCENT DISTRIBUTION			NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR		
	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE
ALL ACUTE CONDITIONS-----	404,350	187,786	216,564	100.0	100.0	100.0	212.0	203.4	220.1
INFECTIVE AND PARASITIC DISEASES----	47,909	22,142	25,767	11.8	11.8	11.9	25.1	24.0	26.2
COMMON CHILDHOOD DISEASES-----	13,905	7,153	6,752	3.4	3.8	3.1	7.3	7.7	6.9
THE VIRUS, N.O.S.-----	25,629	11,515	14,314	6.4	6.1	6.6	13.5	12.5	14.5
OTHER INFECTIVE AND PARASITIC DISEASES-----	8,175	3,474	4,702	2.0	1.8	2.2	4.3	3.8	4.8
RESPIRATORY CONDITIONS-----	240,069	110,592	129,476	59.4	58.9	59.8	125.9	119.8	131.6
UPPER RESPIRATORY CONDITIONS-----	147,628	68,500	78,528	36.4	36.5	36.3	77.1	74.2	79.8
COMMON COLD-----	117,454	54,699	62,755	29.0	29.1	29.0	61.6	59.2	63.8
OTHER ACUTE UPPER RESPIRATORY CONDITIONS-----	29,574	13,801	15,773	7.3	7.3	7.3	15.5	14.9	16.0
INFLUENZA-----	83,343	37,601	45,742	20.6	20.0	21.1	43.7	40.7	46.5
INFLUENZA WITH DIGESTIVE MANIFESTATIONS-----	8,448	3,521	4,927	2.1	1.9	2.3	4.4	3.8	5.0
OTHER INFLUENZA-----	74,895	34,080	40,815	18.5	18.1	18.8	39.3	36.9	41.5
OTHER RESPIRATORY CONDITIONS-----	9,698	4,491	5,206	2.4	2.4	2.4	5.1	4.9	5.3
PNEUMONIA-----	2,448	1,133	1,315	0.6	0.6	0.6	1.3	1.2	1.3
BRONCHITIS-----	4,664	2,176	2,488	1.2	1.2	1.1	2.4	2.4	2.5
OTHER ACUTE RESPIRATORY CONDITIONS-----	2,586	1,182	1,404	0.6	0.6	0.6	1.4	1.3	1.4
DIGESTIVE SYSTEM CONDITIONS-----	19,912	8,948	10,964	4.9	4.8	5.1	10.4	9.7	11.1
DENTAL CONDITIONS-----	8,650	4,147	4,503	2.1	2.2	2.1	4.5	4.5	4.6
FUNCTIONAL AND SYMPTOMATIC UPPER GASTROINTESTINAL DISORDERS, N.E.C.-----	3,950	1,496	2,454	1.0	0.8	1.1	2.1	1.6	2.5
OTHER DIGESTIVE SYSTEM CONDITIONS-----	7,312	3,305	4,008	1.8	1.8	1.9	3.8	3.6	4.1
INJURIES-----	48,357	28,232	20,125	12.0	15.0	9.3	25.4	30.6	20.5
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	14,249	8,431	5,818	3.5	4.5	2.7	7.5	9.1	5.9
FRACTURES AND DISLOCATIONS-----	4,897	3,278	1,619	1.2	1.7	0.7	2.6	3.6	1.6
SPRAINS AND STRAINS-----	9,352	5,153	4,199	2.3	2.7	1.9	4.9	5.6	4.3
OPEN WOUNDS AND LACERATIONS-----	14,439	9,150	5,289	3.6	4.9	2.4	7.6	9.9	5.4
CONTUSIONS AND SUPERFICIAL INJURIES-----	10,025	5,609	4,416	2.5	3.0	2.0	5.3	6.1	4.5
OTHER CURRENT INJURIES-----	9,644	5,041	4,603	2.4	2.7	2.1	5.1	5.5	4.7
ALL OTHER ACUTE CONDITIONS-----	48,103	17,872	30,231	11.9	9.5	14.0	25.2	19.4	30.7
DISEASES OF THE EAR-----	11,376	5,744	5,631	2.8	3.1	2.6	6.0	6.2	5.7
HEADACHES-----	3,929	1,408	2,521	1.0	0.7	1.2	2.1	1.5	2.6
GENITOURINARY DISORDERS-----	6,034	824	5,210	1.5	0.4	2.4	3.2	0.9	5.3
DELIVERIES AND DISORDERS OF PREGNANCY AND THE PUERPERIUM-----	4,743	...	4,743	1.2	...	2.2	2.5	...	4.8
DISEASES OF THE SKIN-----	5,216	2,983	2,233	1.3	1.6	1.0	2.7	3.2	2.3
DISEASES OF THE MUSCULOSKELETAL SYSTEM-----	4,319	1,773	2,546	1.1	0.9	1.2	2.3	1.9	2.6
ALL OTHER ACUTE CONDITIONS-----	12,487	5,140	7,347	3.1	2.7	3.4	6.5	5.6	7.5

NOTES: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 2. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE
	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS			DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR		
ALL ACUTE CONDITIONS-----	1,562,839	720,335	842,504	819.5	780.2	856.3
INFECTIVE AND PARASITIC DISEASES---	197,462	93,535	103,926	103.5	101.3	105.6
COMMON CHILDHOOD DISEASES-----	76,138	39,863	36,275	39.9	43.2	36.9
THE VIRUS, N.O.S.-----	77,792	34,410	43,382	40.8	37.3	44.1
OTHER INFECTIVE AND PARASITIC DISEASES-----	43,532	19,262	24,270	22.8	20.9	24.7
RESPIRATORY CONDITIONS-----	750,360	339,757	410,603	393.5	368.0	417.3
UPPER RESPIRATORY CONDITIONS----	366,714	165,486	201,228	192.3	179.2	204.5
COMMON COLD-----	287,310	127,118	160,192	150.7	137.7	162.8
OTHER ACUTE UPPER RESPIRATORY CONDITIONS-----	79,404	38,368	41,036	41.6	41.6	41.7
INFLUENZA-----	312,770	134,892	177,878	164.0	146.1	180.8
INFLUENZA WITH DIGESTIVE MANIFESTATIONS-----	23,254	8,310	14,944	12.2	9.0	15.2
OTHER INFLUENZA-----	289,516	126,582	162,935	151.8	137.1	165.6
OTHER RESPIRATORY CONDITIONS----	70,876	39,379	31,496	37.2	42.7	32.0
PNEUMONIA-----	36,662	21,105	15,757	19.3	22.9	16.0
BRONCHITIS-----	23,237	11,641	11,596	12.2	12.6	11.6
OTHER ACUTE RESPIRATORY CONDITIONS-----	10,777	6,633	4,143	5.7	7.2	4.2
DIGESTIVE SYSTEM CONDITIONS-----	68,090	29,538	38,552	35.7	32.0	39.2
DENTAL CONDITIONS-----	18,636	9,260	9,376	9.8	10.0	9.5
FUNCTIONAL AND SYMPTOMATIC UPPER GASTROINTESTINAL DISORDERS, N.E.C.-----	8,172	3,313	4,859	4.3	3.6	4.9
OTHER DIGESTIVE SYSTEM CONDITIONS-----	41,282	16,966	24,317	21.6	18.4	24.7
INJURIES-----	322,944	193,431	129,513	169.3	209.5	131.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	176,182	110,928	65,254	92.4	120.2	66.3
FRACTURES AND DISLOCATIONS----	108,735	74,309	34,426	57.0	80.5	35.0
SPRAINS AND STRAINS-----	67,447	36,619	30,828	35.4	39.7	31.3
OPEN WOUNDS AND LACERATIONS-----	47,490	31,244	16,246	24.9	33.8	16.5
CONTUSIONS AND SUPERFICIAL INJURIES-----	54,476	28,484	25,992	28.6	30.9	26.4
OTHER CURRENT INJURIES-----	44,797	22,776	22,021	23.5	24.7	22.4
ALL OTHER ACUTE CONDITIONS-----	223,983	64,072	159,910	117.4	69.4	162.5
DISEASES OF THE EAR-----	32,588	14,764	17,824	17.1	16.0	18.1
HEADACHES-----	3,969	1,802	2,168	2.1	2.0	2.2
GENITOURINARY DISORDERS-----	35,003	3,859	31,144	18.4	4.2	31.7
DELIVERIES AND DISORDERS OF PREGNANCY AND THE PUERPERIUM----	41,885	...	41,885	22.0	...	42.6
DISEASES OF THE SKIN-----	16,386	9,122	7,264	8.6	9.9	7.4
DISEASES OF THE MUSCULOSKELETAL SYSTEM-----	27,124	12,341	14,783	14.2	13.4	15.0
ALL OTHER ACUTE CONDITIONS-----	67,027	22,185	44,842	35.1	24.0	45.6

NOTE: N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 3. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE
	DAYS OF BED DISABILITY IN THOUSANDS			DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR		
ALL ACUTE CONDITIONS-----	697,296	300,603	396,693	365.6	325.6	403.2
INFECTIVE AND PARASITIC DISEASES---	103,492	48,249	55,243	54.3	52.3	56.1
COMMON CHILDHOOD DISEASES-----	34,184	18,089	16,094	17.9	19.6	16.4
THE VIRUS, N.O.S.-----	47,621	21,169	26,452	25.1	22.9	27.1
OTHER INFECTIVE AND PARASITIC DISEASES-----	21,487	8,991	12,496	11.3	9.7	12.7
RESPIRATORY CONDITIONS-----	374,343	166,274	208,069	196.3	180.1	211.5
UPPER RESPIRATORY CONDITIONS-----	153,385	67,805	85,580	80.4	73.4	87.0
COMMON COLD-----	115,620	50,161	65,458	60.6	54.3	66.5
OTHER ACUTE UPPER RESPIRATORY CONDITIONS-----	37,766	17,644	20,122	19.8	19.1	20.5
INFLUENZA-----	178,731	75,710	103,022	93.7	82.0	104.7
INFLUENZA WITH DIGESTIVE MANIFESTATIONS-----	14,659	5,656	9,003	7.7	6.1	9.2
OTHER INFLUENZA-----	164,072	70,053	94,019	86.0	75.9	95.6
OTHER RESPIRATORY CONDITIONS-----	42,227	22,760	19,467	22.1	24.7	19.8
PNEUMONIA-----	23,681	12,832	10,850	12.4	13.9	11.0
BRONCHITIS-----	12,939	6,305	6,634	6.8	6.8	6.7
OTHER ACUTE RESPIRATORY CONDITIONS-----	5,607	3,624	1,983	2.9	3.9	2.0
DIGESTIVE SYSTEM CONDITIONS-----	37,374	15,767	21,606	19.6	17.1	22.0
DENTAL CONDITIONS-----	6,738	3,132	3,606	3.5	3.4	3.7
FUNCTIONAL AND SYMPTOMATIC UPPER GASTROINTESTINAL DISORDERS, N.E.C.-----	4,390	1,970	2,419	2.3	2.1	2.5
OTHER DIGESTIVE SYSTEM CONDITIONS-----	26,246	10,665	15,581	13.8	11.6	15.8
INJURIES-----	89,693	45,925	43,767	47.0	49.7	44.5
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	43,784	25,687	18,097	23.0	27.8	18.4
FRACTURES AND DISLOCATIONS-----	25,722	16,701	9,022	13.5	18.1	9.2
SPRAINS AND STRAINS-----	18,061	8,986	9,075	9.5	9.7	9.2
OPEN WOUNDS AND LACERATIONS-----	10,863	5,819	5,044	5.7	6.3	5.1
CONTUSIONS AND SUPERFICIAL INJURIES-----	17,938	6,769	11,169	9.4	7.3	11.4
OTHER CURRENT INJURIES-----	17,108	7,651	9,457	9.0	8.3	9.6
ALL OTHER ACUTE CONDITIONS-----	92,394	24,386	68,008	48.4	26.4	69.1
DISEASES OF THE EAR-----	12,292	5,013	7,279	6.4	5.4	7.4
HEADACHES-----	1,505	*	*	0.8	*	*
GENITOURINARY DISORDERS-----	19,323	2,499	16,824	10.1	2.7	17.1
DELIVERIES AND DISORDERS OF PREGNANCY AND THE PUERPERIUM-----	19,895	...	19,895	10.4	...	20.2
DISEASES OF THE SKIN-----	2,642	*	*	1.4	*	*
DISEASES OF THE MUSCULOSKELETAL SYSTEM-----	8,482	4,195	4,288	4.4	4.5	4.4
ALL OTHER ACUTE CONDITIONS-----	28,254	11,127	17,127	14.8	12.1	17.4

NOTE: N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 4. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY MEASURES OF THE IMPACT OF ILLNESS, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	MEASURES OF THE IMPACT OF ILLNESS							
	TOTAL	MEDICALLY ATTENDED ONLY	MEDICALLY ATTENDED AND ACTIVITY RESTRICTING	ACTIVITY RESTRICTING ONLY (NOT MEDICALLY ATTENDED)	TOTAL	MEDICALLY ATTENDED ONLY	MEDICALLY ATTENDED AND ACTIVITY RESTRICTING	ACTIVITY RESTRICTING ONLY (NOT MEDICALLY ATTENDED)
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS				NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR			
ALL ACUTE CONDITIONS----	404,350	112,464	150,866	141,020	212.0	59.0	79.1	73.9
INFECTIVE AND PARASITIC DISEASES-----	47,909	6,913	22,905	18,091	25.1	3.6	12.0	9.5
RESPIRATORY CONDITIONS----	240,069	54,769	80,195	105,104	125.9	28.7	42.1	55.1
UPPER RESPIRATORY CONDITIONS-----	147,028	44,841	43,786	58,401	77.1	23.5	23.0	30.6
INFLUENZA-----	83,343	7,299	30,181	45,863	43.7	3.8	15.8	24.0
OTHER RESPIRATORY CONDITIONS-----	9,698	2,629	6,228	841	5.1	1.4	3.3	0.4
DIGESTIVE SYSTEM CONDITIONS-----	19,912	7,713	7,244	4,955	10.4	4.0	3.8	2.6
INJURIES-----	48,357	21,706	20,123	6,528	25.4	11.4	10.6	3.4
ALL OTHER ACUTE CONDITIONS-----	48,103	21,363	20,398	6,342	25.2	11.2	10.7	3.3
<u>MALE</u>								
ALL ACUTE CONDITIONS----	187,786	52,293	70,411	65,081	203.4	56.6	76.3	70.5
INFECTIVE AND PARASITIC DISEASES-----	22,142	3,027	10,833	8,282	24.0	3.3	11.7	9.0
RESPIRATORY CONDITIONS----	110,592	24,696	37,276	48,620	119.8	26.7	40.4	52.7
UPPER RESPIRATORY CONDITIONS-----	68,500	20,826	20,583	27,089	74.2	22.6	22.3	29.3
INFLUENZA-----	37,601	2,964	13,626	21,011	40.7	3.2	14.8	22.8
OTHER RESPIRATORY CONDITIONS-----	4,491	905	3,067	*	4.9	1.0	3.3	*
DIGESTIVE SYSTEM CONDITIONS-----	8,948	3,447	3,412	2,089	9.7	3.7	3.7	2.3
INJURIES-----	28,232	13,511	11,684	3,037	30.6	14.6	12.7	3.3
ALL OTHER ACUTE CONDITIONS-----	17,872	7,613	7,206	3,053	19.4	8.2	7.8	3.3
<u>FEMALE</u>								
ALL ACUTE CONDITIONS----	216,564	60,171	80,455	75,939	220.1	61.2	81.8	77.2
INFECTIVE AND PARASITIC DISEASES-----	25,767	3,886	12,072	9,809	26.2	3.9	12.3	10.0
RESPIRATORY CONDITIONS----	129,476	30,073	42,919	56,484	131.6	30.6	43.6	57.4
UPPER RESPIRATORY CONDITIONS-----	78,528	24,013	23,203	31,311	79.8	24.4	23.6	31.8
INFLUENZA-----	45,742	4,335	16,555	24,852	46.5	4.4	16.8	25.3
OTHER RESPIRATORY CONDITIONS-----	5,206	1,724	3,161	*	5.3	1.8	3.2	*
DIGESTIVE SYSTEM CONDITIONS-----	10,964	4,267	3,832	2,865	11.1	4.3	3.9	2.9
INJURIES-----	20,125	8,195	8,439	3,491	20.5	8.3	8.6	3.5
ALL OTHER ACUTE CONDITIONS-----	30,231	13,750	13,192	3,289	30.7	14.0	13.4	3.3

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 5. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	45 & OVER
BOTH SEXES	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS--	404,350	88,067	106,760	133,653	75,869	212.0	361.4	251.4	197.8	134.8
INFECTIVE AND PARASITIC DISEASES-----	47,909	15,209	15,824	12,567	4,309	25.1	62.4	37.3	18.6	7.7
RESPIRATORY CONDITIONS--	240,069	54,327	65,499	73,941	46,302	125.9	222.9	154.2	109.4	82.3
UPPER RESPIRATORY CONDITIONS-----	147,028	39,718	40,228	40,745	26,337	77.1	163.0	94.7	60.3	46.8
INFLUENZA-----	83,343	11,501	23,185	30,583	18,074	43.7	47.2	54.6	45.3	32.1
OTHER RESPIRATORY CONDITIONS-----	9,698	3,108	2,087	2,613	1,891	5.1	12.8	4.9	3.9	3.4
DIGESTIVE SYSTEM CONDITIONS-----	19,912	3,845	4,051	7,700	4,317	10.4	15.8	9.5	11.4	7.7
INJURIES-----	48,357	6,379	11,485	19,870	10,623	25.4	26.2	27.0	29.4	18.9
ALL OTHER ACUTE CONDITIONS-----	48,103	6,308	9,901	19,576	10,318	25.2	34.1	23.3	29.0	18.3
MALE	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS--	187,786	45,232	54,107	55,982	32,464	203.4	365.0	250.5	174.7	123.5
INFECTIVE AND PARASITIC DISEASES-----	22,142	7,862	8,156	4,305	1,819	24.0	63.4	37.8	13.4	6.9
RESPIRATORY CONDITIONS--	110,592	27,497	31,417	31,197	20,481	119.8	221.9	145.4	97.4	77.9
UPPER RESPIRATORY CONDITIONS-----	68,500	20,330	18,770	17,950	11,451	74.2	164.1	86.9	56.0	43.6
INFLUENZA-----	37,601	5,703	11,368	12,247	8,283	40.7	46.0	52.6	38.2	31.5
OTHER RESPIRATORY CONDITIONS-----	4,491	1,465	1,280	1,000	747	4.9	11.8	5.9	3.1	2.8
DIGESTIVE SYSTEM CONDITIONS-----	8,948	1,915	1,881	3,493	1,658	9.7	15.5	8.7	10.9	6.3
INJURIES-----	28,232	3,405	7,719	12,187	4,921	30.6	27.5	35.7	38.0	18.7
ALL OTHER ACUTE CONDITIONS-----	17,872	4,553	4,934	4,801	3,585	19.4	36.7	22.8	15.0	13.6
FEMALE	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS--	216,564	42,835	52,653	77,671	43,405	220.1	357.6	252.3	218.6	144.7
INFECTIVE AND PARASITIC DISEASES-----	25,767	7,347	7,668	8,262	2,490	26.2	61.3	36.7	23.2	8.3
RESPIRATORY CONDITIONS--	129,476	26,829	34,082	42,744	25,821	131.6	224.0	163.3	120.3	86.1
UPPER RESPIRATORY CONDITIONS-----	78,528	19,389	21,458	22,794	14,887	79.8	161.9	102.8	64.1	49.6
INFLUENZA-----	45,742	5,797	11,817	18,336	9,791	45.5	48.4	56.6	51.6	32.6
OTHER RESPIRATORY CONDITIONS-----	5,206	1,643	807	1,613	1,143	5.3	13.7	3.9	4.5	3.8
DIGESTIVE SYSTEM CONDITIONS-----	10,964	1,930	2,170	4,207	2,658	11.1	16.1	10.4	11.8	8.9
INJURIES-----	20,125	2,974	3,766	7,683	5,702	20.5	24.8	18.0	21.6	19.0
ALL OTHER ACUTE CONDITIONS-----	30,231	3,755	4,967	14,776	6,733	30.7	31.4	23.8	41.6	22.4

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 6. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.]

SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	45 & OVER
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS					DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS--	1,562,839	235,225	357,804	494,511	475,299	819.5	965.2	842.5	731.8	844.3
INFECTIVE AND PARASITIC DISEASES-----	197,462	56,770	73,319	42,463	24,909	103.5	233.0	172.6	62.8	44.2
RESPIRATORY CONDITIONS--	750,360	137,740	192,764	199,653	220,203	393.5	565.2	453.9	295.4	391.2
UPPER RESPIRATORY CONDITIONS-----	366,714	88,498	105,140	85,623	84,453	192.3	363.1	247.6	131.1	150.0
INFLUENZA-----	312,770	32,378	75,316	95,801	109,275	164.0	132.9	177.3	141.8	194.1
OTHER RESPIRATORY CONDITIONS-----	70,876	16,863	12,309	15,229	26,475	37.2	69.2	29.0	22.5	47.0
DIGESTIVE SYSTEM CONDITIONS-----	68,090	8,137	10,258	24,093	25,603	35.7	33.4	24.2	35.7	45.5
INJURIES-----	322,944	9,155	48,941	130,740	134,108	169.3	37.6	115.2	193.5	238.2
ALL OTHER ACUTE CONDITIONS-----	223,983	23,422	32,523	97,561	70,476	117.4	96.1	76.6	144.4	125.2
<u>MALE</u>										
ALL ACUTE CONDITIONS--	720,335	121,295	186,041	205,421	207,578	780.2	978.8	861.3	641.1	789.6
INFECTIVE AND PARASITIC DISEASES-----	93,535	29,556	37,862	14,614	11,504	101.3	238.5	175.3	45.6	43.6
RESPIRATORY CONDITIONS--	339,757	69,585	94,088	81,615	94,469	368.0	561.5	435.6	254.7	359.4
UPPER RESPIRATORY CONDITIONS-----	165,486	44,988	50,037	37,460	33,002	179.2	363.0	231.6	116.9	125.5
INFLUENZA-----	134,892	15,297	35,356	37,336	46,903	146.1	123.4	163.7	116.5	176.4
OTHER RESPIRATORY CONDITIONS-----	39,379	9,300	8,695	5,820	14,564	42.7	75.0	40.3	21.3	55.4
DIGESTIVE SYSTEM CONDITIONS-----	29,538	3,885	5,905	8,211	11,538	32.0	31.4	27.3	25.6	43.9
INJURIES-----	193,431	5,099	32,474	85,238	70,620	209.5	41.1	150.3	266.0	268.6
ALL OTHER ACUTE CONDITIONS-----	64,072	13,170	15,713	15,743	19,447	69.4	106.3	72.7	49.1	74.0
<u>FEMALE</u>										
ALL ACUTE CONDITIONS--	842,504	113,930	171,763	289,090	267,721	856.3	951.2	823.1	813.5	892.3
INFECTIVE AND PARASITIC DISEASES-----	103,926	27,215	35,457	27,850	13,405	105.6	227.2	169.9	78.4	44.7
RESPIRATORY CONDITIONS--	410,603	66,155	98,676	118,038	125,734	417.3	569.0	472.8	332.2	419.1
UPPER RESPIRATORY CONDITIONS-----	201,228	43,510	55,103	51,163	51,451	204.5	363.3	264.0	144.0	171.5
INFLUENZA-----	177,878	17,081	39,960	58,466	62,372	180.8	142.6	191.5	164.5	207.9
OTHER RESPIRATORY CONDITIONS-----	31,496	7,563	3,613	8,409	11,911	32.0	63.1	17.3	23.7	39.7
DIGESTIVE SYSTEM CONDITIONS-----	38,552	4,252	4,353	15,882	14,065	39.2	35.5	20.9	44.7	46.9
INJURIES-----	129,513	4,056	10,467	45,502	53,468	131.6	33.9	78.9	128.0	211.6
ALL OTHER ACUTE CONDITIONS-----	159,910	10,252	16,810	61,818	51,030	162.5	85.6	80.6	236.2	170.1

TABLE 7. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	45 & OVER
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS					DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS-	697,296	108,544	171,501	230,455	196,796	365.6	445.4	403.8	341.0	331.8
INFECTIVE AND PARASITIC DISEASES-----	103,492	27,477	35,511	27,414	13,090	54.3	112.7	83.6	40.6	23.3
RESPIRATORY CONDITIONS----	374,343	61,173	107,495	109,068	96,608	196.3	251.0	253.1	161.4	171.6
UPPER RESPIRATORY CONDITIONS-----	153,385	31,446	49,352	42,385	30,203	80.4	129.0	116.2	62.7	53.7
INFLUENZA-----	178,731	17,867	51,491	57,193	52,179	93.7	73.3	121.2	84.6	92.7
OTHER RESPIRATORY CONDITIONS-----	42,227	11,860	6,651	9,490	14,226	22.1	48.7	15.7	14.0	25.3
DIGESTIVE SYSTEM CONDITIONS-----	37,374	2,827	5,426	13,246	15,872	19.6	11.6	12.8	19.6	28.2
INJURIES-----	89,693	3,922	12,040	40,066	33,666	47.0	16.1	28.3	59.3	59.8
ALL OTHER ACUTE CONDITIONS-----	92,394	13,145	11,029	40,659	27,561	48.4	53.9	26.0	60.2	49.0
<u>MALE</u>										
ALL ACUTE CONDITIONS-	300,603	53,780	82,463	87,936	76,424	325.6	434.0	381.8	274.4	290.7
INFECTIVE AND PARASITIC DISEASES-----	48,249	14,100	18,716	9,229	6,205	52.3	113.8	86.6	28.8	23.6
RESPIRATORY CONDITIONS----	166,274	30,159	49,800	46,839	39,477	180.1	243.4	230.5	146.2	150.2
UPPER RESPIRATORY CONDITIONS-----	67,805	15,547	21,886	19,588	10,783	73.4	125.5	101.3	61.1	41.0
INFLUENZA-----	75,710	8,068	23,523	23,340	20,778	82.0	65.1	108.9	72.8	79.0
OTHER RESPIRATORY CONDITIONS-----	22,760	6,544	4,391	3,910	7,915	24.7	52.8	20.3	12.2	30.1
DIGESTIVE SYSTEM CONDITIONS-----	15,767	*	3,031	4,366	7,186	17.1	*	14.0	13.6	27.3
INJURIES-----	45,925	1,716	6,802	22,713	14,695	49.7	13.8	31.5	70.9	55.9
ALL OTHER ACUTE CONDITIONS-----	24,386	6,622	4,114	4,789	8,862	26.4	53.4	19.0	14.9	33.7
<u>FEMALE</u>										
ALL ACUTE CONDITIONS-	396,693	54,764	89,038	142,519	110,372	403.2	457.2	426.7	401.0	367.9
INFECTIVE AND PARASITIC DISEASES-----	55,243	13,378	16,796	18,185	6,885	56.1	111.7	80.5	51.2	22.9
RESPIRATORY CONDITIONS----	208,069	31,014	57,694	62,230	57,131	211.5	258.9	276.5	175.1	190.4
UPPER RESPIRATORY CONDITIONS-----	85,580	15,898	27,466	22,796	19,419	87.0	132.7	131.6	64.1	64.7
INFLUENZA-----	103,022	9,800	27,968	33,853	31,401	104.7	81.8	134.0	95.3	104.7
OTHER RESPIRATORY CONDITIONS-----	19,467	5,316	2,260	5,580	6,311	19.8	44.4	10.8	15.7	21.0
DIGESTIVE SYSTEM CONDITIONS-----	21,606	1,643	2,395	8,882	8,686	22.0	13.7	11.5	25.0	29.0
INJURIES-----	43,767	2,206	5,238	17,353	18,971	44.5	18.4	25.1	48.8	63.2
ALL OTHER ACUTE CONDITIONS-----	68,008	6,523	6,916	35,870	18,699	69.1	54.5	33.1	100.9	62.3

TABLE 8. DAYS LOST FROM SCHOOL ASSOCIATED WITH ACUTE CONDITIONS AND DAYS LOST FROM SCHOOL PER 100 CHILDREN (6-16) PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE
	DAYS LOST FROM SCHOOL IN THOUSANDS			DAYS LOST FROM SCHOOL PER 100 CHILDREN PER YEAR		
ALL ACUTE CONDITIONS-----	196,525	97,050	99,475	462.7	449.3	476.7
INFECTIVE AND PARASITIC DISEASES-----	43,375	21,671	21,703	102.1	100.3	104.0
RESPIRATORY CONDITIONS-----	122,422	59,398	63,024	288.3	275.0	302.0
UPPER RESPIRATORY CONDITIONS-----	65,102	31,267	33,835	153.3	144.7	162.1
INFLUENZA-----	49,745	22,663	27,082	117.1	104.9	129.8
OTHER RESPIRATORY CONDITIONS-----	7,575	5,468	2,107	17.8	25.3	10.1
DIGESTIVE SYSTEM CONDITIONS-----	5,013	2,640	2,374	11.8	12.2	11.4
INJURIES-----	10,486	6,108	4,378	24.7	28.3	21.0
ALL OTHER ACUTE CONDITIONS-----	15,229	7,234	7,995	35.9	33.5	38.3

TABLE 9. INCIDENCE OF ACUTE CONDITIONS AMONG CURRENTLY EMPLOYED PERSONS AND NUMBER OF ACUTE CONDITIONS PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AGES- 17 & OVER	17-44	45 & OVER	ALL AGES 17 & OVER	17-44	45 & OVER
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CON- DITIONS IN THOUSANDS			NUMBER OF ACUTE CONDITIONS PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR		
ALL ACUTE CONDITIONS-----	120,630	81,872	38,758	165.0	187.3	131.8
INFECTIVE AND PARASITIC DISEASES-----	9,539	6,987	2,552	13.0	16.0	8.7
RESPIRATORY CONDITIONS-----	69,785	45,844	23,941	95.4	104.9	81.4
UPPER RESPIRATORY CONDITIONS-----	38,489	25,061	13,428	52.6	57.3	45.7
INFLUENZA-----	28,646	19,062	9,565	39.2	43.7	32.5
OTHER RESPIRATORY CONDITIONS-----	2,650	1,702	948	3.6	3.9	3.2
DIGESTIVE SYSTEM CONDITIONS-----	7,167	5,174	1,994	9.8	11.8	6.8
INJURIES-----	20,697	14,689	6,008	28.3	33.6	20.4
ALL OTHER ACUTE CONDITIONS-----	13,441	9,178	4,264	18.4	21.0	14.5
<u>MALE</u>						
ALL ACUTE CONDITIONS-----	72,330	49,308	23,022	153.1	174.1	121.7
INFECTIVE AND PARASITIC DISEASES-----	5,335	3,838	1,496	11.3	13.5	7.9
RESPIRATORY CONDITIONS-----	41,332	27,123	14,209	87.5	95.7	75.1
UPPER RESPIRATORY CONDITIONS-----	22,989	15,275	7,714	48.7	53.9	40.8
INFLUENZA-----	16,951	11,025	5,926	35.9	38.9	31.3
OTHER RESPIRATORY CONDITIONS-----	1,391	823	*	2.9	2.9	*
DIGESTIVE SYSTEM CONDITIONS-----	4,284	3,151	1,133	9.1	11.1	6.0
INJURIES-----	14,730	10,750	3,979	31.2	37.9	21.0
ALL OTHER ACUTE CONDITIONS-----	6,650	4,446	2,205	14.1	15.7	11.7
<u>FEMALE</u>						
ALL ACUTE CONDITIONS-----	48,300	32,564	15,736	186.7	211.6	150.1
INFECTIVE AND PARASITIC DISEASES-----	4,204	3,149	1,056	16.3	20.5	10.1
RESPIRATORY CONDITIONS-----	28,454	18,722	9,732	110.0	121.7	92.8
UPPER RESPIRATORY CONDITIONS-----	15,500	9,786	5,714	59.9	63.6	54.5
INFLUENZA-----	11,695	8,056	3,639	45.2	52.4	34.7
OTHER RESPIRATORY CONDITIONS-----	1,259	879	*	4.9	5.7	*
DIGESTIVE SYSTEM CONDITIONS-----	2,884	2,023	861	11.1	13.1	8.2
INJURIES-----	5,967	3,939	2,029	23.1	25.6	19.4
ALL OTHER ACUTE CONDITIONS-----	6,791	4,732	2,059	26.3	30.8	19.6

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 10. DAYS LOST FROM WORK ASSOCIATED WITH ACUTE CONDITIONS AND DAYS LOST FROM WORK PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AGES- 17 & OVER	17-44	45 & OVER	ALL AGES 17 & OVER	17-44	45 & OVER
<u>BOTH SEXES</u>	DAYS LOST FROM WORK IN THOUSANDS			DAYS LOST FROM WORK PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR		
ALL ACUTE CONDITIONS-----	270,784	157,354	113,430	370.3	360.0	385.7
INFECTIVE AND PARASITIC DISEASES-----	20,966	15,063	5,903	28.7	34.5	20.1
RESPIRATORY CONDITIONS-----	119,980	66,546	53,434	164.1	152.2	181.7
UPPER RESPIRATORY CONDITIONS-----	46,867	26,495	20,371	64.1	60.6	69.3
INFLUENZA-----	60,452	34,660	25,792	82.7	79.3	87.7
OTHER RESPIRATORY CONDITIONS-----	12,661	5,391	7,270	17.3	12.3	24.7
DIGESTIVE SYSTEM CONDITIONS-----	12,382	7,072	5,310	16.9	16.2	18.1
INJURIES-----	86,622	51,268	37,354	121.2	117.3	127.0
ALL OTHER ACUTE CONDITIONS-----	28,834	17,405	11,430	39.4	39.8	38.9
<u>MALF</u>						
ALL ACUTE CONDITIONS-----	178,117	100,673	77,445	377.0	355.4	409.3
INFECTIVE AND PARASITIC DISEASES-----	12,267	8,675	3,592	26.0	30.6	19.0
RESPIRATORY CONDITIONS-----	73,209	40,944	32,266	154.9	144.5	170.5
UPPER RESPIRATORY CONDITIONS-----	28,414	16,523	11,891	60.1	58.3	62.8
INFLUENZA-----	36,676	20,516	16,159	77.6	72.4	85.4
OTHER RESPIRATORY CONDITIONS-----	8,120	3,904	4,216	17.2	13.8	22.3
DIGESTIVE SYSTEM CONDITIONS-----	8,495	3,917	4,577	18.0	13.8	24.2
INJURIES-----	70,153	39,185	30,969	148.5	138.3	163.7
ALL OTHER ACUTE CONDITIONS-----	13,993	7,952	6,041	29.6	28.1	31.9
<u>FEMALE</u>						
ALL ACUTE CONDITIONS-----	92,667	56,681	35,986	358.2	368.4	343.2
INFECTIVE AND PARASITIC DISEASES-----	8,699	6,388	2,310	33.6	41.5	22.0
RESPIRATORY CONDITIONS-----	46,771	25,602	21,168	180.8	166.4	201.9
UPPER RESPIRATORY CONDITIONS-----	18,453	9,972	8,480	71.3	64.8	80.9
INFLUENZA-----	23,777	14,144	9,633	91.9	91.9	91.9
OTHER RESPIRATORY CONDITIONS-----	4,541	*	3,055	17.6	*	29.1
DIGESTIVE SYSTEM CONDITIONS-----	3,887	3,155	*	15.0	20.5	*
INJURIES-----	18,469	12,084	6,385	71.4	78.5	60.9
ALL OTHER ACUTE CONDITIONS-----	14,841	9,452	5,389	57.4	61.4	51.4

TABLE 11. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS				NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR			
ALL AGES-----	404,350	263,542	120,705	20,103	212.0	216.0	209.0	183.5
UNDER 5 YEARS-----	73,006	48,949	21,727	2,330	361.2	381.2	339.8	239.0
5-14 YEARS-----	107,840	72,424	29,756	5,660	271.9	290.2	243.3	229.1
15-24 YEARS-----	61,588	41,142	17,603	2,844	209.7	219.2	195.9	176.3
25-44 YEARS-----	86,046	56,731	25,697	3,617	190.4	188.4	197.7	174.2
45-64 YEARS-----	53,529	32,880	16,772	3,877	138.3	132.8	149.2	142.7
65 & OVER YEARS-----	22,340	11,416	9,150	1,775	127.1	108.0	155.0	160.9
<u>MALE</u>								
ALL AGES-----	187,786	121,936	55,212	10,638	203.4	207.5	197.8	188.7
UNDER 5 YEARS-----	37,300	25,086	10,914	1,301	361.5	384.9	331.7	254.1
5-14 YEARS-----	54,526	37,127	14,382	3,017	270.7	294.8	230.0	232.6
15-24 YEARS-----	26,452	17,054	7,824	1,573	189.0	192.0	183.0	188.2
25-44 YEARS-----	37,043	24,065	11,206	1,773	171.6	166.8	182.2	176.9
45-64 YEARS-----	23,023	13,707	7,172	2,144	123.8	115.6	134.7	151.3
65 & OVER YEARS-----	9,441	4,897	3,714	829	122.8	108.9	141.7	144.4
<u>FEMALE</u>								
ALL AGES-----	216,564	141,606	65,493	9,465	220.1	224.0	219.5	178.0
UNDER 5 YEARS-----	35,706	23,864	10,813	1,029	361.0	377.4	348.4	221.8
5-14 YEARS-----	53,314	35,298	15,374	2,642	273.2	285.4	257.3	225.2
15-24 YEARS-----	35,136	24,087	9,778	1,271	228.6	243.7	207.6	163.6
25-44 YEARS-----	49,003	32,667	14,492	1,844	207.6	208.3	211.6	171.7
45-64 YEARS-----	30,506	19,173	9,600	1,733	151.7	148.7	162.1	133.3
65 & OVER YEARS-----	12,900	6,518	5,436	946	130.5	107.3	165.6	178.8

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 12. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR			
ALL AGES-----	1,562,839	1,007,645	471,429	83,765	819.5	825.9	816.2	764.7
UNDER 5 YEARS-----	189,005	130,475	53,824	4,705	935.2	1,016.1	841.8	482.6
5-14 YEARS-----	356,807	241,896	98,806	16,105	899.7	969.2	808.0	652.0
15-24 YEARS-----	212,297	140,227	62,129	9,941	723.0	747.2	691.4	616.3
25-44 YEARS-----	329,432	217,968	93,831	17,633	729.1	723.9	721.7	849.4
45-64 YEARS-----	306,063	187,513	95,592	22,958	790.6	757.5	850.2	845.0
65 & OVER YEARS-----	169,236	89,565	67,247	12,424	962.8	847.3	1,139.0	1,126.4
<u>MALE</u>								
ALL AGES-----	720,335	458,050	220,912	41,373	780.2	779.4	791.4	734.0
UNDER 5 YEARS-----	97,571	66,867	27,765	2,939	945.6	1,026.0	843.9	574.0
5-14 YEARS-----	182,969	124,711	49,394	8,865	908.4	990.4	790.1	683.5
15-24 YEARS-----	91,693	56,811	29,167	5,915	656.7	639.6	682.1	707.5
25-44 YEARS-----	140,324	91,847	41,200	7,277	650.2	636.6	669.7	726.2
45-64 YEARS-----	133,974	79,149	43,032	11,793	720.4	667.5	808.4	832.3
65 & OVER YEARS-----	73,603	38,665	30,354	4,585	957.0	860.0	1,158.1	798.8
<u>FEMALE</u>								
ALL AGES-----	642,504	549,595	250,517	42,392	856.3	869.2	839.5	797.4
UNDER 5 YEARS-----	91,434	63,608	26,059	1,766	924.3	1,005.8	839.5	380.6
5-14 YEARS-----	173,837	117,186	49,412	7,240	890.7	947.6	826.8	617.2
15-24 YEARS-----	120,403	83,415	32,963	4,025	783.3	843.9	699.9	518.0
25-44 YEARS-----	189,108	126,121	52,631	10,356	801.1	804.2	768.6	964.2
45-64 YEARS-----	172,089	108,364	52,560	11,165	855.5	840.4	887.7	858.8
65 & OVER YEARS-----	95,633	50,901	36,893	7,839	967.3	837.9	1,123.8	1,481.9

TABLE 13. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR			
ALL AGES-----	697,296	465,295	203,014	28,988	365.6	381.4	351.5	264.6
UNDER 5 YEARS-----	88,157	59,656	26,868	1,632	436.2	464.6	420.2	167.4
5-14 YEARS-----	167,246	119,417	41,546	6,284	421.7	478.5	339.8	254.4
15-24 YEARS-----	107,204	74,965	25,909	6,330	365.1	399.5	288.3	392.4
25-44 YEARS-----	147,893	99,570	43,040	5,283	327.3	330.7	331.1	254.5
45-64 YEARS-----	122,404	75,640	40,934	5,830	316.2	305.6	364.1	214.6
65 & OVER YEARS-----	64,392	36,048	24,716	3,628	366.3	341.0	418.6	328.9
<u>MALE</u>								
ALL AGES-----	300,603	195,739	90,776	14,088	325.6	333.1	325.2	249.9
UNDER 5 YEARS-----	43,259	28,615	13,534	*	419.3	439.1	411.4	*
5-14 YEARS-----	80,603	57,445	19,779	3,378	400.2	456.2	316.4	260.4
15-24 YEARS-----	42,130	26,401	11,863	3,866	301.1	297.2	277.4	462.4
25-44 YEARS-----	58,187	38,882	17,548	1,757	269.6	269.5	285.2	175.3
45-64 YEARS-----	47,221	28,304	15,670	3,248	253.9	238.7	294.4	229.2
65 & OVER YEARS-----	29,203	16,092	12,382	*	379.7	357.9	472.4	*
<u>FEMALE</u>								
ALL AGES-----	396,693	269,556	112,237	14,900	403.2	426.3	376.1	280.3
UNDER 5 YEARS-----	44,898	31,041	13,334	*	453.9	490.8	429.6	*
5-14 YEARS-----	86,643	61,972	21,767	2,905	444.0	501.1	364.2	247.7
15-24 YEARS-----	65,074	48,563	14,046	2,465	423.4	491.3	298.2	317.2
25-44 YEARS-----	89,706	60,688	25,492	3,525	380.0	387.0	372.3	328.2
45-64 YEARS-----	75,182	47,336	25,264	2,582	373.7	367.1	426.7	198.6
65 & OVER YEARS-----	35,190	19,956	12,335	2,899	355.9	328.5	375.7	548.0

TABLE 14. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
BOTH SEXES	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS				NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR			
ALL ACUTE CONDITIONS-----	404,350	263,542	120,705	20,103	212.0	216.0	209.0	183.5
INFECTIVE AND PARASITIC DISEASES--	47,909	33,523	12,907	1,479	25.1	27.5	22.3	13.5
RESPIRATORY CONDITIONS-----	240,069	154,086	73,404	12,579	125.9	126.3	127.1	114.8
UPPER RESPIRATORY CONDITIONS----	147,028	96,665	42,687	7,677	77.1	79.2	73.9	70.1
INFLUENZA-----	83,343	51,263	27,409	4,672	43.7	42.0	47.5	42.7
OTHER RESPIRATORY CONDITIONS----	9,698	6,159	3,309	*	5.1	5.0	5.7	*
DIGESTIVE SYSTEM CONDITIONS----	19,912	12,955	5,942	1,016	10.4	10.6	10.3	9.3
INJURIES-----	46,357	31,678	13,813	2,866	25.4	26.0	23.9	26.2
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	14,249	9,483	3,799	968	7.5	7.8	6.6	8.8
OPEN WOUNDS AND LACERATIONS----	14,439	9,636	4,034	770	7.6	7.9	7.0	7.0
CONTUSIONS AND SUPERFICIAL INJURIES-----	10,025	6,590	2,780	*	5.3	5.4	4.8	*
OTHER CURRENT INJURIES-----	9,644	5,969	3,201	*	5.1	4.9	5.5	*
ALL OTHER ACUTE CONDITIONS-----	48,103	31,301	14,640	2,163	25.2	25.7	25.3	19.7
MALE								
ALL ACUTE CONDITIONS-----	187,786	121,936	55,212	10,638	203.4	207.5	197.8	188.7
INFECTIVE AND PARASITIC DISEASES--	22,142	15,632	5,702	808	24.0	26.6	20.4	14.3
RESPIRATORY CONDITIONS-----	110,592	70,870	33,146	6,576	119.8	120.6	118.7	116.7
UPPER RESPIRATORY CONDITIONS----	68,500	45,359	18,981	4,160	74.2	77.2	68.0	73.8
INFLUENZA-----	37,601	22,408	12,870	2,323	40.7	38.1	46.1	41.2
OTHER RESPIRATORY CONDITIONS----	4,491	3,103	1,295	*	4.9	5.3	4.6	*
DIGESTIVE SYSTEM CONDITIONS----	8,948	5,715	2,721	*	9.7	9.7	9.7	*
INJURIES-----	28,232	18,369	7,986	1,877	30.6	31.3	28.6	33.3
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	8,431	5,613	2,262	*	9.1	9.6	8.1	*
OPEN WOUNDS AND LACERATIONS----	9,150	5,868	2,734	*	9.9	10.0	9.8	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	5,609	3,664	1,414	*	6.1	6.2	5.1	*
OTHER CURRENT INJURIES-----	5,041	3,224	1,576	*	5.5	5.5	5.6	*
ALL OTHER ACUTE CONDITIONS-----	17,872	11,350	5,657	865	19.4	19.3	20.3	15.3
FEMALE								
ALL ACUTE CONDITIONS-----	216,564	141,606	65,493	9,465	220.1	224.0	219.5	176.0
INFECTIVE AND PARASITIC DISEASES--	25,767	17,891	7,205	*	26.2	28.3	24.1	*
RESPIRATORY CONDITIONS-----	129,476	83,216	40,258	6,003	131.6	131.6	134.9	112.9
UPPER RESPIRATORY CONDITIONS----	78,528	51,306	23,706	3,516	79.8	81.1	79.4	66.1
INFLUENZA-----	45,742	28,854	14,538	2,349	46.5	45.6	48.7	44.2
OTHER RESPIRATORY CONDITIONS----	5,206	3,055	2,014	*	5.3	4.8	6.7	*
DIGESTIVE SYSTEM CONDITIONS----	10,964	7,240	3,221	*	11.1	11.5	10.8	*
INJURIES-----	20,125	13,309	5,827	989	20.5	21.0	19.5	18.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	5,818	3,870	1,536	*	5.9	6.1	5.1	*
OPEN WOUNDS AND LACERATIONS----	5,289	3,768	1,299	*	5.4	6.0	4.4	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	4,416	2,927	1,366	*	4.5	4.6	4.6	*
OTHER CURRENT INJURIES-----	4,603	2,745	1,625	*	4.7	4.3	5.4	*
ALL OTHER ACUTE CONDITIONS-----	30,231	19,951	8,982	1,298	30.7	31.6	30.1	24.4

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 15. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definition of terms are given in Appendix II

SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR			
ALL ACUTE CONDITIONS-----	1,562,839	1,007,645	471,429	83,765	819.5	825.9	816.2	764.7
INFECTIVE AND PARASITIC DISEASES--	197,462	135,181	54,212	8,069	103.5	110.8	93.9	73.7
RESPIRATORY CONDITIONS-----	750,360	490,324	223,114	36,923	393.5	401.9	386.3	337.1
UPPER RESPIRATORY CONDITIONS----	366,714	246,367	99,466	20,881	192.3	201.9	172.2	190.6
INFLUENZA-----	312,770	195,385	102,906	14,480	164.0	160.2	178.2	132.2
OTHER RESPIRATORY CONDITIONS----	70,876	48,572	20,742	1,562	37.2	39.8	35.9	14.3
DIGESTIVE SYSTEM CONDITIONS----	68,090	45,132	20,520	2,439	35.7	37.0	35.5	22.3
INJURIES-----	322,944	202,085	98,660	22,200	169.3	165.6	170.8	202.7
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	176,182	111,659	53,315	11,208	92.4	91.5	92.3	102.3
OPEN WOUNDS AND LACERATIONS----	47,490	27,487	15,723	4,280	24.9	22.5	27.2	39.1
CONTUSIONS AND SUPERFICIAL INJURIES-----	54,476	36,377	13,686	4,413	28.6	29.8	23.7	40.3
OTHER CURRENT INJURIES-----	44,797	26,561	15,937	2,299	23.5	21.8	27.6	21.0
ALL OTHER ACUTE CONDITIONS-----	223,983	134,924	74,924	14,135	117.4	110.6	129.7	129.0
<u>MALE</u>								
ALL ACUTE CONDITIONS-----	720,335	458,050	220,912	41,373	780.2	779.4	791.4	734.0
INFECTIVE AND PARASITIC DISEASES--	93,535	65,227	24,232	4,077	101.3	111.0	86.8	72.3
RESPIRATORY CONDITIONS-----	339,757	216,151	105,252	18,355	368.0	367.8	377.1	325.6
UPPER RESPIRATORY CONDITIONS----	165,486	107,416	47,179	10,891	179.2	182.8	169.0	193.2
INFLUENZA-----	134,892	81,280	46,680	6,933	146.1	138.3	167.2	123.0
OTHER RESPIRATORY CONDITIONS----	39,379	27,455	11,393	*	42.7	46.7	40.8	*
DIGESTIVE SYSTEM CONDITIONS----	29,538	19,750	8,863	*	32.0	33.6	31.8	*
INJURIES-----	193,431	116,702	61,981	14,749	209.5	198.6	222.0	261.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	110,928	67,487	35,588	7,852	120.2	114.8	127.5	139.3
OPEN WOUNDS AND LACERATIONS----	31,244	16,773	11,130	3,340	33.8	28.5	39.9	59.3
CONTUSIONS AND SUPERFICIAL INJURIES-----	28,484	18,858	7,397	2,229	30.9	32.1	26.5	39.5
OTHER CURRENT INJURIES-----	22,776	13,584	7,865	*	24.7	23.1	28.2	*
ALL OTHER ACUTE CONDITIONS-----	64,072	40,221	20,584	3,267	69.4	68.4	73.7	58.0
<u>FEMALE</u>								
ALL ACUTE CONDITIONS-----	842,504	549,595	250,517	42,392	856.3	869.2	839.5	797.4
INFECTIVE AND PARASITIC DISEASES--	103,926	69,954	29,980	3,992	105.6	110.6	100.5	75.1
RESPIRATORY CONDITIONS-----	410,603	274,173	117,861	18,568	417.3	433.6	395.0	349.3
UPPER RESPIRATORY CONDITIONS----	201,228	138,951	52,286	9,990	204.5	219.8	175.2	187.9
INFLUENZA-----	177,878	114,105	56,226	7,547	180.8	180.5	188.4	142.0
OTHER RESPIRATORY CONDITIONS----	31,496	21,116	9,349	*	32.0	33.4	31.3	*
DIGESTIVE SYSTEM CONDITIONS----	38,552	25,382	11,656	1,514	39.2	40.1	39.1	28.5
INJURIES-----	129,513	85,383	36,679	7,451	131.6	135.0	122.9	140.2
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	65,254	44,172	17,726	3,356	66.3	69.9	59.4	63.1
OPEN WOUNDS AND LACERATIONS----	16,246	10,713	4,593	*	16.5	16.9	15.4	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	25,992	17,520	6,288	2,184	26.4	27.7	21.1	41.1
OTHER CURRENT INJURIES-----	22,021	12,978	8,072	*	22.4	20.5	27.0	*
ALL OTHER ACUTE CONDITIONS-----	159,910	94,703	54,340	10,868	162.5	149.8	182.1	204.4

TABLE 16. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
BOTH SEXES	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR			
ALL ACUTE CONDITIONS-----	697,296	465,295	203,014	28,988	365.6	381.4	351.5	264.6
INFECTIVE AND PARASITIC DISEASES--	103,492	74,462	25,972	3,057	54.3	61.0	45.0	27.9
RESPIRATORY CONDITIONS-----	374,343	251,375	109,035	13,933	196.3	206.0	188.8	127.2
UPPER RESPIRATORY CONDITIONS----	153,385	106,536	39,393	7,457	80.4	87.3	68.2	68.1
INFLUENZA-----	178,731	115,935	56,731	6,066	93.7	95.0	98.2	55.4
OTHER RESPIRATORY CONDITIONS----	42,227	28,904	12,911	*	22.1	23.7	22.4	*
DIGESTIVE SYSTEM CONDITIONS----	37,374	25,166	11,206	*	19.6	20.6	19.4	*
INJURIES-----	89,693	57,632	27,283	4,779	47.0	47.2	47.2	43.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	43,784	29,963	12,985	*	23.0	24.6	22.5	*
OPEN WOUNDS AND LACERATIONS----	10,863	6,031	3,904	*	5.7	4.9	6.8	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	17,938	12,685	4,183	*	9.4	10.4	7.2	*
OTHER CURRENT INJURIES-----	17,108	8,952	6,210	1,946	9.0	7.3	10.8	17.8
ALL OTHER ACUTE CONDITIONS----	92,394	56,661	29,517	6,216	48.4	46.4	51.1	56.7
MALL								
ALL ACUTE CONDITIONS-----	300,603	195,739	90,776	14,088	325.6	333.1	325.2	249.9
INFECTIVE AND PARASITIC DISEASES--	48,249	34,042	12,171	2,037	52.3	57.9	43.6	36.1
RESPIRATORY CONDITIONS-----	166,274	108,422	50,611	7,242	180.1	184.5	181.3	128.5
UPPER RESPIRATORY CONDITIONS----	67,805	45,885	17,823	4,097	73.4	78.1	63.8	72.7
INFLUENZA-----	75,710	47,180	25,653	2,877	82.0	80.3	91.9	51.0
OTHER RESPIRATORY CONDITIONS----	22,760	15,356	7,136	*	24.7	26.1	25.6	*
DIGESTIVE SYSTEM CONDITIONS----	15,767	10,421	4,974	*	17.1	17.7	17.8	*
INJURIES-----	45,925	27,904	14,662	3,359	49.7	47.5	52.5	59.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	25,687	16,779	8,105	*	27.8	28.5	29.0	*
OPEN WOUNDS AND LACERATIONS----	5,819	2,363	2,528	*	6.3	4.0	9.1	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	6,769	4,331	1,850	*	7.3	7.4	6.6	*
OTHER CURRENT INJURIES-----	7,651	4,431	2,179	*	8.3	7.5	7.8	*
ALL OTHER ACUTE CONDITIONS----	24,386	14,951	8,358	*	26.4	25.4	29.9	*
FEMALE								
ALL ACUTE CONDITIONS-----	396,693	269,556	112,237	14,900	403.2	426.3	376.1	280.3
INFECTIVE AND PARASITIC DISEASES--	55,243	40,420	13,801	*	56.1	63.9	46.2	*
RESPIRATORY CONDITIONS-----	208,069	142,953	58,424	6,691	211.5	226.1	195.8	125.9
UPPER RESPIRATORY CONDITIONS----	85,580	60,650	21,570	3,360	87.0	95.9	72.3	63.2
INFLUENZA-----	103,022	68,754	31,078	3,189	104.7	108.7	104.1	60.6
OTHER RESPIRATORY CONDITIONS----	19,467	13,548	5,776	*	19.8	21.4	19.4	*
DIGESTIVE SYSTEM CONDITIONS----	21,606	14,744	6,232	*	22.0	23.3	20.9	*
INJURIES-----	43,767	29,728	12,620	*	44.5	47.0	42.3	*
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	18,097	13,184	4,860	*	18.4	20.9	16.4	*
OPEN WOUNDS AND LACERATIONS----	5,044	3,668	*	*	5.1	5.8	*	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	11,169	8,354	2,333	*	11.4	13.2	7.8	*
OTHER CURRENT INJURIES-----	9,457	4,521	4,031	*	9.6	7.2	13.5	*
ALL OTHER ACUTE CONDITIONS----	68,008	41,710	21,160	5,139	69.1	66.0	70.9	96.7

TABLE 17. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY REGION, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR				
ALL AGES-----	404,350	96,300	110,925	123,565	73,560	212.0	202.7	208.8	209.8	235.9
UNDER 5 YEARS-----	73,006	17,705	20,354	22,292	12,655	361.2	383.9	370.4	334.2	368.5
5-14 YEARS-----	107,840	27,004	29,659	31,932	19,245	271.9	291.3	267.7	252.2	289.5
15-24 YEARS-----	61,588	13,804	17,400	19,304	11,080	209.7	202.2	211.0	201.5	235.2
25-44 YEARS-----	86,046	20,223	23,885	25,633	16,305	190.4	176.4	194.0	186.4	212.8
45-64 YEARS-----	53,529	12,756	13,669	16,225	10,878	138.3	122.0	125.0	144.7	177.9
65 & OVER YEARS-----	22,340	4,809	5,957	8,178	3,397	127.1	98.7	117.6	162.8	129.8
<u>MALE</u>										
ALL AGES-----	187,786	43,368	51,079	58,248	35,091	203.4	189.1	197.8	205.8	229.9
UNDER 5 YEARS-----	37,300	9,302	10,201	11,576	6,221	361.5	393.3	368.5	337.3	354.7
5-14 YEARS-----	54,526	13,152	14,532	16,079	10,763	270.7	276.1	261.1	253.5	310.1
15-24 YEARS-----	26,452	5,628	7,245	8,931	4,648	189.0	170.5	184.3	197.0	208.7
25-44 YEARS-----	37,043	8,082	10,910	10,618	7,233	171.6	148.3	183.5	165.9	197.3
45-64 YEARS-----	23,023	5,345	5,887	7,136	4,655	123.8	106.9	110.7	134.3	156.9
65 & OVER YEARS-----	9,441	1,860	2,304	3,707	1,569	122.8	90.3	100.4	171.6	133.2
<u>FEMALE</u>										
ALL AGES-----	216,564	52,932	59,846	65,318	38,469	220.1	215.5	219.1	213.5	241.6
UNDER 5 YEARS-----	35,706	8,403	10,153	10,716	6,434	361.0	374.0	372.5	330.9	382.7
5-14 YEARS-----	53,314	13,852	15,128	15,853	8,482	273.2	307.3	274.3	250.9	267.1
15-24 YEARS-----	35,136	8,176	10,156	10,374	6,431	228.6	231.9	235.3	205.6	259.0
25-44 YEARS-----	49,003	12,141	12,975	14,816	9,071	207.6	201.8	203.9	204.9	227.1
45-64 YEARS-----	30,506	7,411	7,782	9,089	6,223	151.7	135.9	138.5	154.1	197.7
65 & OVER YEARS-----	12,900	2,949	3,653	4,470	1,828	130.5	104.8	131.8	156.2	127.0

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 18. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS					DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR				
ALL AGES-----	1,562,839	375,491	400,141	494,278	292,929	819.5	790.5	753.1	839.3	939.4
UNDER 5 YEARS-----	189,065	52,430	44,942	56,033	33,600	935.2	1,136.8	817.9	870.1	978.5
5-14 YEARS-----	356,007	93,304	92,779	103,080	67,643	899.7	1,006.4	837.4	814.3	1,017.5
15-24 YEARS-----	212,297	46,036	55,074	71,663	39,525	723.0	674.3	667.7	748.0	839.2
25-44 YEARS-----	329,432	76,950	85,400	101,305	65,776	729.1	671.1	693.8	736.8	858.6
45-64 YEARS-----	306,063	68,820	78,089	98,125	61,029	790.6	658.4	714.1	875.3	998.3
65 & OVER YEARS-----	169,236	37,951	43,857	62,072	25,356	962.8	778.8	865.9	1,236.0	968.9
<u>MALE</u>										
ALL AGES-----	720,335	167,603	182,755	229,163	140,814	780.2	730.6	707.7	809.8	922.7
UNDER 5 YEARS-----	97,571	27,653	23,080	28,226	18,612	945.6	1,169.3	833.8	822.4	1,061.1
5-14 YEARS-----	182,969	46,888	48,716	51,618	35,747	908.4	984.2	875.4	813.9	1,029.9
15-24 YEARS-----	91,493	17,225	24,511	33,490	16,668	656.7	521.8	623.4	738.8	748.5
25-44 YEARS-----	140,324	31,930	34,844	45,638	27,912	650.2	586.0	585.9	700.1	761.4
45-64 YEARS-----	133,974	28,667	31,620	43,775	29,912	720.4	573.2	594.6	823.9	1,008.5
65 & OVER YEARS-----	73,603	15,240	19,984	26,417	11,963	957.0	739.8	871.1	1,223.0	1,015.5
<u>FEMALE</u>										
ALL AGES-----	842,504	207,888	217,386	265,115	152,114	856.3	846.3	796.0	866.6	955.4
UNDER 5 YEARS-----	91,434	24,777	21,862	29,808	14,988	924.3	1,102.7	802.0	920.6	891.6
5-14 YEARS-----	173,837	46,416	44,063	51,463	31,896	890.7	1,029.9	799.0	814.5	1,004.3
15-24 YEARS-----	120,403	28,811	30,563	38,173	22,857	783.3	817.1	708.1	756.5	920.5
25-44 YEARS-----	189,108	45,020	50,556	55,667	37,864	801.1	748.2	794.7	769.9	947.8
45-64 YEARS-----	172,089	40,153	46,469	54,350	31,117	855.5	736.3	827.1	921.5	988.5
65 & OVER YEARS-----	95,633	22,712	23,873	35,655	13,392	967.3	807.1	861.5	1,245.8	930.6

TABLE 19. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

SEX AND AGE	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS					DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR				
ALL AGES-----	697,296	167,209	178,191	217,068	134,828	365.6	352.0	335.4	368.6	432.4
UNDER 5 YEARS-----	88,157	20,696	24,433	26,923	16,105	436.2	448.7	444.6	403.6	469.0
5-14 YEARS-----	167,246	42,124	43,093	47,972	34,058	421.7	454.4	388.9	379.0	512.3
15-24 YEARS-----	107,204	23,110	27,313	37,118	19,663	365.1	338.5	331.1	387.5	417.5
25-44 YEARS-----	147,093	37,571	35,779	44,053	30,490	327.3	327.7	290.7	320.4	398.0
45-64 YEARS-----	122,404	27,665	30,247	39,455	25,037	316.2	264.7	276.6	352.0	409.6
65 & OVER YEARS-----	64,392	16,044	17,327	21,547	9,475	366.3	329.2	342.1	429.1	362.1
<u>MALE</u>										
ALL AGES-----	300,603	69,651	76,040	94,143	60,770	325.6	303.6	294.5	332.7	398.2
UNDER 5 YEARS-----	43,259	10,450	12,182	13,104	7,523	419.3	441.9	440.1	381.8	428.9
5-14 YEARS-----	80,603	20,702	20,245	22,294	17,361	400.2	434.6	363.8	351.5	500.2
15-24 YEARS-----	42,130	6,081	11,443	16,248	8,358	301.1	184.2	291.0	358.4	375.3
25-44 YEARS-----	58,187	14,869	13,355	16,635	13,328	269.0	272.9	224.6	255.2	363.6
45-64 YEARS-----	47,221	9,883	10,663	16,812	9,863	253.9	197.6	200.5	316.4	332.5
65 & OVER YEARS-----	29,203	7,665	8,152	9,049	4,336	379.7	372.1	355.4	418.9	368.1
<u>FEMALE</u>										
ALL AGES-----	396,693	97,558	102,152	122,925	74,059	403.2	397.2	374.1	401.8	465.1
UNDER 5 YEARS-----	44,898	10,246	12,251	13,819	8,582	453.9	456.0	449.4	426.8	510.5
5-14 YEARS-----	86,643	21,421	22,848	25,678	16,697	444.0	475.3	414.3	406.4	525.7
15-24 YEARS-----	65,074	17,029	15,870	20,870	11,305	423.4	483.0	367.7	413.6	455.3
25-44 YEARS-----	89,706	22,702	22,424	27,417	17,162	380.0	377.3	352.5	379.2	429.6
45-64 YEARS-----	75,182	17,782	19,584	22,643	15,174	373.7	326.1	348.6	383.9	482.0
65 & OVER YEARS-----	35,190	8,378	9,175	12,498	5,139	355.9	297.7	331.1	436.7	357.1

TABLE 20. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

SEX AND CONDITION GROUP	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
BOTH SEXES	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS-----	404,350	96,300	110,925	123,565	73,560	212.0	202.7	208.0	209.8	235.9
INFECTIVE AND PARASITIC DISEASES-----	47,909	15,625	8,317	17,786	6,181	25.1	32.9	15.7	30.2	19.8
RESPIRATORY CONDITIONS-----	240,069	54,207	70,421	67,523	47,918	125.9	114.1	132.5	114.7	153.7
UPPER RESPIRATORY CONDITIONS----	147,026	41,185	39,128	43,500	23,215	77.1	86.7	73.6	73.9	74.4
INFLUENZA-----	83,343	10,833	28,625	20,361	23,325	43.7	22.8	54.3	34.6	74.8
OTHER RESPIRATORY CONDITIONS----	9,698	2,190	2,467	3,662	1,378	5.1	4.6	4.6	6.2	4.4
DIGESTIVE SYSTEM CONDITIONS-----	19,912	4,909	4,847	7,420	2,736	10.4	10.3	9.1	12.6	8.8
INJURIES-----	48,357	11,966	13,969	14,995	7,427	25.4	25.2	26.3	25.5	23.8
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	14,249	3,484	4,428	4,338	1,998	7.5	7.3	8.3	7.4	6.4
OPEN WOUNDS AND LACERATIONS-----	14,439	3,381	4,050	4,711	2,297	7.6	7.1	7.6	8.0	7.4
CONTUSIONS AND SUPERFICIAL INJURIES-----	10,025	2,474	2,501	3,488	1,562	5.3	5.2	4.7	5.9	5.0
OTHER CURRENT INJURIES-----	9,644	2,627	2,989	2,457	1,570	5.1	5.5	5.6	4.2	5.0
ALL OTHER ACUTE CONDITIONS-----	46,103	9,593	13,372	15,841	9,298	25.2	20.2	25.2	26.9	29.8
MALE										
ALL ACUTE CONDITIONS-----	187,786	43,366	51,079	58,248	35,091	203.4	189.1	197.8	205.8	229.9
INFECTIVE AND PARASITIC DISEASES-----	22,142	7,204	3,761	8,218	2,959	24.0	31.4	14.6	29.0	19.4
RESPIRATORY CONDITIONS-----	110,592	24,451	32,221	31,385	22,535	119.8	106.6	124.8	110.9	147.7
UPPER RESPIRATORY CONDITIONS----	68,500	19,054	18,752	20,348	10,347	74.2	83.1	72.6	71.9	67.8
INFLUENZA-----	37,601	4,491	12,394	9,145	11,571	40.7	19.6	48.0	32.3	75.8
OTHER RESPIRATORY CONDITIONS----	4,491	906	1,076	1,892	*	4.9	3.9	4.2	6.7	*
DIGESTIVE SYSTEM CONDITIONS-----	8,946	1,995	1,953	3,574	1,426	9.7	8.7	7.6	12.6	9.3
INJURIES-----	26,232	6,326	8,425	9,133	4,347	30.6	27.6	32.6	32.3	28.5
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	8,431	1,898	2,773	2,588	1,172	9.1	8.3	10.7	9.1	7.7
OPEN WOUNDS AND LACERATIONS-----	9,150	2,041	2,411	3,273	1,425	9.9	8.9	9.3	11.6	9.3
CONTUSIONS AND SUPERFICIAL INJURIES-----	5,609	1,194	1,576	1,904	935	6.1	5.2	6.1	6.7	6.1
OTHER CURRENT INJURIES-----	5,041	1,193	1,664	1,369	815	5.5	5.2	6.4	4.8	5.3
ALL OTHER ACUTE CONDITIONS-----	17,872	3,393	4,719	5,937	3,823	19.4	14.8	18.3	21.0	25.1
FEMALE										
ALL ACUTE CONDITIONS-----	216,564	52,932	59,846	65,318	38,469	220.1	215.5	219.1	213.5	241.6
INFECTIVE AND PARASITIC DISEASES-----	25,767	8,421	4,556	9,568	3,222	26.2	34.3	16.7	31.3	20.2
RESPIRATORY CONDITIONS-----	129,476	29,756	38,199	36,138	25,383	131.6	121.1	139.9	118.1	159.4
UPPER RESPIRATORY CONDITIONS----	78,526	22,131	20,377	23,152	12,868	79.8	90.1	74.6	75.7	80.8
INFLUENZA-----	45,742	6,341	16,431	11,216	11,754	46.5	25.8	60.2	36.7	73.0
OTHER RESPIRATORY CONDITIONS----	5,206	1,203	1,391	1,770	761	5.3	5.2	5.1	5.8	4.8
DIGESTIVE SYSTEM CONDITIONS-----	10,964	2,915	2,894	3,846	1,310	11.1	11.9	10.6	12.6	8.2
INJURIES-----	20,125	5,640	5,544	5,861	3,080	20.5	23.0	20.3	19.2	19.3
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	5,816	1,507	1,655	1,750	826	5.9	6.5	6.1	5.7	5.2
OPEN WOUNDS AND LACERATIONS-----	5,289	1,340	1,639	1,438	872	5.4	5.5	6.0	4.7	5.5
CONTUSIONS AND SUPERFICIAL INJURIES-----	4,416	1,279	925	1,564	*	4.5	5.2	3.4	5.2	*
OTHER CURRENT INJURIES-----	4,603	1,435	1,325	1,088	755	4.7	5.8	4.9	3.6	4.7
ALL OTHER ACUTE CONDITIONS-----	30,231	6,199	8,653	9,904	5,475	30.7	25.2	31.7	32.4	34.4

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 21. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

SEX AND CONDITION GROUP	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
BOTH SEXES	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS					DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS-----	1,562,839	375,491	400,141	494,278	292,929	819.5	790.5	753.1	839.3	939.4
INFECTIVE AND PARASITIC DISEASES-----	197,462	63,863	37,486	66,553	29,560	103.5	134.4	70.6	113.0	94.8
RESPIRATORY CONDITIONS-----	750,360	177,243	202,480	212,770	157,867	393.5	373.1	381.1	361.3	506.2
UPPER RESPIRATORY CONDITIONS-----	366,714	111,653	96,561	103,418	55,082	192.3	235.0	181.7	175.6	175.5
INFLUENZA-----	312,770	46,419	90,089	84,263	92,000	164.0	97.7	169.6	143.1	295.0
OTHER RESPIRATORY CONDITIONS-----	70,876	19,171	15,830	25,089	10,786	37.2	40.4	29.8	42.6	34.6
DIGESTIVE SYSTEM CONDITIONS-----	68,090	16,639	18,675	23,051	9,725	35.7	35.0	35.1	39.1	31.2
INJURIES-----	322,944	74,192	79,159	115,063	54,530	169.3	156.2	149.0	195.4	174.9
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	176,182	40,261	47,845	58,849	29,207	92.4	84.8	90.0	99.9	93.7
OPEN WOUNDS AND LACERATIONS-----	47,490	9,430	10,128	20,691	7,240	24.9	19.9	19.1	35.1	23.2
CONTUSIONS AND SUPERFICIAL INJURIES-----	54,476	14,266	11,348	18,453	10,409	28.6	30.0	21.4	31.3	33.4
OTHER CURRENT INJURIES-----	44,797	10,215	9,837	17,070	7,675	23.5	21.5	18.5	29.0	24.6
ALL OTHER ACUTE CONDITIONS-----	223,983	43,553	62,341	76,842	41,247	117.4	91.7	117.3	130.5	132.3
MALE										
ALL ACUTE CONDITIONS-----	720,335	167,603	182,755	229,163	140,814	780.2	730.6	707.7	809.8	922.7
INFECTIVE AND PARASITIC DISEASES-----	93,535	27,553	18,577	33,250	14,155	101.3	120.1	71.9	117.5	92.8
RESPIRATORY CONDITIONS-----	339,757	76,808	94,240	97,847	70,863	368.0	334.8	364.9	345.8	464.3
UPPER RESPIRATORY CONDITIONS-----	165,486	47,856	48,463	47,030	22,137	179.2	208.6	187.7	166.2	145.1
INFLUENZA-----	134,892	18,979	38,624	35,418	41,870	146.1	82.7	149.6	125.2	274.4
OTHER RESPIRATORY CONDITIONS-----	39,379	9,972	7,153	15,398	6,857	42.7	43.5	27.7	54.4	44.9
DIGESTIVE SYSTEM CONDITIONS-----	29,538	6,529	7,271	10,459	5,279	32.0	28.5	28.2	37.0	34.6
INJURIES-----	193,431	44,306	46,185	65,855	37,085	209.5	193.1	178.8	232.7	243.0
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	110,928	24,177	30,313	36,296	20,142	120.2	105.4	117.4	128.3	132.0
OPEN WOUNDS AND LACERATIONS-----	31,244	6,842	5,409	14,109	4,884	33.8	29.8	20.9	49.9	32.0
CONTUSIONS AND SUPERFICIAL INJURIES-----	28,484	6,586	6,322	8,533	7,043	30.9	28.7	24.5	30.2	46.2
OTHER CURRENT INJURIES-----	22,776	6,701	4,141	6,918	5,016	24.7	29.2	16.0	24.4	32.9
ALL OTHER ACUTE CONDITIONS-----	64,072	12,407	16,482	21,752	13,432	69.4	54.1	63.8	76.9	86.0
FEMALE										
ALL ACUTE CONDITIONS-----	842,504	207,868	217,386	265,115	152,114	856.3	846.3	796.0	866.6	955.4
INFECTIVE AND PARASITIC DISEASES-----	103,926	36,311	18,909	33,302	15,405	105.6	147.8	69.2	108.9	96.8
RESPIRATORY CONDITIONS-----	410,603	100,436	108,240	114,923	87,003	417.3	408.9	396.4	375.7	546.4
UPPER RESPIRATORY CONDITIONS-----	201,228	63,797	48,098	56,387	32,945	204.5	259.7	176.1	184.3	206.9
INFLUENZA-----	177,878	27,439	51,465	48,845	50,129	180.8	111.7	188.5	159.7	314.8
OTHER RESPIRATORY CONDITIONS-----	31,496	9,199	8,678	9,691	3,929	32.0	37.4	31.8	31.7	24.7
DIGESTIVE SYSTEM CONDITIONS-----	38,552	10,110	11,405	12,591	4,446	39.2	41.2	41.8	41.2	27.9
INJURIES-----	129,513	29,866	32,974	49,208	17,445	131.6	121.7	120.7	160.9	109.6
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	65,254	16,104	17,532	22,553	9,065	66.3	65.6	64.2	73.7	56.9
OPEN WOUNDS AND LACERATIONS-----	16,246	2,588	4,720	6,583	2,355	16.5	10.5	17.3	21.5	14.6
CONTUSIONS AND SUPERFICIAL INJURIES-----	25,992	7,680	5,026	9,921	3,366	26.4	31.3	18.4	32.4	21.1
OTHER CURRENT INJURIES-----	22,021	3,514	5,696	10,152	2,659	22.4	14.3	20.9	33.2	16.7
ALL OTHER ACUTE CONDITIONS-----	159,910	31,146	45,859	55,090	27,816	162.5	126.8	167.9	180.1	174.7

TABLE 22. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

SEX AND CONDITION GROUP	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS					DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR				
ALL ACUTE CONDITIONS-----	697,296	172,209	178,191	217,068	134,828	365.6	352.0	335.4	368.6	432.4
INFECTIVE AND PARASITIC DISEASES-----	103,492	33,906	20,513	35,180	13,893	54.3	71.4	38.6	59.7	44.6
RESPIRATORY CONDITIONS-----	374,343	86,357	100,931	103,928	83,127	196.3	181.8	190.0	176.5	266.6
UPPER RESPIRATORY CONDITIONS----	153,385	47,182	39,702	43,509	22,992	80.4	99.3	74.7	73.9	73.7
INFLUENZA-----	178,731	28,269	50,900	46,112	53,450	93.7	59.5	95.8	78.3	171.4
OTHER RESPIRATORY CONDITIONS----	42,227	10,906	10,329	14,307	6,685	22.1	23.0	19.4	24.3	21.4
DIGESTIVE SYSTEM CONDITIONS-----	37,374	9,677	9,499	12,786	5,411	19.6	20.4	17.9	21.7	17.4
INJURIES-----	89,693	20,205	20,308	32,651	16,529	47.0	42.5	36.2	55.4	53.0
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	43,784	10,338	10,100	13,869	9,477	23.0	21.8	19.0	23.6	30.4
OPEN WOUNDS AND LACERATIONS-----	10,863	2,140	1,816	5,324	1,582	5.7	4.5	3.4	9.0	5.1
CONTUSIONS AND SUPERFICIAL INJURIES-----	17,936	3,881	3,921	7,027	3,110	9.4	8.2	7.4	11.9	10.0
OTHER CURRENT INJURIES-----	17,108	3,846	4,471	6,430	2,360	9.0	8.1	8.4	10.9	7.6
ALL OTHER ACUTE CONDITIONS-----	92,394	17,064	26,940	32,522	15,868	48.4	35.9	50.7	55.2	50.9
<u>MALE</u>										
ALL ACUTE CONDITIONS-----	300,603	69,651	76,040	94,143	60,770	325.6	303.6	294.5	332.7	398.2
INFECTIVE AND PARASITIC DISEASES-----	48,249	14,651	9,566	17,108	6,925	52.3	63.9	37.0	60.5	45.4
RESPIRATORY CONDITIONS-----	166,274	37,020	46,016	46,499	36,739	180.1	161.4	178.2	164.3	240.7
UPPER RESPIRATORY CONDITIONS----	67,805	19,799	20,188	18,892	8,927	73.4	86.3	78.2	66.8	58.5
INFLUENZA-----	75,710	11,796	21,030	18,427	24,457	82.0	51.4	81.4	65.1	160.3
OTHER RESPIRATORY CONDITIONS----	22,760	5,425	4,799	9,181	3,355	24.7	23.6	18.6	32.4	22.0
DIGESTIVE SYSTEM CONDITIONS-----	15,767	3,732	3,613	5,970	2,452	17.1	16.3	14.0	21.1	16.1
INJURIES-----	45,925	10,519	9,819	15,151	10,436	49.7	45.9	38.0	53.5	68.4
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	25,687	6,096	5,394	7,697	6,500	27.8	26.6	20.9	27.2	42.6
OPEN WOUNDS AND LACERATIONS-----	5,819	*	*	3,498	*	6.3	*	*	12.4	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	6,769	*	1,969	2,281	1,527	7.3	*	7.6	8.1	10.0
OTHER CURRENT INJURIES-----	7,651	2,539	1,869	1,575	1,568	8.3	11.1	7.2	5.9	10.3
ALL OTHER ACUTE CONDITIONS-----	24,386	3,728	7,026	9,415	4,217	26.4	16.3	27.2	33.3	27.6
<u>FEMALE</u>										
ALL ACUTE CONDITIONS-----	396,693	97,558	102,152	122,925	74,059	403.2	397.2	374.1	401.8	465.1
INFECTIVE AND PARASITIC DISEASES-----	55,243	19,254	10,948	18,073	6,968	56.1	78.4	40.1	59.1	43.8
RESPIRATORY CONDITIONS-----	206,069	49,337	54,915	57,429	46,387	211.5	200.9	201.1	187.7	291.3
UPPER RESPIRATORY CONDITIONS----	85,580	27,384	19,514	24,617	14,066	87.0	111.5	71.5	80.5	88.3
INFLUENZA-----	103,022	16,473	29,871	27,685	28,992	104.7	67.1	109.4	90.5	182.1
OTHER RESPIRATORY CONDITIONS----	19,467	5,480	5,530	5,127	3,329	19.8	22.3	20.2	16.8	20.9
DIGESTIVE SYSTEM CONDITIONS-----	21,606	5,945	5,886	6,816	2,959	22.0	24.2	21.6	22.3	18.6
INJURIES-----	43,767	9,686	10,489	17,500	6,093	44.5	39.4	38.4	57.2	38.3
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS-----	18,097	4,242	4,706	6,172	2,977	18.4	17.3	17.2	20.2	18.7
OPEN WOUNDS AND LACERATIONS-----	5,044	*	*	1,826	*	5.1	*	*	6.0	*
CONTUSIONS AND SUPERFICIAL INJURIES-----	11,169	2,888	1,952	4,746	1,583	11.4	11.8	7.1	15.5	9.9
OTHER CURRENT INJURIES-----	9,457	*	2,602	4,755	*	9.6	*	9.5	15.5	*
ALL OTHER ACUTE CONDITIONS-----	68,008	13,336	19,914	23,107	11,651	69.1	54.3	72.9	75.5	73.2

TABLE 23. INCIDENCE OF ACUTE CONDITIONS PER QUARTER AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS				NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER QUARTER			
ALL AGES----	68,090	118,759	133,087	84,414	35.8	62.3	69.7	44.1
UNDER 5 YEARS-----	13,343	22,071	22,230	15,363	65.3	108.7	110.3	77.0
5-14 YEARS-----	16,113	30,387	38,724	22,616	40.9	76.8	97.4	56.6
15-24 YEARS-----	11,446	18,728	19,131	12,282	39.4	64.0	64.9	41.4
25-44 YEARS-----	15,187	24,843	27,681	18,334	33.6	55.0	61.3	40.6
45-64 YEARS-----	8,219	15,465	18,292	11,553	21.3	40.0	47.2	29.7
65 & OVER YEARS---	3,781	7,265	7,028	4,266	21.7	41.4	39.9	24.1
<u>MALE</u>								
ALL AGES----	31,375	55,531	61,639	39,240	34.1	60.2	66.7	42.4
UNDER 5 YEARS-----	6,561	11,207	11,297	8,235	62.9	108.1	109.8	80.8
5-14 YEARS-----	8,165	15,348	19,218	11,796	40.8	76.4	95.2	58.2
15-24 YEARS-----	5,079	7,915	8,630	4,829	36.6	56.7	61.4	34.3
25-44 YEARS-----	6,579	11,069	11,534	7,862	30.4	51.3	53.5	36.5
45-64 YEARS-----	3,448	7,024	7,746	4,805	18.6	37.8	41.6	25.7
65 & OVER YEARS---	*	2,967	3,215	*	*	38.6	41.7	*
<u>FEMALE</u>								
ALL AGES----	36,714	63,229	71,447	45,174	37.5	64.4	72.5	45.7
UNDER 5 YEARS-----	6,781	10,864	10,933	7,128	67.8	109.3	110.8	73.0
5-14 YEARS-----	7,949	15,039	19,506	10,821	41.0	77.2	99.7	55.1
15-24 YEARS-----	6,367	10,814	10,502	7,454	42.0	70.7	68.0	47.8
25-44 YEARS-----	8,608	13,774	16,148	10,472	36.5	58.4	68.4	44.4
45-64 YEARS-----	4,771	8,440	10,547	6,748	23.9	42.0	52.3	33.3
65 & OVER YEARS---	*	4,298	3,813	2,551	*	43.6	38.5	25.6

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 24. INCIDENCE OF ACUTE CONDITIONS PER QUARTER AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS				NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER QUARTER			
ALL ACUTE CONDITIONS--	68,090	118,759	133,087	84,414	35.8	62.3	69.7	44.1
INFECTIVE AND PARASITIC DISEASES-----	8,152	11,464	15,787	12,505	4.3	6.0	8.3	6.5
RESPIRATORY CONDITIONS--	27,732	78,602	88,943	44,792	14.6	41.3	46.6	23.4
UPPER RESPIRATORY CONDITIONS-----	20,131	51,722	47,782	27,393	10.6	27.1	25.0	14.3
INFLUENZA-----	6,363	23,974	37,341	15,665	3.3	12.6	19.6	8.2
OTHER RESPIRATORY CONDITIONS-----	*	2,906	3,820	*	*	1.5	2.0	*
DIGESTIVE SYSTEM CONDITIONS-----	5,752	5,281	4,907	3,972	3.0	2.8	2.6	2.1
INJURIES-----	14,577	11,807	10,546	11,427	7.7	6.2	5.5	6.0
ALL OTHER ACUTE CONDITIONS-----	11,877	11,604	12,903	11,719	6.3	6.1	6.8	6.1
<u>MALE</u>								
ALL ACUTE CONDITIONS--	31,375	55,531	61,639	39,240	34.1	60.2	66.7	42.4
INFECTIVE AND PARASITIC DISEASES-----	3,569	5,295	7,171	6,106	3.9	5.7	7.8	6.6
RESPIRATORY CONDITIONS--	12,699	36,931	40,789	20,173	13.8	40.0	44.1	21.8
UPPER RESPIRATORY CONDITIONS-----	9,473	24,881	21,663	12,284	10.3	27.0	23.7	13.3
INFLUENZA-----	2,672	10,679	17,090	7,159	2.9	11.6	18.5	7.7
OTHER RESPIRATORY CONDITIONS-----	*	*	*	*	*	*	*	*
DIGESTIVE SYSTEM CONDITIONS-----	2,754	*	*	*	3.0	*	*	*
INJURIES-----	8,197	6,666	6,343	7,026	8.9	7.2	6.9	7.6
ALL OTHER ACUTE CONDITIONS-----	4,157	4,421	5,018	4,276	4.5	4.8	5.4	4.6
<u>FEMALE</u>								
ALL ACUTE CONDITIONS--	36,714	63,229	71,447	45,174	37.5	64.4	72.5	45.7
INFECTIVE AND PARASITIC DISEASES-----	4,583	6,169	8,616	6,399	4.7	6.3	8.7	6.5
RESPIRATORY CONDITIONS--	15,033	41,671	48,153	24,619	15.4	42.4	48.9	24.9
UPPER RESPIRATORY CONDITIONS-----	10,659	26,841	25,919	15,109	10.9	27.3	26.3	15.3
INFLUENZA-----	3,691	13,295	20,251	8,505	3.8	13.5	20.6	8.6
OTHER RESPIRATORY CONDITIONS-----	*	*	*	*	*	*	*	*
DIGESTIVE SYSTEM CONDITIONS-----	2,998	3,064	2,590	*	3.1	3.1	2.6	*
INJURIES-----	6,380	5,141	4,203	4,401	6.5	5.2	4.3	4.5
ALL OTHER ACUTE CONDITIONS-----	7,720	7,183	7,885	7,443	7.9	7.3	8.0	7.5

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 25. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER			
ALL AGES----	266,737	420,181	530,822	345,099	140.4	220.5	278.0	180.4
UNDER 5 YEARS-----	27,950	61,903	61,083	38,068	136.8	304.8	303.1	190.8
5-14 YEARS-----	47,382	89,722	139,636	80,067	120.3	226.8	351.3	200.5
15-24 YEARS-----	45,921	58,528	62,443	45,405	158.2	200.0	211.8	153.0
25-44 YEARS-----	61,570	87,193	106,539	74,130	136.1	192.9	236.0	164.2
45-64 YEARS-----	51,697	81,298	102,939	70,128	134.3	210.4	265.5	180.2
65 & OVER YEARS---	32,218	41,536	58,182	37,300	184.5	236.6	330.3	211.0
<u>MALE</u>								
ALL AGES----	124,876	200,135	238,265	157,059	135.6	216.9	257.9	169.8
UNDER 5 YEARS-----	13,397	32,819	31,437	19,918	128.5	316.5	305.5	195.5
5-14 YEARS-----	24,080	46,493	69,802	42,595	120.4	231.4	345.8	210.0
15-24 YEARS-----	19,574	23,104	27,559	21,657	141.0	165.5	196.2	153.8
25-44 YEARS-----	30,556	39,842	40,769	29,156	141.4	184.5	189.2	135.3
45-64 YEARS-----	20,774	41,293	44,796	27,112	112.2	222.3	240.6	145.2
65 & OVER YEARS---	16,495	16,584	23,902	16,622	215.4	215.7	310.4	215.5
<u>FEMALE</u>								
ALL AGES----	141,861	220,046	292,557	188,040	144.9	224.0	296.9	190.3
UNDER 5 YEARS-----	14,553	29,084	29,646	18,151	145.5	292.6	300.6	185.9
5-14 YEARS-----	23,302	43,229	69,834	37,472	120.2	222.0	357.1	190.7
15-24 YEARS-----	26,347	35,424	34,884	23,749	173.9	231.6	225.9	152.3
25-44 YEARS-----	31,013	47,351	65,770	44,974	131.4	200.6	278.7	190.6
45-64 YEARS-----	30,923	40,005	58,144	43,017	154.7	199.3	288.5	212.5
65 & OVER YEARS---	15,723	24,953	34,279	20,678	160.4	252.9	345.8	207.5

TABLE 26. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER			
ALL ACUTE CONDITIONS-	266,737	420,181	530,822	345,099	140.4	220.5	278.0	180.4
INFECTIVE AND PARASITIC DISEASES-----	30,110	46,226	69,719	51,406	15.8	24.3	36.5	26.9
RESPIRATORY CONDITIONS---	71,714	216,088	306,331	156,227	37.7	113.4	160.4	81.7
UPPER RESPIRATORY CONDITIONS-----	45,651	121,002	136,423	63,637	24.0	63.5	71.5	33.3
INFLUENZA-----	17,768	74,025	144,313	76,664	9.4	38.9	75.6	40.1
OTHER RESPIRATORY CONDITIONS-----	8,295	21,060	25,595	15,926	4.4	11.1	13.4	8.3
DIGESTIVE SYSTEM CONDITIONS-----	15,851	16,171	21,095	14,974	8.3	8.5	11.0	7.8
INJURIES-----	93,574	85,426	73,095	70,847	49.2	44.8	38.3	37.0
ALL OTHER ACUTE CONDITIONS-----	55,488	56,269	60,580	51,645	29.2	29.5	31.7	27.0
<u>MALE</u>								
ALL ACUTE CONDITIONS-	124,876	200,135	238,265	157,059	135.6	216.9	257.9	169.8
INFECTIVE AND PARASITIC DISEASES-----	13,565	22,064	32,443	25,463	14.7	23.9	35.1	27.5
RESPIRATORY CONDITIONS---	35,119	102,325	133,631	68,683	38.1	110.9	144.6	74.3
UPPER RESPIRATORY CONDITIONS-----	21,053	56,033	56,620	29,580	22.9	62.9	61.5	32.0
INFLUENZA-----	8,201	33,464	61,443	31,784	8.9	36.3	66.5	34.4
OTHER RESPIRATORY CONDITIONS-----	*	10,828	15,369	7,318	*	11.7	16.6	7.9
DIGESTIVE SYSTEM CONDITIONS-----	7,160	7,151	8,285	6,942	7.8	7.7	9.0	7.5
INJURIES-----	54,939	51,577	43,694	43,220	59.7	55.9	47.3	46.7
ALL OTHER ACUTE CONDITIONS-----	14,093	17,018	20,211	12,751	15.3	18.4	21.9	13.8
<u>FEMALE</u>								
ALL ACUTE CONDITIONS-	141,861	220,046	292,557	188,040	144.9	224.0	296.9	190.3
INFECTIVE AND PARASITIC DISEASES-----	16,545	24,162	37,276	25,943	16.9	24.6	37.6	26.3
RESPIRATORY CONDITIONS---	36,596	113,763	172,700	87,544	37.4	115.8	175.3	88.6
UPPER RESPIRATORY CONDITIONS-----	24,598	62,969	79,604	34,057	25.1	64.1	80.8	34.5
INFLUENZA-----	9,567	40,561	82,871	44,879	9.8	41.3	84.1	45.4
OTHER RESPIRATORY CONDITIONS-----	*	10,232	10,226	8,608	*	10.4	10.4	8.7
DIGESTIVE SYSTEM CONDITIONS-----	8,690	9,020	12,810	8,032	8.9	9.2	13.0	8.1
INJURIES-----	38,634	33,850	29,401	27,627	39.5	34.5	29.8	26.0
ALL OTHER ACUTE CONDITIONS-----	41,396	39,251	40,369	38,894	42.3	40.0	41.0	39.4

TABLE 27. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER			
ALL AGES----	99,646	191,761	251,978	153,912	52.4	100.6	132.0	80.5
UNDER 5 YEARS----	11,701	28,836	28,973	18,647	57.3	142.0	143.8	93.5
5-14 YEARS-----	18,872	42,432	68,285	37,657	47.9	107.2	171.8	94.3
15-24 YEARS-----	23,825	30,258	32,328	20,793	82.1	103.4	109.6	70.1
25-44 YEARS-----	19,443	42,066	51,206	35,178	43.0	93.1	113.4	77.9
45-64 YEARS-----	16,550	32,338	44,707	28,809	43.0	83.7	115.3	74.0
65 & OVER YEARS---	9,254	15,831	26,480	12,828	53.0	90.2	157.3	72.6
<u>MALE</u>								
ALL AGES----	45,639	89,037	104,382	61,545	49.6	96.5	113.0	66.5
UNDER 5 YEARS----	*	14,909	14,221	9,182	*	143.8	138.2	90.1
5-14 YEARS-----	10,370	21,082	31,508	17,642	51.9	104.9	156.1	87.0
15-24 YEARS-----	10,307	10,186	13,566	8,070	74.2	72.9	96.6	57.3
25-44 YEARS-----	9,069	19,426	16,743	12,948	42.0	89.9	77.7	60.1
45-64 YEARS-----	6,008	15,850	17,175	8,188	32.4	85.3	92.2	43.9
65 & OVER YEARS---	*	7,583	11,168	*	*	98.6	145.0	*
<u>FEMALE</u>								
ALL AGES----	54,006	102,724	147,596	92,367	55.2	104.6	149.8	93.5
UNDER 5 YEARS----	6,753	13,928	14,752	9,465	67.5	140.1	149.6	96.9
5-14 YEARS-----	8,502	21,350	36,777	20,015	43.9	109.7	188.0	101.9
15-24 YEARS-----	13,519	20,072	18,761	12,723	89.2	131.2	121.5	81.6
25-44 YEARS-----	10,374	22,639	34,463	22,230	43.9	95.9	146.0	94.2
45-64 YEARS-----	10,542	16,488	27,531	20,621	52.7	82.1	136.6	101.9
65 & OVER YEARS---	*	8,247	15,312	7,314	*	83.6	154.5	73.4

TABLE 28. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND CONDITION GROUP	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER			
ALL ACUTE CONDITIONS-	99,646	191,761	251,978	153,912	52.4	100.6	132.0	80.5
INFECTIVE AND PARASITIC DISEASES-----	17,827	23,223	34,335	28,108	9.4	12.2	18.0	14.7
RESPIRATORY CONDITIONS---	32,584	106,057	156,184	79,519	17.1	55.7	81.8	41.6
UPPER RESPIRATORY CONDITIONS-----	17,703	51,352	57,515	26,814	9.3	27.0	30.1	14.0
INFLUENZA-----	10,787	42,789	81,450	43,705	5.7	22.5	42.7	22.8
OTHER RESPIRATORY CONDITIONS-----	*	11,916	17,218	9,000	*	6.3	9.0	4.7
DIGESTIVE SYSTEM CONDITIONS-----	8,003	9,522	12,659	7,190	4.2	5.0	6.6	3.8
INJURIES-----	21,709	26,100	23,018	18,866	11.4	13.7	12.1	9.9
ALL OTHER ACUTE CONDITIONS-----	19,523	26,860	25,783	20,229	10.3	14.1	13.5	10.6
<u>MALE</u>								
ALL ACUTE CONDITIONS-	45,639	89,037	104,382	61,545	49.6	96.5	113.0	66.5
INFECTIVE AND PARASITIC DISEASES-----	8,852	11,445	14,502	13,450	9.6	12.4	15.7	14.5
RESPIRATORY CONDITIONS---	16,319	50,847	65,712	33,397	17.7	55.1	71.1	36.1
UPPER RESPIRATORY CONDITIONS-----	8,317	25,257	22,583	11,648	9.0	27.4	24.4	12.6
INFLUENZA-----	*	19,292	33,042	18,178	*	20.9	35.8	19.7
OTHER RESPIRATORY CONDITIONS-----	*	6,298	10,088	*	*	6.8	10.9	*
DIGESTIVE SYSTEM CONDITIONS-----	*	*	*	*	*	*	*	*
INJURIES-----	12,136	13,873	11,818	8,098	13.2	15.0	12.8	8.8
ALL OTHER ACUTE CONDITIONS-----	*	8,069	7,507	*	*	8.7	8.1	*
<u>FEMALE</u>								
ALL ACUTE CONDITIONS-	54,006	102,724	147,596	92,367	55.2	104.6	149.8	93.5
INFECTIVE AND PARASITIC DISEASES-----	8,975	11,777	19,833	14,657	9.2	12.0	20.1	14.8
RESPIRATORY CONDITIONS---	16,265	55,210	90,471	46,123	16.6	56.2	91.8	46.7
UPPER RESPIRATORY CONDITIONS-----	9,387	26,095	34,932	15,166	9.6	26.6	35.5	15.3
INFLUENZA-----	*	23,496	48,409	25,528	*	23.9	49.1	25.8
OTHER RESPIRATORY CONDITIONS-----	*	*	7,130	*	*	*	7.2	*
DIGESTIVE SYSTEM CONDITIONS-----	*	*	7,817	*	*	*	7.9	*
INJURIES-----	9,573	12,227	11,200	10,768	9.8	12.4	11.4	10.9
ALL OTHER ACUTE CONDITIONS-----	14,740	18,791	18,276	16,201	15.1	19.1	18.5	16.4

TABLE 29. INCIDENCE OF ACUTE RESPIRATORY CONDITIONS AND ASSOCIATED DISABILITY DAYS PER QUARTER AND NUMBER OF ACUTE RESPIRATORY CONDITIONS PER 100 PERSONS PER QUARTER, BY AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
	INCIDENCE OF ACUTE RESPIRATORY CONDITIONS IN THOUSANDS				NUMBER OF ACUTE RESPIRATORY CONDITIONS PER 100 PERSONS PER QUARTER			
ALL AGES----	27,732	78,602	88,943	44,792	14.6	41.3	46.6	23.4
UNDER 5 YEARS-----	6,829	15,767	14,646	7,948	33.4	77.6	72.7	39.8
5-14 YEARS-----	6,912	20,794	26,014	11,683	17.6	52.6	65.5	29.3
15-24 YEARS-----	3,997	11,834	12,564	6,939	13.8	40.4	42.6	23.4
25-44 YEARS-----	5,044	15,672	17,676	9,449	11.2	34.7	39.2	20.9
45-64 YEARS-----	3,346	10,426	13,190	6,291	8.7	27.0	34.0	16.2
65 & OVER YEARS---	*	4,109	4,853	*	*	23.4	27.6	*
	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER			
ALL AGES----	71,714	216,088	306,331	156,227	37.7	113.4	160.4	81.7
UNDER 5 YEARS-----	15,169	40,743	36,966	20,090	74.3	200.6	183.4	100.7
5-14 YEARS-----	18,427	51,996	85,064	35,036	46.8	131.4	214.0	87.8
15-24 YEARS-----	9,242	28,455	34,374	18,317	31.8	97.3	116.6	61.7
25-44 YEARS-----	13,366	38,582	54,222	30,108	29.6	85.4	120.1	66.7
45-64 YEARS-----	9,240	37,689	61,805	32,201	24.0	97.5	159.4	82.7
65 & OVER YEARS---	6,270	18,623	33,900	20,474	35.9	106.1	192.5	115.8
	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER			
ALL AGES----	32,584	106,057	156,184	79,519	17.1	55.7	81.8	41.6
UNDER 5 YEARS-----	6,258	16,919	17,503	9,228	30.6	83.3	86.9	46.3
5-14 YEARS-----	8,652	27,180	46,163	19,350	22.0	68.7	116.2	48.5
15-24 YEARS-----	*	17,311	20,428	11,403	*	59.2	69.3	38.4
25-44 YEARS-----	*	19,638	29,091	17,294	*	43.4	64.4	38.3
45-64 YEARS-----	*	18,091	28,832	15,222	*	46.8	74.3	39.1
65 & OVER YEARS---	*	6,918	14,166	7,023	*	39.4	80.4	39.7

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 30. INCIDENCE OF INJURIES AND ASSOCIATED DISABILITY DAYS PER QUARTER AND NUMBER OF INJURIES PER 100 PERSONS PER QUARTER, BY AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
	INCIDENCE OF INJURIES IN THOUSANDS				NUMBER OF INJURIES PER 100 PERSONS PER QUARTER			
ALL AGES----	14,577	11,807	10,546	11,427	7.7	6.2	5.5	6.0
UNDER 15 YEARS----	4,784	3,246	2,926	4,487	8.0	5.4	4.9	7.5
15-44 YEARS-----	7,233	5,567	5,107	4,383	9.7	7.5	6.8	5.9
45 & OVER YEARS---	2,560	2,994	2,513	2,556	4.6	5.3	4.5	4.5
	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS				DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER			
ALL AGES----	93,574	85,428	73,095	70,847	49.2	44.8	38.3	37.0
UNDER 15 YEARS----	15,401	12,141	8,396	10,816	25.8	20.3	14.0	18.1
15-44 YEARS-----	43,156	34,391	31,790	32,744	58.1	46.2	42.6	43.8
45 & OVER YEARS---	35,016	38,895	32,910	27,287	62.6	69.2	58.4	48.2
	DAYS OF BED DISABILITY IN THOUSANDS				DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER			
ALL AGES----	21,709	26,100	23,018	18,866	11.4	13.7	12.1	9.9
UNDER 15 YEARS----	*	*	*	*	*	*	*	*
15-44 YEARS-----	12,689	11,562	9,914	8,432	17.1	15.5	13.3	11.3
45 & OVER YEARS---	*	9,416	11,180	7,271	*	16.8	19.8	12.8

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 31. POPULATION USED IN OBTAINING RATES SHOWN IN THIS PUBLICATION, BY RESIDENCE, REGION, SEX, AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	TOTAL	ALL SMSAS	NON-SMSA NONFARM	NON-SMSA FARM	NORTH-EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>		POPULATION IN THOUSANDS						
ALL AGES-----	190,716	122,000	57,757	10,954	47,503	53,133	58,891	31,184
UNDER 5 YEARS-----	20,210	12,841	6,394	975	4,612	5,495	6,670	3,434
5-14 YEARS-----	39,658	24,959	12,228	2,470	9,271	11,080	12,659	6,648
15-24 YEARS-----	29,365	18,766	8,986	1,613	6,827	8,248	9,580	4,710
25-44 YEARS-----	45,185	30,109	13,001	2,076	11,466	12,309	13,749	7,661
45 & OVER YEARS-----	56,292	35,324	17,148	3,820	15,327	16,002	16,233	8,730
45-64 YEARS-----	38,713	24,753	11,244	2,717	10,453	10,936	11,210	6,113
65 & OVER YEARS-----	17,578	10,571	5,904	1,103	4,873	5,065	5,022	2,617
UNDER 6 YEARS-----	24,370	15,487	7,706	1,177	5,531	6,635	8,025	4,179
6-16 YEARS-----	42,470	26,619	13,068	2,782	9,913	11,952	13,637	6,968
17-44 YEARS-----	67,579	44,570	19,834	3,175	16,732	18,544	20,996	11,306
<u>MALE</u>								
ALL AGES-----	92,323	58,771	27,914	5,637	22,939	25,824	28,299	15,261
UNDER 5 YEARS-----	10,318	6,517	3,290	512	2,365	2,768	3,432	1,754
5-14 YEARS-----	20,141	12,592	6,252	1,297	4,764	5,565	6,342	3,471
15-24 YEARS-----	13,994	8,882	4,276	836	3,301	3,932	4,533	2,227
25-44 YEARS-----	21,581	14,427	6,152	1,002	5,449	5,947	6,519	3,666
45 & OVER YEARS-----	26,286	16,353	7,944	1,991	7,060	7,612	7,473	4,143
45-64 YEARS-----	18,597	11,857	5,323	1,417	5,001	5,318	5,313	2,966
65 & OVER YEARS-----	7,691	4,496	2,621	574	2,060	2,294	2,160	1,178
UNDER 6 YEARS-----	12,392	7,852	3,929	612	2,850	3,315	4,081	2,146
6-16 YEARS-----	21,601	13,403	6,717	1,481	5,080	6,033	6,862	3,625
17-44 YEARS-----	32,042	21,163	9,325	1,554	7,948	8,864	9,883	5,347
<u>FEMALE</u>								
ALL AGES-----	98,387	63,229	29,842	5,316	24,564	27,309	30,592	15,922
UNDER 5 YEARS-----	9,892	6,324	3,104	464	2,247	2,726	3,238	1,681
5-14 YEARS-----	19,516	12,367	5,976	1,173	4,507	5,515	6,318	3,176
15-24 YEARS-----	15,371	9,884	4,710	777	3,526	4,316	5,046	2,483
25-44 YEARS-----	23,605	15,683	6,848	1,074	6,017	6,362	7,230	3,995
45 & OVER YEARS-----	30,003	18,971	9,204	1,829	8,267	8,390	8,760	4,587
45-64 YEARS-----	20,116	12,895	5,921	1,300	5,453	5,618	5,898	3,148
65 & OVER YEARS-----	9,887	6,075	3,283	529	2,814	2,771	2,862	1,439
UNDER 6 YEARS-----	11,977	7,635	3,777	565	2,680	3,320	3,944	2,033
6-16 YEARS-----	20,869	13,216	6,352	1,301	4,833	5,919	6,775	3,343
17-44 YEARS-----	35,537	23,407	10,510	1,621	8,784	9,681	11,113	5,959

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States, in Current Population Reports, Series P-20, P-25, and P-60.

TABLE 32. POPULATION OF CURRENTLY EMPLOYED PERSONS, BY SEX AND AGE: UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

AGE	BOTH SEXES	MALE	FEMALE
POPULATION IN THOUSANDS			
ALL AGES-17 & OVER-----	73,123	47,252	25,870
17-44-----	43,715	28,329	15,386
45 & OVER-----	29,408	18,923	10,485

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States, in Current Population Reports, Series P-20, P-25, and P-60; and Bureau of Labor Statistics monthly report, Employment and Earnings.

TABLE 33. POPULATION USED IN OBTAINING RATES SHOWN IN THIS PUBLICATION, BY QUARTER, SEX, AND AGE:
UNITED STATES, JULY 1965-JUNE 1966

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
<u>BOTH SEXES</u>	POPULATION IN THOUSANDS			
ALL AGES-----	190,025	190,532	190,923	191,296
UNDER 5 YEARS-----	20,427	20,307	20,152	19,951
5-14 YEARS-----	39,381	39,564	39,744	39,926
15-24 YEARS-----	29,033	29,258	29,487	29,672
25-44 YEARS-----	45,225	45,203	45,148	45,152
45-64 YEARS-----	38,500	38,647	38,779	38,915
65 & OVER YEARS-----	17,459	17,554	17,614	17,679
<u>MALE</u>				
ALL AGES-----	92,101	92,284	92,390	92,488
UNDER 5 YEARS-----	10,428	10,368	10,289	10,186
5-14 YEARS-----	19,999	20,093	20,186	20,281
15-24 YEARS-----	13,885	13,963	14,045	14,079
25-44 YEARS-----	21,616	21,598	21,547	21,556
45-64 YEARS-----	18,515	18,574	18,622	18,672
65 & OVER YEARS-----	7,658	7,689	7,701	7,715
<u>FEMALE</u>				
ALL AGES-----	97,924	98,248	98,533	98,808
UNDER 5 YEARS-----	9,999	9,939	9,863	9,765
5-14 YEARS-----	19,382	19,471	19,558	19,645
15-24 YEARS-----	15,148	15,295	15,442	15,593
25-44 YEARS-----	23,609	23,605	23,600	23,597
45-64 YEARS-----	19,985	20,073	20,157	20,244
65 & OVER YEARS-----	9,801	9,865	9,913	9,964

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States, in Current Population Reports, Series P-20, P-25, and P-60.

APPENDIX I

TECHNICAL NOTES ON METHODS

Background of This Report

This report is one of a series of statistical reports prepared by the National Health Survey. It is based on information collected in a continuing nationwide sample of households in the Health Interview Survey, a major part of the program.

The Health Interview Survey utilizes a questionnaire which, in addition to personal and demographic characteristics, obtains information on illnesses, injuries, chronic conditions and impairments, and other health topics. As data relating to each of these various broad topics are tabulated and analyzed, separate reports are issued which cover one or more of the specific topics. The present report is based on the consolidated sample for 52 weeks of interviewing ending June 1966.

The population covered by the sample for the Health Interview Survey is the civilian, noninstitutional population of the United States living at the time of the interview. The sample does not include members of the Armed Forces, U.S. nationals living in foreign countries, or crews of vessels. It should also be noted that the estimates shown do not represent a complete inventory of injuries for the specified calendar period since no adjustment has been made for persons who incurred injuries during the 2-week-recall period but who died prior to the interview.

Statistical Design of the Health Interview Survey

General plan.—The sampling plan of the survey follows a multistage probability design which permits a continuous sampling of the civilian population of the United States. The first stage of this design consists of drawing a sample of 357 from about 1,900 geographically defined primary sampling units (PSU's) into which the United States has been divided. A PSU is a county, a group of contiguous counties, or a standard metropolitan statistical area.

With no loss in general understanding, the remaining stages can be combined and treated in this discussion as an ultimate stage. Within PSU's then, ultimate stage units called segments are defined in such a manner that each segment contains an expected nine households. A segment consists of a cluster of neighboring households or addresses. Two

general types of segments are used: (1) area segments which are defined geographically, and (2) B segments which are defined from a list of addresses from the Decennial Census and Survey of Construction. Each week a random sample of about 90 segments is drawn. In the approximately 800 households in these segments, household members are interviewed concerning factors related to health.

Since the household members interviewed each week are a representative sample of the population, samples for successive weeks can be combined into larger samples. Thus the design permits both continuous measurement of characteristics of high incidence or prevalence in the population and, through the larger consolidated samples, more detailed analysis of less common characteristics and smaller categories. The continuous collection has administrative and operational advantages as well as technical assets, since it permits field work to be handled with an experienced, stable staff.

Sample size and geographic detail.—The national sample plan for the 12-month period ending June 1966 included about 134,000 persons from 42,000 households in about 4,700 segments.

The overall sample was designed in such a fashion that tabulations could be provided for each of the major geographic regions and for urban and rural sectors of the United States.

Collection of data.—Field operations for the household survey are performed by the Bureau of the Census under specifications established by the National Center for Health Statistics. In accordance with these specifications the Bureau of the Census selects the sample, conducts the field interviewing as an agent of the Center, and performs a manual edit and coding of the questionnaires. The Health Interview Survey, using Center electronic computers, carries out further editing and tabulates the edited data.

Estimating methods.—Each statistic produced by the survey—for example, the number of work-loss days occurring in a specified period—is the result of two stages of ratio estimation. In the first of these, the control factor is the ratio of the 1960 decennial population count to the 1960 estimated population in the National Health Survey's first-stage sample of PSU's. These factors are applied for some 25 color-residence classes.

Later, ratios of sample-produced estimates of the population to official Bureau of the Census figures for current population in about 60 age-sex-color classes are computed, and serve as second-stage factors for ratio estimating.

The effect of the ratio-estimating process is to make the sample more closely representative of the population by age, sex, color, and residence, thus reducing sampling variance.

As noted, each week's sample represents the population living during that week and characteristics of that population. Consolidation of samples over a time period, say a calendar quarter, produces estimates of average characteristics of the U.S. population for that calendar quarter. Similarly, population data for a year are averages of the four quarterly figures.

For statistics measuring the number of occurrences during a specified time period, such as the number of cases of influenza, a similar computational procedure is used, but the statistics have a different interpretation. For these items, the questionnaire asks for the respondent's experience over the 2 calendar weeks prior to the week of interview. In such instances the estimated quarterly total for the statistic is simply 6.5 times the average 2-week estimate produced by the 13 successive samples taken during the period. The annual total is the sum of the four quarters. Thus, the experience of persons *interviewed during a year*—experience which actually occurred for each person in a 2-calendar-week interval prior to week of interview—is treated as though it measured the total of such *experience during the year*. Such interpretation leads to no significant bias.

General Qualifications

Nonresponse.—Data were adjusted for nonresponse by a procedure which imputes to persons in a household which was not interviewed the characteristics of persons in households in the same segment which were interviewed. The total noninterview rate was 5 percent—1 percent was refusal, and the remainder was primarily due to the failure to find any eligible household respondent after repeated trials.

The interview process.—The statistics presented in this report are based on replies secured in interviews of persons in the sampled households. Each person 19 years of age and over, available at the time of interview, was interviewed individually. Proxy respondents within the household were employed for children and for adults not available at the time of the interview, provided the respondent was closely related to the person about whom information was being obtained.

There are limitations to the accuracy of diagnostic and other information collected in household

interviews. For diagnostic information, the household respondent can, at best, pass on to the interviewer only the information the physician has given to the family. For conditions not medically attended, diagnostic information is often no more than a description of symptoms. However, other facts, such as the number of disability days caused by the condition, can be obtained more accurately from household members than from any other source, since only the persons concerned are in a position to report this information.

Rounding of numbers.—The original tabulations on which the data in this report are based show all estimates to the nearest whole unit. All consolidations were made from the original tabulations using the estimates to the nearest unit. In the final published tables the figures are rounded to the nearest thousand, although these are not necessarily accurate to that detail. Devised statistics such as rates and percent distributions are computed after the estimates on which these are based have been rounded to the nearest thousand.

Population figures.—Some of the published tables include population figures for specified categories. Except for certain overall totals by age and sex, which are adjusted to independent estimates, these figures are based on the sample of households in the National Health Survey. These are given primarily to provide denominators for rate computation, and for this purpose are more appropriate for use with the accompanying measures of health characteristics than other population data that may be available. In some instances these will permit users to recombine published data into classes more suitable to their specific needs. With the exception of the overall totals by age and sex, mentioned above, the population figures differ from corresponding figures (which are derived from different sources) published in reports of the Bureau of the Census. For population data for general use, see the official estimates presented in Bureau of the Census reports in the P-20, P-25, and P-60 series.

Reliability of Estimates

Since the estimates are based on a sample, they will differ somewhat from the figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and interviewing personnel and procedures. As in any survey, the results are also subject to measurement error.

The standard error is primarily a measure of sampling variability, that is, the variations that might occur by chance because only a sample of the population is surveyed. As calculated for this report, the standard error also reflects part of the variation which arises in the measurement process. It does

not include estimates of any biases which might lie in the data. The chances are about 68 out of 100 that an estimate from the sample would differ from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference would be less than twice the standard error and about 99 out of 100 that it would be less than 2½ times as large.

The relative standard error of an estimate is obtained by dividing the standard error of the estimate by the estimate itself, and is expressed as a per-

centage of the estimate. Included in this Appendix are charts from which the relative standard errors can be determined for estimates shown in the report. A description of the classes of statistics used in the Health Interview Survey and general rules for determining relative sampling errors are presented in Appendix I of "Current Estimates" (*Vital and Health Statistics*, Series 10, No. 37).

The following guide indicates the appropriate rules and charts to be used in deriving relative standard errors for estimates shown in this report.


Guide to Use of Relative Standard Error Charts

The code shown below identifies the appropriate curve to be used in estimating the relative standard error of the statistic described. The four components of each code describe the statistic as follows: (A)=

aggregate, P=percentage; (2) the number of calendar quarters of data collection; (3) the type of the statistic; and (4) the range of the statistic as described in *Vital and Health Statistics*, Series 10, No. 37.


Statistic	Use:		
	Rule	Code	on page
Number of:			
Persons by residence or region-----	1	A4AN	50
Acute conditions:			
Per quarter-----	1	A1BN	51
Per year-----	1	A4BN	50
Disability days:			
Per quarter-----	1	A1BW	52
Per year-----	1	A4BW	50
Persons in the U.S. population or in any age-sex category thereof-----	Not subject to sampling error		
Percentage distribution of:			
Persons by residence or region-----	2	P4AN-M	53
Disability days in a year-----	2	P4BW	54
Acute conditions in a year-----	2	P4BN-M	55
Incidence rates for acute conditions:			
Per 100 total U.S. population or per 100 persons in any age-sex group of the total U.S. population:			
Per quarter-----	4 (a)	A1BN	51
Per year-----	4 (a)	A4BN	50
Per 100 persons in any other population group per year-----	4 (b)	Numer.: } Denom.: }	A4BN } A4AN } 50 50
Number of disability days:			
Per 100 total U.S. population or per 100 persons in any age-sex group of the total U.S. population:			
Per quarter-----	4 (a)	A1BW	52
Per year-----	4 (a)	A4BW	50
Per 100 persons in any other population group per year-----	4 (b)	Numer.: } Denom.: }	A4BW } A4AN } 50 50
Per case for acute conditions:			
Per quarter-----	4 (b)	Numer.: } Denom.: }	A1BW } A1BN } 52 51
Per year-----	4 (b)	Numer.: } Denom.: }	A4BW } A4AN } 50 50

Relative standard errors for aggregates based on four quarters of data collection
for data of all types and ranges




Example of use of chart: An aggregate of 2,000,000 (on scale at bottom of chart) for a Narrow range Type A statistic (code: A4AN) has a relative standard error of 3.6 percent, (read from scale at left side of chart), or a standard error of 72,000 (3.6 percent of 2,000,000). For a Wide range Type B statistic (code: A4BW), an aggregate of 6,000,000 has a relative error of 16.0 percent or a standard error of 960,000 (16 percent of 6,000,000).

Relative standard errors for aggregates based on one quarter of data for type B data, narrow range




Example of use of chart: An aggregate of 6,000,000 (on scale at bottom of chart) for a Narrow range Type B statistic has a relative standard error of 19.3 percent, read from scale at left side of chart, or a standard error of 1,158,000 (19.3 percent of 6,000,000).

Relative standard errors for aggregates based on one quarter of data collection for type B data, wide range




Example of use of chart: An aggregate of 20,000,000 (on scale at bottom of chart) for a wide range type B statistic has a relative standard error of 16.0 percent, read from scale at left side of chart, or a standard error of 3,200,000 (16.0 percent of 20,000,000).

Relative standard errors for percentages based on four quarters of data collection
 for type A data, Narrow and Medium range
 (Base of percentage shown on curves in millions)




Example of use of chart: An estimate of 20 percent (on scale at bottom of chart) based on an estimate of 10,000,000 has a relative standard error of 3.2 percent (read from the scale at the left side of the chart), the point at which the curve for a base of 10,000,000 intersects the vertical line for 20 percent. The standard error in percentage points is equal to 20 percent X 3.2 percent or 0.64 percentage points.

Relative standard errors for percentages based on four quarters of data collection
for type B data, Wide range
(Base of percentage shown on curves in millions)



Example of use of chart: An estimate of 20 percent (on scale at bottom of chart) based on an estimate of 10,000,000 has a relative standard error of 24.5 percent (read from scale at the left side of the chart), the point at which the curve for a base of 10,000,000 intersects the vertical line for 20 percent. The standard error in percentage points is equal to 20 percent X 24.5 percent or 4.9 percentage points.

Relative standard errors for percentages based on four quarters of data collection
 for type B data, Narrow and Medium range
 (Base of percentage shown on curves in millions)



Example of use of chart: An estimate of 20 percent (on scale at bottom of chart) based on an estimate of 10,000,000 has a relative standard error of 17.0 percent (read from scale at the left side of the chart), the point at which the curve for a base of 10,000,000 intersects the vertical line for 20 percent. The standard error in percentage points is equal to 20 percent X 17.0 percent or 3.4 percentage points.

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Terms Relating to Conditions

Condition.—A morbidity condition, or simply a condition, is any entry on the questionnaire which describes a departure from a state of physical or mental well-being. It results from a positive response to one of a series of "illness-recall" questions. In the coding and tabulating process, conditions are selected or classified according to a number of different criteria, such as, whether they were medically attended; whether they resulted in disability; whether they were acute or chronic; or according to the type of disease, injury, impairment, or symptom reported. For the purposes of each published report or set of tables, only those conditions recorded on the questionnaire which satisfy certain stated criteria are included.

Acute conditions are classified by type according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the code more suitable for a household-interview-type survey.

Acute condition.—An acute condition is defined as a condition which has lasted less than 3 months and which has involved either medical attention or restricted activity. Because of the procedures used to estimate incidence, the acute conditions included in this report are the conditions which had their onset during the 2 weeks prior to the interview week and which involved either medical attention or restricted activity during that 2-week period. However, it excludes certain conditions which are always classified as chronic (listed below) even though the onset occurred within 3 months.

Conditions always classified as chronic:

Asthma	Stomach ulcer
Hay fever	Any other chronic stomach trouble
Tuberculosis	Kidney stones or chronic kidney trouble
Chronic bronchitis	Arthritis or rheumatism
Repeated attacks of sinus trouble	Mental illness
Rheumatic fever	Diabetes
Hardening of the arteries	Thyroid trouble or goiter
High blood pressure	Any allergy
Heart trouble	Epilepsy
Stroke	Chronic nervous trouble
Trouble with varicose veins	Cancer

Hemorrhoids or piles	Chronic skin trouble
Tumor, cyst, or growth	Hernia or rupture
Chronic gallbladder or liver trouble	Prostate trouble
Deafness or serious trouble with hearing	Paralysis of any kind
Serious trouble with seeing, even when wearing glasses	Repeated trouble with back or spine
Cleft palate	Club foot
Any speech defect	Permanent stiffness or any deformity of the foot, leg, fingers, arm, or back
Missing fingers, hand, or arm—toes, foot, or leg	Condition present since birth
Palsy	

Condition groups.—Conditions are classified according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the code more suitable for a household-interview survey. In this report, all tables which have data classified by type of condition employ a five-category regrouping plus several selected subgroups. The International Classification code numbers included in each category are shown below:

Condition Groups	International Classification Code Number (excluding chronic inclusions)
Infective and parasitic diseases-----	021-128
Common childhood diseases-----	055, 065-097, 069
The virus, n.o.c.-----	***
Other infective and parasitic diseases-----	020-055, 057-094, 089, 090-128
Respiratory conditions-----	470-501, 511, 517-525, 527, 783
Upper respiratory conditions-----	470-475, 511, 517
Common cold-----	470
Other acute upper respiratory conditions-----	471-475, 511, 517
Influenza-----	480-483
Influenza with digestive manifestations-----	482
Other influenza-----	480, 481, 482
Other respiratory conditions-----	430-501, 518-525, 527, 783
Pneumonia-----	430-453
Erythema-----	500, 501
Other acute respiratory conditions-----	515-525, 527, 783
Digestive system conditions-----	530-539, 543-553, 570, 571, 575-587, 784, 785
Dental conditions-----	530-525
Functional and symptomatic upper gastrointestinal disorders, n.o.c.-----	514, 784
Other digestive system conditions-----	535-539, 543, 545-553, 570, 571, 573-587, 785
Injuries-----	N890-N895, N893-N895, N900-N904, N906-N909
Fractures, dislocations, sprains, and strains-----	N850-N848
Fracture and dislocations-----	N850-N839
Sprains and strains-----	N840-N848
Open wounds and lacerations-----	N870-N885, N890-N895, N900-N908
Contusions and superficial injuries-----	N910-N929
Other current injuries-----	N850-N859, N930-N934, N936-N939
All other acute conditions-----	All other acute code numbers
Diseases of the ear-----	390-396, 781.3
Headaches-----	781
Contourinary disorders-----	590-637, 766, 789
Deliveries and disorders of pregnancy and the puerperium-----	640-699
Diseases of the skin-----	690-716
Diseases of the musculoskeletal system-----	726-743, 787
All other acute conditions-----	Other acute code numbers

Onset of condition.—A condition is considered to have had its onset when it was first noticed. This could be the time the person first felt sick or became injured, or it could be the time when the person or his family was first told by a physician that he had a condition of which he was previously unaware.

Incidence of conditions.—The incidence of conditions is the estimated number of conditions having their onset in a specified time period. As previously mentioned, minor acute conditions, involving neither restricted activity nor medical attention are excluded from the statistics. The incidence data shown in some reports are further limited to various subclasses of conditions, such as "incidence of conditions involving bed disability."

Activity-restricting condition.—An activity restricting condition is a condition which has caused at least 1 day of restricted activity during the 2 calendar weeks before the interview week. (See definition of "Restricted-activity day.") The incidence of acute activity-restricting conditions is estimated from the number of such conditions reported as having started in the 2-week period, but a condition which did not result in restricted activity until after the end of the 2-week period in which it had its onset is not included.

Bed-disabling condition.—A condition involving at least 1 day of bed disability is called a bed-disabling condition. (See definition of "Bed-disability day.") The incidence of acute bed-disabling conditions is defined in a manner analogous to the incidence of acute activity-restricting conditions.

Medically attended condition.—A condition is considered to be medically attended if a physician has been consulted about it either at its onset or at any time thereafter. Medical attention includes consultation either in person or by telephone for treatment or advice. Advice from the physician transmitted to the patient through the nurse is counted as well as visits to physicians in clinics or hospitals. If during the course of a single visit the physician is consulted about more than one condition for each of several patients, each condition of each patient is counted as medically attended.

Discussions of a child's condition by the physician and a responsible member of the household are considered as medical attention even if the child was not seen at that time.

For the purpose of this definition, the term "physician" includes doctors of medicine and osteopathic physicians.

Terms Relating to Disability

Disability.—Disability is the general term used to describe any temporary or long-term reduction of a person's activity as a result of an acute or chronic condition.

Disability days are classified according to whether they are days of restricted activity, bed-days, work-loss days, or school-loss days. All days of bed disability are, by definition, days of restricted activity. The converse form of this statement is, of course, not true. Days lost from work and days lost from school are also days of restricted activity for the working and school-age populations. Hence, restricted activity is the most inclusive term used in describing disability days.

Condition-days of restricted activity, bed disability, etc.—Condition-days of restricted activity, bed disability, and so forth are days of the various forms of disability associated with any one condition. Since any particular day of disability may be associated with more than one condition, the sum of days for all conditions adds to more than the total number of person-days of disability.

Restricted-activity day.—A day of restricted activity is one on which a person substantially reduces the amount of activity normal for that day because of a specific illness or injury. The type of reduction varies with the age and occupation of the individual as well as with the day of the week or season of the year. Restricted activity covers the range from substantial reduction to complete inactivity for the entire day.

Bed-disability day.—A day of bed disability is one on which a person stays in bed for all or most of the day because of a specific illness or injury. It is considered to be a day only if the period of bed disability includes more than half of the daylight hours. All hospital days for inpatients are considered to be days of bed disability even if the patient was not actually in bed at the hospital.

Work-loss day.—A day lost from work is a normal working day on which a person did not work at his job or business because of a specific illness or injury. If the person's regular work day is less than a whole day and the entire work day was lost, it would be counted as a whole work day lost. The number of days lost from work is determined only for persons 17 years of age or over who reported that at any time during the 2-week period covered by the interview they either worked at, or had a job

or business. (See definition of "Currently employed persons.")

School-loss day.—A day lost from school is a normal school day on which a child did not attend school because of a specific illness or injury. The number of days lost from school is determined only for children 6-16 years of age.

Demographic Terms

Age.—The age recorded for each person is the age at last birthday. Age is recorded in single years and grouped in a variety of distributions depending upon the purpose of the table.

Quarter.—The quarters used by the National Health Survey are actually 13-week periods rather than 3 calendar months. Since each 13-week period begins on a Monday and ends on a Sunday, the actual dates of the beginning and end of each 13-week period may overlap into another calendar quarter. Therefore, the time periods in the table headings are the approximate rather than the precise periods during which the interviewing was conducted.

Currently employed persons.—Currently employed persons are all persons 17 years of age or over who reported that at any time during the 2-week period covered by the interview they either worked at or had a job or business. Current employment includes paid work as an employee of someone else, self-employment in business, farming, or professional practice, and unpaid work in a family business or farm. Persons who were temporarily absent from their job or business because of a temporary illness, vacation, strike, or bad weather are considered as currently employed if they expected to work as soon as the particular event causing their absence no longer existed.

Free-lance workers are considered as having a job if they had a definite arrangement with one or more employees to work for pay according to a weekly or monthly schedule, either full time or part time. Excluded from the currently employed population are such persons who have no definite employment schedule but who work only when their services are needed.

Also excluded from the currently employed population are (1) persons who were not working, even though having a job or business, but were on layoff or looking for work, (2) persons receiving revenue from an enterprise in whose operation they did not participate, (3) persons doing housework or charity work for which they received no pay, and (4) seasonal workers during the unemployment season.

The number of currently employed persons estimated by the National Health Survey (NHS) will differ from the estimates prepared by the Current

Population Survey (CPS), Bureau of the Census, for several reasons. In addition to sampling variability they include three primary conceptual differences, namely: (1) NHS estimates are for persons 17 years of age or over; CPS estimates are for persons 14 years of age or over. (2) NHS uses a 2-week-reference period while CPS uses a 1-week-reference period. (3) NHS is a continuing survey with separate samples taken weekly, while CPS is a monthly sample taken for the survey week which includes the 12th of the month.

Residence.—The place of residence of a member of the civilian, noninstitutional population is classified as inside a standard metropolitan statistical area (SMSA) or outside an SMSA according to farm or nonfarm residence.

Standard metropolitan statistical areas.—The definitions and titles of SMSA's are established by the U.S. Bureau of the Budget with the advice of the Federal Committee on Standard Metropolitan Statistical Areas. There were 212 SMSA's, as defined for the 1960 Decennial Census, for which data may be provided for places of residence in the Health Interview Survey.

The definition of an individual SMSA involves two considerations: first, a city or cities of specified population which constitute the central city and identify the county in which it is located as the central county; and, second, economic and social relationships with contiguous counties (except in New England) which are metropolitan in character, so that the periphery of the specific metropolitan area may be determined. SMSA's are not limited by State boundaries.

Farm and nonfarm residence.—The population residing outside SMSA's is subdivided into the farm population, which comprises all non-SMSA residents living on farms, and the nonfarm population, which comprises the remaining non-SMSA population. The farm population includes persons living on places of 10 acres or more from which sales of farm products amounted to \$50 or more during the previous 12 months or on places of less than 10 acres from which sales of farm products amounted to \$250 or more during the preceding 12 months. Other persons living in non-SMSA territory were classified as nonfarm if their household paid rent for the house but their rent did not include any land used for farming.

Sales of farm products refer to the gross receipts from the sale of field crops, vegetables, fruits, nuts, livestock and livestock products (milk, wool, etc.), poultry and poultry products, and nursery and forest products produced on the place and sold at any time during the preceding 12 months.

Region.—For the purpose of classifying the population by geographic area, the States are grouped into four regions. These regions, which correspond to those used by the Bureau of the Census, are as follows:

<i>Region</i>	<i>States Included</i>
Northeast-----	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania
North Central--	Michigan, Ohio, Indiana, Illinois, Wisconsin, Minnesota, Iowa,

	Missouri, North Dakota, South Dakota, Nebraska, Kansas
South-----	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, Texas
West-----	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Alaska, Washington, Oregon, California, Hawaii



CONDITION (Cont'd.)		REFER RESPONDENT TO TWO-WEEK CALENDAR FOR QUESTIONS 9 - 14			
Ask question 9a for all conditions.	9a. Last week or the week before did his ... cause him to cut down on the things he usually does?	Yes <input type="radio"/>	No <input type="radio"/>	Go to 14a <input type="radio"/>	V <input type="radio"/>
	b. Did he have to cut down for as much as a day?	Yes <input type="radio"/>	No <input type="radio"/>	Go to 14a <input type="radio"/>	V <input type="radio"/>
Ask questions 10 and 11 if "Yes" marked in question 9b.	10. How many days did he have to cut down during that two week period?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	V <input type="radio"/>
	11. During that two week period, how many days did his ... keep him in bed all or most of the day?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	V <input type="radio"/>
Ask question 12 if person is 6 - 16 years old.	12. How many days did his ... keep him from school during that two week period?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	Und. 6 <input type="radio"/>
Ask question 13 if person is 17 years old or over.	13. How many days did his ... keep him from work during that two week period? (For females add) not counting work around the house?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	None <input type="radio"/>
Ask question 14 for all conditions.	14a. When did he first notice his ... during the past 3 months or before that time?			During 3 mos. <input type="radio"/> Before 3 mos. <input type="radio"/> Go to 15 <input type="radio"/>	V <input type="radio"/>
	b. Did he first notice it during the past two weeks or before that time?			Past 2 wks. <input type="radio"/> Before 2 wks. <input type="radio"/> Go to 16 <input type="radio"/>	V <input type="radio"/>
	c. Which week, last week or the week before?			Last week <input type="radio"/> Week before <input type="radio"/>	V <input type="radio"/>
Ask question 15 only if condition was first noticed "Before 3 months."	15. Did -- first notice it during the past 12 months or before that time?			3-12 mos. <input type="radio"/> Before 12 mos. <input type="radio"/>	V <input type="radio"/>
Ask for persons 6 years old or over for whom an eye condition or vision problem (including cataracts and glaucoma) has been reported.	<input type="radio"/> Not an eye condition <input type="radio"/> Not first eye condition <input type="radio"/> Under 6				
	16a. Can -- see well enough to read ordinary newspaper print with glasses?	Yes - Ask 16b <input type="radio"/>	No - Omit 16 b, c <input type="radio"/>		
	b. Can -- see well enough to recognize a friend walking on the other side of the street?	Yes - Omit 16c <input type="radio"/>	No - Ask 16c <input type="radio"/>		
c. How much trouble would you say that -- has in seeing: a great deal, some, or hardly any at all?		Great deal <input type="radio"/>	Some <input type="radio"/>	Hardly any or none <input type="radio"/>	
AA: IF THIS IS A CONDITION ON CARD A OR B, OR STARTED "BEFORE 3 MONTHS," ASK 17-18. OTHERWISE GO TO ITEM BB.					
Ask question 17b if "1" or more days in question 17a and question 11 is blank or marked "None."	17a. ABOUT how many days during the past 12 months has his ... kept him in bed all or most of the day?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	None Go to 18 <input type="radio"/>
	b. Were any of these -- days during last week or the week before?			Yes <input type="radio"/>	No Go to 18 <input type="radio"/>
	c. How many?	Write in and mark <input type="text"/>	Days..	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	
If "Yes" in question 2, ask: If "No" in question 2, omit.	18. ABOUT how many times during the past 12 months has -- seen or talked to a doctor about his...?	Write in and mark <input type="text"/>	Times	<input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	None <input type="radio"/>
BB: Is this the LAST condition for this person? <input type="radio"/> Yes - Ask 19-22 if person has "1" or more conditions past AA <input type="radio"/> No - Go to next condition					
Show Card E, F, G, or H as appropriate based on activity status or age.	19. Please look at each statement on this card (Card E, F, G, H). Then tell me which statement fits -- best in terms of health.			1 2 3 4 - Go to 21 <input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>	V <input type="radio"/>
If 1, 2, or 3 marked in 19, ask: If 4 marked in 19, go to 21.	20. Is this because of any of the conditions you have told me about?	<input type="radio"/> Yes -> Which? _____ (Enter condition numbers) _____ <input type="radio"/> No -> What does cause this limitation? _____ Enter cause _____		WASHINGTON USE Yes <input type="radio"/> No <input type="radio"/>	
	21. Please look at the tan card, Card I. Which one of those statements fits -- best in terms of health?			1 2 3 4 5 6 Step <input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>	V <input type="radio"/>
If 1, 2, 3, 4, or 5 marked in 21, ask: If 6 marked, omit 22 and go to next person.	22. Is this because of any of the conditions you have told me about?	<input type="radio"/> Yes -> Which? _____ (Enter condition numbers) _____ <input type="radio"/> No -> What does cause this limitation? _____ Enter cause _____		WASHINGTON USE Yes <input type="radio"/> No <input type="radio"/>	
					<input type="radio"/> <input type="radio"/>

OUTLINE OF REPORT SERIES FOR VITAL AND HEALTH STATISTICS

Public Health Service Publication No. 1000

- Series 1. Programs and collection procedures.*—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data evaluation and methods research.*—Studies of new statistical methodology including: experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical studies.*—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and committee reports.*—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.*—Statistics on illness, accidental injuries, disability, use of hospital, medical, dental, and other services, and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.*—Data from direct examination, testing, and measurement of national samples of the population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics; and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutional Population Surveys.*—Statistics relating to the health characteristics of persons in institutions, and on medical, nursing, and personal care received, based on national samples of establishments providing these services and samples of the residents or patients.
- Series 13. Data from the Hospital Discharge Survey.*—Statistics relating to discharged patients in short-stay hospitals, based on a sample of patient records in a national sample of hospitals.
- Series 20. Data on mortality.*—Various statistics on mortality other than as included in annual or monthly reports—special analyses by cause of death, age, and other demographic variables, also geographic and time series analyses.
- Series 21. Data on natality, marriage, and divorce.*—Various statistics on natality, marriage, and divorce other than as included in annual or monthly reports—special analyses by demographic variables, also geographic and time series analyses, studies of fertility.
- Series 22. Data from the National Natality and Mortality Surveys.*—Statistics on characteristics of births and deaths not available from the vital records, based on sample surveys stemming from these records, including such topics as mortality by socioeconomic class, medical experience in the last year of life, characteristics of pregnancy, etc.

For a list of titles of reports published in these series, write to: Office of Information
National Center for Health Statistics
U.S. Public Health Service
Washington, D.C. 20201