

Table LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999
 [Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	103,740	100.0	331.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,866	24.9	82.5
2	Malignant neoplasms (C00-C97)	20,233	19.5	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,650	8.3	27.6
4	Cerebrovascular diseases (I60-I69)	5,907	5.7	18.8
5	Diabetes mellitus (E10-E14)	5,182	5.0	16.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,972	2.9	9.5
7	Assault (homicide) (X85-Y09,Y87.1)	2,864	2.8	9.1
8	Chronic lower respiratory diseases (J40-J47)	2,859	2.8	9.1
9	Influenza and pneumonia (J10-J18)	2,246	2.2	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,153	2.1	6.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,958	1.9	6.2
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,695	1.6	5.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,584	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,498	1.4	4.8
15	Septicemia (A40-A41)	1,215	1.2	3.9
...	All other causes (Residual)	16,858	16.3	53.8
Hispanic, both sexes, 1-4 years				
...	All causes	882	100.0	32.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	38.7	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	11.5	3.7
3	Malignant neoplasms (C00-C97)	72	8.2	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	52	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.7	1.2
6	Certain conditions originating in the perinatal period (P00-P96)	18	2.0	*
7	Influenza and pneumonia (J10-J18)	17	1.9	*
8	Septicemia (A40-A41)	14	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
11	Meningitis (G00,G03)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	7	0.8	*
13	Anemias (D50-D64)	6	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
...	All other causes (Residual)	183	20.7	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	464	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	36.6	5.2
2	Malignant neoplasms (C00-C97)	87	18.8	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	6.7	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	29	6.3	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
8	Septicemia (A40-A41)	6	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.1	*
9	Influenza and pneumonia (J10-J18)	5	1.1	*
9	Diseases of appendix (K35-K38)	5	1.1	*
12	Cerebrovascular diseases (I60-I69)	4	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
14	Anemias (D50-D64)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	74	15.9	2.3
Hispanic, both sexes, 10-14 years				
...	All causes	550	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	34.9	7.0
2	Malignant neoplasms (C00-C97)	89	16.2	3.2
3	Assault (homicide) (X85-Y09,Y87.1)	60	10.9	2.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	30	5.5	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	4.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
9	Influenza and pneumonia (J10-J18)	5	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
12	Diabetes mellitus (E10-E14)	4	0.7	*
14	Pneumonitis due to solids and liquids (J69)	3	0.5	*
14	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (Residual)	89	16.2	3.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,871	100.0	67.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	800	42.8	28.8
2	Assault (homicide) (X85-Y09,Y87.1)	482	25.8	17.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	175	9.4	6.3
4	Malignant neoplasms (C00-C97)	94	5.0	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	2.8	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.4	1.0
7	Cerebrovascular diseases (I60-I69)	10	0.5	*
7	Influenza and pneumonia (J10-J18)	10	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-099)	6	0.3	*
12	Septicemia (A40-A41)	5	0.3	*
12	Legal intervention (Y35,Y89.0)	5	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.2	*
...	All other causes (Residual)	182	9.7	6.5
Hispanic, both sexes, 20-24 years				
...	All causes	2,634	100.0	97.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,146	43.5	42.6
2	Assault (homicide) (X85-Y09,Y87.1)	637	24.2	23.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	216	8.2	8.0
4	Malignant neoplasms (C00-C97)	165	6.3	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	2.4	2.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	29	1.1	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.1	1.1
8	Cerebrovascular diseases (I60-I69)	22	0.8	0.8
9	Influenza and pneumonia (J10-J18)	15	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-099)	13	0.5	*
11	Diabetes mellitus (E10-E14)	12	0.5	*
12	Legal intervention (Y35,Y89.0)	8	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	261	9.9	9.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999
 [Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,527	100.0	98.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	936	37.0	36.4
2	Assault (homicide) (X85-Y09,Y87.1)	482	19.1	18.8
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	220	8.7	8.6
4	Malignant neoplasms (C00-C97)	189	7.5	7.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	115	4.6	4.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.7	3.6
7	Cerebrovascular diseases (I60-I69)	46	1.8	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.1	1.1
9	Pregnancy, childbirth and the puerperium (O00-099)	18	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.7	*
11	Diabetes mellitus (E10-E14)	14	0.6	*
12	Influenza and pneumonia (J10-J18)	10	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
12	Legal intervention (Y35,Y89.0)	10	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
...	All other causes (Residual)	331	13.1	12.9
Hispanic, both sexes, 30-34 years				
...	All causes	2,870	100.0	107.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	854	29.8	32.1
2	Assault (homicide) (X85-Y09,Y87.1)	319	11.1	12.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	292	10.2	11.0
4	Malignant neoplasms (C00-C97)	277	9.7	10.4
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	203	7.1	7.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	6.7	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	73	2.5	2.7
8	Cerebrovascular diseases (I60-I69)	56	2.0	2.1
9	Diabetes mellitus (E10-E14)	29	1.0	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.9	1.0
11	Septicemia (A40-A41)	19	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
13	Influenza and pneumonia (J10-J18)	14	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
15	Viral hepatitis (B15-B19)	13	0.5	*
15	Legal intervention (Y35,Y89.0)	13	0.5	*
...	All other causes (Residual)	458	16.0	17.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,624	100.0	140.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	844	23.3	32.7
2	Malignant neoplasms (C00-C97)	471	13.0	18.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	422	11.6	16.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	8.8	12.4
5	Assault (homicide) (X85-Y09,Y87.1)	247	6.8	9.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	228	6.3	8.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	190	5.2	7.4
8	Cerebrovascular diseases (I60-I69)	89	2.5	3.4
9	Diabetes mellitus (E10-E14)	56	1.5	2.2
10	Influenza and pneumonia (J10-J18)	46	1.3	1.8
11	Viral hepatitis (B15-B19)	38	1.0	1.5
12	Chronic lower respiratory diseases (J40-J47)	34	0.9	1.3
13	Septicemia (A40-A41)	30	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-099)	21	0.6	0.8
...	All other causes (Residual)	565	15.6	21.9
Hispanic, both sexes, 40-44 years				
...	All causes	4,174	100.0	192.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	18.3	35.2
2	Malignant neoplasms (C00-C97)	733	17.6	33.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	441	10.6	20.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	433	10.4	20.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	328	7.9	15.2
6	Assault (homicide) (X85-Y09,Y87.1)	185	4.4	8.5
7	Cerebrovascular diseases (I60-I69)	152	3.6	7.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	141	3.4	6.5
9	Diabetes mellitus (E10-E14)	101	2.4	4.7
10	Viral hepatitis (B15-B19)	66	1.6	3.0
11	Chronic lower respiratory diseases (J40-J47)	45	1.1	2.1
12	Septicemia (A40-A41)	44	1.1	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.9	1.8
14	Influenza and pneumonia (J10-J18)	28	0.7	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	1.1
...	All other causes (Residual)	651	15.6	30.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	4,767	100.0	287.6
1	Malignant neoplasms (C00-C97)	1,093	22.9	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	716	15.0	43.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	546	11.5	32.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	423	8.9	25.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	287	6.0	17.3
6	Cerebrovascular diseases (I60-I69)	229	4.8	13.8
7	Diabetes mellitus (E10-E14)	169	3.5	10.2
8	Assault (homicide) (X85-Y09,Y87.1)	133	2.8	8.0
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	121	2.5	7.3
10	Viral hepatitis (B15-B19)	99	2.1	6.0
11	Chronic lower respiratory diseases (J40-J47)	52	1.1	3.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.0	3.0
13	Influenza and pneumonia (J10-J18)	48	1.0	2.9
14	Septicemia (A40-A41)	35	0.7	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.6	1.7
...	All other causes (Residual)	737	15.5	44.5
Hispanic, both sexes, 50-54 years				
...	All causes	5,109	100.0	406.4
1	Malignant neoplasms (C00-C97)	1,423	27.9	113.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,090	21.3	86.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	7.4	30.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	364	7.1	29.0
5	Diabetes mellitus (E10-E14)	244	4.8	19.4
6	Cerebrovascular diseases (I60-I69)	235	4.6	18.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	168	3.3	13.4
8	Viral hepatitis (B15-B19)	104	2.0	8.3
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	89	1.7	7.1
10	Assault (homicide) (X85-Y09,Y87.1)	64	1.3	5.1
11	Chronic lower respiratory diseases (J40-J47)	63	1.2	5.0
12	Septicemia (A40-A41)	57	1.1	4.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.1	4.5
14	Influenza and pneumonia (J10-J18)	56	1.1	4.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	0.7	2.8
...	All other causes (Residual)	680	13.3	54.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	5,495	100.0	589.4
1	Malignant neoplasms (C00-C97)	1,635	29.8	175.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,331	24.2	142.8
3	Diabetes mellitus (E10-E14)	368	6.7	39.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	325	5.9	34.9
5	Cerebrovascular diseases (I60-I69)	286	5.2	30.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	4.5	26.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.7	10.3
8	Chronic lower respiratory diseases (J40-J47)	95	1.7	10.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	89	1.6	9.5
10	Viral hepatitis (B15-B19)	69	1.3	7.4
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	1.2	7.0
12	Septicemia (A40-A41)	63	1.1	6.8
13	Influenza and pneumonia (J10-J18)	58	1.1	6.2
14	Assault (homicide) (X85-Y09,Y87.1)	47	0.9	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	3.3
...	All other causes (Residual)	687	12.5	73.7
Hispanic, both sexes, 60-64 years				
...	All causes	6,824	100.0	913.8
1	Malignant neoplasms (C00-C97)	2,051	30.1	274.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,824	26.7	244.3
3	Diabetes mellitus (E10-E14)	559	8.2	74.9
4	Cerebrovascular diseases (I60-I69)	331	4.9	44.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	313	4.6	41.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	3.2	28.9
7	Chronic lower respiratory diseases (J40-J47)	144	2.1	19.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	2.0	17.9
9	Influenza and pneumonia (J10-J18)	97	1.4	13.0
10	Septicemia (A40-A41)	68	1.0	9.1
11	Viral hepatitis (B15-B19)	58	0.8	7.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	53	0.8	7.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	48	0.7	6.4
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	47	0.7	6.3
15	Aortic aneurysm and dissection (I71)	31	0.5	4.2
...	All other causes (Residual)	850	12.5	113.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	8,843	100.0	1,444.8
1	Malignant neoplasms (C00-C97)	2,598	29.4	424.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,464	27.9	402.6
3	Diabetes mellitus (E10-E14)	721	8.2	117.8
4	Cerebrovascular diseases (I60-I69)	524	5.9	85.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	310	3.5	50.6
6	Chronic lower respiratory diseases (J40-J47)	262	3.0	42.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	2.8	40.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	1.9	28.1
9	Influenza and pneumonia (J10-J18)	143	1.6	23.4
10	Septicemia (A40-A41)	119	1.3	19.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	0.8	11.6
12	Viral hepatitis (B15-B19)	68	0.8	11.1
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	46	0.5	7.5
14	Aortic aneurysm and dissection (I71)	42	0.5	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	6.4
...	All other causes (Residual)	1,019	11.5	166.5
Hispanic, both sexes, 70-74 years				
...	All causes	10,358	100.0	2,115.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,093	29.9	631.6
2	Malignant neoplasms (C00-C97)	2,796	27.0	571.0
3	Diabetes mellitus (E10-E14)	830	8.0	169.5
4	Cerebrovascular diseases (I60-I69)	687	6.6	140.3
5	Chronic lower respiratory diseases (J40-J47)	387	3.7	79.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	273	2.6	55.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	1.9	40.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	1.9	39.8
9	Influenza and pneumonia (J10-J18)	185	1.8	37.8
10	Septicemia (A40-A41)	146	1.4	29.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	87	0.8	17.8
12	Alzheimer's disease (G30)	75	0.7	15.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.7	14.3
14	Viral hepatitis (B15-B19)	61	0.6	12.5
15	Parkinson's disease (G20-G21)	60	0.6	12.3
...	All other causes (Residual)	1,216	11.7	248.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	11,081	100.0	3,154.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,618	32.7	1,030.1
2	Malignant neoplasms (C00-C97)	2,586	23.3	736.2
3	Cerebrovascular diseases (I60-I69)	823	7.4	234.3
4	Diabetes mellitus (E10-E14)	782	7.1	222.6
5	Chronic lower respiratory diseases (J40-J47)	483	4.4	137.5
6	Influenza and pneumonia (J10-J18)	252	2.3	71.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	2.2	68.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	187	1.7	53.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.7	52.1
10	Alzheimer's disease (G30)	166	1.5	47.3
11	Septicemia (A40-A41)	162	1.5	46.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	110	1.0	31.3
13	Parkinson's disease (G20-G21)	97	0.9	27.6
14	Atherosclerosis (I70)	60	0.5	17.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.5	16.2
15	Aortic aneurysm and dissection (I71)	57	0.5	16.2
15	Pneumonitis due to solids and liquids (J69)	57	0.5	16.2
...	All other causes (Residual)	1,162	10.5	330.8
Hispanic, both sexes, 80-84 years				
...	All causes	9,831	100.0	4,838.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,428	34.9	1,687.1
2	Malignant neoplasms (C00-C97)	1,863	19.0	916.9
3	Cerebrovascular diseases (I60-I69)	849	8.6	417.8
4	Diabetes mellitus (E10-E14)	587	6.0	288.9
5	Chronic lower respiratory diseases (J40-J47)	452	4.6	222.5
6	Influenza and pneumonia (J10-J18)	317	3.2	156.0
7	Alzheimer's disease (G30)	196	2.0	96.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	1.9	92.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	1.5	74.3
10	Septicemia (A40-A41)	134	1.4	65.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	122	1.2	60.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	98	1.0	48.2
13	Parkinson's disease (G20-G21)	87	0.9	42.8
14	Atherosclerosis (I70)	70	0.7	34.5
15	Pneumonitis due to solids and liquids (J69)	61	0.6	30.0
...	All other causes (Residual)	1,229	12.5	604.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	8,953	100.0	7,637.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,363	37.6	2,868.7
2	Malignant neoplasms (C00-C97)	1,231	13.7	1,050.1
3	Cerebrovascular diseases (I60-I69)	828	9.2	706.3
4	Diabetes mellitus (E10-E14)	459	5.1	391.5
5	Chronic lower respiratory diseases (J40-J47)	415	4.6	354.0
6	Influenza and pneumonia (J10-J18)	383	4.3	326.7
7	Alzheimer's disease (G30)	244	2.7	208.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	1.9	142.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	1.6	119.4
10	Septicemia (A40-A41)	131	1.5	111.7
11	Pneumonitis due to solids and liquids (J69)	85	0.9	72.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	80	0.9	68.2
13	Atherosclerosis (I70)	79	0.9	67.4
14	Parkinson's disease (G20-G21)	63	0.7	53.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	51	0.6	43.5
...	All other causes (Residual)	1,234	13.8	1,052.6
Hispanic, both sexes, 90-94 years				
...	All causes	5,662	100.0	10,756.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,349	41.5	4,462.6
2	Malignant neoplasms (C00-C97)	588	10.4	1,117.1
3	Cerebrovascular diseases (I60-I69)	505	8.9	959.4
4	Influenza and pneumonia (J10-J18)	296	5.2	562.3
5	Chronic lower respiratory diseases (J40-J47)	249	4.4	473.1
6	Diabetes mellitus (E10-E14)	179	3.2	340.1
7	Alzheimer's disease (G30)	174	3.1	330.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.0	210.9
9	Septicemia (A40-A41)	83	1.5	157.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	1.5	157.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	80	1.4	152.0
12	Pneumonitis due to solids and liquids (J69)	67	1.2	127.3
13	Atherosclerosis (I70)	61	1.1	115.9
14	Parkinson's disease (G20-G21)	36	0.6	68.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	49.4
...	All other causes (Residual)	775	13.7	1,472.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	2,285	100.0	13,791.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,050	46.0	6,337.5
2	Cerebrovascular diseases (I60-I69)	170	7.4	1,026.1
3	Malignant neoplasms (C00-C97)	150	6.6	905.4
3	Influenza and pneumonia (J10-J18)	150	6.6	905.4
5	Chronic lower respiratory diseases (J40-J47)	91	4.0	549.3
6	Alzheimer's disease (G30)	74	3.2	446.6
7	Diabetes mellitus (E10-E14)	60	2.6	362.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.1	289.7
9	Septicemia (A40-A41)	37	1.6	223.3
10	Atherosclerosis (I70)	35	1.5	211.3
11	Pneumonitis due to solids and liquids (J69)	28	1.2	169.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.2	169.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	0.8	*
14	Nutritional deficiencies (E40-E64)	15	0.7	*
15	Anemias (D50-D64)	11	0.5	*
...	All other causes (Residual)	319	14.0	1,925.4
Hispanic, both sexes, 100 years and over				
...	All causes	458	100.0	16,721.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	45.0	7,521.0
2	Influenza and pneumonia (J10-J18)	47	10.3	1,716.0
3	Cerebrovascular diseases (I60-I69)	33	7.2	1,204.8
4	Malignant neoplasms (C00-C97)	26	5.7	949.3
5	Alzheimer's disease (G30)	9	2.0	*
5	Atherosclerosis (I70)	9	2.0	*
5	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	2.0	*
9	Septicemia (A40-A41)	6	1.3	*
9	Diabetes mellitus (E10-E14)	6	1.3	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.3	*
9	Pneumonitis due to solids and liquids (J69)	6	1.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
14	Nutritional deficiencies (E40-E64)	4	0.9	*
15	Parkinson's disease (G20-G21)	2	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	*
...	All other causes (Residual)	73	15.9	2,665.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	57,991	100.0	367.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,554	23.4	86.0
2	Malignant neoplasms (C00-C97)	10,668	18.4	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,578	11.3	41.7
4	Cerebrovascular diseases (I60-I69)	2,809	4.8	17.8
5	Assault (homicide) (X85-Y09,Y87.1)	2,399	4.1	15.2
6	Diabetes mellitus (E10-E14)	2,335	4.0	14.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,143	3.7	13.6
8	Chronic lower respiratory diseases (J40-J47)	1,547	2.7	9.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,525	2.6	9.7
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,430	2.5	9.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,192	2.1	7.6
12	Influenza and pneumonia (J10-J18)	1,092	1.9	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	814	1.4	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	756	1.3	4.8
15	Septicemia (A40-A41)	605	1.0	3.8
...	All other causes (Residual)	8,544	14.7	54.2
Hispanic, male, 1-4 years				
...	All causes	481	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	41.0	14.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.1	4.5
3	Malignant neoplasms (C00-C97)	34	7.1	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	25	5.2	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.2	1.4
6	Influenza and pneumonia (J10-J18)	10	2.1	*
7	Septicemia (A40-A41)	8	1.7	*
8	Meningitis (G00,G03)	5	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
10	Anemias (D50-D64)	4	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	92	19.1	6.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 5-9 years			
...	All causes	266	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	40.2	6.5
2	Malignant neoplasms (C00-C97)	45	16.9	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.0	*
3	Assault (homicide) (X85-Y09,Y87.1)	16	6.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.9	*
7	Influenza and pneumonia (J10-J18)	4	1.5	*
8	Septicemia (A40-A41)	3	1.1	*
8	Cerebrovascular diseases (I60-I69)	3	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
12	Diseases of appendix (K35-K38)	2	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
15	Shigellosis and amebiasis (A03,A06)	1	0.4	*
15	Meningococcal infection (A39)	1	0.4	*
15	Anemias (D50-D64)	1	0.4	*
15	Meningitis (G00,G03)	1	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	39	14.7	2.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 10-14 years			
...	All causes	326	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	36.8	8.6
2	Malignant neoplasms (C00-C97)	47	14.4	3.4
3	Assault (homicide) (X85-Y09,Y87.1)	43	13.2	3.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	21	6.4	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.7	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
9	Cerebrovascular diseases (I60-I69)	3	0.9	*
11	Diabetes mellitus (E10-E14)	2	0.6	*
11	Influenza and pneumonia (J10-J18)	2	0.6	*
11	Diseases of appendix (K35-K38)	2	0.6	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Septicemia (A40-A41)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	39	12.0	2.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 15-19 years			
...	All causes	1,432	100.0	99.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	608	42.5	42.1
2	Assault (homicide) (X85-Y09,Y87.1)	425	29.7	29.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	146	10.2	10.1
4	Malignant neoplasms (C00-C97)	55	3.8	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.2	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.2	*
7	Cerebrovascular diseases (I60-I69)	7	0.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
8	Influenza and pneumonia (J10-J18)	5	0.3	*
8	Legal intervention (Y35,Y89.0)	5	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.3	*
12	Septicemia (A40-A41)	3	0.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.1	*
...	All other causes (Residual)	113	7.9	7.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 20-24 years			
...	All causes	2,115	100.0	151.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	967	45.7	69.3
2	Assault (homicide) (X85-Y09,Y87.1)	565	26.7	40.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	192	9.1	13.8
4	Malignant neoplasms (C00-C97)	103	4.9	7.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.0	3.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.8	*
7	Cerebrovascular diseases (I60-I69)	14	0.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.6	*
9	Diabetes mellitus (E10-E14)	8	0.4	*
9	Legal intervention (Y35,Y89.0)	8	0.4	*
11	Influenza and pneumonia (J10-J18)	6	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.1	*
14	Tuberculosis (A16-A19)	2	0.1	*
14	Septicemia (A40-A41)	2	0.1	*
14	Anemias (D50-D64)	2	0.1	*
14	Aortic aneurysm and dissection (I71)	2	0.1	*
14	Diseases of appendix (K35-K38)	2	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.1	*
...	All other causes (Residual)	158	7.5	11.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	1,956	100.0	148.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	787	40.2	59.7
2	Assault (homicide) (X85-Y09,Y87.1)	428	21.9	32.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	187	9.6	14.2
4	Malignant neoplasms (C00-C97)	98	5.0	7.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	87	4.4	6.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.9	5.8
7	Cerebrovascular diseases (I60-I69)	25	1.3	1.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.8	*
9	Legal intervention (Y35,Y89.0)	10	0.5	*
10	Diabetes mellitus (E10-E14)	8	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
12	Influenza and pneumonia (J10-J18)	6	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	205	10.5	15.5
Hispanic, male, 30-34 years				
...	All causes	2,128	100.0	154.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	707	33.2	51.4
2	Assault (homicide) (X85-Y09,Y87.1)	270	12.7	19.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	227	10.7	16.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	177	8.3	12.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	6.7	10.3
6	Malignant neoplasms (C00-C97)	124	5.8	9.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	56	2.6	4.1
8	Cerebrovascular diseases (I60-I69)	37	1.7	2.7
9	Diabetes mellitus (E10-E14)	13	0.6	*
9	Legal intervention (Y35,Y89.0)	13	0.6	*
11	Septicemia (A40-A41)	11	0.5	*
11	Viral hepatitis (B15-B19)	11	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.5	*
14	Influenza and pneumonia (J10-J18)	9	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
...	All other causes (Residual)	307	14.4	22.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,601	100.0	195.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	692	26.6	52.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	338	13.0	25.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	9.0	17.7
4	Assault (homicide) (X85-Y09,Y87.1)	201	7.7	15.1
5	Malignant neoplasms (C00-C97)	198	7.6	14.9
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	188	7.2	14.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	154	5.9	11.6
8	Cerebrovascular diseases (I60-I69)	49	1.9	3.7
9	Influenza and pneumonia (J10-J18)	34	1.3	2.6
10	Diabetes mellitus (E10-E14)	31	1.2	2.3
11	Viral hepatitis (B15-B19)	30	1.2	2.3
12	Septicemia (A40-A41)	20	0.8	1.5
13	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
15	Legal intervention (Y35,Y89.0)	10	0.4	*
...	All other causes (Residual)	395	15.2	29.7
Hispanic, male, 40-44 years				
...	All causes	2,875	100.0	262.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	615	21.4	56.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	11.7	30.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	328	11.4	30.0
4	Malignant neoplasms (C00-C97)	306	10.6	28.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	267	9.3	24.4
6	Assault (homicide) (X85-Y09,Y87.1)	153	5.3	14.0
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	119	4.1	10.9
8	Cerebrovascular diseases (I60-I69)	97	3.4	8.9
9	Viral hepatitis (B15-B19)	54	1.9	4.9
10	Diabetes mellitus (E10-E14)	53	1.8	4.8
11	Septicemia (A40-A41)	31	1.1	2.8
12	Chronic lower respiratory diseases (J40-J47)	22	0.8	2.0
13	Influenza and pneumonia (J10-J18)	20	0.7	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
...	All other causes (Residual)	437	15.2	39.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	3,213	100.0	392.4
1	Malignant neoplasms (C00-C97)	539	16.8	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	531	16.5	64.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	443	13.8	54.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	354	11.0	43.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	224	7.0	27.4
6	Cerebrovascular diseases (I60-I69)	136	4.2	16.6
7	Diabetes mellitus (E10-E14)	105	3.3	12.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	104	3.2	12.7
9	Assault (homicide) (X85-Y09,Y87.1)	100	3.1	12.2
10	Viral hepatitis (B15-B19)	76	2.4	9.3
11	Influenza and pneumonia (J10-J18)	32	1.0	3.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.9	3.7
13	Septicemia (A40-A41)	22	0.7	2.7
14	Chronic lower respiratory diseases (J40-J47)	21	0.7	2.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.5	*
...	All other causes (Residual)	480	14.9	58.6
Hispanic, male, 50-54 years				
...	All causes	3,284	100.0	542.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	817	24.9	135.0
2	Malignant neoplasms (C00-C97)	698	21.3	115.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	293	8.9	48.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	291	8.9	48.1
5	Cerebrovascular diseases (I60-I69)	149	4.5	24.6
6	Diabetes mellitus (E10-E14)	147	4.5	24.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	130	4.0	21.5
8	Viral hepatitis (B15-B19)	77	2.3	12.7
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	2.0	10.7
10	Assault (homicide) (X85-Y09,Y87.1)	50	1.5	8.3
11	Influenza and pneumonia (J10-J18)	39	1.2	6.4
12	Septicemia (A40-A41)	34	1.0	5.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	5.6
14	Chronic lower respiratory diseases (J40-J47)	31	0.9	5.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	0.6	*
...	All other causes (Residual)	410	12.5	67.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	3,409	100.0	780.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	924	27.1	211.6
2	Malignant neoplasms (C00-C97)	901	26.4	206.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	257	7.5	58.9
4	Diabetes mellitus (E10-E14)	202	5.9	46.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	5.4	41.9
6	Cerebrovascular diseases (I60-I69)	157	4.6	36.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	76	2.2	17.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.7	13.3
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	55	1.6	12.6
10	Chronic lower respiratory diseases (J40-J47)	50	1.5	11.5
11	Viral hepatitis (B15-B19)	43	1.3	9.8
12	Septicemia (A40-A41)	37	1.1	8.5
12	Assault (homicide) (X85-Y09,Y87.1)	37	1.1	8.5
14	Influenza and pneumonia (J10-J18)	32	0.9	7.3
15	Aortic aneurysm and dissection (I71)	17	0.5	*
...	All other causes (Residual)	380	11.1	87.0
Hispanic, male, 60-64 years				
...	All causes	4,081	100.0	1,195.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,167	28.6	341.7
2	Malignant neoplasms (C00-C97)	1,159	28.4	339.4
3	Diabetes mellitus (E10-E14)	278	6.8	81.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	224	5.5	65.6
5	Cerebrovascular diseases (I60-I69)	197	4.8	57.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	3.7	44.2
7	Chronic lower respiratory diseases (J40-J47)	85	2.1	24.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.9	23.1
9	Influenza and pneumonia (J10-J18)	48	1.2	14.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	42	1.0	12.3
11	Septicemia (A40-A41)	41	1.0	12.0
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	1.0	11.4
13	Viral hepatitis (B15-B19)	32	0.8	9.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.7	8.5
15	Aortic aneurysm and dissection (I71)	24	0.6	7.0
...	All other causes (Residual)	486	11.9	142.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,026	100.0	1,841.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,522	30.3	557.5
2	Malignant neoplasms (C00-C97)	1,469	29.2	538.1
3	Diabetes mellitus (E10-E14)	331	6.6	121.2
4	Cerebrovascular diseases (I60-I69)	273	5.4	100.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	198	3.9	72.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	3.2	59.3
7	Chronic lower respiratory diseases (J40-J47)	136	2.7	49.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	1.7	30.8
9	Influenza and pneumonia (J10-J18)	82	1.6	30.0
10	Septicemia (A40-A41)	67	1.3	24.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	42	0.8	15.4
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	41	0.8	15.0
13	Viral hepatitis (B15-B19)	32	0.6	11.7
14	Aortic aneurysm and dissection (I71)	30	0.6	11.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.6	10.3
...	All other causes (Residual)	529	10.5	193.8
Hispanic, male, 70-74 years				
...	All causes	5,757	100.0	2,704.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,833	31.8	861.2
2	Malignant neoplasms (C00-C97)	1,576	27.4	740.4
3	Diabetes mellitus (E10-E14)	378	6.6	177.6
4	Cerebrovascular diseases (I60-I69)	368	6.4	172.9
5	Chronic lower respiratory diseases (J40-J47)	237	4.1	111.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.8	76.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	2.2	59.2
8	Influenza and pneumonia (J10-J18)	105	1.8	49.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.6	44.2
10	Septicemia (A40-A41)	67	1.2	31.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	42	0.7	19.7
12	Aortic aneurysm and dissection (I71)	37	0.6	17.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	16.0
13	Alzheimer's disease (G30)	34	0.6	16.0
15	Parkinson's disease (G20-G21)	33	0.6	15.5
...	All other causes (Residual)	631	11.0	296.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	5,860	100.0	3,913.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,957	33.4	1,306.8
2	Malignant neoplasms (C00-C97)	1,436	24.5	958.9
3	Cerebrovascular diseases (I60-I69)	423	7.2	282.5
4	Diabetes mellitus (E10-E14)	336	5.7	224.4
5	Chronic lower respiratory diseases (J40-J47)	278	4.7	185.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	2.3	88.8
7	Influenza and pneumonia (J10-J18)	129	2.2	86.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	1.7	66.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.6	62.1
10	Septicemia (A40-A41)	90	1.5	60.1
11	Alzheimer's disease (G30)	67	1.1	44.7
12	Parkinson's disease (G20-G21)	60	1.0	40.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	56	1.0	37.4
14	Pneumonitis due to solids and liquids (J69)	34	0.6	22.7
15	Aortic aneurysm and dissection (I71)	31	0.5	20.7
...	All other causes (Residual)	637	10.9	425.4
Hispanic, male, 80-84 years				
...	All causes	4,457	100.0	5,696.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,526	34.2	1,950.4
2	Malignant neoplasms (C00-C97)	967	21.7	1,236.0
3	Cerebrovascular diseases (I60-I69)	335	7.5	428.2
4	Chronic lower respiratory diseases (J40-J47)	245	5.5	313.1
5	Diabetes mellitus (E10-E14)	226	5.1	288.9
6	Influenza and pneumonia (J10-J18)	155	3.5	198.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.8	104.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.8	101.0
9	Alzheimer's disease (G30)	61	1.4	78.0
10	Septicemia (A40-A41)	54	1.2	69.0
11	Parkinson's disease (G20-G21)	53	1.2	67.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	43	1.0	55.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.0	55.0
14	Aortic aneurysm and dissection (I71)	34	0.8	43.5
15	Pneumonitis due to solids and liquids (J69)	31	0.7	39.6
...	All other causes (Residual)	523	11.7	668.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	3,483	100.0	8,542.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,250	35.9	3,065.6
2	Malignant neoplasms (C00-C97)	564	16.2	1,383.2
3	Cerebrovascular diseases (I60-I69)	284	8.2	696.5
4	Chronic lower respiratory diseases (J40-J47)	216	6.2	529.7
5	Diabetes mellitus (E10-E14)	160	4.6	392.4
6	Influenza and pneumonia (J10-J18)	155	4.5	380.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.5	215.8
8	Alzheimer's disease (G30)	72	2.1	176.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	1.9	164.3
10	Septicemia (A40-A41)	46	1.3	112.8
11	Pneumonitis due to solids and liquids (J69)	39	1.1	95.6
12	Atherosclerosis (I70)	37	1.1	90.7
13	Parkinson's disease (G20-G21)	31	0.9	76.0
14	Aortic aneurysm and dissection (I71)	29	0.8	71.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.7	63.8
...	All other causes (Residual)	419	12.0	1,027.6
Hispanic, male, 90-94 years				
...	All causes	1,960	100.0	11,534.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	741	37.8	4,360.9
2	Malignant neoplasms (C00-C97)	271	13.8	1,594.9
3	Cerebrovascular diseases (I60-I69)	183	9.3	1,077.0
4	Chronic lower respiratory diseases (J40-J47)	119	6.1	700.3
5	Influenza and pneumonia (J10-J18)	118	6.0	694.4
6	Diabetes mellitus (E10-E14)	46	2.3	270.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.3	270.7
8	Alzheimer's disease (G30)	38	1.9	223.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.9	217.7
10	Septicemia (A40-A41)	28	1.4	164.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	1.4	164.8
12	Pneumonitis due to solids and liquids (J69)	22	1.1	129.5
13	Atherosclerosis (I70)	18	0.9	*
14	Parkinson's disease (G20-G21)	14	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.5	*
...	All other causes (Residual)	241	12.3	1,418.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	703	100.0	13,992.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	286	40.7	5,692.7
2	Malignant neoplasms (C00-C97)	59	8.4	1,174.4
3	Cerebrovascular diseases (I60-I69)	50	7.1	995.2
4	Influenza and pneumonia (J10-J18)	49	7.0	975.3
5	Chronic lower respiratory diseases (J40-J47)	48	6.8	955.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	3.0	418.0
7	Alzheimer's disease (G30)	18	2.6	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.1	*
9	Pneumonitis due to solids and liquids (J69)	13	1.8	*
10	Diabetes mellitus (E10-E14)	11	1.6	*
10	Atherosclerosis (I70)	11	1.6	*
12	Septicemia (A40-A41)	10	1.4	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.0	*
14	Nutritional deficiencies (E40-E64)	6	0.9	*
15	Anemias (D50-D64)	5	0.7	*
...	All other causes (Residual)	94	13.4	1,871.0
Hispanic, male, 100 years and over				
...	All causes	107	100.0	14,439.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	43.9	6,342.8
2	Influenza and pneumonia (J10-J18)	12	11.2	*
3	Cerebrovascular diseases (I60-I69)	10	9.3	*
4	Malignant neoplasms (C00-C97)	8	7.5	*
5	Chronic lower respiratory diseases (J40-J47)	5	4.7	*
6	Atherosclerosis (I70)	4	3.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
7	Nutritional deficiencies (E40-E64)	1	0.9	*
7	Alzheimer's disease (G30)	1	0.9	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.9	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
7	Hyperplasia of prostate (N40)	1	0.9	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	0.9	*
...	All other causes (Residual)	14	13.1	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	45,749	100.0	293.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,312	26.9	79.0
2	Malignant neoplasms (C00-C97)	9,565	20.9	61.4
3	Cerebrovascular diseases (I60-I69)	3,098	6.8	19.9
4	Diabetes mellitus (E10-E14)	2,847	6.2	18.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,072	4.5	13.3
6	Chronic lower respiratory diseases (J40-J47)	1,312	2.9	8.4
7	Influenza and pneumonia (J10-J18)	1,154	2.5	7.4
8	Certain conditions originating in the perinatal period (P00-P96)	961	2.1	6.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	829	1.8	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	770	1.7	4.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	742	1.6	4.8
12	Alzheimer's disease (G30)	667	1.5	4.3
13	Septicemia (A40-A41)	610	1.3	3.9
14	Assault (homicide) (X85-Y09,Y87.1)	465	1.0	3.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	433	0.9	2.8
...	All other causes (Residual)	7,912	17.3	50.8
Hispanic, female, 1-4 years				
...	All causes	401	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	35.9	10.7
2	Malignant neoplasms (C00-C97)	38	9.5	2.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	9.5	2.8
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.2	*
5	Certain conditions originating in the perinatal period (P00-P96)	13	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.0	*
8	Influenza and pneumonia (J10-J18)	7	1.7	*
9	Septicemia (A40-A41)	6	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Meningitis (G00,G03)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	83	20.7	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 5-9 years			
...	All causes	198	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	31.8	4.0
2	Malignant neoplasms (C00-C97)	42	21.2	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	7.6	*
4	Assault (homicide) (X85-Y09,Y87.1)	13	6.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	3.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.0	*
8	Septicemia (A40-A41)	3	1.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.5	*
8	Diseases of appendix (K35-K38)	3	1.5	*
11	Pneumonitis due to solids and liquids (J69)	2	1.0	*
12	Anemias (D50-D64)	1	0.5	*
12	Cerebrovascular diseases (I60-I69)	1	0.5	*
12	Influenza and pneumonia (J10-J18)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	16.2	2.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 10-14 years			
...	All causes	224	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	32.1	5.4
2	Malignant neoplasms (C00-C97)	42	18.8	3.1
3	Assault (homicide) (X85-Y09,Y87.1)	17	7.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.8	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	9	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.7	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
7	Influenza and pneumonia (J10-J18)	3	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
11	Diabetes mellitus (E10-E14)	2	0.9	*
11	Cerebrovascular diseases (I60-I69)	2	0.9	*
11	Pneumonitis due to solids and liquids (J69)	2	0.9	*
14	Meningococcal infection (A39)	1	0.4	*
14	Septicemia (A40-A41)	1	0.4	*
14	Anemias (D50-D64)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Diseases of appendix (K35-K38)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	41	18.3	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 15-19 years			
...	All causes	439	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	43.7	14.4
2	Assault (homicide) (X85-Y09,Y87.1)	57	13.0	4.3
3	Malignant neoplasms (C00-C97)	39	8.9	2.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	6.6	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.8	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	2.3	*
7	Pregnancy, childbirth and the puerperium (O00-099)	6	1.4	*
8	Influenza and pneumonia (J10-J18)	5	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
10	Meningitis (G00,G03)	3	0.7	*
10	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Septicemia (A40-A41)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Diabetes mellitus (E10-E14)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	56	12.8	4.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 20-24 years			
...	All causes	519	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	34.5	13.8
2	Assault (homicide) (X85-Y09,Y87.1)	72	13.9	5.6
3	Malignant neoplasms (C00-C97)	62	11.9	4.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	4.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	3.1	*
7	Pregnancy, childbirth and the puerperium (O00-099)	13	2.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.3	*
9	Influenza and pneumonia (J10-J18)	9	1.7	*
10	Cerebrovascular diseases (I60-I69)	8	1.5	*
11	Diabetes mellitus (E10-E14)	4	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Septicemia (A40-A41)	2	0.4	*
13	Viral hepatitis (B15-B19)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Anemias (D50-D64)	2	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
...	All other causes (Residual)	83	16.0	6.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	571	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	26.1	11.9
2	Malignant neoplasms (C00-C97)	91	15.9	7.3
3	Assault (homicide) (X85-Y09,Y87.1)	54	9.5	4.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	5.8	2.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	28	4.9	2.2
6	Cerebrovascular diseases (I60-I69)	21	3.7	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	18	3.2	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.3	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
11	Diabetes mellitus (E10-E14)	6	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.1	*
13	Septicemia (A40-A41)	4	0.7	*
13	Influenza and pneumonia (J10-J18)	4	0.7	*
15	Viral hepatitis (B15-B19)	3	0.5	*
...	All other causes (Residual)	114	20.0	9.1
Hispanic, female, 30-34 years				
...	All causes	742	100.0	57.7
1	Malignant neoplasms (C00-C97)	153	20.6	11.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	19.8	11.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	65	8.8	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	6.9	4.0
5	Assault (homicide) (X85-Y09,Y87.1)	49	6.6	3.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	26	3.5	2.0
7	Cerebrovascular diseases (I60-I69)	19	2.6	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	17	2.3	*
9	Diabetes mellitus (E10-E14)	16	2.2	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.3	*
12	Septicemia (A40-A41)	8	1.1	*
12	Pregnancy, childbirth and the puerperium (O00-099)	8	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
15	Influenza and pneumonia (J10-J18)	5	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
...	All other causes (Residual)	140	18.9	10.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,023	100.0	81.5
1	Malignant neoplasms (C00-C97)	273	26.7	21.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	14.9	12.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	8.3	6.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	84	8.2	6.7
5	Assault (homicide) (X85-Y09,Y87.1)	46	4.5	3.7
6	Cerebrovascular diseases (I60-I69)	40	3.9	3.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	3.9	3.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	36	3.5	2.9
9	Diabetes mellitus (E10-E14)	25	2.4	2.0
10	Pregnancy, childbirth and the puerperium (O00-099)	21	2.1	1.7
11	Chronic lower respiratory diseases (J40-J47)	20	2.0	1.6
12	Influenza and pneumonia (J10-J18)	12	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
14	Septicemia (A40-A41)	10	1.0	*
15	Viral hepatitis (B15-B19)	8	0.8	*
...	All other causes (Residual)	160	15.6	12.7
Hispanic, female, 40-44 years				
...	All causes	1,299	100.0	121.4
1	Malignant neoplasms (C00-C97)	427	32.9	39.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	11.4	13.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	105	8.1	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	8.0	9.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	61	4.7	5.7
6	Cerebrovascular diseases (I60-I69)	55	4.2	5.1
7	Diabetes mellitus (E10-E14)	48	3.7	4.5
8	Assault (homicide) (X85-Y09,Y87.1)	32	2.5	3.0
9	Chronic lower respiratory diseases (J40-J47)	23	1.8	2.1
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	1.7	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.5	1.9
12	Septicemia (A40-A41)	13	1.0	*
13	Viral hepatitis (B15-B19)	12	0.9	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.9	*
15	Influenza and pneumonia (J10-J18)	8	0.6	*
...	All other causes (Residual)	209	16.1	19.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	1,554	100.0	185.2
1	Malignant neoplasms (C00-C97)	554	35.6	66.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	11.9	22.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	6.6	12.3
4	Cerebrovascular diseases (I60-I69)	93	6.0	11.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	69	4.4	8.2
6	Diabetes mellitus (E10-E14)	64	4.1	7.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	63	4.1	7.5
8	Assault (homicide) (X85-Y09,Y87.1)	33	2.1	3.9
9	Chronic lower respiratory diseases (J40-J47)	31	2.0	3.7
10	Viral hepatitis (B15-B19)	23	1.5	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.3	2.4
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	17	1.1	*
13	Influenza and pneumonia (J10-J18)	16	1.0	*
14	Septicemia (A40-A41)	13	0.8	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	0.8	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.8	*
...	All other causes (Residual)	244	15.7	29.1
Hispanic, female, 50-54 years				
...	All causes	1,825	100.0	280.0
1	Malignant neoplasms (C00-C97)	725	39.7	111.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	273	15.0	41.9
3	Diabetes mellitus (E10-E14)	97	5.3	14.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	4.9	13.7
5	Cerebrovascular diseases (I60-I69)	86	4.7	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	71	3.9	10.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	38	2.1	5.8
8	Chronic lower respiratory diseases (J40-J47)	32	1.8	4.9
9	Viral hepatitis (B15-B19)	27	1.5	4.1
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	1.3	3.7
11	Septicemia (A40-A41)	23	1.3	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.3	3.5
13	Influenza and pneumonia (J10-J18)	17	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.9	*
15	Assault (homicide) (X85-Y09,Y87.1)	14	0.8	*
...	All other causes (Residual)	270	14.8	41.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	2,086	100.0	420.9
1	Malignant neoplasms (C00-C97)	734	35.2	148.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	407	19.5	82.1
3	Diabetes mellitus (E10-E14)	166	8.0	33.5
4	Cerebrovascular diseases (I60-I69)	129	6.2	26.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	68	3.3	13.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	3.2	13.5
7	Chronic lower respiratory diseases (J40-J47)	45	2.2	9.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.8	7.7
9	Septicemia (A40-A41)	26	1.2	5.2
9	Viral hepatitis (B15-B19)	26	1.2	5.2
9	Influenza and pneumonia (J10-J18)	26	1.2	5.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.6	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	10	0.5	*
15	Assault (homicide) (X85-Y09,Y87.1)	10	0.5	*
...	All other causes (Residual)	286	13.7	57.7
Hispanic, female, 60-64 years				
...	All causes	2,743	100.0	676.9
1	Malignant neoplasms (C00-C97)	892	32.5	220.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	657	24.0	162.1
3	Diabetes mellitus (E10-E14)	281	10.2	69.3
4	Cerebrovascular diseases (I60-I69)	134	4.9	33.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	89	3.2	22.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	2.4	16.0
7	Chronic lower respiratory diseases (J40-J47)	59	2.2	14.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.0	13.6
9	Influenza and pneumonia (J10-J18)	49	1.8	12.1
10	Septicemia (A40-A41)	27	1.0	6.7
11	Viral hepatitis (B15-B19)	26	0.9	6.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24	0.9	5.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.4	*
15	Pneumonitis due to solids and liquids (J69)	8	0.3	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	8	0.3	*
...	All other causes (Residual)	344	12.5	84.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	3,817	100.0	1,125.8
1	Malignant neoplasms (C00-C97)	1,129	29.6	333.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	942	24.7	277.8
3	Diabetes mellitus (E10-E14)	390	10.2	115.0
4	Cerebrovascular diseases (I60-I69)	251	6.6	74.0
5	Chronic lower respiratory diseases (J40-J47)	126	3.3	37.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	112	2.9	33.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.3	26.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.2	24.5
9	Influenza and pneumonia (J10-J18)	61	1.6	18.0
10	Septicemia (A40-A41)	52	1.4	15.3
11	Viral hepatitis (B15-B19)	36	0.9	10.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.8	8.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.7	7.7
14	Aortic aneurysm and dissection (I71)	12	0.3	*
15	Pneumonitis due to solids and liquids (J69)	9	0.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.2	*
...	All other causes (Residual)	462	12.1	136.3
Hispanic, female, 70-74 years				
...	All causes	4,601	100.0	1,662.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,260	27.4	455.2
2	Malignant neoplasms (C00-C97)	1,220	26.5	440.7
3	Diabetes mellitus (E10-E14)	452	9.8	163.3
4	Cerebrovascular diseases (I60-I69)	319	6.9	115.2
5	Chronic lower respiratory diseases (J40-J47)	150	3.3	54.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	111	2.4	40.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	2.2	37.2
8	Influenza and pneumonia (J10-J18)	80	1.7	28.9
9	Septicemia (A40-A41)	79	1.7	28.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.5	24.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	45	1.0	16.3
12	Alzheimer's disease (G30)	41	0.9	14.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.8	13.0
14	Viral hepatitis (B15-B19)	34	0.7	12.3
15	Parkinson's disease (G20-G21)	27	0.6	9.8
...	All other causes (Residual)	575	12.5	207.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	5,221	100.0	2,591.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,661	31.8	824.4
2	Malignant neoplasms (C00-C97)	1,150	22.0	570.8
3	Diabetes mellitus (E10-E14)	446	8.5	221.4
4	Cerebrovascular diseases (I60-I69)	400	7.7	198.5
5	Chronic lower respiratory diseases (J40-J47)	205	3.9	101.7
6	Influenza and pneumonia (J10-J18)	123	2.4	61.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	2.0	52.6
8	Alzheimer's disease (G30)	99	1.9	49.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.8	46.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	1.6	41.2
11	Septicemia (A40-A41)	72	1.4	35.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	54	1.0	26.8
13	Parkinson's disease (G20-G21)	37	0.7	18.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.6	16.4
15	Atherosclerosis (I70)	30	0.6	14.9
...	All other causes (Residual)	628	12.0	311.7
Hispanic, female, 80-84 years				
...	All causes	5,374	100.0	4,300.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,902	35.4	1,522.2
2	Malignant neoplasms (C00-C97)	896	16.7	717.1
3	Cerebrovascular diseases (I60-I69)	514	9.6	411.4
4	Diabetes mellitus (E10-E14)	361	6.7	288.9
5	Chronic lower respiratory diseases (J40-J47)	207	3.9	165.7
6	Influenza and pneumonia (J10-J18)	162	3.0	129.7
7	Alzheimer's disease (G30)	135	2.5	108.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	2.0	86.4
9	Septicemia (A40-A41)	80	1.5	64.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	79	1.5	63.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.3	55.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.0	44.0
13	Atherosclerosis (I70)	40	0.7	32.0
14	Parkinson's disease (G20-G21)	34	0.6	27.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	24.8
...	All other causes (Residual)	701	13.0	561.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	5,470	100.0	7,154.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,113	38.6	2,763.8
2	Malignant neoplasms (C00-C97)	667	12.2	872.4
3	Cerebrovascular diseases (I60-I69)	544	9.9	711.5
4	Diabetes mellitus (E10-E14)	299	5.5	391.1
5	Influenza and pneumonia (J10-J18)	228	4.2	298.2
6	Chronic lower respiratory diseases (J40-J47)	199	3.6	260.3
7	Alzheimer's disease (G30)	172	3.1	225.0
8	Septicemia (A40-A41)	85	1.6	111.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.4	103.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	1.3	95.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	54	1.0	70.6
12	Pneumonitis due to solids and liquids (J69)	46	0.8	60.2
13	Atherosclerosis (I70)	42	0.8	54.9
14	Parkinson's disease (G20-G21)	32	0.6	41.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.6	40.5
...	All other causes (Residual)	806	14.7	1,054.2
Hispanic, female, 90-94 years				
...	All causes	3,702	100.0	10,385.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,608	43.4	4,511.2
2	Cerebrovascular diseases (I60-I69)	322	8.7	903.4
3	Malignant neoplasms (C00-C97)	317	8.6	889.3
4	Influenza and pneumonia (J10-J18)	178	4.8	499.4
5	Alzheimer's disease (G30)	136	3.7	381.5
6	Diabetes mellitus (E10-E14)	133	3.6	373.1
7	Chronic lower respiratory diseases (J40-J47)	130	3.5	364.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	1.8	182.4
9	Septicemia (A40-A41)	55	1.5	154.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.4	145.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	1.2	129.1
12	Pneumonitis due to solids and liquids (J69)	45	1.2	126.2
13	Atherosclerosis (I70)	43	1.2	120.6
14	Parkinson's disease (G20-G21)	22	0.6	61.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.5	56.1
...	All other causes (Residual)	530	14.3	1,486.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	1,582	100.0	13,704.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	764	48.3	6,618.2
2	Cerebrovascular diseases (I60-I69)	120	7.6	1,039.5
3	Influenza and pneumonia (J10-J18)	101	6.4	874.9
4	Malignant neoplasms (C00-C97)	91	5.8	788.3
5	Alzheimer's disease (G30)	56	3.5	485.1
6	Diabetes mellitus (E10-E14)	49	3.1	424.5
7	Chronic lower respiratory diseases (J40-J47)	43	2.7	372.5
8	Septicemia (A40-A41)	27	1.7	233.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.7	233.9
10	Atherosclerosis (I70)	24	1.5	207.9
11	Pneumonitis due to solids and liquids (J69)	15	0.9	*
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	0.8	*
14	Nutritional deficiencies (E40-E64)	9	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	7	0.4	*
...	All other causes (Residual)	217	13.7	1,879.8
Hispanic, female, 100 years and over				
...	All causes	351	100.0	17,567.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	45.3	7,958.0
2	Influenza and pneumonia (J10-J18)	35	10.0	1,751.8
3	Cerebrovascular diseases (I60-I69)	23	6.6	1,151.2
4	Malignant neoplasms (C00-C97)	18	5.1	*
5	Alzheimer's disease (G30)	8	2.3	*
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.3	*
7	Septicemia (A40-A41)	6	1.7	*
7	Diabetes mellitus (E10-E14)	6	1.7	*
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.7	*
7	Pneumonitis due to solids and liquids (J69)	6	1.7	*
11	Atherosclerosis (I70)	5	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.1	*
14	Nutritional deficiencies (E40-E64)	3	0.9	*
15	Parkinson's disease (G20-G21)	2	0.6	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	*
...	All other causes (Residual)	56	16.0	2,802.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,953,197	100.0	996.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	607,704	31.1	310.0
2	Malignant neoplasms (C00-C97)	456,315	23.4	232.8
3	Cerebrovascular diseases (I60-I69)	138,716	7.1	70.8
4	Chronic lower respiratory diseases (J40-J47)	111,620	5.7	56.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,326	3.8	37.4
6	Influenza and pneumonia (J10-J18)	54,280	2.8	27.7
7	Diabetes mellitus (E10-E14)	49,347	2.5	25.2
8	Alzheimer's disease (G30)	40,835	2.1	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26,418	1.4	13.5
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,534	1.3	12.5
11	Septicemia (A40-A41)	23,023	1.2	11.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,602	1.0	10.0
13	Aortic aneurysm and dissection (I71)	13,934	0.7	7.1
14	Parkinson's disease (G20-G21)	13,475	0.7	6.9
15	Atherosclerosis (I70)	13,286	0.7	6.8
...	All other causes (Residual)	286,782	14.7	146.3
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,819	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	37.0	11.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	284	10.1	3.0
3	Malignant neoplasms (C00-C97)	281	10.0	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	156	5.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	3.3	1.0
6	Influenza and pneumonia (J10-J18)	58	2.1	0.6
7	Septicemia (A40-A41)	47	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	37	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	27	1.0	0.3
11	Meningitis (G00,G03)	26	0.9	0.3
12	Meningococcal infection (A39)	24	0.9	0.3
13	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	658	23.3	6.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,990	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	843	42.4	6.6
2	Malignant neoplasms (C00-C97)	311	15.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	5.7	0.9
4	Assault (homicide) (X85-Y09,Y87.1)	78	3.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	3.3	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	2.0	0.3
7	Influenza and pneumonia (J10-J18)	31	1.6	0.2
8	Septicemia (A40-A41)	29	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	19	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	17	0.9	*
12	Meningococcal infection (A39)	9	0.5	*
13	Diabetes mellitus (E10-E14)	8	0.4	*
13	Meningitis (G00,G03)	8	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	382	19.2	3.0
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,498	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,065	42.6	8.2
2	Malignant neoplasms (C00-C97)	297	11.9	2.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	173	6.9	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	5.2	1.0
5	Assault (homicide) (X85-Y09,Y87.1)	98	3.9	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.6	0.7
7	Chronic lower respiratory diseases (J40-J47)	32	1.3	0.2
8	Influenza and pneumonia (J10-J18)	31	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.0	0.2
10	Cerebrovascular diseases (I60-I69)	21	0.8	0.2
11	Septicemia (A40-A41)	18	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
13	Anemias (D50-D64)	12	0.5	*
14	Diabetes mellitus (E10-E14)	11	0.4	*
15	Meningitis (G00,G03)	10	0.4	*
...	All other causes (Residual)	468	18.7	3.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	8,490	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,840	57.0	36.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,169	13.8	8.9
3	Malignant neoplasms (C00-C97)	476	5.6	3.6
4	Assault (homicide) (X85-Y09,Y87.1)	409	4.8	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	269	3.2	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	1.7	1.1
7	Chronic lower respiratory diseases (J40-J47)	60	0.7	0.5
8	Influenza and pneumonia (J10-J18)	44	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	37	0.4	0.3
10	Diabetes mellitus (E10-E14)	36	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	0.2
12	Septicemia (A40-A41)	21	0.2	0.2
13	Meningococcal infection (A39)	20	0.2	0.2
14	Anemias (D50-D64)	16	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.2	*
...	All other causes (Residual)	904	10.6	6.9
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	9,378	100.0	78.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,570	48.7	38.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,661	17.7	14.0
3	Malignant neoplasms (C00-C97)	594	6.3	5.0
4	Assault (homicide) (X85-Y09,Y87.1)	553	5.9	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	3.4	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	68	0.7	0.6
8	Diabetes mellitus (E10-E14)	58	0.6	0.5
9	Influenza and pneumonia (J10-J18)	51	0.5	0.4
10	Septicemia (A40-A41)	50	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	50	0.5	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.3	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.2
15	Pregnancy, childbirth and the puerperium (O00-099)	18	0.2	*
...	All other causes (Residual)	1,179	12.6	9.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	10,048	100.0	82.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,854	38.4	31.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,776	17.7	14.6
3	Malignant neoplasms (C00-C97)	953	9.5	7.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	542	5.4	4.5
5	Assault (homicide) (X85-Y09,Y87.1)	531	5.3	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	188	1.9	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	1.5	1.2
8	Diabetes mellitus (E10-E14)	122	1.2	1.0
9	Cerebrovascular diseases (I60-I69)	100	1.0	0.8
10	Influenza and pneumonia (J10-J18)	75	0.7	0.6
11	Septicemia (A40-A41)	58	0.6	0.5
12	Chronic lower respiratory diseases (J40-J47)	57	0.6	0.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	45	0.4	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.4	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.3	0.3
...	All other causes (Residual)	1,523	15.2	12.5
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	13,938	100.0	103.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,001	28.7	29.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,151	15.4	15.9
3	Malignant neoplasms (C00-C97)	1,729	12.4	12.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,213	8.7	9.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	606	4.3	4.5
6	Assault (homicide) (X85-Y09,Y87.1)	604	4.3	4.5
7	Diabetes mellitus (E10-E14)	241	1.7	1.8
8	Cerebrovascular diseases (I60-I69)	201	1.4	1.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	198	1.4	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	1.3	1.4
11	Influenza and pneumonia (J10-J18)	124	0.9	0.9
12	Septicemia (A40-A41)	97	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	89	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	0.5
...	All other causes (Residual)	2,359	16.9	17.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	23,345	100.0	144.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,186	22.2	32.1
2	Malignant neoplasms (C00-C97)	4,025	17.2	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,998	12.8	18.6
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,699	11.6	16.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,044	4.5	6.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	712	3.0	4.4
7	Assault (homicide) (X85-Y09,Y87.1)	696	3.0	4.3
8	Cerebrovascular diseases (I60-I69)	514	2.2	3.2
9	Diabetes mellitus (E10-E14)	462	2.0	2.9
10	Influenza and pneumonia (J10-J18)	283	1.2	1.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	216	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	167	0.7	1.0
13	Septicemia (A40-A41)	165	0.7	1.0
14	Viral hepatitis (B15-B19)	131	0.6	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	0.5	0.8
...	All other causes (Residual)	3,924	16.8	24.3
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	34,285	100.0	208.0
1	Malignant neoplasms (C00-C97)	7,823	22.8	47.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,938	17.3	36.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,658	16.5	34.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,770	8.1	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,436	4.2	8.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,038	3.0	6.3
7	Cerebrovascular diseases (I60-I69)	892	2.6	5.4
8	Diabetes mellitus (E10-E14)	769	2.2	4.7
9	Assault (homicide) (X85-Y09,Y87.1)	631	1.8	3.8
10	Chronic lower respiratory diseases (J40-J47)	366	1.1	2.2
11	Influenza and pneumonia (J10-J18)	354	1.0	2.1
12	Viral hepatitis (B15-B19)	333	1.0	2.0
13	Septicemia (A40-A41)	274	0.8	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	0.6	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	0.6	1.2
...	All other causes (Residual)	5,597	16.3	34.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	45,974	100.0	312.7
1	Malignant neoplasms (C00-C97)	13,404	29.2	91.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,762	21.2	66.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,789	10.4	32.6
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,468	5.4	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,170	4.7	14.8
6	Cerebrovascular diseases (I60-I69)	1,302	2.8	8.9
7	Diabetes mellitus (E10-E14)	1,197	2.6	8.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	766	1.7	5.2
9	Chronic lower respiratory diseases (J40-J47)	747	1.6	5.1
10	Viral hepatitis (B15-B19)	580	1.3	3.9
11	Influenza and pneumonia (J10-J18)	503	1.1	3.4
12	Assault (homicide) (X85-Y09,Y87.1)	455	1.0	3.1
13	Septicemia (A40-A41)	421	0.9	2.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	0.6	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	265	0.6	1.8
...	All other causes (Residual)	6,873	14.9	46.7
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	60,834	100.0	471.6
1	Malignant neoplasms (C00-C97)	21,540	35.4	167.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,061	24.8	116.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,603	5.9	27.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,260	3.7	17.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,019	3.3	15.7
6	Cerebrovascular diseases (I60-I69)	1,841	3.0	14.3
7	Diabetes mellitus (E10-E14)	1,743	2.9	13.5
8	Chronic lower respiratory diseases (J40-J47)	1,625	2.7	12.6
9	Influenza and pneumonia (J10-J18)	606	1.0	4.7
10	Septicemia (A40-A41)	594	1.0	4.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	520	0.9	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	445	0.7	3.5
13	Viral hepatitis (B15-B19)	378	0.6	2.9
14	Assault (homicide) (X85-Y09,Y87.1)	359	0.6	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	295	0.5	2.3
...	All other causes (Residual)	7,945	13.1	61.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	78,170	100.0	764.4
1	Malignant neoplasms (C00-C97)	30,407	38.9	297.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,346	26.0	199.0
3	Chronic lower respiratory diseases (J40-J47)	3,555	4.5	34.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,968	3.8	29.0
5	Cerebrovascular diseases (I60-I69)	2,649	3.4	25.9
6	Diabetes mellitus (E10-E14)	2,382	3.0	23.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,013	2.6	19.7
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,501	1.9	14.7
9	Influenza and pneumonia (J10-J18)	797	1.0	7.8
10	Septicemia (A40-A41)	749	1.0	7.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	645	0.8	6.3
12	Aortic aneurysm and dissection (I71)	450	0.6	4.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	357	0.5	3.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	275	0.4	2.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	262	0.3	2.6
...	All other causes (Residual)	8,814	11.3	86.2
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	102,623	100.0	1,225.8
1	Malignant neoplasms (C00-C97)	40,381	39.3	482.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,604	26.9	329.7
3	Chronic lower respiratory diseases (J40-J47)	6,231	6.1	74.4
4	Cerebrovascular diseases (I60-I69)	3,662	3.6	43.7
5	Diabetes mellitus (E10-E14)	3,292	3.2	39.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,487	2.4	29.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,104	2.1	25.1
8	Influenza and pneumonia (J10-J18)	1,097	1.1	13.1
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,079	1.1	12.9
10	Septicemia (A40-A41)	1,043	1.0	12.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	991	1.0	11.8
12	Aortic aneurysm and dissection (I71)	745	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	542	0.5	6.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	411	0.4	4.9
15	Alzheimer's disease (G30)	288	0.3	3.4
...	All other causes (Residual)	10,666	10.4	127.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	148,130	100.0	1,942.0
1	Malignant neoplasms (C00-C97)	54,833	37.0	718.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,913	27.6	536.4
3	Chronic lower respiratory diseases (J40-J47)	11,099	7.5	145.5
4	Cerebrovascular diseases (I60-I69)	6,342	4.3	83.1
5	Diabetes mellitus (E10-E14)	4,820	3.3	63.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,875	1.9	37.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,191	1.5	28.7
8	Influenza and pneumonia (J10-J18)	1,843	1.2	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,738	1.2	22.8
10	Septicemia (A40-A41)	1,646	1.1	21.6
11	Aortic aneurysm and dissection (I71)	1,373	0.9	18.0
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,050	0.7	13.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	846	0.6	11.1
14	Alzheimer's disease (G30)	731	0.5	9.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	683	0.5	9.0
...	All other causes (Residual)	15,147	10.2	198.6
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	219,360	100.0	2,997.8
1	Malignant neoplasms (C00-C97)	72,286	33.0	987.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,837	28.6	858.7
3	Chronic lower respiratory diseases (J40-J47)	18,301	8.3	250.1
4	Cerebrovascular diseases (I60-I69)	11,990	5.5	163.9
5	Diabetes mellitus (E10-E14)	6,844	3.1	93.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,693	1.7	50.5
7	Influenza and pneumonia (J10-J18)	3,638	1.7	49.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,980	1.4	40.7
9	Septicemia (A40-A41)	2,494	1.1	34.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,291	1.0	31.3
11	Aortic aneurysm and dissection (I71)	2,175	1.0	29.7
12	Alzheimer's disease (G30)	2,090	1.0	28.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,430	0.7	19.5
14	Parkinson's disease (G20-G21)	1,336	0.6	18.3
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,199	0.5	16.4
...	All other causes (Residual)	23,776	10.8	324.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	288,018	100.0	4,602.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,727	30.8	1,417.7
2	Malignant neoplasms (C00-C97)	77,288	26.8	1,234.9
3	Chronic lower respiratory diseases (J40-J47)	22,770	7.9	363.8
4	Cerebrovascular diseases (I60-I69)	20,866	7.2	333.4
5	Diabetes mellitus (E10-E14)	8,523	3.0	136.2
6	Influenza and pneumonia (J10-J18)	6,686	2.3	106.8
7	Alzheimer's disease (G30)	5,195	1.8	83.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,946	1.7	79.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,440	1.5	70.9
10	Septicemia (A40-A41)	3,628	1.3	58.0
11	Aortic aneurysm and dissection (I71)	2,814	1.0	45.0
12	Parkinson's disease (G20-G21)	2,811	1.0	44.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,027	0.7	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,899	0.7	30.3
15	Pneumonitis due to solids and liquids (J69)	1,758	0.6	28.1
...	All other causes (Residual)	33,640	11.7	537.5
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	316,326	100.0	7,546.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,460	33.7	2,539.8
2	Malignant neoplasms (C00-C97)	64,007	20.2	1,527.0
3	Cerebrovascular diseases (I60-I69)	28,312	9.0	675.4
4	Chronic lower respiratory diseases (J40-J47)	21,892	6.9	522.3
5	Influenza and pneumonia (J10-J18)	10,099	3.2	240.9
6	Alzheimer's disease (G30)	9,258	2.9	220.9
7	Diabetes mellitus (E10-E14)	8,298	2.6	198.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,811	1.8	138.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,064	1.6	120.8
10	Septicemia (A40-A41)	4,089	1.3	97.6
11	Parkinson's disease (G20-G21)	3,783	1.2	90.2
12	Pneumonitis due to solids and liquids (J69)	2,603	0.8	62.1
13	Aortic aneurysm and dissection (I71)	2,593	0.8	61.9
14	Atherosclerosis (I70)	2,346	0.7	56.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,230	0.7	53.2
...	All other causes (Residual)	39,481	12.5	941.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	292,962	100.0	12,853.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	108,389	37.0	4,755.5
2	Malignant neoplasms (C00-C97)	42,207	14.4	1,851.8
3	Cerebrovascular diseases (I60-I69)	29,995	10.2	1,316.0
4	Chronic lower respiratory diseases (J40-J47)	15,486	5.3	679.4
5	Influenza and pneumonia (J10-J18)	11,998	4.1	526.4
6	Alzheimer's disease (G30)	10,953	3.7	480.6
7	Diabetes mellitus (E10-E14)	6,280	2.1	275.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,537	1.9	242.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,807	1.6	210.9
10	Septicemia (A40-A41)	3,974	1.4	174.4
11	Parkinson's disease (G20-G21)	3,124	1.1	137.1
12	Pneumonitis due to solids and liquids (J69)	3,086	1.1	135.4
13	Atherosclerosis (I70)	3,050	1.0	133.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,380	0.8	104.4
15	Aortic aneurysm and dissection (I71)	1,929	0.7	84.6
...	All other causes (Residual)	39,767	13.6	1,744.7
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	191,334	100.0	19,435.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,983	40.2	7,819.7
2	Cerebrovascular diseases (I60-I69)	20,832	10.9	2,116.0
3	Malignant neoplasms (C00-C97)	18,092	9.5	1,837.7
4	Influenza and pneumonia (J10-J18)	9,993	5.2	1,015.1
5	Alzheimer's disease (G30)	8,422	4.4	855.5
6	Chronic lower respiratory diseases (J40-J47)	6,810	3.6	691.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,625	1.9	368.2
8	Diabetes mellitus (E10-E14)	3,190	1.7	324.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,126	1.6	317.5
10	Atherosclerosis (I70)	2,725	1.4	276.8
11	Septicemia (A40-A41)	2,452	1.3	249.1
12	Pneumonitis due to solids and liquids (J69)	2,317	1.2	235.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,818	1.0	184.7
14	Parkinson's disease (G20-G21)	1,387	0.7	140.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	967	0.5	98.2
...	All other causes (Residual)	28,595	14.9	2,904.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	73,705	100.0	25,936.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,757	43.1	11,175.0
2	Cerebrovascular diseases (I60-I69)	7,676	10.4	2,701.1
3	Malignant neoplasms (C00-C97)	4,716	6.4	1,659.5
4	Influenza and pneumonia (J10-J18)	4,610	6.3	1,622.2
5	Alzheimer's disease (G30)	3,189	4.3	1,122.2
6	Chronic lower respiratory diseases (J40-J47)	1,928	2.6	678.4
7	Atherosclerosis (I70)	1,440	2.0	506.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	1.7	443.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,140	1.5	401.2
10	Diabetes mellitus (E10-E14)	934	1.3	328.7
11	Pneumonitis due to solids and liquids (J69)	919	1.2	323.4
12	Septicemia (A40-A41)	917	1.2	322.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	775	1.1	272.7
14	Nutritional deficiencies (E40-E64)	312	0.4	109.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	294	0.4	103.5
...	All other causes (Residual)	11,838	16.1	4,165.7
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	15,326	100.0	32,216.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,064	46.1	14,849.1
2	Cerebrovascular diseases (I60-I69)	1,315	8.6	2,764.2
3	Influenza and pneumonia (J10-J18)	1,237	8.1	2,600.3
4	Malignant neoplasms (C00-C97)	621	4.1	1,305.4
5	Alzheimer's disease (G30)	542	3.5	1,139.3
6	Atherosclerosis (I70)	402	2.6	845.0
7	Chronic lower respiratory diseases (J40-J47)	297	1.9	624.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	1.5	473.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	1.3	409.9
10	Pneumonitis due to solids and liquids (J69)	186	1.2	391.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	172	1.1	361.6
12	Septicemia (A40-A41)	136	0.9	285.9
13	Diabetes mellitus (E10-E14)	129	0.8	271.2
14	Nutritional deficiencies (E40-E64)	76	0.5	159.8
15	Anemias (D50-D64)	67	0.4	140.8
...	All other causes (Residual)	2,662	17.4	5,595.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	944,913	100.0	984.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	293,206	31.0	305.5
2	Malignant neoplasms (C00-C97)	236,384	25.0	246.3
3	Chronic lower respiratory diseases (J40-J47)	55,314	5.9	57.6
4	Cerebrovascular diseases (I60-I69)	51,968	5.5	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,113	4.9	48.1
6	Diabetes mellitus (E10-E14)	23,177	2.5	24.2
7	Influenza and pneumonia (J10-J18)	23,097	2.4	24.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,620	2.1	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,793	1.4	13.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	12,613	1.3	13.1
11	Alzheimer's disease (G30)	12,253	1.3	12.8
12	Septicemia (A40-A41)	9,863	1.0	10.3
13	Aortic aneurysm and dissection (I71)	8,551	0.9	8.9
14	Parkinson's disease (G20-G21)	7,648	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	6,642	0.7	6.9
...	All other causes (Residual)	125,671	13.3	131.0
White, non-Hispanic, male, 1-4 years				
...	All causes	1,605	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	38.4	12.7
2	Malignant neoplasms (C00-C97)	160	10.0	3.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	9.6	3.2
4	Assault (homicide) (X85-Y09,Y87.1)	83	5.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	1.1
6	Septicemia (A40-A41)	25	1.6	0.5
7	Certain conditions originating in the perinatal period (P00-P96)	24	1.5	0.5
8	Influenza and pneumonia (J10-J18)	22	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Meningococcal infection (A39)	15	0.9	*
10	Meningitis (G00,G03)	15	0.9	*
10	Cerebrovascular diseases (I60-I69)	15	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
...	All other causes (Residual)	377	23.5	7.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999
 [Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	1,115	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	45.3	7.7
2	Malignant neoplasms (C00-C97)	180	16.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	4.8	0.8
4	Assault (homicide) (X85-Y09,Y87.1)	41	3.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	3.2	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.7	*
7	Influenza and pneumonia (J10-J18)	17	1.5	*
8	Septicemia (A40-A41)	15	1.3	*
9	Cerebrovascular diseases (I60-I69)	13	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Meningococcal infection (A39)	4	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
12	Diabetes mellitus (E10-E14)	4	0.4	*
12	Meningitis (G00,G03)	4	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	195	17.5	3.0
White, non-Hispanic, male, 10-14 years				
...	All causes	1,528	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	700	45.8	10.6
2	Malignant neoplasms (C00-C97)	165	10.8	2.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	140	9.2	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	4.3	1.0
5	Assault (homicide) (X85-Y09,Y87.1)	52	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	3.1	0.7
7	Chronic lower respiratory diseases (J40-J47)	23	1.5	0.3
8	Influenza and pneumonia (J10-J18)	19	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.9	*
10	Septicemia (A40-A41)	9	0.6	*
11	Cerebrovascular diseases (I60-I69)	8	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.4	*
14	Anemias (D50-D64)	6	0.4	*
...	All other causes (Residual)	259	17.0	3.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,799	100.0	85.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,250	56.0	48.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	978	16.9	14.5
3	Malignant neoplasms (C00-C97)	283	4.9	4.2
3	Assault (homicide) (X85-Y09,Y87.1)	283	4.9	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	167	2.9	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	1.6	1.4
7	Chronic lower respiratory diseases (J40-J47)	32	0.6	0.5
8	Diabetes mellitus (E10-E14)	21	0.4	0.3
8	Cerebrovascular diseases (I60-I69)	21	0.4	0.3
8	Influenza and pneumonia (J10-J18)	21	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
12	Septicemia (A40-A41)	11	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
14	Meningococcal infection (A39)	8	0.1	*
14	Anemias (D50-D64)	8	0.1	*
...	All other causes (Residual)	602	10.4	8.9
White, non-Hispanic, male, 20-24 years				
...	All causes	6,878	100.0	112.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,499	50.9	57.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,438	20.9	23.6
3	Assault (homicide) (X85-Y09,Y87.1)	361	5.2	5.9
4	Malignant neoplasms (C00-C97)	346	5.0	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	2.8	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	1.3
7	Septicemia (A40-A41)	33	0.5	0.5
8	Diabetes mellitus (E10-E14)	31	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	30	0.4	0.5
9	Influenza and pneumonia (J10-J18)	30	0.4	0.5
11	Chronic lower respiratory diseases (J40-J47)	27	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.3	*
14	Legal intervention (Y35,Y89.0)	15	0.2	*
15	Aortic aneurysm and dissection (I71)	9	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.1	*
...	All other causes (Residual)	742	10.8	12.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	7,056	100.0	116.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,953	41.9	48.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,478	20.9	24.3
3	Malignant neoplasms (C00-C97)	510	7.2	8.4
4	Assault (homicide) (X85-Y09,Y87.1)	360	5.1	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	4.7	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	139	2.0	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.1	1.3
8	Diabetes mellitus (E10-E14)	63	0.9	1.0
9	Cerebrovascular diseases (I60-I69)	48	0.7	0.8
10	Influenza and pneumonia (J10-J18)	44	0.6	0.7
11	Septicemia (A40-A41)	33	0.5	0.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.4	0.5
13	Legal intervention (Y35,Y89.0)	28	0.4	0.5
14	Chronic lower respiratory diseases (J40-J47)	26	0.4	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.3	0.3
...	All other causes (Residual)	915	13.0	15.0
White, non-Hispanic, male, 30-34 years				
...	All causes	9,393	100.0	139.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,979	31.7	44.3
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,727	18.4	25.7
3	Malignant neoplasms (C00-C97)	831	8.8	12.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	821	8.7	12.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	504	5.4	7.5
6	Assault (homicide) (X85-Y09,Y87.1)	393	4.2	5.8
7	Diabetes mellitus (E10-E14)	140	1.5	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	122	1.3	1.8
9	Cerebrovascular diseases (I60-I69)	103	1.1	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	1.1	1.5
11	Influenza and pneumonia (J10-J18)	75	0.8	1.1
12	Septicemia (A40-A41)	54	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	40	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	0.6
...	All other causes (Residual)	1,426	15.2	21.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	15,144	100.0	187.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,770	24.9	46.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,134	14.1	26.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,098	13.9	25.9
4	Malignant neoplasms (C00-C97)	1,789	11.8	22.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	880	5.8	10.9
6	Assault (homicide) (X85-Y09,Y87.1)	455	3.0	5.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	445	2.9	5.5
8	Diabetes mellitus (E10-E14)	289	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	252	1.7	3.1
10	Influenza and pneumonia (J10-J18)	148	1.0	1.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.8	1.5
12	Septicemia (A40-A41)	89	0.6	1.1
13	Viral hepatitis (B15-B19)	88	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	77	0.5	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.5	0.9
...	All other causes (Residual)	2,439	16.1	30.1
White, non-Hispanic, male, 40-44 years				
...	All causes	22,147	100.0	268.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,456	20.1	54.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,179	18.9	50.7
3	Malignant neoplasms (C00-C97)	3,636	16.4	44.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,112	9.5	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,033	4.7	12.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	888	4.0	10.8
7	Cerebrovascular diseases (I60-I69)	460	2.1	5.6
8	Diabetes mellitus (E10-E14)	458	2.1	5.6
9	Assault (homicide) (X85-Y09,Y87.1)	426	1.9	5.2
10	Viral hepatitis (B15-B19)	245	1.1	3.0
11	Influenza and pneumonia (J10-J18)	186	0.8	2.3
12	Chronic lower respiratory diseases (J40-J47)	162	0.7	2.0
13	Septicemia (A40-A41)	154	0.7	1.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	0.5	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	0.5	1.3
...	All other causes (Residual)	3,525	15.9	42.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-49 years				
...	All causes	29,448	100.0	403.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,501	25.5	102.7
2	Malignant neoplasms (C00-C97)	6,622	22.5	90.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,531	12.0	48.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,876	6.4	25.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,647	5.6	22.5
6	Diabetes mellitus (E10-E14)	760	2.6	10.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	666	2.3	9.1
8	Cerebrovascular diseases (I60-I69)	664	2.3	9.1
9	Viral hepatitis (B15-B19)	423	1.4	5.8
10	Chronic lower respiratory diseases (J40-J47)	386	1.3	5.3
11	Assault (homicide) (X85-Y09,Y87.1)	328	1.1	4.5
12	Influenza and pneumonia (J10-J18)	309	1.0	4.2
13	Septicemia (A40-A41)	234	0.8	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	0.6	2.2
15	Aortic aneurysm and dissection (I71)	145	0.5	2.0
...	All other causes (Residual)	4,194	14.2	57.4
White, non-Hispanic, male, 50-54 years				
...	All causes	37,703	100.0	593.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,246	29.8	177.0
2	Malignant neoplasms (C00-C97)	10,966	29.1	172.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,611	6.9	41.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,718	4.6	27.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,512	4.0	23.8
6	Diabetes mellitus (E10-E14)	1,041	2.8	16.4
7	Cerebrovascular diseases (I60-I69)	998	2.6	15.7
8	Chronic lower respiratory diseases (J40-J47)	788	2.1	12.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	470	1.2	7.4
10	Influenza and pneumonia (J10-J18)	361	1.0	5.7
11	Septicemia (A40-A41)	317	0.8	5.0
12	Viral hepatitis (B15-B19)	294	0.8	4.6
13	Assault (homicide) (X85-Y09,Y87.1)	251	0.7	3.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	0.6	3.8
15	Aortic aneurysm and dissection (I71)	195	0.5	3.1
...	All other causes (Residual)	4,691	12.4	73.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	47,570	100.0	955.2
1	Malignant neoplasms (C00-C97)	16,371	34.4	328.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,675	30.8	294.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,040	4.3	41.0
4	Chronic lower respiratory diseases (J40-J47)	1,830	3.8	36.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,467	3.1	29.5
6	Cerebrovascular diseases (I60-I69)	1,452	3.1	29.2
7	Diabetes mellitus (E10-E14)	1,379	2.9	27.7
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,141	2.4	22.9
9	Influenza and pneumonia (J10-J18)	436	0.9	8.8
10	Septicemia (A40-A41)	381	0.8	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	0.8	7.3
12	Aortic aneurysm and dissection (I71)	353	0.7	7.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	224	0.5	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.4	4.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	181	0.4	3.6
...	All other causes (Residual)	5,075	10.7	101.9
White, non-Hispanic, male, 60-64 years				
...	All causes	61,122	100.0	1,521.3
1	Malignant neoplasms (C00-C97)	22,287	36.5	554.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,791	30.7	467.7
3	Chronic lower respiratory diseases (J40-J47)	3,261	5.3	81.2
4	Cerebrovascular diseases (I60-I69)	2,027	3.3	50.5
5	Diabetes mellitus (E10-E14)	1,821	3.0	45.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,690	2.8	42.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,418	2.3	35.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	860	1.4	21.4
9	Influenza and pneumonia (J10-J18)	613	1.0	15.3
10	Aortic aneurysm and dissection (I71)	575	0.9	14.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	538	0.9	13.4
12	Septicemia (A40-A41)	523	0.9	13.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	336	0.5	8.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	229	0.4	5.7
15	Pneumonitis due to solids and liquids (J69)	163	0.3	4.1
...	All other causes (Residual)	5,990	9.8	149.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	86,299	100.0	2,433.4
1	Malignant neoplasms (C00-C97)	30,626	35.5	863.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,627	30.9	750.8
3	Chronic lower respiratory diseases (J40-J47)	5,865	6.8	165.4
4	Cerebrovascular diseases (I60-I69)	3,320	3.8	93.6
5	Diabetes mellitus (E10-E14)	2,568	3.0	72.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,805	2.1	50.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,428	1.7	40.3
8	Influenza and pneumonia (J10-J18)	1,075	1.2	30.3
9	Aortic aneurysm and dissection (I71)	992	1.1	28.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	922	1.1	26.0
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	862	1.0	24.3
12	Septicemia (A40-A41)	847	1.0	23.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	503	0.6	14.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	361	0.4	10.2
15	Alzheimer's disease (G30)	338	0.4	9.5
...	All other causes (Residual)	8,160	9.5	230.1
White, non-Hispanic, male, 70-74 years				
...	All causes	122,856	100.0	3,780.8
1	Malignant neoplasms (C00-C97)	39,892	32.5	1,227.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,072	31.0	1,171.6
3	Chronic lower respiratory diseases (J40-J47)	9,694	7.9	298.3
4	Cerebrovascular diseases (I60-I69)	6,017	4.9	185.2
5	Diabetes mellitus (E10-E14)	3,498	2.8	107.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,157	1.8	66.4
7	Influenza and pneumonia (J10-J18)	2,012	1.6	61.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,610	1.3	49.5
9	Aortic aneurysm and dissection (I71)	1,459	1.2	44.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,318	1.1	40.6
11	Septicemia (A40-A41)	1,264	1.0	38.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,007	0.8	31.0
13	Alzheimer's disease (G30)	925	0.8	28.5
14	Parkinson's disease (G20-G21)	888	0.7	27.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	823	0.7	25.3
...	All other causes (Residual)	12,220	9.9	376.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	149,671	100.0	5,712.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,308	32.3	1,843.6
2	Malignant neoplasms (C00-C97)	41,239	27.6	1,573.8
3	Chronic lower respiratory diseases (J40-J47)	11,705	7.8	446.7
4	Cerebrovascular diseases (I60-I69)	9,436	6.3	360.1
5	Diabetes mellitus (E10-E14)	4,116	2.8	157.1
6	Influenza and pneumonia (J10-J18)	3,454	2.3	131.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,632	1.8	100.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,359	1.6	90.0
9	Alzheimer's disease (G30)	2,113	1.4	80.6
10	Parkinson's disease (G20-G21)	1,780	1.2	67.9
11	Aortic aneurysm and dissection (I71)	1,774	1.2	67.7
12	Septicemia (A40-A41)	1,762	1.2	67.2
13	Pneumonitis due to solids and liquids (J69)	1,063	0.7	40.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,021	0.7	39.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,002	0.7	38.2
...	All other causes (Residual)	15,907	10.6	607.1
White, non-Hispanic, male, 80-84 years				
...	All causes	146,857	100.0	9,286.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,114	34.1	3,169.1
2	Malignant neoplasms (C00-C97)	32,296	22.0	2,042.3
3	Cerebrovascular diseases (I60-I69)	11,182	7.6	707.1
4	Chronic lower respiratory diseases (J40-J47)	10,672	7.3	674.9
5	Influenza and pneumonia (J10-J18)	4,801	3.3	303.6
6	Diabetes mellitus (E10-E14)	3,606	2.5	228.0
7	Alzheimer's disease (G30)	3,319	2.3	209.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,919	2.0	184.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,548	1.7	161.1
10	Parkinson's disease (G20-G21)	2,225	1.5	140.7
11	Septicemia (A40-A41)	1,776	1.2	112.3
12	Aortic aneurysm and dissection (I71)	1,469	1.0	92.9
13	Pneumonitis due to solids and liquids (J69)	1,429	1.0	90.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,011	0.7	63.9
15	Atherosclerosis (I70)	966	0.7	61.1
...	All other causes (Residual)	16,524	11.3	1,044.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	111,859	100.0	15,318.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,681	36.4	5,571.1
2	Malignant neoplasms (C00-C97)	19,315	17.3	2,645.1
3	Cerebrovascular diseases (I60-I69)	9,548	8.5	1,307.6
4	Chronic lower respiratory diseases (J40-J47)	7,142	6.4	978.1
5	Influenza and pneumonia (J10-J18)	4,862	4.3	665.8
6	Alzheimer's disease (G30)	3,120	2.8	427.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,368	2.1	324.3
8	Diabetes mellitus (E10-E14)	2,310	2.1	316.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,144	1.9	293.6
10	Parkinson's disease (G20-G21)	1,593	1.4	218.2
11	Pneumonitis due to solids and liquids (J69)	1,492	1.3	204.3
12	Septicemia (A40-A41)	1,396	1.2	191.2
13	Atherosclerosis (I70)	981	0.9	134.3
14	Aortic aneurysm and dissection (I71)	917	0.8	125.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	778	0.7	106.5
...	All other causes (Residual)	13,212	11.8	1,809.3
White, non-Hispanic, male, 90-94 years				
...	All causes	55,125	100.0	21,460.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,346	38.7	8,309.9
2	Malignant neoplasms (C00-C97)	7,148	13.0	2,782.7
3	Cerebrovascular diseases (I60-I69)	4,884	8.9	1,901.3
4	Influenza and pneumonia (J10-J18)	3,160	5.7	1,230.2
5	Chronic lower respiratory diseases (J40-J47)	2,795	5.1	1,088.1
6	Alzheimer's disease (G30)	1,719	3.1	669.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,251	2.3	487.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,161	2.1	452.0
9	Pneumonitis due to solids and liquids (J69)	908	1.6	353.5
10	Diabetes mellitus (E10-E14)	876	1.6	341.0
11	Atherosclerosis (I70)	702	1.3	273.3
12	Septicemia (A40-A41)	659	1.2	256.5
13	Parkinson's disease (G20-G21)	590	1.1	229.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	413	0.7	160.8
15	Aortic aneurysm and dissection (I71)	369	0.7	143.7
...	All other causes (Residual)	7,144	13.0	2,781.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 95-99 years				
...	All causes	15,589	100.0	26,212.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,460	41.4	10,862.4
2	Malignant neoplasms (C00-C97)	1,519	9.7	2,554.2
3	Cerebrovascular diseases (I60-I69)	1,286	8.2	2,162.4
4	Influenza and pneumonia (J10-J18)	1,139	7.3	1,915.2
5	Chronic lower respiratory diseases (J40-J47)	666	4.3	1,119.9
6	Alzheimer's disease (G30)	451	2.9	758.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	362	2.3	608.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	334	2.1	561.6
9	Pneumonitis due to solids and liquids (J69)	323	2.1	543.1
10	Atherosclerosis (I70)	279	1.8	469.1
11	Septicemia (A40-A41)	184	1.2	309.4
12	Diabetes mellitus (E10-E14)	166	1.1	279.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	0.8	196.7
14	Parkinson's disease (G20-G21)	98	0.6	164.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	122.7
15	Aortic aneurysm and dissection (I71)	73	0.5	122.7
...	All other causes (Residual)	2,059	13.2	3,462.2
White, non-Hispanic, male, 100 years and over				
...	All causes	2,365	100.0	30,016.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	989	41.8	12,552.4
2	Influenza and pneumonia (J10-J18)	249	10.5	3,160.3
3	Malignant neoplasms (C00-C97)	179	7.6	2,271.9
4	Cerebrovascular diseases (I60-I69)	168	7.1	2,132.3
5	Chronic lower respiratory diseases (J40-J47)	92	3.9	1,167.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.3	685.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	2.2	672.7
8	Atherosclerosis (I70)	52	2.2	660.0
9	Pneumonitis due to solids and liquids (J69)	44	1.9	558.4
10	Alzheimer's disease (G30)	40	1.7	507.7
11	Diabetes mellitus (E10-E14)	22	0.9	279.2
12	Septicemia (A40-A41)	18	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
15	Anemias (D50-D64)	12	0.5	*
...	All other causes (Residual)	364	15.4	4,619.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	1,008,284	100.0	1,007.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314,498	31.2	314.2
2	Malignant neoplasms (C00-C97)	219,931	21.8	219.7
3	Cerebrovascular diseases (I60-I69)	86,748	8.6	86.7
4	Chronic lower respiratory diseases (J40-J47)	56,306	5.6	56.3
5	Influenza and pneumonia (J10-J18)	31,183	3.1	31.2
6	Alzheimer's disease (G30)	28,582	2.8	28.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,213	2.7	27.2
8	Diabetes mellitus (E10-E14)	26,170	2.6	26.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,625	1.4	13.6
10	Septicemia (A40-A41)	13,160	1.3	13.1
11	Atherosclerosis (I70)	8,559	0.8	8.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,653	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	6,989	0.7	7.0
14	Pneumonitis due to solids and liquids (J69)	6,497	0.6	6.5
15	Parkinson's disease (G20-G21)	5,827	0.6	5.8
...	All other causes (Residual)	155,343	15.4	155.2
White, non-Hispanic, female, 1-4 years				
...	All causes	1,214	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	35.0	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	10.7	2.8
3	Malignant neoplasms (C00-C97)	121	10.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	73	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J10-J18)	36	3.0	0.8
7	Septicemia (A40-A41)	22	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Meningitis (G00,G03)	11	0.9	*
13	Meningococcal infection (A39)	9	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
...	All other causes (Residual)	281	23.1	6.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	875	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	38.6	5.4
2	Malignant neoplasms (C00-C97)	131	15.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	6.9	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	37	4.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.3	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.4	0.3
7	Septicemia (A40-A41)	14	1.6	*
7	Influenza and pneumonia (J10-J18)	14	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	1.0	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningococcal infection (A39)	5	0.6	*
12	Anemias (D50-D64)	5	0.6	*
14	Diabetes mellitus (E10-E14)	4	0.5	*
14	Meningitis (G00,G03)	4	0.5	*
...	All other causes (Residual)	188	21.5	3.0
White, non-Hispanic, female, 10-14 years				
...	All causes	970	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	37.6	5.8
2	Malignant neoplasms (C00-C97)	132	13.6	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	6.7	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	46	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.4	0.7
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	3.4	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.3	*
7	Cerebrovascular diseases (I60-I69)	13	1.3	*
9	Influenza and pneumonia (J10-J18)	12	1.2	*
10	Septicemia (A40-A41)	9	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Diabetes mellitus (E10-E14)	6	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	207	21.3	3.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,691	100.0	42.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,590	59.1	25.0
2	Malignant neoplasms (C00-C97)	193	7.2	3.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	191	7.1	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	126	4.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	102	3.8	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	1.9	0.8
7	Chronic lower respiratory diseases (J40-J47)	28	1.0	0.4
8	Influenza and pneumonia (J10-J18)	23	0.9	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
10	Cerebrovascular diseases (I60-I69)	16	0.6	*
11	Diabetes mellitus (E10-E14)	15	0.6	*
12	Meningococcal infection (A39)	12	0.4	*
13	Septicemia (A40-A41)	10	0.4	*
14	Anemias (D50-D64)	8	0.3	*
14	Pregnancy, childbirth and the puerperium (O00-099)	8	0.3	*
...	All other causes (Residual)	300	11.1	4.7
White, non-Hispanic, female, 20-24 years				
...	All causes	2,500	100.0	43.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,071	42.8	18.4
2	Malignant neoplasms (C00-C97)	248	9.9	4.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	223	8.9	3.8
4	Assault (homicide) (X85-Y09,Y87.1)	192	7.7	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	5.0	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	2.0	0.9
7	Cerebrovascular diseases (I60-I69)	38	1.5	0.7
8	Diabetes mellitus (E10-E14)	27	1.1	0.5
9	Chronic lower respiratory diseases (J40-J47)	23	0.9	0.4
10	Influenza and pneumonia (J10-J18)	21	0.8	0.4
11	Pregnancy, childbirth and the puerperium (O00-099)	18	0.7	*
12	Septicemia (A40-A41)	17	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
...	All other causes (Residual)	412	16.5	7.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	2,992	100.0	49.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	901	30.1	14.8
2	Malignant neoplasms (C00-C97)	443	14.8	7.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	298	10.0	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	7.1	3.5
5	Assault (homicide) (X85-Y09,Y87.1)	171	5.7	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	2.4	1.2
7	Diabetes mellitus (E10-E14)	59	2.0	1.0
8	Cerebrovascular diseases (I60-I69)	52	1.7	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.6	0.8
10	Influenza and pneumonia (J10-J18)	31	1.0	0.5
10	Chronic lower respiratory diseases (J40-J47)	31	1.0	0.5
12	Pregnancy, childbirth and the puerperium (O00-099)	29	1.0	0.5
13	Septicemia (A40-A41)	25	0.8	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
...	All other causes (Residual)	581	19.4	9.6
White, non-Hispanic, female, 30-34 years				
...	All causes	4,545	100.0	67.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,022	22.5	15.1
2	Malignant neoplasms (C00-C97)	898	19.8	13.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	424	9.3	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	392	8.6	5.8
5	Assault (homicide) (X85-Y09,Y87.1)	211	4.6	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	102	2.2	1.5
7	Diabetes mellitus (E10-E14)	101	2.2	1.5
8	Cerebrovascular diseases (I60-I69)	98	2.2	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	1.8	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	76	1.7	1.1
11	Influenza and pneumonia (J10-J18)	49	1.1	0.7
11	Chronic lower respiratory diseases (J40-J47)	49	1.1	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	45	1.0	0.7
14	Septicemia (A40-A41)	43	0.9	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.7	0.5
...	All other causes (Residual)	917	20.2	13.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	8,201	100.0	101.7
1	Malignant neoplasms (C00-C97)	2,236	27.3	27.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,416	17.3	17.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	864	10.5	10.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	601	7.3	7.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	267	3.3	3.3
6	Cerebrovascular diseases (I60-I69)	262	3.2	3.2
7	Assault (homicide) (X85-Y09,Y87.1)	241	2.9	3.0
8	Diabetes mellitus (E10-E14)	173	2.1	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	164	2.0	2.0
10	Influenza and pneumonia (J10-J18)	135	1.6	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	90	1.1	1.1
13	Septicemia (A40-A41)	76	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.6	0.6
15	Viral hepatitis (B15-B19)	43	0.5	0.5
...	All other causes (Residual)	1,485	18.1	18.4
White, non-Hispanic, female, 40-44 years				
...	All causes	12,138	100.0	147.4
1	Malignant neoplasms (C00-C97)	4,187	34.5	50.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,482	12.2	18.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,479	12.2	18.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	658	5.4	8.0
5	Cerebrovascular diseases (I60-I69)	432	3.6	5.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	403	3.3	4.9
7	Diabetes mellitus (E10-E14)	311	2.6	3.8
8	Assault (homicide) (X85-Y09,Y87.1)	205	1.7	2.5
9	Chronic lower respiratory diseases (J40-J47)	204	1.7	2.5
10	Influenza and pneumonia (J10-J18)	168	1.4	2.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	150	1.2	1.8
12	Septicemia (A40-A41)	120	1.0	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.8	1.2
14	Viral hepatitis (B15-B19)	88	0.7	1.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.7	1.0
...	All other causes (Residual)	2,072	17.1	25.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	16,526	100.0	223.4
1	Malignant neoplasms (C00-C97)	6,782	41.0	91.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,261	13.7	30.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,258	7.6	17.0
4	Cerebrovascular diseases (I60-I69)	638	3.9	8.6
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	592	3.6	8.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	523	3.2	7.1
7	Diabetes mellitus (E10-E14)	437	2.6	5.9
8	Chronic lower respiratory diseases (J40-J47)	361	2.2	4.9
9	Influenza and pneumonia (J10-J18)	194	1.2	2.6
10	Septicemia (A40-A41)	187	1.1	2.5
11	Viral hepatitis (B15-B19)	157	1.0	2.1
12	Assault (homicide) (X85-Y09,Y87.1)	127	0.8	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	0.7	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	0.7	1.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.6	1.4
...	All other causes (Residual)	2,676	16.2	36.2
White, non-Hispanic, female, 50-54 years				
...	All causes	23,131	100.0	353.5
1	Malignant neoplasms (C00-C97)	10,574	45.7	161.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,815	16.5	58.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	992	4.3	15.2
4	Cerebrovascular diseases (I60-I69)	843	3.6	12.9
5	Chronic lower respiratory diseases (J40-J47)	837	3.6	12.8
6	Diabetes mellitus (E10-E14)	702	3.0	10.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	542	2.3	8.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	507	2.2	7.7
9	Septicemia (A40-A41)	277	1.2	4.2
10	Influenza and pneumonia (J10-J18)	245	1.1	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	0.9	3.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.5	1.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.5	1.8
14	Assault (homicide) (X85-Y09,Y87.1)	108	0.5	1.7
15	Viral hepatitis (B15-B19)	84	0.4	1.3
...	All other causes (Residual)	3,158	13.7	48.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-59 years				
...	All causes	30,600	100.0	583.3
1	Malignant neoplasms (C00-C97)	14,036	45.9	267.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,671	18.5	108.1
3	Chronic lower respiratory diseases (J40-J47)	1,725	5.6	32.9
4	Cerebrovascular diseases (I60-I69)	1,197	3.9	22.8
5	Diabetes mellitus (E10-E14)	1,003	3.3	19.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	928	3.0	17.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	546	1.8	10.4
8	Septicemia (A40-A41)	368	1.2	7.0
9	Influenza and pneumonia (J10-J18)	361	1.2	6.9
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	360	1.2	6.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	280	0.9	5.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	3.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	0.4	2.4
14	Aortic aneurysm and dissection (I71)	97	0.3	1.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	94	0.3	1.8
...	All other causes (Residual)	3,651	11.9	69.6
White, non-Hispanic, female, 60-64 years				
...	All causes	41,501	100.0	953.0
1	Malignant neoplasms (C00-C97)	18,094	43.6	415.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,813	21.2	202.4
3	Chronic lower respiratory diseases (J40-J47)	2,970	7.2	68.2
4	Cerebrovascular diseases (I60-I69)	1,635	3.9	37.5
5	Diabetes mellitus (E10-E14)	1,471	3.5	33.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	1.9	18.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	686	1.7	15.8
8	Septicemia (A40-A41)	520	1.3	11.9
9	Influenza and pneumonia (J10-J18)	484	1.2	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	1.1	10.4
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	219	0.5	5.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	206	0.5	4.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	182	0.4	4.2
14	Aortic aneurysm and dissection (I71)	170	0.4	3.9
15	Alzheimer's disease (G30)	149	0.4	3.4
...	All other causes (Residual)	4,652	11.2	106.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 65-69 years				
...	All causes	61,831	100.0	1,515.0
1	Malignant neoplasms (C00-C97)	24,207	39.2	593.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,286	23.1	350.0
3	Chronic lower respiratory diseases (J40-J47)	5,234	8.5	128.2
4	Cerebrovascular diseases (I60-I69)	3,022	4.9	74.0
5	Diabetes mellitus (E10-E14)	2,252	3.6	55.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,070	1.7	26.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	816	1.3	20.0
8	Septicemia (A40-A41)	799	1.3	19.6
9	Influenza and pneumonia (J10-J18)	768	1.2	18.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	763	1.2	18.7
11	Alzheimer's disease (G30)	393	0.6	9.6
12	Aortic aneurysm and dissection (I71)	381	0.6	9.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	343	0.6	8.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	322	0.5	7.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	188	0.3	4.6
...	All other causes (Residual)	6,987	11.3	171.2
White, non-Hispanic, female, 70-74 years				
...	All causes	96,504	100.0	2,372.3
1	Malignant neoplasms (C00-C97)	32,394	33.6	796.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,765	25.7	608.8
3	Chronic lower respiratory diseases (J40-J47)	8,607	8.9	211.6
4	Cerebrovascular diseases (I60-I69)	5,973	6.2	146.8
5	Diabetes mellitus (E10-E14)	3,346	3.5	82.3
6	Influenza and pneumonia (J10-J18)	1,626	1.7	40.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,536	1.6	37.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,370	1.4	33.7
9	Septicemia (A40-A41)	1,230	1.3	30.2
10	Alzheimer's disease (G30)	1,165	1.2	28.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	973	1.0	23.9
12	Aortic aneurysm and dissection (I71)	716	0.7	17.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	607	0.6	14.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	569	0.6	14.0
15	Parkinson's disease (G20-G21)	448	0.5	11.0
...	All other causes (Residual)	11,179	11.6	274.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	138,347	100.0	3,802.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,419	29.2	1,110.9
2	Malignant neoplasms (C00-C97)	36,049	26.1	990.8
3	Cerebrovascular diseases (I60-I69)	11,430	8.3	314.2
4	Chronic lower respiratory diseases (J40-J47)	11,065	8.0	304.1
5	Diabetes mellitus (E10-E14)	4,407	3.2	121.1
6	Influenza and pneumonia (J10-J18)	3,232	2.3	88.8
7	Alzheimer's disease (G30)	3,082	2.2	84.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,314	1.7	63.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,081	1.5	57.2
10	Septicemia (A40-A41)	1,866	1.3	51.3
11	Aortic aneurysm and dissection (I71)	1,040	0.8	28.6
12	Parkinson's disease (G20-G21)	1,031	0.7	28.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,006	0.7	27.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	996	0.7	27.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	897	0.6	24.7
...	All other causes (Residual)	17,432	12.6	479.1
White, non-Hispanic, female, 80-84 years				
...	All causes	169,469	100.0	6,492.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,346	33.2	2,158.6
2	Malignant neoplasms (C00-C97)	31,711	18.7	1,214.8
3	Cerebrovascular diseases (I60-I69)	17,130	10.1	656.2
4	Chronic lower respiratory diseases (J40-J47)	11,220	6.6	429.8
5	Alzheimer's disease (G30)	5,939	3.5	227.5
6	Influenza and pneumonia (J10-J18)	5,298	3.1	203.0
7	Diabetes mellitus (E10-E14)	4,692	2.8	179.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,892	1.7	110.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,516	1.5	96.4
10	Septicemia (A40-A41)	2,313	1.4	88.6
11	Parkinson's disease (G20-G21)	1,558	0.9	59.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,418	0.8	54.3
13	Atherosclerosis (I70)	1,380	0.8	52.9
14	Pneumonitis due to solids and liquids (J69)	1,174	0.7	45.0
15	Aortic aneurysm and dissection (I71)	1,124	0.7	43.1
...	All other causes (Residual)	22,758	13.4	871.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 85-89 years				
...	All causes	181,103	100.0	11,691.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,708	37.4	4,371.0
2	Malignant neoplasms (C00-C97)	22,892	12.6	1,477.8
3	Cerebrovascular diseases (I60-I69)	20,447	11.3	1,320.0
4	Chronic lower respiratory diseases (J40-J47)	8,344	4.6	538.7
5	Alzheimer's disease (G30)	7,833	4.3	505.7
6	Influenza and pneumonia (J10-J18)	7,136	3.9	460.7
7	Diabetes mellitus (E10-E14)	3,970	2.2	256.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,169	1.7	204.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,663	1.5	171.9
10	Septicemia (A40-A41)	2,578	1.4	166.4
11	Atherosclerosis (I70)	2,069	1.1	133.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,686	0.9	108.8
13	Pneumonitis due to solids and liquids (J69)	1,594	0.9	102.9
14	Parkinson's disease (G20-G21)	1,531	0.8	98.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,051	0.6	67.8
...	All other causes (Residual)	26,432	14.6	1,706.3
White, non-Hispanic, female, 90-94 years				
...	All causes	136,209	100.0	18,720.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,637	40.8	7,646.6
2	Cerebrovascular diseases (I60-I69)	15,948	11.7	2,191.8
3	Malignant neoplasms (C00-C97)	10,944	8.0	1,504.1
4	Influenza and pneumonia (J10-J18)	6,833	5.0	939.1
5	Alzheimer's disease (G30)	6,703	4.9	921.2
6	Chronic lower respiratory diseases (J40-J47)	4,015	2.9	551.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,374	1.7	326.3
8	Diabetes mellitus (E10-E14)	2,314	1.7	318.0
9	Atherosclerosis (I70)	2,023	1.5	278.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,965	1.4	270.1
11	Septicemia (A40-A41)	1,793	1.3	246.4
12	Pneumonitis due to solids and liquids (J69)	1,409	1.0	193.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,405	1.0	193.1
14	Parkinson's disease (G20-G21)	797	0.6	109.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	622	0.5	85.5
...	All other causes (Residual)	21,427	15.7	2,944.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	58,116	100.0	25,862.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,297	43.5	11,257.7
2	Cerebrovascular diseases (I60-I69)	6,390	11.0	2,843.7
3	Influenza and pneumonia (J10-J18)	3,471	6.0	1,544.7
4	Malignant neoplasms (C00-C97)	3,197	5.5	1,422.7
5	Alzheimer's disease (G30)	2,738	4.7	1,218.5
6	Chronic lower respiratory diseases (J40-J47)	1,262	2.2	561.6
7	Atherosclerosis (I70)	1,161	2.0	516.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	926	1.6	412.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	778	1.3	346.2
10	Diabetes mellitus (E10-E14)	768	1.3	341.8
11	Septicemia (A40-A41)	733	1.3	326.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	658	1.1	292.8
13	Pneumonitis due to solids and liquids (J69)	596	1.0	265.2
14	Nutritional deficiencies (E40-E64)	263	0.5	117.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	221	0.4	98.3
...	All other causes (Residual)	9,657	16.6	4,297.6
White, non-Hispanic, female, 100 years and over				
...	All causes	12,961	100.0	32,653.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,075	46.9	15,305.0
2	Cerebrovascular diseases (I60-I69)	1,147	8.8	2,889.7
3	Influenza and pneumonia (J10-J18)	988	7.6	2,489.1
4	Alzheimer's disease (G30)	502	3.9	1,264.7
5	Malignant neoplasms (C00-C97)	442	3.4	1,113.5
6	Atherosclerosis (I70)	350	2.7	881.8
7	Chronic lower respiratory diseases (J40-J47)	205	1.6	516.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.3	433.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	156	1.2	393.0
10	Pneumonitis due to solids and liquids (J69)	142	1.1	357.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.1	355.2
12	Septicemia (A40-A41)	118	0.9	297.3
13	Diabetes mellitus (E10-E14)	107	0.8	269.6
14	Nutritional deficiencies (E40-E64)	70	0.5	176.4
15	Anemias (D50-D64)	55	0.4	138.6
...	All other causes (Residual)	2,291	17.7	5,771.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	281,979	100.0	852.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,713	27.6	234.8
2	Malignant neoplasms (C00-C97)	61,409	21.8	185.6
3	Cerebrovascular diseases (I60-I69)	18,731	6.6	56.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,533	4.4	37.9
5	Diabetes mellitus (E10-E14)	11,830	4.2	35.7
6	Chronic lower respiratory diseases (J40-J47)	7,840	2.8	23.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,751	2.7	23.4
8	Assault (homicide) (X85-Y09,Y87.1)	7,536	2.7	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,659	2.4	20.1
10	Influenza and pneumonia (J10-J18)	5,794	2.1	17.5
11	Septicemia (A40-A41)	5,772	2.0	17.4
12	Certain conditions originating in the perinatal period (P00-P96)	5,000	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,708	1.3	11.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,793	1.0	8.4
15	Alzheimer's disease (G30)	2,325	0.8	7.0
...	All other causes (Residual)	44,585	15.8	134.7
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,283	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	33.4	20.6
2	Assault (homicide) (X85-Y09,Y87.1)	151	11.8	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	10.7	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.8	2.4
5	Malignant neoplasms (C00-C97)	47	3.7	2.3
6	Influenza and pneumonia (J10-J18)	40	3.1	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	29	2.3	1.4
8	Anemias (D50-D64)	20	1.6	1.0
9	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
11	Septicemia (A40-A41)	16	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
13	Meningitis (G00,G03)	11	0.9	*
14	Cerebrovascular diseases (I60-I69)	10	0.8	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
...	All other causes (Residual)	289	22.5	13.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	869	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	45.1	13.2
2	Malignant neoplasms (C00-C97)	84	9.7	2.8
3	Assault (homicide) (X85-Y09,Y87.1)	65	7.5	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	5.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.7	1.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	26	3.0	0.9
7	Chronic lower respiratory diseases (J40-J47)	19	2.2	*
8	Anemias (D50-D64)	17	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
10	Influenza and pneumonia (J10-J18)	8	0.9	*
11	Septicemia (A40-A41)	7	0.8	*
12	Meningitis (G00,G03)	4	0.5	*
12	Diseases of appendix (K35-K38)	4	0.5	*
14	Cerebrovascular diseases (I60-I69)	3	0.3	*
15	Meningococcal infection (A39)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.2	*
...	All other causes (Residual)	141	16.2	4.8
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	902	100.0	30.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	34.3	10.6
2	Malignant neoplasms (C00-C97)	95	10.5	3.2
3	Assault (homicide) (X85-Y09,Y87.1)	80	8.9	2.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	6.2	1.9
5	Chronic lower respiratory diseases (J40-J47)	47	5.2	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	5.0	1.5
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	3.2	1.0
8	Anemias (D50-D64)	16	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.7	*
10	Cerebrovascular diseases (I60-I69)	12	1.3	*
11	Diabetes mellitus (E10-E14)	8	0.9	*
12	Septicemia (A40-A41)	7	0.8	*
12	Influenza and pneumonia (J10-J18)	7	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
...	All other causes (Residual)	165	18.3	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,813	100.0	97.3
1	Assault (homicide) (X85-Y09,Y87.1)	1,117	39.7	38.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	764	27.2	26.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	176	6.3	6.1
4	Malignant neoplasms (C00-C97)	136	4.8	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	4.6	4.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	1.5	1.5
7	Chronic lower respiratory diseases (J40-J47)	36	1.3	1.2
8	Anemias (D50-D64)	23	0.8	0.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.7	0.7
10	Cerebrovascular diseases (I60-I69)	18	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-099)	18	0.6	*
12	Influenza and pneumonia (J10-J18)	14	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
15	Legal intervention (Y35,Y89.0)	10	0.4	*
...	All other causes (Residual)	286	10.2	9.9
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,167	100.0	163.0
1	Assault (homicide) (X85-Y09,Y87.1)	1,635	39.2	64.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	930	22.3	36.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	287	6.9	11.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	5.0	8.1
5	Malignant neoplasms (C00-C97)	178	4.3	7.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	2.6	4.3
7	Anemias (D50-D64)	61	1.5	2.4
8	Chronic lower respiratory diseases (J40-J47)	42	1.0	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.6
10	Influenza and pneumonia (J10-J18)	35	0.8	1.4
11	Diabetes mellitus (E10-E14)	31	0.7	1.2
12	Pregnancy, childbirth and the puerperium (O00-099)	28	0.7	1.1
13	Septicemia (A40-A41)	27	0.6	1.1
14	Cerebrovascular diseases (I60-I69)	22	0.5	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.9
...	All other causes (Residual)	512	12.3	20.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,390	100.0	177.4
1	Assault (homicide) (X85-Y09,Y87.1)	1,262	28.7	51.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	841	19.2	34.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	446	10.2	18.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	318	7.2	12.8
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	258	5.9	10.4
6	Malignant neoplasms (C00-C97)	227	5.2	9.2
7	Chronic lower respiratory diseases (J40-J47)	67	1.5	2.7
8	Anemias (D50-D64)	55	1.3	2.2
9	Cerebrovascular diseases (I60-I69)	49	1.1	2.0
10	Diabetes mellitus (E10-E14)	48	1.1	1.9
11	Influenza and pneumonia (J10-J18)	47	1.1	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.9	1.6
13	Septicemia (A40-A41)	37	0.8	1.5
13	Pregnancy, childbirth and the puerperium (O00-099)	37	0.8	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.6	1.1
...	All other causes (Residual)	630	14.4	25.5
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,602	100.0	222.1
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,033	18.4	41.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	897	16.0	35.6
3	Assault (homicide) (X85-Y09,Y87.1)	860	15.4	34.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	595	10.6	23.6
5	Malignant neoplasms (C00-C97)	433	7.7	17.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	239	4.3	9.5
7	Diabetes mellitus (E10-E14)	112	2.0	4.4
8	Cerebrovascular diseases (I60-I69)	100	1.8	4.0
9	Anemias (D50-D64)	84	1.5	3.3
10	Chronic lower respiratory diseases (J40-J47)	73	1.3	2.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.1	2.5
12	Influenza and pneumonia (J10-J18)	60	1.1	2.4
13	Septicemia (A40-A41)	48	0.9	1.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	38	0.7	1.5
15	Pregnancy, childbirth and the puerperium (O00-099)	33	0.6	1.3
...	All other causes (Residual)	934	16.7	37.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	8,563	100.0	311.6
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,458	17.0	53.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,246	14.6	45.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,063	12.4	38.7
4	Malignant neoplasms (C00-C97)	1,049	12.3	38.2
5	Assault (homicide) (X85-Y09,Y87.1)	730	8.5	26.6
6	Cerebrovascular diseases (I60-I69)	267	3.1	9.7
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	244	2.8	8.9
8	Diabetes mellitus (E10-E14)	192	2.2	7.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.5	4.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	1.4	4.3
11	Influenza and pneumonia (J10-J18)	116	1.4	4.2
12	Septicemia (A40-A41)	109	1.3	4.0
13	Chronic lower respiratory diseases (J40-J47)	93	1.1	3.4
14	Anemias (D50-D64)	66	0.8	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	0.8	2.4
...	All other causes (Residual)	1,618	18.9	58.9
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	12,207	100.0	466.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,260	18.5	86.3
2	Malignant neoplasms (C00-C97)	2,000	16.4	76.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,696	13.9	64.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	9.7	45.1
5	Assault (homicide) (X85-Y09,Y87.1)	575	4.7	22.0
6	Cerebrovascular diseases (I60-I69)	541	4.4	20.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	324	2.7	12.4
8	Diabetes mellitus (E10-E14)	310	2.5	11.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.6	7.3
10	Influenza and pneumonia (J10-J18)	189	1.5	7.2
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	186	1.5	7.1
12	Septicemia (A40-A41)	170	1.4	6.5
13	Chronic lower respiratory diseases (J40-J47)	165	1.4	6.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	110	0.9	4.2
15	Viral hepatitis (B15-B19)	94	0.8	3.6
...	All other causes (Residual)	2,213	18.1	84.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	15,131	100.0	706.9
1	Malignant neoplasms (C00-C97)	3,278	21.7	153.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,239	21.4	151.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,322	8.7	61.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,138	7.5	53.2
5	Cerebrovascular diseases (I60-I69)	780	5.2	36.4
6	Diabetes mellitus (E10-E14)	513	3.4	24.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	507	3.4	23.7
8	Assault (homicide) (X85-Y09,Y87.1)	364	2.4	17.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	1.8	12.6
10	Chronic lower respiratory diseases (J40-J47)	256	1.7	12.0
11	Septicemia (A40-A41)	251	1.7	11.7
12	Influenza and pneumonia (J10-J18)	220	1.5	10.3
13	Viral hepatitis (B15-B19)	160	1.1	7.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	144	1.0	6.7
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	144	1.0	6.7
...	All other causes (Residual)	2,546	16.8	119.0
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	16,246	100.0	1,004.9
1	Malignant neoplasms (C00-C97)	4,471	27.5	276.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,209	25.9	260.3
3	Cerebrovascular diseases (I60-I69)	882	5.4	54.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	765	4.7	47.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	735	4.5	45.5
6	Diabetes mellitus (E10-E14)	706	4.3	43.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	398	2.4	24.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	2.1	21.4
9	Chronic lower respiratory diseases (J40-J47)	300	1.8	18.6
10	Septicemia (A40-A41)	267	1.6	16.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	226	1.4	14.0
12	Influenza and pneumonia (J10-J18)	205	1.3	12.7
13	Assault (homicide) (X85-Y09,Y87.1)	193	1.2	11.9
14	Viral hepatitis (B15-B19)	117	0.7	7.2
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	82	0.5	5.1
...	All other causes (Residual)	2,344	14.4	145.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	18,176	100.0	1,470.9
1	Malignant neoplasms (C00-C97)	5,587	30.7	452.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,202	28.6	421.0
3	Cerebrovascular diseases (I60-I69)	1,017	5.6	82.3
4	Diabetes mellitus (E10-E14)	948	5.2	76.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	562	3.1	45.5
6	Chronic lower respiratory diseases (J40-J47)	472	2.6	38.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	421	2.3	34.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	378	2.1	30.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	363	2.0	29.4
10	Septicemia (A40-A41)	329	1.8	26.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	221	1.2	17.9
12	Influenza and pneumonia (J10-J18)	208	1.1	16.8
13	Assault (homicide) (X85-Y09,Y87.1)	115	0.6	9.3
14	Aortic aneurysm and dissection (I71)	82	0.5	6.6
15	Viral hepatitis (B15-B19)	73	0.4	5.9
...	All other causes (Residual)	2,198	12.1	177.9
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	20,887	100.0	2,060.8
1	Malignant neoplasms (C00-C97)	6,735	32.2	664.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,144	29.4	606.2
3	Cerebrovascular diseases (I60-I69)	1,252	6.0	123.5
4	Diabetes mellitus (E10-E14)	1,188	5.7	117.2
5	Chronic lower respiratory diseases (J40-J47)	613	2.9	60.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	2.4	49.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	2.3	47.9
8	Septicemia (A40-A41)	387	1.9	38.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	328	1.6	32.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	285	1.4	28.1
11	Influenza and pneumonia (J10-J18)	278	1.3	27.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	192	0.9	18.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.4	8.3
14	Assault (homicide) (X85-Y09,Y87.1)	79	0.4	7.8
15	Aortic aneurysm and dissection (I71)	70	0.3	6.9
...	All other causes (Residual)	2,266	10.8	223.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	25,217	100.0	2,802.3
1	Malignant neoplasms (C00-C97)	7,888	31.3	876.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,622	30.2	847.0
3	Cerebrovascular diseases (I60-I69)	1,529	6.1	169.9
4	Diabetes mellitus (E10-E14)	1,438	5.7	159.8
5	Chronic lower respiratory diseases (J40-J47)	899	3.6	99.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	698	2.8	77.6
7	Septicemia (A40-A41)	520	2.1	57.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	1.8	50.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	342	1.4	38.0
10	Influenza and pneumonia (J10-J18)	327	1.3	36.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	250	1.0	27.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	132	0.5	14.7
13	Aortic aneurysm and dissection (I71)	126	0.5	14.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	13.8
15	Pneumonitis due to solids and liquids (J69)	90	0.4	10.0
...	All other causes (Residual)	2,778	11.0	308.7
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	30,350	100.0	4,227.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,397	31.0	1,308.9
2	Malignant neoplasms (C00-C97)	8,737	28.8	1,217.0
3	Cerebrovascular diseases (I60-I69)	2,263	7.5	315.2
4	Diabetes mellitus (E10-E14)	1,682	5.5	234.3
5	Chronic lower respiratory diseases (J40-J47)	1,191	3.9	165.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	856	2.8	119.2
7	Septicemia (A40-A41)	692	2.3	96.4
8	Influenza and pneumonia (J10-J18)	553	1.8	77.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	489	1.6	68.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	422	1.4	58.8
11	Alzheimer's disease (G30)	187	0.6	26.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	182	0.6	25.4
13	Pneumonitis due to solids and liquids (J69)	174	0.6	24.2
14	Aortic aneurysm and dissection (I71)	145	0.5	20.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.4	18.4
...	All other causes (Residual)	3,248	10.7	452.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	32,618	100.0	6,031.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,398	31.9	1,922.8
2	Malignant neoplasms (C00-C97)	8,347	25.6	1,543.5
3	Cerebrovascular diseases (I60-I69)	2,715	8.3	502.1
4	Diabetes mellitus (E10-E14)	1,635	5.0	302.3
5	Chronic lower respiratory diseases (J40-J47)	1,351	4.1	249.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	911	2.8	168.5
7	Septicemia (A40-A41)	792	2.4	146.5
8	Influenza and pneumonia (J10-J18)	659	2.0	121.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	528	1.6	97.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	499	1.5	92.3
11	Alzheimer's disease (G30)	358	1.1	66.2
12	Pneumonitis due to solids and liquids (J69)	225	0.7	41.6
13	Aortic aneurysm and dissection (I71)	164	0.5	30.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	162	0.5	30.0
15	Atherosclerosis (I70)	152	0.5	28.1
...	All other causes (Residual)	3,722	11.4	688.3
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	28,016	100.0	8,702.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,410	33.6	2,923.0
2	Malignant neoplasms (C00-C97)	5,845	20.9	1,815.6
3	Cerebrovascular diseases (I60-I69)	2,599	9.3	807.3
4	Diabetes mellitus (E10-E14)	1,360	4.9	422.5
5	Chronic lower respiratory diseases (J40-J47)	1,013	3.6	314.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	839	3.0	260.6
7	Influenza and pneumonia (J10-J18)	751	2.7	233.3
8	Septicemia (A40-A41)	721	2.6	224.0
9	Alzheimer's disease (G30)	525	1.9	163.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	478	1.7	148.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	426	1.5	132.3
12	Pneumonitis due to solids and liquids (J69)	244	0.9	75.8
13	Atherosclerosis (I70)	160	0.6	49.7
14	Aortic aneurysm and dissection (I71)	139	0.5	43.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	41.3
...	All other causes (Residual)	3,373	12.0	1,047.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,831	100.0	13,485.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,483	35.6	4,800.4
2	Malignant neoplasms (C00-C97)	3,905	16.4	2,209.8
3	Cerebrovascular diseases (I60-I69)	2,435	10.2	1,377.9
4	Diabetes mellitus (E10-E14)	970	4.1	548.9
5	Influenza and pneumonia (J10-J18)	869	3.6	491.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	727	3.1	411.4
7	Chronic lower respiratory diseases (J40-J47)	694	2.9	392.7
8	Septicemia (A40-A41)	673	2.8	380.8
9	Alzheimer's disease (G30)	567	2.4	320.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	416	1.7	235.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	349	1.5	197.5
12	Pneumonitis due to solids and liquids (J69)	233	1.0	131.8
13	Atherosclerosis (I70)	190	0.8	107.5
14	Aortic aneurysm and dissection (I71)	118	0.5	66.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.5	61.1
...	All other causes (Residual)	3,094	13.0	1,750.8
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	13,993	100.0	16,009.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,242	37.5	5,997.6
2	Malignant neoplasms (C00-C97)	1,703	12.2	1,948.5
3	Cerebrovascular diseases (I60-I69)	1,452	10.4	1,661.3
4	Influenza and pneumonia (J10-J18)	671	4.8	767.7
5	Diabetes mellitus (E10-E14)	494	3.5	565.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	2.8	454.2
7	Septicemia (A40-A41)	394	2.8	450.8
8	Alzheimer's disease (G30)	362	2.6	414.2
9	Chronic lower respiratory diseases (J40-J47)	337	2.4	385.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	266	1.9	304.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	1.4	225.4
12	Pneumonitis due to solids and liquids (J69)	176	1.3	201.4
13	Atherosclerosis (I70)	162	1.2	185.4
14	Nutritional deficiencies (E40-E64)	63	0.5	72.1
15	Anemias (D50-D64)	57	0.4	65.2
...	All other causes (Residual)	2,020	14.4	2,311.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,229	100.0	18,195.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,546	40.9	7,437.3
2	Cerebrovascular diseases (I60-I69)	614	9.9	1,793.6
3	Malignant neoplasms (C00-C97)	531	8.5	1,551.1
4	Influenza and pneumonia (J10-J18)	311	5.0	908.5
5	Alzheimer's disease (G30)	182	2.9	531.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.8	505.4
7	Septicemia (A40-A41)	162	2.6	473.2
8	Diabetes mellitus (E10-E14)	141	2.3	411.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	125	2.0	365.1
10	Chronic lower respiratory diseases (J40-J47)	110	1.8	321.3
11	Pneumonitis due to solids and liquids (J69)	95	1.5	277.5
12	Atherosclerosis (I70)	91	1.5	265.8
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	1.3	236.6
14	Nutritional deficiencies (E40-E64)	46	0.7	134.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	87.6
...	All other causes (Residual)	991	15.9	2,894.9
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	1,842	100.0	25,253.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	785	42.6	10,762.3
2	Cerebrovascular diseases (I60-I69)	143	7.8	1,960.5
3	Malignant neoplasms (C00-C97)	121	6.6	1,658.9
4	Influenza and pneumonia (J10-J18)	112	6.1	1,535.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.9	740.3
6	Septicemia (A40-A41)	52	2.8	712.9
7	Alzheimer's disease (G30)	48	2.6	658.1
8	Atherosclerosis (I70)	39	2.1	534.7
9	Diabetes mellitus (E10-E14)	37	2.0	507.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	1.9	479.8
11	Chronic lower respiratory diseases (J40-J47)	25	1.4	342.7
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.0	*
13	Pneumonitis due to solids and liquids (J69)	17	0.9	*
14	Nutritional deficiencies (E40-E64)	14	0.8	*
15	Anemias (D50-D64)	12	0.7	*
...	All other causes (Residual)	329	17.9	4,510.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	143,883	100.0	918.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,077	25.8	236.6
2	Malignant neoplasms (C00-C97)	32,542	22.6	207.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,629	6.0	55.1
4	Cerebrovascular diseases (I60-I69)	7,823	5.4	49.9
5	Assault (homicide) (X85-Y09,Y87.1)	6,131	4.3	39.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,377	3.7	34.3
7	Diabetes mellitus (E10-E14)	4,718	3.3	30.1
8	Chronic lower respiratory diseases (J40-J47)	4,452	3.1	28.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,984	2.1	19.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,818	2.0	18.0
11	Influenza and pneumonia (J10-J18)	2,786	1.9	17.8
12	Septicemia (A40-A41)	2,594	1.8	16.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,812	1.3	11.6
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,630	1.1	10.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,550	1.1	9.9
...	All other causes (Residual)	20,960	14.6	133.7
Black, non-Hispanic, male, 1-4 years				
...	All causes	734	100.0	69.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	35.4	24.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	10.8	7.5
3	Assault (homicide) (X85-Y09,Y87.1)	74	10.1	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.7	2.6
5	Influenza and pneumonia (J10-J18)	25	3.4	2.4
6	Malignant neoplasms (C00-C97)	24	3.3	2.3
7	Certain conditions originating in the perinatal period (P00-P96)	19	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.5	*
9	Anemias (D50-D64)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
13	Meningitis (G00,G03)	6	0.8	*
14	Cerebrovascular diseases (I60-I69)	4	0.5	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	3	0.4	*
...	All other causes (Residual)	158	21.5	15.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, male, 5-9 years			
...	All causes	508	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	50.0	16.9
2	Malignant neoplasms (C00-C97)	40	7.9	2.7
3	Assault (homicide) (X85-Y09,Y87.1)	34	6.7	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	4.7	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.5	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.8	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.6	*
8	Anemias (D50-D64)	7	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
10	Influenza and pneumonia (J10-J18)	5	1.0	*
11	Septicemia (A40-A41)	4	0.8	*
12	Meningitis (G00,G03)	3	0.6	*
13	Diseases of appendix (K35-K38)	2	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	76	15.0	5.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, male, 10-14 years			
...	All causes	573	100.0	38.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	37.2	14.3
2	Assault (homicide) (X85-Y09,Y87.1)	55	9.6	3.7
3	Malignant neoplasms (C00-C97)	54	9.4	3.6
4	Chronic lower respiratory diseases (J40-J47)	33	5.8	2.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	5.1	2.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.7	1.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	3.8	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.6	*
8	Anemias (D50-D64)	9	1.6	*
10	Cerebrovascular diseases (I60-I69)	6	1.0	*
11	Diabetes mellitus (E10-E14)	5	0.9	*
12	Septicemia (A40-A41)	3	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Influenza and pneumonia (J10-J18)	3	0.5	*
15	Meningococcal infection (A39)	2	0.3	*
15	Meningitis (G00,G03)	2	0.3	*
15	Diseases of appendix (K35-K38)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	94	16.4	6.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	2,133	100.0	145.2
1	Assault (homicide) (X85-Y09,Y87.1)	969	45.4	66.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	26.6	38.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	152	7.1	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	4.2	6.1
5	Malignant neoplasms (C00-C97)	84	3.9	5.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	1.2	1.8
7	Chronic lower respiratory diseases (J40-J47)	23	1.1	1.6
8	Anemias (D50-D64)	13	0.6	*
9	Legal intervention (Y35,Y89.0)	10	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
12	Cerebrovascular diseases (I60-I69)	7	0.3	*
13	Influenza and pneumonia (J10-J18)	6	0.3	*
14	Meningococcal infection (A39)	5	0.2	*
14	Meningitis (G00,G03)	5	0.2	*
...	All other causes (Residual)	159	7.5	10.8
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,153	100.0	249.7
1	Assault (homicide) (X85-Y09,Y87.1)	1,464	46.4	115.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	713	22.6	56.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	256	8.1	20.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	3.9	9.7
5	Malignant neoplasms (C00-C97)	97	3.1	7.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	44	1.4	3.5
7	Anemias (D50-D64)	37	1.2	2.9
8	Chronic lower respiratory diseases (J40-J47)	29	0.9	2.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.8	1.9
10	Legal intervention (Y35,Y89.0)	21	0.7	1.7
11	Influenza and pneumonia (J10-J18)	17	0.5	*
12	Diabetes mellitus (E10-E14)	15	0.5	*
13	Cerebrovascular diseases (I60-I69)	12	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Septicemia (A40-A41)	11	0.3	*
...	All other causes (Residual)	279	8.8	22.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,038	100.0	257.1
1	Assault (homicide) (X85-Y09,Y87.1)	1,070	35.2	90.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	647	21.3	54.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	268	8.8	22.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	229	7.5	19.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	6.3	16.2
6	Malignant neoplasms (C00-C97)	113	3.7	9.6
7	Chronic lower respiratory diseases (J40-J47)	34	1.1	2.9
8	Anemias (D50-D64)	29	1.0	2.5
9	Influenza and pneumonia (J10-J18)	25	0.8	2.1
10	Cerebrovascular diseases (I60-I69)	24	0.8	2.0
10	Legal intervention (Y35,Y89.0)	24	0.8	2.0
12	Diabetes mellitus (E10-E14)	21	0.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.6	*
14	Septicemia (A40-A41)	14	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.5	*
...	All other causes (Residual)	318	10.5	26.9
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,595	100.0	304.9
1	Assault (homicide) (X85-Y09,Y87.1)	701	19.5	59.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	18.5	56.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	651	18.1	55.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	355	9.9	30.1
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	203	5.6	17.2
6	Malignant neoplasms (C00-C97)	181	5.0	15.4
7	Diabetes mellitus (E10-E14)	64	1.8	5.4
8	Cerebrovascular diseases (I60-I69)	48	1.3	4.1
9	Anemias (D50-D64)	43	1.2	3.6
10	Influenza and pneumonia (J10-J18)	33	0.9	2.8
10	Chronic lower respiratory diseases (J40-J47)	33	0.9	2.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.8	2.5
13	Septicemia (A40-A41)	27	0.8	2.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	22	0.6	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.5	*
...	All other causes (Residual)	520	14.5	44.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	5,104	100.0	396.8
1	Human immunodeficiency virus (HIV) disease (B20-B24)	957	18.8	74.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	773	15.1	60.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	14.8	58.6
4	Assault (homicide) (X85-Y09,Y87.1)	548	10.7	42.6
5	Malignant neoplasms (C00-C97)	382	7.5	29.7
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	215	4.2	16.7
7	Cerebrovascular diseases (I60-I69)	123	2.4	9.6
8	Diabetes mellitus (E10-E14)	107	2.1	8.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	75	1.5	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.3	5.2
11	Influenza and pneumonia (J10-J18)	58	1.1	4.5
12	Septicemia (A40-A41)	55	1.1	4.3
13	Chronic lower respiratory diseases (J40-J47)	44	0.9	3.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.7	2.9
15	Anemias (D50-D64)	35	0.7	2.7
...	All other causes (Residual)	874	17.1	67.9
Black, non-Hispanic, male, 40-44 years				
...	All causes	7,280	100.0	596.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,392	19.1	114.0
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,200	16.5	98.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	874	12.0	71.6
4	Malignant neoplasms (C00-C97)	873	12.0	71.5
5	Assault (homicide) (X85-Y09,Y87.1)	432	5.9	35.4
6	Cerebrovascular diseases (I60-I69)	264	3.6	21.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	218	3.0	17.8
8	Diabetes mellitus (E10-E14)	169	2.3	13.8
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	138	1.9	11.3
10	Influenza and pneumonia (J10-J18)	115	1.6	9.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	1.5	8.7
12	Septicemia (A40-A41)	99	1.4	8.1
13	Chronic lower respiratory diseases (J40-J47)	77	1.1	6.3
14	Viral hepatitis (B15-B19)	64	0.9	5.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	60	0.8	4.9
...	All other causes (Residual)	1,199	16.5	98.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,266	100.0	948.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,024	21.8	207.3
2	Malignant neoplasms (C00-C97)	1,687	18.2	172.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	992	10.7	101.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	864	9.3	88.5
5	Cerebrovascular diseases (I60-I69)	409	4.4	41.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	357	3.9	36.6
7	Diabetes mellitus (E10-E14)	290	3.1	29.7
8	Assault (homicide) (X85-Y09,Y87.1)	285	3.1	29.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.8	16.7
10	Influenza and pneumonia (J10-J18)	140	1.5	14.3
11	Septicemia (A40-A41)	136	1.5	13.9
12	Chronic lower respiratory diseases (J40-J47)	126	1.4	12.9
13	Viral hepatitis (B15-B19)	125	1.3	12.8
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	114	1.2	11.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	83	0.9	8.5
...	All other causes (Residual)	1,471	15.9	150.6
Black, non-Hispanic, male, 50-54 years				
...	All causes	9,760	100.0	1,349.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,692	27.6	372.1
2	Malignant neoplasms (C00-C97)	2,411	24.7	333.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	577	5.9	79.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	560	5.7	77.4
5	Cerebrovascular diseases (I60-I69)	491	5.0	67.9
6	Diabetes mellitus (E10-E14)	351	3.6	48.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	292	3.0	40.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	1.9	26.3
9	Chronic lower respiratory diseases (J40-J47)	158	1.6	21.8
10	Assault (homicide) (X85-Y09,Y87.1)	146	1.5	20.2
11	Septicemia (A40-A41)	143	1.5	19.8
12	Influenza and pneumonia (J10-J18)	137	1.4	18.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	135	1.4	18.7
14	Viral hepatitis (B15-B19)	90	0.9	12.4
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	69	0.7	9.5
...	All other causes (Residual)	1,318	13.5	182.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	10,627	100.0	1,970.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,245	30.5	601.6
2	Malignant neoplasms (C00-C97)	3,117	29.3	577.9
3	Cerebrovascular diseases (I60-I69)	541	5.1	100.3
4	Diabetes mellitus (E10-E14)	457	4.3	84.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	3.9	77.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	331	3.1	61.4
7	Chronic lower respiratory diseases (J40-J47)	254	2.4	47.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	254	2.4	47.1
9	Septicemia (A40-A41)	173	1.6	32.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	1.6	31.7
11	Influenza and pneumonia (J10-J18)	125	1.2	23.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	1.1	21.1
13	Assault (homicide) (X85-Y09,Y87.1)	99	0.9	18.4
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	54	0.5	10.0
15	Aortic aneurysm and dissection (I71)	52	0.5	9.6
...	All other causes (Residual)	1,222	11.5	226.5
Black, non-Hispanic, male, 60-64 years				
...	All causes	11,866	100.0	2,748.4
1	Malignant neoplasms (C00-C97)	3,816	32.2	883.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,662	30.9	848.2
3	Cerebrovascular diseases (I60-I69)	694	5.8	160.7
4	Diabetes mellitus (E10-E14)	529	4.5	122.5
5	Chronic lower respiratory diseases (J40-J47)	370	3.1	85.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	2.7	75.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	2.1	59.1
8	Septicemia (A40-A41)	219	1.8	50.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	199	1.7	46.1
10	Influenza and pneumonia (J10-J18)	157	1.3	36.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	142	1.2	32.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	140	1.2	32.4
13	Assault (homicide) (X85-Y09,Y87.1)	60	0.5	13.9
14	Aortic aneurysm and dissection (I71)	44	0.4	10.2
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	41	0.3	9.5
...	All other causes (Residual)	1,214	10.2	281.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,720	100.0	3,567.2
1	Malignant neoplasms (C00-C97)	4,423	32.2	1,150.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,226	30.8	1,098.8
3	Cerebrovascular diseases (I60-I69)	782	5.7	203.3
4	Diabetes mellitus (E10-E14)	630	4.6	163.8
5	Chronic lower respiratory diseases (J40-J47)	533	3.9	138.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	310	2.3	80.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	297	2.2	77.2
8	Septicemia (A40-A41)	274	2.0	71.2
9	Influenza and pneumonia (J10-J18)	198	1.4	51.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	166	1.2	43.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	165	1.2	42.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	99	0.7	25.7
13	Aortic aneurysm and dissection (I71)	73	0.5	19.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.5	16.9
15	Pneumonitis due to solids and liquids (J69)	59	0.4	15.3
...	All other causes (Residual)	1,420	10.3	369.2
Black, non-Hispanic, male, 70-74 years				
...	All causes	15,539	100.0	5,236.8
1	Malignant neoplasms (C00-C97)	4,805	30.9	1,619.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,790	30.8	1,614.3
3	Cerebrovascular diseases (I60-I69)	1,075	6.9	362.3
4	Chronic lower respiratory diseases (J40-J47)	705	4.5	237.6
5	Diabetes mellitus (E10-E14)	632	4.1	213.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	2.5	128.4
7	Septicemia (A40-A41)	335	2.2	112.9
8	Influenza and pneumonia (J10-J18)	320	2.1	107.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	1.9	100.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	183	1.2	61.7
11	Pneumonitis due to solids and liquids (J69)	109	0.7	36.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	107	0.7	36.1
13	Alzheimer's disease (G30)	81	0.5	27.3
14	Aortic aneurysm and dissection (I71)	74	0.5	24.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.4	21.2
...	All other causes (Residual)	1,580	10.2	532.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	15,706	100.0	7,455.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,725	30.1	2,242.9
2	Malignant neoplasms (C00-C97)	4,603	29.3	2,185.0
3	Cerebrovascular diseases (I60-I69)	1,169	7.4	554.9
4	Chronic lower respiratory diseases (J40-J47)	810	5.2	384.5
5	Diabetes mellitus (E10-E14)	588	3.7	279.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	2.8	206.5
7	Septicemia (A40-A41)	343	2.2	162.8
8	Influenza and pneumonia (J10-J18)	336	2.1	159.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	1.8	133.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	204	1.3	96.8
11	Pneumonitis due to solids and liquids (J69)	135	0.9	64.1
12	Alzheimer's disease (G30)	130	0.8	61.7
13	Atherosclerosis (I70)	78	0.5	37.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	35.6
15	Parkinson's disease (G20-G21)	67	0.4	31.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.4	31.8
...	All other causes (Residual)	1,659	10.6	787.5
Black, non-Hispanic, male, 80-84 years				
...	All causes	11,846	100.0	10,546.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,691	31.2	3,286.1
2	Malignant neoplasms (C00-C97)	3,000	25.3	2,670.9
3	Cerebrovascular diseases (I60-I69)	931	7.9	828.9
4	Chronic lower respiratory diseases (J40-J47)	592	5.0	527.1
5	Diabetes mellitus (E10-E14)	428	3.6	381.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	368	3.1	327.6
7	Influenza and pneumonia (J10-J18)	352	3.0	313.4
8	Septicemia (A40-A41)	305	2.6	271.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	1.7	178.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	177	1.5	157.6
11	Alzheimer's disease (G30)	160	1.4	142.4
12	Pneumonitis due to solids and liquids (J69)	126	1.1	112.2
13	Atherosclerosis (I70)	64	0.5	57.0
14	Aortic aneurysm and dissection (I71)	62	0.5	55.2
15	Parkinson's disease (G20-G21)	56	0.5	49.9
...	All other causes (Residual)	1,333	11.3	1,186.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,736	100.0	15,711.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,831	32.4	5,091.6
2	Malignant neoplasms (C00-C97)	1,887	21.6	3,393.8
3	Cerebrovascular diseases (I60-I69)	735	8.4	1,321.9
4	Chronic lower respiratory diseases (J40-J47)	362	4.1	651.1
5	Influenza and pneumonia (J10-J18)	352	4.0	633.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	3.3	525.2
7	Diabetes mellitus (E10-E14)	278	3.2	500.0
8	Septicemia (A40-A41)	234	2.7	420.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.9	291.4
10	Alzheimer's disease (G30)	157	1.8	282.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	131	1.5	235.6
12	Pneumonitis due to solids and liquids (J69)	103	1.2	185.2
13	Atherosclerosis (I70)	68	0.8	122.3
14	Parkinson's disease (G20-G21)	47	0.5	84.5
15	Aortic aneurysm and dissection (I71)	44	0.5	79.1
...	All other causes (Residual)	1,053	12.1	1,893.9
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,134	100.0	17,053.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,433	34.7	5,911.5
2	Malignant neoplasms (C00-C97)	710	17.2	2,928.9
3	Cerebrovascular diseases (I60-I69)	363	8.8	1,497.5
4	Influenza and pneumonia (J10-J18)	224	5.4	924.1
5	Chronic lower respiratory diseases (J40-J47)	171	4.1	705.4
6	Diabetes mellitus (E10-E14)	121	2.9	499.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.8	470.3
8	Septicemia (A40-A41)	106	2.6	437.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	1.9	325.9
10	Alzheimer's disease (G30)	75	1.8	309.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	1.5	251.6
12	Pneumonitis due to solids and liquids (J69)	58	1.4	239.3
13	Atherosclerosis (I70)	39	0.9	160.9
14	Parkinson's disease (G20-G21)	22	0.5	90.8
15	Nutritional deficiencies (E40-E64)	19	0.5	*
...	All other causes (Residual)	539	13.0	2,223.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,451	100.0	17,446.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	38.1	6,649.0
2	Malignant neoplasms (C00-C97)	198	13.6	2,380.7
3	Cerebrovascular diseases (I60-I69)	110	7.6	1,322.6
4	Influenza and pneumonia (J10-J18)	81	5.6	973.9
5	Chronic lower respiratory diseases (J40-J47)	52	3.6	625.2
6	Septicemia (A40-A41)	37	2.5	444.9
7	Alzheimer's disease (G30)	31	2.1	372.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	2.1	360.7
8	Pneumonitis due to solids and liquids (J69)	30	2.1	360.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.1	360.7
11	Atherosclerosis (I70)	23	1.6	276.5
12	Diabetes mellitus (E10-E14)	22	1.5	264.5
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	1.5	264.5
14	Nutritional deficiencies (E40-E64)	13	0.9	*
15	Aortic aneurysm and dissection (I71)	8	0.6	*
...	All other causes (Residual)	211	14.5	2,537.0
Black, non-Hispanic, male, 100 years and over				
...	All causes	307	100.0	20,912.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	41.7	8,719.3
2	Malignant neoplasms (C00-C97)	33	10.7	2,248.0
3	Influenza and pneumonia (J10-J18)	19	6.2	*
4	Cerebrovascular diseases (I60-I69)	17	5.5	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	3.9	*
6	Atherosclerosis (I70)	10	3.3	*
6	Chronic lower respiratory diseases (J40-J47)	10	3.3	*
8	Alzheimer's disease (G30)	7	2.3	*
9	Diabetes mellitus (E10-E14)	5	1.6	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.6	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	1.3	*
12	Septicemia (A40-A41)	3	1.0	*
13	Nutritional deficiencies (E40-E64)	2	0.7	*
13	Pneumonitis due to solids and liquids (J69)	2	0.7	*
13	Hyperplasia of prostate (N40)	2	0.7	*
...	All other causes (Residual)	48	15.6	3,269.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	138,096	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,636	29.4	233.3
2	Malignant neoplasms (C00-C97)	28,867	20.9	165.7
3	Cerebrovascular diseases (I60-I69)	10,908	7.9	62.6
4	Diabetes mellitus (E10-E14)	7,112	5.2	40.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,904	2.8	22.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,675	2.7	21.1
7	Chronic lower respiratory diseases (J40-J47)	3,388	2.5	19.5
8	Septicemia (A40-A41)	3,178	2.3	18.2
9	Influenza and pneumonia (J10-J18)	3,008	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,374	1.7	13.6
11	Certain conditions originating in the perinatal period (P00-P96)	2,182	1.6	12.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,158	1.6	12.4
13	Alzheimer's disease (G30)	1,636	1.2	9.4
14	Assault (homicide) (X85-Y09,Y87.1)	1,405	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	981	0.7	5.6
...	All other causes (Residual)	22,684	16.4	130.2
Black, non-Hispanic, female, 1-4 years				
...	All causes	549	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	30.6	16.4
2	Assault (homicide) (X85-Y09,Y87.1)	77	14.0	7.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.6	5.7
4	Malignant neoplasms (C00-C97)	23	4.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.0	2.2
6	Influenza and pneumonia (J10-J18)	15	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	7	1.3	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	Cerebrovascular diseases (I60-I69)	6	1.1	*
13	Meningitis (G00,G03)	5	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	123	22.4	12.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 5-9 years			
...	All causes	361	100.0	24.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	38.2	9.5
2	Malignant neoplasms (C00-C97)	44	12.2	3.0
3	Assault (homicide) (X85-Y09,Y87.1)	31	8.6	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	6.9	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.9	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	12	3.3	*
7	Anemias (D50-D64)	10	2.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.7	*
10	Septicemia (A40-A41)	3	0.8	*
10	Influenza and pneumonia (J10-J18)	3	0.8	*
12	Cerebrovascular diseases (I60-I69)	2	0.6	*
12	Diseases of appendix (K35-K38)	2	0.6	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Diabetes mellitus (E10-E14)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	60	16.6	4.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 10-14 years			
...	All causes	329	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	29.2	6.7
2	Malignant neoplasms (C00-C97)	41	12.5	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	8.2	1.9
4	Assault (homicide) (X85-Y09,Y87.1)	25	7.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	5.5	*
6	Chronic lower respiratory diseases (J40-J47)	14	4.3	*
7	Anemias (D50-D64)	7	2.1	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.8	*
9	Cerebrovascular diseases (I60-I69)	6	1.8	*
11	Septicemia (A40-A41)	4	1.2	*
11	Influenza and pneumonia (J10-J18)	4	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.2	*
14	Diabetes mellitus (E10-E14)	3	0.9	*
15	Meningococcal infection (A39)	2	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
15	Pneumonitis due to solids and liquids (J69)	2	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	57	17.3	4.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999
 [Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	680	100.0	47.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	28.8	13.8
2	Assault (homicide) (X85-Y09,Y87.1)	148	21.8	10.4
3	Malignant neoplasms (C00-C97)	52	7.6	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	5.6	2.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	3.5	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	18	2.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.9	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	Anemias (D50-D64)	10	1.5	*
12	Diabetes mellitus (E10-E14)	8	1.2	*
12	Influenza and pneumonia (J10-J18)	8	1.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
15	Meningitis (G00,G03)	4	0.6	*
...	All other causes (Residual)	111	16.3	7.8
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,014	100.0	78.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	21.4	16.8
2	Assault (homicide) (X85-Y09,Y87.1)	171	16.9	13.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	8.5	6.7
4	Malignant neoplasms (C00-C97)	81	8.0	6.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	65	6.4	5.0
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	31	3.1	2.4
7	Pregnancy, childbirth and the puerperium (O00-099)	28	2.8	2.2
8	Anemias (D50-D64)	24	2.4	1.9
9	Influenza and pneumonia (J10-J18)	18	1.8	*
10	Septicemia (A40-A41)	16	1.6	*
10	Diabetes mellitus (E10-E14)	16	1.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.6	*
13	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
14	Cerebrovascular diseases (I60-I69)	10	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
...	All other causes (Residual)	212	20.9	16.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,352	100.0	104.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	14.3	15.0
2	Assault (homicide) (X85-Y09,Y87.1)	192	14.2	14.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	178	13.2	13.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	9.4	9.8
5	Malignant neoplasms (C00-C97)	114	8.4	8.8
6	Pregnancy, childbirth and the puerperium (O00-099)	37	2.7	2.9
7	Chronic lower respiratory diseases (J40-J47)	33	2.4	2.6
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	2.1	2.2
9	Diabetes mellitus (E10-E14)	27	2.0	2.1
10	Anemias (D50-D64)	26	1.9	2.0
11	Cerebrovascular diseases (I60-I69)	25	1.8	1.9
12	Septicemia (A40-A41)	23	1.7	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.7	1.8
14	Influenza and pneumonia (J10-J18)	22	1.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.0	*
...	All other causes (Residual)	288	21.3	22.3
Black, non-Hispanic, female, 30-34 years				
...	All causes	2,007	100.0	149.5
1	Human immunodeficiency virus (HIV) disease (B20-B24)	382	19.0	28.4
2	Malignant neoplasms (C00-C97)	252	12.6	18.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	12.0	17.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	11.5	17.2
5	Assault (homicide) (X85-Y09,Y87.1)	159	7.9	11.8
6	Cerebrovascular diseases (I60-I69)	52	2.6	3.9
7	Diabetes mellitus (E10-E14)	48	2.4	3.6
8	Anemias (D50-D64)	41	2.0	3.1
9	Chronic lower respiratory diseases (J40-J47)	40	2.0	3.0
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	36	1.8	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.6	2.5
11	Pregnancy, childbirth and the puerperium (O00-099)	33	1.6	2.5
13	Influenza and pneumonia (J10-J18)	27	1.3	2.0
14	Septicemia (A40-A41)	21	1.0	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.8	*
...	All other causes (Residual)	396	19.7	29.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	3,459	100.0	236.7
1	Malignant neoplasms (C00-C97)	667	19.3	45.6
2	Human immunodeficiency virus (HIV) disease (B20-B24)	501	14.5	34.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	473	13.7	32.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	8.9	21.1
5	Assault (homicide) (X85-Y09,Y87.1)	182	5.3	12.5
6	Cerebrovascular diseases (I60-I69)	144	4.2	9.9
7	Diabetes mellitus (E10-E14)	85	2.5	5.8
8	Influenza and pneumonia (J10-J18)	58	1.7	4.0
9	Septicemia (A40-A41)	54	1.6	3.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	54	1.6	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.5	3.5
12	Chronic lower respiratory diseases (J40-J47)	49	1.4	3.4
13	Anemias (D50-D64)	31	0.9	2.1
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	0.8	2.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	0.8	1.9
...	All other causes (Residual)	744	21.5	50.9
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,927	100.0	352.7
1	Malignant neoplasms (C00-C97)	1,127	22.9	80.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	868	17.6	62.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	496	10.1	35.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	6.3	22.0
5	Cerebrovascular diseases (I60-I69)	277	5.6	19.8
6	Assault (homicide) (X85-Y09,Y87.1)	143	2.9	10.2
7	Diabetes mellitus (E10-E14)	141	2.9	10.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	106	2.2	7.6
9	Chronic lower respiratory diseases (J40-J47)	88	1.8	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.7	6.2
11	Influenza and pneumonia (J10-J18)	74	1.5	5.3
12	Septicemia (A40-A41)	71	1.4	5.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	1.0	3.6
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	48	1.0	3.4
15	Anemias (D50-D64)	38	0.8	2.7
...	All other causes (Residual)	1,006	20.4	72.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	5,865	100.0	503.9
1	Malignant neoplasms (C00-C97)	1,591	27.1	136.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,215	20.7	104.4
3	Cerebrovascular diseases (I60-I69)	371	6.3	31.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	330	5.6	28.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	4.7	23.5
6	Diabetes mellitus (E10-E14)	223	3.8	19.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.6	12.9
8	Chronic lower respiratory diseases (J40-J47)	130	2.2	11.2
9	Septicemia (A40-A41)	115	2.0	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	1.8	9.1
11	Influenza and pneumonia (J10-J18)	80	1.4	6.9
12	Assault (homicide) (X85-Y09,Y87.1)	79	1.3	6.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	1.0	5.2
14	Anemias (D50-D64)	38	0.6	3.3
15	Viral hepatitis (B15-B19)	35	0.6	3.0
...	All other causes (Residual)	1,067	18.2	91.7
Black, non-Hispanic, female, 50-54 years				
...	All causes	6,486	100.0	726.1
1	Malignant neoplasms (C00-C97)	2,060	31.8	230.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,517	23.4	169.8
3	Cerebrovascular diseases (I60-I69)	391	6.0	43.8
4	Diabetes mellitus (E10-E14)	355	5.5	39.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	2.9	21.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	175	2.7	19.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	2.4	17.5
8	Chronic lower respiratory diseases (J40-J47)	142	2.2	15.9
9	Septicemia (A40-A41)	124	1.9	13.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.6	11.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	91	1.4	10.2
12	Influenza and pneumonia (J10-J18)	68	1.0	7.6
13	Assault (homicide) (X85-Y09,Y87.1)	47	0.7	5.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.4	3.2
15	Viral hepatitis (B15-B19)	27	0.4	3.0
...	All other causes (Residual)	1,010	15.6	113.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	7,549	100.0	1,084.2
1	Malignant neoplasms (C00-C97)	2,470	32.7	354.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,957	25.9	281.1
3	Diabetes mellitus (E10-E14)	491	6.5	70.5
4	Cerebrovascular diseases (I60-I69)	476	6.3	68.4
5	Chronic lower respiratory diseases (J40-J47)	218	2.9	31.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	2.7	29.7
7	Septicemia (A40-A41)	156	2.1	22.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	1.9	20.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	109	1.4	15.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	107	1.4	15.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	90	1.2	12.9
12	Influenza and pneumonia (J10-J18)	83	1.1	11.9
13	Anemias (D50-D64)	30	0.4	4.3
13	Aortic aneurysm and dissection (I71)	30	0.4	4.3
15	Viral hepatitis (B15-B19)	29	0.4	4.2
...	All other causes (Residual)	952	12.6	136.7
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,021	100.0	1,550.5
1	Malignant neoplasms (C00-C97)	2,919	32.4	501.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,482	27.5	426.6
3	Diabetes mellitus (E10-E14)	659	7.3	113.3
4	Cerebrovascular diseases (I60-I69)	558	6.2	95.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.7	42.3
6	Chronic lower respiratory diseases (J40-J47)	243	2.7	41.8
7	Septicemia (A40-A41)	168	1.9	28.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	1.8	27.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	143	1.6	24.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.4	22.2
11	Influenza and pneumonia (J10-J18)	121	1.3	20.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.6	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	7.7
14	Aortic aneurysm and dissection (I71)	26	0.3	4.5
15	Viral hepatitis (B15-B19)	24	0.3	4.1
...	All other causes (Residual)	1,045	11.6	179.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	11,497	100.0	2,231.3
1	Malignant neoplasms (C00-C97)	3,465	30.1	672.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,396	29.5	659.1
3	Diabetes mellitus (E10-E14)	808	7.0	156.8
4	Cerebrovascular diseases (I60-I69)	747	6.5	145.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	388	3.4	75.3
6	Chronic lower respiratory diseases (J40-J47)	366	3.2	71.0
7	Septicemia (A40-A41)	246	2.1	47.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	177	1.5	34.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.4	30.5
10	Influenza and pneumonia (J10-J18)	129	1.1	25.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	84	0.7	16.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.5	11.5
13	Aortic aneurysm and dissection (I71)	53	0.5	10.3
14	Viral hepatitis (B15-B19)	38	0.3	7.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.3	6.4
...	All other causes (Residual)	1,351	11.8	262.2
Black, non-Hispanic, female, 70-74 years				
...	All causes	14,811	100.0	3,516.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,607	31.1	1,093.8
2	Malignant neoplasms (C00-C97)	3,932	26.5	933.6
3	Cerebrovascular diseases (I60-I69)	1,188	8.0	282.1
4	Diabetes mellitus (E10-E14)	1,050	7.1	249.3
5	Chronic lower respiratory diseases (J40-J47)	486	3.3	115.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	475	3.2	112.8
7	Septicemia (A40-A41)	357	2.4	84.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	239	1.6	56.7
9	Influenza and pneumonia (J10-J18)	233	1.6	55.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	1.3	45.1
11	Alzheimer's disease (G30)	106	0.7	25.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	75	0.5	17.8
13	Aortic aneurysm and dissection (I71)	71	0.5	16.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	16.4
15	Pneumonitis due to solids and liquids (J69)	65	0.4	15.4
...	All other causes (Residual)	1,668	11.3	396.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,912	100.0	5,123.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,673	33.5	1,718.5
2	Malignant neoplasms (C00-C97)	3,744	22.1	1,134.2
3	Cerebrovascular diseases (I60-I69)	1,546	9.1	468.3
4	Diabetes mellitus (E10-E14)	1,047	6.2	317.2
5	Chronic lower respiratory diseases (J40-J47)	541	3.2	163.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	2.8	144.2
7	Septicemia (A40-A41)	449	2.7	136.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	324	1.9	98.2
9	Influenza and pneumonia (J10-J18)	323	1.9	97.8
10	Alzheimer's disease (G30)	228	1.3	69.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	1.3	65.7
12	Aortic aneurysm and dissection (I71)	102	0.6	30.9
13	Pneumonitis due to solids and liquids (J69)	90	0.5	27.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	26.4
15	Atherosclerosis (I70)	74	0.4	22.4
...	All other causes (Residual)	1,991	11.8	603.1
Black, non-Hispanic, female, 80-84 years				
...	All causes	16,170	100.0	7,714.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,719	35.4	2,728.4
2	Malignant neoplasms (C00-C97)	2,845	17.6	1,357.3
3	Cerebrovascular diseases (I60-I69)	1,668	10.3	795.8
4	Diabetes mellitus (E10-E14)	932	5.8	444.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	2.9	224.7
6	Chronic lower respiratory diseases (J40-J47)	421	2.6	200.9
7	Septicemia (A40-A41)	416	2.6	198.5
8	Influenza and pneumonia (J10-J18)	399	2.5	190.4
9	Alzheimer's disease (G30)	365	2.3	174.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	301	1.9	143.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	1.4	107.3
12	Pneumonitis due to solids and liquids (J69)	118	0.7	56.3
13	Atherosclerosis (I70)	96	0.6	45.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.5	40.1
15	Aortic aneurysm and dissection (I71)	77	0.5	36.7
...	All other causes (Residual)	2,033	12.6	969.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,095	100.0	12,463.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,652	37.4	4,666.6
2	Malignant neoplasms (C00-C97)	2,018	13.4	1,666.2
3	Cerebrovascular diseases (I60-I69)	1,700	11.3	1,403.6
4	Diabetes mellitus (E10-E14)	692	4.6	571.4
5	Influenza and pneumonia (J10-J18)	517	3.4	426.9
6	Septicemia (A40-A41)	439	2.9	362.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	2.9	359.2
8	Alzheimer's disease (G30)	410	2.7	338.5
9	Chronic lower respiratory diseases (J40-J47)	332	2.2	274.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	285	1.9	235.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.2	154.4
12	Pneumonitis due to solids and liquids (J69)	130	0.9	107.3
13	Atherosclerosis (I70)	122	0.8	100.7
14	Aortic aneurysm and dissection (I71)	74	0.5	61.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	57.8
...	All other causes (Residual)	2,032	13.5	1,677.7
Black, non-Hispanic, female, 90-94 years				
...	All causes	9,859	100.0	15,609.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,809	38.6	6,030.6
2	Cerebrovascular diseases (I60-I69)	1,089	11.0	1,724.2
3	Malignant neoplasms (C00-C97)	993	10.1	1,572.2
4	Influenza and pneumonia (J10-J18)	447	4.5	707.7
5	Diabetes mellitus (E10-E14)	373	3.8	590.6
6	Septicemia (A40-A41)	288	2.9	456.0
7	Alzheimer's disease (G30)	287	2.9	454.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	283	2.9	448.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	205	2.1	324.6
10	Chronic lower respiratory diseases (J40-J47)	166	1.7	262.8
11	Atherosclerosis (I70)	123	1.2	194.7
12	Pneumonitis due to solids and liquids (J69)	118	1.2	186.8
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	1.2	186.8
14	Anemias (D50-D64)	48	0.5	76.0
15	Nutritional deficiencies (E40-E64)	44	0.4	69.7
...	All other causes (Residual)	1,468	14.9	2,324.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	4,778	100.0	18,436.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,993	41.7	7,690.2
2	Cerebrovascular diseases (I60-I69)	504	10.5	1,944.7
3	Malignant neoplasms (C00-C97)	333	7.0	1,284.9
4	Influenza and pneumonia (J10-J18)	230	4.8	887.5
5	Alzheimer's disease (G30)	151	3.2	582.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	3.0	551.8
7	Septicemia (A40-A41)	125	2.6	482.3
8	Diabetes mellitus (E10-E14)	119	2.5	459.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	95	2.0	366.6
10	Atherosclerosis (I70)	68	1.4	262.4
11	Pneumonitis due to solids and liquids (J69)	65	1.4	250.8
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	1.2	227.7
13	Chronic lower respiratory diseases (J40-J47)	58	1.2	223.8
14	Nutritional deficiencies (E40-E64)	33	0.7	127.3
15	Anemias (D50-D64)	25	0.5	96.5
...	All other causes (Residual)	777	16.3	2,998.1
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,535	100.0	26,347.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	657	42.8	11,277.0
2	Cerebrovascular diseases (I60-I69)	126	8.2	2,162.7
3	Influenza and pneumonia (J10-J18)	93	6.1	1,596.3
4	Malignant neoplasms (C00-C97)	88	5.7	1,510.5
5	Septicemia (A40-A41)	49	3.2	841.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.7	720.9
7	Alzheimer's disease (G30)	41	2.7	703.7
8	Diabetes mellitus (E10-E14)	32	2.1	549.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	2.0	532.1
10	Atherosclerosis (I70)	29	1.9	497.8
11	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
11	Pneumonitis due to solids and liquids (J69)	15	1.0	*
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	0.9	*
14	Anemias (D50-D64)	12	0.8	*
14	Nutritional deficiencies (E40-E64)	12	0.8	*
...	All other causes (Residual)	279	18.2	4,788.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable