

Table LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	103,740	100.0	331.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,866	24.9	82.5
2	Malignant neoplasms (C00-C97)	20,233	19.5	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,650	8.3	27.6
4	Cerebrovascular diseases (I60-I69)	5,907	5.7	18.8
5	Diabetes mellitus (E10-E14)	5,182	5.0	16.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,972	2.9	9.5
7	Assault (homicide) (X85-Y09,Y87.1)	2,864	2.8	9.1
8	Chronic lower respiratory diseases (J40-J47)	2,859	2.8	9.1
9	Influenza and pneumonia (J10-J18)	2,246	2.2	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,153	2.1	6.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,958	1.9	6.2
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,695	1.6	5.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,584	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,498	1.4	4.8
15	Septicemia (A40-A41)	1,215	1.2	3.9
...	All other causes (Residual)	16,858	16.3	53.8
Hispanic, both sexes, 1-4 years				
...	All causes	882	100.0	32.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	38.7	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	11.5	3.7
3	Malignant neoplasms (C00-C97)	72	8.2	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	52	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.7	1.2
6	Certain conditions originating in the perinatal period (P00-P96)	18	2.0	*
7	Influenza and pneumonia (J10-J18)	17	1.9	*
8	Septicemia (A40-A41)	14	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
11	Meningitis (G00,G03)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	7	0.8	*
13	Anemias (D50-D64)	6	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
...	All other causes (Residual)	183	20.7	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,014	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	362	35.7	6.1
2	Malignant neoplasms (C00-C97)	176	17.4	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	89	8.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	5.6	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	3.6	0.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	30	3.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.7	*
9	Influenza and pneumonia (J10-J18)	10	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.9	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
13	Septicemia (A40-A41)	8	0.8	*
13	Diseases of appendix (K35-K38)	8	0.8	*
15	Pneumonitis due to solids and liquids (J69)	5	0.5	*
...	All other causes (Residual)	171	16.9	2.9
Hispanic, both sexes, 15-24 years				
...	All causes	4,505	100.0	82.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,946	43.2	35.6
2	Assault (homicide) (X85-Y09,Y87.1)	1,119	24.8	20.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	391	8.7	7.1
4	Malignant neoplasms (C00-C97)	259	5.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	2.6	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.2	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.7	0.6
8	Cerebrovascular diseases (I60-I69)	32	0.7	0.6
9	Influenza and pneumonia (J10-J18)	25	0.6	0.5
10	Pregnancy, childbirth and the puerperium (O00-099)	19	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
12	Diabetes mellitus (E10-E14)	14	0.3	*
13	Legal intervention (Y35,Y89.0)	13	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.3	*
15	Septicemia (A40-A41)	9	0.2	*
...	All other causes (Residual)	446	9.9	8.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	5,397	100.0	103.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,790	33.2	34.2
2	Assault (homicide) (X85-Y09,Y87.1)	801	14.8	15.3
3	Malignant neoplasms (C00-C97)	466	8.6	8.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	423	7.8	8.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	407	7.5	7.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	286	5.3	5.5
7	Cerebrovascular diseases (I60-I69)	102	1.9	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	80	1.5	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.0	1.0
10	Diabetes mellitus (E10-E14)	43	0.8	0.8
11	Chronic lower respiratory diseases (J40-J47)	31	0.6	0.6
12	Septicemia (A40-A41)	26	0.5	0.5
12	Pregnancy, childbirth and the puerperium (O00-099)	26	0.5	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	0.5
15	Influenza and pneumonia (J10-J18)	24	0.4	0.5
...	All other causes (Residual)	813	15.1	15.5
Hispanic, both sexes, 35-44 years				
...	All causes	7,798	100.0	164.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,607	20.6	33.8
2	Malignant neoplasms (C00-C97)	1,204	15.4	25.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	855	11.0	18.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	761	9.8	16.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	518	6.6	10.9
6	Assault (homicide) (X85-Y09,Y87.1)	432	5.5	9.1
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	369	4.7	7.8
8	Cerebrovascular diseases (I60-I69)	241	3.1	5.1
9	Diabetes mellitus (E10-E14)	157	2.0	3.3
10	Viral hepatitis (B15-B19)	104	1.3	2.2
11	Chronic lower respiratory diseases (J40-J47)	79	1.0	1.7
12	Septicemia (A40-A41)	74	0.9	1.6
12	Influenza and pneumonia (J10-J18)	74	0.9	1.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	0.8	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	0.4	0.7
...	All other causes (Residual)	1,227	15.7	25.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	9,876	100.0	338.8
1	Malignant neoplasms (C00-C97)	2,516	25.5	86.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,806	18.3	62.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	926	9.4	31.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	787	8.0	27.0
5	Cerebrovascular diseases (I60-I69)	464	4.7	15.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	455	4.6	15.6
7	Diabetes mellitus (E10-E14)	413	4.2	14.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	210	2.1	7.2
9	Viral hepatitis (B15-B19)	203	2.1	7.0
10	Assault (homicide) (X85-Y09,Y87.1)	197	2.0	6.8
11	Chronic lower respiratory diseases (J40-J47)	115	1.2	3.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.1	3.7
13	Influenza and pneumonia (J10-J18)	104	1.1	3.6
14	Septicemia (A40-A41)	92	0.9	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	0.6	2.2
...	All other causes (Residual)	1,417	14.3	48.6
Hispanic, both sexes, 55-64 years				
...	All causes	12,319	100.0	733.7
1	Malignant neoplasms (C00-C97)	3,686	29.9	219.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,155	25.6	187.9
3	Diabetes mellitus (E10-E14)	927	7.5	55.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	638	5.2	38.0
5	Cerebrovascular diseases (I60-I69)	617	5.0	36.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	3.8	27.8
7	Chronic lower respiratory diseases (J40-J47)	239	1.9	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	1.9	13.7
9	Influenza and pneumonia (J10-J18)	155	1.3	9.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	137	1.1	8.2
11	Septicemia (A40-A41)	131	1.1	7.8
12	Viral hepatitis (B15-B19)	127	1.0	7.6
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	112	0.9	6.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	82	0.7	4.9
15	Assault (homicide) (X85-Y09,Y87.1)	69	0.6	4.1
...	All other causes (Residual)	1,548	12.6	92.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	19,201	100.0	1,742.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,557	28.9	504.4
2	Malignant neoplasms (C00-C97)	5,394	28.1	489.6
3	Diabetes mellitus (E10-E14)	1,551	8.1	140.8
4	Cerebrovascular diseases (I60-I69)	1,211	6.3	109.9
5	Chronic lower respiratory diseases (J40-J47)	649	3.4	58.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	583	3.0	52.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	440	2.3	39.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	1.9	33.5
9	Influenza and pneumonia (J10-J18)	328	1.7	29.8
10	Septicemia (A40-A41)	265	1.4	24.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	0.8	14.3
12	Viral hepatitis (B15-B19)	129	0.7	11.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.6	9.9
14	Alzheimer's disease (G30)	100	0.5	9.1
15	Aortic aneurysm and dissection (I71)	95	0.5	8.6
...	All other causes (Residual)	2,263	11.8	205.4
Hispanic, both sexes, 75-84 years				
...	All causes	20,912	100.0	3,771.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,046	33.7	1,270.8
2	Malignant neoplasms (C00-C97)	4,449	21.3	802.4
3	Cerebrovascular diseases (I60-I69)	1,672	8.0	301.6
4	Diabetes mellitus (E10-E14)	1,369	6.5	246.9
5	Chronic lower respiratory diseases (J40-J47)	935	4.5	168.6
6	Influenza and pneumonia (J10-J18)	569	2.7	102.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	426	2.0	76.8
8	Alzheimer's disease (G30)	362	1.7	65.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	334	1.6	60.2
10	Septicemia (A40-A41)	296	1.4	53.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	285	1.4	51.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	232	1.1	41.8
13	Parkinson's disease (G20-G21)	184	0.9	33.2
14	Atherosclerosis (I70)	130	0.6	23.4
15	Pneumonitis due to solids and liquids (J69)	118	0.6	21.3
...	All other causes (Residual)	2,505	12.0	451.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	17,358	100.0	9,175.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,968	40.1	3,683.4
2	Malignant neoplasms (C00-C97)	1,995	11.5	1,054.6
3	Cerebrovascular diseases (I60-I69)	1,536	8.8	812.0
4	Influenza and pneumonia (J10-J18)	876	5.0	463.1
5	Chronic lower respiratory diseases (J40-J47)	764	4.4	403.9
6	Diabetes mellitus (E10-E14)	704	4.1	372.1
7	Alzheimer's disease (G30)	501	2.9	264.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	331	1.9	175.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	1.5	137.4
10	Septicemia (A40-A41)	257	1.5	135.9
11	Pneumonitis due to solids and liquids (J69)	186	1.1	98.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	185	1.1	97.8
13	Atherosclerosis (I70)	184	1.1	97.3
14	Parkinson's disease (G20-G21)	110	0.6	58.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.4	39.1
...	All other causes (Residual)	2,427	14.0	1,283.0
Hispanic, male, all ages				
...	All causes	57,991	100.0	367.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,554	23.4	86.0
2	Malignant neoplasms (C00-C97)	10,668	18.4	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,578	11.3	41.7
4	Cerebrovascular diseases (I60-I69)	2,809	4.8	17.8
5	Assault (homicide) (X85-Y09,Y87.1)	2,399	4.1	15.2
6	Diabetes mellitus (E10-E14)	2,335	4.0	14.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,143	3.7	13.6
8	Chronic lower respiratory diseases (J40-J47)	1,547	2.7	9.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,525	2.6	9.7
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,430	2.5	9.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,192	2.1	7.6
12	Influenza and pneumonia (J10-J18)	1,092	1.9	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	814	1.4	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	756	1.3	4.8
15	Septicemia (A40-A41)	605	1.0	3.8
...	All other causes (Residual)	8,544	14.7	54.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	481	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	41.0	14.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.1	4.5
3	Malignant neoplasms (C00-C97)	34	7.1	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	25	5.2	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.2	1.4
6	Influenza and pneumonia (J10-J18)	10	2.1	*
7	Septicemia (A40-A41)	8	1.7	*
8	Meningitis (G00,G03)	5	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
10	Anemias (D50-D64)	4	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	92	19.1	6.6
Hispanic, male, 5-14 years				
...	All causes	592	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	38.3	7.4
2	Malignant neoplasms (C00-C97)	92	15.5	3.0
3	Assault (homicide) (X85-Y09,Y87.1)	59	10.0	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	4.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.9	0.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	21	3.5	0.7
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.2	*
10	Cerebrovascular diseases (I60-I69)	6	1.0	*
10	Influenza and pneumonia (J10-J18)	6	1.0	*
12	Septicemia (A40-A41)	4	0.7	*
12	Diseases of appendix (K35-K38)	4	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Meningococcal infection (A39)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	84	14.2	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,547	100.0	124.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,575	44.4	55.5
2	Assault (homicide) (X85-Y09,Y87.1)	990	27.9	34.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	338	9.5	11.9
4	Malignant neoplasms (C00-C97)	158	4.5	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	2.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.8	1.1
7	Cerebrovascular diseases (I60-I69)	21	0.6	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.6	0.7
9	Legal intervention (Y35,Y89.0)	13	0.4	*
10	Influenza and pneumonia (J10-J18)	11	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
12	Diabetes mellitus (E10-E14)	8	0.2	*
14	Septicemia (A40-A41)	5	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.1	*
...	All other causes (Residual)	283	8.0	10.0
Hispanic, male, 25-34 years				
...	All causes	4,084	100.0	151.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,494	36.6	55.4
2	Assault (homicide) (X85-Y09,Y87.1)	698	17.1	25.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	364	8.9	13.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	314	7.7	11.7
5	Malignant neoplasms (C00-C97)	222	5.4	8.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	5.4	8.1
7	Cerebrovascular diseases (I60-I69)	62	1.5	2.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.5	2.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.6	0.9
10	Legal intervention (Y35,Y89.0)	23	0.6	0.9
11	Diabetes mellitus (E10-E14)	21	0.5	0.8
12	Influenza and pneumonia (J10-J18)	15	0.4	*
13	Septicemia (A40-A41)	14	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
...	All other causes (Residual)	524	12.8	19.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,476	100.0	226.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,307	23.9	53.9
2	Human immunodeficiency virus (HIV) disease (B20-B24)	666	12.2	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	572	10.4	23.6
4	Malignant neoplasms (C00-C97)	504	9.2	20.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	421	7.7	17.4
6	Assault (homicide) (X85-Y09,Y87.1)	354	6.5	14.6
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	307	5.6	12.7
8	Cerebrovascular diseases (I60-I69)	146	2.7	6.0
9	Viral hepatitis (B15-B19)	84	1.5	3.5
9	Diabetes mellitus (E10-E14)	84	1.5	3.5
11	Influenza and pneumonia (J10-J18)	54	1.0	2.2
12	Septicemia (A40-A41)	51	0.9	2.1
13	Chronic lower respiratory diseases (J40-J47)	36	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.6	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.9
...	All other causes (Residual)	838	15.3	34.6
Hispanic, male, 45-54 years				
...	All causes	6,497	100.0	456.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,348	20.7	94.7
2	Malignant neoplasms (C00-C97)	1,237	19.0	86.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	734	11.3	51.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	647	10.0	45.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	354	5.4	24.9
6	Cerebrovascular diseases (I60-I69)	285	4.4	20.0
7	Diabetes mellitus (E10-E14)	252	3.9	17.7
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	169	2.6	11.9
9	Viral hepatitis (B15-B19)	153	2.4	10.7
10	Assault (homicide) (X85-Y09,Y87.1)	150	2.3	10.5
11	Influenza and pneumonia (J10-J18)	71	1.1	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.0	4.5
13	Septicemia (A40-A41)	56	0.9	3.9
14	Chronic lower respiratory diseases (J40-J47)	52	0.8	3.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	0.5	2.5
...	All other causes (Residual)	890	13.7	62.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	7,490	100.0	962.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,091	27.9	268.7
2	Malignant neoplasms (C00-C97)	2,060	27.5	264.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	481	6.4	61.8
4	Diabetes mellitus (E10-E14)	480	6.4	61.7
5	Cerebrovascular diseases (I60-I69)	354	4.7	45.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	334	4.5	42.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	1.8	17.6
8	Chronic lower respiratory diseases (J40-J47)	135	1.8	17.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	118	1.6	15.2
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	94	1.3	12.1
11	Influenza and pneumonia (J10-J18)	80	1.1	10.3
12	Septicemia (A40-A41)	78	1.0	10.0
13	Viral hepatitis (B15-B19)	75	1.0	9.6
14	Assault (homicide) (X85-Y09,Y87.1)	56	0.7	7.2
15	Aortic aneurysm and dissection (I71)	41	0.5	5.3
...	All other causes (Residual)	876	11.7	112.6
Hispanic, male, 65-74 years				
...	All causes	10,783	100.0	2,219.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,355	31.1	690.6
2	Malignant neoplasms (C00-C97)	3,045	28.2	626.7
3	Diabetes mellitus (E10-E14)	709	6.6	145.9
4	Cerebrovascular diseases (I60-I69)	641	5.9	131.9
5	Chronic lower respiratory diseases (J40-J47)	373	3.5	76.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	360	3.3	74.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	2.7	59.3
8	Influenza and pneumonia (J10-J18)	187	1.7	38.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	178	1.7	36.6
10	Septicemia (A40-A41)	134	1.2	27.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	84	0.8	17.3
12	Aortic aneurysm and dissection (I71)	67	0.6	13.8
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	63	0.6	13.0
14	Viral hepatitis (B15-B19)	59	0.5	12.1
15	Parkinson's disease (G20-G21)	52	0.5	10.7
...	All other causes (Residual)	1,188	11.0	244.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	10,317	100.0	4,525.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,483	33.8	1,527.7
2	Malignant neoplasms (C00-C97)	2,403	23.3	1,054.0
3	Cerebrovascular diseases (I60-I69)	758	7.3	332.5
4	Diabetes mellitus (E10-E14)	562	5.4	246.5
5	Chronic lower respiratory diseases (J40-J47)	523	5.1	229.4
6	Influenza and pneumonia (J10-J18)	284	2.8	124.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.1	93.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.8	79.8
9	Septicemia (A40-A41)	144	1.4	63.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	136	1.3	59.7
11	Alzheimer's disease (G30)	128	1.2	56.1
12	Parkinson's disease (G20-G21)	113	1.1	49.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	99	1.0	43.4
14	Aortic aneurysm and dissection (I71)	65	0.6	28.5
14	Pneumonitis due to solids and liquids (J69)	65	0.6	28.5
...	All other causes (Residual)	1,160	11.2	508.8
Hispanic, male, 85 years and over				
...	All causes	6,253	100.0	9,842.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,324	37.2	3,658.0
2	Malignant neoplasms (C00-C97)	902	14.4	1,419.8
3	Cerebrovascular diseases (I60-I69)	527	8.4	829.5
4	Chronic lower respiratory diseases (J40-J47)	388	6.2	610.7
5	Influenza and pneumonia (J10-J18)	334	5.3	525.7
6	Diabetes mellitus (E10-E14)	217	3.5	341.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	2.5	245.5
8	Alzheimer's disease (G30)	129	2.1	203.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	1.9	188.9
10	Septicemia (A40-A41)	84	1.3	132.2
11	Pneumonitis due to solids and liquids (J69)	74	1.2	116.5
12	Atherosclerosis (I70)	70	1.1	110.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	1.0	96.0
14	Parkinson's disease (G20-G21)	49	0.8	77.1
15	Aortic aneurysm and dissection (I71)	37	0.6	58.2
...	All other causes (Residual)	781	12.5	1,229.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	45,749	100.0	293.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,312	26.9	79.0
2	Malignant neoplasms (C00-C97)	9,565	20.9	61.4
3	Cerebrovascular diseases (I60-I69)	3,098	6.8	19.9
4	Diabetes mellitus (E10-E14)	2,847	6.2	18.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,072	4.5	13.3
6	Chronic lower respiratory diseases (J40-J47)	1,312	2.9	8.4
7	Influenza and pneumonia (J10-J18)	1,154	2.5	7.4
8	Certain conditions originating in the perinatal period (P00-P96)	961	2.1	6.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	829	1.8	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	770	1.7	4.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	742	1.6	4.8
12	Alzheimer's disease (G30)	667	1.5	4.3
13	Septicemia (A40-A41)	610	1.3	3.9
14	Assault (homicide) (X85-Y09,Y87.1)	465	1.0	3.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	433	0.9	2.8
...	All other causes (Residual)	7,912	17.3	50.8
Hispanic, female, 1-4 years				
...	All causes	401	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	35.9	10.7
2	Malignant neoplasms (C00-C97)	38	9.5	2.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	9.5	2.8
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.2	*
5	Certain conditions originating in the perinatal period (P00-P96)	13	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.0	*
8	Influenza and pneumonia (J10-J18)	7	1.7	*
9	Septicemia (A40-A41)	6	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Meningitis (G00,G03)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	83	20.7	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	422	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	32.0	4.6
2	Malignant neoplasms (C00-C97)	84	19.9	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	30	7.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	6.6	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.1	*
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	9	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J10-J18)	4	0.9	*
10	Pneumonitis due to solids and liquids (J69)	4	0.9	*
10	Diseases of appendix (K35-K38)	4	0.9	*
14	Cerebrovascular diseases (I60-I69)	3	0.7	*
15	Anemias (D50-D64)	2	0.5	*
15	Diabetes mellitus (E10-E14)	2	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	76	18.0	2.6
Hispanic, female, 15-24 years				
...	All causes	958	100.0	36.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	38.7	14.1
2	Assault (homicide) (X85-Y09,Y87.1)	129	13.5	4.9
3	Malignant neoplasms (C00-C97)	101	10.5	3.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	53	5.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.5	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	2.7	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	19	2.0	*
8	Influenza and pneumonia (J10-J18)	14	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Diabetes mellitus (E10-E14)	6	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
13	Septicemia (A40-A41)	4	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
13	Meningitis (G00,G03)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	150	15.7	5.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,313	100.0	51.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	22.5	11.7
2	Malignant neoplasms (C00-C97)	244	18.6	9.6
3	Assault (homicide) (X85-Y09,Y87.1)	103	7.8	4.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	93	7.1	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	5.1	2.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	59	4.5	2.3
7	Cerebrovascular diseases (I60-I69)	40	3.0	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	2.2	1.1
9	Pregnancy, childbirth and the puerperium (O00-099)	26	2.0	1.0
10	Diabetes mellitus (E10-E14)	22	1.7	0.9
11	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.4	*
13	Septicemia (A40-A41)	12	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.8	*
...	All other causes (Residual)	264	20.1	10.4
Hispanic, female, 35-44 years				
...	All causes	2,322	100.0	99.9
1	Malignant neoplasms (C00-C97)	700	30.1	30.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	12.9	12.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	189	8.1	8.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	8.1	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	97	4.2	4.2
6	Cerebrovascular diseases (I60-I69)	95	4.1	4.1
7	Assault (homicide) (X85-Y09,Y87.1)	78	3.4	3.4
8	Diabetes mellitus (E10-E14)	73	3.1	3.1
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	62	2.7	2.7
10	Chronic lower respiratory diseases (J40-J47)	43	1.9	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.3	1.3
12	Pregnancy, childbirth and the puerperium (O00-099)	25	1.1	1.1
13	Septicemia (A40-A41)	23	1.0	1.0
14	Viral hepatitis (B15-B19)	20	0.9	0.9
14	Influenza and pneumonia (J10-J18)	20	0.9	0.9
...	All other causes (Residual)	377	16.2	16.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	3,379	100.0	226.7
1	Malignant neoplasms (C00-C97)	1,279	37.9	85.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	458	13.6	30.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	5.7	12.9
4	Cerebrovascular diseases (I60-I69)	179	5.3	12.0
5	Diabetes mellitus (E10-E14)	161	4.8	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	140	4.1	9.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	101	3.0	6.8
8	Chronic lower respiratory diseases (J40-J47)	63	1.9	4.2
9	Viral hepatitis (B15-B19)	50	1.5	3.4
10	Assault (homicide) (X85-Y09,Y87.1)	47	1.4	3.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.3	2.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	41	1.2	2.8
13	Septicemia (A40-A41)	36	1.1	2.4
14	Influenza and pneumonia (J10-J18)	33	1.0	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.9	1.9
...	All other causes (Residual)	527	15.6	35.4
Hispanic, female, 55-64 years				
...	All causes	4,829	100.0	536.0
1	Malignant neoplasms (C00-C97)	1,626	33.7	180.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,064	22.0	118.1
3	Diabetes mellitus (E10-E14)	447	9.3	49.6
4	Cerebrovascular diseases (I60-I69)	263	5.4	29.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	157	3.3	17.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	2.7	14.7
7	Chronic lower respiratory diseases (J40-J47)	104	2.2	11.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.9	10.3
9	Influenza and pneumonia (J10-J18)	75	1.6	8.3
10	Septicemia (A40-A41)	53	1.1	5.9
11	Viral hepatitis (B15-B19)	52	1.1	5.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	42	0.9	4.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.6	3.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.4	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	18	0.4	*
...	All other causes (Residual)	636	13.2	70.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	8,418	100.0	1,366.8
1	Malignant neoplasms (C00-C97)	2,349	27.9	381.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,202	26.2	357.5
3	Diabetes mellitus (E10-E14)	842	10.0	136.7
4	Cerebrovascular diseases (I60-I69)	570	6.8	92.6
5	Chronic lower respiratory diseases (J40-J47)	276	3.3	44.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	223	2.6	36.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	2.3	31.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	1.8	24.7
9	Influenza and pneumonia (J10-J18)	141	1.7	22.9
10	Septicemia (A40-A41)	131	1.6	21.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	0.9	12.0
12	Viral hepatitis (B15-B19)	70	0.8	11.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.7	10.1
14	Alzheimer's disease (G30)	49	0.6	8.0
15	Parkinson's disease (G20-G21)	32	0.4	5.2
...	All other causes (Residual)	1,054	12.5	171.1
Hispanic, female, 75-84 years				
...	All causes	10,595	100.0	3,245.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,563	33.6	1,091.5
2	Malignant neoplasms (C00-C97)	2,046	19.3	626.8
3	Cerebrovascular diseases (I60-I69)	914	8.6	280.0
4	Diabetes mellitus (E10-E14)	807	7.6	247.2
5	Chronic lower respiratory diseases (J40-J47)	412	3.9	126.2
6	Influenza and pneumonia (J10-J18)	285	2.7	87.3
7	Alzheimer's disease (G30)	234	2.2	71.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	2.0	65.6
9	Septicemia (A40-A41)	152	1.4	46.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	1.4	46.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.4	45.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	133	1.3	40.7
13	Parkinson's disease (G20-G21)	71	0.7	21.7
14	Atherosclerosis (I70)	70	0.7	21.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.6	19.6
...	All other causes (Residual)	1,329	12.5	407.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	11,105	100.0	8,838.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,644	41.8	3,696.2
2	Malignant neoplasms (C00-C97)	1,093	9.8	869.9
3	Cerebrovascular diseases (I60-I69)	1,009	9.1	803.1
4	Influenza and pneumonia (J10-J18)	542	4.9	431.4
5	Diabetes mellitus (E10-E14)	487	4.4	387.6
6	Chronic lower respiratory diseases (J40-J47)	376	3.4	299.3
7	Alzheimer's disease (G30)	372	3.3	296.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.6	139.3
9	Septicemia (A40-A41)	173	1.6	137.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	1.3	111.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	124	1.1	98.7
12	Atherosclerosis (I70)	114	1.0	90.7
13	Pneumonitis due to solids and liquids (J69)	112	1.0	89.1
14	Parkinson's disease (G20-G21)	61	0.5	48.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	40.6
...	All other causes (Residual)	1,632	14.7	1,298.9
White, non-Hispanic, both sexes, all ages				
...	All causes	1,953,197	100.0	996.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	607,704	31.1	310.0
2	Malignant neoplasms (C00-C97)	456,315	23.4	232.8
3	Cerebrovascular diseases (I60-I69)	138,716	7.1	70.8
4	Chronic lower respiratory diseases (J40-J47)	111,620	5.7	56.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,326	3.8	37.4
6	Influenza and pneumonia (J10-J18)	54,280	2.8	27.7
7	Diabetes mellitus (E10-E14)	49,347	2.5	25.2
8	Alzheimer's disease (G30)	40,835	2.1	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26,418	1.4	13.5
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,534	1.3	12.5
11	Septicemia (A40-A41)	23,023	1.2	11.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,602	1.0	10.0
13	Aortic aneurysm and dissection (I71)	13,934	0.7	7.1
14	Parkinson's disease (G20-G21)	13,475	0.7	6.9
15	Atherosclerosis (I70)	13,286	0.7	6.8
...	All other causes (Residual)	286,782	14.7	146.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,819	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	37.0	11.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	284	10.1	3.0
3	Malignant neoplasms (C00-C97)	281	10.0	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	156	5.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	3.3	1.0
6	Influenza and pneumonia (J10-J18)	58	2.1	0.6
7	Septicemia (A40-A41)	47	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	37	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	27	1.0	0.3
11	Meningitis (G00,G03)	26	0.9	0.3
12	Meningococcal infection (A39)	24	0.9	0.3
13	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	658	23.3	6.9
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,488	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,908	42.5	7.4
2	Malignant neoplasms (C00-C97)	608	13.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	5.4	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	176	3.9	0.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	174	3.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	3.5	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	1.5	0.3
8	Influenza and pneumonia (J10-J18)	62	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	55	1.2	0.2
10	Septicemia (A40-A41)	47	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	40	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	31	0.7	0.1
13	Diabetes mellitus (E10-E14)	19	0.4	*
14	Meningitis (G00,G03)	18	0.4	*
15	Anemias (D50-D64)	17	0.4	*
...	All other causes (Residual)	867	19.3	3.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	17,868	100.0	71.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,410	52.7	37.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,830	15.8	11.3
3	Malignant neoplasms (C00-C97)	1,070	6.0	4.3
4	Assault (homicide) (X85-Y09,Y87.1)	962	5.4	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	585	3.3	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	277	1.6	1.1
7	Chronic lower respiratory diseases (J40-J47)	110	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	105	0.6	0.4
9	Influenza and pneumonia (J10-J18)	95	0.5	0.4
10	Diabetes mellitus (E10-E14)	94	0.5	0.4
11	Septicemia (A40-A41)	71	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.2	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.2	0.1
15	Anemias (D50-D64)	28	0.2	0.1
...	All other causes (Residual)	2,103	11.8	8.4
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,986	100.0	93.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,855	32.7	30.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,927	16.4	15.3
3	Malignant neoplasms (C00-C97)	2,682	11.2	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,755	7.3	6.8
5	Assault (homicide) (X85-Y09,Y87.1)	1,135	4.7	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	794	3.3	3.1
7	Diabetes mellitus (E10-E14)	363	1.5	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	334	1.4	1.3
9	Cerebrovascular diseases (I60-I69)	301	1.3	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	243	1.0	0.9
11	Influenza and pneumonia (J10-J18)	199	0.8	0.8
12	Septicemia (A40-A41)	155	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	146	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.4	0.4
...	All other causes (Residual)	3,882	16.2	15.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	57,630	100.0	176.6
1	Malignant neoplasms (C00-C97)	11,848	20.6	36.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,844	18.8	33.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,936	15.5	27.4
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,469	9.5	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,148	3.7	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,082	3.6	6.4
7	Cerebrovascular diseases (I60-I69)	1,406	2.4	4.3
8	Assault (homicide) (X85-Y09,Y87.1)	1,327	2.3	4.1
9	Diabetes mellitus (E10-E14)	1,231	2.1	3.8
10	Influenza and pneumonia (J10-J18)	637	1.1	2.0
11	Chronic lower respiratory diseases (J40-J47)	533	0.9	1.6
12	Viral hepatitis (B15-B19)	464	0.8	1.4
13	Septicemia (A40-A41)	439	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	426	0.7	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	319	0.6	1.0
...	All other causes (Residual)	9,521	16.5	29.2
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	106,808	100.0	387.0
1	Malignant neoplasms (C00-C97)	34,944	32.7	126.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,823	23.2	89.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,392	7.9	30.4
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,487	4.2	16.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,430	4.1	16.1
6	Cerebrovascular diseases (I60-I69)	3,143	2.9	11.4
7	Diabetes mellitus (E10-E14)	2,940	2.8	10.7
8	Chronic lower respiratory diseases (J40-J47)	2,372	2.2	8.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,286	1.2	4.7
10	Influenza and pneumonia (J10-J18)	1,109	1.0	4.0
11	Septicemia (A40-A41)	1,015	1.0	3.7
12	Viral hepatitis (B15-B19)	958	0.9	3.5
13	Assault (homicide) (X85-Y09,Y87.1)	814	0.8	2.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	717	0.7	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	514	0.5	1.9
...	All other causes (Residual)	14,864	13.9	53.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	180,793	100.0	972.1
1	Malignant neoplasms (C00-C97)	70,788	39.2	380.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,950	26.5	257.8
3	Chronic lower respiratory diseases (J40-J47)	9,786	5.4	52.6
4	Cerebrovascular diseases (I60-I69)	6,311	3.5	33.9
5	Diabetes mellitus (E10-E14)	5,674	3.1	30.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,455	3.0	29.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,117	2.3	22.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,580	1.4	13.9
9	Influenza and pneumonia (J10-J18)	1,894	1.0	10.2
10	Septicemia (A40-A41)	1,792	1.0	9.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,636	0.9	8.8
12	Aortic aneurysm and dissection (I71)	1,195	0.7	6.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	899	0.5	4.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	686	0.4	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	473	0.3	2.5
...	All other causes (Residual)	19,557	10.8	105.2
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	367,490	100.0	2,458.9
1	Malignant neoplasms (C00-C97)	127,119	34.6	850.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,750	28.2	694.2
3	Chronic lower respiratory diseases (J40-J47)	29,400	8.0	196.7
4	Cerebrovascular diseases (I60-I69)	18,332	5.0	122.7
5	Diabetes mellitus (E10-E14)	11,664	3.2	78.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,568	1.8	43.9
7	Influenza and pneumonia (J10-J18)	5,481	1.5	36.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,718	1.3	31.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	4,482	1.2	30.0
10	Septicemia (A40-A41)	4,140	1.1	27.7
11	Aortic aneurysm and dissection (I71)	3,548	1.0	23.7
12	Alzheimer's disease (G30)	2,821	0.8	18.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,276	0.6	15.2
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,249	0.6	15.0
15	Parkinson's disease (G20-G21)	1,807	0.5	12.1
...	All other causes (Residual)	39,135	10.6	261.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	604,344	100.0	5,783.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	195,187	32.3	1,867.8
2	Malignant neoplasms (C00-C97)	141,295	23.4	1,352.1
3	Cerebrovascular diseases (I60-I69)	49,178	8.1	470.6
4	Chronic lower respiratory diseases (J40-J47)	44,662	7.4	427.4
5	Diabetes mellitus (E10-E14)	16,821	2.8	161.0
6	Influenza and pneumonia (J10-J18)	16,785	2.8	160.6
7	Alzheimer's disease (G30)	14,453	2.4	138.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,757	1.8	102.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,504	1.6	90.9
10	Septicemia (A40-A41)	7,717	1.3	73.8
11	Parkinson's disease (G20-G21)	6,594	1.1	63.1
12	Aortic aneurysm and dissection (I71)	5,407	0.9	51.7
13	Pneumonitis due to solids and liquids (J69)	4,361	0.7	41.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,001	0.7	38.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,940	0.7	37.7
...	All other causes (Residual)	73,682	12.2	705.1
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	573,327	100.0	15,945.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	224,193	39.1	6,235.4
2	Malignant neoplasms (C00-C97)	65,636	11.4	1,825.5
3	Cerebrovascular diseases (I60-I69)	59,818	10.4	1,663.7
4	Influenza and pneumonia (J10-J18)	27,838	4.9	774.2
5	Chronic lower respiratory diseases (J40-J47)	24,521	4.3	682.0
6	Alzheimer's disease (G30)	23,106	4.0	642.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,647	1.9	296.1
8	Diabetes mellitus (E10-E14)	10,533	1.8	293.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,268	1.6	257.8
10	Atherosclerosis (I70)	7,617	1.3	211.8
11	Septicemia (A40-A41)	7,479	1.3	208.0
12	Pneumonitis due to solids and liquids (J69)	6,508	1.1	181.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,145	0.9	143.1
14	Parkinson's disease (G20-G21)	4,822	0.8	134.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,133	0.5	87.1
...	All other causes (Residual)	83,063	14.5	2,310.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	944,913	100.0	984.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	293,206	31.0	305.5
2	Malignant neoplasms (C00-C97)	236,384	25.0	246.3
3	Chronic lower respiratory diseases (J40-J47)	55,314	5.9	57.6
4	Cerebrovascular diseases (I60-I69)	51,968	5.5	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,113	4.9	48.1
6	Diabetes mellitus (E10-E14)	23,177	2.5	24.2
7	Influenza and pneumonia (J10-J18)	23,097	2.4	24.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,620	2.1	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,793	1.4	13.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	12,613	1.3	13.1
11	Alzheimer's disease (G30)	12,253	1.3	12.8
12	Septicemia (A40-A41)	9,863	1.0	10.3
13	Aortic aneurysm and dissection (I71)	8,551	0.9	8.9
14	Parkinson's disease (G20-G21)	7,648	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	6,642	0.7	6.9
...	All other causes (Residual)	125,671	13.3	131.0
White, non-Hispanic, male, 1-4 years				
...	All causes	1,605	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	38.4	12.7
2	Malignant neoplasms (C00-C97)	160	10.0	3.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	9.6	3.2
4	Assault (homicide) (X85-Y09,Y87.1)	83	5.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	1.1
6	Septicemia (A40-A41)	25	1.6	0.5
7	Certain conditions originating in the perinatal period (P00-P96)	24	1.5	0.5
8	Influenza and pneumonia (J10-J18)	22	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Meningococcal infection (A39)	15	0.9	*
10	Meningitis (G00,G03)	15	0.9	*
10	Cerebrovascular diseases (I60-I69)	15	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
...	All other causes (Residual)	377	23.5	7.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,643	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	45.6	9.2
2	Malignant neoplasms (C00-C97)	345	13.1	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	141	5.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	4.5	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	93	3.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.2	0.6
7	Influenza and pneumonia (J10-J18)	36	1.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	36	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.2	0.2
10	Septicemia (A40-A41)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	21	0.8	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
13	Meningitis (G00,G03)	11	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.4	*
15	Diabetes mellitus (E10-E14)	9	0.3	*
...	All other causes (Residual)	462	17.5	3.5
White, non-Hispanic, male, 15-24 years				
...	All causes	12,677	100.0	98.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,749	53.2	52.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,416	19.1	18.8
3	Assault (homicide) (X85-Y09,Y87.1)	644	5.1	5.0
4	Malignant neoplasms (C00-C97)	629	5.0	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	357	2.8	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	1.4	1.4
7	Chronic lower respiratory diseases (J40-J47)	59	0.5	0.5
8	Diabetes mellitus (E10-E14)	52	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	51	0.4	0.4
9	Influenza and pneumonia (J10-J18)	51	0.4	0.4
11	Septicemia (A40-A41)	44	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.2	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.2	0.2
14	Legal intervention (Y35,Y89.0)	20	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,361	10.7	10.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,449	100.0	128.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,932	36.1	46.3
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,205	19.5	25.0
3	Malignant neoplasms (C00-C97)	1,341	8.2	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,152	7.0	9.0
5	Assault (homicide) (X85-Y09,Y87.1)	753	4.6	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	643	3.9	5.0
7	Diabetes mellitus (E10-E14)	203	1.2	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	1.1	1.4
9	Cerebrovascular diseases (I60-I69)	151	0.9	1.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	151	0.9	1.2
11	Influenza and pneumonia (J10-J18)	119	0.7	0.9
12	Septicemia (A40-A41)	87	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	66	0.4	0.5
14	Legal intervention (Y35,Y89.0)	62	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.4	0.5
...	All other causes (Residual)	2,346	14.3	18.3
White, non-Hispanic, male, 35-44 years				
...	All causes	37,291	100.0	228.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,949	21.3	48.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,590	17.7	40.3
3	Malignant neoplasms (C00-C97)	5,425	14.5	33.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,210	11.3	25.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,768	4.7	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,478	4.0	9.0
7	Assault (homicide) (X85-Y09,Y87.1)	881	2.4	5.4
8	Diabetes mellitus (E10-E14)	747	2.0	4.6
9	Cerebrovascular diseases (I60-I69)	712	1.9	4.4
10	Influenza and pneumonia (J10-J18)	334	0.9	2.0
11	Viral hepatitis (B15-B19)	333	0.9	2.0
12	Septicemia (A40-A41)	243	0.7	1.5
13	Chronic lower respiratory diseases (J40-J47)	239	0.6	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	0.5	1.2
...	All other causes (Residual)	5,964	16.0	36.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	67,151	100.0	491.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,747	27.9	137.2
2	Malignant neoplasms (C00-C97)	17,588	26.2	128.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,142	9.1	45.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,388	5.0	24.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,365	5.0	24.6
6	Diabetes mellitus (E10-E14)	1,801	2.7	13.2
7	Cerebrovascular diseases (I60-I69)	1,662	2.5	12.2
8	Chronic lower respiratory diseases (J40-J47)	1,174	1.7	8.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,136	1.7	8.3
10	Viral hepatitis (B15-B19)	717	1.1	5.2
11	Influenza and pneumonia (J10-J18)	670	1.0	4.9
12	Assault (homicide) (X85-Y09,Y87.1)	579	0.9	4.2
13	Septicemia (A40-A41)	551	0.8	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	0.6	3.0
15	Aortic aneurysm and dissection (I71)	340	0.5	2.5
...	All other causes (Residual)	8,885	13.2	65.0
White, non-Hispanic, male, 55-64 years				
...	All causes	108,692	100.0	1,208.0
1	Malignant neoplasms (C00-C97)	38,658	35.6	429.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,466	30.8	371.9
3	Chronic lower respiratory diseases (J40-J47)	5,091	4.7	56.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,730	3.4	41.5
5	Cerebrovascular diseases (I60-I69)	3,479	3.2	38.7
6	Diabetes mellitus (E10-E14)	3,200	2.9	35.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,885	2.7	32.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,001	1.8	22.2
9	Influenza and pneumonia (J10-J18)	1,049	1.0	11.7
10	Aortic aneurysm and dissection (I71)	928	0.9	10.3
11	Septicemia (A40-A41)	904	0.8	10.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	903	0.8	10.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	536	0.5	6.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	410	0.4	4.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	331	0.3	3.7
...	All other causes (Residual)	11,121	10.2	123.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	209,155	100.0	3,077.6
1	Malignant neoplasms (C00-C97)	70,518	33.7	1,037.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,699	30.9	952.0
3	Chronic lower respiratory diseases (J40-J47)	15,559	7.4	228.9
4	Cerebrovascular diseases (I60-I69)	9,337	4.5	137.4
5	Diabetes mellitus (E10-E14)	6,066	2.9	89.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,962	1.9	58.3
7	Influenza and pneumonia (J10-J18)	3,087	1.5	45.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,746	1.3	40.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,532	1.2	37.3
10	Aortic aneurysm and dissection (I71)	2,451	1.2	36.1
11	Septicemia (A40-A41)	2,111	1.0	31.1
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,869	0.9	27.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,326	0.6	19.5
14	Alzheimer's disease (G30)	1,263	0.6	18.6
15	Parkinson's disease (G20-G21)	1,182	0.6	17.4
...	All other causes (Residual)	20,447	9.8	300.9
White, non-Hispanic, male, 75-84 years				
...	All causes	296,528	100.0	7,057.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,422	33.2	2,342.5
2	Malignant neoplasms (C00-C97)	73,535	24.8	1,750.2
3	Chronic lower respiratory diseases (J40-J47)	22,377	7.5	532.6
4	Cerebrovascular diseases (I60-I69)	20,618	7.0	490.7
5	Influenza and pneumonia (J10-J18)	8,255	2.8	196.5
6	Diabetes mellitus (E10-E14)	7,722	2.6	183.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,551	1.9	132.1
8	Alzheimer's disease (G30)	5,432	1.8	129.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,907	1.7	116.8
10	Parkinson's disease (G20-G21)	4,005	1.4	95.3
11	Septicemia (A40-A41)	3,538	1.2	84.2
12	Aortic aneurysm and dissection (I71)	3,243	1.1	77.2
13	Pneumonitis due to solids and liquids (J69)	2,492	0.8	59.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,013	0.7	47.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,740	0.6	41.4
...	All other causes (Residual)	32,678	11.0	777.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	184,938	100.0	17,539.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,476	37.6	6,589.0
2	Malignant neoplasms (C00-C97)	28,161	15.2	2,670.7
3	Cerebrovascular diseases (I60-I69)	15,886	8.6	1,506.6
4	Chronic lower respiratory diseases (J40-J47)	10,695	5.8	1,014.3
5	Influenza and pneumonia (J10-J18)	9,410	5.1	892.4
6	Alzheimer's disease (G30)	5,330	2.9	505.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,006	2.2	379.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,721	2.0	352.9
9	Diabetes mellitus (E10-E14)	3,374	1.8	320.0
10	Pneumonitis due to solids and liquids (J69)	2,767	1.5	262.4
11	Parkinson's disease (G20-G21)	2,287	1.2	216.9
12	Septicemia (A40-A41)	2,257	1.2	214.0
13	Atherosclerosis (I70)	2,014	1.1	191.0
14	Aortic aneurysm and dissection (I71)	1,364	0.7	129.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,240	0.7	117.6
...	All other causes (Residual)	22,950	12.4	2,176.5
White, non-Hispanic, female, all ages				
...	All causes	1,008,284	100.0	1,007.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314,498	31.2	314.2
2	Malignant neoplasms (C00-C97)	219,931	21.8	219.7
3	Cerebrovascular diseases (I60-I69)	86,748	8.6	86.7
4	Chronic lower respiratory diseases (J40-J47)	56,306	5.6	56.3
5	Influenza and pneumonia (J10-J18)	31,183	3.1	31.2
6	Alzheimer's disease (G30)	28,582	2.8	28.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,213	2.7	27.2
8	Diabetes mellitus (E10-E14)	26,170	2.6	26.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,625	1.4	13.6
10	Septicemia (A40-A41)	13,160	1.3	13.1
11	Atherosclerosis (I70)	8,559	0.8	8.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,653	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	6,989	0.7	7.0
14	Pneumonitis due to solids and liquids (J69)	6,497	0.6	6.5
15	Parkinson's disease (G20-G21)	5,827	0.6	5.8
...	All other causes (Residual)	155,343	15.4	155.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,214	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	35.0	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	10.7	2.8
3	Malignant neoplasms (C00-C97)	121	10.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	73	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J10-J18)	36	3.0	0.8
7	Septicemia (A40-A41)	22	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Meningitis (G00,G03)	11	0.9	*
13	Meningococcal infection (A39)	9	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
...	All other causes (Residual)	281	23.1	6.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,845	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	38.1	5.6
2	Malignant neoplasms (C00-C97)	263	14.3	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	6.8	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	83	4.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	3.9	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.8	0.3
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	1.8	0.3
8	Influenza and pneumonia (J10-J18)	26	1.4	0.2
9	Septicemia (A40-A41)	23	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	19	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	19	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.9	*
13	Anemias (D50-D64)	11	0.6	*
14	Diabetes mellitus (E10-E14)	10	0.5	*
15	Meningococcal infection (A39)	8	0.4	*
...	All other causes (Residual)	400	21.7	3.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,191	100.0	42.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,661	51.3	21.9
2	Malignant neoplasms (C00-C97)	441	8.5	3.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	414	8.0	3.4
4	Assault (homicide) (X85-Y09,Y87.1)	318	6.1	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	4.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	2.0	0.8
7	Cerebrovascular diseases (I60-I69)	54	1.0	0.4
8	Chronic lower respiratory diseases (J40-J47)	51	1.0	0.4
9	Influenza and pneumonia (J10-J18)	44	0.8	0.4
10	Diabetes mellitus (E10-E14)	42	0.8	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	0.2
12	Septicemia (A40-A41)	27	0.5	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	719	13.9	5.9
White, non-Hispanic, female, 25-34 years				
...	All causes	7,537	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,923	25.5	15.0
2	Malignant neoplasms (C00-C97)	1,341	17.8	10.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	722	9.6	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	603	8.0	4.7
5	Assault (homicide) (X85-Y09,Y87.1)	382	5.1	3.0
6	Diabetes mellitus (E10-E14)	160	2.1	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	2.1	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	151	2.0	1.2
9	Cerebrovascular diseases (I60-I69)	150	2.0	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.2	0.7
11	Influenza and pneumonia (J10-J18)	80	1.1	0.6
11	Chronic lower respiratory diseases (J40-J47)	80	1.1	0.6
13	Pregnancy, childbirth and the puerperium (O00-099)	74	1.0	0.6
14	Septicemia (A40-A41)	68	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.7	0.4
...	All other causes (Residual)	1,499	19.9	11.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	20,339	100.0	124.8
1	Malignant neoplasms (C00-C97)	6,423	31.6	39.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,895	14.2	17.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,346	11.5	14.4
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,259	6.2	7.7
5	Cerebrovascular diseases (I60-I69)	694	3.4	4.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	670	3.3	4.1
7	Diabetes mellitus (E10-E14)	484	2.4	3.0
8	Assault (homicide) (X85-Y09,Y87.1)	446	2.2	2.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	314	1.5	1.9
10	Influenza and pneumonia (J10-J18)	303	1.5	1.9
11	Chronic lower respiratory diseases (J40-J47)	294	1.4	1.8
12	Septicemia (A40-A41)	196	1.0	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	1.0	1.2
14	Viral hepatitis (B15-B19)	131	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	0.6	0.8
...	All other causes (Residual)	3,557	17.5	21.8
White, non-Hispanic, female, 45-54 years				
...	All causes	39,657	100.0	284.5
1	Malignant neoplasms (C00-C97)	17,356	43.8	124.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,076	15.3	43.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,250	5.7	16.1
4	Cerebrovascular diseases (I60-I69)	1,481	3.7	10.6
5	Chronic lower respiratory diseases (J40-J47)	1,198	3.0	8.6
6	Diabetes mellitus (E10-E14)	1,139	2.9	8.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,099	2.8	7.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	2.7	7.6
9	Septicemia (A40-A41)	464	1.2	3.3
10	Influenza and pneumonia (J10-J18)	439	1.1	3.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	0.8	2.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	243	0.6	1.7
13	Viral hepatitis (B15-B19)	241	0.6	1.7
14	Assault (homicide) (X85-Y09,Y87.1)	235	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	218	0.5	1.6
...	All other causes (Residual)	5,842	14.7	41.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	72,101	100.0	751.0
1	Malignant neoplasms (C00-C97)	32,130	44.6	334.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,484	20.1	150.9
3	Chronic lower respiratory diseases (J40-J47)	4,695	6.5	48.9
4	Cerebrovascular diseases (I60-I69)	2,832	3.9	29.5
5	Diabetes mellitus (E10-E14)	2,474	3.4	25.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,725	2.4	18.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,232	1.7	12.8
8	Septicemia (A40-A41)	888	1.2	9.2
9	Influenza and pneumonia (J10-J18)	845	1.2	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	733	1.0	7.6
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	579	0.8	6.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	363	0.5	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	276	0.4	2.9
14	Aortic aneurysm and dissection (I71)	267	0.4	2.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	0.3	2.3
...	All other causes (Residual)	8,353	11.6	87.0
White, non-Hispanic, female, 65-74 years				
...	All causes	158,335	100.0	1,942.9
1	Malignant neoplasms (C00-C97)	56,601	35.7	694.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,051	24.7	479.2
3	Chronic lower respiratory diseases (J40-J47)	13,841	8.7	169.8
4	Cerebrovascular diseases (I60-I69)	8,995	5.7	110.4
5	Diabetes mellitus (E10-E14)	5,598	3.5	68.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,606	1.6	32.0
7	Influenza and pneumonia (J10-J18)	2,394	1.5	29.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,186	1.4	26.8
9	Septicemia (A40-A41)	2,029	1.3	24.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,736	1.1	21.3
11	Alzheimer's disease (G30)	1,558	1.0	19.1
12	Aortic aneurysm and dissection (I71)	1,097	0.7	13.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	950	0.6	11.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	891	0.6	10.9
15	Parkinson's disease (G20-G21)	625	0.4	7.7
...	All other causes (Residual)	18,177	11.5	223.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-84 years				
...	All causes	307,816	100.0	4,926.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,765	31.4	1,548.6
2	Malignant neoplasms (C00-C97)	67,760	22.0	1,084.4
3	Cerebrovascular diseases (I60-I69)	28,560	9.3	457.1
4	Chronic lower respiratory diseases (J40-J47)	22,285	7.2	356.6
5	Diabetes mellitus (E10-E14)	9,099	3.0	145.6
6	Alzheimer's disease (G30)	9,021	2.9	144.4
7	Influenza and pneumonia (J10-J18)	8,530	2.8	136.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,206	1.7	83.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,597	1.5	73.6
10	Septicemia (A40-A41)	4,179	1.4	66.9
11	Parkinson's disease (G20-G21)	2,589	0.8	41.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,414	0.8	38.6
13	Atherosclerosis (I70)	2,190	0.7	35.0
14	Aortic aneurysm and dissection (I71)	2,164	0.7	34.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,988	0.6	31.8
...	All other causes (Residual)	40,469	13.1	647.6
White, non-Hispanic, female, 85 years and over				
...	All causes	388,389	100.0	15,284.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	154,717	39.8	6,088.7
2	Cerebrovascular diseases (I60-I69)	43,932	11.3	1,728.9
3	Malignant neoplasms (C00-C97)	37,475	9.6	1,474.8
4	Influenza and pneumonia (J10-J18)	18,428	4.7	725.2
5	Alzheimer's disease (G30)	17,776	4.6	699.6
6	Chronic lower respiratory diseases (J40-J47)	13,826	3.6	544.1
7	Diabetes mellitus (E10-E14)	7,159	1.8	281.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,641	1.7	261.3
9	Atherosclerosis (I70)	5,603	1.4	220.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,547	1.4	218.3
11	Septicemia (A40-A41)	5,222	1.3	205.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,905	1.0	153.7
13	Pneumonitis due to solids and liquids (J69)	3,741	1.0	147.2
14	Parkinson's disease (G20-G21)	2,535	0.7	99.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,924	0.5	75.7
...	All other causes (Residual)	59,958	15.4	2,359.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	281,979	100.0	852.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,713	27.6	234.8
2	Malignant neoplasms (C00-C97)	61,409	21.8	185.6
3	Cerebrovascular diseases (I60-I69)	18,731	6.6	56.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,533	4.4	37.9
5	Diabetes mellitus (E10-E14)	11,830	4.2	35.7
6	Chronic lower respiratory diseases (J40-J47)	7,840	2.8	23.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,751	2.7	23.4
8	Assault (homicide) (X85-Y09,Y87.1)	7,536	2.7	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,659	2.4	20.1
10	Influenza and pneumonia (J10-J18)	5,794	2.1	17.5
11	Septicemia (A40-A41)	5,772	2.0	17.4
12	Certain conditions originating in the perinatal period (P00-P96)	5,000	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,708	1.3	11.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,793	1.0	8.4
15	Alzheimer's disease (G30)	2,325	0.8	7.0
...	All other causes (Residual)	44,585	15.8	134.7
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,283	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	33.4	20.6
2	Assault (homicide) (X85-Y09,Y87.1)	151	11.8	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	10.7	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.8	2.4
5	Malignant neoplasms (C00-C97)	47	3.7	2.3
6	Influenza and pneumonia (J10-J18)	40	3.1	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	29	2.3	1.4
8	Anemias (D50-D64)	20	1.6	1.0
9	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
11	Septicemia (A40-A41)	16	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
13	Meningitis (G00,G03)	11	0.9	*
14	Cerebrovascular diseases (I60-I69)	10	0.8	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
...	All other causes (Residual)	289	22.5	13.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,771	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	701	39.6	11.9
2	Malignant neoplasms (C00-C97)	179	10.1	3.0
3	Assault (homicide) (X85-Y09,Y87.1)	145	8.2	2.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	5.9	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	4.3	1.3
6	Chronic lower respiratory diseases (J40-J47)	66	3.7	1.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.3	0.7
8	Anemias (D50-D64)	33	1.9	0.6
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	1.6	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.0	*
11	Cerebrovascular diseases (I60-I69)	15	0.8	*
11	Influenza and pneumonia (J10-J18)	15	0.8	*
13	Septicemia (A40-A41)	14	0.8	*
14	Diabetes mellitus (E10-E14)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
...	All other causes (Residual)	317	17.9	5.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,980	100.0	128.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,752	39.4	50.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,694	24.3	31.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	463	6.6	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	4.8	6.2
5	Malignant neoplasms (C00-C97)	314	4.5	5.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	130	1.9	2.4
7	Anemias (D50-D64)	84	1.2	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	1.5
9	Chronic lower respiratory diseases (J40-J47)	78	1.1	1.4
10	Influenza and pneumonia (J10-J18)	49	0.7	0.9
11	Pregnancy, childbirth and the puerperium (O00-099)	46	0.7	0.8
12	Diabetes mellitus (E10-E14)	43	0.6	0.8
13	Cerebrovascular diseases (I60-I69)	40	0.6	0.7
14	Septicemia (A40-A41)	32	0.5	0.6
15	Legal intervention (Y35,Y89.0)	31	0.4	0.6
...	All other causes (Residual)	806	11.5	14.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	9,992	100.0	200.0
1	Assault (homicide) (X85-Y09,Y87.1)	2,122	21.2	42.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,738	17.4	34.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,479	14.8	29.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	913	9.1	18.3
5	Malignant neoplasms (C00-C97)	660	6.6	13.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	497	5.0	9.9
7	Diabetes mellitus (E10-E14)	160	1.6	3.2
8	Cerebrovascular diseases (I60-I69)	149	1.5	3.0
9	Chronic lower respiratory diseases (J40-J47)	140	1.4	2.8
10	Anemias (D50-D64)	139	1.4	2.8
11	Influenza and pneumonia (J10-J18)	107	1.1	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.0	2.1
13	Septicemia (A40-A41)	85	0.9	1.7
14	Pregnancy, childbirth and the puerperium (O00-099)	70	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	0.6	1.2
...	All other causes (Residual)	1,570	15.7	31.4
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	20,770	100.0	387.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,506	16.9	65.3
2	Human immunodeficiency virus (HIV) disease (B20-B24)	3,154	15.2	58.8
3	Malignant neoplasms (C00-C97)	3,049	14.7	56.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,245	10.8	41.8
5	Assault (homicide) (X85-Y09,Y87.1)	1,305	6.3	24.3
6	Cerebrovascular diseases (I60-I69)	808	3.9	15.1
7	Diabetes mellitus (E10-E14)	502	2.4	9.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	453	2.2	8.4
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	430	2.1	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	310	1.5	5.8
11	Influenza and pneumonia (J10-J18)	305	1.5	5.7
12	Septicemia (A40-A41)	279	1.3	5.2
13	Chronic lower respiratory diseases (J40-J47)	258	1.2	4.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	175	0.8	3.3
15	Anemias (D50-D64)	142	0.7	2.6
...	All other causes (Residual)	3,849	18.5	71.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	31,377	100.0	835.2
1	Malignant neoplasms (C00-C97)	7,749	24.7	206.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,448	23.7	198.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,057	6.6	54.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,903	6.1	50.7
5	Cerebrovascular diseases (I60-I69)	1,662	5.3	44.2
6	Diabetes mellitus (E10-E14)	1,219	3.9	32.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	905	2.9	24.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	615	2.0	16.4
9	Assault (homicide) (X85-Y09,Y87.1)	557	1.8	14.8
10	Chronic lower respiratory diseases (J40-J47)	556	1.8	14.8
11	Septicemia (A40-A41)	518	1.7	13.8
12	Influenza and pneumonia (J10-J18)	425	1.4	11.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	370	1.2	9.8
14	Viral hepatitis (B15-B19)	277	0.9	7.4
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	226	0.7	6.0
...	All other causes (Residual)	4,890	15.6	130.2
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	39,063	100.0	1,736.7
1	Malignant neoplasms (C00-C97)	12,322	31.5	547.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,346	29.0	504.4
3	Cerebrovascular diseases (I60-I69)	2,269	5.8	100.9
4	Diabetes mellitus (E10-E14)	2,136	5.5	95.0
5	Chronic lower respiratory diseases (J40-J47)	1,085	2.8	48.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,047	2.7	46.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	879	2.3	39.1
8	Septicemia (A40-A41)	716	1.8	31.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	691	1.8	30.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	613	1.6	27.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	506	1.3	22.5
12	Influenza and pneumonia (J10-J18)	486	1.2	21.6
13	Assault (homicide) (X85-Y09,Y87.1)	194	0.5	8.6
14	Aortic aneurysm and dissection (I71)	152	0.4	6.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	148	0.4	6.6
...	All other causes (Residual)	4,473	11.5	198.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	55,567	100.0	3,434.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,019	30.6	1,052.0
2	Malignant neoplasms (C00-C97)	16,625	29.9	1,027.6
3	Cerebrovascular diseases (I60-I69)	3,792	6.8	234.4
4	Diabetes mellitus (E10-E14)	3,120	5.6	192.9
5	Chronic lower respiratory diseases (J40-J47)	2,090	3.8	129.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,554	2.8	96.1
7	Septicemia (A40-A41)	1,212	2.2	74.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	1.7	58.3
9	Influenza and pneumonia (J10-J18)	880	1.6	54.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	764	1.4	47.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	432	0.8	26.7
12	Aortic aneurysm and dissection (I71)	271	0.5	16.8
13	Pneumonitis due to solids and liquids (J69)	264	0.5	16.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	256	0.5	15.8
15	Alzheimer's disease (G30)	249	0.4	15.4
...	All other causes (Residual)	6,096	11.0	376.8
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	60,634	100.0	7,028.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,808	32.7	2,296.0
2	Malignant neoplasms (C00-C97)	14,192	23.4	1,645.1
3	Cerebrovascular diseases (I60-I69)	5,314	8.8	616.0
4	Diabetes mellitus (E10-E14)	2,995	4.9	347.2
5	Chronic lower respiratory diseases (J40-J47)	2,364	3.9	274.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,750	2.9	202.9
7	Septicemia (A40-A41)	1,513	2.5	175.4
8	Influenza and pneumonia (J10-J18)	1,410	2.3	163.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,006	1.7	116.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	925	1.5	107.2
11	Alzheimer's disease (G30)	883	1.5	102.4
12	Pneumonitis due to solids and liquids (J69)	469	0.8	54.4
13	Atherosclerosis (I70)	312	0.5	36.2
14	Aortic aneurysm and dissection (I71)	303	0.5	35.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	295	0.5	34.2
...	All other causes (Residual)	7,095	11.7	822.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	45,895	100.0	15,015.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,056	37.2	5,580.3
2	Malignant neoplasms (C00-C97)	6,260	13.6	2,048.1
3	Cerebrovascular diseases (I60-I69)	4,644	10.1	1,519.4
4	Influenza and pneumonia (J10-J18)	1,963	4.3	642.2
5	Diabetes mellitus (E10-E14)	1,642	3.6	537.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,351	2.9	442.0
7	Septicemia (A40-A41)	1,281	2.8	419.1
8	Chronic lower respiratory diseases (J40-J47)	1,166	2.5	381.5
9	Alzheimer's disease (G30)	1,159	2.5	379.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	842	1.8	275.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	646	1.4	211.4
12	Pneumonitis due to solids and liquids (J69)	521	1.1	170.5
13	Atherosclerosis (I70)	482	1.1	157.7
14	Nutritional deficiencies (E40-E64)	219	0.5	71.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	194	0.4	63.5
...	All other causes (Residual)	6,469	14.1	2,116.5
Black, non-Hispanic, male, all ages				
...	All causes	143,883	100.0	918.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,077	25.8	236.6
2	Malignant neoplasms (C00-C97)	32,542	22.6	207.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,629	6.0	55.1
4	Cerebrovascular diseases (I60-I69)	7,823	5.4	49.9
5	Assault (homicide) (X85-Y09,Y87.1)	6,131	4.3	39.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,377	3.7	34.3
7	Diabetes mellitus (E10-E14)	4,718	3.3	30.1
8	Chronic lower respiratory diseases (J40-J47)	4,452	3.1	28.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,984	2.1	19.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,818	2.0	18.0
11	Influenza and pneumonia (J10-J18)	2,786	1.9	17.8
12	Septicemia (A40-A41)	2,594	1.8	16.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,812	1.3	11.6
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,630	1.1	10.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,550	1.1	9.9
...	All other causes (Residual)	20,960	14.6	133.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	734	100.0	69.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	35.4	24.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	10.8	7.5
3	Assault (homicide) (X85-Y09,Y87.1)	74	10.1	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.7	2.6
5	Influenza and pneumonia (J10-J18)	25	3.4	2.4
6	Malignant neoplasms (C00-C97)	24	3.3	2.3
7	Certain conditions originating in the perinatal period (P00-P96)	19	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.5	*
9	Anemias (D50-D64)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
13	Meningitis (G00,G03)	6	0.8	*
14	Cerebrovascular diseases (I60-I69)	4	0.5	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	3	0.4	*
...	All other causes (Residual)	158	21.5	15.0
Black, non-Hispanic, male, 5-14 years				
...	All causes	1,081	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	43.2	15.6
2	Malignant neoplasms (C00-C97)	94	8.7	3.1
3	Assault (homicide) (X85-Y09,Y87.1)	89	8.2	3.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	4.9	1.8
5	Chronic lower respiratory diseases (J40-J47)	46	4.3	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	23	2.1	0.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	2.0	0.7
9	Anemias (D50-D64)	16	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
11	Influenza and pneumonia (J10-J18)	8	0.7	*
12	Septicemia (A40-A41)	7	0.6	*
12	Cerebrovascular diseases (I60-I69)	7	0.6	*
14	Diabetes mellitus (E10-E14)	5	0.5	*
14	Meningitis (G00,G03)	5	0.5	*
...	All other causes (Residual)	185	17.1	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,286	100.0	193.5
1	Assault (homicide) (X85-Y09,Y87.1)	2,433	46.0	89.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,281	24.2	46.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	408	7.7	14.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	212	4.0	7.8
5	Malignant neoplasms (C00-C97)	181	3.4	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	52	1.0	1.9
6	Chronic lower respiratory diseases (J40-J47)	52	1.0	1.9
8	Anemias (D50-D64)	50	0.9	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	0.9	1.8
10	Legal intervention (Y35,Y89.0)	31	0.6	1.1
11	Influenza and pneumonia (J10-J18)	23	0.4	0.8
12	Diabetes mellitus (E10-E14)	19	0.4	*
12	Cerebrovascular diseases (I60-I69)	19	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
15	Septicemia (A40-A41)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
...	All other causes (Residual)	432	8.2	15.8
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,633	100.0	281.0
1	Assault (homicide) (X85-Y09,Y87.1)	1,771	26.7	75.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,313	19.8	55.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	919	13.9	38.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	546	8.2	23.1
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	432	6.5	18.3
6	Malignant neoplasms (C00-C97)	294	4.4	12.5
7	Diabetes mellitus (E10-E14)	85	1.3	3.6
8	Anemias (D50-D64)	72	1.1	3.0
8	Cerebrovascular diseases (I60-I69)	72	1.1	3.0
10	Chronic lower respiratory diseases (J40-J47)	67	1.0	2.8
11	Influenza and pneumonia (J10-J18)	58	0.9	2.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	0.7	2.0
13	Septicemia (A40-A41)	41	0.6	1.7
13	Legal intervention (Y35,Y89.0)	41	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.5	1.4
...	All other causes (Residual)	843	12.7	35.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	12,384	100.0	493.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,165	17.5	86.3
2	Human immunodeficiency virus (HIV) disease (B20-B24)	2,157	17.4	86.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,628	13.1	64.9
4	Malignant neoplasms (C00-C97)	1,255	10.1	50.0
5	Assault (homicide) (X85-Y09,Y87.1)	980	7.9	39.1
6	Cerebrovascular diseases (I60-I69)	387	3.1	15.4
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	353	2.9	14.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	293	2.4	11.7
9	Diabetes mellitus (E10-E14)	276	2.2	11.0
10	Influenza and pneumonia (J10-J18)	173	1.4	6.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	1.4	6.9
12	Septicemia (A40-A41)	154	1.2	6.1
13	Chronic lower respiratory diseases (J40-J47)	121	1.0	4.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	97	0.8	3.9
15	Viral hepatitis (B15-B19)	79	0.6	3.2
...	All other causes (Residual)	2,093	16.9	83.5
Black, non-Hispanic, male, 45-54 years				
...	All causes	19,026	100.0	1,119.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,716	24.8	277.4
2	Malignant neoplasms (C00-C97)	4,098	21.5	241.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,552	8.2	91.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,441	7.6	84.8
5	Cerebrovascular diseases (I60-I69)	900	4.7	52.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	649	3.4	38.2
7	Diabetes mellitus (E10-E14)	641	3.4	37.7
8	Assault (homicide) (X85-Y09,Y87.1)	431	2.3	25.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	1.9	20.8
10	Chronic lower respiratory diseases (J40-J47)	284	1.5	16.7
11	Septicemia (A40-A41)	279	1.5	16.4
12	Influenza and pneumonia (J10-J18)	277	1.5	16.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	218	1.1	12.8
14	Viral hepatitis (B15-B19)	215	1.1	12.6
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	183	1.0	10.8
...	All other causes (Residual)	2,789	14.7	164.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	22,493	100.0	2,316.1
1	Malignant neoplasms (C00-C97)	6,933	30.8	713.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,907	30.7	711.2
3	Cerebrovascular diseases (I60-I69)	1,235	5.5	127.2
4	Diabetes mellitus (E10-E14)	986	4.4	101.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	742	3.3	76.4
6	Chronic lower respiratory diseases (J40-J47)	624	2.8	64.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	471	2.1	48.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	453	2.0	46.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	426	1.9	43.9
10	Septicemia (A40-A41)	392	1.7	40.4
11	Influenza and pneumonia (J10-J18)	282	1.3	29.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.1	26.4
13	Assault (homicide) (X85-Y09,Y87.1)	159	0.7	16.4
14	Aortic aneurysm and dissection (I71)	96	0.4	9.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	95	0.4	9.8
...	All other causes (Residual)	2,436	10.8	250.8
Black, non-Hispanic, male, 65-74 years				
...	All causes	29,259	100.0	4,294.4
1	Malignant neoplasms (C00-C97)	9,228	31.5	1,354.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,016	30.8	1,323.3
3	Cerebrovascular diseases (I60-I69)	1,857	6.3	272.6
4	Diabetes mellitus (E10-E14)	1,262	4.3	185.2
5	Chronic lower respiratory diseases (J40-J47)	1,238	4.2	181.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	691	2.4	101.4
7	Septicemia (A40-A41)	609	2.1	89.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	2.0	87.5
9	Influenza and pneumonia (J10-J18)	518	1.8	76.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	348	1.2	51.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	273	0.9	40.1
12	Pneumonitis due to solids and liquids (J69)	168	0.6	24.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	160	0.5	23.5
14	Aortic aneurysm and dissection (I71)	147	0.5	21.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.4	18.8
...	All other causes (Residual)	3,020	10.3	443.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	27,552	100.0	8,530.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,416	30.5	2,605.7
2	Malignant neoplasms (C00-C97)	7,603	27.6	2,353.9
3	Cerebrovascular diseases (I60-I69)	2,100	7.6	650.2
4	Chronic lower respiratory diseases (J40-J47)	1,402	5.1	434.1
5	Diabetes mellitus (E10-E14)	1,016	3.7	314.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	803	2.9	248.6
7	Influenza and pneumonia (J10-J18)	688	2.5	213.0
8	Septicemia (A40-A41)	648	2.4	200.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	1.8	149.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	381	1.4	118.0
11	Alzheimer's disease (G30)	290	1.1	89.8
12	Pneumonitis due to solids and liquids (J69)	261	0.9	80.8
13	Atherosclerosis (I70)	142	0.5	44.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	38.4
14	Aortic aneurysm and dissection (I71)	124	0.5	38.4
...	All other causes (Residual)	3,071	11.1	950.8
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,628	100.0	16,321.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,945	33.8	5,517.3
2	Malignant neoplasms (C00-C97)	2,828	19.3	3,155.3
3	Cerebrovascular diseases (I60-I69)	1,225	8.4	1,366.8
4	Influenza and pneumonia (J10-J18)	676	4.6	754.2
5	Chronic lower respiratory diseases (J40-J47)	595	4.1	663.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	3.1	499.8
7	Diabetes mellitus (E10-E14)	426	2.9	475.3
8	Septicemia (A40-A41)	380	2.6	424.0
9	Alzheimer's disease (G30)	270	1.8	301.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	268	1.8	299.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	226	1.5	252.2
12	Pneumonitis due to solids and liquids (J69)	193	1.3	215.3
13	Atherosclerosis (I70)	140	1.0	156.2
14	Parkinson's disease (G20-G21)	74	0.5	82.6
15	Aortic aneurysm and dissection (I71)	65	0.4	72.5
...	All other causes (Residual)	1,869	12.8	2,085.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	138,096	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,636	29.4	233.3
2	Malignant neoplasms (C00-C97)	28,867	20.9	165.7
3	Cerebrovascular diseases (I60-I69)	10,908	7.9	62.6
4	Diabetes mellitus (E10-E14)	7,112	5.2	40.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,904	2.8	22.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,675	2.7	21.1
7	Chronic lower respiratory diseases (J40-J47)	3,388	2.5	19.5
8	Septicemia (A40-A41)	3,178	2.3	18.2
9	Influenza and pneumonia (J10-J18)	3,008	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,374	1.7	13.6
11	Certain conditions originating in the perinatal period (P00-P96)	2,182	1.6	12.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,158	1.6	12.4
13	Alzheimer's disease (G30)	1,636	1.2	9.4
14	Assault (homicide) (X85-Y09,Y87.1)	1,405	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	981	0.7	5.6
...	All other causes (Residual)	22,684	16.4	130.2
Black, non-Hispanic, female, 1-4 years				
...	All causes	549	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	30.6	16.4
2	Assault (homicide) (X85-Y09,Y87.1)	77	14.0	7.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.6	5.7
4	Malignant neoplasms (C00-C97)	23	4.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.0	2.2
6	Influenza and pneumonia (J10-J18)	15	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	7	1.3	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	Cerebrovascular diseases (I60-I69)	6	1.1	*
13	Meningitis (G00,G03)	5	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	123	22.4	12.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	690	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	33.9	8.1
2	Malignant neoplasms (C00-C97)	85	12.3	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	56	8.1	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	7.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.6	1.1
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	0.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	18	2.6	*
8	Anemias (D50-D64)	17	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
9	Cerebrovascular diseases (I60-I69)	8	1.2	*
11	Septicemia (A40-A41)	7	1.0	*
11	Influenza and pneumonia (J10-J18)	7	1.0	*
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Diabetes mellitus (E10-E14)	4	0.6	*
...	All other causes (Residual)	130	18.8	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,694	100.0	62.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	24.4	15.2
2	Assault (homicide) (X85-Y09,Y87.1)	319	18.8	11.8
3	Malignant neoplasms (C00-C97)	133	7.9	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	7.3	4.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	78	4.6	2.9
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	55	3.2	2.0
7	Pregnancy, childbirth and the puerperium (O00-099)	46	2.7	1.7
8	Anemias (D50-D64)	34	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.9	1.2
10	Influenza and pneumonia (J10-J18)	26	1.5	1.0
10	Chronic lower respiratory diseases (J40-J47)	26	1.5	1.0
12	Diabetes mellitus (E10-E14)	24	1.4	0.9
13	Cerebrovascular diseases (I60-I69)	21	1.2	0.8
14	Septicemia (A40-A41)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.9	*
...	All other causes (Residual)	329	19.4	12.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,359	100.0	127.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	560	16.7	21.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	12.7	16.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	367	10.9	13.9
4	Malignant neoplasms (C00-C97)	366	10.9	13.9
5	Assault (homicide) (X85-Y09,Y87.1)	351	10.4	13.3
6	Cerebrovascular diseases (I60-I69)	77	2.3	2.9
7	Diabetes mellitus (E10-E14)	75	2.2	2.8
8	Chronic lower respiratory diseases (J40-J47)	73	2.2	2.8
9	Pregnancy, childbirth and the puerperium (O00-099)	70	2.1	2.7
10	Anemias (D50-D64)	67	2.0	2.5
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	1.9	2.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.7	2.1
13	Influenza and pneumonia (J10-J18)	49	1.5	1.9
14	Septicemia (A40-A41)	44	1.3	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	1.1
...	All other causes (Residual)	686	20.4	26.0
Black, non-Hispanic, female, 35-44 years				
...	All causes	8,386	100.0	293.4
1	Malignant neoplasms (C00-C97)	1,794	21.4	62.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,341	16.0	46.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	997	11.9	34.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	7.4	21.6
5	Cerebrovascular diseases (I60-I69)	421	5.0	14.7
6	Assault (homicide) (X85-Y09,Y87.1)	325	3.9	11.4
7	Diabetes mellitus (E10-E14)	226	2.7	7.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	160	1.9	5.6
9	Chronic lower respiratory diseases (J40-J47)	137	1.6	4.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	1.6	4.8
11	Influenza and pneumonia (J10-J18)	132	1.6	4.6
12	Septicemia (A40-A41)	125	1.5	4.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	78	0.9	2.7
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	77	0.9	2.7
15	Anemias (D50-D64)	69	0.8	2.4
...	All other causes (Residual)	1,750	20.9	61.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-54 years				
...	All causes	12,351	100.0	600.4
1	Malignant neoplasms (C00-C97)	3,651	29.6	177.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,732	22.1	132.8
3	Cerebrovascular diseases (I60-I69)	762	6.2	37.0
4	Diabetes mellitus (E10-E14)	578	4.7	28.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	505	4.1	24.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	462	3.7	22.5
7	Chronic lower respiratory diseases (J40-J47)	272	2.2	13.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	2.1	12.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	256	2.1	12.4
10	Septicemia (A40-A41)	239	1.9	11.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	152	1.2	7.4
12	Influenza and pneumonia (J10-J18)	148	1.2	7.2
13	Assault (homicide) (X85-Y09,Y87.1)	126	1.0	6.1
14	Anemias (D50-D64)	64	0.5	3.1
15	Viral hepatitis (B15-B19)	62	0.5	3.0
...	All other causes (Residual)	2,080	16.8	101.1
Black, non-Hispanic, female, 55-64 years				
...	All causes	16,570	100.0	1,296.5
1	Malignant neoplasms (C00-C97)	5,389	32.5	421.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,439	26.8	347.3
3	Diabetes mellitus (E10-E14)	1,150	6.9	90.0
4	Cerebrovascular diseases (I60-I69)	1,034	6.2	80.9
5	Chronic lower respiratory diseases (J40-J47)	461	2.8	36.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	2.7	35.4
7	Septicemia (A40-A41)	324	2.0	25.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	1.8	23.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	250	1.5	19.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	238	1.4	18.6
11	Influenza and pneumonia (J10-J18)	204	1.2	16.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	142	0.9	11.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.4	5.7
14	Aortic aneurysm and dissection (I71)	56	0.3	4.4
15	Viral hepatitis (B15-B19)	53	0.3	4.1
...	All other causes (Residual)	1,999	12.1	156.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-74 years				
...	All causes	26,308	100.0	2,809.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,003	30.4	854.6
2	Malignant neoplasms (C00-C97)	7,397	28.1	789.9
3	Cerebrovascular diseases (I60-I69)	1,935	7.4	206.6
4	Diabetes mellitus (E10-E14)	1,858	7.1	198.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	863	3.3	92.2
6	Chronic lower respiratory diseases (J40-J47)	852	3.2	91.0
7	Septicemia (A40-A41)	603	2.3	64.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	416	1.6	44.4
9	Influenza and pneumonia (J10-J18)	362	1.4	38.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	347	1.3	37.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	159	0.6	17.0
12	Alzheimer's disease (G30)	137	0.5	14.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.5	13.7
14	Aortic aneurysm and dissection (I71)	124	0.5	13.2
15	Pneumonitis due to solids and liquids (J69)	96	0.4	10.3
...	All other causes (Residual)	3,028	11.5	323.3
Black, non-Hispanic, female, 75-84 years				
...	All causes	33,082	100.0	6,129.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,392	34.4	2,110.8
2	Malignant neoplasms (C00-C97)	6,589	19.9	1,220.8
3	Cerebrovascular diseases (I60-I69)	3,214	9.7	595.5
4	Diabetes mellitus (E10-E14)	1,979	6.0	366.7
5	Chronic lower respiratory diseases (J40-J47)	962	2.9	178.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	947	2.9	175.5
7	Septicemia (A40-A41)	865	2.6	160.3
8	Influenza and pneumonia (J10-J18)	722	2.2	133.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	625	1.9	115.8
10	Alzheimer's disease (G30)	593	1.8	109.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	1.3	81.9
12	Pneumonitis due to solids and liquids (J69)	208	0.6	38.5
13	Aortic aneurysm and dissection (I71)	179	0.5	33.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	171	0.5	31.7
15	Atherosclerosis (I70)	170	0.5	31.5
...	All other causes (Residual)	4,024	12.2	745.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	31,267	100.0	14,474.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,111	38.7	5,606.5
2	Malignant neoplasms (C00-C97)	3,432	11.0	1,588.8
3	Cerebrovascular diseases (I60-I69)	3,419	10.9	1,582.7
4	Influenza and pneumonia (J10-J18)	1,287	4.1	595.8
5	Diabetes mellitus (E10-E14)	1,216	3.9	562.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	903	2.9	418.0
7	Septicemia (A40-A41)	901	2.9	417.1
8	Alzheimer's disease (G30)	889	2.8	411.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	616	2.0	285.2
10	Chronic lower respiratory diseases (J40-J47)	571	1.8	264.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	1.2	175.0
12	Atherosclerosis (I70)	342	1.1	158.3
13	Pneumonitis due to solids and liquids (J69)	328	1.0	151.8
14	Nutritional deficiencies (E40-E64)	155	0.5	71.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.4	64.8
...	All other causes (Residual)	4,579	14.6	2,119.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable