

Table LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	103,740	100.0	331.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,866	24.9	82.5
2	Malignant neoplasms (C00-C97)	20,233	19.5	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,650	8.3	27.6
4	Cerebrovascular diseases (I60-I69)	5,907	5.7	18.8
5	Diabetes mellitus (E10-E14)	5,182	5.0	16.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,972	2.9	9.5
7	Assault (homicide) (X85-Y09,Y87.1)	2,864	2.8	9.1
8	Chronic lower respiratory diseases (J40-J47)	2,859	2.8	9.1
9	Influenza and pneumonia (J10-J18)	2,246	2.2	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,153	2.1	6.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,958	1.9	6.2
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,695	1.6	5.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,584	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,498	1.4	4.8
15	Septicemia (A40-A41)	1,215	1.2	3.9
...	All other causes (Residual)	16,858	16.3	53.8
Hispanic, both sexes, 1-4 years				
...	All causes	882	100.0	32.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	38.7	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	11.5	3.7
3	Malignant neoplasms (C00-C97)	72	8.2	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	52	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.7	1.2
6	Certain conditions originating in the perinatal period (P00-P96)	18	2.0	*
7	Influenza and pneumonia (J10-J18)	17	1.9	*
8	Septicemia (A40-A41)	14	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
11	Meningitis (G00,G03)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	7	0.8	*
13	Anemias (D50-D64)	6	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
...	All other causes (Residual)	183	20.7	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,014	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	362	35.7	6.1
2	Malignant neoplasms (C00-C97)	176	17.4	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	89	8.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	5.6	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	3.6	0.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	30	3.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.7	*
9	Influenza and pneumonia (J10-J18)	10	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.9	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
13	Septicemia (A40-A41)	8	0.8	*
13	Diseases of appendix (K35-K38)	8	0.8	*
15	Pneumonitis due to solids and liquids (J69)	5	0.5	*
...	All other causes (Residual)	171	16.9	2.9
Hispanic, both sexes, 15-24 years				
...	All causes	4,505	100.0	82.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,946	43.2	35.6
2	Assault (homicide) (X85-Y09,Y87.1)	1,119	24.8	20.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	391	8.7	7.1
4	Malignant neoplasms (C00-C97)	259	5.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	2.6	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.2	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.7	0.6
8	Cerebrovascular diseases (I60-I69)	32	0.7	0.6
9	Influenza and pneumonia (J10-J18)	25	0.6	0.5
10	Pregnancy, childbirth and the puerperium (O00-099)	19	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
12	Diabetes mellitus (E10-E14)	14	0.3	*
13	Legal intervention (Y35,Y89.0)	13	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.3	*
15	Septicemia (A40-A41)	9	0.2	*
...	All other causes (Residual)	446	9.9	8.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	13,195	100.0	132.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,397	25.7	34.0
2	Malignant neoplasms (C00-C97)	1,670	12.7	16.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,262	9.6	12.6
4	Assault (homicide) (X85-Y09,Y87.1)	1,233	9.3	12.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,047	7.9	10.5
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	792	6.0	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	598	4.5	6.0
8	Cerebrovascular diseases (I60-I69)	343	2.6	3.4
9	Diabetes mellitus (E10-E14)	200	1.5	2.0
10	Viral hepatitis (B15-B19)	120	0.9	1.2
11	Chronic lower respiratory diseases (J40-J47)	110	0.8	1.1
12	Septicemia (A40-A41)	100	0.8	1.0
13	Influenza and pneumonia (J10-J18)	98	0.7	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.6	0.8
...	All other causes (Residual)	2,054	15.6	20.6
Hispanic, both sexes, 45-64 years				
...	All causes	22,195	100.0	483.2
1	Malignant neoplasms (C00-C97)	6,202	27.9	135.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,961	22.4	108.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,425	6.4	31.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,392	6.3	30.3
5	Diabetes mellitus (E10-E14)	1,340	6.0	29.2
6	Cerebrovascular diseases (I60-I69)	1,081	4.9	23.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	592	2.7	12.9
8	Chronic lower respiratory diseases (J40-J47)	354	1.6	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	337	1.5	7.3
10	Viral hepatitis (B15-B19)	330	1.5	7.2
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	322	1.5	7.0
12	Assault (homicide) (X85-Y09,Y87.1)	266	1.2	5.8
13	Influenza and pneumonia (J10-J18)	259	1.2	5.6
14	Septicemia (A40-A41)	223	1.0	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	146	0.7	3.2
...	All other causes (Residual)	2,965	13.4	64.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	57,471	100.0	3,114.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,571	34.1	1,060.6
2	Malignant neoplasms (C00-C97)	11,838	20.6	641.5
3	Cerebrovascular diseases (I60-I69)	4,419	7.7	239.5
4	Diabetes mellitus (E10-E14)	3,624	6.3	196.4
5	Chronic lower respiratory diseases (J40-J47)	2,348	4.1	127.2
6	Influenza and pneumonia (J10-J18)	1,773	3.1	96.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,126	2.0	61.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,034	1.8	56.0
9	Alzheimer's disease (G30)	963	1.7	52.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	941	1.6	51.0
11	Septicemia (A40-A41)	818	1.4	44.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	575	1.0	31.2
13	Parkinson's disease (G20-G21)	378	0.7	20.5
14	Atherosclerosis (I70)	368	0.6	19.9
15	Pneumonitis due to solids and liquids (J69)	361	0.6	19.6
...	All other causes (Residual)	7,334	12.8	397.4
Hispanic, male, all ages				
...	All causes	57,991	100.0	367.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,554	23.4	86.0
2	Malignant neoplasms (C00-C97)	10,668	18.4	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,578	11.3	41.7
4	Cerebrovascular diseases (I60-I69)	2,809	4.8	17.8
5	Assault (homicide) (X85-Y09,Y87.1)	2,399	4.1	15.2
6	Diabetes mellitus (E10-E14)	2,335	4.0	14.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,143	3.7	13.6
8	Chronic lower respiratory diseases (J40-J47)	1,547	2.7	9.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,525	2.6	9.7
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,430	2.5	9.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,192	2.1	7.6
12	Influenza and pneumonia (J10-J18)	1,092	1.9	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	814	1.4	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	756	1.3	4.8
15	Septicemia (A40-A41)	605	1.0	3.8
...	All other causes (Residual)	8,544	14.7	54.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	481	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	41.0	14.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.1	4.5
3	Malignant neoplasms (C00-C97)	34	7.1	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	25	5.2	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.2	1.4
6	Influenza and pneumonia (J10-J18)	10	2.1	*
7	Septicemia (A40-A41)	8	1.7	*
8	Meningitis (G00,G03)	5	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
10	Anemias (D50-D64)	4	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	92	19.1	6.6
Hispanic, male, 5-14 years				
...	All causes	592	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	38.3	7.4
2	Malignant neoplasms (C00-C97)	92	15.5	3.0
3	Assault (homicide) (X85-Y09,Y87.1)	59	10.0	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	4.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.9	0.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	21	3.5	0.7
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.2	*
10	Cerebrovascular diseases (I60-I69)	6	1.0	*
10	Influenza and pneumonia (J10-J18)	6	1.0	*
12	Septicemia (A40-A41)	4	0.7	*
12	Diseases of appendix (K35-K38)	4	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Meningococcal infection (A39)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	84	14.2	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,547	100.0	124.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,575	44.4	55.5
2	Assault (homicide) (X85-Y09,Y87.1)	990	27.9	34.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	338	9.5	11.9
4	Malignant neoplasms (C00-C97)	158	4.5	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	2.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.8	1.1
7	Cerebrovascular diseases (I60-I69)	21	0.6	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.6	0.7
9	Legal intervention (Y35,Y89.0)	13	0.4	*
10	Influenza and pneumonia (J10-J18)	11	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
12	Diabetes mellitus (E10-E14)	8	0.2	*
14	Septicemia (A40-A41)	5	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.1	*
...	All other causes (Residual)	283	8.0	10.0
Hispanic, male, 25-44 years				
...	All causes	9,560	100.0	186.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,801	29.3	54.7
2	Assault (homicide) (X85-Y09,Y87.1)	1,052	11.0	20.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	980	10.3	19.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	791	8.3	15.5
5	Malignant neoplasms (C00-C97)	726	7.6	14.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	671	7.0	13.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	483	5.1	9.4
8	Cerebrovascular diseases (I60-I69)	208	2.2	4.1
9	Diabetes mellitus (E10-E14)	105	1.1	2.1
10	Viral hepatitis (B15-B19)	95	1.0	1.9
11	Influenza and pneumonia (J10-J18)	69	0.7	1.3
12	Septicemia (A40-A41)	65	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	49	0.5	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.4	0.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.4	0.8
...	All other causes (Residual)	1,384	14.5	27.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	13,987	100.0	635.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,439	24.6	156.2
2	Malignant neoplasms (C00-C97)	3,297	23.6	149.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,128	8.1	51.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,068	7.6	48.5
5	Diabetes mellitus (E10-E14)	732	5.2	33.2
6	Cerebrovascular diseases (I60-I69)	639	4.6	29.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	472	3.4	21.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	263	1.9	11.9
9	Viral hepatitis (B15-B19)	228	1.6	10.4
10	Assault (homicide) (X85-Y09,Y87.1)	206	1.5	9.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.4	9.1
12	Chronic lower respiratory diseases (J40-J47)	187	1.3	8.5
13	Influenza and pneumonia (J10-J18)	151	1.1	6.9
14	Septicemia (A40-A41)	134	1.0	6.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	0.5	3.4
...	All other causes (Residual)	1,767	12.6	80.2
Hispanic, male, 65 years and over				
...	All causes	27,353	100.0	3,518.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,162	33.5	1,178.6
2	Malignant neoplasms (C00-C97)	6,350	23.2	816.9
3	Cerebrovascular diseases (I60-I69)	1,926	7.0	247.8
4	Diabetes mellitus (E10-E14)	1,488	5.4	191.4
5	Chronic lower respiratory diseases (J40-J47)	1,284	4.7	165.2
6	Influenza and pneumonia (J10-J18)	805	2.9	103.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	590	2.2	75.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	2.0	70.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	527	1.9	67.8
10	Septicemia (A40-A41)	362	1.3	46.6
11	Alzheimer's disease (G30)	308	1.1	39.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	244	0.9	31.4
13	Parkinson's disease (G20-G21)	214	0.8	27.5
14	Pneumonitis due to solids and liquids (J69)	170	0.6	21.9
15	Aortic aneurysm and dissection (I71)	169	0.6	21.7
...	All other causes (Residual)	3,208	11.7	412.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	45,749	100.0	293.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,312	26.9	79.0
2	Malignant neoplasms (C00-C97)	9,565	20.9	61.4
3	Cerebrovascular diseases (I60-I69)	3,098	6.8	19.9
4	Diabetes mellitus (E10-E14)	2,847	6.2	18.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,072	4.5	13.3
6	Chronic lower respiratory diseases (J40-J47)	1,312	2.9	8.4
7	Influenza and pneumonia (J10-J18)	1,154	2.5	7.4
8	Certain conditions originating in the perinatal period (P00-P96)	961	2.1	6.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	829	1.8	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	770	1.7	4.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	742	1.6	4.8
12	Alzheimer's disease (G30)	667	1.5	4.3
13	Septicemia (A40-A41)	610	1.3	3.9
14	Assault (homicide) (X85-Y09,Y87.1)	465	1.0	3.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	433	0.9	2.8
...	All other causes (Residual)	7,912	17.3	50.8
Hispanic, female, 1-4 years				
...	All causes	401	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	35.9	10.7
2	Malignant neoplasms (C00-C97)	38	9.5	2.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	9.5	2.8
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.2	*
5	Certain conditions originating in the perinatal period (P00-P96)	13	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.0	*
8	Influenza and pneumonia (J10-J18)	7	1.7	*
9	Septicemia (A40-A41)	6	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Meningitis (G00,G03)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	83	20.7	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	422	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	32.0	4.6
2	Malignant neoplasms (C00-C97)	84	19.9	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	30	7.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	6.6	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.1	*
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	9	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J10-J18)	4	0.9	*
10	Pneumonitis due to solids and liquids (J69)	4	0.9	*
10	Diseases of appendix (K35-K38)	4	0.9	*
14	Cerebrovascular diseases (I60-I69)	3	0.7	*
15	Anemias (D50-D64)	2	0.5	*
15	Diabetes mellitus (E10-E14)	2	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	76	18.0	2.6
Hispanic, female, 15-24 years				
...	All causes	958	100.0	36.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	38.7	14.1
2	Assault (homicide) (X85-Y09,Y87.1)	129	13.5	4.9
3	Malignant neoplasms (C00-C97)	101	10.5	3.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	53	5.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.5	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	2.7	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	19	2.0	*
8	Influenza and pneumonia (J10-J18)	14	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Diabetes mellitus (E10-E14)	6	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
13	Septicemia (A40-A41)	4	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
13	Meningitis (G00,G03)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	150	15.7	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	3,635	100.0	74.8
1	Malignant neoplasms (C00-C97)	944	26.0	19.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	16.4	12.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	282	7.8	5.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	256	7.0	5.3
5	Assault (homicide) (X85-Y09,Y87.1)	181	5.0	3.7
6	Cerebrovascular diseases (I60-I69)	135	3.7	2.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	121	3.3	2.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	115	3.2	2.4
9	Diabetes mellitus (E10-E14)	95	2.6	2.0
10	Chronic lower respiratory diseases (J40-J47)	61	1.7	1.3
11	Pregnancy, childbirth and the puerperium (O00-099)	51	1.4	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	1.3	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.2	0.9
14	Septicemia (A40-A41)	35	1.0	0.7
15	Influenza and pneumonia (J10-J18)	29	0.8	0.6
...	All other causes (Residual)	644	17.7	13.2
Hispanic, female, 45-64 years				
...	All causes	8,208	100.0	343.2
1	Malignant neoplasms (C00-C97)	2,905	35.4	121.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,522	18.5	63.6
3	Diabetes mellitus (E10-E14)	608	7.4	25.4
4	Cerebrovascular diseases (I60-I69)	442	5.4	18.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	3.9	13.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	297	3.6	12.4
7	Chronic lower respiratory diseases (J40-J47)	167	2.0	7.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	1.7	5.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	120	1.5	5.0
10	Influenza and pneumonia (J10-J18)	108	1.3	4.5
11	Viral hepatitis (B15-B19)	102	1.2	4.3
12	Septicemia (A40-A41)	89	1.1	3.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	0.9	3.0
14	Assault (homicide) (X85-Y09,Y87.1)	60	0.7	2.5
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	59	0.7	2.5
...	All other causes (Residual)	1,198	14.6	50.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	30,118	100.0	2,820.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,409	34.6	974.7
2	Malignant neoplasms (C00-C97)	5,488	18.2	513.9
3	Cerebrovascular diseases (I60-I69)	2,493	8.3	233.4
4	Diabetes mellitus (E10-E14)	2,136	7.1	200.0
5	Chronic lower respiratory diseases (J40-J47)	1,064	3.5	99.6
6	Influenza and pneumonia (J10-J18)	968	3.2	90.6
7	Alzheimer's disease (G30)	655	2.2	61.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	580	1.9	54.3
9	Septicemia (A40-A41)	456	1.5	42.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	444	1.5	41.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	414	1.4	38.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	331	1.1	31.0
13	Atherosclerosis (I70)	207	0.7	19.4
14	Pneumonitis due to solids and liquids (J69)	191	0.6	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	177	0.6	16.6
...	All other causes (Residual)	4,105	13.6	384.4
White, non-Hispanic, both sexes, all ages				
...	All causes	1,953,197	100.0	996.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	607,704	31.1	310.0
2	Malignant neoplasms (C00-C97)	456,315	23.4	232.8
3	Cerebrovascular diseases (I60-I69)	138,716	7.1	70.8
4	Chronic lower respiratory diseases (J40-J47)	111,620	5.7	56.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,326	3.8	37.4
6	Influenza and pneumonia (J10-J18)	54,280	2.8	27.7
7	Diabetes mellitus (E10-E14)	49,347	2.5	25.2
8	Alzheimer's disease (G30)	40,835	2.1	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26,418	1.4	13.5
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,534	1.3	12.5
11	Septicemia (A40-A41)	23,023	1.2	11.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,602	1.0	10.0
13	Aortic aneurysm and dissection (I71)	13,934	0.7	7.1
14	Parkinson's disease (G20-G21)	13,475	0.7	6.9
15	Atherosclerosis (I70)	13,286	0.7	6.8
...	All other causes (Residual)	286,782	14.7	146.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,819	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	37.0	11.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	284	10.1	3.0
3	Malignant neoplasms (C00-C97)	281	10.0	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	156	5.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	3.3	1.0
6	Influenza and pneumonia (J10-J18)	58	2.1	0.6
7	Septicemia (A40-A41)	47	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	37	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	27	1.0	0.3
11	Meningitis (G00,G03)	26	0.9	0.3
12	Meningococcal infection (A39)	24	0.9	0.3
13	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	658	23.3	6.9
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,488	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,908	42.5	7.4
2	Malignant neoplasms (C00-C97)	608	13.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	5.4	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	176	3.9	0.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	174	3.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	3.5	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	1.5	0.3
8	Influenza and pneumonia (J10-J18)	62	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	55	1.2	0.2
10	Septicemia (A40-A41)	47	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	40	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	31	0.7	0.1
13	Diabetes mellitus (E10-E14)	19	0.4	*
14	Meningitis (G00,G03)	18	0.4	*
15	Anemias (D50-D64)	17	0.4	*
...	All other causes (Residual)	867	19.3	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	17,868	100.0	71.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,410	52.7	37.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,830	15.8	11.3
3	Malignant neoplasms (C00-C97)	1,070	6.0	4.3
4	Assault (homicide) (X85-Y09,Y87.1)	962	5.4	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	585	3.3	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	277	1.6	1.1
7	Chronic lower respiratory diseases (J40-J47)	110	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	105	0.6	0.4
9	Influenza and pneumonia (J10-J18)	95	0.5	0.4
10	Diabetes mellitus (E10-E14)	94	0.5	0.4
11	Septicemia (A40-A41)	71	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.2	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.2	0.1
15	Anemias (D50-D64)	28	0.2	0.1
...	All other causes (Residual)	2,103	11.8	8.4
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	81,616	100.0	140.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,699	22.9	32.1
2	Malignant neoplasms (C00-C97)	14,530	17.8	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,691	13.1	18.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	9,396	11.5	16.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,876	3.5	4.9
6	Assault (homicide) (X85-Y09,Y87.1)	2,462	3.0	4.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,391	2.9	4.1
8	Cerebrovascular diseases (I60-I69)	1,707	2.1	2.9
9	Diabetes mellitus (E10-E14)	1,594	2.0	2.7
10	Influenza and pneumonia (J10-J18)	836	1.0	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	760	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	679	0.8	1.2
13	Septicemia (A40-A41)	594	0.7	1.0
14	Viral hepatitis (B15-B19)	515	0.6	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	0.5	0.7
...	All other causes (Residual)	13,451	16.5	23.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	287,601	100.0	622.5
1	Malignant neoplasms (C00-C97)	105,732	36.8	228.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,773	25.3	157.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,847	4.8	30.0
4	Chronic lower respiratory diseases (J40-J47)	12,158	4.2	26.3
5	Cerebrovascular diseases (I60-I69)	9,454	3.3	20.5
6	Diabetes mellitus (E10-E14)	8,614	3.0	18.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	8,547	3.0	18.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	7,067	2.5	15.3
9	Influenza and pneumonia (J10-J18)	3,003	1.0	6.5
10	Septicemia (A40-A41)	2,807	1.0	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,353	0.8	5.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,655	0.6	3.6
13	Aortic aneurysm and dissection (I71)	1,635	0.6	3.5
14	Viral hepatitis (B15-B19)	1,423	0.5	3.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,393	0.5	3.0
...	All other causes (Residual)	35,140	12.2	76.1
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,545,161	100.0	5,329.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	523,130	33.9	1,804.5
2	Malignant neoplasms (C00-C97)	334,050	21.6	1,152.3
3	Cerebrovascular diseases (I60-I69)	127,328	8.2	439.2
4	Chronic lower respiratory diseases (J40-J47)	98,583	6.4	340.0
5	Influenza and pneumonia (J10-J18)	50,104	3.2	172.8
6	Alzheimer's disease (G30)	40,380	2.6	139.3
7	Diabetes mellitus (E10-E14)	39,018	2.5	134.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,972	1.8	96.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,490	1.5	81.0
10	Septicemia (A40-A41)	19,336	1.3	66.7
11	Parkinson's disease (G20-G21)	13,223	0.9	45.6
12	Atherosclerosis (I70)	12,738	0.8	43.9
13	Pneumonitis due to solids and liquids (J69)	12,269	0.8	42.3
14	Aortic aneurysm and dissection (I71)	11,992	0.8	41.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,873	0.7	37.5
...	All other causes (Residual)	200,675	13.0	692.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	944,913	100.0	984.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	293,206	31.0	305.5
2	Malignant neoplasms (C00-C97)	236,384	25.0	246.3
3	Chronic lower respiratory diseases (J40-J47)	55,314	5.9	57.6
4	Cerebrovascular diseases (I60-I69)	51,968	5.5	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,113	4.9	48.1
6	Diabetes mellitus (E10-E14)	23,177	2.5	24.2
7	Influenza and pneumonia (J10-J18)	23,097	2.4	24.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,620	2.1	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,793	1.4	13.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	12,613	1.3	13.1
11	Alzheimer's disease (G30)	12,253	1.3	12.8
12	Septicemia (A40-A41)	9,863	1.0	10.3
13	Aortic aneurysm and dissection (I71)	8,551	0.9	8.9
14	Parkinson's disease (G20-G21)	7,648	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	6,642	0.7	6.9
...	All other causes (Residual)	125,671	13.3	131.0
White, non-Hispanic, male, 1-4 years				
...	All causes	1,605	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	38.4	12.7
2	Malignant neoplasms (C00-C97)	160	10.0	3.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	9.6	3.2
4	Assault (homicide) (X85-Y09,Y87.1)	83	5.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	1.1
6	Septicemia (A40-A41)	25	1.6	0.5
7	Certain conditions originating in the perinatal period (P00-P96)	24	1.5	0.5
8	Influenza and pneumonia (J10-J18)	22	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Meningococcal infection (A39)	15	0.9	*
10	Meningitis (G00,G03)	15	0.9	*
10	Cerebrovascular diseases (I60-I69)	15	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
...	All other causes (Residual)	377	23.5	7.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,643	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	45.6	9.2
2	Malignant neoplasms (C00-C97)	345	13.1	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	141	5.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	4.5	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	93	3.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.2	0.6
7	Influenza and pneumonia (J10-J18)	36	1.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	36	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.2	0.2
10	Septicemia (A40-A41)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	21	0.8	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
13	Meningitis (G00,G03)	11	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.4	*
15	Diabetes mellitus (E10-E14)	9	0.3	*
...	All other causes (Residual)	462	17.5	3.5
White, non-Hispanic, male, 15-24 years				
...	All causes	12,677	100.0	98.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,749	53.2	52.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,416	19.1	18.8
3	Assault (homicide) (X85-Y09,Y87.1)	644	5.1	5.0
4	Malignant neoplasms (C00-C97)	629	5.0	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	357	2.8	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	1.4	1.4
7	Chronic lower respiratory diseases (J40-J47)	59	0.5	0.5
8	Diabetes mellitus (E10-E14)	52	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	51	0.4	0.4
9	Influenza and pneumonia (J10-J18)	51	0.4	0.4
11	Septicemia (A40-A41)	44	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.2	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.2	0.2
14	Legal intervention (Y35,Y89.0)	20	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,361	10.7	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	53,740	100.0	184.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,881	25.8	47.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,742	14.4	26.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	7,415	13.8	25.4
4	Malignant neoplasms (C00-C97)	6,766	12.6	23.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,411	4.5	8.3
6	Assault (homicide) (X85-Y09,Y87.1)	1,634	3.0	5.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,629	3.0	5.6
8	Diabetes mellitus (E10-E14)	950	1.8	3.3
9	Cerebrovascular diseases (I60-I69)	863	1.6	3.0
10	Influenza and pneumonia (J10-J18)	453	0.8	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	408	0.8	1.4
12	Viral hepatitis (B15-B19)	362	0.7	1.2
13	Septicemia (A40-A41)	330	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	305	0.6	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	248	0.5	0.9
...	All other causes (Residual)	8,343	15.5	28.6
White, non-Hispanic, male, 45-64 years				
...	All causes	175,843	100.0	776.1
1	Malignant neoplasms (C00-C97)	56,246	32.0	248.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,213	29.7	230.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,872	5.6	43.6
4	Chronic lower respiratory diseases (J40-J47)	6,265	3.6	27.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,250	3.6	27.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,389	3.1	23.8
7	Cerebrovascular diseases (I60-I69)	5,141	2.9	22.7
8	Diabetes mellitus (E10-E14)	5,001	2.8	22.1
9	Influenza and pneumonia (J10-J18)	1,719	1.0	7.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,467	0.8	6.5
11	Septicemia (A40-A41)	1,455	0.8	6.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,309	0.7	5.8
13	Aortic aneurysm and dissection (I71)	1,268	0.7	5.6
14	Viral hepatitis (B15-B19)	1,025	0.6	4.5
15	Assault (homicide) (X85-Y09,Y87.1)	872	0.5	3.8
...	All other causes (Residual)	20,351	11.6	89.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	690,621	100.0	5,730.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	232,597	33.7	1,929.9
2	Malignant neoplasms (C00-C97)	172,214	24.9	1,428.9
3	Chronic lower respiratory diseases (J40-J47)	48,631	7.0	403.5
4	Cerebrovascular diseases (I60-I69)	45,841	6.6	380.4
5	Influenza and pneumonia (J10-J18)	20,752	3.0	172.2
6	Diabetes mellitus (E10-E14)	17,162	2.5	142.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,519	2.0	112.2
8	Alzheimer's disease (G30)	12,025	1.7	99.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,160	1.6	92.6
10	Septicemia (A40-A41)	7,906	1.1	65.6
11	Parkinson's disease (G20-G21)	7,474	1.1	62.0
12	Aortic aneurysm and dissection (I71)	7,058	1.0	58.6
13	Pneumonitis due to solids and liquids (J69)	6,119	0.9	50.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4,725	0.7	39.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,548	0.7	37.7
...	All other causes (Residual)	78,890	11.4	654.6
White, non-Hispanic, female, all ages				
...	All causes	1,008,284	100.0	1,007.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314,498	31.2	314.2
2	Malignant neoplasms (C00-C97)	219,931	21.8	219.7
3	Cerebrovascular diseases (I60-I69)	86,748	8.6	86.7
4	Chronic lower respiratory diseases (J40-J47)	56,306	5.6	56.3
5	Influenza and pneumonia (J10-J18)	31,183	3.1	31.2
6	Alzheimer's disease (G30)	28,582	2.8	28.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,213	2.7	27.2
8	Diabetes mellitus (E10-E14)	26,170	2.6	26.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,625	1.4	13.6
10	Septicemia (A40-A41)	13,160	1.3	13.1
11	Atherosclerosis (I70)	8,559	0.8	8.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,653	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	6,989	0.7	7.0
14	Pneumonitis due to solids and liquids (J69)	6,497	0.6	6.5
15	Parkinson's disease (G20-G21)	5,827	0.6	5.8
...	All other causes (Residual)	155,343	15.4	155.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,214	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	35.0	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	10.7	2.8
3	Malignant neoplasms (C00-C97)	121	10.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	73	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J10-J18)	36	3.0	0.8
7	Septicemia (A40-A41)	22	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Meningitis (G00,G03)	11	0.9	*
13	Meningococcal infection (A39)	9	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
...	All other causes (Residual)	281	23.1	6.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,845	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	38.1	5.6
2	Malignant neoplasms (C00-C97)	263	14.3	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	6.8	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	83	4.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	3.9	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.8	0.3
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	1.8	0.3
8	Influenza and pneumonia (J10-J18)	26	1.4	0.2
9	Septicemia (A40-A41)	23	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	19	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	19	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.9	*
13	Anemias (D50-D64)	11	0.6	*
14	Diabetes mellitus (E10-E14)	10	0.5	*
15	Meningococcal infection (A39)	8	0.4	*
...	All other causes (Residual)	400	21.7	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,191	100.0	42.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,661	51.3	21.9
2	Malignant neoplasms (C00-C97)	441	8.5	3.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	414	8.0	3.4
4	Assault (homicide) (X85-Y09,Y87.1)	318	6.1	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	4.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	2.0	0.8
7	Cerebrovascular diseases (I60-I69)	54	1.0	0.4
8	Chronic lower respiratory diseases (J40-J47)	51	1.0	0.4
9	Influenza and pneumonia (J10-J18)	44	0.8	0.4
10	Diabetes mellitus (E10-E14)	42	0.8	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	0.2
12	Septicemia (A40-A41)	27	0.5	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	719	13.9	5.9
White, non-Hispanic, female, 25-44 years				
...	All causes	27,876	100.0	95.6
1	Malignant neoplasms (C00-C97)	7,764	27.9	26.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,818	17.3	16.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,949	10.6	10.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,981	7.1	6.8
5	Cerebrovascular diseases (I60-I69)	844	3.0	2.9
6	Assault (homicide) (X85-Y09,Y87.1)	828	3.0	2.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	762	2.7	2.6
8	Diabetes mellitus (E10-E14)	644	2.3	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	465	1.7	1.6
10	Influenza and pneumonia (J10-J18)	383	1.4	1.3
11	Chronic lower respiratory diseases (J40-J47)	374	1.3	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	352	1.3	1.2
13	Septicemia (A40-A41)	264	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	0.7	0.6
15	Viral hepatitis (B15-B19)	153	0.5	0.5
...	All other causes (Residual)	5,108	18.3	17.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	111,758	100.0	474.7
1	Malignant neoplasms (C00-C97)	49,486	44.3	210.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,560	18.4	87.3
3	Chronic lower respiratory diseases (J40-J47)	5,893	5.3	25.0
4	Cerebrovascular diseases (I60-I69)	4,313	3.9	18.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,975	3.6	16.9
6	Diabetes mellitus (E10-E14)	3,613	3.2	15.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,297	2.1	9.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,678	1.5	7.1
9	Septicemia (A40-A41)	1,352	1.2	5.7
10	Influenza and pneumonia (J10-J18)	1,284	1.1	5.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	0.9	4.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	581	0.5	2.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	468	0.4	2.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	399	0.4	1.7
15	Viral hepatitis (B15-B19)	398	0.4	1.7
...	All other causes (Residual)	14,417	12.9	61.2
White, non-Hispanic, female, 65 years and over				
...	All causes	854,540	100.0	5,044.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290,533	34.0	1,715.2
2	Malignant neoplasms (C00-C97)	161,836	18.9	955.4
3	Cerebrovascular diseases (I60-I69)	81,487	9.5	481.1
4	Chronic lower respiratory diseases (J40-J47)	49,952	5.8	294.9
5	Influenza and pneumonia (J10-J18)	29,352	3.4	173.3
6	Alzheimer's disease (G30)	28,355	3.3	167.4
7	Diabetes mellitus (E10-E14)	21,856	2.6	129.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,453	1.7	85.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,330	1.4	72.8
10	Septicemia (A40-A41)	11,430	1.3	67.5
11	Atherosclerosis (I70)	8,367	1.0	49.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,210	0.8	42.6
13	Pneumonitis due to solids and liquids (J69)	6,150	0.7	36.3
14	Parkinson's disease (G20-G21)	5,749	0.7	33.9
15	Aortic aneurysm and dissection (I71)	4,934	0.6	29.1
...	All other causes (Residual)	120,546	14.1	711.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	281,979	100.0	852.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,713	27.6	234.8
2	Malignant neoplasms (C00-C97)	61,409	21.8	185.6
3	Cerebrovascular diseases (I60-I69)	18,731	6.6	56.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,533	4.4	37.9
5	Diabetes mellitus (E10-E14)	11,830	4.2	35.7
6	Chronic lower respiratory diseases (J40-J47)	7,840	2.8	23.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,751	2.7	23.4
8	Assault (homicide) (X85-Y09,Y87.1)	7,536	2.7	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,659	2.4	20.1
10	Influenza and pneumonia (J10-J18)	5,794	2.1	17.5
11	Septicemia (A40-A41)	5,772	2.0	17.4
12	Certain conditions originating in the perinatal period (P00-P96)	5,000	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,708	1.3	11.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,793	1.0	8.4
15	Alzheimer's disease (G30)	2,325	0.8	7.0
...	All other causes (Residual)	44,585	15.8	134.7
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,283	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	33.4	20.6
2	Assault (homicide) (X85-Y09,Y87.1)	151	11.8	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	10.7	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.8	2.4
5	Malignant neoplasms (C00-C97)	47	3.7	2.3
6	Influenza and pneumonia (J10-J18)	40	3.1	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	29	2.3	1.4
8	Anemias (D50-D64)	20	1.6	1.0
9	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
11	Septicemia (A40-A41)	16	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
13	Meningitis (G00,G03)	11	0.9	*
14	Cerebrovascular diseases (I60-I69)	10	0.8	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
...	All other causes (Residual)	289	22.5	13.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,771	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	701	39.6	11.9
2	Malignant neoplasms (C00-C97)	179	10.1	3.0
3	Assault (homicide) (X85-Y09,Y87.1)	145	8.2	2.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	5.9	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	4.3	1.3
6	Chronic lower respiratory diseases (J40-J47)	66	3.7	1.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.3	0.7
8	Anemias (D50-D64)	33	1.9	0.6
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	1.6	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.0	*
11	Cerebrovascular diseases (I60-I69)	15	0.8	*
11	Influenza and pneumonia (J10-J18)	15	0.8	*
13	Septicemia (A40-A41)	14	0.8	*
14	Diabetes mellitus (E10-E14)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
...	All other causes (Residual)	317	17.9	5.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,980	100.0	128.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,752	39.4	50.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,694	24.3	31.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	463	6.6	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	4.8	6.2
5	Malignant neoplasms (C00-C97)	314	4.5	5.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	130	1.9	2.4
7	Anemias (D50-D64)	84	1.2	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	1.5
9	Chronic lower respiratory diseases (J40-J47)	78	1.1	1.4
10	Influenza and pneumonia (J10-J18)	49	0.7	0.9
11	Pregnancy, childbirth and the puerperium (O00-099)	46	0.7	0.8
12	Diabetes mellitus (E10-E14)	43	0.6	0.8
13	Cerebrovascular diseases (I60-I69)	40	0.6	0.7
14	Septicemia (A40-A41)	32	0.5	0.6
15	Legal intervention (Y35,Y89.0)	31	0.4	0.6
...	All other causes (Residual)	806	11.5	14.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	30,762	100.0	296.8
1	Human immunodeficiency virus (HIV) disease (B20-B24)	4,633	15.1	44.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,419	14.4	42.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,983	12.9	38.4
4	Malignant neoplasms (C00-C97)	3,709	12.1	35.8
5	Assault (homicide) (X85-Y09,Y87.1)	3,427	11.1	33.1
6	Cerebrovascular diseases (I60-I69)	957	3.1	9.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	927	3.0	8.9
8	Diabetes mellitus (E10-E14)	662	2.2	6.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	507	1.6	4.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	413	1.3	4.0
11	Influenza and pneumonia (J10-J18)	412	1.3	4.0
12	Chronic lower respiratory diseases (J40-J47)	398	1.3	3.8
13	Septicemia (A40-A41)	364	1.2	3.5
14	Anemias (D50-D64)	281	0.9	2.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	212	0.7	2.0
...	All other causes (Residual)	5,458	17.7	52.7
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	70,440	100.0	1,172.8
1	Malignant neoplasms (C00-C97)	20,071	28.5	334.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,794	26.7	312.9
3	Cerebrovascular diseases (I60-I69)	3,931	5.6	65.4
4	Diabetes mellitus (E10-E14)	3,355	4.8	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,950	4.2	49.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,670	3.8	44.5
7	Chronic lower respiratory diseases (J40-J47)	1,641	2.3	27.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,596	2.3	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,494	2.1	24.9
10	Septicemia (A40-A41)	1,234	1.8	20.5
11	Influenza and pneumonia (J10-J18)	911	1.3	15.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	876	1.2	14.6
13	Assault (homicide) (X85-Y09,Y87.1)	751	1.1	12.5
14	Viral hepatitis (B15-B19)	395	0.6	6.6
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	342	0.5	5.7
...	All other causes (Residual)	9,429	13.4	157.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	162,096	100.0	5,817.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,883	33.2	1,934.0
2	Malignant neoplasms (C00-C97)	37,077	22.9	1,330.8
3	Cerebrovascular diseases (I60-I69)	13,750	8.5	493.5
4	Diabetes mellitus (E10-E14)	7,757	4.8	278.4
5	Chronic lower respiratory diseases (J40-J47)	5,620	3.5	201.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,655	2.9	167.1
7	Influenza and pneumonia (J10-J18)	4,253	2.6	152.6
8	Septicemia (A40-A41)	4,006	2.5	143.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,612	1.6	93.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,514	1.6	90.2
11	Alzheimer's disease (G30)	2,291	1.4	82.2
12	Pneumonitis due to solids and liquids (J69)	1,254	0.8	45.0
13	Atherosclerosis (I70)	944	0.6	33.9
14	Aortic aneurysm and dissection (I71)	763	0.5	27.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	745	0.5	26.7
...	All other causes (Residual)	19,972	12.3	716.8
Black, non-Hispanic, male, all ages				
...	All causes	143,883	100.0	918.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,077	25.8	236.6
2	Malignant neoplasms (C00-C97)	32,542	22.6	207.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,629	6.0	55.1
4	Cerebrovascular diseases (I60-I69)	7,823	5.4	49.9
5	Assault (homicide) (X85-Y09,Y87.1)	6,131	4.3	39.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,377	3.7	34.3
7	Diabetes mellitus (E10-E14)	4,718	3.3	30.1
8	Chronic lower respiratory diseases (J40-J47)	4,452	3.1	28.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,984	2.1	19.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,818	2.0	18.0
11	Influenza and pneumonia (J10-J18)	2,786	1.9	17.8
12	Septicemia (A40-A41)	2,594	1.8	16.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,812	1.3	11.6
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,630	1.1	10.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,550	1.1	9.9
...	All other causes (Residual)	20,960	14.6	133.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	734	100.0	69.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	35.4	24.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	10.8	7.5
3	Assault (homicide) (X85-Y09,Y87.1)	74	10.1	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.7	2.6
5	Influenza and pneumonia (J10-J18)	25	3.4	2.4
6	Malignant neoplasms (C00-C97)	24	3.3	2.3
7	Certain conditions originating in the perinatal period (P00-P96)	19	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.5	*
9	Anemias (D50-D64)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
13	Meningitis (G00,G03)	6	0.8	*
14	Cerebrovascular diseases (I60-I69)	4	0.5	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	3	0.4	*
...	All other causes (Residual)	158	21.5	15.0
Black, non-Hispanic, male, 5-14 years				
...	All causes	1,081	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	43.2	15.6
2	Malignant neoplasms (C00-C97)	94	8.7	3.1
3	Assault (homicide) (X85-Y09,Y87.1)	89	8.2	3.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	4.9	1.8
5	Chronic lower respiratory diseases (J40-J47)	46	4.3	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	23	2.1	0.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	2.0	0.7
9	Anemias (D50-D64)	16	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
11	Influenza and pneumonia (J10-J18)	8	0.7	*
12	Septicemia (A40-A41)	7	0.6	*
12	Cerebrovascular diseases (I60-I69)	7	0.6	*
14	Diabetes mellitus (E10-E14)	5	0.5	*
14	Meningitis (G00,G03)	5	0.5	*
...	All other causes (Residual)	185	17.1	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,286	100.0	193.5
1	Assault (homicide) (X85-Y09,Y87.1)	2,433	46.0	89.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,281	24.2	46.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	408	7.7	14.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	212	4.0	7.8
5	Malignant neoplasms (C00-C97)	181	3.4	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	52	1.0	1.9
6	Chronic lower respiratory diseases (J40-J47)	52	1.0	1.9
8	Anemias (D50-D64)	50	0.9	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	0.9	1.8
10	Legal intervention (Y35,Y89.0)	31	0.6	1.1
11	Influenza and pneumonia (J10-J18)	23	0.4	0.8
12	Diabetes mellitus (E10-E14)	19	0.4	*
12	Cerebrovascular diseases (I60-I69)	19	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
15	Septicemia (A40-A41)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
...	All other causes (Residual)	432	8.2	15.8
Black, non-Hispanic, male, 25-44 years				
...	All causes	19,017	100.0	390.6
1	Human immunodeficiency virus (HIV) disease (B20-B24)	3,076	16.2	63.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,941	15.5	60.4
3	Assault (homicide) (X85-Y09,Y87.1)	2,751	14.5	56.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,711	14.3	55.7
5	Malignant neoplasms (C00-C97)	1,549	8.1	31.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	785	4.1	16.1
7	Cerebrovascular diseases (I60-I69)	459	2.4	9.4
8	Diabetes mellitus (E10-E14)	361	1.9	7.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	320	1.7	6.6
10	Influenza and pneumonia (J10-J18)	231	1.2	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	1.2	4.5
12	Septicemia (A40-A41)	195	1.0	4.0
13	Chronic lower respiratory diseases (J40-J47)	188	1.0	3.9
14	Anemias (D50-D64)	145	0.8	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	0.6	2.3
...	All other causes (Residual)	2,971	15.6	61.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	41,519	100.0	1,554.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,623	28.0	435.1
2	Malignant neoplasms (C00-C97)	11,031	26.6	413.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,183	5.3	81.7
4	Cerebrovascular diseases (I60-I69)	2,135	5.1	79.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,023	4.9	75.7
6	Diabetes mellitus (E10-E14)	1,627	3.9	60.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,102	2.7	41.3
8	Chronic lower respiratory diseases (J40-J47)	908	2.2	34.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	779	1.9	29.2
10	Septicemia (A40-A41)	671	1.6	25.1
11	Assault (homicide) (X85-Y09,Y87.1)	590	1.4	22.1
12	Influenza and pneumonia (J10-J18)	559	1.3	20.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	474	1.1	17.7
14	Viral hepatitis (B15-B19)	280	0.7	10.5
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	278	0.7	10.4
...	All other causes (Residual)	5,256	12.7	196.8
Black, non-Hispanic, male, 65 years and over				
...	All causes	71,439	100.0	6,530.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,377	31.3	2,045.5
2	Malignant neoplasms (C00-C97)	19,659	27.5	1,797.1
3	Cerebrovascular diseases (I60-I69)	5,182	7.3	473.7
4	Chronic lower respiratory diseases (J40-J47)	3,235	4.5	295.7
5	Diabetes mellitus (E10-E14)	2,704	3.8	247.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,942	2.7	177.5
7	Influenza and pneumonia (J10-J18)	1,882	2.6	172.0
8	Septicemia (A40-A41)	1,637	2.3	149.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,347	1.9	123.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	955	1.3	87.3
11	Alzheimer's disease (G30)	672	0.9	61.4
12	Pneumonitis due to solids and liquids (J69)	622	0.9	56.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	386	0.5	35.3
14	Atherosclerosis (I70)	362	0.5	33.1
15	Aortic aneurysm and dissection (I71)	336	0.5	30.7
...	All other causes (Residual)	8,141	11.4	744.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	138,096	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,636	29.4	233.3
2	Malignant neoplasms (C00-C97)	28,867	20.9	165.7
3	Cerebrovascular diseases (I60-I69)	10,908	7.9	62.6
4	Diabetes mellitus (E10-E14)	7,112	5.2	40.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,904	2.8	22.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,675	2.7	21.1
7	Chronic lower respiratory diseases (J40-J47)	3,388	2.5	19.5
8	Septicemia (A40-A41)	3,178	2.3	18.2
9	Influenza and pneumonia (J10-J18)	3,008	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,374	1.7	13.6
11	Certain conditions originating in the perinatal period (P00-P96)	2,182	1.6	12.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,158	1.6	12.4
13	Alzheimer's disease (G30)	1,636	1.2	9.4
14	Assault (homicide) (X85-Y09,Y87.1)	1,405	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	981	0.7	5.6
...	All other causes (Residual)	22,684	16.4	130.2
Black, non-Hispanic, female, 1-4 years				
...	All causes	549	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	30.6	16.4
2	Assault (homicide) (X85-Y09,Y87.1)	77	14.0	7.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.6	5.7
4	Malignant neoplasms (C00-C97)	23	4.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.0	2.2
6	Influenza and pneumonia (J10-J18)	15	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	7	1.3	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	Cerebrovascular diseases (I60-I69)	6	1.1	*
13	Meningitis (G00,G03)	5	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	123	22.4	12.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	690	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	33.9	8.1
2	Malignant neoplasms (C00-C97)	85	12.3	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	56	8.1	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	7.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.6	1.1
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	0.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	18	2.6	*
8	Anemias (D50-D64)	17	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
9	Cerebrovascular diseases (I60-I69)	8	1.2	*
11	Septicemia (A40-A41)	7	1.0	*
11	Influenza and pneumonia (J10-J18)	7	1.0	*
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Diabetes mellitus (E10-E14)	4	0.6	*
...	All other causes (Residual)	130	18.8	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,694	100.0	62.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	24.4	15.2
2	Assault (homicide) (X85-Y09,Y87.1)	319	18.8	11.8
3	Malignant neoplasms (C00-C97)	133	7.9	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	7.3	4.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	78	4.6	2.9
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	55	3.2	2.0
7	Pregnancy, childbirth and the puerperium (O00-099)	46	2.7	1.7
8	Anemias (D50-D64)	34	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.9	1.2
10	Influenza and pneumonia (J10-J18)	26	1.5	1.0
10	Chronic lower respiratory diseases (J40-J47)	26	1.5	1.0
12	Diabetes mellitus (E10-E14)	24	1.4	0.9
13	Cerebrovascular diseases (I60-I69)	21	1.2	0.8
14	Septicemia (A40-A41)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.9	*
...	All other causes (Residual)	329	19.4	12.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, 25-44 years				
...	All causes	11,745	100.0	213.8
1	Malignant neoplasms (C00-C97)	2,160	18.4	39.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,708	14.5	31.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,557	13.3	28.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	8.9	19.0
5	Assault (homicide) (X85-Y09,Y87.1)	676	5.8	12.3
6	Cerebrovascular diseases (I60-I69)	498	4.2	9.1
7	Diabetes mellitus (E10-E14)	301	2.6	5.5
8	Chronic lower respiratory diseases (J40-J47)	210	1.8	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	1.6	3.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	187	1.6	3.4
11	Influenza and pneumonia (J10-J18)	181	1.5	3.3
12	Septicemia (A40-A41)	169	1.4	3.1
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	142	1.2	2.6
14	Anemias (D50-D64)	136	1.2	2.5
15	Pregnancy, childbirth and the puerperium (O00-099)	110	0.9	2.0
...	All other causes (Residual)	2,475	21.1	45.0
Black, non-Hispanic, female, 45-64 years				
...	All causes	28,921	100.0	867.1
1	Malignant neoplasms (C00-C97)	9,040	31.3	271.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,171	24.8	215.0
3	Cerebrovascular diseases (I60-I69)	1,796	6.2	53.8
4	Diabetes mellitus (E10-E14)	1,728	6.0	51.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	767	2.7	23.0
6	Chronic lower respiratory diseases (J40-J47)	733	2.5	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	715	2.5	21.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	647	2.2	19.4
9	Septicemia (A40-A41)	563	1.9	16.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	494	1.7	14.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	402	1.4	12.1
12	Influenza and pneumonia (J10-J18)	352	1.2	10.6
13	Assault (homicide) (X85-Y09,Y87.1)	161	0.6	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	4.0
15	Viral hepatitis (B15-B19)	115	0.4	3.4
...	All other causes (Residual)	4,104	14.2	123.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 1999

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	90,657	100.0	5,357.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,506	34.8	1,861.9
2	Malignant neoplasms (C00-C97)	17,418	19.2	1,029.3
3	Cerebrovascular diseases (I60-I69)	8,568	9.5	506.3
4	Diabetes mellitus (E10-E14)	5,053	5.6	298.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,713	3.0	160.3
6	Chronic lower respiratory diseases (J40-J47)	2,385	2.6	140.9
7	Influenza and pneumonia (J10-J18)	2,371	2.6	140.1
8	Septicemia (A40-A41)	2,369	2.6	140.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,657	1.8	97.9
10	Alzheimer's disease (G30)	1,619	1.8	95.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,167	1.3	69.0
12	Pneumonitis due to solids and liquids (J69)	632	0.7	37.3
13	Atherosclerosis (I70)	582	0.6	34.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	439	0.5	25.9
15	Aortic aneurysm and dissection (I71)	427	0.5	25.2
...	All other causes (Residual)	11,751	13.0	694.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable