

IV Antimicrobial Starts & NHSN Reporting: Do You Know What Counts?

Vannessa Cox, MBA, Public Health Analyst II

July 2025

Objectives

- Define what qualifies as an IV antimicrobial starts in NHSN.
- Identify where the antimicrobial start must occur.
- Accurately count and report qualifying starts.

What is an IV Antimicrobial Start?

- The administration of an intravenous (IV) antimicrobial treatment in an outpatient dialysis setting.
- Must be new or restarted after 21 days.
- Includes IV antiviral and antifungal treatments.

Where Must It Occur?

Location requirements:

- Must be administered in the outpatient dialysis facility.
- Medications given in emergency rooms, inpatient units, or infusion centers do not count.
- Temporary holding areas within a hospital are also excluded.

What Counts vs. What Does Not Count?

Does Count

- First dose of a new IV antimicrobial treatment.
- Restart of the same treatment after 21+ days without it.
- Any reason for administration.

Does NOT Count

- Continued administration of the same antimicrobial treatment.
- Oral antibiotics.
- Antimicrobial flushes.
- Administration that occurs outside the dialysis facility that does not qualify (e.g., hospital or ER).

How to Count Multiple Starts?

- One start per drug per patient per month
 - If a patient receives two qualifying starts in July (e.g., vancomycin and cefazolin): report both.
 - If patient restarts the same treatment within 21 days: do not report again.

Handling Unusual IV Antimicrobial Start Situations

Restart scenario:

- A patient stopped cefazolin on June 1 and received it again on July 10 at your dialysis facility - this counts as a new start.

Transfer scenario:

A patient received vancomycin in the hospital, and another dose was given in your dialysis facility on discharge day – this does NOT count unless it is a new start.

Precautionary dose scenario:

- A patient receives a new IV antimicrobial start at your dialysis clinic before any culture results this counts if it meets the 21-day rule and is not a continuation.
- Must be authorized by nephrologist or by a licensed provider.

Common Errors to Avoid

- Counting oral antimicrobials.
- Including hospital only administrations.
- Reporting maintenance doses multiple times.
- Forgetting the 21-day restart gap.

Reporting Tips & Reminders

- Double check the 21-day restart rule.
- Verify administration date and medication name.
- Confirm the medication was administered through an IV.
- New IV antimicrobial in dialysis outpatient unit is reportable.
- Transfer from hospital only count if newly started in your dialysis outpatient facility.
- Clarify route and consult if unsure.

Resources and Support

- NHSN Dialysis Event Surveillance Protocol
- NHSN Enrollment Guide
- Home Dialysis Center Practices Survey
- Submit <u>Service Now ticket</u>
- NHSN CDC Dialysis team NHSN@cdc.gov
- SAMS Helpdesk <u>SAMShelp@cdc.gov</u>

Thank You

For more information, contact CDC/ATSDR 1-800-CDC-INFO (232-4636)

TTY: 1-888-232-6348 www.cdc.gov www.atsdr.cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention and the Agency for Toxic Substances and Disease Registry.

