was involved. This information is intended for use in advancing knowledge needed to protect workers. Comments regarding this presentation may be submitted to the NIOSH Docket Office This presentation should not be considered a final statement of NIOSH policy or of any agency or individual who

National Personal Protective Technology Laboratory

(Gas Mask / APR) Use Guidelines **CBRN Air-Purifying Respirator**

Holiday Inn Select, Pittsburgh South Pittsburgh, PA

Mike Bergman

EG&G Technical Services, Inc.

July 19, 2005





GBRN Respirator Standards

to represent any agency disseminated by the National determination or policy. The findings and conclusions in this Institute for Occupational Safety and Health and should not be construed presentation have not been formally





CBRN APR Overview

- Statement of Standard for Chemical, Respirator (APR), Dated March 7, 2003 Biological, Radiological, and Nuclear (CBRN) Full Facepiece Air Purifying
- 14G approval under 42 CFR part 84
- 139 identified CBRN threat protections
- NIOSH assigned protection factor = 50







CBRN APR Guidelines

- NIOSH Cautions and Limitations specific to CBRN use
- **CBRN Respirator Use Life (CRUL)**
- Canister Capacity (Cap) Selection
- **CBRN Canister interchangeability**
- A crisis provision when supplies are limited
- Escape contingency from IDLH environments
- based on 5 min gas life testing at 100 lpm
- Industrial use / CBRN use





OSHA Respiratory Protection Standard Requirements - 29 CFR 1910.134

- Determination of medical fitness
- Fit testing of tight-fitting facepieces
- Procedures and schedules for cleaning, disinfecting, storing, inspecting, repairing, discarding, and otherwise maintaining respirators
- Canister change schedule for gases and vapors





CBRN APR Use Conditions

- All of the following conditions must be met
- Types of inhalation hazards and concentrations have been identified
- CBRN canister is capable of removing the hazard
- Oxygen is known to be at least 19.5% by volume
- less than the APR maximum use concentration (MUC) Contaminant concentrations are less than IDLH and
- vapors Canister change schedule established for gases and
- Use complies with all NIOSH Cautions and Limitations





Selection Matrix

- and protective clothing selection Interim guidelines for respirator
- Available on OSHA website www.osha.gov (follow emergency response links)
- airborne concentrations for PPE selection is given at defined blister agents and nerve agents







NPPTL Research to Practice

CBRN APR Canister Sticker Label

 Shows "Cap" level, protections, approval number, and cautions and limitations



TO THE YEAR

ANYWHERE USA

Canister PAN 12345XYZ Pricesion CBRN Cap 1



CAMPONIA VALUE CONTROL CONTROL

2. CAUTIONS AND UBITATIONS

After the same an administrative continues was that 12 figures the specific for a second continues of the second continues of

WAS applied an partient that to execute their best medianes or according with ACSA. COM, and other application Chiefe to that a projection, and/or measurement in partie of advances in the and manufacture of their expectant If apply apply Special and other their advances in advances in advances apply. When the Charles manufacture determined the manufacture determined the projection of the contraction before determined apply.

I CREW CAUTIONS AND LIMITATIONS

The expected of the section of the s





CBRN APR Threat Protections

- Minimum of 139 CBRN threat protections
- Chemical (110)
- Biological (13)
- Radiological and nuclear (16)
- CBRN canister is tested using 11 Test Representative Agents (TRA)
- 10 gases and 1 particulate aerosol
- Challenge concentration of TRA gases are multiples of IDLH of the TRA





Canister Gas/Vapor Service Life

Service life

- The length of time required for an air-purifying element to reach a specific effluent concentration

(2004 NIOSH Respirator Selection Logic)

Factors which affect canister service life*

- Quality and amount of sorbent
- Packing uniformity and density
- Breathing rate
- Relative humidity
- Temperature
- Contaminant concentration
- Affinity of gas or vapor for the sorbent
- Presence of other gases and vapors

^{*} The Occupational Environment: Its Evaluation and Control (Respiratory Protection, Sec 5: Chapter 36). Fairfax, VA: American Industrial Hygiene Association Press, 1997.





Canister Capacity (Cap) Selection

6 "Cap" levels based on NIOSH gas life testing

Cap 6	Cap 5	Cap 4	Cap 3	Cap 2	Cap 1	CBRN Canister Cap designations
%X	6X	4X	3X	2X		Relative gas/vapor service life compared to a Cap 1 canister*

*At similar exposure conditions (temperature, relative humidity, work rate, and contaminant concentration)





OSHA Requirement

- 29 CFR 1910.134 (d)(3)(iii)(B)(2)
- Based on objective information or data that will ensure that canisters are changed before the end of their service life
- Applies to gases and vapors, not particulates
- Where no end-of-service-life indicator (ESLI) is appropriate, a change schedule is required
- CBRN APR are not currently approved with ESLI





CBRN Canister interchangeability

A crisis provision

- Use of a different manufacturer's CBRN canister under restricted supply conditions
- Possible by the requirement of standardized threads and interface connectors
- Decision to proceed with interchangeability is the responsibility of the incident commander or other commanding authority under crisis conditions
- Assembly with a canister other than specified in the approval assembly matrix is not in its NIOSH-approved configuration





CBRN Use / Industrial Use

- different approved respirator configurations The same facepiece part number may be part of
- An approved configuration will specify either a CBRN canister or an industrial canister (e.g. P100)
- packaging until needed for a CBRN response The CBRN canister should not be used for routine industrial use and should remain in its sealed
- Always properly maintain the system so it can be used for a CBRN response





CBRN Respirator Use Life (CRUL)

Use Guidelines

- UU: "The respirator should not be used beyond 8 hours than 2 hours encountered, the respirator should not be used for more possibility of agent permeation. If liquid exposure is after initial exposure to chemical warfare agents to avoid
- CRUL time includes decontamination time
- Dispose entire system (facepiece, canister, and accessories) at end of CRUL time





CBRN Respirator Use Life (CRUL)

Chemical warfare agents applicable to (CRUL):

(G and V agents) **Nerve Agents** including:

- GB (Sarin)
- GA (Tabun)
-) (Soman)

GF (Cyclohexyl Sarin)

Blister Agents including: (mustards, Lewisite)

- H (sulfur mustard)
- HD (distilled sulfur mustard)
- Nitrogen mustards
- (HN-1, HN-2 and HN-3)
- Lewisite (L, L-1, L-2 and L-3)



Methods

- CBRN Respirator Use Life (CRUL)
- Software (mathematical models)
- Manufacturer's test data
- Rules of Thumb (not to be used as a sole method)





- Methods (cont.)
- CBRN Respirator Use Life (CRUL)
- In a CWA environment the 8-hour (vapor) and 2-hour of a longer calculated canister service life (liquid) maximum system use life applies regardless





- Methods (cont.)
- Software (OSHA website)
- OSHA Advisor Genius 1998
- Breakthrough Program 2003
- Corrects for relative humidity
- Both programs calculate change schedules for individual organic vapors only
- Manufacturer's software





- Methods (cont.)
- Manufacturer's test data (for a specific chemical)
- The manufacturer may have service life information for a specific chemical





- Methods (cont.) Rules of Thumb for organic vapors*
- If the chemical's boiling point is > 70 °C and the service life of 8 hours at a normal work rate concentration is less than 200 ppm you can expect a
- Service life is inversely proportional to work rate
- Reducing concentration by a factor of 10 will increase service life by a factor of 5
- Humidity above 85% will reduce service life by 50%

^{*} The Occupational Environment: Its Evaluation and Control (Respiratory Protection, Sec 5: Chapter 36). Fairfax, VA: American Industrial Hygiene Association Press, 1997.





NIOSH/NPPTL Contact Information

NIOSH, NPPTL 626 Cochrans Mill Road Pittsburgh PA 15236 P.O. Box 18070





Mike Bergman vzb3@cdc.gov EG&G Technical Services, Inc. NIOSH NPPTL Support



1-800-35-NIOSH

http://www.cdc.gov/niosh/npptl/default.html

any agency determination or policy. Mention of any company or product does not constitute by the National Institute for Occupational Safety and Health and should not be construed to represent DISCLAIMER: The findings and conclusions in this presentation have not been formally disseminated endorsement by NIOSH or NPPTL







CBRN Threats

- Chemical (gases, vapors, liquids, particulates)
- Chemical warfare agents
- Toxic industrial chemicals / Toxic industrial materials (TICs / TIMs)
- **Biological (particulates)**
- Micro-organisms (disease-causing bacteria and viruses) and biological toxins
- Radiological (particulates)
- Particulates carrying radiation dispersed by a radiological dispersive device (RDD) or "dirty bomb"
- Nuclear (particulates)
- Particulates carrying radiation dispersed from a detonation involving nuclear fuel, a nuclear weapon, or weapon's component





GBRN Protection

- CBRN respirators meet new design and performance standards, including special CBRN tests
- Live agent tests (LAT)
- System testing against GB (Sarin) and HD (distilled sulfur mustard)
- Laboratory respirator protection level (LRPL)
- Quantitative corn-oil, fit-factor test using human test subjects
- Canister gas life tests
- Address high physiological demand, need for large sorbent capacity, and protection for multiple respiratory hazards





Canister Capacity

Gas Life Laboratory Test Times

Capacity # 6	Capacity # 5	Capacity # 4	Capacity # 3	Capacity # 2	Capacity # 1	Capacity Level
120	90	60	45	30	15	Test Time (min)
Test Concentration X 120	Test Concentration X 90	Test Concentration X 60	Test Concentration X 45	Test Concentration X 30	Test Concentration X 15	Capacity (ppm-min)





Test Representative Agents

- Organic vapor family cyclohexane
- Acid gas family — SO₂, H₂S, CNCL, COCl₂, HCN
- Base gas family ammonia
- Hydride family — phosphine
- Nitrogen oxide family — nitrogen dioxide
- Formaldehyde family — formaldehyde
- Particulate family DOP





61 Organic vapor family

with vapor pressures less than that of cyclohexane

Acid gas family

Base gas family

Hydride family

Nitrogen oxide family

Formaldehyde family (only member of family)

32 Particulate family





Organic Vapor Family

chloroacetyl chloride	chloroacetophenone	chloroacetonitrile	chloroacetone	bromobenzylcyanide	bromoacetone	allyl chlorocarbonate	allyl alcohol	acrylonitrile	acetone cyanohydrin
methanesulfonyl chloride	lewisite	iso-propyl chloroformate	iso-butyl chloroformate	hexaethyl tetraphosphate	hexachlorocyclopentadiene	ethylene dibromide	ethyl phosphorodichloridate	ethyl chlorothioformate	ethyl chloroformate
tetraethyl dithiopyrophosphate	tert-octyl mercapatan	tabun	soman	sec-butyl chloroformate	sarin	phosgene oximedichloroforoxime	phenyldichloroarsine	phenylcarbylamine chloride	phenyl mercapatan





Organic Vapor Family (Continued)

chloropicrin	methyl orthosilicate	tetraethyl lead
chloropivaloyl chloride	methyl parathion	tetramethyl lead
crotonaldehyde	mustard, lewisite mixture	tetranitromethane
cyclohexyl methyphosphonate	nitrogen mustard HN-1	trimethoxysilane
dibenz-(b,f)-1,4-oxazepine	nitrogen mustard HN-2	trimethylacetyl chloride
diketene	nitrogen mustard HN-3	V-Sub X
dimethyl sulfate	n-propyl chloroformate	diphosgene
diphenylchloroarsine	o-chlorobenzylidene malononitrile	o-ethyl-s- (2isopropyaminoethyl)methyl phosphonthiolate
diphenylcyanoarsine	parathion	ethyl phosphonothioicdichloride
distilled mustard	perchloromethyl mercaptan	methyl phosphonic dichloride
		phosphorus oxychloride





Acid Gas Family

boron trichloride boron tribromide

boron trifluoride

bromine

bromine trifluoride bromine chloride

carbonyl fluoride

chlorine

chlorine pentafluoride

chlorine trifluoride

chlorosulfonic acid

cyanogen chloride

dichlorosilane

dichloride ethyl phosphonous

fluorine

hydrogen bromide

hydrogen chloride

hydrogen cyanide

hydrogen fluoride

hydrogen iodide

hydrogen sulfide

phosgene

phosphorus trichloride

silicon tetrafluoride

sulfur dioxide

sulfuric acid sulfur trioxide

sulfuryl chloride

titanium tetrachloride

tungsten hexafluoride

bromine pentafluoride*

hydrogen selenide*





Nitrogen Oxide Family	nitric acid	nitric acid, fuming	Nitrogen dioxide	nitrogen tetraoxide	nitrogen trioxide
Base Gas Family	allyl amine	ammonia	dimethyl hydrazine, 1,2	methyl hydrazine	
Hydride Family	arsine	germane	phosphine	stibine	
Particulate Family Formaldehyde Family	adamsite	sodium azide	Sodium fluoroacetate	13 Biological agents	16 Radiological / Nuclear agents
Formaldehyde Family	formaldehyde				





Particulate Family Biological Agents

Anthrax

Brucellosis

Glanders

Pneumonic plague

Tularemia

Q Fever

Smallpox

Viral hemorrhagic fevers

T-2 mycotoxins

Botulism

Ricin

Staphylococcus enterotoxin B

Venezuelan equine encephalitis





Radiological and Nuclear Agents Particulate Family

Hydrogen 3

Carbon 14

Phosphorous 32

Cobalt 60

Nickel 63

Strontium 90

Technetium 99m

lodine 131

Cesium 137

Promethium 147

Thallium 204

Radium 226

Thorium 232

Uranium 235 & 238

Plutonium 239

Americium 241



