

National Immunization Survey

A User's Guide for the 2015 Public-Use Data File

Centers for Disease Control and Prevention

**National Center for Immunization
and Respiratory Diseases**

Presented by:

NORC at the University of Chicago

September 2016

Acknowledgments

The development and production of the NIS public-use data files is a team effort that has included contributions from many individuals (listed in alphabetical order) in the three organizations:

National Center for Immunization and Respiratory Diseases, CDC – Laurie Elam-Evans, Holly Hill, James Singleton, Phil Smith, and Larry Wilkinson.

NORC at the University of Chicago – Ken Copeland, Nicholas Davis, Can Geng, Vicki Pineau, Benjamin Skalland, Xian Tao, Bess Welch, Kirk Wolter, and Michael Zeddies.

Table of Contents

| | |
|---|-----------|
| 1. Introduction..... | 1 |
| 2. Sample Design | 7 |
| 2.1. The NIS RDD Telephone Survey | 7 |
| 2.2. The NIS Provider Record Check | 9 |
| 2.3. Summary of Data Collection | 10 |
| 2.4. Informed Consent, Security, and Confidentiality of Information..... | 14 |
| 3. Content of NIS Questionnaires | 15 |
| 3.1. Content of the Household Questionnaire..... | 15 |
| 3.2. Content of the Immunization History Questionnaire (IHQ)..... | 18 |
| 4. Data Preparation and Processing Procedures..... | 19 |
| 4.1. Data Preparation | 19 |
| 4.1.1. Editing in the CATI System | 20 |
| 4.1.2. Post-CATI Edits | 20 |
| 4.1.3. Editing of Provider Data..... | 21 |
| 4.2. Limitations of Data Editing Procedures..... | 22 |
| 4.3. Variable-Naming Conventions | 23 |
| 4.4. Missing Value Codes..... | 23 |
| 4.5. Imputation for Item Non-Response | 23 |
| 4.6. Vaccine-Specific Recoding of Verbatim Responses | 25 |
| 4.7. Composite Variables..... | 25 |
| 4.8. Sub-Sets of the NIS Data..... | 28 |
| 4.9. Confidentiality and Disclosure Avoidance..... | 28 |
| 5. Quality Control and Quality Assurance Procedures | 28 |
| 6. Sampling Weights | 30 |
| 6.1. Base Sampling Weight | 31 |
| 6.2. Adjustments for Non-Resolution of Telephone Numbers, Screener Non-Response, and Interview Non-Response | 32 |
| 6.3. Adjustment for Multiple Telephone Lines and Deriving Annual Weights..... | 33 |
| 6.4. Calibration | 34 |
| 6.5. Adjustment for Provider Non-Response..... | 36 |
| 6.6. Sampling Weights for Territories | 39 |
| 7. Contents of the Public-Use Data File | 41 |
| 7.1. Section 1: ID, Weight, and Flag Variables | 44 |
| 7.2. Section 2: Household-Reported Vaccination and Chickenpox Information..... | 45 |

| | | |
|------------|--|-----------|
| 7.3. | Section 3: Demographic, Socio-Economic, and Other Household/Child Information..... | 45 |
| 7.4. | Section 4: Geographic Variables | 47 |
| 7.5. | Section 5: Number of Providers Identified and Consent Variables | 47 |
| 7.6. | Section 6: Number of Responding Providers Variables | 47 |
| 7.7. | Section 7: Characteristics of Providers Variables..... | 47 |
| 7.8. | Section 8: Provider-Reported Up-To-Date Vaccination Variables | 49 |
| | 7.8.1. Hib Up-To-Date Variables | 51 |
| | 7.8.2. Rotavirus Up-To-Date Variables..... | 55 |
| 7.9. | Section 9: Provider-Reported Age-At-Vaccination Variables..... | 56 |
| 7.10. | Section 10: Health Insurance Module Variables | 57 |
| 8. | Analytic and Reporting Guidelines | 60 |
| 8.1. | Use of NIS Sampling Weights..... | 60 |
| 8.2. | Estimation and Analysis | 62 |
| | 8.2.1. Estimating Vaccination Coverage Rates | 62 |
| | 8.2.2. Estimating Standard Errors of Vaccination Coverage Rates | 63 |
| 8.3. | Combining Multiple Years of NIS Data..... | 63 |
| | 8.3.1. Estimation of Multi-Year Means | 63 |
| | 8.3.2. Estimation of Multi-Year Contrasts..... | 67 |
| 9. | Summary Tables | 72 |
| 10. | Assessment of Total Survey Error..... | 73 |
| 11. | Limitations..... | 76 |
| 12. | Citations for NIS Data..... | 77 |
| 13. | References..... | 82 |

Appendices

| | |
|--|-----|
| Appendix A: Glossary of Abbreviations and Terms..... | 85 |
| Appendix B: Summary Statistics for Sampling Weights by Estimation Area..... | 88 |
| Appendix C: Flags for Inconsistent Values in the Breastfeeding Data..... | 92 |
| Appendix D: Programs for Estimation: Examples of the Use of SUDAAN, SAS, and R to Estimate Vaccination Coverage Rates and Their Standard Errors, and an Example of the Production of a Cross-Tabulation and Chart..... | 93 |
| Appendix E: Alphabetical Listing of Variables that are in the 2004-2015 Public-Use Data Files..... | 126 |
| Appendix F: Summary Tables | 164 |
| Appendix G: Trends in NIS Response Rates and Vaccination Coverage Rates, 1995-2015..... | 173 |
| Appendix H: Vaccine Type Codes..... | 180 |
| Appendix I: Key NIS Response Rates by Area | 181 |

List of Tables and Figures

| | | |
|------------|--|-----|
| Table 1: | Selected Operational Results of Q1/2015-Q4/2015 NIS Data Collection (Excluding territories)..... | 13 |
| Table 2: | Content of the Household Interview, National Immunization Survey, 2015 | 16 |
| Table 3: | Distribution of Age (in Days) at the Birth Dose of Hepatitis B Vaccine, National Immunization Survey, 2015 | 25 |
| Table 4: | Weighted Distribution of Children by Race/Ethnicity and Corresponding Combined Vaccine Series 4:3:1:3*:3:1:4, Pneumococcal, and Varicella Vaccination Coverage Estimates, National Immunization Survey, 2015 | 27 |
| Table 5: | NIS Variables Commonly Used in Analyses or for Published Estimates..... | 42 |
| Table 6: | Vaccine Categories and Vaccine Types, National Immunization Survey, 2015..... | 52 |
| Table 7: | Up-To-Date Variables for Hib, National Immunization Survey, 2009-2015..... | 54 |
| Table 8: | Up-To-Date Variables for Vaccine Series Including Hib, National Immunization Survey, 2009-2015 | 55 |
| Figure 1: | Question Flow for the Eight Health Insurance Questions Used to Create the Health Insurance Variables Included in the PUF | 59 |
| Table 9: | Summary of Weights and Stratum Variables, National Immunization Survey, 2015..... | 61 |
| Table 10: | Cross-Walk Between ITRUEIAP, ESTIAP, ESTIAP06-ESTIAP15, and Common Denominator Estimation Area (CDIAP), National Immunization Survey, 2015..... | 68 |
| Figure 2: | Comparison of Estimated Mean Total Error for 4:3:1:3:3:1 Series, ≥ 1 MMR, and ≥ 4 DTaP by Survey Year, National Immunization Survey, 2009-2012 | 74 |
| Figure 3: | Comparison of Estimated Mean Total Error for 4:3:1:3:3:1 Series, ≥ 1 MMR, and ≥ 4 DTaP, Excluding and Including the Under-Reporting Error Component, National Immunization Survey, 2012 | 75 |
| Table B.1: | Distribution of Dual-Frame Sampling Weights for Children with Completed Household Interviews, National Immunization Survey, 2015 | 88 |
| Table B.2: | Distribution of Dual-Frame Sampling Weights for Children with Adequate Provider Data, National Immunization Survey, 2015 | 90 |
| Table E.1 | Alphabetical Listing of Variables that are in the 2004-2015 Public-Use Data Files | 126 |
| Table F.1: | Estimated Population Totals and Sample Sizes of Children 19-35 Months of Age by State and Estimation Area, National Immunization Survey, 2015 | 164 |
| Table F.2: | Estimated Population Totals and Sample Sizes for Age Group by Maternal Education, National Immunization Survey, 2015 | 166 |
| Table F.3: | Estimated Population Totals and Sample Sizes for Age Group by Poverty Status, National Immunization Survey, 2015 | 167 |
| Table F.4: | Estimated Population Totals and Sample Sizes for Race/Ethnicity by Poverty Status, National Immunization Survey, 2015 | 168 |
| Table F.5: | Estimated Population Totals and Sample Sizes for Age Group by Race/Ethnicity, National Immunization Survey, 2015 | 169 |
| Table F.6: | Estimated Population Totals and Sample Sizes for Age Group by Gender, National Immunization Survey, 2015 | 170 |
| Table F.7: | Estimated Vaccination Coverage with Individual Vaccines and Selected Vaccination Series Among Children 19-35 Months of Age by State and Estimation Area, National Immunization Survey Q1/2015-Q4/2015..... | 171 |

| | | |
|-------------|---|-----|
| Table G.1: | Key Indicators from Landline Sample Household and Provider Data Collection by Survey Year, National Immunization Survey, 1995-2015 | 173 |
| Figure G.1: | Trends in Landline Sample Key Indicators from Household and Provider Data Collection by Survey Year, National Immunization Survey, 1995-2015 | 174 |
| Table G.2: | Key Indicators from Cell-Phone Sample Household and Provider Data Collection by Survey Year, National Immunization Survey, 2011-2015 | 175 |
| Figure G.2: | Trends in Cell-Phone Sample Key Indicators from Household and Provider Data Collection by Survey Year, National Immunization Survey, 2011-2015 | 175 |
| Table G.3: | CASRO Response Rate for the Combined Landline and Cell-Phone Samples by Survey Year, National Immunization Survey, 2011-2015 | 177 |
| Figure G.3: | Trend in CASRO Response Rate for the Combined Landline and Cell-Phone Samples by Survey Year, National Immunization Survey, 2011-2015 | 177 |
| Table G.4: | Vaccine-Specific Coverage Levels Among Children Age 19-35 Months in the United States by Survey Year, National Immunization Survey, 1995-2015 | 178 |
| Figure G.4: | Trends in Vaccine-Specific Coverage Levels among Children 19-35 Months of Age in the United States by Survey Year, National Immunization Survey, 1995-2015..... | 179 |
| Table H.1: | 2015 NIS Vaccine Type Codes | 180 |
| Table I.1: | Key Indicators for the Landline Sample by Estimation Area, National Immunization Survey, 2015..... | 181 |
| Table I.2: | Key Indicators for the Cell-Phone Sample by Estimation Area, National Immunization Survey, 2015 | 183 |
| Table I.3: | CASRO Response Rate for the Combined Landline and Cell-Phone Samples by Estimation Area, National Immunization Survey, 2015 | 185 |

Convention for Bolding Text

The Data User's Guide uses **bold** font to highlight substantive changes in the methodology or study design from last year's Guide.

This page intentionally blank.

1. Introduction

In 1992, the Childhood Immunization Initiative (CII) (CDC 1994) was established to 1) improve the delivery of vaccines to children; 2) reduce the cost of vaccines for parents; 3) enhance awareness, partnerships, and community participation; 4) improve vaccinations and their use; and 5) monitor vaccination coverage and occurrences of disease. Subsequently, the Healthy People 2020 objectives established the targets of having at least 90% of 2-year-old children fully vaccinated with most recommended vaccines (targets are 85% for HepA and the birth dose of HepB, and 80% for rotavirus) and 80% of 2-year-old children vaccinated with the basic immunization series. To fulfill the CII mandate of monitoring vaccination coverage and marking progress toward achieving those objectives, the National Immunization Survey (NIS) has been implemented by the National Center for Immunization and Respiratory Diseases (NCIRD) and the National Center for Health Statistics (NCHS) of the Centers for Disease Control and Prevention (CDC) since 1994.

The target population for the NIS is children aged 19 through 35 months living in households in the United States at the time of the interview. The official coverage estimates reported from the NIS are rates of being up-to-date with respect to the requisite numbers of doses of all recommended vaccines (Robinson 2016). These vaccines and their recommended numbers of doses are:

- diphtheria and tetanus toxoids and acellular pertussis vaccine, diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids vaccine (DTaP/DTP/DT) – 4 doses;
- poliovirus vaccine (polio) – 3 doses;
- measles, mumps, and rubella vaccine (MMR) – 1 dose;
- *Haemophilus influenzae* type b vaccine (Hib) – 3 or 4 doses depending on product type;
- hepatitis B vaccine (Hep B) – 3 doses;
- varicella zoster (chicken pox) vaccine (varicella) – 1 dose;

- pneumococcal conjugate vaccine (PCV) – 4 or 5 doses depending on vaccine type; (Infants and children who have received ≥ 1 dose of PCV7 should complete the immunization series with PCV13. A single supplemental dose of PCV13 is recommended for all children aged 14-59 months who have received 4 doses of PCV7 or another age-appropriate, complete PCV7 schedule.)
- hepatitis A vaccine (Hep A) – 2 doses;
- influenza vaccine; (For the recommended number of doses of influenza vaccine and other vaccines, see <http://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/index.html>.)
- rotavirus vaccine (RV) – 2 or 3 doses depending on product type.

In addition to these vaccines, interest focuses on the combined vaccine series 4:3:1:3*:3:1:4 (4+ DTaP/DTP/DT; 3+ polio; 1+ measles-containing vaccine (MCV); full series Hib, i.e., 3 or 4 doses depending on type of vaccine received; 3+ Hep B; 1+ varicella at or after 12 months of age; and 4+ PCV).

The NIS collects data on each of these vaccines. Varicella vaccine was added in Quarter 3, 1996, pneumococcal conjugate vaccine in Quarter 4, 2000, influenza vaccine and hepatitis A vaccine in Quarter 1, 2003, and rotavirus vaccine in Quarter 3, 2007. The remainder of the vaccines have been included in the NIS from its start in 1994. In October 2000, the Advisory Committee on Immunization Practices recommended that all children aged 2 to 23 months receive 4 doses of pneumococcal conjugate vaccine (CDC 2000). Influenza vaccine was recommended for children aged 6 to 23 months starting with the 2004-05 season (CDC 2003) and for all children ≥ 6 months starting in 2010.

The NIS uses random digit dialing (RDD) telephone survey methodology to identify households containing children in the target age range, and interviews are conducted with the adult who is most knowledgeable about the child's vaccinations. With consent of the child's parent or guardian, the NIS also contacts (by mail) the child's health care provider(s) to request information on vaccinations from the

child's medical records. Since 2005, NIS sampling, data collection, and weighting operations have been conducted by NORC at the University of Chicago.

Samples of telephone numbers are drawn independently, for each calendar quarter, within selected geographical areas, or strata. **In 2015, there are 61 geographic strata for which vaccination coverage levels can be estimated (see Table F.1), including 7 local areas; the remaining 54 estimation areas are either an entire state, the District of Columbia, a territory (the U.S. Virgin Islands, Guam, and Puerto Rico), or a “rest of state” area. For states with “rest of state” or local estimation areas, we can also produce estimates for the whole-state area. This design makes it possible to produce annual estimates of vaccination coverage levels for each state or territory, each “rest of state” area, the District of Columbia, and for each of the 7 local estimation areas with a specified degree of precision (a coefficient of variation of approximately 7.5%).** Further, by using the same data collection methodology and survey instruments in all estimation areas, the NIS produces comparable vaccination coverage levels among estimation areas and over time.

Data for the U.S. Virgin Islands and Guam are not included in the 2015 public-use data file to protect respondent confidentiality, as the sampling fractions were large in these small-population areas. Interested researchers can access data for the U.S. Virgin Islands and Guam by submitting a proposal and working through the NCHS Research Data Center. The link and guidelines for developing a proposal are located at the following URL: www.cdc.gov/rdc.

When the NIS was established in 1994, 78 areas were chosen for sampling strata, including the 50 states, 6 urban areas that receive federal Section 317 immunization grants (Bexar County, TX; Chicago, IL; District of Columbia; Houston, TX; New York City; Philadelphia County, PA), and 22 other urban areas. These areas were called “Immunization Action Plan” (IAP) areas in reference to plans developed to improve vaccination coverage following the resurgence of measles during 1989-1991. In 2005 and 2006, selected non-grantee IAP areas were “rotated off” (i.e., sample design no longer ensured adequate sample

size to produce estimates for the area), and replaced by new areas “rotated on” (i.e., sample design ensured adequate sample size to produce estimates for the area). Starting in 2007, the base NIS geographic strata included 56 areas (6 grantee urban areas and 50 state or “rest of state” areas). In addition, starting in 2007, state immunization programs could choose city/county areas of interest to have sample design that ensured adequate sample size to produce estimates for the area, using their grant funds. **In 2015, two additional area were chosen: El Paso County, TX, and Hidalgo County, TX. NIS data were also collected in the U.S. Virgin Islands, Guam, and Puerto Rico in 2015; as noted throughout this report, several of the sampling, data collection, and estimation procedures differed for these territories when compared to the rest of the United States, including the creation of separate survey weight variables for analysis that is to include territories.**

The $61 = 56 + 2 + 3$ (territories) areas are called *estimation areas*. Table 10 in Section 8 shows a cross-walk of estimation areas between years.

To maintain consistency with past NIS public-use data files, or PUFs, variable names and descriptions continue to use the term “IAP” to designate areas included as strata, which was the term used prior to 2008. The changing geographic strata over time will not cause a problem with bias in estimation of state and national coverage levels since the geographic strata are nested within state.

In 2015, the NIS utilized a dual-frame sampling design with independent samples drawn from landline and cell-phone sampling frames. The cell-phone component was added to the survey in 2011 in order to address the rapid rise of cell-phone-only households. **Preliminary results from the July-December 2015 National Health Interview Survey (NHIS) indicate that the number of households with only cell phones continues to increase. Approximately 57.7% of all children under 18 years of age—over 42 million children—live in households with access to only wireless telephones (Blumberg and Luke 2016). Several of the sampling, data collection, and estimation procedures differ for the cell-phone sample as compared to the landline sample, as noted throughout this report.**

For the 2015 NIS landline sample, the household interviews began on January 26, 2015 and ended on February 14, 2016. For the 2015 NIS cell phone sample, the household interviews began on January 26, 2015 and ended on February 15, 2016. Provider data collection extended from February 2015 to April 2016 for both sample sources. A total sample (including sample from territories) of approximately 14.5 million telephone numbers (6.3 million landline and 8.2 million cell-phone) yielded household interviews for 28,639 children (4,873 landline and 23,766 cell-phone), 16,000 of whom (2,898 landline and 13,102 cell-phone) had adequate provider data (provider-reported vaccination data adequate to determine whether the child was up-to-date with respect to the recommended immunization schedule). The 2015 NIS public-use data file (which does not include data for the U.S. Virgin Islands and Guam) contains data for the 27,592 children with completed household interviews, and more extensive data (e.g. provider-reported vaccinations and facility data) for the 15,391 children with adequate provider data (including 180 unvaccinated children).

In 2012, to reduce the length of the household interview, decrease respondent burden, and potentially improve response rates, the NIS household questionnaire was modified. Official NIS vaccination rate estimates are based on the provider-reported vaccination histories for each child. Among children with data received from vaccination providers identified in the household interview, it must be determined which children have “adequate provider data,” that is, which children have provider data adequate to determine whether the child is up to date with respect to the recommended immunization schedule. Beginning in 2012, questions that were previously used to define adequate provider data were no longer available. With this questionnaire change, it was no longer possible to use the same definition of adequate provider data as was used prior to 2012, and so beginning in 2012 all children with any provider-reported vaccination data are considered to have adequate provider data. See the user’s guide for the 2014 NIS public-use data file (NCHS 2015) for more detail about this change and its impact.

The weights included in this public-use data file afford the data analyst the capability of conducting several different types of analyses, depending on interests and aims. One can choose to analyze all children with completed household interviews or only the subset of children for whom the provider-reported data are adequate. Also, one can choose to include or exclude children who reside in territories in the analysis. Previous NIS public-use data files have also provided analysts with these capabilities.

The 2015 NIS public-use data file includes only dual-frame weights. Dual-frame estimates are the best estimates for the NIS in terms of minimizing any bias due to the incompleteness of the landline sampling frame. Section 6 of this user's guide provides information about the creation of the weight variables included in the 2015 NIS public-use data file, and Section 8 provides guidance for their use.

Published tables of vaccination coverage estimates for 2015 will be available on the National Center for Immunization and Respiratory Diseases website, <http://www.cdc.gov/vaccines/imz-managers/coverage/nis/child/index.html>. An article summarizing key findings from the NIS data, as published in the *Morbidity and Mortality Weekly Report (MMWR)*, will also be available on this website. The accompanying codebook (NCIRD 2016) documents the contents of the 2015 NIS public-use data file. For reference, Appendix E (Alphabetical Listing of Variables in the 2004-2015 Public-Use Data Files) provides a full list of variables in the 2015 and previous NIS public-use data files.

Additional information on the NIS is available at: <http://www.cdc.gov/vaccines/imz-managers/nis/about.html>.

For additional information on the NIS public-use data file, please contact the NCIRD Information Dissemination Staff:

Information Dissemination Staff, NCIRD
1600 Clifton Road
Atlanta, GA 30333
E-mail: cdcinfo@cdc.gov
Internet: <http://www.cdc.gov/vaccines/nis>

2. Sample Design

The NIS uses two phases of data collection to obtain vaccination information for a large national probability sample of young children: an RDD telephone survey designed to identify households with children 19 through 35 months of age, followed by the Provider Record Check, a mailed survey to children's vaccination providers. This section summarizes these two phases of data collection. Other descriptions of the sample design are given by Ezzati-Rice et al. (1995), Zell et al. (2000), and Smith et al. (2001a, 2005).

2.1. The NIS RDD Telephone Survey

The NIS RDD telephone survey phase uses independent, quarterly samples of telephone numbers. Sampling frames were provided by Marketing Systems Group (MSG). Landline telephone and cell-phone numbers were sampled within estimation areas in each quarter of 2015. **Table F.1 (in Appendix F) lists the 61 estimation areas for the 2015 NIS by state and shows the estimated number of children living in each state and estimation area in 2015.**

The NIS uses the list-assisted method of RDD (Lepkowski 1988) to sample landline telephone numbers. This method selects a random sample of telephone numbers from "banks" of 100 consecutive telephone numbers (e.g., 773-256-0000 to 773-256-0099) that contain at least one directory-listed residential telephone number. Because directory listings are not available for cell phones, the NIS cell-phone sample did not use the list-assisted method of RDD, but rather used RDD without list-assistance. That is, the cell-phone sample was selected from all banks of cell-phone numbers, not just those containing at least one directory-listed residential telephone number. **Directory listings were also unavailable for landline sample in the U.S. Virgin Islands, Guam, and Puerto Rico, so the landline and cell-phone samples for these areas were selected without list-assistance using simple random sampling.**

The target sample size of completed telephone interviews in each estimation area is designed to achieve an approximately equal coefficient of variation of 7.5% for an estimator of vaccination coverage derived

from provider-reported vaccination histories, given a true coverage parameter of 50%. Landline telephone and cell-phone sample sizes were chosen such that the two samples combined meet the target coefficient of variation of 7.5%.

In 2015, including the U.S. territory samples, 55.9% of children (59.5% of landline sample children and 55.1% of cell-phone sample children) with a completed household interview were determined to have adequate provider data. Excluding territories, this proportion was 56.2% (59.7% for the landline sample and 55.5% for the cell-phone sample). The percentage of children with adequate provider data varies among the non-territory estimation areas (from 48.8% in New Jersey to 68.9% in Hidalgo County, TX). Among the U.S. territories, the percentages were 56.4% in the U.S. Virgin Islands, 60.4% in Guam, and 36.3% in Puerto Rico. The phrase “adequate provider data” means that sufficient vaccination history information was obtained from the provider(s) to determine whether the child is up-to-date with respect to the recommended vaccination schedule. Starting with the 2002 NIS public-use data file, the definition of children with adequate provider data was expanded to include unvaccinated children. These are children for whom either (1) the respondent reported during the household interview that the child had received no vaccinations and has no providers, or (2) the respondent reported during the household interview that the child had received no vaccinations but has one or more providers, and those providers all reported administering no vaccinations. An NCHS Series 2 Report on the statistical methodology of the NIS (Smith et al. 2005) includes details of how unvaccinated children are included in the estimates of vaccine coverage. This report can be viewed at http://www.cdc.gov/nchs/data/series/sr_02/sr02_138.pdf. This modification to the NIS produces only small changes in vaccination coverage for estimation areas and states, because the number of unvaccinated children in the sample is very small (only 186 in 2015). As described in the introduction, the definition of adequate provider was modified in 2012 to include all children with provider-reported vaccination data, plus unvaccinated children.

The design and implementation of the NIS landline sample involves four procedures. First, statistical models predict the number of sample telephone numbers needed in each estimation area to meet the target precision requirements. Second, the sample for an estimation area is divided into random sub-samples called replicates. By releasing replicates as needed, it is possible to spread the interviews for each sampling area evenly across the entire calendar quarter. Third, an automated procedure eliminates a portion of the non-working and non-residential telephone numbers, plus numbers on the NIS do-not-call list, from the sample before the interviewers dial them. Fourth, the sample telephone numbers are matched against a national database of residential telephone numbers in order to obtain usable mailing addresses for as many sample households as possible. To promote participation in the NIS, an advance letter is sent to these addresses approximately two weeks prior to calling to conduct the household interview.

The design and implementation of the cell-phone sample differs from that of the landline sample in two ways:

- Prior to 2014, there was no process to remove non-working and non-residential cell-phone numbers before dialing them. Beginning in 2014 and continuing in 2015, an automated process was implemented to remove cell-phone numbers flagged as having no recent activity and that were therefore very likely to be non-working cell phones.
- Cell-phone numbers were not matched to an external database to obtain mailing addresses. Cell-phone sample cases were not sent advance letters.

2.2. The NIS Provider Record Check

At the end of the household interview, consent to contact the child's vaccination provider(s) is requested from the parent/guardian. When oral consent is obtained, each provider is mailed an immunization history questionnaire. This mail survey portion of the NIS is the Provider Record Check (PRC). The Provider Record Check is conducted in the same manner for both landline and cell-phone sample cases.

The instructions ask vaccination providers to mail or fax the immunization history questionnaire back upon completion. Two weeks after the initial mailing, a telephone call is made to providers who have still not responded, to remind and encourage them to complete the form and either mail or fax the information back. In some instances, provider-reported vaccination histories are completed over the telephone. The data from the questionnaires are edited, entered, cleaned, and merged with the household information from the RDD survey to produce a child level record.

2.3. Summary of Data Collection

Table 1 presents selected operational results of NIS data collection for calendar year 2015. To facilitate comparisons with prior years, the numbers in Table 1 are presented separately for the landline and cell-phone samples, and exclude the U.S. territory samples. **Children aged 19 through 35 months during 2015 data collection were born between January 2012 and May 2014.**

The landline RDD sample (in replicates that were released for use) consisted of 5,990,464 telephone numbers. Of those, 3,483,747 were eliminated before release to the telephone centers by the automated procedure as non-working numbers, non-residential numbers, or numbers on the NIS do-not-call list. The remaining 2,506,717 numbers were sent to the telephone centers to be dialed, and 458,086 households were identified, as shown in Rows C and F. Among the identified households, 412,023 (89.9%) were successfully screened. Of these, 5,462 (1.3%) contained one or more age-eligible children. Among these households, 4,387 (80.3%) completed the household interview.

The cell-phone sample (in replicates that were released for use) consisted of 7,867,079 telephone numbers. Of those, 1,994,640 were eliminated before release to the telephone centers by the automated procedure as inactive cell phones or numbers on the NIS do-not-call list. The remaining 5,872,439 numbers were sent to the telephone centers to be dialed, and 1,143,468 active personal cell-phone numbers (APCNs) were identified, as shown in Row F. Among the identified APCNs,

906,520 (79.3%) were successfully screened. Of these, 30,246 (3.3%) were deemed eligible for the NIS interview. Among the identified eligible respondents, 21,838 (72.2%) completed the interview.

A standard approach for measuring response rates in telephone surveys has been defined by the Council of American Survey Research Organizations (CASRO 1982). The CASRO response rate is equivalent to “RR3” of AAPOR Standard Definitions (AAPOR 2015). **In 2015, the CASRO response rate (Row J, Table 1) for the landline sample was 59.1%. The CASRO response rate equals the product of the resolution rate (81.9%, Row E), the screening completion rate (89.9%, Row G), and the interview completion rate among eligible households (80.3%, Row I).** The resolution rate is the percentage of the total telephone numbers selected that are classifiable as non-working, non-residential, or residential. The screening completion rate is the percentage of known households that are successfully screened for the presence of age-eligible children. The interview completion rate is the percentage of households with one or more age-eligible children who complete the household interview.

The CASRO response rate (Row J) for the cell-phone sample in 2015 was 32.2%. As with the landline sample, it equals the product of the resolution rate (56.3%, Row E), the screening completion rate (79.3%, Row G), and the interview completion rate among eligible households (72.2%, Row I).

The CASRO response rate (Row J) for the combined landline and cell-phone sample was 34.9% in 2015. See footnote 6 of Table 1 for a description of the calculation of the combined CASRO response rate.

Row K of Table 1 shows that household interviews were completed on behalf of 4,522 age-eligible children in the landline sample and 22,453 children in the cell-phone sample. Rows L through O give results for the Provider Record Check phase. Specifically, Row L gives the rate of obtaining oral consent from household respondents to contact their children’s vaccination providers – 67.0% for landline sample cases and 63.4% for cell-phone sample cases in 2015.

The number of immunization history questionnaires mailed to vaccination providers exceeds the number of completed interviews for children with consent because some children have more than one vaccination provider. **Of the questionnaires mailed to providers of children from the landline sample, 3,489 (94.7%, Row N) were returned. Among the children with completed household interviews, 2,700 (59.7%, Row O) had adequate vaccination histories based on provider reporting (2,671) or were determined to be unvaccinated (29). The other 40.3% of children lacked adequate provider data for a variety of reasons, such as the parent did not give consent to contact the child's provider(s), the provider(s) did not have records for the child, or the provider(s) did not report the vaccination history.**

Of the questionnaires mailed to providers of children from the cell-phone sample, 16,653 (94.4%, Row N) were returned. Among the cell-phone sample children with completed household interviews, 12,467 (55.5%, Row O) had adequate vaccination histories based on provider reporting (12,318) or had no vaccinations based on household reporting (149). The other 44.5% of children lacked adequate provider data for a variety of reasons, such as the parent did not give consent to contact the child's provider(s), or the provider(s) did not have medical records for the child.

In 2015, data from the Health Insurance Module (HIM) were collected. Among the 4,522 children in the landline sample with completed household interviews, 3,149 (69.6%, Row P) completed the HIM. Among the 22,453 children in the cell-phone sample with completed household interviews, 14,620 (65.1%, Row P) completed the HIM.

For each estimation area and each state, Table F.1 (see Appendix F) shows the number of children with completed household interviews and the number of children with adequate provider data.

Table 1: Selected Operational Results of Q1/2015-Q4/2015 NIS Data Collection (Excluding territories)

| Row | Key Indicator | Landline Sample | | Cell-Phone Sample | | Combined Samples | | Formula |
|------------------------|--|--|---------|--|---------|--|---------|--------------------|
| | | Number | Percent | Number | Percent | Number | Percent | |
| Household Phase | | | | | | | | |
| A | Total Selected Telephone Numbers in Released Replicates | 5,990,464 | -- | 7,867,079 | -- | 13,857,543 | -- | |
| B | Phone Numbers Resolved before Computer-Assisted Telephone Interviewing | 3,483,747 | 58.2% | 1,994,640 | 25.4% | 5,478,387 | 39.5% | B/A |
| C | Total Phone Numbers Released to Telephone Centers | 2,506,717 | -- | 5,872,439 | -- | 8,379,156 | | A-B |
| D | Advance Letters Mailed | 986,263 | 39.3% | 0 | 0.0% | 986,263 | 11.8% | D/C |
| E | Resolved Phone Numbers ¹ – Resolution Rate | 4,904,614 | 81.9% | 4,430,414 | 56.3% | 9,335,028 | 67.4% | E/A |
| F | Households Identified – WRN/APCN Rate ² | 458,086 | 9.3% | 1,143,468 | 25.8% | 1,601,554 | 17.2% | F/E |
| G | Households Successfully Screened ³ – Screener Completion Rate | 412,023 | 89.9% | 906,520 | 79.3% | 1,318,543 | 82.3% | G/F |
| H | Eligible Households – Eligibility Rate ⁴ | 5,462 | 1.3% | 30,246 | 3.3% | 35,708 | 2.7% | H/G |
| I | Households with Completed Household Interviews – Interview Completion Rate | 4,387 | 80.3% | 21,838 | 72.2% | 26,225 | 73.4% | I/H |
| J | CASRO Response Rate ⁵ | -- | 59.1% | -- | 32.2% | | 34.9% | E*G*I ⁶ |
| K | Age-Eligible Children with Completed Household Interviews ⁷ | 4,522 | -- | 22,453 | -- | 26,975 | -- | |
| Provider Phase | | | | | | | | |
| L | Children with Consent to Contact Vaccination Providers | 3,028 | 67.0% | 14,243 | 63.4% | 17,271 | 64.0% | L/K |
| M | Immunization History Questionnaires Mailed to Providers | 3,685 | -- | 17,636 | -- | 21,321 | -- | |
| N | Immunization History Questionnaires Returned from Providers | 3,489 | 94.7% | 16,653 | 94.4% | 20,142 | 94.5% | N/M |
| O | Children with Adequate Provider Data | 2,700 (includes 29 unvaccinated children) | 59.7% | 12,467 (includes 149 unvaccinated children) | 55.5% | 15,167 (includes 178 unvaccinated children) | 56.2% | O/K |
| Modules | | | | | | | | |
| P | Age-Eligible Children with Completed Household Interview and Completed Health Insurance Module | 3,149 | 69.6% | 14,620 | 65.1% | 17,769 | 65.9% | P/K |

¹ Includes phone numbers resolved before CATI (Row 2).

² For the landline sample, this is the working residential number (WRN) rate; for the cell-phone sample, it is the active personal cell-phone number (APCN) rate.

³ For the landline sample, this is the age-eligibility screener; for the cell-phone sample, it is a combination of the screener for non-minor-only cell phone status and the age-eligibility screener.

⁴ For the landline sample, this is the age-eligibility rate; for the cell-phone sample, it reflects a combination of the non-minor-only cell-phone rate and the age-eligibility rate.

⁵ CASRO, Council of American Survey Research Organizations.

⁶ The response rate is the number of households with a completed household interview divided by the estimated number of eligible households in the sample. Within each sample type (landline or cell phone), the number of eligible households was estimated using the CASRO assumptions; these assumptions are that the rate of households among the unresolved telephone numbers is the same as the observed rate of households among the resolved telephone numbers, and the rate of eligible households among unscreened households is the same as the observed rate of eligible households among screened households. Under these assumptions, within each sample type the CASRO response rate is equal to the product of the resolution rate, the screener completion rate, and the interview completion rate. For the combined samples, we have defined the CASRO response rate as the total number of households with a completed interview divided by the estimated total number of eligible households across both sample types, where the estimated total number of eligible households is equal to the sum of the estimated number of eligible households in the landline sample (using CASRO assumptions) and the estimated number of eligible households in the cell-phone sample (using CASRO assumptions).

⁷ Rows K-P reflect the removal of children with an ineligible best date of birth.

2.4. Informed Consent, Security, and Confidentiality of Information

The advance letter, introduction to the telephone survey, and oral consent assure the respondent of the confidentiality of his/her responses and the voluntary nature of the survey. Informed consent is obtained from the person in the household most knowledgeable about the eligible child's vaccination history (generally the parent or guardian of the child). Informed consent to contact the child's vaccination provider(s) is obtained at the end of the interview.

Information in the NIS is collected and processed under high security. To ensure privacy of the respondents and confidentiality of sensitive information, NCHS and NCIRD have established standards for release of data from this survey. All CDC staff and contractor staff involved with the NIS sign the NCHS and NCIRD confidentiality agreement and follow instructions to prevent disclosure.

All information in the NIS is collected under strict confidentiality and can be used only for research [Section 308(d) of the Public Health Service Act, 42 U.S. Code 242m(d) and the Privacy Act of 1974 (5 U.S. Code 552a)]. Prior to public release, the contents of the public-use data file go through extensive review by the NCHS Disclosure Review Board to protect participant privacy as well as data confidentiality.

3. Content of NIS Questionnaires

This section describes the questionnaires used in the 2015 NIS telephone interview of households and in the NIS Provider Record Check.

3.1. Content of the Household Questionnaire

The computer-assisted telephone interview (CATI) questionnaire used in the RDD phase of NIS data collection consists of two parts: a screener to identify households with children aged 19 through 35 months and an interview portion. The questionnaire is modeled on the Immunization Supplement to the National Health Interview Survey (NHIS) (NCHS 1999). The NIS CATI questionnaire has been translated into Spanish, and Language Line Services (formerly part of AT&T) is used for real-time translation into many other languages (Wall et al. 1995). Table 2 summarizes the content of each section of the NIS household interview. The CATI questionnaire is available at <http://www.cdc.gov/vaccines/imz-managers/nis/datasets.html>.

In the screener, the purpose of the survey is explained to the respondent, and the household is screened to determine whether it contains any children aged 19 through 35 months (any child who was or would be aged 19 through 35 months during the calendar quarter is eligible). If the household has an eligible child, the respondent is asked whether he/she is the most knowledgeable person for the child's vaccination history. If the respondent indicates that another person in the household is more knowledgeable, the interviewer asks to speak to him/her at that time. If that person is unavailable to be interviewed, the interview proceeds to Section MR, the name of the most knowledgeable person is recorded, and a "callback" is scheduled for a later date. For the cell-phone sample, prior to screening for age-eligibility, the household is screened to ensure that the cell-phone is used by an adult (i.e., to ensure it is not a minor-only cell phone). If the household has more than one age-eligible child, data are collected for each eligible child.

Table 2: Content of the Household Interview, National Immunization Survey, 2015

| Questionnaire Section | Content of Section |
|----------------------------------|---|
| Section S | Screening questions to determine eligibility, roster of eligible children, availability of shot records |
| Section MR | Most-knowledgeable-respondent callback questions |
| Section A (Guam only) | Vaccination history (asked if shot records are available) |
| Section B | Ever vaccinated and flu vaccination questions |
| Section C | Demographic and socioeconomic questions |
| Section D | Provider information and request for consent to contact the eligible child's vaccination provider(s) |
| Section E | Health Insurance Module (HIM) |

Prior to Q1/2012, the person being interviewed was asked during the screener section whether he/she had a written record (shot card) of the child's vaccination history, and whether it was easily accessible. If a shot card was available, the respondent was asked to provide information directly from it in Section A. However, beginning in Q1/2012, Section A and most of Section B were eliminated from the regular questionnaire, and therefore all interviews proceeded directly to a reduced form of Section B, which asks the respondent to recall information about the child's influenza vaccinations. **In 2015, Section A was reinstated for Guam respondents. As described above in the Introduction, Guam and U.S. Virgin Islands data are not included in the 2015 public-use data file.**

Section C obtains information that includes relationship of respondent to the child, race and Hispanic origin of the child, household income, educational attainment of the mother, and other information on the socioeconomic characteristics of the household and its eligible children. This section is asked of all respondents upon completion of Section A or Section B.

In Section D of the NIS household interview, identifying information (such as name, address, and telephone number) for the child's vaccination provider(s) is requested, as well as the full names of the

child(ren) and the respondent, so that NIS personnel can contact the provider(s) and identify the child(ren) whose immunization information the NIS is requesting. After this information is obtained, consent to contact the child's vaccination provider(s) is requested. When oral consent and sufficient identifying information are obtained, the immunization history questionnaire is mailed to the child's vaccination provider(s).

Beginning in 2006, a Health Insurance Module (HIM) was administered upon completion of Section D to collect data regarding the types of medical insurance coverage the child has had since birth. If a respondent provided consent to contact medical providers and completed Section D, he/she flowed directly into the HIM. If, however, consent or any other critical provider question was refused, the call was terminated; only upon callback on which consent was granted or a second refusal given within Section D was the respondent asked the HIM. See Section 7.10 of this user's guide for information on the HIM variables included on the public-use data file.

Some changes were made to the NIS questionnaire during 2015. These are listed below.

Year references for income questions were updated to refer to the previous year, 2014. That is, the question text at CFAMINC, C13_DON'T_KNOW, and C13_REFUSED was updated to ask about 2014 income rather than 2013.

Changes and additions were made throughout the year to the questions about flu vaccinations in Section B; the flu question responses are not included on the PUF, and so the PUF contents are not affected.

“Living with partner” was added as a response option to question C7 beginning in Q1/2015.

Data were collected in the U.S. Virgin Islands, Guam, and Puerto Rico in 2015 (landline and cell-phone sample). Section A was added and administered to Guam sample respondents that had shot

records available. A few minor changes were made to the instrument for the Puerto Rico sample, such as adding a reference to *municipio* of residence.

3.2. Content of the Immunization History Questionnaire (IHQ)

The immunization history questionnaire mailed to the vaccination providers is designed to be simple and brief, to minimize provider burden and encourage survey participation. The structure and content of this form were initially derived from the National Immunization Provider Record Check Study (NHIS/NIPRCS), which collected and reconciled vaccination data from the providers of respondents to the Immunization Supplement to the National Health Interview Survey. The immunization history questionnaire consists of two double-sided pages. Page 1 includes space for a label that gives the child's name, date of birth, and gender. The remainder of page 1 contains questions about the facility and vaccination provider. Page 2 gives instructions for filling out the shot grid, which appears on page 3. Page 4 thanks the vaccination provider for providing the information, and lists websites and telephone numbers that can be used to obtain more information about the NIS and the National Center for Immunization and Respiratory Diseases. The Immunization History Questionnaire is available at <http://www.cdc.gov/vaccines/imz-managers/nis/datasets.html>.

No changes were made to the English version of the NIS IHQ during 2015. For Puerto Rico, a Spanish-translated version of the NIS IHQ was used and this version differed slightly from the IHQ used in 2015 for other estimation areas in a few ways. Question 5b, which asks if the practice has been deputized to administer Vaccines for Children vaccines to underinsured children, was removed from the Puerto Rico IHQ, and the response options for Indian Health Service and Pharmacy at Question 5c were removed. These changes were made to the Puerto Rico IHQ because the material was not relevant to Puerto Rico.

4. Data Preparation and Processing Procedures

The household and provider data collection in the NIS incorporate extensive data preparation and processing procedures. During the household interview, the CATI system supports reconciliation of critical errors as interviewers enter the data. After completion of interviewing for a quarter, post-CATI editing and data cleaning produce a final interview data file. The editing of the provider data begins with a manual review of returned immunization history questionnaires, data entry of the questionnaires, and cleaning of the provider data file. After the provider data are merged with the household interview data and responses from multiple providers for a child are consolidated into a child-level data record, the editing continues. A quality assurance check is performed, from all sources of the date-of-birth information, to ensure that the provider completed the questionnaire for the correct child and to confirm age-eligibility. Editing of the provider-reported vaccination dates then attempts to resolve specific types of discrepancies in the provider data. The end product is an analytic file containing household and provider data for use in estimating vaccination coverage.

4.1. Data Preparation

The editing and cleaning of NIS data involve several steps. First, the CATI system enables interviewers to reconcile potential errors while the respondent is on the telephone. Further cleaning and editing take place in a post-CATI clean-up stage, involving an extensive review of data values, cross tabulations, and the recoding of verbatim responses for race and ethnicity. The next step involves the creation of numerous composite variables. Provider data are cleaned in a separate step. After these steps have been completed, imputations are performed for item non-response on selected variables, and weights are calculated. The procedures and rules of the National Health Interview Survey serve as the standard in all stages of data editing and cleaning (<http://www.cdc.gov/nchs/nhis.htm>).

4.1.1. *Editing in the CATI System*

The CATI software checks consistency across data elements and does not allow interviewers to enter invalid values. Catching potential errors early increases the efficiency of post-survey data cleaning and processing.

To prevent an overly complicated CATI system, out-of-range and inconsistent responses produce a warning screen, allowing the interviewer to correct real time errors. This allows the interviewer to reconcile errors while respondent is on the telephone. CATI warning screens focus on items critical to the survey, such as those that determine a child's eligibility (e.g., date of birth).

A CATI system cannot simultaneously incorporate every possible type of error check and maximize system performance. To reconcile this trade-off, post-CATI edits are used to resolve problems that do not require access to the respondent, as well as unanticipated logic problems that appear in the data.

4.1.2. *Post-CATI Edits*

The post-CATI editing process produces final, cleaned data files for each quarter. The steps in this process, implemented after all data collection activities for a quarter are completed, are described below.

Initial Post-CATI Edits and File Creation

After completion of interviewing each quarter, the raw data are extracted from the CATI data system and used to create two files: the sample file and the interview data file. The sample file contains one record for each sample telephone number and summary information for telephone numbers and households. The interview data file contains one record for each eligible sample child and all data reported for the child during the household survey.

Following creation of these two files, a preliminary analysis of each file identifies out-of-range values and extraneous codes. The first check verifies the eligibility status of children. Once the required corrections are verified, invalid values are replaced with either an appropriate data value or a missing value code.

Frequency Review

After the pre-programmed edits are run, frequency distributions of all variables in each file are produced and reviewed. Each variable's range of values is examined for any invalid values or unusual distributions. If blank values exist for a variable, they are checked to see whether they are allowable and whether they occur in excessive numbers. Any problems are investigated and corrected as appropriate.

File Crosschecks

Crosscheck programs ensure that cases exist across files in a consistent manner. Specifically, checks ensure that each case in the interview data file is also present in the sample file and that each case in the sample file was released to the telephone center. Checks also ensure that no duplicate households exist in the sample file and no duplicate children exist in the interview data file.

When all checks have been performed, the final quarterly interview data file is created. Programmers and statisticians then create composite variables constructed from basic variables for each child. Sampling weights (described in Section 6 of this Guide) are added to each record.

4.1.3. *Editing of Provider Data*

Six to eight weeks after the close of household data collection for a quarter, the majority of the immunization history questionnaires have been collected from providers. The data from the hard-copy questionnaires are entered and independently re-entered to provide 100% verification. The provider data file is cleaned, in a similar fashion to the household data file, for out-of-range values and consistency. A computer program back-codes "other shot" verbatim responses into the proper vaccine category (e.g., Engerix B counts as Hep B, and Tetramune counts as DTP and Hib). These translations come from a file that contains all such verbatim responses ever encountered in the NIS. Also, the provider data file is checked for duplicate records, and exact duplicates are removed. If the provider data contain a date of birth, gender, or name for the child that differs from the household interview for that child, the questionnaire is re-examined to see whether it may have been filled out for the incorrect child. Provider data that appear to have been filled out for the wrong child are removed from the provider database.

When a child has data from multiple providers, decision rules are applied to produce the most complete picture of the child's vaccination history.

Once these data have been cleaned, they are combined with the household data file. Information from up to five providers can be added to a child's record. If more than one provider reported vaccination data for the child, the data from the multiple provider reports are combined into a single history for the child, called the "synthesized provider-reported vaccination history." The determination of whether the child is up-to-date for recommended vaccines and vaccine series is based on the child's synthesized provider-reported vaccination history.

Many variables in the household data file are checked against or verified with the provider data file. For example, a child's date of birth as recorded by the provider is checked against the date of birth as given by the household, to verify that the provider was reporting for that specific child and to form a "best" date of birth for the child. All children with at least one provider-reported vaccination are considered to have adequate provider data.

4.2. Limitations of Data Editing Procedures

Although data editing procedures were used for the NIS, the data user should be aware that some inconsistent data might remain in the public-use data file. The variables that indicate whether a child is up-to-date on each vaccine or series (on which the estimates of vaccination coverage are based) are derived from provider-reported data, and the NIS does not re-contact households or providers to attempt to reconcile potential discrepancies in provider-reported vaccination dates or to resolve date-of-birth reporting errors. However, beginning with the 1999 NIS, the provider-reported data are manually reviewed and edited to correct specific reporting errors. The *National Immunization Survey: Guide to Quality Control Procedures* (CDC 2002) discusses the change in editing procedures in more detail. Some children with adequate provider data may have incomplete vaccination histories. These incomplete histories arise from three primary sources: 1) the household does not identify all vaccination providers, 2)

some but not all providers respond with vaccination data, and 3) all identified providers respond with vaccination data but fail to list all the vaccinations in the child's medical record. Even with these limitations, the NIS overall is a rich source of data for assessment of up-to-date status and age-appropriate vaccination. Also, NIS is the only source to provide comparable provider-reported vaccination data across states and local areas in the United States.

4.3. Variable-Naming Conventions

The names of variables follow a systematic pattern as much as possible. The codebook for the public-use data file groups the variables into ten broad categories according to the source of the data (household or provider) and the content of the variable (NCIRD 2016). See Section 7 of this report for detailed information on the contents of the public-use data file.

4.4. Missing Value Codes

Missing value codes for each variable can be found in the codebook (NCIRD 2016). For household variables, the missing value codes usually are 77 for DON'T KNOW and 99 for REFUSED. Some household variables may also contain blanks, if the question was not asked. The variables developed from the immunization history questionnaire generally do not have specific missing value codes.

4.5. Imputation for Item Non-Response

The NIS uses imputation primarily to replace missing values in the socioeconomic and demographic variables used in weighting. Missing values of these variables are imputed for all children with a completed household interview – i.e., all children appearing on the public-use data file. A sequential hot-deck method is used to assign imputed values (Ford 1983). Class variables are used to separate respondents into cells. Donors and recipients must agree on the categories of the class variables, which include the estimation area. Within the categories of the class variables, respondents are sorted by variables related to the variable to be imputed. The last case with an observed value is used as the donor for up to four recipients. The “Notes” line for each variable in the codebook (NCIRD 2016) identifies

variables that contain imputed values. These variables include the gender, Hispanic origin, race, and first-born status of the child, and the education level, age group, marital status, and mobility status of the mother.

The count of vaccinations for a specific vaccine is based on the number of unique vaccination *dates* reported by the child’s provider(s). In filling out the immunization history questionnaire a provider may not know the date of the first dose of hepatitis B, which is typically given at birth. The provider does, however, have the option of checking the “Given at Birth” box for the first dose of hepatitis B. If it was checked “yes” and the date of the birth dose of hepatitis B was not reported, a program assigns the date of the birth dose for this vaccine. A value is imputed from the distribution of provider-reported dates for the birth dose of hepatitis B. The birth dose for this imputation is defined as being given in the first 7 days of life--between the date of birth (i.e., 0 days) and the date of birth plus 6 days. This imputation procedure was first implemented for Quarter 1, 2000 – Quarter 4, 2000. **For 2015 (excluding territories), a total of 36 children had the date of the birth dose of hepatitis B assigned using the above procedure (see HEP_FLAG).**

Table 3 shows the distribution of age in days at the birth dose of hepatitis B for children in 2015 with a provider-reported birth dose. A similar table is included in the 2000-2014 data user’s guides. For 1997, 1998, and 1999, Section 5 of the data user’s guide provides information on the distribution of age in days for the birth dose of hepatitis B vaccine and gives guidance on imputing age in days at birth dose for children with a missing date, but for whom the provider checked the box indicating that a dose was administered at birth (see HEP_BRTH).

Table 3: Distribution of Age (in Days) at the Birth Dose of Hepatitis B Vaccine, National Immunization Survey, 2015

| Age in Days at Birth Dose | Unweighted Percentage Of Birth Doses* |
|----------------------------------|--|
| 0 | 58.3 |
| 1 | 26.9 |
| 2 | 9.0 |
| 3 | 2.4 |
| 4 | 1.5 |
| 5 | 0.9 |
| 6+ | 1.0 |

* Excludes territories.

4.6. Vaccine-Specific Recoding of Verbatim Responses

On the IHQ, providers can list vaccinations in the “other” section of the IHQ shot grid. After data collection, they are reclassified into the listed categories, if possible, using a vaccination recoding table. This table is reviewed by National Center for Immunization and Respiratory Diseases personnel to ensure the shots are recoded into the appropriate category or categories (for combination shots).

4.7. Composite Variables

A number of composite variables (constructed from basic variables) are created and included in the NIS public-use data file. Composite variables assist users and data analysts by eliminating duplication of effort and making NIS data easier to use.

Since the initial years of NIS data collection, the household composite variables have included up-to-date status on individual vaccinations, race of child, household income, and up-to-date status on several vaccination series. Many of these household composite variables are included in the NIS public-use data file. See Section 7 of this report for information on the key variables that are included.

In Quarter 3, 1999, the NIS race questions (see questions C3, C9 and C10 in the household questionnaire) were expanded to include Alaska Native, Native Hawaiian, and Pacific Islander, implementing the revised Office of Management and Budget (OMB) standards for classification of race and ethnicity (http://www.whitehouse.gov/omb/fedreg_1997standards). The composite race variables in the 2002 through present NIS public-use data files, however, contain only three categories: white alone; black alone; and all other races alone/multiple races. (The variable RACE_K classifies each child into one of these three categories, while the variable RACEETHK includes a separate “Hispanic” category.) The “all other races alone” category includes Asian, American Indian or Alaska Native, Native Hawaiian or Pacific Islander, and other races. If more than one race was selected during administration of the child race questions, the child is classified as multi-racial. Because of small sample sizes and risk of disclosure within estimation areas, the 2002 through present NIS public-use data files do not contain any variables with separate multiple-race categories. Rather, the multi-racial children are included in the “all other races” category. Table 4 shows some characteristics of the current race/ethnicity categories.

Table 4: Weighted Distribution of Children by Race/Ethnicity and Corresponding Combined Vaccine Series 4:3:1:3*:3:1:4*, Pneumococcal, and Varicella Vaccination Coverage Estimates, National Immunization Survey, 2015

| Race/Ethnicity Classification | Weighted Distribution of Children aged 19-35 Months in U.S. Estimate (%) | Weighted Percentage 4:3:1:3*:3:1:4 UTD Estimate (%) (Standard Error (%)) | Weighted Percentage 4+ Pneumococcal Estimate (%) (Standard Error (%)) | Weighted Percentage 1+ Varicella at 12+ Months Estimate (%) (Standard Error (%)) |
|---|---|---|--|---|
| Hispanic | 26.67 | 71.69 (1.64) | 84.01 (1.27) | 92.67 (0.91) |
| Non-Hispanic white only | 48.36 | 72.68 (0.90) | 85.01 (0.74) | 91.16 (0.55) |
| Non-Hispanic black only | 12.56 | 69.06 (1.84) | 81.43 (1.49) | 91.77 (1.05) |
| Non-Hispanic American Indian or Alaska Native only | 1.00 | 68.17 (4.59) | 77.13 (4.05) | 87.75 (3.05) |
| Non-Hispanic Asian only | 5.16 | 77.90 (2.52) | 84.96 (2.12) | 93.44 (1.50) |
| Non-Hispanic Native Hawaiian or Pacific Islander only | 0.25 | 71.75 (7.48) | 76.77 (6.75) | 91.82 (3.47) |
| Multiracial | 6.01 | 73.73 (2.34) | 83.66 (1.86) | 92.09 (1.26) |
| Non-Hispanic white/black | 2.76 | 68.43 (3.66) | 80.20 (3.23) | 91.53 (2.11) |
| Non-Hispanic white/American Indian or Alaska Native | 0.87 | 73.25 (4.87) | 82.22 (4.35) | 93.49 (2.59) |
| Non-Hispanic white/Asian | 1.33 | 79.64 (3.58) | 87.72 (2.96) | 90.45 (2.80) |
| Non-Hispanic other combination | 1.05 | 80.54 (7.22) | 88.80 (3.78) | 94.46 (1.82) |

Note: UTD = up-to-date. Weighted by PROVWT_D. Children with an unknown Hispanic origin and/or race were imputed by a hot-deck method. This table includes both landline and cell-phone interviews, but excludes territories.

* 4+ DTaP/DTP/DT; 3+ polio; 1+ MCV; full series Hib, i.e., 3 or 4 doses depending on type of vaccine received; 3+ Hep B; 1+ varicella at or after 12 months of age; and 4+ PCV

4.8. Sub-Sets of the NIS Data

The NIS public-use data file contains data for all eligible children who have a completed household interview. An interview is considered complete if the respondent completed Section C of the questionnaire. As explained in Section 6 of this guide, each child with a completed household interview is assigned a weight (**RDDWT_D for the United States, excluding territories; RDDWT_D_TERR for the United States, including territories**) for use in estimation.

The NIS uses the synthesized provider-reported vaccination histories to form the estimates of vaccination coverage because the provider data are considered more accurate than household-reported data. Thus, the most important sub-set of the data consists of children with adequate provider data. For these children, one or more providers returned an immunization history questionnaire that included vaccination data. Unvaccinated children are also considered to have adequate provider data. As discussed in Section 7 below, the PDAT variable identifies the children with adequate provider data (PDAT=1). These children have a separate weight (**PROVWT_D for the United States, excluding territories; PROVWT_D_TERR for the United States, including territories**), which should be used to form estimates of vaccination coverage (see Section 6).

4.9. Confidentiality and Disclosure Avoidance

To prevent identification of participants in the NIS and the resulting disclosure of information, certain items from the questionnaires are not included in the public-use data file. In addition, some of the released variables either are top- or bottom-coded, or have their categories collapsed. Variable labels indicate which variables have been re-coded in these ways.

5. Quality Control and Quality Assurance Procedures

A major contributor to NIS data quality is its sample management system, which in 2015 managed over 480 sample frame by estimation area by quarter samples and used a number of performance measures to

track their progress toward completion. Important aspects of the quality assurance program for the RDD component of the NIS included on-line interviewer monitoring; on-line provider look-ups in a database system integrated with the CATI system, including names, addresses, and telephone numbers of vaccination providers; and automated range-edits and consistency checks. These and other quality assurance procedures contributed to a reduction in total data collection cost by minimizing interviewer labor and overall burden to respondents. Khare et al. (2000), Khare et al. (2001), and the *National Immunization Survey: Guide to Quality Control Procedures* (CDC 2002) describe quality assurance procedures.

The Provider Record Check component used quality control measures at four junctions: prior to mailing packets to providers; during the telephone prompting effort; during the editing of returned questionnaires; and during and after their data entry. The final quality assurance activities are implemented during post-processing of the returned questionnaires or vaccination records. All returned questionnaires were examined to identify and correct any obvious errors prior to data entry and then key-entered with 100% verification. The keying error rate is estimated, by way of a second verification process, to be less than 1%.

6. Sampling Weights

Each of the two phases of data collection results in a separate sampling weight for each child that has data at that phase. The RDD-phase sampling weights permit analyses of data for children with completed household interviews. Each child with adequate provider data (the sub-set on which official estimates of vaccination coverage are based) has a provider-phase sampling weight. **The dual-frame RDD-phase sampling weight variable for producing estimates for children with completed household interviews in the United States excluding territories is called RDDWT_D; and the dual-frame RDD-phase weight variable for producing estimates for the United States including territories is called RDDWT_D_TERR. The dual-frame provider-phase sampling weight variable for producing estimates for children with adequate provider data in the United States excluding territories is called PROVWT_D; and the dual-frame provider-phase weight variable for producing estimates for the United States including territories is called PROVWT_D_TERR.** See Section 8 of this user's guide for more information about the weights included in the data file and the proper way to use them.

As discussed below, revisions in weighting methodology were made on various occasions and the names of the weight variables were also changed to keep track of the revisions. The RDD-phase sampling weights were called HY_WGT in 1995-2001, RDD_WT in 2002, WGT_RDD in 2003 and 2004, RDDWT in 2005-2008, RDDWT/RDDWTVI from 2009-2010, RDDWT_LL/RDDWTVI_LL/RDDWT_D in 2011, RDDWT_D/RDDWTVI_D in 2012, RDDWT_D/RDDWTVIGU_D in 2013, **and RDDWT_D/RDDWT_D_TERR in 2014-2015.** The provider-phase sampling weights were called W0 in 1995-2001, WT in 2002, WGT in 2003 and 2004, PROVWT in 2005-2008, PROVWT/PROVWTVI from 2009-2010, PROVWT_LL/PROVWTVI_LL/PROVWT_D in 2011, PROVWT_D/PROVWTVI_D in 2012, PROVWT_D/PROVWTVIGU_D in 2013, **and PROVWT_D/PROVWT_D_TERR in 2014-2015.**

A sampling weight may be interpreted as the approximate number of children in the target population that a child in the sample represents. Thus, for example, the sum of the sampling weights of children who are up-to-date (on a particular vaccine or series of vaccines) yields an estimate of the total number of children in the target population who are up-to-date. Dividing this sum by the total of the sampling weights for all children gives an estimate of the corresponding vaccination coverage rate.

This section describes how these weights are developed and adjusted so as to achieve an accurate representation of the target population. The base weights reflect each telephone number's probability of being selected into the sample; the adjustments take into account non-resolution of residential/non-residential/non-working status of a telephone number, non-response to the screener and household interviews, number of telephone lines in the household, combination of landline and cell-phone sample sources, raking for differential coverage rates and non-coverage of households that do not have telephones, non-response by providers, and a final raking adjustment. Note that when deriving dual-frame weights, initial adjustments described below are performed separately for the landline and cell-phones samples, and then both samples are combined and further adjustments are performed on the combined samples.

6.1. Base Sampling Weight

In each quarterly NIS sample, each child with a completed household interview receives a base sampling weight. For all four quarters of the landline and cell-phone samples, the base sampling weight is equal to the total of telephone numbers in the sampling frame for the estimation area divided by the total of telephone numbers that were randomly sampled from that sampling frame and released for interview during that quarter.

6.2. Adjustments for Non-Resolution of Telephone Numbers, Screener Non-Response, and Interview Non-Response

Non-response occurs in population-based surveys when potential respondents refuse to participate, are not available at the time of the interview, or could not be reached during the survey period. Thus, the sum of the base sampling weights of children with completed household interviews will underestimate the size of the target population in the estimation area, because not all sampled households respond to all stages of data collection up to the household interview. As a result, the base sampling weights must be adjusted so they accurately reflect the number of children in the target population that each sampled child with a completed household interview represents.

Some sampled households with age-eligible children fail to complete the household interview because of unit non-response: for some telephone numbers, it is never determined whether or not the number is a working residential number despite multiple call attempts; for some households it is never determined whether or not the household contains age-eligible children; and some households with age-eligible children do not complete the household interview. To compensate for these three types of unit non-response, the sampling weights of children with a completed household interview are adjusted to account for the estimated number of age-eligible children in households whose telephone numbers are never resolved, the estimated number of age-eligible children in households that fail to complete the screening interview, and the number of identified age-eligible children for whom the household interview is not completed. For the landline sample, each of these adjustments is carried out within each estimation area by forming weighting cells based on the residential directory-listed status of the sample telephone number, percent of the population that is white in the telephone exchange, and MSA status of the telephone exchange (i.e., weighting cells were formed from directory-listed versus non-directory-listed telephone number; telephone exchanges with 75% or higher white population versus telephone exchanges with less than 75% white population; and MSA/non-MSA status). For the cell-phone sample, each of these adjustments is carried out within estimation area by forming weighting cells based on MSA/non-MSA status of the wire center associated with the cell-phone number. Each cell in each stage of

adjustment must have sufficient resolved/responding cases (usually 20, but 15 for interview non-response) at that stage of adjustment; cells with a deficient number of responding cases are collapsed with neighboring cells. The order of the variables in cell collapsing for the landline sample is MSA status, percent of population that is white, and directory listed status of the telephone number, and for the cell-phone sample, both MSA categories are collapsed if either of the cells have a deficient number of responding cases. Once the adjustment cells are formed, the weights of the unresolved/non-responding records from the previous adjustment step are distributed to the weights of the resolved/responding records within each cell.

6.3. Adjustment for Multiple Telephone Lines and Deriving Annual Weights

Once the non-response-adjusted interview weights for households are computed, these weights are adjusted for additional telephone lines in the household. Because households with multiple telephone lines have a greater chance of being sampled, for the landline sample each child's household interview weight is adjusted by dividing it by the total number of residential landlines reported in the household (up to a maximum of 3), and for the cell-phone sample each child's household interview weight is adjusted by dividing it by the total number of cell-phones used by parents or guardians (up to a maximum of 3). Prior to 2005, the adjustment for multiple telephone lines was made by adjusting the base sampling weights before making any other adjustments. Beginning in 2005, the adjustment for multiple telephone lines has been shifted after the interview non-response adjustment, because the information on the number of telephone lines in a household is available only for households with completed household interviews. This shifts the adjustment for multiple telephone lines to the point where the information about the number of telephone lines is actually collected.

Up to the previous step, the sampling weights are adjusted separately for each quarter and sample type (landline, cell-phone), and the weights in each quarter pertain to the target population. However, annual vaccination coverage estimates are obtained from data for four consecutive quarters, so the weights in

each quarterly file are adjusted when the data from the four quarters of the landline and cell-phone samples are combined. The adjustment factor is proportional to the number of households with completed household interviews in each quarter within sample type (landline, cell-phone) and estimation area.

6.4. Calibration

Next, survey weights are calibrated to population control totals as described below. The control totals used for the NIS are derived from current natality data from the National Center for Health Statistics (NCHS 2012, 2013). Because the Vital Statistics data give the counts of all live births in the United States, regardless of whether the household has telephone service, the control totals include all eligible children. The control total for each raking dimension is derived from the NCHS natality files from 2012 and 2013 (children born between July 1, 2012 and November 30, 2013 would have been 19 through 35 months on June 30, 2015). Use of the natality data to form the required population control totals for the NIS has three limitations: 1) the natality file provides a universe of live births and therefore does not reflect infant mortality; 2) the natality file does not include children born outside the United States who immigrate to this country before reaching ages 19 through 35 months; and 3) the natality file records residence at time of birth, and some children may move from one estimation area to another by the time they reach 19 through 35 months of age. Adjustments are made to the natality data to account for these three factors. **For 2015, the methodology is similar to that for 2014 except instead of using three-year American Community Survey Public-Use Microdata Sample (PUMS) data, the combined 2012, 2013, and 2014 one-year American Community Survey PUMS data files were used to make the immigration and migration adjustments.**

The proportions of 19 through 35 month old children by detailed telephone status (cell-phone-only, landline and cell-phone dual user, landline-only, phoneless) within each estimation area were derived using a similar small area modeling approach as described in Blumberg et al. 2011. These modeled telephone status estimates are applied to the population control total for the estimation area to estimate the control totals by detailed telephone status within the estimation area.

Survey weights for the landline and cell-phone samples must be integrated to provide dual-frame weights for the full target population of age-eligible children. The landline and cell-phone sampling frames overlap in coverage of children in landline and cell-phone dual use households and exclude children in phoneless households. The critical issues associated with combining the landline and cell-phone samples are: a) adjustment for overlap of the landline and cell-phone samples; and b) adjustment for noncoverage of children in phoneless households.

Prior to combining the landline and cell-phone samples, survey weights are adjusted to agree with independent estimates of the population total by telephone status. Adjustments to population totals by telephone status are made separately for the landline sample and the cell-phone sample (with the overlap adjusted for in the next step, as explained below). Since the proportion of children living in landline-only households is continuously decreasing (and thus, the number of landline-only sample cases is small), children in dual landline and cell-phone households from the landline sample are combined with children in landline-only households and then adjusted to the combined population estimate of children living in dual user and landline-only households within each estimation area. In the cell-phone sample, in each estimation area children in dual landline and cell-phone households are weighted to represent children living in dual landline and cell-phone households in the estimation area, and children in cell-phone-only households are weighted to represent children in cell-phone-only households in the estimation area.

Next, since the landline and cell-phone sampling frames overlap in coverage of children in landline and cell-phone dual-use households, children in dual-user households from both samples are combined based on the effective number of children with a completed household interview within each sample type (landline, cell-phone), and are weighted to represent children in dual-use households within each estimation area. Finally, children in phoneless households, which are excluded from the dual-frame sample, are accounted for in the raking step described below.

To reduce sampling variability and improve the precision of estimation, extreme weights are trimmed and then recalibrated to control totals. RDD sampling weight values exceeding the median weight plus three times the interquartile range of the weights within an estimation area are truncated and then recalibrated to control totals. This is done by up to five iterations. This weight trimming prevents children with unusually large weights from having an unusually large impact on immunization coverage estimates.

The final step in adjusting the RDD sampling weights is a raking adjustment (Deming 1943) of the trimmed, telephone status adjusted weights. The raking procedure uses estimation area-level control totals for maternal education categories, maternal race/ethnicity, age group of the child, gender of the child, and telephone status. Briefly, raking takes each variable in turn and applies a proportional adjustment to the current weights of the children who belong to the same category of the variable. After a number of iterations over all the variables, the raked weights have totals that match all the desired control totals. Raking makes it possible to incorporate additional variables into the weighting and to use more detailed categories for those variables. Smith et al. (2005) gives the details of various aspects of the NIS estimation procedures.

The sampling weights after all the foregoing adjustments constitute the “RDD sampling weights” (**RDDWT_D for the United States excluding territories; RDDWT_D_TERR for the United States including territories**).

6.5. Adjustment for Provider Non-Response

Among the 26,975 children with a completed household interview from the landline and cell-phone samples (excluding territories), 15,167 (56.2%) had adequate provider data. Starting with the 2002 NIS public-use data file, the definition of children with adequate provider data includes unvaccinated children. These are children for whom the respondent reported during the household interview that the child had received no vaccination and has no providers, or for whom one or more providers were reported but those providers reported administering no vaccinations. Among the 15,167 children with adequate provider

data, 178 were unvaccinated children. Failure to obtain adequate provider data for the remaining 43.8% was attributable to:

- parent or guardian not identifying any providers or not giving consent to contact the child’s vaccination provider(s) (35.7%);
- consent to contact vaccination providers obtained but no providers returned the immunization history questionnaire (4.7%); and
- one or more providers returned the immunization history questionnaire, but no providers reported any vaccination data (3.4%).

The 11,808 children for whom a household interview was completed but adequate provider data were not obtained are classified as “partial non-responders” because they have only a partial response to the NIS as a whole.

Empirical results suggest that children with adequate provider data have characteristics believed to be associated with a greater likelihood of being up-to-date, compared with children who had missing provider data. Specifically, children with adequate provider data are more likely to live in households that have higher total family income, have a white mother, and live outside a central city of a Metropolitan Statistical Area. Also, a child with missing provider data is less likely to live in the state where the mother lived when the child was born. These factors indicate a potential lack of continuity of health care, and are associated with lower vaccination rates (Coronado et al. 2000). If no adjustment is made to the RDD sampling weights to account for these differences, estimated vaccination coverage rates may be biased.

To reduce potential bias in estimators of vaccination coverage attributable to partial non-response, a weighting-class adjustment is used in each estimation area (Brick and Kalton 1996). This adjustment involves three steps. In the first step, sampled children are classified according to the quintile of their estimated probabilities of having adequate provider data. In the statistical literature these probabilities are

called response propensities (Rosenbaum and Rubin 1983, 1984; Rosenbaum 1987). Children who have similar response propensities will also be similar with respect to variables that are strongly associated with the probability of having adequate provider data. In this important respect, children in each class are comparable. Because of this comparability, any sub-sample of children in a class may represent all children in the class. Therefore, the weighting-class adjustment uses the children with adequate provider data to represent all children in the class. An NCHS Series 2 Report on the statistical methodology of the NIS (Smith et al. 2005) includes details of the methodology for forming weighting classes based on propensity scores. This report can be viewed at http://www.cdc.gov/nchs/data/series/sr_02/sr02_138.pdf.

In the second step of this weighting-class adjustment, within each class an adjustment factor redistributes the RDD sample weights of the children with missing provider data to the weights of the children who have adequate provider data. These adjusted sampling weights of children with adequate provider data are initial non-response-adjusted provider-phase weights. The model for children with adequate provider data includes significant main effects, and also significant two-way interactions between sample type (landline, cell-phone) and all other variables.

Within an estimation area, the sums of non-response adjusted weights of children with adequate provider data for the various levels of important socio-demographic variables (such as race/ethnicity) may not be equal to corresponding population totals. To reduce bias attributable to these differences, raking was used in the third step to adjust the non-response adjusted weights to match estimation area control totals.

Control totals for these variables were estimated using the weighted totals from the sample of children with completed household interviews. Smith et al. (2001b, 2005) describe the development of this approach in more detail. Similar to the RDD weighting, the extreme weights exceeding the median weight plus three times the interquartile range of the weights within an estimation area are truncated and then recalibrated to control totals. These raked weights of children with adequate provider data are called “final provider-phase weights” (**PROVWT_D for the United States excluding territories; PROVWT_D_TERR for the United States including territories**). Because of the comparability of

children within each weighting class, any estimate that uses data only from the children with adequate provider data along with their provider-phase sampling weights, will have less bias attributable to differences between children with adequate provider data and children with missing provider data.

Appendix B summarizes the distribution of the sampling weights (**RDDWT_D, PROVWT_D, RDDWT_D_TERR, and PROVWT_D_TERR**) in each estimation area.

NIS public-use data files for 1995 to 2001 do not include sampling weights that account for the effect of unvaccinated children. An assessment of the effect of accounting for unvaccinated children for the period 1995 to 2003 was made. Weights were calculated for each year with and without unvaccinated children and the vaccination coverage estimates compared. Details of this assessment and the results are available in the user's guide for the 2004 NIS public-use data file. At the national level, accounting for unvaccinated children had very little effect on the estimates of 4:3:1:3 vaccination coverage. Within estimation areas also, the two coverage estimates differed little. The largest difference (in either direction) was most often around 2 percentage points. Differences of that magnitude are small relative to the standard errors of the estimates. Although accounting for unvaccinated children has a small effect on estimates of vaccination coverage, data users who use the public-use data files to examine estimation area-level trends over time are advised to interpret the results with appropriate caution.

6.6. Sampling Weights for Territories

The NIS weighting process was followed as closely as possible for territories. Due to differences in the availability of external data sources for territories, slight changes were necessary to accurately estimate vaccination rates for this area. These differences are stated below.

In step 6.2, each of the non-response adjustments for territories was done at the estimation area level. That is, no weighting cells were formed for territories.

For the U.S. Virgin Islands, 2013 natality data were not available, so 2011 and 2012 data (NCHS 2011, 2012) were used instead of 2012 and 2013 data to create the control totals for the U.S. Virgin Islands. Similar to the dual-frame weights for the United States excluding territories, the final step in adjusting the RDD sampling weights for territories is a raking adjustment. For Guam and Puerto Rico, a different set of race/ethnicity categories were used for post-stratification and raking adjustments than were used in other areas. The three Guam race/ethnicity categories were: Chamorro/Guamanian, Other Asian/Pacific Islander, and All Other. The two Puerto Rico race/ethnicity categories were: White and All Other.

After sampling weights were calculated for all children in all estimation areas, including territories, they were stored in the variables RDDWT_D_TERR and PROVWT_D_TERR. These weight variables permit one to conduct analysis of all estimation areas, including territories. The weight variables RDDWT_D and PROVWT_D are equal to RDDWT_D_TERR and PROVWT_D_TERR for all children except those in territories, for whom the value of these weight variables is blank or missing. RDDWT_D and PROVWT_D permit one to conduct analysis of all estimation areas, excluding territories.

7. Contents of the Public-Use Data File

The NIS public-use data file contains a record for each eligible child for whom Section C of the household interview was completed, along with household-reported information about the child and the child's mother. For children with Immunization History Questionnaires (IHQs) returned by one or more providers, the file also contains provider characteristic variables, as well as variables based on the child's synthesized provider-reported vaccination history: the age of the child at each vaccination, the number of each type of vaccination received, and indicators of whether the child is up-to-date with respect to various recommended vaccines and vaccine series.

The public-use data file consists of ten sections, the contents of which are described below in detail. For additional information, users are encouraged to consult the codebook (NCIRD 2016). The codebook is divided into the ten sections described below and contains variable names, labels, and response frequencies (for categorical variables). For select variables, the codebook also gives additional information about the variable in the "Notes" field.

Table 5 lists key NIS variables commonly used in analyses. A full list of variables appearing on the 2004-2015 NIS public-use data files appears in Appendix E, along with the reason for the addition, subtraction, or modification of the variables in 2005-2015. Information on changes made between 1995-2004 can be found in the *Alphabetical Listing of Variables that are Not Available in All Public-Use Data Files, National Immunization Survey, 1995-2004*. www.cdc.gov/nchs/data/nis/pufvariables1995to2004.pdf

Table 5: NIS Variables Commonly Used in Analyses or for Published Estimates

| Variables | Categories |
|--|--|
| ID Variables | |
| SEQNUMC – unique child ID variable | |
| SEQNUMHH – unique household ID variable | |
| Geographic Variables | |
| ESTIAP15 – estimation area number (introduced in 2008; ITRUEIAP used through 2004; ESTIAP in 2005; ESTIAP06 in 2006; ESTIAP07 in 2007; ESTIAP08 in 2008; ESTIAP09 in 2009; ESTIAP10 in 2010; ESTIAP11 in 2011; ESTIAP12 in 2012; ESTIAP13 in 2013; ESTIAP14 in 2014; ESTIAP15 in 2015) | |
| STATE – state FIPS code | |
| CEN_REG – census region | Northeast Midwest South West |
| Child Demographic Variables | |
| AGEGRP – age category of child | 19-23 months 24-29 months 30-35 months |
| RACEETHK – race/ethnicity of child (introduced in 2002; RACEKIDR used in 1995-2001) | Hispanic White alone, non-Hispanic Black alone, non-Hispanic All other races alone and multi-racial, non-Hispanic |
| SEX – gender of child | Male Female |
| FRSTBRN – firstborn status of the child | No Yes |
| Mother Demographic Variables | |
| EDUC1 – education of the mother | <12 years 12 years >12 years, not a college graduate College graduate |
| MARITAL2 – marital status of mother (Living with partner response option added to questionnaire in 2015) | Currently married Never married, widowed, divorced, separated, or deceased, living with partner |
| M_AGEGRP – age group of mother | <=19 years 20-29 years 30 years or older |
| Poverty Variables | |
| INCPOV1 – poverty status (introduced in 2005; INCPOV1R used through 2004) | At or above poverty level, income > \$75,000 At or above poverty level, income <= \$75,000 Below poverty level Not determined |
| INCPORAR – income-to-poverty ratio (introduced in 2005; INCPORAT used through 2004) | |
| WIC Variables | |

| Variables | Categories |
|--|--------------------|
| CWIC_01 – child ever participated in WIC program | Yes |
| | No |
| | Never heard of WIC |
| | Don't know |
| | Refused |
| CWIC_02 – child currently participating in WIC program | Missing |
| | Yes |
| | No |
| | Don't know |
| | Refused |
| Breastfeeding Variables | |
| CBF_01 – child ever fed breast milk | Missing |
| | Yes |
| | No |
| | Don't know |
| BF_ENDR06 – length of time in days child was fed breast milk | Refused |
| BF_EXCLR06 – length of time in days child was exclusively fed breast milk or formula (introduced in 2006) | Missing |
| BF_FORMR08 – age in days when child was first fed formula (introduced in 2008; BF_FORMR06 used in 2006 and 2007) | |
| Chicken Pox Variables | |
| HAD_CPOX – did child ever have chicken pox (introduced in 2005; I_HADCPX used through 2004) | Yes |
| | No |
| | Don't know |
| | Refused |
| | Missing |
| AGECPOXR – age in months when child had chicken pox (introduced in 2005; IAGECPXR used through 2004) | 0-6 months |
| | 7-12 months |
| | 13-18 months |
| | 19-24 months |
| | 25-30 months |
| | 31 months or older |
| Presence of Provider Data Variables | |
| PDAT – adequate provider data indicator | Yes |
| | No |
| Number of Provider-Reported Doses of Vaccine Variables | |
| P_NUMDTP – total number of DTP/DTaP doses | |
| P_NUMPOL – total number of polio doses | |
| P_NUMMMR – total number of MCV doses | |
| P_NUMHIB – total number of Hib doses | |
| P_NUMHEP – total number of hepatitis B doses | |
| P_NUMVRC – total number of varicella doses | |
| P_NUMPCV – total number of pneumococcal doses | |
| P_NUMFLU – total number of seasonal influenza doses | |
| P_NUMHEA – total number of hepatitis A doses | |
| P_NUMROT – total number of rotavirus doses | |
| Provider Characteristic Variables | |

| Variables | Categories |
|---|---|
| PROV_FAC – provider facility type | All public facilities All hospital facilities All private facilities All military/other facilities Mixed types Unknown |
| VFC_ORDER – do child’s providers order vaccines for children from state/local health department? (introduced in 2006) | All providers Some but not all providers No providers Unknown |
| REGISTRY – provider(s) reported child’s vaccination(s) to state or community immunization registry | All providers Some but not all providers No providers Unknown |

Before describing the sections of the public-use data file below, we first summarize the differences between the 2014 and 2015 NIS public-use data files:

- Because the 2015 estimation areas differ from those used in prior years, a new 2015 estimation area variable has been added (ESTIAP15) and the 2014 estimation area variable (ESTIAP14) has been dropped.
- A new response option was added for mother’s marital status, “Living with partner”. A value of “2” on the recoded marital status variable MARITAL2 now represents “never married; widowed; divorced; separated; deceased; living with partner”.

7.1. Section 1: ID, Weight, and Flag Variables

SEQNUMHH and SEQNUMC are the unique household and child identifiers, respectively. PDAT indicates which children are considered to have adequate provider data. As described in Section 6 of this report, RDDWT_D/RDDWT_D_TERR and PROVWT_D/PROVWT_D_TERR are the final household- and provider-phase weights, respectively. PROVWT_D/PROVWT_D_TERR should be used when analyzing the provider-reported data, i.e., the variables in Sections 7, 8, and 9 of the NIS public-use data file.

7.2. Section 2: Household-Reported Vaccination and Chickenpox Information

Section 2 of the public-use data file contains variables derived from the information collected in Section B of the household questionnaire. In particular, it contains variables indicating whether the respondent reported that the child has had chicken pox disease (**HAD_CPOX**) and the child's age in months at chicken pox disease (**AGECPOXR**).

7.3. Section 3: Demographic, Socio-Economic, and Other Household/Child Information

Section 3 of the NIS public-use data file consists of information collected during the household screening interview and Section C of the household main interview. To protect confidentiality, many of these variables have been collapsed, top-coded, or bottom-coded from the original, fully-detailed versions; the variable labels (see the public-use data file codebook) indicate which variables have been collapsed or recoded.

AGEGRP is the age of the child in months in three categories (19-23 months, 24-29 months, 30-35 months), based on the child's best date of birth and the eligibility date. **SEX** gives the gender of the child, and **FRSTBRN** indicates whether the child is the first born, with missing values of these variables imputed. The language in which the interview was conducted is stored in variable **LANGUAGE**, and **C5R** gives the relationship of the respondent to the child.

The breastfeeding variables include whether the child was ever fed breast milk (**CBF_01**), length of time in days the child was fed breast milk (**BF_ENDR06**), the age in days when the child was first fed formula (**BF_FORMR08**), and the length of time in days the child was exclusively fed breast milk or formula (**BF_EXCLR06**). Two types of inconsistencies arise in the breastfeeding data: 1) duration of any breastfeeding can exceed age of the child, and 2) age when the child was first fed formula can exceed the age of the child. **BFENDFL06** is set equal to 1 when **BF_ENDR06** exceeds the age of the child (with a buffer), and **BFFORMFL06** is set equal to 1 when **BF_FORMR08** exceeds the age of the child (with a

buffer). Appendix C provides details on how the flags were created. Data users are cautioned to review Appendix C before analyzing any of the breastfeeding variables.

The WIC variables include whether the child ever participated in the WIC program (**CWIC_01**) and whether the child is currently participating (**CWIC_02**).

C1R and **CHILDNM** give the number of people and children, respectively, in the household. The child's Hispanic origin indicator, race with three categories, and race/ethnicity with four categories are presented in variables **I_HISP_K**, **RACE_K**, and **RACEETHK**, respectively; for each of these variables, missing values have been imputed. The age, education level, and marital status of the mother of the child are stored in variables **M_AGEGRP**, **EDUC1**, and **MARITAL2** (married vs. not married), with missing values imputed.

The categorized total combined income for the child's family is given by **INCQ298A**. **INCPOV1** gives the family's poverty status (at or above poverty, income > \$75,000; at or above poverty, income <= \$75,000; below poverty; unknown), and **INCPORAR** gives the ratio of the family's income to the poverty level. Household tenure is given by **RENT_OWN**.

The number of landline telephone numbers in the household, the number of working cell phones household members have available for personal use, and the number of these cell phones that are usually used by parents or guardians are given by **NUM_PHONE**, **NUM_CELLS_HH**, and **NUM_CELLS_PARENTS**, respectively.

Variable **CEN_REG** gives the census region of the respondent's current residence, and **MOBIL_I** indicates whether the mother's current state of residence is the same as her state of residence at the time of the child's birth.

7.4. Section 4: Geographic Variables

Variables **ESTIAP15** and **STATE** give the 2015 estimation area and state of residence, respectively, for each child. **EST_GRANT** indicates which of the 50 states, District of Columbia, and 5 local areas that receive federal Section 317 immunization grants (Bexar County, TX; City of Chicago, IL; City of Houston, TX; New York City; Philadelphia County, PA) the child resides in.

7.5. Section 5: Number of Providers Identified and Consent Variables

Variable **D7** indicates whether the respondent gave consent to contact the child's providers. If **D7=1**, then consent was granted; if **D7=2** then consent was explicitly denied; and if **D7** is missing, consent was not granted because the respondent broke off the interview before being explicitly asked for consent.

Variable **D6R** gives the number of providers identified by the respondent. Note that sometimes respondents report erroneous provider counts and sometimes report the same provider more than one time, and **D6R** does not reflect cleaning or de-duplication of the initially-reported provider count.

7.6. Section 6: Number of Responding Providers Variables

Variable **N_PRVR** indicates the number of providers returning IHQs with vaccination information for the child. That is, **N_PRVR** is the number of IHQs that were returned for the child that contain information on the IHQ shot grid.

7.7. Section 7: Characteristics of Providers Variables

The variables in this section of the public-use data file summarize the information collected in IHQ questions 5c, 6, and 7 across the child's providers who returned IHQs containing vaccination (i.e., shot grid) data.

PROV_FAC indicates the facility type of the child's vaccination providers based on responses to IHQ question 5c. If all of the child's providers that returned IHQs containing shot grid data (see Section 6 variable **N_PRVR**) reported the facility type to be:

- a public health department-operated clinic, community health center, or rural health clinic, then PROV_FAC=1 (all public facilities);
- a hospital-based clinic, then PROV_FAC=2 (all hospital facilities);
- a private practice, then PROV_FAC=3 (all private facilities);
- a military health care facility, WIC clinic, school-based health center, pharmacy, or other type of facility, then PROV_FAC=4 (all military/WIC/school/pharmacy or other facilities).

If the responses of providers that returned IHQs containing shot grid data fell into more than one of the above bulleted categories, PROV_FAC=5 (mixed); otherwise, if at least one of the child's providers returned an IHQ containing shot grid data, PROV_FAC=6 (unknown). If none of the child's providers returned an IHQ containing shot grid data, PROV_FAC is set to missing.

The Vaccines For Children (VFC) program is a federally-funded program that provides vaccines at no cost to children who might not otherwise be vaccinated because of inability to pay (<http://www.cdc.gov/vaccines/programs/vfc/index.html>). CDC buys vaccines at a discount and distributes them to awardees—i.e., state health departments and certain local and territorial public health agencies—which in turn distribute them at no charge to those private physicians' offices and public health clinics registered as VFC providers. **VFC_ORDER**, based on responses to IHQ question 6, indicates whether the child's vaccination providers order vaccines from a state or local health department to administer to children. If all of the child's providers that returned IHQs containing shot grid data (see Section 6 variable N_PRVR) reported that they order vaccines from a state or local health department to administer to children, then VFC_ORDER=1 (all providers); if at least one of the child's providers that returned an IHQ containing shot grid data reported that the practice orders vaccines from a state or local health department to administer to children and the child's other providers that returned IHQs containing shot grid data reported either that they did not order such vaccines or that they did not know whether or not they did, then VFC_ORDER=2 (some but possibly or definitely not all providers); if all of the child's

providers that returned IHQs containing shot grid data reported that they do not order vaccines from a state or local health department to administer to children, then VFC_ORDER=3 (no providers); if none of the conditions for VFC_ORDER=1, 2, or 3 was met but at least one of the child's providers returned an IHQ containing shot grid data, VFC_ORDER=4 (unknown). If none of the child's providers returned an IHQ containing shot grid data, VFC_ORDER is set to missing. Note that having a provider that orders VFC vaccine does not imply that the child is VFC-entitled; providers enrolled in the VFC program could also vaccinate children who are not VFC-entitled.

REGISTRY is based on responses to IHQ question 7 and indicates whether the child's vaccination providers reported the child's vaccinations to a community or state immunization registry (also known as Immunization Information Systems, or IIS). If all of the child's providers that returned IHQs containing shot grid data (see Section 6 variable N_PRVR) indicated that they reported to a registry, then REGISTRY=1 (all providers); if at least one of the child's providers that returned an IHQ containing shot grid data indicated that the practice reported to a registry and the child's other providers that returned IHQs containing shot grid data indicated that they did not report to a registry, that they did not know whether or not they reported to a registry, or that the question is not applicable, then REGISTRY=2 (some but possibly or definitely not all providers); if all of the child's providers that returned IHQs containing shot grid data indicated that they did not report to a registry or that the question is not applicable, then REGISTRY=3 (no providers); if none of the conditions for REGISTRY=1, 2, or 3 was met but at least one of the child's providers returned an IHQ containing shot grid data, REGISTRY=4 (unknown). If none of the child's providers returned an IHQ containing shot grid data, REGISTRY is set to missing.

7.8. Section 8: Provider-Reported Up-To-Date Vaccination Variables

This section contains vaccination count and up-to-date variables based on the child's synthesized provider-reported vaccination history. To facilitate data processing and to accommodate the large and continually growing number of vaccination types covered by the NIS, the provider-reported vaccination data are organized around the concept of vaccine categories and vaccine types within vaccine category.

The vaccine categories correspond to the sections of the IHQ shot grid, and the vaccine types correspond to the type boxes on the IHQ shot grid. (For each vaccine category, an “unknown” vaccine type is created for vaccinations that are reported without a type box being checked. Also, a few vaccine types, such as Measles-Mumps, arise through the backcoding of shots initially reported in the “other” section of the IHQ shot grid.) Table 6 shows the vaccine categories and types for the 2015 NIS. Note that a single vaccination can fall into more than one vaccine category; for example, an MMR-Varicella vaccination is part of both the Measles-containing and Varicella-containing vaccine categories. (The full list of vaccine type codes can also be found in Appendix H.)

For each vaccine category, Section 8 of the public-use data file contains a variable typically named **P_NUMYYY** – where “YYY” is the vaccine category abbreviation given in Table 6 – that stores the number of vaccinations in that vaccine category in the child’s synthesized provider-reported vaccination history. For each vaccine type in Table 6, Section 8 also contains a variable that stores the number of vaccinations of that vaccine type in the child’s synthesized provider-reported vaccination history. For example, **P_NUMDHI** is the number of DTaP/HepB/IPV shots in the child’s history.

This section of the public-use data file also contains up-to-date indicators for a variety of recommended vaccines and vaccine series. These variables’ names typically begin with “**P_UTD**”. Additional variables indicate whether the child is up-to-date for various vaccine series. For example, **P_UTD431** indicates whether the child has received 4 or more DTaP/DTP shots, 3 or more polio shots, and one or more measles-containing shots. The variable labels indicate what is needed to be considered up-to-date for each variable, and the “Notes” field in the codebook shows the vaccine type codes (see Table 6) being included when determining whether the child is up-to-date.

Note that it is possible that the administration of the NIS interview itself prompts some respondents to vaccinate their children following the interview; to ensure that the vaccination rate estimates aren’t artificially boosted because of this, the synthesized vaccination history count and up-to-date variables in

this section of the public-use data file count only vaccinations received before the date the household interview was completed.

7.8.1. *Hib Up-To-Date Variables*

A Hib vaccine shortage and interim recommendation to suspend the booster dose for healthy children occurred December 2007 to September 2009 (CDC 2010). Furthermore, the NIS has historically considered children to be up-to-date for Hib if the child had 3 or more doses of any Hib-containing vaccine, but for some Hib vaccine product types, 4 doses are required. Because the NIS has historically not distinguished between product types for Hib vaccine, children who received 3 doses of a vaccine product that required 4 doses were misclassified as up-to-date for Hib (CDC 2010).

Because of the Hib vaccine shortage and because of the dependence of the Hib recommendation on product type, in 2009 the IHQ was modified to capture the manufacturer of the Hib vaccinations the child has received. Beginning with the 2009 NIS public-use data file, new up-to-date variables were added to indicate up-to-date status based on Hib recommendation (i.e., the primary series recommended during the shortage vs. the full series) and on the Hib manufacturer.

Table 6: Vaccine Categories and Vaccine Types, National Immunization Survey, 2015

| Vaccine Category Abbreviation | Vaccination Category Description | Vaccine Type Code | Vaccine Type Description |
|--------------------------------------|---|--------------------------|---------------------------------------|
| DTP | DTaP/DTP-containing vaccine | 03 | DTaP/DTP-containing, unknown type |
| | | 04 | DTaP |
| | | 07 | DTaP-Hib |
| | | 08 | DTaP-HepB-IPV |
| | | D3 | DTaP-IPV-Hib |
| POL or POLIO | Polio-containing vaccine | 08 | DTaP-HepB-IPV |
| | | 20 | OPV |
| | | 21 | IPV |
| | | 22 | Polio-containing, unknown type |
| | | D3 | DTaP-IPV-Hib |
| MCV or MMR | Measles-containing vaccine | 30 | MMR |
| | | 31 | Measles only |
| | | 32 | Measles-mumps |
| | | 33 | Measles-rubella |
| | | MM | Measles-containing, unknown type |
| | | VM | MMR-Varicella |
| HIB | Hib-containing vaccine | 07 | DTaP-Hib |
| | | 43 | HepB-Hib |
| | | 44 | Hib-only, unknown type |
| | | D3 | DTaP-IPV-Hib |
| | | HG | Hib-only (GSK) |
| | | HI | Hib-containing, unknown type |
| | | HM | Hib-only (Merck) |
| | | HS | Hib-only (Sanofi) |
| | | HY | Hib-MenCY |
| HEPB or HEP | Hepatitis B-containing vaccine | 08 | DTaP-HepB-IPV |
| | | 43 | HepB-Hib |
| | | 60 | HepB-only |
| | | HB | HepB-containing, unknown type |
| VRC | Varicella-containing vaccine | VA | Varicella-containing, unknown type |
| | | VM | MMR-Varicella |
| | | VO | Varicella-only |
| PCV | Pneumococcal-containing vaccine | 70 | Conjugate-unknown |
| | | 71 | Polysaccharide |
| | | 72 | Pneumococcal-containing, unknown type |
| | | 73 | Conjugate-7 |
| | | 74 | Conjugate-13 |

| Vaccine Category Abbreviation | Vaccination Category Description | Vaccine Type Code | Vaccine Type Description |
|--------------------------------------|---|--------------------------|---------------------------------|
| HEPA or HEA | Hepatitis A-containing vaccine | HA | Hepatitis A |
| FLU | Seasonal influenza vaccine | FL | Seasonal flu, unknown type |
| | | FM | Seasonal flu spray |
| | | FN | Injected seasonal flu |
| MP | Mumps-only vaccine | MP | Mumps-only |
| MPRB or MPR | Mumps-Rubella-only vaccine | MB | Mumps-Rubella-only |
| RB | Rubella-only vaccine | RB | Rubella-only |
| ROT | Rotavirus-containing vaccine | RG | Rotarix [®] (GSK) |
| | | RM | RotaTeq [®] (Merck) |
| | | RO | Rotavirus, unknown type |

Table 7 shows the Hib up-to-date variables appearing on the public-use-date file beginning in 2009: in addition to the existing up-to-date indicator based on 3+ Hib of any type (P_UTDHIB), an indicator based on the “shortage” (i.e., primary series) recommendations accounting for manufacturer (3+ Hib of any type or 2+ Hib of Merck types) and an indicator based on the “routine” (i.e., full series) recommendations accounting for manufacturer (4+ Hib of any type or 2 Hib of Merck types followed by 1 Hib of any type) were added. Table 8 shows the up-to-date series variables that include Hib appearing on the public-use-date file beginning in 2009: in addition to the existing vaccine series up-to-date variables based on 3+ Hib of any type (PUTD4313, PUT43133, PU431331, PU4313313, PU4313314), variables based on the “routine” (i.e., full series) Hib recommendations accounting for manufacturer (4+ Hib of any type or 2 Hib of Merck types followed by 1 Hib of any type) were added (P_UTD431H_ROUT_S, P_UTD431H3_ROUT_S, P_UTD431H31_ROUT_S, P_UTD431H313_ROUT_S, P_UTD431H314_ROUT_S).

Note that for these Hib up-to-date variables that account for the manufacturer, if the manufacturer is unknown because the provider failed to check a type box on the IHQ, it has been assumed that the manufacturer of the Hib vaccine is not Merck; that is, these variables are based on a “strict” treatment of Hib vaccinations of unknown type, erring on the side of classifying the child as not up-to-date.

Beginning with the 2010 NIS public-use data file, two new vaccination series up-to-date indicators were added that ignore the Hib component altogether. These are PU431_31 (indicates up-to-date status as measured by PU431331, but excluding the Hib component) and PU431_314 (indicates up-to-date status as measured by PU4313314, but excluding the Hib component).

Table 7: Up-To-Date Variables for Hib, National Immunization Survey, 2009-2015

| Name | Description | Up-To-Date Criteria |
|------------------|--|--|
| P_UTDHIB | Historical UTD flag for Hib. | 3+ of any type (07,43,44,D3,HG,HI,HM,HS,HY) |
| P_UTDHIB_SHORT_S | UTD flag for Hib-shortage (i.e., primary series) recommendation, accounting for manufacturer. New starting 2009. | 3+ of any type (07,43,44,D3,HG,HI,HM,HS,HY) OR 2+ Merck types (HM,43) |
| P_UTDHIB_ROUT_S | UTD flag for routine (i.e., full series) Hib recommendation, accounting for manufacturer. New starting 2009. | 4+ of any type (07,43,44,D3,HG,HI,HM,HS,HY) OR 2 Merck types (HM,43) followed by 1 of any type (07,43,44,D3,HG,HI,HM,HS,HY) |

Table 8: Up-To-Date Variables for Vaccine Series Including Hib, National Immunization Survey, 2009-2015

| Name | Description |
|---------------------|--|
| PUTD4313 | UTD flag for the 4:3:1:3 series using the 3+ any type UTD definition for HIB |
| P_UTD431H_ROUT_S | UTD flag for the 4:3:1:3 series using the routine (i.e., full series) UTD definition for HIB |
| PUT43133 | UTD flag for the 4:3:1:3:3 series using the 3+ any type UTD definition for HIB |
| P_UTD431H3_ROUT_S | UTD flag for the 4:3:1:3:3 series using the routine (i.e., full series) UTD definition for HIB |
| PU431331 | UTD flag for the 4:3:1:3:3:1 series using the 3+ any type UTD definition for HIB |
| P_UTD431H31_ROUT_S | UTD flag for the 4:3:1:3:3:1 series using the routine (i.e., full series) UTD definition for HIB |
| PU4313313 | UTD flag for the 4:3:1:3:3:1:3 series using the 3+ any type UTD definition for HIB |
| P_UTD431H313_ROUT_S | UTD flag for the 4:3:1:3:3:1:3 series using the routine (i.e., full series) UTD definition for HIB |
| PU4313314 | UTD flag for the 4:3:1:3:3:1:4 series using the 3+ any type UTD definition for HIB |
| P_UTD431H314_ROUT_S | UTD flag for the 4:3:1:3:3:1:4 series using the routine (i.e., full series) UTD definition for HIB |

7.8.2. Rotavirus Up-To-Date Variables

The up-to-date status for Rotavirus vaccine depends on the manufacturer of the vaccines received; the requirement is two or more doses of Rotarix[®] (GSK) or three or more doses of Rotavirus vaccine of any type. Beginning with the 2009 NIS public-use data file, an up-to-date variable for Rotavirus vaccine (P_UTDROT_S) was added to indicate up-to-date status, accounting for the manufacturer (3+ Rotavirus doses of any type or 2+ Rotarix[®] doses).

Note that for this Rotavirus up-to-date variable, if the manufacturer is unknown because the provider failed to check a type box on the IHQ, it has been assumed that the Rotavirus vaccine dose is not Rotarix[®]; that is, this variable is based on a “strict” treatment of Rotavirus vaccinations of unknown type, erring on the side of classifying the child as not up-to-date.

7.9. Section 9: Provider-Reported Age-At-Vaccination Variables

This section contains variables storing the child’s age in days and months at each vaccination in the synthesized provider-reported vaccination history, along with the vaccine types of those vaccinations.

For each vaccine category, variables named **DYYY1 - DYYY9** and **YYY_AGE1 - YYY_AGE9** store the age in days and months, respectively, of the child when the vaccination was administered for up to nine vaccinations in the child’s synthesized provider-reported vaccination history, where “YYY” is the vaccine category abbreviation given in Table 6. For vaccine categories that contain multiple vaccine types, variables **XYYTY1 - XYYTY9** give the corresponding vaccine type code (see Table 6).

Unlike the vaccination count and up-to-date variables in Section 8 of the public-use data file, the variables in Section 9 include vaccinations given both before and after the household interview was completed. If desired, users can limit the Section 9 variables to only those before the household interview date by examining the corresponding Section 8 “P_NUM” variable and limiting the analysis of the Section 9 variables to only the first *n* variables, where *n* is equal to the number of vaccinations in the vaccine category before the household interview date as indicated by the corresponding “P_NUM” variable.

Users of the public-use data file should be aware that the age-at-vaccination variables included in Section 9 may contain a small number of vaccination ages that are implausible according to the recommended immunization schedules (<http://www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html>). Such ages may arise if a medical provider inadvertently records an erroneous vaccination date or if a vaccination date is incorrectly transcribed onto an IHQ. The quality control procedures of the NIS address implausible ages to every extent possible. Suspicious dates are manually reviewed and corrected if there is evidence either from the household interview or from another provider that the date is incorrect. In rare cases, however, when there is no further information with which to correct the reported vaccination date, the vaccination is treated as having actually occurred and the implausible age at vaccination persists on the data file. The data user should consider these issues in deciding how to analyze the NIS data.

7.10. Section 10: Health Insurance Module Variables

The Health Insurance Module (HIM) (Section E) was introduced in 2006 to gather information on the health insurance coverage of the child. HIM data were included in the NIS public-use data file for the first time in 2007. The NIS public-use data file contains seven variables as follows:


- INS_1 – “Is child covered by health insurance provided through employer or union?”;
- INS_2 – “Is child covered by any MEDICAID plan?”;
- INS_3 – “Is child covered by S-CHIP?”;
- INS_3A – “Is child covered by any MEDICAID plan or S-CHIP?”;
- INS_4_5 – “Is the child covered by Indian Health Service, Military Health Care, TRICARE, CHAMPUS, or CHAMP-VA?”;
- INS_6 – “Is child covered by any other health insurance or health care plan?”; and
- INS_11 - “Anytime when child was not covered by health insurance?”

Note that INS_4_5 combines the responses at questions INS_4 and INS_5. Each variable has “Yes”, “No”, “Don’t Know”, and “Refused” as response options. Also, users will encounter blanks or missing values in each variable. There are several reasons for the missing values. First, in order to reach the HIM section, the respondent must first finish Section D. Since the NIS public-use data file contains records for all respondents completing Section C, and because some of these Section C respondents did not complete Section D, some records are for respondents who did not reach the HIM. Second, there is a possibility that the respondent began the HIM but broke off the interview before finishing. Finally, there are skip patterns in the module. That is, depending on the respondent’s answers to previous questions, certain questions may be skipped. Figure 1 illustrates the flow of the eight questions used to create the seven HIM variables included in the NIS public-use data file.

The first question (INS_1) was asked of all respondents who reached the HIM. If the name of the Medicaid and S-CHIP programs were the same in the child's state, the respondent skipped to INS_3A; if the names of the Medicaid and S-CHIP programs were different in the child's state, the respondent was instead asked questions INS_2 and INS_3. **Puerto Rico respondents were asked INS_2, but then skipped to INS_5; U.S. Virgin Islands and Guam respondents skipped from INS_1 to INS_5. Thus, question INS_4 was asked of all HIM respondents outside the territories, whereas INS_5 and INS_6 were asked of all HIM respondents.** Based on the respondent's answers to previous HIM questions (some of which are not included in the public-use data file), if it was determined that the child currently had health insurance or if the child's insurance status was unknown, the respondent was asked if the child was ever uninsured at question INS_11.

Prior to 2012, the variable VFC_I indicated VFC-eligibility. The variable VFC_I was dropped from the PUF beginning 2012 due to changes to Page 1 of the NIS IHQ. Entitlement to the Vaccines for Children (VFC) program is determined by a number of factors. A child is entitled if 1) the child is covered by Medicaid, 2) the child is uninsured, 3) the child is of American Indian or Alaska Native race, or 4) the child is underinsured and has received vaccinations from a Federally Qualified Health Center (FQHC). The first three criteria were unaffected by the change to Page 1 of the IHQ. For the fourth criterion, the approach for ascertaining if a provider was a FQHC was changed on the IHQ in 2012. While CDC evaluates the accuracy of the provider-reported FQHC status, the VFC_I variable remains dropped from the PUF. Medicaid and uninsured components of VFC entitlement can be analyzed using other health insurance module variables.

Figure 1: Question Flow for the Eight Health Insurance Questions Used to Create the Health Insurance Variables Included in the PUF



8. Analytic and Reporting Guidelines

Data from the NIS public-use data file can be used to produce national, state, and estimation area estimates of vaccination coverage using the PROVWT_D weight (PROVWT_D_TERR if territories are to be included).

Information in the data file can also be used to calculate standard errors of the vaccination coverage estimates that reflect the complex sample design of the NIS. The sample is stratified by the two sample frames and the 61 estimation areas. The stratum identifier (STRATUM) and the coded household identifier (SEQNUMHH) are key variables for obtaining standard errors for estimation area, state, and national estimates of vaccination coverage rates. The estimation area variable ESTIAP15 defines mutually exclusive and exhaustive geographic areas, while STRATUM is a combination of the estimation area variable for that year and the sampling frame (landline or cell-phone).

Demographic and socioeconomic variables in the file can be used to obtain national vaccination coverage estimates for sub-groups of the population. Data users should, however, be aware that estimates for such sub-groups at the state or estimation area level will generally have large standard errors because of small sample sizes. The CDC standard for precision of sub-group estimates is that the ratio of the standard error to the estimate should be less than or equal to 0.3, and each analytic cell should contain at least 30 respondents.

8.1. Use of NIS Sampling Weights

The 2015 NIS public-use data file contains two sets of child level weights. **The RDDWT_D variable gives the household-phase weight for all children 19 through 35 months in the United States excluding territories (RDDWT_D_TERR if territories are to be included).** These weights should be used to form estimates from children with completed household interviews. The weights reflect the stratified sample design and also have been adjusted for unit non-response, for the number of telephone

lines in the household, for combining the landline and cell-phone samples, for post-stratification to population control totals, and for the exclusion of households without telephones.

The weight variables that apply to children with adequate provider data are

PROVWT_D/PROVWT_D_TERR with stratum variable STRATUM. These weights should be used to form estimates of vaccination coverage. Each child with adequate provider data (PDAT = 1) has a positive value for PROVWT_D/PROVWT_D_TERR. Starting with the 2002 file, the definition of children with adequate provider data was expanded to include unvaccinated children (as discussed in Section 2). Table 9 presents a summary of the appropriate weights and stratum variables to use for various types of analyses.

Table 9: Summary of Weights and Stratum Variables, National Immunization Survey, 2015

| Weight Variable | Population* | Sample Frame | Strata | Stratum Variable |
|------------------------|---|---------------------|--------------------------------|-------------------------|
| RDDWT_D_TERR | United States including territories | Dual Frame | Sample Type by Estimation Area | STRATUM |
| RDDWT_D | United States excluding territories | Dual Frame | Sample Type by Estimation Area | STRATUM |
| PROVWT_D_TERR | United States including territories, children with adequate provider data | Dual Frame | Sample Type by Estimation Area | STRATUM |
| PROVWT_D | United States excluding territories, children with adequate provider data | Dual Frame | Sample Type by Estimation Area | STRATUM |

* Each weight will contain a missing value for all records that are not included in the population covered by the weight.

The 2015 NIS public-use data file does not contain any provider-level weights. The NIS does not sample providers directly; rather, they are included in the survey through the children they vaccinate. A user of the file should not attempt provider-level analyses (e.g., estimate the percentage of providers in the U.S. that are private providers), because the NIS sample was not designed for that purpose.

8.2. Estimation and Analysis

8.2.1. Estimating Vaccination Coverage Rates

Vaccination coverage rates are ratio estimators, as described in the statistical literature on methods for complex sample surveys. Because of the adjustment to the sampling weights for provider-phase non-response, statistical analyses require only data from children with adequate provider data (PDAT = 1), along with their final provider sampling weights (PROVWT_D/PROVWT_D_TERR). To summarize the statistical methodology by which vaccination coverage rates and their standard errors are obtained from these data, let Y_{hij} be an indicator, for the j th child with adequate provider data in the i th sampled household in the h th stratum of the NIS sampling design, equal to 1 if the child is up-to-date according to the provider data and 0 otherwise. Also, let W_{hij} denote the value of PROVWT_D/PROVWT_D_TERR

for this child. Then, letting $\hat{Y}_h = \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} W_{hij} Y_{hij}$ and $\hat{T}_h = \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} W_{hij}$, the national estimator of the

vaccination coverage rate may be expressed as

$$\hat{\theta} = \frac{\sum_{h=1}^L \hat{Y}_h}{\sum_{h=1}^L \hat{T}_h}$$

where L denotes the number of strata, n_h denotes the number of sampled households containing children with adequate provider data in the h th stratum, and m_{hi} denotes the number of age-eligible children with adequate provider data in the i th household in the h th stratum.

Letting L instead denote the number of strata in a state, the above formula can also be used to calculate vaccination coverage rates for states (regardless of whether the state contains only one or more than one stratum).

8.2.2. *Estimating Standard Errors of Vaccination Coverage Rates*

The Taylor-series method can be used to estimate the sampling variance of vaccination coverage rates for

the U.S., the states, and estimation areas. Letting $Z_{hij} = \frac{W_{hij}(Y_{hij} - \hat{\theta})}{\sum_{h=1}^L \hat{T}_h}$, $Z_{hi} = \sum_{j=1}^{m_{hi}} Z_{hij}$, and $\bar{Z}_h = \frac{\sum_{i=1}^{n_h} Z_{hi}}{n_h}$

yields an estimator of the variance of the estimated vaccination coverage rate, $\hat{\theta}$, equal to

$$v(\hat{\theta}) = \sum_{h=1}^L \frac{n_h}{n_h - 1} \sum_{i=1}^{n_h} (Z_{hi} - \bar{Z}_h)^2$$

(Wolter, 2007) The standard error is the square root of the variance. The estimation of standard errors for estimates of vaccination coverage rates in the NIS can be implemented in specialized statistical software such as SUDAAN (Research Triangle Institute 2008), SAS (SAS Institute Inc. 2003), R (Lumley, 2010), and Stata (Stata Corporation 2009). Appendix D gives several examples of the use of SAS, R, and SUDAAN to estimate vaccination coverage rates and their standard errors for estimation areas and states. For all procedures, the option of with-replacement sampling of primary sampling units within stratum is used, because the sampling fractions for households within an estimation area are all quite small. For all estimates, the variable STRATUM is used as the stratum variable and the household identifier (SEQNUMHH) is used as the primary sampling unit identifier. The data file should be sorted first on STRATUM and then on SEQNUMHH before running the programs for SUDAAN and SAS.

8.3. **Combining Multiple Years of NIS Data**

8.3.1. *Estimation of Multi-Year Means*

With release of the 2015 NIS public-use data file, twenty years of public-use NIS data are now available. The precision of estimates of vaccination coverage for sub-domains (e.g., by race/ethnicity of child) within estimation areas or states can be improved by combining two or more years of NIS data. Data users should, however, be aware that estimates from combined years of NIS data represent an average over two or more years. Although combining several years of NIS data will yield a larger sample size for

estimation areas and states, the composition of the population in a geographic area may change over time, making interpretation of the results difficult. Furthermore, if vaccination administration schedules or vaccination coverage changes over time, the estimate of vaccination coverage for the combined time period applies to a hypothetical population that existed at the middle of the time period, making interpretation of the results even more difficult. Given the use of independent RDD samples in the NIS, it is also possible that a child could appear in more than one public-use data file.

To estimate a multi-year mean for a given NIS variable, the weights in each participating file (RDD-phase weights HY_WGT in 1995-2001, RDD_WT in 2002, WGT_RDD in 2003-2004, RDDWT in 2005-2010, RDDWT_D/RDDWT_LL in 2011, RDDWT_D/RDDWTVI_D in 2012, RDDWT_D/RDDWTVIGU_D in 2013, **and RDDWT_D/RDDWT_D_TERR in 2014-2015**; and provider-phase weights W0 in 1995-2001, WT in 2002, WGT in 2003-2004, PROVWT in 2005-2010, PROVWT_D/PROVWT_LL in 2011, PROVWT_D/PROVWTVI_D in 2012, PROVWT_D/PROVWTVIGU_D in 2013, **and PROVWT_D/PROVWT_D_TERR in 2014-2015**, should be divided by the number of years being combined. For example, if data for 2013, 2014, and 2015 for children in the United States (excluding territories) with adequate provider data are to be combined, then the weights that exclude the territories in the three files – **called PROVWT_D in each of 2013-2015** – should be divided by 3 to obtain revised weights, which should be saved as a new variable, say NEWWT. It is necessary to use NEWWT in the analysis to obtain correct weighted estimates for children aged 19 through 35 months. Furthermore, the child and household ID numbers (SEQNUMC and SEQNUMHH) in the files are unique only within a year, not across years. It is important for the user to create revised, unique ID numbers when combining data from multiple years.

The following SAS code can be used:

```
YRSEQC = 1 * (YEAR || SEQNUMC);
```

```
YRSEQHH = 1 * (YEAR || SEQNUMHH);
```

YEAR is the 4-digit year variable for the NIS data year (e.g., 2015).

To produce valid estimates of sampling variability and valid confidence intervals for multi-year coverage rates and other multi-year means, it is necessary to use specialized software such as SAS or SUDAAN.

The years 2005 to 2015 bring an important new complication for variance estimation not encountered in previous NIS years, because some traditional estimation areas were removed and other new areas were defined and introduced to the survey (see Section 2 above for more information about rotating estimation areas). The variance strata for 2004 and all prior years are defined by the variable ITRUEIAP, while the variance strata for 2005-2015 are defined by the variables ESTIAP for 2005, ESTIAP06 for 2006, ESTIAP07 for 2007, ESTIAP08 for 2008, ESTIAP09 for 2009, ESTIAP10 for 2010, STRATUM_D/ESTIAP11 for 2011, and STRATUM for 2012-2015, with STRATUM_D and STRATUM being a combination of the estimation area variable for that year and the sampling frame (landline or cell-phone). The estimation area variables ITRUEIAP, ESTIAP, and ESTIAP06-ESTIAP15 define mutually exclusive and exhaustive geographic areas. However, they are not exactly the same areas. For example, Boston and Rest of Massachusetts are each estimation areas in 2006, 2004, and all prior years, while statewide Massachusetts is an estimation area in 2005 and 2007-2015. Other areas, such as New York City and Rest of New York, are estimation areas in all years, including 2005-2015.

To make inferences concerning multi-year means, the user must take two actions. First, he/she must define and save a new stratum variable with a common name for all years included in the analysis. Second, he/she must define a common set of estimation domains that can be supported by each of the files included in the multi-year analysis. To take these actions, the user should follow the following seven-step procedure (or its equivalent):

- i. Compute and save the new, common variance-stratum variable for each year participating in the analysis. The variable should be defined by the equation

$$\text{STRATUMV} = \text{ITRUEIAP}, \text{ for children in the 2004 or prior years' public-use data files}$$

= ESTIAP, for children in the 2005 public-use data file

= ESTIAP06, for children in the 2006 public-use data file

= ESTIAP07, for children in the 2007 public-use data file

= ESTIAP08, for children in the 2008 public-use data file

= ESTIAP09, for children in the 2009 public-use data file

= ESTIAP10, for children in the 2010 public-use data file

= STRATUM_D if using PROVWT_D or
 ESTIAP11 if using PROVWT_LL, for children in the 2011 public-use data file

= STRATUM, for children in the 2012, 2013, 2014, and 2015 public-use data files

- ii. Compute and save the new, common weight variable, NEWWT, as instructed above for each year participating in the analysis.
- iii. Compute and save the new, unique child and household identification numbers, YRSEQC and YRSEQHH, as instructed above for each year participating in the analysis.
- iv. Compute and save a variable defining the common estimation domains to be studied for each year participating in the analysis. For example, one could use the CDIAP (Common Denominator Estimation Area) variable set forth in Table 10 or states as geographic domains.
- v. Merge the multiple files into one consolidated file in a format compatible with the specialized software to be used.
- vi. Sort the consolidated file by YEAR, STRATUMV, and YRSEQHH.
- vii. Run the specialized software on the consolidated file, computing estimates, variance estimates, and confidence intervals. For SUDAAN users, sampling levels or stages may be specified by the statement

NEST YEAR STRATUMV YRSEQHH / PSULEV = 3;

the specification of weights by

WEIGHT NEWWT;

and the specification of estimation domains, for example, by the two statements

```
CLASS YEAR CDIAP STATE;  
TABLES CDIAP;
```

or

```
CLASS YEAR CDIAP STATE;  
TABLES STATE;
```

8.3.2. Estimation of Multi-Year Contrasts

Considerations similar to those for multi-year means arise in the estimation of contrasts between NIS years. For example, a typical contrast of interest would be the difference between the immunization coverage parameters in 2014 and in 2015.

To make inferences concerning a multi-year contrast, the user will need to work with the original weights reported on the files and store them in a common variable. One must not divide the original weights by the number of years included in the contrast. For example, one may define the new, common weight variable as

```
NEWWT2    =    PROVWT_D/PROVWT_LL    , if the child is in the 2011 PUF.  
          =    PROVWT_D                , if the child is in the 2012, 2013, 2014, or 2015  
          PUF.
```

The user should follow the seven-step procedure set forth in the section on multi-year means, using NEWWT2 in lieu of NEWWT. In SUDAAN, the user should also specify the contrast of interest through use of a CONTRAST statement or an appropriate regression model. For example, to compare the 4:3:1:3:3:1 up-to-date estimate from 2014 to the 2015 estimate, SUDAAN users can use the following WEIGHT, VAR, and CONTRAST statements:

```
WEIGHT NEWWT2;  
VAR PU431331;  
CONTRAST YEAR = (-1 1);
```

Table 10: Cross-Walk Between ITRUEIAP, ESTIAP, ESTIAP06-ESTIAP15, and Common Denominator Estimation Area (CDIAP), National Immunization Survey, 2015

| CDIAP | Area Name | ITRUEIAP (1995-2004) | ESTIAP (2005) | ESTIAP06 (2006) | ESTIAP07 (2007) | ESTIAP08 (2008) | ESTIAP09 (2009) | ESTIAP10 (2010) | ESTIAP11 (2011) | ESTIAP12 (2012) | ESTIAP13 (2013) | ESTIAP14 (2014) | ESTIAP15 (2015) |
|------------|--------------------------|-------------------------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Alabama | | | | | | | | | | | | | |
| 20 | AL-Jefferson County | 21 | 21 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 |
| 20 | AL-Rest of State | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 |
| 74 | Alaska | 74 | 74 | 74 | 74 | 74 | 74 | 74 | 74 | 74 | 74 | 74 | 74 |
| Arizona | | | | | | | | | | | | | |
| 66 | AZ-Maricopa County | 67 | 67 | 67 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 |
| 66 | AZ-Rest of State | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 | 66 |
| 46 | Arkansas | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 | 46 |
| California | | | | | | | | | | | | | |
| 68 | CA-Fresno County | 68 | 68 | 84 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-Los Angeles County | 69 | 69 | 69 | 69 | 69 | 69 | 69 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-Northern CA | 68 | 68 | 85 | 68 | 85 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-San Diego County | 71 | 68 | 71 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-Santa Clara County | 70 | 68 | 70 | 68 | 70 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-San Bernardino County | 68 | 80 | 68 | 80 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-Alameda County | 68 | 79 | 68 | 79 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| 68 | CA-Rest of State | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 | 68 |
| Colorado | | | | | | | | | | | | | |
| 60 | CO-Denver | 60 | 81 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 |
| 60 | CO-Rest of State | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 | 60 |
| 1 | Connecticut | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 13 | Delaware | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 | 13 |
| 12 | District of Columbia | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 |
| Florida | | | | | | | | | | | | | |
| 22 | FL-Miami-Dade County | 24 | 22 | 24 | 24 | 24 | 22 | 22 | 22 | 22 | 22 | 22 | 22 |
| 22 | FL-Duval County | 23 | 23 | 23 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 |
| 22 | FL-Orange County | 22 | 22 | 22 | 22 | 91 | 22 | 22 | 22 | 22 | 22 | 22 | 22 |
| 22 | FL-Rest of State | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 | 22 |
| Georgia | | | | | | | | | | | | | |

| CDIAP | Area Name | ITRUEIAP (1995-2004) | ESTIAP (2005) | ESTIAP06 (2006) | ESTIAP07 (2007) | ESTIAP08 (2008) | ESTIAP09 (2009) | ESTIAP10 (2010) | ESTIAP11 (2011) | ESTIAP12 (2012) | ESTIAP13 (2013) | ESTIAP14 (2014) | ESTIAP15 (2015) |
|-------|-----------------------------------|-------------------------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| 25 | GA-Fulton/DeKalb Counties | 26 | 26 | 26 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 |
| 25 | GA-Rest of State | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 |
| 72 | Hawaii | 72 | 72 | 72 | 72 | 72 | 72 | 72 | 72 | 72 | 72 | 72 | 72 |
| 75 | Idaho | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 | 75 |
| | Illinois | | | | | | | | | | | | |
| 35 | IL-City of Chicago | 35 | 35 | 35 | 35 | 35 | 35 | 35 | 35 | 35 | 35 | 35 | 35 |
| 34 | IL-Madison and St. Clair Counties | 34 | 34 | 34 | 34 | 92 | 34 | 34 | 34 | 34 | 34 | 34 | 34 |
| 34 | IL-Rest of State | 34 | 34 | 34 | 34 | 34 | 34 | 34 | 34 | 34 | 34 | 34 | 34 |
| | Indiana | | | | | | | | | | | | |
| 36 | IN-Lake County | 36 | 36 | 36 | 36 | 36 | 96 | 36 | 36 | 36 | 36 | 36 | 36 |
| 36 | IN-Marion County | 37 | 36 | 37 | 37 | 36 | 37 | 36 | 36 | 36 | 36 | 36 | 36 |
| 36 | IN-Rest of State | 36 | 36 | 36 | 36 | 36 | 36 | 36 | 36 | 36 | 36 | 36 | 36 |
| 56 | Iowa | 56 | 56 | 56 | 56 | 56 | 56 | 56 | 56 | 56 | 56 | 56 | 56 |
| | Kansas | | | | | | | | | | | | |
| 57 | KS-Eastern KS | 57 | 57 | 86 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 |
| 57 | KS-Rest of State | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 |
| 27 | Kentucky | 27 | 27 | 27 | 27 | 27 | 27 | 27 | 27 | 27 | 27 | 27 | 27 |
| | Louisiana | | | | | | | | | | | | |
| 47 | LA-Orleans Parish | 48 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 |
| 47 | LA-Rest of State | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 | 47 |
| 4 | Maine | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| | Maryland | | | | | | | | | | | | |
| 14 | MD-City of Baltimore | 15 | 15 | 15 | 14 | 15 | 15 | 14 | 14 | 14 | 14 | 14 | 14 |
| 14 | MD-Prince George's County | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 103 | 14 | 14 | 14 | 14 |
| 14 | MD-Rest of State | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 |
| | Massachusetts | | | | | | | | | | | | |
| 2 | MA-City of Boston | 3 | 2 | 3 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 2 | MA-Rest of State | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| | Michigan | | | | | | | | | | | | |
| 38 | MI-City of Detroit | 39 | 39 | 39 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 |
| 38 | MI-Rest of State | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 |
| | Minnesota | | | | | | | | | | | | |
| 40 | MN-Twin Cities | 40 | 40 | 40 | 40 | 93 | 40 | 40 | 40 | 40 | 40 | 40 | 40 |

| CDIAP | Area Name | ITRUEIAP (1995-2004) | ESTIAP (2005) | ESTIAP06 (2006) | ESTIAP07 (2007) | ESTIAP08 (2008) | ESTIAP09 (2009) | ESTIAP10 (2010) | ESTIAP11 (2011) | ESTIAP12 (2012) | ESTIAP13 (2013) | ESTIAP14 (2014) | ESTIAP15 (2015) |
|-------|-----------------------------|-------------------------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| 40 | MN-Rest of State | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 | 40 |
| 28 | Mississippi | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 | 28 |
| | Missouri | | | | | | | | | | | | |
| 58 | MO-St. Louis County/City | 58 | 82 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 |
| 58 | MO-Rest of State | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 | 58 |
| 61 | Montana | 61 | 61 | 61 | 61 | 61 | 61 | 61 | 61 | 61 | 61 | 61 | 61 |
| 59 | Nebraska | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 |
| | Nevada | | | | | | | | | | | | |
| 73 | NV-Clark County | 73 | 83 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 |
| 73 | NV-Rest of State | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 | 73 |
| 5 | New Hampshire | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| | New Jersey | | | | | | | | | | | | |
| 8 | NJ-City of Newark | 9 | 9 | 9 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 8 | NJ-Rest of State | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| | New Mexico | | | | | | | | | | | | |
| 49 | NM-Southern NM | 49 | 49 | 88 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 |
| 49 | NM-Rest of State | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 | 49 |
| | New York | | | | | | | | | | | | |
| 11 | NY-City of New York | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 |
| 10 | NY-Rest of State | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 29 | North Carolina | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 | 29 |
| 62 | North Dakota | 62 | 62 | 62 | 62 | 62 | 62 | 62 | 62 | 62 | 62 | 62 | 62 |
| | Ohio | | | | | | | | | | | | |
| 41 | OH-Cuyahoga County | 42 | 42 | 42 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 |
| 41 | OH-Franklin County | 43 | 43 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 |
| 41 | OH-Rest of State | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 | 41 |
| 50 | Oklahoma | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |
| 76 | Oregon | 76 | 76 | 76 | 76 | 76 | 76 | 76 | 76 | 76 | 76 | 76 | 76 |
| | Pennsylvania | | | | | | | | | | | | |
| 16 | PA-Allegheny County | 16 | 16 | 87 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 |
| 17 | PA-Philadelphia County | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 | 17 |
| 16 | PA-Rest of State | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 | 16 |

| CDIAP | Area Name | ITRUEIAP (1995-2004) | ESTIAP (2005) | ESTIAP06 (2006) | ESTIAP07 (2007) | ESTIAP08 (2008) | ESTIAP09 (2009) | ESTIAP10 (2010) | ESTIAP11 (2011) | ESTIAP12 (2012) | ESTIAP13 (2013) | ESTIAP14 (2014) | ESTIAP15 (2015) |
|-----------|--------------------------|-------------------------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| 6 | Rhode Island | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| 30 | South Carolina | 30 | 30 | 30 | 30 | 30 | 30 | 30 | 30 | 30 | 30 | 30 | 30 |
| 63 | South Dakota | 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 | 63 |
| | Tennessee | | | | | | | | | | | | |
| 31 | TN-Davidson County | 33 | 33 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| 31 | TN-Shelby County | 32 | 32 | 32 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| 31 | TN-Rest of State | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| | Texas | | | | | | | | | | | | |
| 55 | TX-Bexar County | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 |
| 54 | TX-City of Houston | 54 | 54 | 54 | 54 | 54 | 54 | 54 | 54 | 54 | 54 | 54 | 54 |
| 51 | TX-Dallas County | 52 | 52 | 52 | 52 | 52 | 52 | 52 | 52 | 52 | 51 | 51 | 51 |
| 53 | TX-El Paso County | 53 | 53 | 53 | 53 | 53 | 53 | 53 | 53 | 53 | 53 | 53 | 53 |
| 51 | TX-Hidalgo County | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 107 |
| 51 | TX-Rest of State | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 | 51 |
| 64 | Utah | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 | 64 |
| 7 | Vermont | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 18 | Virginia | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 |
| | Washington* | | | | | | | | | | | | |
| 77 | WA-Eastern WA | 77 | 77 | 771 | 77 | 774 | 774 | 97 | 77 | 77 | 77 | 77 | 77 |
| 77 | WA-Western WA | 77 | 77 | 77 | 773 | 774 | 774 | 102 | 77 | 77 | 77 | 77 | 77 |
| 77 | WA-King County | 78 | 78 | 78 | 77 | 77 | 77 | 102 | 77 | 77 | 77 | 77 | 77 |
| 77 | WA-Rest of State | 77 | 77 | 772 | 77 | 77 | 77 | - | 77 | 77 | 77 | 77 | 77 |
| 19 | West Virginia | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| | Wisconsin | | | | | | | | | | | | |
| 44 | WI-Milwaukee County | 45 | 45 | 45 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 |
| 44 | WI-Rest of State | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 | 44 |
| 65 | Wyoming | 65 | 65 | 65 | 65 | 65 | 65 | 65 | 65 | 65 | 65 | 65 | 65 |
| - | Puerto Rico | - | - | - | - | - | - | - | - | - | - | 106 | 106 |

* The estimation area WA-Eastern WA was introduced in 2006, and while this estimation area also existed in 2010, the county definition of the area changed, making cross-year comparisons inadvisable. The estimation area WA-Western WA, introduced in 2007, presents the same issue. The counties included in the area changed (e.g., in 2010 it included King County). Analysis of Washington state data across years should use the entire state as the “Common Denominator”.

9. Summary Tables

Appendix F contains seven tables. Appendix Table F.1 lists the 61 estimation areas for the 2015 NIS by state. At the national level and for each state and estimation area, it provides the estimated population total of children aged 19 through 35 months of age in 2015, and (from 2015 NIS data collection) the number of children with completed household interviews and number of children with adequate provider data.

Appendix Tables F.2 through F.6 summarize pairs of variables: age group of child by maternal education (Appendix Table F.2), age group by family poverty status (Appendix Table F.3), race/ethnicity by family poverty status (Appendix Table F.4), age group by race/ethnicity (Appendix Table F.5), and age group by gender (Appendix Table F.6). Each of these tables gives the unweighted and weighted counts of children who have completed household interviews and the unweighted and weighted counts of children with adequate provider data.

Appendix Table F.7 presents estimates of vaccination coverage and symmetric 95% confidence intervals obtained from SUDAAN. The data user should obtain the same estimates from the 2015 NIS public-use data file.

Appendix G contains four tables and time-series charts. Table G.1 and Figure G.1 show key components of the NIS response rates and the CASRO response rates for the landline sample by year of the survey. Table G.2 and Figure G.2 show key components of the NIS response rates and the CASRO response rates for the cell-phone sample by year of the survey. Table G.3 and Figure G.3 show the CASRO response rates for the combined landline and cell-phone samples. Table G.4 and Figure G.4 show vaccination coverage estimates since 1995.

Appendix H shows the vaccine type codes used in the 2015 NIS public-use data file.

Appendix I presents key response rate components and the overall CASRO response rate by estimation area in the 2015 NIS landline and cell-phone samples.

10. Assessment of Total Survey Error

Assessing the validity of the NIS estimates of vaccination coverage is a critical and ongoing aspect of the NIS surveillance program. CDC frequently conducts evaluation studies and controlled experiments to understand the causes and impacts of sampling and nonsampling errors on the estimates and enable formulation of methodological refinements that have the demonstrated capacity to improve data quality. As landline phone use decreased and cell phone use increased dramatically over the past 10 years, and the NIS transitioned from a single-frame landline RDD sampling design to a dual-frame landline and cell phone RDD design, CDC has monitored the NIS estimates utilizing a Total Survey Error (TSE) approach.

TSE is the sum of the errors that arise at every step of a survey, including both sampling error and nonsampling errors such as coverage, nonresponse, and measurement errors (Mulry and Spencer, 1991). Pooling information from multiple evaluations of their precision and accuracy, we have conducted TSE analyses for the 2009-2012 NIS and NIS-Teen data (Molinari et al. 2011, Pineau et al. 2012; Pineau et al. 2013; Wolter et al. 2017 *expected*), including components for coverage error and nonresponse error. Figure 2 charts the estimated mean total error for NIS vaccination coverage rate estimators using the base/design weights (i.e., unadjusted for nonresponse and not calibrated to population totals) and using the final weights. For the 4:3:1:3:3:1 vaccine series, ≥ 1 MMR, and ≥ 4 DTaP, total error is smaller in 2010-2012 than in 2009. Final weights appear to reduce bias for 4:3:1:3:3:1 series vaccination coverage rate estimator but not for 1+ MMR or 4+ DTaP.

For 2012, the TSE model was extended to include a form of measurement error called “provider under-reporting” error, in addition to the coverage and nonresponse error components previously included in the model (Wolter et al. 2017 *expected*). Sometimes also called “underascertainment,” provider under-reporting error arises when a child with adequate provider data is truly vaccinated but is reported as

unvaccinated in the child’s provider-reported vaccination history. Under-reporting error can occur if the household respondent fails to nominate all of the child’s vaccination providers, if one or more of the child’s nominated vaccination providers fails to report a vaccination history for the child, or if one or more of the child’s nominated providers reports a vaccination history but fails to report all of the vaccinations the child has received. Figure 3 presents the estimated mean total error for 2012 NIS vaccination coverage rate estimators first excluding and then including the under-reporting error component in the TSE model. The addition of under-reporting error into the TSE model results in negative estimates of TSE for the two vaccines and the vaccine series; that is, the results suggest that the NIS vaccination coverage rate estimates may be lower than the true vaccination coverage rates for these vaccines and this series, and that the total error is largely due to under-reporting error.

Figure 2: Comparison of Estimated Mean Total Error for 4:3:1:3:3:1 Series, ≥ 1 MMR, and ≥ 4 DTaP by Survey Year, National Immunization Survey, 2009-2012





Figure 3: Comparison of Estimated Mean Total Error for 4:3:1:3:3:1 Series, ≥ 1 MMR, and ≥ 4 DTaP, Excluding and Including the Under-Reporting Error Component, National Immunization Survey, 2012



11. Limitations

The findings in this report are subject to at least four limitations. First, because NIS is a telephone survey, results are weighted to be representative of all children aged 19 through 35 months. Although statistical adjustments were made to account for non-response and households without telephones, some bias might remain. Second, underestimates of vaccination coverage might have resulted from the exclusive use of provider-reported vaccination histories because completeness of these records is unknown. Third, although national estimates of vaccination coverage are precise, estimates for state and local areas should be interpreted with caution because their sample sizes are smaller and their confidence intervals generally are wider than those for national estimates. Finally, analysis of trends across data years that span from 2010 and earlier to 2011-2015 are subject to potential bias that may remain after weighting adjustments because of the switch from landline to dual landline and cell-phone frames in 2011. In addition, analysis of trends across data years that span from 2011 to 2015 are subject to potential bias that may remain after weighting adjustments because of the expansions and reductions of the share of the total sample that came from the cell-phone frame across these years.

12. Citations for NIS Data

In publications, please acknowledge the original data source. The citation for the 2015 NIS public-use data file is:

U.S. Department of Health and Human Services (DHHS). National Center for Health Statistics. The 2015 National Immunization Survey, Hyattsville, MD: Centers for Disease Control and Prevention, 2016.

Information about the NIS is located at <http://www.cdc.gov/vaccines/imz-managers/nis/about.html>.

The NIS public-use data files are located at <http://www.cdc.gov/vaccines/imz-managers/nis/datasets.html>.

Please place the acronym “NIS” in the titles, keywords, or abstracts of journal articles and other publications in order to facilitate retrieval of such materials in bibliographic searches.

The following publications use NIS data, published from 2010 or later:

2016

Cardemil CV, Cullen KA, Harris L, Greby SM, Santibanez TA. Factors associated with provider reporting of child and adolescent vaccination history to Immunization Information Systems: Results from the National Immunization Survey, 2006-2012. *J Public Health Management Practice* 2016;22(3):245-254.

Chen W, Elam-Evans LD, Hill HA, Yankey D. Employment and socioeconomic factors associated with children’s up-to-date vaccination status. *Clinical Pediatrics* 2016. (in press)

Curran D, Terlinden A, Poirrier J-E, Masseria C, Krishnarajah G. Vaccine timeliness: A cost analysis of the potential implications of delayed pertussis vaccination in the US. *Pediatr Infect Dis J* 2016;35(5):542-547.

Hu T, Decker SL, Chou S-Y. Medicaid pay for performance programs and childhood immunization status. *Am J Prev Med* 2016;50(5S1):S51-S57.

Kurosky SK, Davis KL, Krishnarajah G. Completion and compliance of childhood vaccinations in the United States. *Vaccine* 2016 Jan 12;34(3):387-394.

Murphy TV, Denniston MM, Hill HA, McDonald M, Kleven M, Elam-Evans LD, Nelson NP, Iskander J, Ward JD. Progress toward eliminating hepatitis A disease in the United States. *MMWR Suppl* 2016;65(1):29-41.

Santibanez TA, Grohskopf LA, Zhai Y, Kahn KE. Complete influenza vaccination trends for children six to twenty-three months. *Pediatrics* 2016;137(3):e20153280.

Varan AK, Rodriguez-Lainz A, Hill HA, Elam-Evans LD, Yankey D, Li Q. Vaccination coverage disparities between foreign-born and U.S.-born children aged 19-35 months, United States, 2010-2012. *J Immigrant Minority Health* 2016 (in press).

Walsh B, Doherty E, O'Neill C. Since the start of the Vaccines for Children program, uptake has increased, and most disparities have decreased. *Health Affairs* 2016;35(2):356-364.

Zhao Z, Smith PJ, Hill HA. Evaluation of potentially achievable vaccination coverage with simultaneous administration of vaccines among children in the United States. *Vaccine* 2016;34:3030-3036.

2015

Crouch E, Dickes LA. A prediction model of childhood immunization rates. *Appl Health Econ Health Policy* 2015 Apr;13(2):243-251.

Dunn, AC, Black CL, Arnold J, Brodine S, Waalen J, Binkin N. Childhood vaccination coverage rates among military dependents in the United States. *Pediatrics* 2015 May;135(5):e1148-56.

Hill HA, Elam-Evans LD, Yankey D, Singleton JA, Kolasa M. National, state, and selected local area vaccination coverage among children aged 19-35 months – United States, 2014. *MMWR* 2015;64(33):889-896.

Joyce T, Reeder J. Changes in Breastfeeding among WIC participants following implementation of the new food package. *Matern Child Health J* 2015 Apr;19(4):868-76.

Smith PJ, Marcuse EK, Seward JF, Zhao Z, Orenstein WA. Children and adolescents unvaccinated against measles: geographic clustering, parents' beliefs, and missed opportunities. *Public Health Rep* 2015 Sep-Oct;130(5):485-504.

Srivastav A, Zhai Y, Santibanez TA, Kahn KE, Smith PJ, Singleton JA. Influenza vaccination coverage of Vaccine for Children (VFC)-entitled versus privately insured children, United States, 2011-2013. *Vaccine* 2015. To appear.

Wolter KM, Tao X, Montgomery R, Smith PJ. Optimum allocation for a dual-frame telephone survey. *Survey Methodology* 2015 Dec;41(2):389-401.

2014

Elam-Evans LD, Yankey D, Singleton JA, Kolasa M. National, state, and selected local area vaccination coverage among children aged 19-35 months - United States, 2013. *MMWR* 2014;63(34):741-8.

Johnson NB, Hayes LD, Brown K, Hoo EC, Ethier KA. CDC National Health Report: leading causes of morbidity and mortality and associated behavioral risk and protective factors--United States, 2005-2013 *MMWR* 2014;63(Suppl-4):1-27.

Santibanez TA, Lu PJ, O'Halloran A, Meghani A, Grabowsky M, Singleton JA. Trends in childhood influenza vaccination coverage—U.S., 2004-2012. *Public Health Rep* 2014;129(5):417-27.

Thomas TN, Kolasa MS, Zhang F, Shefer AM. Assessing immunization interventions in the Women, Infants, and Children (WIC) Program. *Am J Prev Med* 2014 Sep 10. [Epub ahead of print: DOI: 10.1016/j.amepre.2014.06.017]

Walker AT, Smith PJ, Kolasa M. Reduction of racial/ethnic disparities in vaccination coverage, 1995-2011. *MMWR Surveill Summ* 2014 Apr 18;63 Suppl 1:7-12.

Whitney CG, Zhou F, Singleton J, Schuchat A. Benefits from immunization during the vaccines for children program era - United States, 1994-2013. *MMWR* 2014 Apr 25;63(16):352-5.

Yang YT, Debold V. A longitudinal analysis of the effect of nonmedical exemption law and vaccine uptake on vaccine-targeted disease rates. *Am J Public Health* 2014;104:371-377.

Zhao Z, Smith PJ, Yankey D, Copeland KR. Calculating adjusted survival functions for complex sample survey data and application to vaccination coverage studies with National Immunization Survey. *British Journal of Mathematics & Computer Science* 2014;4(18):2686-2698.

2013

Allen JA, Li R, Scanlon KS, Perrine CG, Chen J, Odom E, Black C. Progress in increasing breastfeeding and reducing racial/ethnic differences — United States, 2000–2008 births. *MMWR* 2013 Feb 8;62(5):77-80.

Black CL, Yankey D, Kolasa M. National, state, and local area vaccination coverage among children aged 19–35 months — United States, 2012. *MMWR* 2013 Sept 13;62(36):733-740.

Schuller KA, Probst JC. Factors associated with influenza vaccination among US children in 2008. *J Infect Public Health* 2013;6(2):80-88.

Zhao Z, Murphy TV. Which newborns missed the hepatitis B birth dose vaccination among U.S. children? *Preventive Medicine*. 2013, 57, 613-7.

Zhao Z, Smith P. Trends in vaccination coverage disparities among children, United States, 2001-2010. *Vaccine*. 2013, 31(19), 2324-2327

2012

Black CL, Yankey D, Kolasa M. National, state, and local area vaccination coverage among children aged 19-35 months – United States, 2011. *MMWR* 2012 Sept 7;61(35):689-696.

Bundy DG, Solomon BS, Kim JM, Miller MR. Accuracy and usefulness of the HEDIS childhood immunization measures. *Pediatrics* 2012;129(4):648-656.

Groom AV, Santibanez TA, Bryan RT. Vaccination coverage among American Indian and Alaska Native Children, 2006 – 2010. *Pediatrics* 2012;130(6):e1592-e1599.

Jensen E. Participation in the Supplemental Nutrition Program for Women, Infants, and Children (WIC) and breastfeeding: national, regional, and state level analyses. *Matern Child Health J* 2012;16:624-631.

Ransom J, Schaff K, Kan L. Is there an association between local health department organizational and administrative factors and childhood immunization coverage rates? *J Health Hum Serv Adm* 2012 Spring;34(4):418-455.

Santibanez TA, Shefer A, Briere EC, Cohn AC, Groom AV. Effects of a nationwide Hib vaccine shortage on vaccination coverage. *Vaccine* 2012;30:941-947.

Thompson KM, Wallace GS, Tebbens RJD, Smith PJ, Barskey AE, Pallansch MA, Gallagher KM, Alexander JP, Armstrong GL, Cochi SL, Wassilak SGF. Trends in the risk of U.S. polio outbreaks and poliovirus vaccine availability for response. *Public Health Rep* 2012;127:23-37.

Zhao Z, Murphy TV. The association of hepatitis B vaccine supply policy with timing of receipt of the first dose of hepatitis B vaccination. *Open Journal of Statistics* 2012;2:429-434.

2011

Black CL, Wooten KG, Yankey D, Kolasa M. National and state vaccination coverage among children aged 19-35 months – United States, 2010. *MMWR* 2011 Sept 2;60(34):1157-1163.

Byrd KK, Santibanez TA, Chaves SS. Predictors of hepatitis A vaccination among young children in the United States. *Vaccine* 2011;29:3254-3259.

Dozier AM, McKee KS. State breastfeeding worksite statutes... breastfeeding rates ... and..... *Breastfeed Med* 2011;6(5):319-324.

Flaherman VJ, Chien AT, McCulloch CE, Dudley RA. Breastfeeding rates differ significantly by method used: a cause for concern for public health measurement. *Breastfeed Med* 2011;6(1):31-35.

Molinari NM, Wolter KM, Skalland B, Montgomery R, Khare M, Smith PJ, Barron ML, Copeland K, Santos K, Singleton JA. Quantifying bias in a health survey: modeling total survey error in the National Immunization Survey. *Stat Med* 2011;30:505-514.

Smith PJ, Humiston SG, Marcuse EK, Zhao Z, Dorell CG, Howes C, Hibbs B. Parental delay or refusal of vaccine doses, childhood vaccination coverage at 24 months of age, and the health belief model. *Public Health Rep.* 2011 Jul-Aug;126 Suppl 2:135-46.

Smith PJ, Lindley MC, Rodewald LE. Vaccination coverage among U.S. Children aged 19-35 months entitled by the Vaccines for Children Program, 2009. *Public Health Rep* 2011 Jul-Aug;126 Suppl 2:109-23.

Smith PJ, Singleton JA. County-level trends in vaccination coverage among children aged 19-35 months – United States, 1995 – 2008. *MMWR Surveill Summ* 2011 Apr 29;60(4):1-86.

Smith PJ, Singleton JA. County-level trends in vaccination coverage for selected counties in the U.S., children aged 19-35 months, United States, 1995-2008: attainment of the healthy people 2010 objectives and association with county-level indices of access to care, economic conditions, and demographic composition. *Morbidity and Mortality Weekly Report. Surveillance Summaries.* 2011;60(4):1-86.

Smith PJ, Wood D, Darden PM. Highlights of historical events leading to national surveillance of vaccination coverage in the United States. *Public Health Reports.* 2011;(Suppl 2):3-12.

Zhao Z. Power of tests for comparing trend curves with application to National Immunization Survey (NIS). *Stat Med* 2011;30:531-540.

Zhao Z, Murphy TV, Jacques-Carroll L. Progress in newborn hepatitis B vaccination by birth year cohorts – 1998-2007, USA. *Vaccine* 2011;30:14-20.

2010

Cohen SA, Ahmed S, Klassen AC, Agree EM, Louis TA, Naumova EN. Childhood Hib vaccination and pneumonia and influenza burden in US seniors. *Vaccine* 2010;28:4462-4469.

Committee on Practice and Ambulatory Medicine and Council on Community Pediatrics. Increasing immunization coverage. *Pediatrics* 2010;125:1295-1304.

Groom H, Kennedy A, Evans V, Fasano N. Qualitative analysis of immunization programs with most improved childhood vaccination coverage from 2001 to 2004. *J Public Health Management Practice* 2010;16(1):E1-E8.

Kennedy A, Groom H, Evans V, Fasano N. A qualitative analysis of immunization programs with sustained high coverage, 200-2005. *J Public Health Management Practice* 2010;16(1):E9-E17.

McElligott JT, Darden PM. Are patient-held vaccination records associated with improved vaccination coverage rates? *Pediatrics* 2010;125(3):e467-e472.

Mennito SH, Darden PM. Impact of practice policies on pediatric immunization rates. *J Pediatr* 2010;156:618-622.

Santibanez TA, Singleton JA, Shefer A, Cohn A. Changes in measurement of *Haemophilus influenzae* serotype b (Hib) vaccination coverage – National Immunization Survey, United States, 2009. *MMWR* 2010 Aug 27;59(33):1069-1072.

Scanlon KS, Grummer-Strawn L, Li R, Chen J. Racial and ethnic differences in breastfeeding initiation and duration, by state – National Immunization Survey, United States, 2004-2008. *MMWR* 2010 Mar 26;59(11):327-334.

Smith PJ, Humiston SG, Parnell T, Vannice KS, Salmon DA. The association between intentional delay of vaccine administration and timely childhood vaccination coverage. *Public Health Rep* 2010;125:534-541.

Wooten KG, Kolasa M, Singleton JA, Shefer A. National, state, and local area vaccination coverage among children aged 19-35 months – United States, 2009. *MMWR* 2010 Sep 17;59(36):1171-1177.

Zhao Z, Luman ET. Progress Toward eliminating disparities in vaccination coverage among U.S. children, 2000–2008. *Am J Prev Med* 2010;38(2):127–137.

13. References

The American Association for Public Opinion Research (2015). *Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys*.

Blumberg, S.J. and Luke, J.V. (2016). Wireless substitution: Early release of estimates from the National Health Interview Survey, July-December 2015. National Center for Health Statistics. <http://www.cdc.gov/nchs/data/nhis/earlyrelease/wireless201605.pdf>

Blumberg, S.J., Luke, J.V., Ganesh, N., Davern, M.E., Boudreaux, M.H. and Soderberg, K. (2011). Wireless substitution: State-level estimates from the National Health Interview Survey, January 2007–June 2010. National Center for Health Statistics. <http://www.cdc.gov/nchs/data/nhsr/nhsr039.pdf>

Brick, J.M. and Kalton, G. (1996). Handling missing data in survey research. *Statistical Methods in Medical Research*, 5:215–238.

Council of American Survey Research Organizations (1982). On the Definition of Response Rates: A Special Report of the CASRO Task Force on Completion Rates. Council of American Survey Research Organizations: <http://www.casro.org>.

Centers for Disease Control and Prevention (1994). Reported vaccine-preventable diseases - United States, 1993, and the Childhood Immunization Initiative. *MMWR*, 43:57-60.

Centers for Disease Control and Prevention (2000). Prevention of pneumococcal disease among infants and young children using a pneumococcal conjugate vaccine. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*, 49(RR-9):1-35.

Centers for Disease Control and Prevention (2002). *National Immunization Survey: Guide to Quality Control Procedures*. <http://www.cdc.gov/nchs/data/nis/qcman.pdf>.

Centers for Disease Control and Prevention (2003). Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*, 52(RR-8):1-34.

Centers for Disease Control and Prevention (2010). Changes in measurement of *Haemophilus influenzae* serotype b (Hib) vaccination coverage—National Immunization Survey, United States, 2009. *MMWR* 2010;59:1069--72. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5933a3.htm>.

Coronado, V.G., Maes, E.F., Rodewald, L.E., Chu, S., Battaglia, M.P., Hoaglin, D.C., Merced, N.L., Yusuf, H., Cordero, J.F., and Orenstein, W.A. (2000). Risk factors for underimmunization among 19-35 month-old children in the United States: National Immunization Survey, July 1996-June 1998. Unpublished manuscript, Centers for Disease Control and Prevention, Atlanta.

Deming, W.E. (1943). *Statistical Adjustment of Data*. New York: Wiley.

Ezzati-Rice, T.M., Zell, E.R., Battaglia, M.P., Ching, P.L.Y.H., and Wright, R.A. (1995). The design of the National Immunization Survey. *1995 Proceedings of the Section on Survey Research Methods*, Alexandria, VA: American Statistical Association, pp. 668-672.

- Ford, B.L. (1983). An overview of hot-deck procedures, in: *Incomplete data in sample surveys*, Madow W. G., Olkin I., Rubin D. B. (Eds.), Academic Press, New York, pp. 185-207.
- Khare, M., Battaglia, M.P., Huggins, V.J., Stokley, S., Hoaglin, D.C., Wright, R.A., and Rodén, A.-S. (2000). Accuracy of vaccination dates reported by immunization providers in the National Immunization Survey. *2000 Proceedings of the Section on Survey Research Methods*. Alexandria, VA: American Statistical Association, pp. 665-670.
- Khare, M., Battaglia, M.P., Stokley, S., Wright, R.A., and Huggins, V.J. (2001). Quality of immunization histories reported in the National Immunization Survey. *Proceedings of the International Conference on Quality in Official Statistics (CD-ROM)*. Stockholm: Statistics Sweden.
- Lepkowski, J.M. (1988). Telephone sampling methods in the United States. *Telephone Survey Methodology*. Edited by Groves, R.M., Biemer, P.P., Lyberg, L.E., Massey, J.T., Nicholls, W.L., and Waksberg, J. New York: John Wiley & Sons, pp. 73-98.
- Lumley, T. (2010). Survey Analysis in R. <http://faculty.washington.edu/tlumley/survey/>
- Molinari, N., Wolter, K., Skalland, B., Montgomery, R., Khare, M., Smith, P., and Singleton, J. (2011). *Quantifying Bias in a Health Survey: Modeling Total Survey Error in the National Immunization Survey*. *Statistics in Medicine*, 505-515.
- Mulry, M. H. and Spencer, B. C. (1991). Total Error in PES Estimates of Population. *Journal of the American Statistical Association*, 839-863.
- National Center for Health Statistics (1999). *National Health Interview Survey: Research for the 1995-2004 Redesign*. Vital and Health Statistics, Series 2, No. 126 (DHHS publication no. (PHS) 99-1326). Hyattsville, MD: National Center for Health Statistics.
- National Center for Health Statistics. (2011). *Natality Data, Public-Use Data Files*. http://www.cdc.gov/nchs/data_access/vitalstatsonline.htm.
- National Center for Health Statistics. (2012). *Natality Data, Public-Use Data Files*. http://www.cdc.gov/nchs/data_access/vitalstatsonline.htm.
- National Center for Health Statistics. (2013). *Natality Data, Public-Use Data Files*. http://www.cdc.gov/nchs/data_access/vitalstatsonline.htm.
- National Center for Health Statistics. (2015). *National Immunization Survey 2014 Public-Use Data File: Documentation, Codebook and Frequencies*. Hyattsville, MD.
- National Center for Immunization and Respiratory Diseases. (2016). *National Immunization Survey 2015 Public-Use Data File: Documentation, Codebook and Frequencies*. Hyattsville, MD.
- Pineau, V., Wolter, K., Skalland, B., Zeng, W., Zhao, Z. and Khare, M. (2012). *Modeling Total Survey error in the 2010 National Immunization survey (NIS): Pre-School Children and Teens*. Presented at the 2012 American Statistical Association (ASA) Joint Statistical Meetings, San Diego, CA.
- Pineau, V., Wolter, K., Skalland, B., Zeng, W., Black, C., Dorell, C., Khare, M., and Yankey, D. (2013). *Modeling Total Survey Error in the 2011 national Immunization Survey (NIS): Pre-School Children and*

Teens. Presented at the 2013 American Statistical Association (ASA) Joint Statistical Meetings, Montreal, Canada.

Research Triangle Institute (2008). *SUDAAN Language Manual, Release 9.0*. Research Triangle Park, NC: Research Triangle Institute.

Robinson, C.L. (2016). Advisory committee on immunization practices recommended immunization schedules for persons aged 0 through 18 years--United States, 2016. *MMWR* 65(4): 86-87.

Rosenbaum, P.R. (1987). Model-based direct adjustment. *Journal of the American Statistical Association*, 82:387-394.

Rosenbaum, P.R. and Rubin, D.B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70:41-55.

Rosenbaum, P.R. and Rubin, D.B. (1984). Reducing bias in observational studies using subclassification on the propensity score. *Journal of the American Statistical Association*, 79:516-534.

SAS Institute Inc. (2003). *SAS/STAT User's Guide, Version 8*. Cary, NC: SAS Institute Inc.

Smith, P.J., Battaglia, M.P., Huggins, V.J., Hoaglin, D.C., Rodén, A.-S., Khare, M., Ezzati-Rice, T.M., and Wright, R.A. (2001a). Overview of the sampling design and statistical methods used in the National Immunization Survey. *American Journal of Preventive Medicine*, 20(4S):17-24.

Smith, P.J., Rao, J.N.K., Battaglia, M.P., Ezzati-Rice, T.M., Daniels, D., and Khare, M. (2001b). *Compensating for Provider Non-response Using Response Propensities to Form Adjustment Cells: The National Immunization Survey*. Vital and Health Statistics, Series 2, No. 133 (DHHS publication no. (PHS) 2001-1333). Hyattsville, MD: National Center for Health Statistics.

Smith, P.J., Hoaglin, D.C., Battaglia, M.P., Khare, M., and Barker, L.E. (2005). *Statistical Methodology of the National Immunization Survey: 1994-2002*. National Center for Health Statistics. Vital Health Stat 2(138).

StataCorp (2005). *Stata Statistical Software: Release 9*. College Station, TX: StataCorp LP.

Wall, T.P., Kochanek, K.M., Fitti, J.E., and Zell, E.R. (1995). The use of real time translation services in RDD telephone surveys. Presented at the 1995 Conference of the American Association for Public Opinion Research, Fort Lauderdale, FL.

Wolter, K. M. (2007). *Introduction to Variance Estimation*. New York, NY: Springer-Verlag.

Wolter, K., Pineau, V., Skalland, B., Zeng, W., Singleton, J., Khare, M., Zhao, Z., Yankey, D., and Smith, P. (2017 *expected*). Total survey error assessment for socio-demographic subgroups in the 2012 U.S. National Immunization Survey. In Biemer, P., De Leeuw, E., Edwards, B., Kreuter, F., Lyberg, L., Tucker, C., and West, B. (Eds.) *Total Survey Error in Practice*, John Wiley & Sons, Inc., Hoboken, NJ, USA..

Zell, E.R., Ezzati-Rice, T.M., Battaglia, M.P., and Wright, R.A. (2000). National Immunization Survey: The methodology of a vaccination surveillance system. *Public Health Reports*, 115(1):65-77.

Appendix A: Glossary of Abbreviations and Terms

| | |
|---------------------------------|--|
| 3:3:1 | The series of 3 or more DTaP vaccinations, 3 or more polio vaccinations, and 1 or more MCV vaccinations |
| 4:3:1 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, and 1 or more MCV vaccinations |
| 4:3:1:3 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, and 3 or more Hib vaccinations of any type |
| 4:3:1:3* (routine Hib) | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, and 3 or 4 Hib vaccinations depending on manufacturer (routine recommendation) |
| 4:3:1:3:3 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or more Hib vaccinations of any type, and 3 or more hepatitis B vaccinations |
| 4:3:1:3*:3 (routine Hib) | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or 4 Hib vaccinations depending on manufacturer (routine recommendation), and 3 or more hepatitis B vaccinations |
| 4:3:1:3:3:1 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or more Hib vaccinations of any type, 3 or more hepatitis B vaccinations, and 1 or more varicella vaccinations given at age 12 months or older |
| 4:3:1:3*:3:1 (routine Hib) | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or 4 Hib vaccinations depending on manufacturer (routine recommendation), 3 or more hepatitis B vaccinations, and 1 or more varicella vaccinations given at age 12 months or older |
| 4:3:1:3:3:1:3 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or more Hib vaccinations of any type, 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 3 or more pneumococcal vaccinations |
| 4:3:1:3*:3:1:3 (routine Hib) | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or 4 Hib vaccinations depending on manufacturer (routine recommendation), 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 3 or more pneumococcal vaccinations |
| 4:3:1:3:3:1:4 | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or more Hib vaccinations of any type, 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 4 or more pneumococcal vaccinations |

| | |
|---------------------------------|--|
| 4:3:1:3*:3:1:4 (routine Hib) | The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MCV vaccinations, 3 or 4 Hib vaccinations depending on manufacturer (routine recommendation), 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 4 or more pneumococcal vaccinations |
| CATI | Computer-assisted telephone interviewing |
| CDC | Centers for Disease Control and Prevention |
| CII | Childhood Immunization Initiative |
| DOB | Date of birth |
| DTaP | Diphtheria and tetanus toxoids and acellular pertussis |
| DTP | Diphtheria and tetanus toxoids and pertussis |
| DT | Diphtheria and tetanus toxoids |
| FLU | Seasonal influenza |
| H1N | Monovalent 2009 H1N1 influenza |
| Hep A | Hepatitis A |
| Hep B | Hepatitis B |
| Hib | <i>Haemophilus influenzae</i> type b |
| Hib routine recommendation | Four or more doses of Hib vaccine of any type, or two or more doses of Hib vaccine of Merck types followed by one dose of Hib vaccine of any type |
| Hib shortage recommendation | Three or more doses of Hib vaccine of any type or two or more doses of Hib vaccine of Merck types |
| IAP | Immunization Action Plan |
| IHQ | Immunization history questionnaire |
| IPV | Inactivated poliovirus vaccine |
| MCV | Measles-containing vaccine |
| MMR | Measles, mumps, and rubella |
| NCHS | National Center for Health Statistics |
| NCIRD | National Center for Immunization and Respiratory Diseases |
| NIS | National Immunization Survey |

| | |
|------|----------------------------------|
| NHIS | National Health Interview Survey |
| NIP | National Immunization Program |
| OPV | Oral poliovirus vaccine |
| PCV | Pneumococcal conjugate vaccine |
| PRC | Provider Record Check |
| PUF | Public-use (Data) File |
| RDD | Random digit dialing |
| ROT | Rotavirus |
| SC | Shot card |
| UTD | Up-to-date |
| VFC | Vaccines for Children |
| VRC | Varicella |

Appendix B: Summary Statistics for Sampling Weights by Estimation Area

Table B.1: Distribution of Dual-Frame Sampling Weights* for Children with Completed Household Interviews, National Immunization Survey, 2015

| State/Estimation Area | n | Sum | Minimum | Maximum | Mean | Coefficient of Variation |
|-----------------------|--------|------------|---------|---------|---------|--------------------------|
| U.S. National† | 26,975 | 5701539.58 | 2.49 | 5856.42 | 211.36 | 152.40 |
| Alabama | 494 | 84042.20 | 18.12 | 534.23 | 170.13 | 69.71 |
| Alaska | 498 | 15311.61 | 9.33 | 71.47 | 30.75 | 44.26 |
| Arizona | 479 | 125138.52 | 20.05 | 785.29 | 261.25 | 73.59 |
| Arkansas | 419 | 54196.27 | 15.97 | 408.59 | 129.35 | 78.00 |
| California | 646 | 724479.75 | 2.72 | 5856.42 | 1121.49 | 116.29 |
| Colorado | 440 | 95511.01 | 6.16 | 608.81 | 217.07 | 63.54 |
| Connecticut | 469 | 54155.77 | 24.99 | 313.84 | 115.47 | 62.84 |
| Delaware | 410 | 15716.58 | 10.98 | 97.89 | 38.33 | 51.43 |
| District of Columbia | 576 | 12398.05 | 5.04 | 57.95 | 21.52 | 58.59 |
| Florida | 546 | 315013.51 | 4.08 | 2155.42 | 576.95 | 81.42 |
| Georgia | 540 | 186272.22 | 5.73 | 1131.67 | 344.95 | 75.82 |
| Hawaii | 469 | 25965.60 | 10.63 | 158.15 | 55.36 | 46.95 |
| Idaho | 985 | 222541.86 | 7.67 | 647.00 | 225.93 | 60.61 |
| Illinois | 320 | 56511.16 | 9.06 | 486.16 | 176.60 | 54.92 |
| IL-City of Chicago | 665 | 166030.70 | 7.67 | 647.00 | 249.67 | 58.77 |
| IL-Rest of State | 342 | 30722.98 | 11.20 | 199.78 | 89.83 | 35.78 |
| Indiana | 426 | 121057.24 | 9.40 | 779.35 | 284.17 | 56.75 |
| Iowa | 444 | 55272.78 | 15.93 | 339.31 | 124.49 | 52.43 |
| Kansas | 367 | 58367.35 | 21.06 | 383.41 | 159.04 | 47.62 |
| Kentucky | 437 | 78299.08 | 16.10 | 446.13 | 179.17 | 53.08 |
| Louisiana | 427 | 88668.64 | 12.01 | 569.50 | 207.65 | 56.60 |
| Maine | 386 | 18614.44 | 9.50 | 118.10 | 48.22 | 64.51 |
| Maryland | 658 | 105465.39 | 17.19 | 640.48 | 160.28 | 97.02 |
| Massachusetts | 521 | 102929.55 | 7.45 | 635.30 | 197.56 | 69.43 |
| Michigan | 427 | 162006.92 | 7.12 | 1117.42 | 379.41 | 61.91 |
| Minnesota | 521 | 100888.34 | 8.01 | 508.60 | 193.64 | 55.95 |
| Mississippi | 464 | 54235.40 | 14.16 | 327.67 | 116.89 | 64.91 |
| Missouri | 401 | 106676.79 | 6.80 | 771.46 | 266.03 | 62.75 |
| Montana | 390 | 17100.50 | 7.50 | 129.34 | 43.85 | 63.47 |
| Nebraska | 1,119 | 334940.01 | 7.43 | 840.98 | 299.32 | 54.57 |
| Nevada | 564 | 167964.60 | 7.83 | 768.38 | 297.81 | 50.26 |
| New Hampshire | 555 | 166975.41 | 7.43 | 840.98 | 300.86 | 58.58 |
| New Jersey | 374 | 37665.32 | 9.25 | 269.45 | 100.71 | 51.69 |
| New Mexico | 377 | 51393.29 | 15.03 | 387.23 | 136.32 | 54.59 |
| New York | 372 | 19116.11 | 14.11 | 118.93 | 51.39 | 44.47 |
| NY-City of New York | 557 | 153666.94 | 6.28 | 784.93 | 275.88 | 68.83 |
| NY-Rest of State | 400 | 39050.01 | 9.05 | 309.14 | 97.63 | 69.32 |
| North Carolina | 483 | 173794.97 | 5.61 | 1088.16 | 359.82 | 68.51 |
| North Dakota | 398 | 15261.69 | 12.76 | 99.81 | 38.35 | 48.98 |
| Ohio | 418 | 198461.09 | 52.54 | 1299.35 | 474.79 | 56.85 |
| Oklahoma | 376 | 75919.74 | 13.69 | 557.45 | 201.91 | 57.74 |
| Oregon | 341 | 67036.19 | 19.96 | 488.97 | 196.59 | 50.53 |
| Pennsylvania | 1,039 | 204791.65 | 4.12 | 1172.48 | 197.10 | 128.81 |

| State/Estimation Area | n | Sum | Minimum | Maximum | Mean | Coefficient of Variation |
|------------------------------|----------|------------|----------------|----------------|-------------|---------------------------------|
| PA-Philadelphia County | 487 | 33050.31 | 8.02 | 171.48 | 67.87 | 52.06 |
| PA-Rest of State | 552 | 171741.34 | 4.12 | 1172.48 | 311.13 | 97.77 |
| Rhode Island | 463 | 16313.23 | 7.31 | 89.78 | 35.23 | 56.77 |
| South Carolina | 487 | 82138.06 | 8.07 | 496.30 | 168.66 | 66.60 |
| South Dakota | 399 | 17989.42 | 10.41 | 135.79 | 45.09 | 58.72 |
| Tennessee | 2,586 | 568417.14 | 6.68 | 1332.10 | 219.81 | 123.81 |
| Texas | 428 | 39136.32 | 20.67 | 232.89 | 91.44 | 59.10 |
| TX-Bexar County | 360 | 71255.42 | 6.68 | 558.53 | 197.93 | 59.88 |
| TX-City of Houston | 348 | 20393.13 | 9.42 | 160.04 | 58.60 | 53.75 |
| TX-El Paso County | 370 | 23985.20 | 13.14 | 194.24 | 64.82 | 48.42 |
| TX-Hidalgo County | 1,080 | 413647.08 | 12.43 | 1332.10 | 383.01 | 90.81 |
| TX-Rest of State | 372 | 117279.88 | 7.90 | 856.16 | 315.27 | 56.87 |
| Utah | 379 | 73265.59 | 6.44 | 594.20 | 193.31 | 63.54 |
| Vermont | 409 | 8540.46 | 6.88 | 53.12 | 20.88 | 45.37 |
| Virginia | 655 | 149555.84 | 2.49 | 1134.16 | 228.33 | 138.71 |
| Washington | 526 | 127236.49 | 5.51 | 757.19 | 241.89 | 70.36 |
| West Virginia | 453 | 29917.36 | 15.15 | 179.29 | 66.04 | 56.79 |
| Wisconsin | 389 | 98535.31 | 11.40 | 648.83 | 253.30 | 55.71 |
| Wyoming | 421 | 10205.41 | 6.09 | 59.66 | 24.24 | 51.41 |
| U.S. Virgin Islands | 580 | 1956.18 | 1.00 | 9.75 | 3.37 | 60.70 |
| Guam | 467 | 5058.12 | 2.89 | 29.73 | 10.83 | 51.56 |
| Puerto Rico | 617 | 50147.68 | 7.58 | 241.03 | 81.28 | 64.64 |

* Distribution of RDDWT_D_TERR.

† Excludes territories.

Table B.2: Distribution of Dual-Frame Sampling Weights* for Children with Adequate Provider Data, National Immunization Survey, 2015

| State/Estimation Area | n | Sum | Minimum | Maximum | Mean | Coefficient of Variation |
|------------------------|--------|------------|---------|----------|---------|--------------------------|
| U.S. National† | 15,167 | 5701539.58 | 2.88 | 10146.43 | 375.92 | 157.96 |
| Alabama | 272 | 84042.20 | 34.01 | 1037.67 | 308.98 | 81.00 |
| Alaska | 295 | 15311.61 | 10.94 | 139.53 | 51.90 | 55.93 |
| Arizona | 266 | 125138.52 | 32.25 | 1471.44 | 470.45 | 71.58 |
| Arkansas | 229 | 54196.27 | 24.97 | 760.01 | 236.66 | 69.68 |
| California | 345 | 724479.75 | 5.81 | 10146.43 | 2099.94 | 115.04 |
| Colorado | 246 | 95511.01 | 13.05 | 1155.94 | 388.26 | 65.45 |
| Connecticut | 250 | 54155.77 | 31.47 | 662.34 | 216.62 | 70.73 |
| Delaware | 232 | 15716.58 | 18.36 | 187.50 | 67.74 | 52.49 |
| District of Columbia | 321 | 12398.05 | 8.66 | 112.74 | 38.62 | 67.91 |
| Florida | 273 | 315013.51 | 12.46 | 4150.20 | 1153.90 | 79.82 |
| Georgia | 286 | 186272.22 | 7.83 | 2270.88 | 651.30 | 83.68 |
| Hawaii | 261 | 25965.60 | 14.71 | 278.73 | 99.49 | 57.61 |
| Idaho | 559 | 222541.86 | 15.89 | 1372.45 | 398.11 | 64.58 |
| Illinois | 188 | 56511.16 | 15.89 | 824.42 | 300.59 | 54.61 |
| IL-City of Chicago | 371 | 166030.70 | 19.86 | 1372.45 | 447.52 | 62.72 |
| IL-Rest of State | 209 | 30722.98 | 26.86 | 371.18 | 147.00 | 43.79 |
| Indiana | 231 | 121057.24 | 13.71 | 1517.24 | 524.06 | 65.42 |
| Iowa | 279 | 55272.78 | 26.35 | 553.23 | 198.11 | 53.60 |
| Kansas | 228 | 58367.35 | 27.98 | 677.68 | 256.00 | 50.64 |
| Kentucky | 241 | 78299.08 | 45.55 | 946.74 | 324.89 | 59.03 |
| Louisiana | 223 | 88668.64 | 22.24 | 1199.60 | 397.62 | 72.67 |
| Maine | 221 | 18614.44 | 14.04 | 286.16 | 84.23 | 73.13 |
| Maryland | 361 | 105465.39 | 36.14 | 1095.56 | 292.15 | 95.10 |
| Massachusetts | 307 | 102929.55 | 14.98 | 1112.50 | 335.28 | 71.32 |
| Michigan | 254 | 162006.92 | 19.74 | 2044.04 | 637.82 | 63.45 |
| Minnesota | 289 | 100888.34 | 14.66 | 1058.13 | 349.09 | 60.95 |
| Mississippi | 233 | 54235.40 | 14.79 | 780.21 | 232.77 | 78.42 |
| Missouri | 226 | 106676.79 | 10.41 | 1408.63 | 472.02 | 65.73 |
| Montana | 251 | 17100.50 | 12.72 | 201.39 | 68.13 | 63.96 |
| Nebraska | 566 | 334940.01 | 13.11 | 1845.71 | 591.77 | 62.49 |
| Nevada | 291 | 167964.60 | 13.11 | 1845.71 | 577.20 | 62.04 |
| New Hampshire | 275 | 166975.41 | 15.26 | 1822.61 | 607.18 | 62.89 |
| New Jersey | 231 | 37665.32 | 15.37 | 451.28 | 163.05 | 56.41 |
| New Mexico | 214 | 51393.29 | 43.71 | 706.19 | 240.16 | 56.18 |
| New York | 195 | 19116.11 | 29.15 | 234.33 | 98.03 | 46.21 |
| NY-City of New York | 272 | 153666.94 | 15.01 | 1839.92 | 564.95 | 77.49 |
| NY-Rest of State | 244 | 39050.01 | 12.60 | 543.39 | 160.04 | 72.52 |
| North Carolina | 261 | 173794.97 | 14.03 | 2334.24 | 665.88 | 69.71 |
| North Dakota | 234 | 15261.69 | 16.76 | 174.98 | 65.22 | 51.48 |
| Ohio | 219 | 198461.09 | 102.16 | 2922.31 | 906.22 | 68.63 |
| Oklahoma | 220 | 75919.74 | 18.92 | 1049.34 | 345.09 | 63.84 |
| Oregon | 192 | 67036.19 | 35.96 | 1023.25 | 349.15 | 55.53 |
| Pennsylvania | 575 | 204791.65 | 13.58 | 1934.41 | 356.16 | 123.75 |
| PA-Philadelphia County | 272 | 33050.31 | 18.32 | 347.32 | 121.51 | 59.47 |
| PA-Rest of State | 303 | 171741.34 | 13.58 | 1934.41 | 566.80 | 91.75 |
| Rhode Island | 272 | 16313.23 | 9.54 | 176.80 | 59.98 | 60.77 |
| South Carolina | 259 | 82138.06 | 18.63 | 980.28 | 317.14 | 65.37 |
| South Dakota | 219 | 17989.42 | 16.67 | 249.80 | 82.14 | 61.54 |

| State/Estimation Area | n | Sum | Minimum | Maximum | Mean | Coefficient of Variation |
|------------------------------|----------|------------|----------------|----------------|-------------|---------------------------------|
| Tennessee | 1,500 | 568417.14 | 9.58 | 2419.46 | 378.94 | 129.72 |
| Texas | 224 | 39136.32 | 41.28 | 441.39 | 174.72 | 52.37 |
| TX-Bexar County | 214 | 71255.42 | 12.66 | 987.02 | 332.97 | 56.63 |
| TX-City of Houston | 211 | 20393.13 | 9.58 | 261.50 | 96.65 | 53.31 |
| TX-El Paso County | 255 | 23985.20 | 22.72 | 257.47 | 94.06 | 49.24 |
| TX-Hidalgo County | 596 | 413647.08 | 18.63 | 2419.46 | 694.04 | 92.49 |
| TX-Rest of State | 210 | 117279.88 | 11.54 | 1640.26 | 558.48 | 67.42 |
| Utah | 247 | 73265.59 | 10.57 | 914.27 | 296.62 | 68.25 |
| Vermont | 262 | 8540.46 | 10.59 | 83.87 | 32.60 | 44.56 |
| Virginia | 373 | 149555.84 | 2.88 | 1687.22 | 400.95 | 132.25 |
| Washington | 309 | 127236.49 | 11.62 | 1374.37 | 411.77 | 74.67 |
| West Virginia | 241 | 29917.36 | 18.71 | 369.77 | 124.14 | 72.90 |
| Wisconsin | 245 | 98535.31 | 16.35 | 1159.22 | 402.18 | 60.42 |
| Wyoming | 219 | 10205.41 | 6.09 | 138.25 | 46.60 | 61.74 |
| U.S. Virgin Islands | 327 | 1956.18 | 1.01 | 19.31 | 5.98 | 70.38 |
| Guam | 282 | 5058.12 | 4.10 | 49.96 | 17.94 | 52.87 |
| Puerto Rico | 224 | 50147.68 | 25.12 | 726.38 | 223.87 | 72.90 |

* Distribution of PROVWT_D_TERR.

† Excludes territories.

Appendix C: Flags for Inconsistent Values in the Breastfeeding Data

Two different types of inconsistency can arise in breastfeeding data. The first is that the duration of any breastfeeding can exceed the age of the child, and the second is that the age of the child when first fed formula can exceed the age of child. BF_ENDR06 stores the duration of any breastfeeding, and BF_ENDFL06 flags the inconsistency; BF_FORMR08 stores the age of the child when first fed formula, and BF_FORMFL06 flags the inconsistency.

1. Both BF_ENDR06 and BF_FORMR08 are formulated using the following conversion factors:

if unit=1(days) then BF_ENDR06 = number x 1
if unit=2(weeks) then BF_ENDR06 = number x 7
if unit=3(months) then BF_ENDR06 = number x 30.4375
if unit=4(years) then BF_ENDR06 = number x 365.25

if unit=1(days) then BF_FORMR08 = number x 1
if unit=2(weeks) then BF_FORMR08 = number x 7
if unit=3(months) then BF_FORMR08 = number x 30.4375
if unit=4(years) then BF_FORMR08 = number x 365.25

2. Flagging BF_ENDR06 when the duration of any breastfeeding exceeds the age in days with a buffer for different units:

if unit=1(days) flag when BF_ENDR06 > age + 1
if unit=2(weeks) flag when BF_ENDR06 > age + 3
if unit=3(months) flag when BF_ENDR06 > age + 15
if unit=4(years) flag when BF_ENDR06 > age + 182

The different buffers allow for the impact of rounding durations upward in the specified units (for example, 50 days might be reported as 2 months).

3. Flagging BF_FORMR08 when the age when first fed formula exceeds the age in days with a buffer for different units:

if unit=1(days) flag when BF_FORMR08 > age + 1
if unit=2(weeks) flag when BF_FORMR08 > age + 3
if unit=3(months) flag when BF_FORMR08 > age + 15
if unit=4(years) flag when BF_FORMR08 > age + 182

The different buffers allow for the impact of rounding durations upward in the specified units (for example, 50 days might be reported as 2 months).

Appendix D: Programs for Estimation: Examples of the Use of SUDAAN, SAS, and R to Estimate Vaccination Coverage Rates and Their Standard Errors, and an Example of the Production of a Cross-Tabulation and Chart

- I. SUDAAN (RTI, 2008) Page 94
- II. SAS (SAS, 2003) Page 107
- III. 'R' (Lumley, 2009) Page 118

A. SUDAAN

```
*****.
title1 'SUD_IAP.SAS';
*****
THIS PROGRAM WILL PRODUCE ESTIMATION AREA ESTIMATES AND STANDARD ERRORS FOR
P_UTD431H314_ROUT_S USING SAS CALLABLE SUDAAN.
SUDAAN NOTES:
1. ALL VARIABLES USED MUST BE NUMERIC.
2. VARIABLES IN THE SUBGROUP STATEMENT MUST HAVE VALUES 1,2,..K
WHERE K IS THE NUMBER OF LEVELS FOR EACH VARIABLE.
3. DATA MUST BE SORTED ACCORDING TO THE SAMPLE DESIGN VARIABLES
(STRATUM AND PRIMARY SAMPLING UNIT), SPECIFIED IN THE
NEST STATEMENT.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;
%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let estiap=estiap15; * --- ESTIMATION AREA VARIABLE TO USE ---*;
%let wt=provwt_d; * --- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. Use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;

Proc format;
/*
THE FOLLOWING FORMAT WILL BE USED FOR P_UTD431H314_ROUT_S.
ORIGINAL VALUES OF P_UTD431H314_ROUT_S ARE 1,0.
MUST BE CONVERTED TO 1,2 IN SUDAAN.
*/
value putd431h314f
1='4:3:1:H:3:1:4 Up-to-Date'
2='Not 4:3:1:H:3:1:4 Up-to-Date';

value estiapf
. = "Missing"
0 = "US Total"
1 = "CT"
2 = "MA"
4 = "ME"
5 = "NH"
6 = "RI"
7 = "VT"
8 = "NJ"
10 = "NY-Rest of State"
11 = "NY-City of New York"
```

```
12 = "DC"  
13 = "DE"  
14 = "MD"  
16 = "PA-Rest of State"  
17 = "PA-Philadelphia County"  
18 = "VA"  
19 = "WV"  
20 = "AL"  
22 = "FL"  
25 = "GA"  
27 = "KY"  
28 = "MS"  
29 = "NC"  
30 = "SC"  
31 = "TN"  
34 = "IL-Rest of State"  
35 = "IL-City of Chicago"  
36 = "IN"  
38 = "MI"  
40 = "MN"  
41 = "OH"  
44 = "WI"  
46 = "AR"  
47 = "LA"  
49 = "NM"  
50 = "OK"  
51 = "TX-Rest of State"  
53 = "TX-El Paso County"  
54 = "TX-City of Houston"  
55 = "TX-Bexar County"  
56 = "IA "  
57 = "KS"  
58 = "MO"  
59 = "NE"  
60 = "CO"  
61 = "MT"  
62 = "ND"  
63 = "SD"  
64 = "UT"  
65 = "WY"  
66 = "AZ"  
68 = "CA"  
72 = "HI"  
73 = "NV"  
74 = "AK"  
75 = "ID"  
76 = "OR"  
77 = "WA "  
106 = "Puerto Rico"  
107 = "TX-Hidalgo County"  
;  
run;
```

```

data sud_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S &estiap &wt &strat);
if P_UTD431H314_ROUT_S=0 then P_UTD431H314_ROUT_S=2; *--- CONVERT P_UTD431H314_ROUT_S=0 TO
P_UTD431H314_ROUT_S=2 ---*;
nseqnumh=1*seqnumhh; *---CONVERT HOUSEHOLD ID SEQNUMHH FROM CHARACTER TO NUMERIC ---*;
run;
*=== SORT BY NEST VARIABLES: STRATUM (STRATUM) NSEQNUMH (PRIMARY SAMPLING UNIT) ===*;
proc sort;
by &strat nseqnumh;
run;
proc crosstab data=sud_file filetype=sas design=wr;
weight &wt;
nest &strat nseqnumh;
subgroup &estiap P_UTD431H314_ROUT_S ;
levels 107 2 ;
tables &estiap * P_UTD431H314_ROUT_S ;
print nsum wsum rowper serow/style=nchs ;
rtitle "4:3:1:H:3:1:4 ESTIMATES BY Estimation Area";
rformat &estiap estiapf.;
rformat P_UTD431H314_ROUT_S putd431h314f.;
output rowper serow/filename=sud_est filetype=sas replace;
run;
proc print data=sud_est(where=(P_UTD431H314_ROUT_S=1 and rowper ne .)) noobs label;
format &estiap estiapf.;
var &estiap rowper serow ;
label
rowper='Percent 4:3:1:H:3:1:4 Up-to-Date'
serow='Standard Error'
;
title "4:3:1:H:3:1:4 ESTIMATES BY Estimation Area";
run;

*****.
title1 'SUDSTATE.SAS';
*****
THIS PROGRAM WILL PRODUCE STATE ESTIMATES AND STANDARD ERRORS
FOR P_UTD431H314_ROUT_S USING SAS CALLABLE SUDAAN.
NOTE : THE STATE VARIABLE IS BASED ON FIPSTATE CODES ,THERE ARE
NO STATES WITH FIPS CODES 3,7,14,43,52,57-71,73-78.
SUDAAN NOTES:
1. ALL VARIABLES USED MUST BE NUMERIC.
2. VARIABLES IN THE SUBGROUP STATEMENT MUST HAVE VALUES 1,2,..K
WHERE K IS THE NUMBER OF LEVELS FOR EACH VARIABLE.
3. DATA MUST BE SORTED ACCORDING TO THE SAMPLE DESIGN VARIABLES
(STRATUM AND PRIMARY SAMPLING UNIT), SPECIFIED IN THE
NEST STATEMENT.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;

```

```

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;
%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=provwt_d; *--- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. Use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;

```

PROC FORMAT;

```

/*
THE FOLLOWING FORMAT WILL BE USED FOR P_UTD431H314_ROUT_S.
ORIGINAL VALUES OF P_UTD431H314_ROUT_S ARE 1,0.
MUST BE CONVERTED TO 1,2 IN SUDAAN.
*/

```

```

value putd431h314f
1='4:3:1:H:3:1:4 Up-to-Date'
2='Not 4:3:1:H:3:1:4 Up-to-Date'
;

```

```

value statef
0='U.S. Total'
1='Alabama '
2='Alaska '
4='Arizona '
5='Arkansas '
6='California '
8='Colorado '
9='Connecticut '
10='Delaware '
11='District of Columbia'
12='Florida '
13='Georgia '
15='Hawaii '
16='Idaho '
17='Illinois '
18='Indiana '
19='Iowa '
20='Kansas '
21='Kentucky '
22='Louisiana '
23='Maine '
24='Maryland '
25='Massachusetts '
26='Michigan '
27='Minnesota '
28='Mississippi '
29='Missouri '
30='Montana '
31='Nebraska '
32='Nevada '
33='New Hampshire '
34='New Jersey '
35='New Mexico '

```

```

36 ='New York '
37 ='North Carolina '
38 ='North Dakota '
39 ='Ohio '
40 ='Oklahoma '
41 ='Oregon '
42 ='Pennsylvania '
44 ='Rhode Island '
45 ='South Carolina '
46 ='South Dakota '
47 ='Tennessee '
48 ='Texas '
49 ='Utah '
50 ='Vermont '
51 ='Virginia '
53 ='Washington '
54 ='West Virginia '
55 ='Wisconsin '
56 ='Wyoming '
72 ='Puerto Rico '
;
run;
data sud_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S state &wt &strat);
if P_UTD431H314_ROUT_S=0 then P_UTD431H314_ROUT_S=2; *** CONVERT P_UTD431H314_ROUT_S=0 TO
P_UTD431H314_ROUT_S=2 ***;
nseqnumh=1*seqnumhh; *** CONVERT HOUSEHOLD ID SEQNUMH FROM CHARACTER TO NUMERIC ***;
run;
*=== SORT BY NEST VARIABLES: STRATUM (STRATUM) NSEQNUMH (PRIMARY SAMPLING UNIT) ===*;
proc sort;
by &strat nseqnumh;
run;
proc crosstab data=sud_file filetype=sas design=wr;
weight &wt;
nest &strat nseqnumh;
subgroup state P_UTD431H314_ROUT_S ;
levels 56 2 ;
tables state * P_UTD431H314_ROUT_S ;
print nsum wsum rowper serow/style=nchs ;
rtile "4:3:1:H:3:1:4 ESTIMATES BY STATE";
rformat state statef.;
rformat P_UTD431H314_ROUT_S putd431h314f.;
output rowper serow / filename=sud_est2 filetype=sas replace;
run;
*** EXCLUDE 3,7,14,43,52,57-71,73-78 THERE ARE NO STATES WITH THESE FIPS CODES *** ;
option spool;
proc print data=sud_est2(where=(P_UTD431H314_ROUT_S=1
& state notin (3,7,14,43,52) & not(57<=state<=71) & not(73<=state<=78))) label noobs;
format state statef.;
var state rowper serow ;
label
rowper='Percent 4:3:1:H:3:1:4 Up-to-Date'

```

```

serow='Standard Error'
;
title "4:3:1:H:3:1:4 ESTIMATES BY STATE";
run;
option nospool;

*****;
title1 'PROG_3.SAS';
*****
THIS PROGRAM WILL PRODUCE A TABLE OF HAD_CPOX BY STATE FOR ALL RDD
COMPLETES USING RDDWT_D. THE PROGRAM USES SAS CALLABLE SUDAAN.
SUDAAN NOTES:
1. ALL VARIABLES USED MUST BE NUMERIC.
2. VARIABLES IN THE SUBGROUP STATEMENT MUST HAVE VALUES 1,2,..K
WHERE K IS THE NUMBER OF LEVELS FOR EACH VARIABLE.
3. DATA MUST BE SORTED ACCORDING TO THE SAMPLE DESIGN VARIABLES
(STRATUM AND PRIMARY SAMPLING UNIT), SPECIFIED IN THE
NEST STATEMENT.
*****;
options ps=78 ls=90 obs= max;
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;
%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=rddwt_d; * --- WEIGHT TO USE (RDDWT_D is the dual-frame weight excluding territories. Use
RDDWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;

PROC FORMAT;
/*
THE FOLLOWING FORMAT WILL BE USED FOR HAD_CPOX.
*/
value hadcpoxf
1='Yes'
2='No'
;
value statef
0='U.S. Total '
1='Alabama '
2='Alaska '
4='Arizona '
5='Arkansas '
6='California '
8='Colorado '
9='Connecticut '
10='Delaware '

```

```

11 ='District of Columbia'
12 ='Florida '
13 ='Georgia '
15 ='Hawaii '
16 ='Idaho '
17 ='Illinois '
18 ='Indiana '
19 ='Iowa '
20 ='Kansas '
21 ='Kentucky '
22 ='Louisiana '
23 ='Maine '
24 ='Maryland '
25 ='Massachusetts '
26 ='Michigan '
27 ='Minnesota '
28 ='Mississippi '
29 ='Missouri '
30 ='Montana '
31 ='Nebraska '
32 ='Nevada '
33 ='New Hampshire '
34 ='New Jersey '
35 ='New Mexico '
36 ='New York '
37 ='North Carolina '
38 ='North Dakota '
39 ='Ohio '
40 ='Oklahoma '
41 ='Oregon '
42 ='Pennsylvania '
44 ='Rhode Island '
45 ='South Carolina '
46 ='South Dakota '
47 ='Tennessee '
48 ='Texas '
49 ='Utah '
50 ='Vermont '
51 ='Virginia '
53 ='Washington '
54 ='West Virginia '
55 ='Wisconsin '
56 ='Wyoming '
72 ='Puerto Rico '
;
run;
data sud_file;
set &in_file(keep= seqnumhh seqnumc state had_cpox &wt &strat);
nseqnumh=1*seqnumhh; *** CONVERT HOUSEHOLD ID SEQNUMH FROM CHARACTER TO NUMERIC ***;
run;
*=== SORT BY NEST VARIABLES: STRATUM (STRATUM) NSEQNUMH (PRIMARY SAMPLING UNIT) ===*;
proc sort;

```

```

by &strat nseqnumh;
run;
proc crosstab data=sud_file filetype=sas design=wr;
weight &wt;
nest &strat nseqnumh;
subgroup state had_cpox ;
levels 56 2 ;
tables state * had_cpox ;
print nsum wsum rowper serow/style=nchs ;
rtitle "HAD_CPOX ESTIMATES BY STATE";
rtitle "WEIGHT = &WT";
rformat state statef.;
rformat had_cpox hadcpoxf.;
output rowper serow / filename=sud_est3 filetype=sas replace;
run;
*** EXCLUDE 3,7,14,43,52,57-71,73-78 THERE ARE NO STATES WITH THESE FIPS CODES *** ;
option spool;
proc print data=sud_est3(where=(had_cpox=1
& state notin (3,7,14,43,52) & not(57<=state<=71) & not(73<=state<=78))) label noobs;
format state statef.;
var state rowper serow ;
label
rowper='Percent HAD_CPOX = Yes'
serow='Standard Error'
;
title "CHILD HAD CHICKEN POX BY STATE";
run;
option nospool;

```

```

*****.
title1 'PROG_4.SAS';
*****
TABLE OF P_UTD431H314_ROUT_S BY INCPOV1 BY RACE_K. SAVE % UTD
ESTIMATES (NOT S.E.'S) FOR USE IN THE PROGRAM CHART_4.
THIS PROGRAM WILL PRODUCE ESTIMATES USING SAS CALLABLE SUDAAN.
SUDAAN NOTES:
1. ALL VARIABLES USED MUST BE NUMERIC.
2. VARIABLES IN THE SUBGROUP STATEMENT MUST HAVE VALUES 1,2,..K
WHERE K IS THE NUMBER OF LEVELS FOR EACH VARIABLE.
3. DATA MUST BE SORTED ACCORDING TO THE SAMPLE DESIGN VARIABLES
(STRATUM AND PRIMARY SAMPLING UNIT), SPECIFIED IN THE
NEST STATEMENT.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;

```

```
libname out 'c:\nispuf15'; *--- SPECIFY THE PATH FOR WHERE YOU WANT THE CHART OUTPUT TO GO ---*;

%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=provwt_d; *--- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. Use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;
%let qtr_lab=Q1/2015 - Q4/2015; *NIS 4 QUARTER PERIOD*;
```

PROC FORMAT;

```
/*
THE FOLLOWING FORMAT WILL BE USED FOR P_UTD431H314_ROUT_S.
ORIGINAL VALUES OF P_UTD431H314_ROUT_S ARE 1,0.
MUST BE CONVERTED TO 1,2 IN SUDAAN.
*/
```

```
value putd431h314f
```

```
1='4:3:1:H:3:1:4 Up-to-date'
```

```
2='Not 4:3:1:H:3:1:4 Up-to-date'
```

```
;
```

```
VALUE RACE KF
```

```
1 = "WHITE ONLY"
```

```
2 = "BLACK ONLY"
```

```
3 = "OTHER AND MULTIPLE RACE"
```

```
;
```

```
VALUE INCPVR2F
```

```
1 = "ABOVE, > $75,000"
```

```
2 = "ABOVE, <= $75,000"
```

```
3 = "BELOW"
```

```
4 = "UNKNOWN"
```

```
;
```

```
value statef
```

```
0='U.S. Total '
```

```
1='Alabama '
```

```
2='Alaska '
```

```
4='Arizona '
```

```
5='Arkansas '
```

```
6='California '
```

```
8='Colorado '
```

```
9='Connecticut '
```

```
10='Delaware '
```

```
11='District of Columbia'
```

```
12='Florida '
```

```
13='Georgia '
```

```
15='Hawaii '
```

```
16='Idaho '
```

```
17='Illinois '
```

```
18='Indiana '
```

```
19='Iowa '
```

```
20='Kansas '
```

```
21='Kentucky '
```

```
22='Louisiana '
```

```
23='Maine '
```

```

24 ='Maryland '
25 ='Massachusetts '
26 ='Michigan '
27 ='Minnesota '
28 ='Mississippi '
29 ='Missouri '
30 ='Montana '
31 ='Nebraska '
32 ='Nevada '
33 ='New Hampshire '
34 ='New Jersey '
35 ='New Mexico '
36 ='New York '
37 ='North Carolina '
38 ='North Dakota '
39 ='Ohio '
40 ='Oklahoma '
41 ='Oregon '
42 ='Pennsylvania '
44 ='Rhode Island '
45 ='South Carolina '
46 ='South Dakota '
47 ='Tennessee '
48 ='Texas '
49 ='Utah '
50 ='Vermont '
51 ='Virginia '
53 ='Washington '
54 ='West Virginia '
55 ='Wisconsin '
56 ='Wyoming '
72 ='Puerto Rico '
;
run;
data sud_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S race_k incpov1 &wt &strat);
nseqnumh=1*seqnumhh; *** CONVERT HOUSEHOLD ID SEQNUMH FROM CHARACTER TO NUMERIC ***;
if P_UTD431H314_ROUT_S=0 then P_UTD431H314_ROUT_S=2; *** CONVERT P_UTD431H314_ROUT_S=0 TO
P_UTD431H314_ROUT_S=2 ***;
run;
*=== SORT BY NEST VARIABLES: STRATUM (STRATUM) NSEQNUMH (PRIMARY SAMPLING UNIT) ===*;
proc sort;
by &strat nseqnumh;
run;
proc freq;
tables P_UTD431H314_ROUT_S incpov1 race_k;
run;
title3 "Table 4A. &qtr_lab: Unweighted Frequencies";
proc crosstab data=sud_file filetype=sas design=wr;
weight &wt;
nest &strat nseqnumh;
subgroup incpov1 race_k P_UTD431H314_ROUT_S ;

```

```

levels 4 3 2 ;
tables (incpov1 * race_k * P_UTD431H314_ROUT_S) ;
print nsum wsum rowper="4:3:1:H:3:1:4 Up-to-Date (ROWPER)"
serow="Standard Error (SEROW)" /style=nchs ;
rtitle "Table 4B. &qtr_lab, Percent 4:3:1:H:3:1:4 Up-to-Date and Estimated Standard Errors";
rtitle "WEIGHT = &WT";
rformat P_UTD431H314_ROUT_S putd431h314f.;
rformat incpov1 incpvr2f.;
rformat race_k race_kf.;
output rowper serow / filename=sud_est4 filetype=sas replace;
run;
data out.sud_est4;
set sud_est4(where=(P_UTD431H314_ROUT_S=1 & incpov1 > 0 & race_k > 0));
keep incpov1 race_k rowper serow;
label rowper='4:3:1:H:3:1:4 Up-to-Date';
format rowper 5.2;
format serow 5.2;
run;
proc print data=out.sud_est4 label;
format race_k race_kf.;
format incpov1 incpvr2f.;
title "&qtr_lab: 4:3:1:H:3:1:4 ESTIMATES AND STANDARD ERRORS BY INCPOV1 BY RACE_K";
run;

*****.
title1 'SAS_GRAPH_4.SAS';
*****
THIS PROGRAM BUILDS OFF OF THE PROGRAM SAS_PROG_4. IT PRODUCES A CHART OF
P_UTD431H314_ROUT_S BY INCPOV1 BY RACE_K. IT CREATES A BAR CHART IN SAS GRAPH FOR
THE 4X3 = 12 CELLS. THE OUTPUT OF THE FOLLOWING EXAMPLE IS ATTACHED AT THE
END.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;

%let out='c:\nispuf15'; *--- SPECIFY THE PATH FOR WHERE YOU WANT THE CHART
OUTPUT TO GO ---*;

%let in_file=dd.sud_est4; *--- NAME OF SAS DATASET OUTPUT FROM PROG_4 ---*;
%let qtr_lab=Q1/2015 - Q4/2015; *NIS 4 QUARTER PERIOD*;
PROC FORMAT;
VALUE INCPVR2F
1 = "ABOVE, > $75,000"
2 = "ABOVE, <= $75,000"
3 = "BELOW"
4 = "UNKNOWN"
;
VALUE RACE KF
1 = "WHITE ONLY"
2 = "BLACK ONLY"
3 = "OTHER/MULT RACE"

```


```

;
run;
data sud_est4;
set &in file;
format rowper 3.
race_k race_kf.
incpov1 incpvr2f.
;
label
race_k = 'Race of Child'
incpov1 = 'Poverty Status'
;
filename odsout &out;
ods listing close;
/* SET THE GRAPHICS ENVIRONMENT */
goptions reset=global gunit=pct border
fext=swissb htitle=4 htext=1.5
device=gif
;
ods html body='graph 4 sud.html' path=odsout;
TITLE1 HEIGHT=3 "Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4 ";
TITLE2 HEIGHT=3 "by Race and Poverty Status, National Immunization Survey, 2015";
footnote j=r 'graph_4sud';
pattern1 value = solid color = blue;
pattern2 value = x3 color = green;
pattern3 value = l3 color = red;
pattern4 value = empty color = lib;
axis width = 3;
run;
proc gchart data=sud_est4;
vbar race_k
/frame
discrete
sumvar=rowper
group=incpov1
gspace = 5
gaxis = axis
raxis = axis
name = 'graph_4_sud'
patternid = midpoint
;
run;
quit;
ods html close;
ods listing;
ods html close;
ods listing;

```

Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4 by Race and Poverty Status, National Immunization Survey, 2015

4:3:1:H:3:1:4 Up-to-Date



graph_4su d

B. SAS

```
*****;
title1 'SAS_IAP.SAS';
*****
THIS PROGRAM WILL PRODUCE ESTIMATION AREA ESTIMATES AND STANDARD ERRORS
FOR P_UTD431H314_ROUT_S USING SAS.
*****;
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS STORED ---*;

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;
%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let estiap=estiap15; * --- ESTIMATION AREA VARIABLE TO USE ---*;
%let wt=provwt_d; * --- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. Use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;
proc format;
value putd431h314f
0='Not 4:3:1:H:3:1:4 Up-To-Date'
1='4:3:1:H:3:1:4 Up-To-Date';

value estiapf
. = "Missing"
0 = "US Total"
1 = "CT"
2 = "MA"
4 = "ME"
5 = "NH"
6 = "RI"
7 = "VT"
8 = "NJ"
10 = "NY-Rest of State"
11 = "NY-City of New York"
12 = "DC"
13 = "DE"
14 = "MD"
16 = "PA-Rest of State"
17 = "PA-Philadelphia County"
18 = "VA"
19 = "WV"
20 = "AL"
22 = "FL"
25 = "GA"
27 = "KY"
28 = "MS"
29 = "NC"
30 = "SC"
```

```

31 = "TN"
34 = "IL-Rest of State"
35 = "IL-City of Chicago"
36 = "IN "
38 = "MI"
40 = "MN"
41 = "OH"
44 = "WI"
46 = "AR"
47 = "LA"
49 = "NM"
50 = "OK"
51 = "TX-Rest of State"
53 = "TX-El Paso County"
54 = "TX-City of Houston"
55 = "TX-Bexar County"
56 = "IA"
57 = "KS"
58 = "MO"
59 = "NE"
60 = "CO"
61 = "MT"
62 = "ND"
63 = "SD"
64 = "UT"
65 = "WY"
66 = "AZ"
68 = "CA"
72 = "HI"
73 = "NV"
74 = "AK"
75 = "ID"
76 = "OR"
77 = "WA"
106 = "Puerto Rico"
107 = "TX-Hidalgo County"
;
run;
data sas_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S &estiap &wt &strat);
run;
proc sort data = sas_file;
by &estiap;
run;
title '4:3:1:H:3:1:4 ESTIMATES BY Estimation Area';
ods output Statistics=sas_est;
proc surveymeans data = sas_file nobc sum mean stderr;
stratum &strat;
cluster seqnumhh;
weight &wt;
class P_UTD431H314_ROUT_S;
var P_UTD431H314_ROUT_S;

```

```

by &estiap;
format P_UTD431H314_ROUT_S putd431h314f.;
format &estiap estiapf.;
run;
data sas_est;
set sas_est;
mean = mean*100; *CONVERT TO PERCENT ESTIMATES;
stderr = stderr*100;
run;
proc print data=sas_est(where=(varlevel='4:3:1:H:3:1:4 Up-To-Date')) noobs
label;
format &estiap estiapf.;
format mean stderr 5.2;
var &estiap mean stderr;
label
mean='Percent 4:3:1:H:3:1:4 Up-to-Date'
stderr='Standard Error';
title "4:3:1:H:3:1:4 Estimates by Estimation Area";
run;

*****.
title1 'SASSTATE.SAS';
*****
THIS PROGRAM WILL PRODUCE STATE ESTIMATES AND STANDARD ERRORS
FOR P_UTD431H314_ROUT_S USING SAS.
NOTE : THE STATE VARIABLE IS BASED ON FIPSTATE CODES ,THERE ARE
NO STATES WITH FIPS CODES 3,7,14,43,52,57-71,73-77.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;
*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;

%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=provwt_d; * --- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. Use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;
proc format;
value putd431h314f
0='Not 4:3:1:H:3:1:4 Up-To-Date'
1='4:3:1:H:3:1:4 Up-To-Date';
value statef
. = "Missing"
0 = 'U.S. Total '
1 = 'Alabama '
2 = 'Alaska '
4 = 'Arizona '
5 = 'Arkansas '
6 = 'California '

```

```

8='Colorado '
9='Connecticut '
10='Delaware '
11='District of Columbia'
12='Florida '
13='Georgia '
15='Hawaii '
16='Idaho '
17='Illinois '
18='Indiana '
19='Iowa '
20='Kansas '
21='Kentucky '
22='Louisiana '
23='Maine '
24='Maryland '
25='Massachusetts '
26='Michigan '
27='Minnesota '
28='Mississippi '
29='Missouri '
30='Montana '
31='Nebraska '
32='Nevada '
33='New Hampshire '
34='New Jersey '
35='New Mexico '
36='New York '
37='North Carolina '
38='North Dakota '
39='Ohio '
40='Oklahoma '
41='Oregon '
42='Pennsylvania '
44='Rhode Island '
45='South Carolina '
46='South Dakota '
47='Tennessee '
48='Texas '
49='Utah '
50='Vermont '
51='Virginia '
53='Washington '
54='West Virginia '
55='Wisconsin '
56='Wyoming '
72='Puerto Rico '
;
run;
data sas_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S state &wt &strat);
run;

```

```

proc sort data = sas_file;
by state;
title1 '4:3:1:H:3:1:4 ESTIMATES BY STATE';
ods output Statistics=sas_est2;
run;
proc surveymeans data = sas_file nobks sum mean stderr;
stratum &strat;
cluster seqnumhh;
weight &wt;
class P_UTD431H314_ROUT_S;
var P_UTD431H314_ROUT_S;
by state;
format P_UTD431H314_ROUT_S putd431h314f.;
format state statef.;
run;
data sas_est2;
set sas_est2;
mean = mean*100; *CONVERT TO PERCENT ESTIMATES;
stderr = stderr*100;
run;
proc print data=sas_est2(where=(varlevel='4:3:1:H:3:1:4 Up-To-Date')) nobks
label;
format state statef.;
format mean stderr 5.2;
var state mean stderr;
label
mean='Percent 4:3:1:H:3:1:4 Up-to-Date'
stderr='Standard Error';
title "4:3:1:H:3:1:4 ESTIMATES BY STATE";
run;

*****.
title1 'SAS_PROG_3.SAS';
*****
THIS PROGRAM WILL PRODUCE A TABLE OF HAD_CPOX BY STATE FOR ALL RDD
COMPLETES USING RDDWT. THE PROGRAM USES SAS.
*****;
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;
*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;

%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=rddwt_d; *--- WEIGHT TO USE (RDDWT_D is the dual-frame weight excluding territories. Use
RDDWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;
PROC FORMAT;
value hadcpoxf
1='Yes'

```

```
2='No'  
;  
value statef  
0='U.S. Total '  
1='Alabama '  
2='Alaska '  
4='Arizona '  
5='Arkansas '  
6='California '  
8='Colorado '  
9='Connecticut '  
10='Delaware '  
11='District of Columbia'  
12='Florida '  
13='Georgia '  
15='Hawaii '  
16='Idaho '  
17='Illinois '  
18='Indiana '  
19='Iowa '  
20='Kansas '  
21='Kentucky '  
22='Louisiana '  
23='Maine '  
24='Maryland '  
25='Massachusetts '  
26='Michigan '  
27='Minnesota '  
28='Mississippi '  
29='Missouri '  
30='Montana '  
31='Nebraska '  
32='Nevada '  
33='New Hampshire '  
34='New Jersey '  
35='New Mexico '  
36='New York '  
37='North Carolina '  
38='North Dakota '  
39='Ohio '  
40='Oklahoma '  
41='Oregon '  
42='Pennsylvania '  
44='Rhode Island '  
45='South Carolina '  
46='South Dakota '  
47='Tennessee '  
48='Texas '  
49='Utah '  
50='Vermont '  
51='Virginia '  
53='Washington '
```

```

54='West Virginia '
55='Wisconsin '
56='Wyoming '
72='Puerto Rico '
;
run;
data sas_file;
set &in_file(keep= seqnumhh seqnumc state had_cpox &wt &strat);
run;
proc sort data = sas_file;
by state;
title1 'HAD_CPOX ESTIMATES BY STATE';
ods output Statistics=sas_est3;
run;
proc surveymeans data = sas_file nobsum mean stderr;
stratum &strat;
cluster seqnumhh;
weight &wt;
class had_cpox;
var had_cpox;
by state;
format had_cpox hadcpoxf.;
format state statef.;
run;
data sas_est3;
set sas_est3;
mean = mean*100; *CONVERT TO PERCENT ESTIMATES;
stderr = stderr*100;
run;
proc print data=sas_est3(where=(varlevel='Yes')) nobsum label;
format state statef.;
format mean stderr 5.2;
var state mean stderr;
label
mean='Percent HAD_CPOX = Yes'
stderr='Standard Error';
title "CHILD HAD CHICKEN POX BY ESTIMATION AREA";
run;

*****;
title1 'SAS_PROG_4.SAS';
*****;
TABLE OF P_UTD431H314_ROUT_S BY INCPOV1 BY RACE_K. SAVE % UTD
ESTIMATES (NOT S.E.'S) FOR USE IN THE PROGRAM SAS_GRAPH_4.
THIS PROGRAM WILL PRODUCE ESTIMATES USING SAS.
*****;
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;
libname library 'c:\nispuf15'; *--- IF DATASET WAS CREATED WITH FORMATS
STORED ---*;

```

```

*--- PERMANENTLY SPECIFY PATH TO LIBRARY ---*;
*--- OTHERWISE COMMENT THIS STATEMENT OUT ---*;

libname out 'c:\nispuf15'; *--- SPECIFY THE PATH FOR
WHERE YOU WANT THE CHART OUTPUT TO GO ---*;

%let in_file=dd.nispuf15; *--- NAME OF SAS DATASET ---*;
%let wt=provwt_d; *--- WEIGHT TO USE (PROVWT_D is the dual-frame weight excluding territories. use
PROVWT_D_TERR to include territories) ---*;
%let strat=stratum; * --- STRATUM VARIABLE TO USE FOR VARIANCE ESTIMATION;
%let qtr lab=Q1/2015 - Q4/2015; *NIS 4 QUARTER PERIOD*;
PROC FORMAT;
value putd431h314f
0='Not 4:3:1:H:3:1:4 Up-To-Date'
1='4:3:1:H:3:1:4 Up-To-Date'
;
VALUE RACE_KF
1 = "WHITE ONLY"
2 = "BLACK ONLY"
3 = "OTHER AND MULTIPLE RACE"
;
VALUE INCPVR2F
1 = "ABOVE, > $75,000"
2 = "ABOVE, <= $75,000"
3 = "BELOW"
4 = "UNKNOWN"
;
run;
data sas_file;
set &in_file(keep= seqnumhh seqnumc P_UTD431H314_ROUT_S race_k incpov1 &wt &strat);
run;
proc sort data = sas_file;
by incpov1 race_k;
run;
proc freq;
tables P_UTD431H314_ROUT_S incpov1 race_k;
title1 "Table 4A. &qtr_lab: Unweighted Frequencies";
run;
data sas_file;
set sas_file;
if P_UTD431H314_ROUT_S < 0 | incpov1 < 0 | race_k < 0 | &wt. < 0 then delete;
run;
proc surveymeans data = sas_file nobsum mean stderr;
ods output Domain=sas_est4;
stratum &strat;
cluster seqnumhh;
weight &wt;
class P_UTD431H314_ROUT_S;
var P_UTD431H314_ROUT_S;
domain incpov1*race_k;
format P_UTD431H314_ROUT_S putd431h314f.;
format incpov1 incpvr2f.;

```

```

format race_k race_kf.;
run;
data sas_est4;
set sas est4;
mean = mean*100; *CONVERT TO PERCENT ESTIMATES;
stderr = stderr*100;
run;
proc print data=sas_est4(where=(varlevel='4:3:1:H:3:1:4 Up-To-Date')) noobs
label;
format incpov1 incpvr2f.;
format race k race kf.;
format mean stderr 5.2;
var incpov1 race_k mean stderr;
label
mean='4:3:1:H:3:1:4 Up-To-Date'
stderr='Standard Error';
title1 "Table 4B. &qtr_lab, Percent 4:3:1:H:3:1:4 Up-to-Date and Estimated
Standard Errors";
run;
data out.sas_est4;
set sas_est4(where=(varlevel='4:3:1:H:3:1:4 Up-To-Date'));
keep incpov1 race_k mean;
label mean='4:3:1:H:3:1:4 Up-to-Date';
format mean 5.2;
run;

*****.
title1 'SAS_GRAPH_4.SAS';
*****
THIS PROGRAM BUILDS OFF OF THE PROGRAM SAS_PROG_4. IT PRODUCES A CHART OF
P_UTD431H314_ROUT_$ BY INCPOV1 BY RACE_K. IT CREATES A BAR CHART IN SAS GRAPH FOR
THE 4X3 = 12 CELLS. THE OUTPUT OF THE FOLLOWING EXAMPLE IS ATTACHED AT THE
END.
*****.
options ps=78 ls=90 obs= max;

libname dd 'c:\nispuf15'; *--- SPECIFY PATH TO SAS DATASET ---*;

%let out='c:\nispuf15'; *--- SPECIFY THE PATH FOR WHERE YOU WANT THE CHART
OUTPUT TO GO ---*;

%let in_file=dd.sas_est4; *--- NAME OF SAS DATASET OUTPUT FROM PROG_4 ---
*;
%let qtr_lab=Q1/2015 - Q4/2015; *NIS 4 QUARTER PERIOD*;
PROC FORMAT;
VALUE INCPVR2F
1 = "ABOVE, > $75,000"
2 = "ABOVE, <= $75,000"
3 = "BELOW"
4 = "UNKNOWN"
;
VALUE RACE_KF

```


```

1 = "WHITE ONLY"
2 = "BLACK ONLY"
3 = "OTHER/MULT RACE"
;
run;
data sas_est4;
set &in file;
format mean 3.
race_k race_kf.
incpov1 incpvr2f.
;
label
race_k = 'Race of Child'
incpov1 = 'Poverty Status'
;
filename odsout &out;
ods listing close;
/* SET THE GRAPHICS ENVIRONMENT */
goptions reset=global gunit=pct border
ftext=swissb htitle=4 htext=1.5
device=gif
;
ods html body='graph_4.html' path=odsout;
TITLE1 HEIGHT=3 "Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4 ";
TITLE2 HEIGHT=3 "by Race and Poverty Status, National Immunization Survey, 2015";
footnote j=r 'graph_4';
pattern1 value = solid color = blue;
pattern2 value = x3 color = green;
pattern3 value = l3 color = red;
pattern4 value = empty color = lib;
axis width = 3;
run;
proc gchart data=sas_est4;
vbar race_k
/frame
discrete
sumvar=mean
group=incpov1
gspace = 5
gaxis = axis
raxis = axis
name = 'graph_4'
patternid = midpoint
;
run;
quit;
ods html close;
ods listing;

```

Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4 by Race and Poverty Status, National Immunization Survey, 2015

4:3:1:H:3:1:4 Up-to-Date



graph_4

C. 'R'

```
#####
title <- "R_IAP.R"
#####
#THIS PROGRAM WILL PRODUCE ESTIMATION AREA ESTIMATES AND STANDARD ERRORS
#FOR P_UTD431H314_ROUT_S USING R.
#
#R NOTES:
#1. R IS CASE SENSITIVE.
#2. A FILE PATH IS SEPERATED BY SLASH(/)
#####
library(survey) #TO USE svydesign(), svymean(), and svyby()
library(Hmisc) #TO USE prn()

dd <- "c:/nispuf15" #"path-to-dataset"

#--- NAME OF R DATASET ---#
in.file <- paste(dd,"NISPUF15.RData",sep="")
#---READ R DATASET---#
load(in.file)
#---FORMAT---#
UTD431H314levels=c(0,1)
UTD431H314labels=c("NOT 4:3:1:H:3:1:4 UTD", "4:3:1:H:3:1:4 UTD")
ESTIAPlevels=c(0, 1, 2, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 22, 25, 27, 28, 29, 30, 31, 34, 35, 36, 38, 40, 41, 44, 46, 47, 49, 50, 51, 52,
53, 54, 55, 56, 57, 58, 59, 60, 61, 63, 64, 65, 66, 68, 72, 73, 74, 75, 76, 77, 106, 107)
ESTIAPlabels=c("US Total", "CT", "MA", "ME", "NH", "RI", "VT", "NJ", "NY-Rest of State", "NY-City of New York", "DC", "DE", "MD", "PA-
Rest of State", "PA-Philadelphia County", "VA", "WV", "AL", "FL", "GA", "KY", "MS", "NC", "SC", "TN", "IL-Rest of State", "IL-City of
Chicago", "IN", "MI", "MN", "OH", "WI", "AR", "LA", "NM", "OK", "TX-Rest of State", "TX-El Paso County", "TX-City of Houston", "TX-Bexar
County", "IA", "KS", "MO", "NE", "CO", "MT", "ND", "SD", "UT", "WY", "AZ", "CA", "HI", "NV", "AK", "ID", "OR", "WA", "Puerto Rico",
"TX-Hidalgo County")

#---PROVWT_D WILL BE USED AS A WEIGHT (PROVWT_D IS THE DUAL-FRAME WEIGHT EXCLUDING TERRITORIES. USE
PROVWT_D_TERR TO INCLUDE TERRITORIES)---#
#---STRATUM WILL BE USED AS A STRATUM VARIABLE FOR VARIANCE ESTIMATION ---#
R_FILE <- subset(NISPUF15, select=c(SEQNUMHH, SEQNUMC, P_UTD431H314_ROUT_S, ESTIAP15,
PROVWT_D, STRATUM))
names(R_FILE) <- c("SEQNUMHH", "SEQNUMC", "P_UTD431H314_ROUT_S", "ESTIAP", "WT", "STRATUM")
R_FILE <- na.omit(R_FILE)
#---ASSIGN LABELS---#
R_FILE$P_UTD431H314_ROUT_S <- factor(R_FILE$P_UTD431H314_ROUT_S, levels=UTD431H314levels,
labels=UTD431H314labels)
R_FILE$ESTIAP <- factor(R_FILE$ESTIAP, levels=ESTIAPlevels,
labels=ESTIAPlabels)

#---SPECIFY A SAMPLING DESIGN AND FORCE WT AS NUMERIC---#
svydsg <- svydesign(id=~SEQNUMHH, strata=~STRATUM, weights=~(as.numeric(WT)),
data=R_FILE)

#---U.S. TOTAL ESTIMATES AND STANDARD ERRORS---#
r_nation <- svymean(~P_UTD431H314_ROUT_S, svydsg)
PERCENT_UTD <- round(r_nation*100,2) #CONVERT INTO PERCENT ESTIMATES(MEAN)
SE_UTD <- round(SE(r_nation)*100,2) #CONVERT INTO PERCENT ESTIMATES(SE)
r_nation_est <- cbind(PERCENT_UTD, SE_UTD)
title <- "PERCENT 4:3:1:H:3:1:4 ESTIMATES AT A NATIONAL LEVEL"
prn(r_nation_est, title)

#---ESTIMATION AREA ESTIMATES AND STANDARD ERRORS---#
r_est <- svyby(~P_UTD431H314_ROUT_S, ~ESTIAP, svydsg, svymean)
r_est[,-c(1)] <- round(r_est[,-c(1)]*100,2) #CONVERT INTO PERCENT ESTIMATES
```

```

r_est <- subset(r_est, select=c(1,3,5))

#SELECT ESTIMATES FOR UP-TO-DATE CASES
names(r_est) <- c("ESTIMATION AREA", "PERCENT 4:3:1:H:3:1:4 UTD", "STANDARD ERROR UTD")
title <- "PERCENT 4:3:1:H:3:1:4 ESTIMATES BY ESTIMATION AREA"
prn(r_est, title)

#####
title <- "R_STATE.R"
#####
#THIS PROGRAM WILL PRODUCE STATE ESTIMATES AND STANDARD ERRORS
#FOR P_UTD431H314_ROUT_S USING R.
#
#NOTE : THE STATE VARIABLE IS BASED ON FIPSTATE CODES ,THERE ARE
#NO STATES WITH FIPS CODES 3,7,14,43,52,57-71,73-78.
#
#R NOTES:
#1. R IS CASE SENSITIVE.
#2. A FILE PATH IS SEPERATED BY SLASH(/)
#####
library(survey) #TO USE svydesign(), svymean(), and svyby()
library(Hmisc) #TO USE prn()

dd <- "c:/nispuf15" #"path-to-data"

#--- NAME OF R DATASET ---#
in.file <- paste(dd,"NISPUF15.RData",sep="")
#---READ R DATASET---#
load(in.file)
#---FORMAT---#
UTD431H314levels=c(0,1)
UTD431H314labels=c("NOT 4:3:1:H:3:1:4 UTD", "4:3:1:H:3:1:4 UTD")
STATElevels=c(1, 2, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 16, 17,
18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35,
36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 53,
54, 55, 56, 72)
STATElabels=c(
"ALABAMA",
"ALASKA",
"ARIZONA",
"ARKANSAS",
"CALIFORNIA",
"COLORADO",
"CONNECTICUT",
"DELAWARE",
"DISTRICT OF COLUMBIA",
"FLORIDA",
"GEORGIA",
"HAWAII",
"IDAHO",
"ILLINOIS",
"INDIANA",
"IOWA",
"KANSAS",
"KENTUCKY",
"LOUISIANA",
"MAINE",
"MARYLAND",
"MASSACHUSETTS",
"MICHIGAN",

```

```

"MINNESOTA",
"MISSISSIPPI",
"MISSOURI",
"MONTANA",
"NEBRASKA",
"NEVADA",
"NEW HAMPSHIRE",
"NEW JERSEY",
"NEW MEXICO",
"NEW YORK",
"NORTH CAROLINA",
"NORTH DAKOTA",
"OHIO",
"OKLAHOMA",
"OREGON",
"PENNSYLVANIA",
"RHODE ISLAND",
"SOUTH CAROLINA",
"SOUTH DAKOTA",
"TENNESSEE",
"TEXAS",
"UTAH",
"VERMONT",
"VIRGINIA",
"WASHINGTON",
"WEST VIRGINIA",
"WISCONSIN",
"WYOMING",
"PUERTO RICO"
)
#---PROVWT_D WILL BE USED AS A WEIGHT (PROVWT_D IS THE DUAL-FRAME WEIGHT EXCLUDING TERRITORIES USE
PROVWT_D_TERR TO INCLUDE TERRITORIES)---#
#---STRATUM WILL BE USED AS A STRATUM VARIABLE FOR VARIANCE ESTIMATION ---#
R_FILE <- subset(NISPUF15, select=c(SEQNUMHH, SEQNUMC, P_UTD431H314_ROUT_S,
STATE, PROVWT_D, STRATUM))
names(R_FILE) <- c("SEQNUMHH", "SEQNUMC", "P_UTD431H314_ROUT_S", "STATE",
"WT", "STRATUM")
R_FILE <- na.omit(R_FILE)
#---ASSIGN LABELS---#
R_FILE$P_UTD431H314_ROUT_S <- factor(R_FILE$P_UTD431H314_ROUT_S, levels=UTD431H314levels,
labels=UTD431H314labels)
R_FILE$STATE <- factor(R_FILE$STATE, levels=STATElevels,
labels=STATElabels)
#---SPECIFY A SAMPLING DESIGN AND FORCE WT AS NUMERIC---#
svydsq <- svydesign(id=~SEQNUMHH, strata=~STRATUM, weights=~(as.numeric(WT)),
data=R_FILE)
#---STATE ESTIMATES AND STANDARD ERRORS---#
r_est2 <- svyby(~P_UTD431H314_ROUT_S, ~STATE, svydsq, svymean)
r_est2[-c(1)] <- round(r_est2[-c(1)]*100,2) #CONVERT INTO PERCENT ESTIMATES
r_est2 <- subset(r_est2, select=c(1,3,5)) #SELECT ESTIMATES FOR UP-TO-DATE CASES
names(r_est2) <- c("STATE", "PERCENT 4:3:1:H:3:1:4 UTD", "STANDARD ERROR UTD")
prn(r_est2, '4:3:1:H:3:1:4 ESTIMATES BY STATE')

#####
title <- "R_PROG_3.R"
#####
#THIS PROGRAM WILL PRODUCE A TABLE OF HAD_CPOX BY STATE FOR ALL RDD
#COMPLETES USING RDDWT_D. THE PROGRAM USES R.
#
#R NOTES:
#1. R IS CASE SENSITIVE.
#2. A FILE PATH IS SEPERATED BY SLASH(/)
#####

```

```

library(survey) #TO USE svydesign(), svymean(), and svyby()
library(Hmisc) #TO USE prn()
library(prettyR) #TO USE freq()

dd <- "c:/nispuf15" #"path-to-dataset"

#--- NAME OF R DATASET ---#
in.file <- paste(dd,"NISPUF15.RData",sep="")
#---READ R DATASET---#
load(in.file)
#---FORMAT---#
HAD_CPOXlevels=c(1,2,77,99)
HAD_CPOXlabels=c("YES", "NO", "DON'T KNOW", "REFUSED")
STATElevels=c(1, 2, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 16, 17,
18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35,
36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 53,
54, 55, 56, 72)
STATElabels=c(
"ALABAMA",
"ALASKA",
"ARIZONA",
"ARKANSAS",
"CALIFORNIA",
"COLORADO",
"CONNECTICUT",
"DELAWARE",
"DISTRICT OF COLUMBIA",
"FLORIDA",
"GEORGIA",
"HAWAII",
"IDAHO",
"ILLINOIS",
"INDIANA",
"IOWA",
"KANSAS",
"KENTUCKY",
"LOUISIANA",
"MAINE",
"MARYLAND",
"MASSACHUSETTS",
"MICHIGAN",
"MINNESOTA",
"MISSISSIPPI",
"MISSOURI",
"MONTANA",
"NEBRASKA",
"NEVADA",
"NEW HAMPSHIRE",
"NEW JERSEY",
"NEW MEXICO",
"NEW YORK",
"NORTH CAROLINA",
"NORTH DAKOTA",
"OHIO",
"OKLAHOMA",
"OREGON",
"PENNSYLVANIA",
"RHODE ISLAND",
"SOUTH CAROLINA",
"SOUTH DAKOTA",
"TENNESSEE",
"TEXAS",

```

```

"UTAH",
"VERMONT",
"VIRGINIA",
"WASHINGTON",
"WEST VIRGINIA",
"WISCONSIN",
"WYOMING",
"PUERTO RICO"
)
#---RDDWT_D WILL BE USED AS A WEIGHT (RDDWT_D IS THE DUAL-FRAME WEIGHT EXCLUDING TERRITORIES. USE
RDDWT_D_TERR TO INCLUDE TERRITORIES)---#
#---STRATUM WILL BE USED AS A STRATUM VARIABLE FOR VARIANCE ESTIMATION ---#
R_FILE <- subset(NISPUF15, select=c(SEQNUMHH, SEQNUMC, STATE,
HAD_CPOX, RDDWT_D, STRATUM))
names(R_FILE) <- c("SEQNUMHH", "SEQNUMC", "STATE", "HAD_CPOX",
"WT", "STRATUM")

#---ASSIGN LABELS---#
R_FILE$HAD_CPOX <- factor(R_FILE$HAD_CPOX, levels=HAD_CPOXlevels,
labels=HAD_CPOXlabels)
R_FILE$STATE <- factor(R_FILE$STATE, levels=STATElevels,
labels=STATElabels)
R_FILE <- na.omit(R_FILE)
summary(R_FILE$HAD_CPOX)

#---SPECIFY A SAMPLING DESIGN AND FORCE WT AS NUMERIC---#
svydsg <- svydesign(id=~SEQNUMHH, strata=~STRATUM, weights=~(as.numeric(WT)),
data=R_FILE)

#---U.S. TOTAL ESTIMATES AND STANDARD ERRORS---#
r_nation <- svymean(~HAD_CPOX, svydsg)
PERCENT_UTD <- round(r_nation*100,2) #CONVERT INTO PERCENT ESTIMATES(MEAN)
SE_UTD <- round(SE(r_nation)*100,2) #CONVERT INTO PERCENT ESTIMATES(SE)
r_nation_est3 <- cbind(PERCENT_UTD, SE_UTD)
prn(r_nation_est3, "PERCENT HAD_CPOX = YES ESTIMATES AT A NATIONAL
LEVEL\n")

#---HAD_CPOX = YES ESTIMATES BY STATE---#
r_est3 <- svyby(~HAD_CPOX, ~STATE, svydsg, svymean)
r_est3[,c(1)] <- round(r_est3[,c(1)]*100,2) #CONVERT INTO PERCENT ESTIMATES
r_est3 <- subset(r_est3, select=c(1,2,6)) #SELECT ESTIMATES FOR HAD_CPOX=YES
names(r_est3) <- c("STATE", "PERCENT HAD_CPOX=YES", "STANDARD ERROR
HAD_CPOX=Y")
prn(r_est3, 'PERCENT HAD_CPOX ESTIMATES BY STATE')

#####
title <- "PROG_4.R"
#####
#TABLE OF P_UTD431H314_ROUT_S BY INCPOV1 BY RACE_K. SAVE % UTD
#ESTIMATES (NOT S.E.'S) FOR USE IN THE PROGRAM GRAPH_4.
#
#THIS PROGRAM WILL PRODUCE ESTIMATES USING R.
#
#R NOTES:
#1. R IS CASE SENSITIVE.
#2. A FILE PATH IS SEPERATED BY SLASH(/)
#####
library(survey) #TO USE svydesign(), svymean(), and svyby()
library(Hmisc) #TO USE prn()

dd <- "c:/nispuf15" #"path-to-dataset"

```

```
out <- "c:/nispuf15" #"path-to-output"
```

```
### NAME OF R DATASET ###
in.file <- paste(dd, "/NISPUF15.RData", sep="")
### READ R DATASET ###
load(in.file)
### FORMAT ###
UTD431H314levels=c(0,1)
UTD431H314labels=c("NOT 4:3:1:H:3:1:4 UTD", "4:3:1:H:3:1:4 UTD")
RACE_PUFlevels=c(1,2,3)
RACE_PUFlabels=c("WHITE ONLY", "BLACK ONLY", "OTHER + MULTIPLE RACE")
INCPOVlevels=c(1,2,3,4)
INCPOVlabels=c("ABOVE POVERTY, > $75K", "ABOVE POVERTY, <= $75K", "BELOW POVERTY", "UNKNOWN")
### PROVWT_D WILL BE USED AS A WEIGHT (PROVWT_D IS THE DUAL-FRAME WEIGHT EXCLUDING TERRITORIES. USE
PROVWT_D_TERR TO INCLUDE TERRITORIES) ###
### STRATUM WILL BE USED AS A STRATUM VARIABLE FOR VARIANCE ESTIMATION ###
R_FILE <- subset(NISPUF15, select=c(SEQNUMHH, SEQNUMC, P_UTD431H314_ROUT_S, RACE_K, INCPOV1, PROVWT_D, STRATUM))
names(R_FILE) <- c("SEQNUMHH", "SEQNUMC", "P_UTD431H314_ROUT_S", "RACE_K", "INCPOV1", "WT", "STRATUM")
### ASSIGN LABELS ###
R_FILE$P_UTD431H314_ROUT_S <- factor(R_FILE$P_UTD431H314_ROUT_S, levels=UTD431H314levels, labels=UTD431H314labels,
exclude=NULL)
R_FILE$RACE_K <- factor(R_FILE$RACE_K, levels=RACE_PUFlevels, labels=RACE_PUFlabels, exclude=NULL)
R_FILE$INCPOV1 <- factor(R_FILE$INCPOV1, levels=INCPOVlevels, labels=INCPOVlabels, exclude=NULL)
### UNWEIGHTED FREQUENCIES ###
unwt_freq <- function(UNWT.VAR) {#FUNCTION TO PRINT UNWEIGHTED FREQUENCIES
unwt.tab <- wtd.table(UNWT.VAR, weights= NULL, type='table')
unwtd.freq <- data.frame(cbind(
unwt.tab, round(unwt.tab/sum(unwt.tab)*100,2),
cumsum(unwt.tab), cumsum(round(unwt.tab/sum(unwt.tab)*100,2))))
names(unwtd.freq) <- c("Frequency", "Percent", "Cumulative Frequency", "Cumulative Percent")
unwtd.title <- paste("Table 4A. Q1/2015 - Q4/2015", 'UNWEIGHTED FREQUENCIES', label(UNWT.VAR), sep="\n")
label(unwtd.freq) <- unwtd.title
print(unwtd.freq)
}
unwt_freq(R_FILE$P_UTD431H314_ROUT_S)
unwt_freq(R_FILE$INCPOV1)
unwt_freq(R_FILE$RACE_K)
R_FILE <- na.omit(R_FILE)
### SPECIFY A SAMPLING DESIGN AND FORCE WT AS NUMERIC ###
svydsg <- svydesign(id=~SEQNUMHH, strata=~STRATUM, weights=~(as.numeric(WT)),
data=R_FILE)
### PERCENT 4:3:1:H:3:1:4 UP-TO-DATE AND ESTIMATED STANDARD ERRORS ###
r_est4 <- svyby(~P_UTD431H314_ROUT_S, ~RACE_K+INCPOV1, svydsg, svymean)
r_est4[, -c(1,2)] <- round(r_est4[, -c(1,2)]*100,2) #CONVERT INTO PERCENT ESTIMATES
r_est4 <- subset(r_est4, select=c(1,2,4,6)) #SELECT ESTIMATES FOR UP-TODATE CASES
names(r_est4) <- c("RACE", "INCOME", "PERCENT_UTD", "STANDARD_ERROR_UTD")
title <- "Table 4B. Q1/2015 - Q4/2015, Percent 4:3:1:H:3:1:4 UTD and Estimated Standard Errors"
prn(r_est4, title)
### SAVE PERCENT UP-TO-DATE ESTIMATES FOR USE IN THE PROGRAM GRAPH_4 ###
r_est4 <- subset(r_est4, select=c(RACE, INCOME, PERCENT_UTD))
save(r_est4, file=paste(out, "/r_est4", sep=""))

#####
title <- "GRAPH_4.R"
#####
#THIS PROGRAM BUILDS OFF OF THE PROGRAM PROG_4. IT PRODUCES A CHART OF
#P_UTD431H314_ROUT_S BY INCPOV1 BY RACE_K. IT CREATES A BAR CHART IN R GRAPH FOR
#THE 4X3 = 12 CELLS.
```

```

#R NOTES:
#1. R IS CASE SENSITIVE.
#2. A FILE PATH IS SEPERATED BY SLASH(/)
#####
library(survey) #TO USE svydesign(), svymean(), and svyby()
library(Hmisc) #TO USE prn()


dd <- "c:/nispuf15" #---SPECIFY PATH TO R DATASET THAT WAS THE OUTPUT OF R_PROG_4---#

out <- "c:/nispuf15" #---SPECIFY THE PATH FOR WHERE YOU WANT THE CHART OUTPUT TO GO---#

#---NAME OF R DATASET OUTPUT FROM R_PROG_4---#
in.file <- paste(dd,"/r_est4",sep="")
#---READ R DATASET---#
load(in.file)
#---BARChart---#
#NOTE:R DOES NOT SUPPORT CREATING A HTML FILE CONTAINING A BARChart#
#CREATE A DATA MATRIX FOR DRAWING A BARChart#
utd431H314 <- matrix(r_est4$PERCENT_UTD, nrow=3, ncol=4, byrow=F, dimnames=list(levels(r_est4$RACE), levels(r_est4$INCOME)))
#CREATE GRAPH_4.GIF#
barplot(utd431H314, beside=TRUE, space=c(0.2,1),
col = c("wheat", "lightpink2", "forestgreen"),
axis.lty = 1,
sub="(Graph 4 using 'R')", cex.sub=1, ylim=c(0,120),
xlab="Poverty Status",
ylab="4:3:1:H:3:1:4 Up-To-Date (%)", cex=1, cex.names=1, border=NA)
legend("top", rownames(utd431H314), col=c("wheat", "lightpink2",
"forestgreen"), title="Race of Child", pch=15, cex=1)
title1 <- "Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4 \n"
title2 <- "by Race and Poverty Status, National Immunization Survey, 2015\n"
mtext(paste(title1,title2), cex=1.3)

```

Percentage of Children Up-to-date with Vaccine Series 4:3:1:H:3:1:4
by Race and Poverty Status, National Immunization Survey, 2015



Appendix E: Alphabetical Listing of Variables that are in the 2004-2015 Public-Use Data Files

Table E.1 Alphabetical Listing of Variables that are in the 2004-2015 Public-Use Data Files*

| Variable Name | Variable Label [†] | Year of Data Collection | | | | | | | | | | | | | Notes [§] |
|---------------|---|-------------------------|------|------|------|------|------|------|------|------|------|------|------|---|--------------------|
| | | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | | |
| AGECPOXR | AGE IN MONTHS AT CHICKEN POX DISEASE (RECODE) | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces IAGECPXR starting 2005. This version is not imputed. | |
| AGEGRP | AGE CATEGORY OF CHILD (19-23, 24-29, 30-35 MO) (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| ALL4SHOT | HH REPORT OF 4:3:1:3 UP-TO-DATE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. | |
| BF_ENDR | DURATION OF BREAST FEEDING IN DAYS (TOPCODE) | Y | Y | | | | | | | | | | | Dropped starting in 2006 because of question wording change. Replaced by BF_ENDR06. | |
| BF_ENDR06 | DURATION OF BREAST FEEDING IN DAYS (RECODE) | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces BF_ENDR starting 2006. | |
| BF_EXCLR | DURATION OF EXCLUSIVE BREAST FEEDING IN DAYS (TOPCODE) | Y | Y | | | | | | | | | | | Dropped starting in 2006 because of question wording change. Replaced by BF_EXCLR06. | |
| BF_EXCLR06 | DURATION OF EXCLUSIVE BREAST/FORMULA FEEDING IN DAYS (RECODE) | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces BF_EXCLR starting 2006. | |
| BF_FORMR06 | AGE IN DAYS WHEN CHILD FIRST FED FORMULA (TOPCODE) | | | Y | Y | | | | | | | | | Question CBF_03_X added starting 2006. Replaced by BF_FORMR06 starting 2008. | |
| BF_FORMR08 | AGE IN DAYS WHEN CHILD FIRST FED FORMULA (RECODE) | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Replaces BF_FORMR06 to add a "never fed formula" code. | |
| BFENDFL | DURATION OF BREAST FEEDING EXCEEDS CHILD AGE IN DAYS, WITH BUFFER | Y | Y | | | | | | | | | | | Dropped starting in 2006 because of question wording change. Replaced by BFENDFL06. | |
| BFENDFL06 | DURATION OF BREAST FEEDING EXCEEDS CHILD AGE IN DAYS, WITH BUFFER | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces BFENDFL starting 2006. | |
| BFEXCLFL | DURATION OF EXCLUSIVE BREAST FEEDING EXCEEDS TOTAL BREASTFEEDING, WITH BUFFER | Y | Y | | | | | | | | | | | Dropped starting in 2006 because question wording change does not allow it to be derived. | |
| BFFORMFL06 | AGE IN DAYS WHEN CHILD FIRST FED FORMULA EXCEEDS CHILD AGE IN DAYS, WITH BUFFER | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Question CBF_03_X added starting 2006. | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| C_431 | HH REPORT OF 4:3:1 UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_4313 | HH REPORT OF 4:3:1:3 UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_DTP | HH REPORT OF 4+ DT-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_HEP | HH REPORT OF 3+ HEPATITIS B-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_HIB | HH REPORT OF 3+ HIB-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_MMR | HH REPORT OF 1+ MEASLES-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_POL | HH REPORT OF 3+ POLIO-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C_VRC | HH REPORT OF 1+ VARICELLA-CONTAINING UP-TO-DATE BY SHOT CARD USE | Y | Y | | | | | | | | | | | Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| C1R | NUMBER OF PEOPLE IN HOUSEHOLD (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| C5R | RELATIONSHIP OF RESPONDENT TO CHILD (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| CBF_01 | WAS CHILD EVER BREAST FED OR FED BREAST MILK? | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| CEN_REG | CENSUS REGION BASED ON TRUE STATE OF RESIDENCE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| CHILDNM | NUMBER OF CHILDREN LESS THAN 18 YEARS IN HH (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| CWIC_01 | CHILD EVER RECEIVED WIC BENEFITS? | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| CWIC_02 | CHILD CURRENTLY RECEIVING WIC BENEFITS? | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| D6R | NUMBER OF VACCINATION PROVIDERS IDENTIFIED BY RESPONDENT (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| D7 | CONSENT TO OBTAIN CHILD'S IMMUNIZATION RECORDS FROM PROVIDERS | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP1 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DDTP2 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP3 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP4 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP5 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP6 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP7 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP8 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DDTP9 | AGE IN DAYS OF PROV-REPTD DT-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DFLU1 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU2 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU3 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU4 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU5 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU6 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU7 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU8 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DFLU9 | AGE IN DAYS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DH1N1 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #1 | | | | | | | | Y | Y | Y | | | Introduced in 2010. Removed 2013. |
| DH1N2 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #2 | | | | | | | | Y | Y | Y | | | Introduced in 2010. Removed 2013. |
| DH1N3 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #3 | | | | | | | | Y | Y | Y | | | Introduced in 2010. Removed 2013. |
| DH1N4 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #4 | | | | | | | | Y | Y | Y | | | Introduced in 2010. Removed 2013. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DH1N5 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #5 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| DH1N6 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #6 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| DH1N7 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #7 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| DH1N8 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #8 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| DH1N9 | AGE IN DAYS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #9 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| DHEPA1 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA2 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA3 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA4 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA5 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA6 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA7 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA8 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPA9 | AGE IN DAYS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DHEPB1 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB2 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB3 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB4 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB5 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB6 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB7 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DHEPB8 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHEPB9 | AGE IN DAYS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DHIB1 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB2 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB3 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB4 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB5 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB6 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB7 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB8 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DHIB9 | AGE IN DAYS OF PROV-REPTD HIB-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DISPCODE | NIS PROVIDER RECORD-CHECK DISPOSITION CODE | Y | Y | Y | Y | Y | Y | Y | Y | | | | | Dropped starting in 2012. |
| DMMR1 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMMR2 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMMR3 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMMR4 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMMR5 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMMR6 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMMR7 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |

| Variable Name | Variable Label [†] | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DMMR8 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMMR9 | AGE IN DAYS OF PROV-REPTD MEASLES-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMP1 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMP2 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMP3 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMP4 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMP5 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMP6 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMP7 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMP8 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMP9 | AGE IN DAYS OF PROV-REPTD MUMPS-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMPRB1 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMPRB2 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMPRB3 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMPRB4 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DMPRB5 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMPRB6 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DMPRB7 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMPRB8 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DMPRB9 | AGE IN DAYS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DPCV1 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV2 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV3 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV4 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV5 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV6 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV7 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV8 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPCV9 | AGE IN DAYS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DPOLIO1 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO2 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO3 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO4 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO5 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO6 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO7 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DPOLIO8 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DPOLIO9 | AGE IN DAYS OF PROV-REPTD POLIO-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DRB1 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB2 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB3 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB4 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB5 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB6 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB7 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB8 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DRB9 | AGE IN DAYS OF PROV-REPTD RUBELLA-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DROT1 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT2 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT3 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT4 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT5 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT6 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT7 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT8 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DROT9 | AGE IN DAYS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DTP_SOUR | SHOT CARD USED FOR DTP REPORTING | Y | | | | | | | | | | | | Dropped starting in 2005 because this variable is redundant with variable SHOTCARD. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| DTP1_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP2_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP3_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP4_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP5_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP6_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP7_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP8_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DTP9_AGE | AGE IN MONTHS OF PROV-REPTD DT-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DVRC1 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DVRC2 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DVRC3 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DVRC4 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| DVRC5 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DVRC6 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DVRC7 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DVRC8 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| DVRC9 | AGE IN DAYS OF PROV-REPTD VARICELLA-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| EDUC1 | EDUCATION OF MOTHER CATEGORIES (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ | |
|-----------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|--------------------|
| ENTRY2 | CHILD LIVES IN STATE WITH HEPATITIS B STATE ENTRY LAW FOR DAY CARE/HEAD START (2001-2002 SCHOOL YEAR) | Y | | | | | | | | | | | | Dropped starting in 2005. | |
| ESTIAP | ESTIMATION IAP AREA OF RESIDENCE | | Y | | | | | | | | | | | New estimation area variable starting in 2005. Replaced ITRUEIAP. | |
| ESTIAP06 | ESTIMATION IAP AREA OF RESIDENCE | | | Y | | | | | | | | | | New starting 2006 because estimation areas were modified. | |
| ESTIAP07 | ESTIMATION AREA OF RESIDENCE | | | | Y | | | | | | | | | New starting 2007 because estimation areas were modified. | |
| ESTIAP08 | ESTIMATION AREA OF RESIDENCE | | | | | Y | | | | | | | | New starting 2008 because estimation areas were modified. | |
| ESTIAP09 | ESTIMATION AREA OF RESIDENCE | | | | | | Y | | | | | | | New starting 2009 because estimation areas were modified. | |
| ESTIAP10 | ESTIMATION AREA OF RESIDENCE | | | | | | | Y | | | | | | New starting 2010 because estimation areas were modified. | |
| ESTIAP11 | ESTIMATION AREA OF RESIDENCE | | | | | | | | Y | | | | | New starting 2011 because estimation areas were modified. | |
| ESTIAP12 | ESTIMATION AREA OF RESIDENCE | | | | | | | | | Y | | | | New starting 2012 because estimation areas were modified. | |
| ESTIAP13 | ESTIMATION AREA OF RESIDENCE | | | | | | | | | | Y | | | New starting 2013 because estimation areas were modified. | |
| ESTIAP14 | ESTIMATION AREA OF RESIDENCE | | | | | | | | | | | Y | | New starting 2014 because estimation areas were modified. | |
| ESTIAP15 | ESTIMATION AREA OF RESIDENCE | | | | | | | | | | | | Y | New starting 2015 because estimation areas were modified. | |
| EST_GRANT | AREA OF RESIDENCE ACCORDING TO THE 56 ORIGINAL CORE GRANTEE AREAS | | | | | | | | | | Y | Y | Y | Y | New starting 2012. |
| FLU1_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU2_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU3_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU4_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU5_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU6_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU7_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |
| FLU8_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|--|
| FLU9_AGE | AGE IN MONTHS OF PROV-REPTD SEASONAL FLU-CONTAINING VACCINATION #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| FRSTBRN | FIRST BORN STATUS OF CHILD | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| FUL2_MMR | HOUSEHOLD REPORT OF 1+ MMR AT ANY AGE | Y | | | | | | | | | | | | Replaced by FULL_MMR starting in 2005. |
| FULL_CPO | HH REPORT OF 1+ VARICELLA-CONTAINING SHOT AT ANY AGE | Y | Y | | | | | | | | | | | Starting 2005, a code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| FULL_DTP | HH REPORT OF 4+ DT-CONTAINING SHOT | Y | Y | | | | | | | | | | | Starting 2005, a code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| FULL_HEP | HH REPORT OF 3+ HEPATITIS B-CONTAINING SHOTS | Y | Y | | | | | | | | | | | Starting 2005, a code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| FULL_HIB | HH REPORT OF 3+ HIB-CONTAINING SHOTS | Y | Y | | | | | | | | | | | Starting 2005, a code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| FULL_MMR | HH REPORT OF 1+ MEASLES-CONTAINING SHOT AT ANY AGE | | Y | | | | | | | | | | | Replaced FUL2_MMR starting in 2005. A code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| FULL_POL | HH REPORT OF 3+ POLIO-CONTAINING SHOTS | Y | Y | | | | | | | | | | | Starting 2005, a code of 88 added for children with unknown UTD status. Dropped starting in 2006 because no longer possible to derive due to shortened Section B. |
| H1N1_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #1 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N2_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #2 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N3_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #3 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| H1N4_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #4 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N5_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #5 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N6_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #6 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N7_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #7 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N8_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #8 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| H1N9_AGE | AGE IN MONTHS OF PROV-REPTD MONOVALENT 2009 H1N1 FLU VACCINATION #9 | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| HAD_CPOX | CHILD EVER HAD CHICKEN POX DISEASE? | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces I_HADCPX starting in 2005. This version is not imputed. |
| HEA1_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA2_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA3_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA4_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA5_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA6_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA7_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA8_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEA9_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS A-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| HEP_BRTH | HEPATITIS B-CONTAINING SHOT GIVEN AT BIRTH FLAG | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP_FLAG | HEPATITIS B BIRTH SHOT DATE IMPUTATION FLAG | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| HEP1_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP2_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP3_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP4_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP5_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP6_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP7_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP8_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HEP9_AGE | AGE IN MONTHS OF PROV-REPTD HEPATITIS B-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| HH_DTP | HH REPORT OF NUMBER OF DT-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| HH_FLU | HH REPORT OF NUMBER OF SEASONAL FLU VACCINATIONS RECEIVED IN THE 12 MONTHS PRIOR TO INTERVIEW | | | | Y | Y | | Y | | | | | | FLU questions added to the HH questionnaire starting in 2007. Dropped in 2009 due to mid-year questionnaire changes. Reinstated in 2010. Dropped again in 2011 due to mid-year questionnaire changes. |
| HH_H1N | HH REPORT OF NUMBER OF MONOVALENT 2009 H1N1 FLU VACCINATIONS RECEIVED IN THE 12 MONTHS PRIOR TO INTERVIEW | | | | | | | Y | | | | | | H1N1 flu questions added to the HH questionnaire starting in 2009. Introduced in the PUF in 2010. Dropped in 2011 due to mid-year questionnaire changes. |
| HH_HEPB | HH REPORT OF NUMBER OF HEPATITIS B-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| HH_HIB | HH REPORT OF NUMBER OF HIB-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|--|
| HH_MCV | HH REPORT OF NUMBER OF MEASLES-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| HH_POL | HH REPORT OF NUMBER OF POLIO-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| HH_VRC | HH REPORT OF NUMBER OF VARICELLA-CONTAINING SHOTS RECEIVED | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| HIB1_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB2_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB3_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB4_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB5_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB6_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB7_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB8_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| HIB9_AGE | AGE IN MONTHS OF PROV-REPTD HIB-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| HUTD4313 | HOUSEHOLD REPORT OF 4:3:1:3 UTD (UP-TO-DATE) | Y | | | | | | | | | | | | Dropped starting in 2005 because this variable is redundant with variable ALL4SHOT. |
| I_HADCPX | DID CHILD EVER HAVE CHICKEN POX? | Y | | | | | | | | | | | | Replaced by HAD_CPOX starting in 2005. |
| I_HISP_K | HISPANIC ORIGIN OF CHILD | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| IAGECPXR | AGE IN MONTHS WHEN CHILD HAD CHICKEN POX (RECODE) | Y | | | | | | | | | | | | Replaced by AGECPXR starting in 2005. |
| INCPORAR | INCOME TO POVERTY RATIO (RECODE) | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces INCPORAT starting 2005. INCPORAT used categories whereas INCPORAR is continuous. INCPORAR has been top- and bottom-coded. |

| Variable Name | Variable Label [†] | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|--|
| INCPORAT | INCOME TO POVERTY RATIO | Y | | | | | | | | | | | | Replaced by INCPORAR starting in 2005. |
| INCPOV1 | POVERTY STATUS | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces INCPOV1R starting in 2005. INCPOV1R used two categories whereas INCPOV1 uses three. |
| INCPOV1R | POVERTY STATUS (RECODE) | Y | | | | | | | | | | | | Replaced by INCPOV1 starting in 2005. |
| INCQ298A | FAMILY INCOME CATEGORIES (RECODE) | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces INCQ298R starting in 2005. INCQ298A uses different categories than were used by INCQ298R. |
| INCQ298R | FAMILY INCOME CATEGORIES (RECODE) | Y | | | | | | | | | | | | Replaced by INCQ298A starting in 2005. |
| INOPHONR | LENGTH OF INTERRUPTION IN TELEPHONE SERVICE IN DAYS (RECODE) | Y | Y | Y | Y | Y | Y | | | | | | | Removed in 2010 due to questionnaire change. |
| INS_1 | IS CHILD COVERED BY HEALTH INSURANCE PROVIDED THROUGH EMPLOYER OR UNION? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INS_11 | ANY TIME WHEN CHILD WAS NOT COVERED BY ANY HEALTH INSURANCE? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INS_2 | IS CHILD COVERED BY ANY MEDICAID PLAN? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INS_3 | IS CHILD COVERED BY S-CHIP? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INS_3A | IS CHILD COVERED BY ANY MEDICAID PLAN OR S-CHIP? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INS_4 | IS CHILD COVERED BY INDIAN HEALTH SERVICE? | | | | Y | Y | | | | | | | | Replaced by INS_4_5 starting 2009. |
| INS_4_5 | IS CHILD COVERED BY INDIAN HEALTH SERVICE, MILITARY HEALTH CARE, TRICARE, CHAMPUS, OR CHAMP-VA? | | | | | | Y | Y | Y | Y | Y | Y | Y | Replaces INS_4 and INS_5 starting 2009. |
| INS_5 | IS CHILD COVERED BY MILITARY HEALTH CARE, TRICARE, CHAMPUS, OR CHAMP-VA? | | | | Y | Y | | | | | | | | Replaced by INS_4_5 starting 2009. |
| INS_6 | IS CHILD COVERED BY ANY OTHER HEALTH INSURANCE OR HEALTH CARE PLAN? | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| INTRP | PHONE INTERRUPTION OF 7 DAYS OR MORE IN PAST YEAR? | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ITRUEIAP | IAP AREA OF CURRENT RESIDENCE | Y | | | | | | | | | | | | The new estimation area variable starting in 2005 is ESTIAP. |
| LANGUAGE | LANGUAGE IN WHICH INTERVIEW WAS CONDUCTED | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| M_AGEGRP | AGE OF MOTHER CATEGORIES (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MARITAL | MARITAL STATUS OF MOTHER CATEGORIES (RECODE) | Y | Y | Y | Y | Y | | | | | | | | Replaced by MARITAL2 starting 2009. |
| MARITAL2 | MARITAL STATUS OF MOTHER (RECODE) | | | | | | Y | Y | Y | Y | Y | Y | Y | Replaces MARITAL starting 2009. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| MMR1_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MMR2_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MMR3_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MMR4_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MMR5_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MMR6_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MMR7_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MMR8_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MMR9_AGE | AGE IN MONTHS OF PROV-REPTD MEASLES-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MOBIL | GEOGRAPHIC MOBILITY STATUS: STATE OF RESIDENCE OF CHILD AT BIRTH VERSUS CURRENT STATE | Y | | | | | | | | | | | | Replaced by MOBIL_I starting in 2005. |
| MOBIL_I | GEOGRAPHIC MOBILITY STATUS: STATE OF RESIDENCE OF CHILD AT BIRTH VERSUS CURRENT STATE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Replaces MOBIL starting in 2005. This version is imputed. |
| MP1_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MP2_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MP3_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MP4_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MP5_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MP6_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|-------------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| MP7_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MP8_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MP9_AGE | AGE IN MONTHS OF PROV-REPTD MUMPS-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MPR1_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MPR2_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MPR3_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MPR4_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| MPR5_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MPR6_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MPR7_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MPR8_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| MPR9_AGE | AGE IN MONTHS OF PROV-REPTD (MUMPS/RUBELLA)-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| N_PRVR | NUMBER OF PROVIDERS RESPONDING WITH VACCINATION DATA FOR CHILD (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| NUM_CELLS_PARENTS | NUMBER OF WORKING CELL PHONES USUALLY USED BY PARENTS OR GUARDIANS | | | | | | Y | Y | Y | Y | Y | Y | Y | |
| NUM_CELLS_HH | NUMBER OF WORKING CELL PHONES HOUSEHOLD MEMBERS HAVE AVAILABLE FOR PERSONAL USE | | | | | | Y | Y | Y | Y | Y | Y | Y | |
| NUM_PHONE | NUMBER OF RESIDENTIAL TELEPHONE NUMBERS IN HOUSEHOLD (EXCLUDING CELL PHONES) | | | | | | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|--|
| P_NUHEPX | NUMBER OF HEPATITIS B-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUHIBX | NUMBER OF HIB-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUHPHB | NUMBER OF HEPATITIS B/HIB COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMIL | NUMBER OF MONOVALENT 2009 H1N1 FLU VACCINATIONS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| P_NUMIM | NUMBER OF MONOVALENT 2009 H1N1 FLU SPRAY VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| P_NUMIN | NUMBER OF INJECTED MONOVALENT 2009 H1N1 FLU VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| P_NUMDAH | NUMBER OF DTAP/HIB COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMDHB | NUMBER OF DTP/HIB CONTAINING SHOTS DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | | | | | | | | Dropped in 2009 due to change to IHQ shotgrid. |
| P_NUMDHI | NUMBER OF DTAP/HEPB/IPV COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMDHM | NUMBER OF DTP/HIB COMBO SHOTS DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | | | | | | | | Dropped in 2009 due to change to IHQ shotgrid. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_NUMDIH | NUMBER OF DTAP/IPV/HIB COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to change to IHQ shotgrid. |
| P_NUMDTA | NUMBER OF DTAP-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMDTM | NUMBER OF DT-ONLY SHOTS DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | | | | | | | | Dropped in 2009 due to change to IHQ shotgrid. |
| P_NUMDTP | NUMBER OF DT-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMFLU | NUMBER OF SEASONAL FLU-CONTAINING VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMFLUL | NUMBER OF SEASONAL FLU-CONTAINING VACCINATIONS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| P_NUMFLUM | NUMBER OF SEASONAL FLU SPRAY VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| P_NUMFLUN | NUMBER OF INJECTED SEASONAL FLU VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| P_NUMH1N | NUMBER OF MONOVALENT 2009 H1N1 FLU VACCINATIONS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_NUMH2 | NUMBER OF HIB-SANOFI or HIB-GLAXOSMITHKLINE SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | | | | | | | Added in 2009 due to change to IHQ shotgrid. Replaced in 2010 by P_NUMHG and P_NUMHS. |
| P_NUMHEA | NUMBER OF HEPATITIS A-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMHEN | NUMBER OF HEPATITIS B-CONTAINING SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMHEP | NUMBER OF HEPATITIS B-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMHG | NUMBER OF HIB-GLAXOSMITHKLINE SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_NUMHHY | NUMBER OF HIB-MENCY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | | | | | Y | Y | Added in 2014 due to change in IHQ shotgrid. |
| P_NUMHIB | NUMBER OF HIB-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMHIN | NUMBER OF HIB-CONTAINING SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMHION | NUMBER OF HIB-ONLY SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to change to IHQ shotgrid. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|--|
| P_NUMHM | NUMBER OF HIB-MERCK SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to change to IHQ shotgrid. |
| P_NUMHS | NUMBER OF HIB-SANOFI SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_NUMIPV | NUMBER OF IPV-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMCN | NUMBER OF MEASLES-CONTAINING SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMMR | NUMBER OF MEASLES-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMMRX | NUMBER OF MMR-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMMX | NUMBER OF MMR-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMP | NUMBER OF MUMPS-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMPR | NUMBER OF (MUMPS/RUBELLA)-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---------------------|
| P_NUMMRV | NUMBER OF MMR/VARICELLA COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMS | NUMBER OF MEASLES-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMSM | NUMBER OF MEASLES/MUMPS COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMMSR | NUMBER OF MEASLES/RUBELLA COMBO SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMOLN | NUMBER OF POLIO SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMOPV | NUMBER OF OPV-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMPCC | NUMBER OF PCV CONJUGATE SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMPCC13 | NUMBER OF PNEUMOCOCCAL CONJUGATE-13 SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_NUMPCC7 | NUMBER OF PNEUMOCOCCAL CONJUGATE-7 SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_NUMPCCN | NUMBER OF PNEUMOCOCCAL CONJUGATE SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_NUMPCN | NUMBER OF PCV SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMPCP | NUMBER OF PCV POLYSACCHARIDE SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMPCV | NUMBER OF PNEUMOCOCCAL-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMPOL | NUMBER OF POLIO-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMRB | NUMBER OF RUBELLA-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMRG | NUMBER OF ROTARIX-GSK SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Starting in 2009, rotavirus type boxes were added to the IHQ shot grid. |
| P_NUMRM | NUMBER OF ROTATEQ-MERCK SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Starting in 2009, rotavirus type boxes were added to the IHQ shot grid. |
| P_NUMRO | NUMBER OF ROTAVIRUS SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Starting in 2009, rotavirus type boxes were added to the IHQ shot grid. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|------------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_NUMROT | NUMBER OF ROTAVIRUS-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMTPM | NUMBER OF DTP-ONLY SHOTS DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | | | | | | | | Dropped in 2009 due to change to IHQ shotgrid. |
| P_NUMTPN | NUMBER OF DT-CONTAINING SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMVRC | NUMBER OF VARICELLA-CONTAINING SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMVRN | NUMBER OF VARICELLA-CONTAINING SHOTS OF UNKNOWN TYPE BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_NUMVRX | NUMBER OF VARICELLA-ONLY SHOTS BY 36 MONTHS OF AGE DETERMINED FROM PROVIDER INFO, EXCLUDING ANY VACCINATIONS AFTER THE HH INTERVIEW DATE. | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_U12VRC | UTD (UP-TO-DATE) FLAG FOR PROVIDER 1+ VARICELLA-CONTAINING SHOT AT 12+ MONTHS, BY 36 MONTHS OF AGE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTD331 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3:3:1 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTD431 | <i>UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE.</i> | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTD431H_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1:3* BY 36 MONTHS OF AGE, USING THE ROUTINE, STRICT DEFINITION OF HIB UTD, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_UTD431H3_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1:3*~3 BY 36 MONTHS OF AGE, USING THE ROUTINE, STRICT DEFINITION OF HIB UTD, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTD431H31_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1:3*~3:1 BY 36 MONTHS OF AGE (INCLUDES 1+ VARICELLA-CONTAINING AT AGE 12+ MTHS) USING THE ROUTINE, STRICT DEFINITION OF HIB UTD, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTD431H313_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1:3*~3:1:3 BY 36 MONTHS OF AGE (INCLUDES 1+ VARICELLA-CONTAINING AT AGE 12+ MTHS) USING THE ROUTINE, STRICT DEFINITION OF HIB UTD, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTD431H314_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4:3:1:3*~3:1:4 BY 36 MONTHS OF AGE (INCLUDES 1+ VARICELLA-CONTAINING AT AGE 12+ MTHS) USING THE ROUTINE, STRICT DEFINITION OF HIB UTD, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTDFL1 | UTD FLAG FOR PROVIDER SEASONAL INFLUENZA VARIABLE 1 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | | Removed 2014. |
| P_UTDFL2 | UTD FLAG FOR PROVIDER SEASONAL INFLUENZA VARIABLE 2 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | | | Removed 2014. |
| P_UTDFL3 | UTD FLAG FOR PROVIDER SEASONAL INFLUENZA VARIABLE 3 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | Y | Y | Y | Y | Y | Y | Y | | | Removed 2014. |
| P_UTDH1N_1 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 1+ MONOVALENT 2009 H1N1 FLU VACCINATION BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE AND EXCLUDING VACCINATIONS GIVEN PRIOR TO 10/5/2009. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|------------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| P_UTDH1N_2 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 2+ MONOVALENT 2009 H1N1 FLU VACCINATIONS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE AND EXCLUDING VACCINATIONS GIVEN PRIOR TO 10/5/2009. | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| P_UTDHEP | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ HEPATITIS B-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDHEPA1 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 1+ HEPATITIS A-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| P_UTDHEPA2 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 2+ HEPATITIS A-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_UTDHIB | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ HIB-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDHIB_ROUT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ HIB DOSES BY 36 MONTHS OF AGE, BASED ON THE ROUTINE (NON-SHORTAGE) HIB RECOMMENDATIONS AND A STRICT TREATMENT OF HIB SHOTS OF UNKNOWN TYPE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTDHIB_SHORT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ HIB DOSES BY 36 MONTHS OF AGE, BASED ON THE HIB SHORTAGE RECOMMENDATIONS AND A STRICT TREATMENT OF HIB SHOTS OF UNKNOWN TYPE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | Added in 2009 due to new Hib vaccination recommendations. |
| P_UTDMCV | UTD (UP-TO-DATE) FLAG FOR PROVIDER 1+ MEASLES-CONTAINING SHOT BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDMMX | UTD FLAG FOR PROVIDER 1+ MMR COMBO SHOT BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---------------------|
| P_UTDPC3 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ PNEUMOCOCCAL-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDPCV | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4+ PNEUMOCOCCAL-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDPCVB13 | UTD (UP-TO-DATE) INDICATOR FOR PROVIDER 1+ PNEUMOCOCCAL VACCINATIONS OF TYPE CONJUGATE 13, GIVEN 4+ DOSES OF TYPE CONJUGATE 7, BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| P_UTDPOL | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ POLIO-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDROT_S | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ ROTAVIRUS DOSES BY 36 MONTHS OF AGE, BASED ON A STRICT TREATMENT OF ROTAVIRUS VACCINATIONS OF UNKNOWN TYPE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDTP3 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 3+ DT-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| P_UTDTP4 | UTD (UP-TO-DATE) FLAG FOR PROVIDER 4+ DT-CONTAINING SHOTS BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV1_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV2_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV3_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV4_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV5_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV6_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| PCV7_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV8_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PCV9_AGE | AGE IN MONTHS OF PROV-REPTD PNEUMOCOCCAL-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| PDAT | CHILD HAS ADEQUATE PROVIDER DATA | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL1_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL2_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL3_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL4_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL5_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL6_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL7_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL8_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| POL9_AGE | AGE IN MONTHS OF PROV-REPTD POLIO-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| PROV_FAC | PROVIDER FACILITY TYPES | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PROVWT | WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN (EXCLUDING U.S. VIRGIN ISLANDS) | | Y | Y | Y | Y | Y | Y | | | | | | Removed in 2011 due to additional of dual-frame weights. Replaced by PROVWT_LL. |
| PROVWT_D | DUAL-FRAME WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| PROVWT_LL | LANDLINE-FRAME WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN (EXCLUDING U.S. VIRGIN ISLANDS) | | | | | | | | Y | | | | | Removed in 2012. |
| PROVWTVI | WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN (INCLUDING U.S. VIRGIN ISLANDS) | | | | | | Y | Y | | | | | | Removed in 2011 due to additional of dual-frame weights. Replaced by PROVWTVI_LL. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---------------------------|
| PROVWTVI_D | COMBINATION OF THE DUAL-FRAME WEIGHT FOR CHILDREN IN THE U.S. PROPER AND LANDLINE WEIGHT FOR CHILDREN IN THE U.S. VIRGIN ISLANDS FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN | | | | | | | | | Y | | | | Added 2012. Removed 2013. |
| PROVWTVIGU_D | THE DUAL-FRAME WEIGHT FOR CHILDREN IN THE U.S. PROPER, THE U.S. VIRGIN ISLANDS AND GUAM FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN | | | | | | | | | | Y | | | Added 2013. Removed 2014. |
| PROVWT_D_TERR | THE DUAL-FRAME WEIGHT FOR CHILDREN IN THE U.S. PROPER AND TERRITORIES FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN | | | | | | | | | | | Y | Y | Added 2014. |
| PROVWTVI_LL | LANDLINE-FRAME WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN (INCLUDING U.S. VIRGIN ISLANDS) | | | | | | | | Y | | | | | Added 2011. Removed 2012. |
| PU431_31 | UTD FLAG FOR PROVIDER 4:3:1::3:1 (4:3:1:3:3:1 EXCLUDING HIB; INCLUDES 1+ VARICELLA AT AGE 12+ MTHS) BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| PU431_314 | UTD FLAG FOR PROVIDER 4:3:1::3:1:4 (4:3:1:3:3:1:4 EXCLUDING HIB; INCLUDES 1+ VARICELLA AT AGE 12+ MTHS) BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | | | | Y | Y | Y | Y | Y | Y | Introduced in 2010. |
| PU431331 | UTD FLAG FOR PROVIDER 4:3:1:3:3:1 (INCLUDES 1+ VARICELLA AT AGE 12+ MTHS) BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PU4313313 | UTD FLAG FOR PROVIDER 4:3:1:3:3:1:3 (INCLUDES 1+ VARICELLA AT AGE 12+ MTHS) BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PU4313314 | UTD FLAG FOR PROVIDER 4:3:1:3:3:1:4 (INCLUDES 1+ VARICELLA AT AGE 12+ MTHS) BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|--------------------|---|------|------|------|------|------|------|------|------|------|------|------|----------|--|
| PUT43133 | UTD FLAG FOR PROVIDER 4:3:1:3:3 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| PUTD4313 | UTD FLAG FOR PROVIDER 4:3:1:3 BY 36 MONTHS OF AGE, EXCLUDING ANY VACCINATIONS AFTER THE HOUSEHOLD INTERVIEW DATE. | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| Q5WEB1 | INTEREST IN IHQ ON WEBSITE PROVIDER #1 | Y | | | | | | | | | | | | Question was not asked starting in 2005. |
| Q5WEB2 | INTEREST IN IHQ ON WEBSITE PROVIDER #2 | Y | | | | | | | | | | | | Question was not asked starting in 2005. |
| Q5WEB3 | INTEREST IN IHQ ON WEBSITE PROVIDER #3 | Y | | | | | | | | | | | | Question was not asked starting in 2005. |
| Q5WEB4 | INTEREST IN IHQ ON WEBSITE PROVIDER #4 | Y | | | | | | | | | | | | Question was not asked starting in 2005. |
| Q5WEB5 | INTEREST IN IHQ ON WEBSITE PROVIDER #5 | Y | | | | | | | | | | | | Question was not asked starting in 2005. |
| RACE_K | RACE OF CHILD (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RACEETHK | RACE/ETHNICITY OF CHILD (RECODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RACEETHK_VI | RACE/ETHNICITY OF CHILD FOR U.S. VIRGIN ISLANDS (RECODE) | | | | | | | | | | | | Y | Added in 2015 for U.S. Virgin Islands. |
| RB1_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB2_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB3_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB4_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB5_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB6_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB7_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB8_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RB9_AGE | AGE IN MONTHS OF PROV-REPTD RUBELLA-ONLY SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| RDDWT | HH-PHASE CHILD INTERVIEW WEIGHT (EXCLUDING U.S. VIRGIN ISLANDS) | | Y | Y | Y | Y | Y | Y | | | | | | Removed in 2011 due to additional of dual-frame weights. Replaced by RDDWT_LL. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|--|
| RDDWT_D | DUAL-FRAME HH-PHASE CHILD INTERVIEW WEIGHT | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| RDDWT_LL | LANDLINE-FRAME HH-PHASE CHILD INTERVIEW WEIGHT (EXCLUDING U.S. VIRGIN ISLANDS) | | | | | | | | Y | | | | | Added 2011. Removed 2012. |
| RDDWTVI | HH-PHASE CHILD INTERVIEW WEIGHT (INCLUDING U.S. VIRGIN ISLANDS) | | | | | | Y | Y | | | | | | Removed in 2011 due to additional of dual-frame weights. Replaced by RDDWTVI_LL. |
| RDDWTVI_D | COMBINATION OF THE DUAL-FRAME HH-PHASE WEIGHT FOR HOUSEHOLDS IN THE U.S. PROPER AND LANDLINE HH-PHASE WEIGHT FOR HOUSEHOLDS IN THE U.S. VIRGIN ISLANDS | | | | | | | | | Y | | | | Added 2012. Removed 2013. |
| RDDWTVIGU_D | THE DUAL-FRAME HH-PHASE WEIGHT FOR HOUSEHOLDS IN THE U.S. PROPER, THE U.S. VIRGIN ISLANDS AND GUAM | | | | | | | | | | Y | | | Added 2013. Removed 2014. |
| RDDWT_D_TERR | THE DUAL-FRAME HH-PHASE WEIGHT FOR HOUSEHOLDS IN THE U.S. PROPER AND TERRITORIES | | | | | | | | | | | Y | Y | Added 2014. |
| RDDWTVI_LL | LANDLINE-FRAME HH-PHASE CHILD INTERVIEW WEIGHT (INCLUDING U.S. VIRGIN ISLANDS) | | | | | | | | Y | | | | | Added 2011. Removed 2012. |
| REGISTRY | CHILD'S PROVIDERS REPORTED CHILD'S VACCINATIONS TO IMMUNIZATION REGISTRY | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| RENT_OWN | IS HOME OWNED/BEING BOUGHT, RENTED, OR OCCUPIED BY SOME OTHER ARRANGEMENT? | | | | | | Y | Y | Y | Y | Y | Y | Y | |
| ROT1_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT2_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT3_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT4_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT5_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #5 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT6_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #6 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT7_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #7 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| ROT8_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #8 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| ROT9_AGE | AGE IN MONTHS OF PROV-REPTD ROTAVIRUS-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| SC_431 | HH SHOT CARD REPORT OF 4:3:1 UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_4313 | HH SHOT CARD REPORT OF 4:3:1:3 UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_43133 | HH SHOT CARD REPORT OF 4:3:1:3:3 UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_DTP | HH SHOT CARD REPORT OF 4+ DT-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_HEPB | HH SHOT CARD REPORT OF 3+ HEPATITIS B-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_HIB | HH SHOT CARD REPORT OF 3+ HIB-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_MCV | HH SHOT CARD REPORT OF 1+ MEASLES-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_POL | HH SHOT CARD REPORT OF 3+ POLIO-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SC_VRC | HH SHOT CARD REPORT OF 1+ VARICELLA-CONTAINING UP-TO-DATE | | | Y | Y | Y | Y | Y | Y | | | | | Added in 2006 as a partial replacement for the "FULL" and "C_" variables. Dropped in 2012 due to questionnaire changes. |
| SEQNUMC | UNIQUE CHILD IDENTIFIER | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| SEQNUMHH | UNIQUE HOUSEHOLD IDENTIFIER | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| SEX | GENDER OF CHILD | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| SHORT | Q1/2004 SHORT QUESTIONNAIRE STUDY FLAG | Y | | | | | | | | | | | | There was no short questionnaire study in 2005. |
| STATE | TRUE STATE OF RESIDENCE (STATE FIPS CODE) | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| STRATUM | STRATUM VARIABLE FOR DUAL-FRAME VARIANCE ESTIMATION | | | | | | | | | Y | Y | Y | Y | Added 2012. Equal to sample frame by estimation area. |
| STRATUM_D | STRATUM VARIABLE FOR DUAL-FRAME VARIANCE ESTIMATION | | | | | | | | Y | | | | | Added 2011. Equal to sample frame by estimation area. |
| TEL_SAMPFRAME | SAMPLE FRAME INDICATOR | | | | | | | | Y | | | | | Added 2011. Removed 2012. |
| U1D_HEP | BIRTH DOSE HEPATITIS B-CONTAINING GIVEN FROM BIRTH TO DAY 1 FLAG | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| U2D_HEP | BIRTH DOSE HEPATITIS B-CONTAINING GIVEN FROM BIRTH TO DAY 2 FLAG | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| U3D_HEP | BIRTH DOSE HEPATITIS B-CONTAINING GIVEN FROM BIRTH TO DAY 3 FLAG | | | | | | | | Y | Y | Y | Y | Y | Added 2011. |
| VFC_I | DERIVED: IS CHILD VFC ELIGIBLE? | | | | | | Y | Y | Y | | | | | Removed in 2012 |
| VFC_ORDER | DO CHILD'S PROVIDERS ORDER VACCINES FROM STATE/LOCAL HEALTH DEPT? | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| VFC_PRO | PARTICIPATION OF CHILD'S PROVIDERS IN VACCINES FOR CHILDREN PROGRAM | Y | Y | | | | | | | | | | | Question was not asked starting in 2006. |
| VRC1_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #1 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| VRC2_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #2 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| VRC3_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #3 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| VRC4_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #4 | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| VRC5_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #5 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| VRC6_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #6 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| VRC7_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #7 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| VRC8_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #8 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| VRC9_AGE | AGE IN MONTHS OF PROV-REPTD VARICELLA-CONTAINING SHOT #9 | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| WGT | NEW WEIGHT FOR CHILDREN WITH ADEQUATE PROVIDER DATA AND UNVACCINATED CHILDREN | Y | | | | | | | | | | | | Replaced by PROVWT starting in 2005. |
| WGT_RDD | RDD CHILD INTERVIEW WEIGHT | Y | | | | | | | | | | | | Replaced by RDDWT starting in 2005. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| XDTPTY1 | DT-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY2 | DT-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY3 | DT-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY4 | DT-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY5 | DT-CONTAINING VACCINATION #5 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY6 | DT-CONTAINING VACCINATION #6 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY7 | DT-CONTAINING VACCINATION #7 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY8 | DT-CONTAINING VACCINATION #8 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XDTPTY9 | DT-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XFLUTY1 | SEASONAL FLU-CONTAINING VACCINATION #1 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY2 | SEASONAL FLU-CONTAINING VACCINATION #2 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY3 | SEASONAL FLU-CONTAINING VACCINATION #3 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY4 | SEASONAL FLU-CONTAINING VACCINATION #4 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY5 | SEASONAL FLU-CONTAINING VACCINATION #5 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY6 | SEASONAL FLU-CONTAINING VACCINATION #6 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY7 | SEASONAL FLU-CONTAINING VACCINATION #7 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XFLUTY8 | SEASONAL FLU-CONTAINING VACCINATION #8 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes§ |
|---------------|---|------|------|------|------|------|------|------|------|------|------|------|------|---|
| XFLUTY9 | SEASONAL FLU-CONTAINING VACCINATION #9 TYPE CODE | | | | | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2008, influenza type boxes were added to the IHQ shot grid. |
| XH1NTY1 | MONOVALENT 2009 H1N1 FLU VACCINATION #1 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY2 | MONOVALENT 2009 H1N1 FLU VACCINATION #2 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY3 | MONOVALENT 2009 H1N1 FLU VACCINATION #3 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY4 | MONOVALENT 2009 H1N1 FLU VACCINATION #4 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY5 | MONOVALENT 2009 H1N1 FLU VACCINATION #5 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY6 | MONOVALENT 2009 H1N1 FLU VACCINATION #6 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY7 | MONOVALENT 2009 H1N1 FLU VACCINATION #7 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY8 | MONOVALENT 2009 H1N1 FLU VACCINATION #8 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XH1NTY9 | MONOVALENT 2009 H1N1 FLU VACCINATION #9 TYPE CODE | | | | | | | Y | Y | Y | | | | Introduced in 2010. Removed 2013. |
| XHEPTY1 | HEPATITIS B-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY2 | HEPATITIS B-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY3 | HEPATITIS B-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY4 | HEPATITIS B-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY5 | HEPATITIS B-CONTAINING VACCINATION #5 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY6 | HEPATITIS B-CONTAINING VACCINATION #6 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY7 | HEPATITIS B-CONTAINING VACCINATION #7 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY8 | HEPATITIS B-CONTAINING VACCINATION #8 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHEPTY9 | HEPATITIS B-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XHIBTY1 | HIB-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY2 | HIB-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| XHIBTY3 | HIB-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY4 | HIB-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY5 | HIB-CONTAINING VACCINATION #5 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY6 | HIB-CONTAINING VACCINATION #6 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY7 | HIB-CONTAINING VACCINATION #7 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY8 | HIB-CONTAINING VACCINATION #8 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XHIBTY9 | HIB-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XMMRTY1 | MEASLES-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XMMRTY2 | MEASLES-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XMMRTY3 | MEASLES-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XMMRTY4 | MEASLES-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XMMRTY5 | MEASLES-CONTAINING VACCINATION #5 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XMMRTY6 | MEASLES-CONTAINING VACCINATION #6 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XMMRTY7 | MEASLES-CONTAINING VACCINATION #7 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XMMRTY8 | MEASLES-CONTAINING VACCINATION #8 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XMMRTY9 | MEASLES-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XPCVTY1 | PNEUMOCOCCAL-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY2 | PNEUMOCOCCAL-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY3 | PNEUMOCOCCAL-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

| Variable Name | Variable Label† | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|---|
| XPCVTY4 | PNEUMOCOCCAL-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY5 | PNEUMOCOCCAL-CONTAINING VACCINATION #5 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY6 | PNEUMOCOCCAL-CONTAINING VACCINATION #6 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY7 | PNEUMOCOCCAL-CONTAINING VACCINATION #7 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY8 | PNEUMOCOCCAL-CONTAINING VACCINATION #8 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPCVTY9 | PNEUMOCOCCAL-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XPOLTY1 | POLIO-CONTAINING VACCINATION #1 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY2 | POLIO-CONTAINING VACCINATION #2 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY3 | POLIO-CONTAINING VACCINATION #3 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY4 | POLIO-CONTAINING VACCINATION #4 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY5 | POLIO-CONTAINING VACCINATION #5 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY6 | POLIO-CONTAINING VACCINATION #6 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY7 | POLIO-CONTAINING VACCINATION #7 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY8 | POLIO-CONTAINING VACCINATION #8 TYPE CODE | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |
| XPOLTY9 | POLIO-CONTAINING VACCINATION #9 TYPE CODE | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Starting in 2005, nine shot variables are included for each vaccine category. |
| XROTTY1 | ROTAVIRUS-CONTAINING VACCINATION #1 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY2 | ROTAVIRUS-CONTAINING VACCINATION #2 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY3 | ROTAVIRUS-CONTAINING VACCINATION #3 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY4 | ROTAVIRUS-CONTAINING VACCINATION #4 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY5 | ROTAVIRUS-CONTAINING VACCINATION #5 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY6 | ROTAVIRUS-CONTAINING VACCINATION #6 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |

| Variable Name | Variable Label [†] | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Notes [§] |
|---------------|--|------|------|------|------|------|------|------|------|------|------|------|------|--|
| XROTTY7 | ROTAVIRUS-CONTAINING VACCINATION #7 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY8 | ROTAVIRUS-CONTAINING VACCINATION #8 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XROTTY9 | ROTAVIRUS-CONTAINING VACCINATION #9 TYPE CODE | | | | | | Y | Y | Y | Y | Y | Y | Y | Rotavirus vaccination types were added to the IHQ starting 2009. |
| XVRCTY1 | VARICELLA-CONTAINING VACCINATION #1 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY2 | VARICELLA-CONTAINING VACCINATION #2 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY3 | VARICELLA-CONTAINING VACCINATION #3 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY4 | VARICELLA-CONTAINING VACCINATION #4 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY5 | VARICELLA-CONTAINING VACCINATION #5 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY6 | VARICELLA-CONTAINING VACCINATION #6 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY7 | VARICELLA-CONTAINING VACCINATION #7 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY8 | VARICELLA-CONTAINING VACCINATION #8 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| XVRCTY9 | VARICELLA-CONTAINING VACCINATION #9 TYPE CODE | | | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Varicella vaccination types were added to the IHQ starting 2006. |
| YEAR | YEAR OF INTERVIEW | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | Y | |

* For a list of variables that appeared in one or more (but not all) NIS public-use data files from 1995-2004, see "Alphabetical Listing of Variables that are Not Available in All Public-Use Data Files, National Immunization Survey, 1995-2004": www.cdc.gov/nchs/data/nis/pufvariables1995to2004.pdf

† If the variable appeared in the 2015 NIS public-use data file, then the 2015 label is given; otherwise the label from the most recent NIS public-use data file in which the variable appeared is given.

§ Starting in 2005, a code of 77 is used for "Don't Know" responses and a code of 99 is used for "Refused" responses.

Appendix F: Summary Tables

Table F.1: Estimated Population Totals and Sample Sizes of Children 19-35 Months of Age by State and Estimation Area, National Immunization Survey, 2015

| State/Estimation Area | ESTIAP | Estimated Population Total of Children | Number of Children with Complete Household Interviews | Number of Children with Adequate Provider Data | Percent of Children with Adequate Provider Data |
|-----------------------|--------|--|---|--|---|
| U.S. National* | | 5,701,540 | 26,975 | 15,167 | 56.2 |
| Alabama | 20 | 84,042 | 494 | 272 | 55.1 |
| Alaska | 74 | 15,312 | 498 | 295 | 59.2 |
| Arizona | 66 | 125,139 | 479 | 266 | 55.5 |
| Arkansas | 46 | 54,196 | 419 | 229 | 54.7 |
| California | 68 | 724,480 | 646 | 345 | 53.4 |
| Colorado | 60 | 95,511 | 440 | 246 | 55.9 |
| Connecticut | 1 | 54,156 | 469 | 250 | 53.3 |
| Delaware | 13 | 15,717 | 410 | 232 | 56.6 |
| District of Columbia | 12 | 12,398 | 576 | 321 | 55.7 |
| Florida | 22 | 315,014 | 546 | 273 | 50.0 |
| Georgia | 25 | 186,272 | 540 | 286 | 53.0 |
| Hawaii | 72 | 25,966 | 469 | 261 | 55.7 |
| Idaho | 75 | 30,723 | 342 | 209 | 61.1 |
| Illinois | | 222,542 | 985 | 559 | 56.8 |
| IL-City of Chicago | 35 | 56,511 | 320 | 188 | 58.8 |
| IL-Rest of State | 34 | 166,031 | 665 | 371 | 55.8 |
| Indiana | 36 | 121,057 | 426 | 231 | 54.2 |
| Iowa | 56 | 55,273 | 444 | 279 | 62.8 |
| Kansas | 57 | 58,367 | 367 | 228 | 62.1 |
| Kentucky | 27 | 78,299 | 437 | 241 | 55.1 |
| Louisiana | 47 | 88,669 | 427 | 223 | 52.2 |
| Maine | 4 | 18,614 | 386 | 221 | 57.3 |
| Maryland | 14 | 105,465 | 658 | 361 | 54.9 |
| Massachusetts | 2 | 102,930 | 521 | 307 | 58.9 |
| Michigan | 38 | 162,007 | 427 | 254 | 59.5 |
| Minnesota | 40 | 100,888 | 521 | 289 | 55.5 |
| Mississippi | 28 | 54,235 | 464 | 233 | 50.2 |
| Missouri | 58 | 106,677 | 401 | 226 | 56.4 |
| Montana | 61 | 17,101 | 390 | 251 | 64.4 |
| Nebraska | 59 | 37,665 | 374 | 231 | 61.8 |
| Nevada | 73 | 51,393 | 377 | 214 | 56.8 |
| New Hampshire | 5 | 19,116 | 372 | 195 | 52.4 |
| New Jersey | 8 | 153,667 | 557 | 272 | 48.8 |

| State/Estimation Area | ESTIAP | Estimated Population Total of Children | Number of Children with Complete Household Interviews | Number of Children with Adequate Provider Data | Percent of Children with Adequate Provider Data |
|------------------------------|---------------|---|--|---|--|
| New Mexico | 49 | 39,050 | 400 | 244 | 61.0 |
| New York | | 334,940 | 1,119 | 566 | 50.6 |
| NY-City of New York | 11 | 167,965 | 564 | 291 | 51.6 |
| NY-Rest of State | 10 | 166,975 | 555 | 275 | 49.5 |
| North Carolina | 29 | 173,795 | 483 | 261 | 54.0 |
| North Dakota | 62 | 15,262 | 398 | 234 | 58.8 |
| Ohio | 41 | 198,461 | 418 | 219 | 52.4 |
| Oklahoma | 50 | 75,920 | 376 | 220 | 58.5 |
| Oregon | 76 | 67,036 | 341 | 192 | 56.3 |
| Pennsylvania | | 204,792 | 1,039 | 575 | 55.3 |
| PA-Philadelphia County | 17 | 33,050 | 487 | 272 | 55.9 |
| PA-Rest of State | 16 | 171,741 | 552 | 303 | 54.9 |
| Rhode Island | 6 | 16,313 | 463 | 272 | 58.7 |
| South Carolina | 30 | 82,138 | 487 | 259 | 53.2 |
| South Dakota | 63 | 17,989 | 399 | 219 | 54.9 |
| Tennessee | 31 | 117,280 | 372 | 210 | 56.5 |
| Texas | | 568,417 | 2,586 | 1,500 | 58.0 |
| TX-Bexar County | 55 | 39,136 | 428 | 224 | 52.3 |
| TX-City of Houston | 54 | 71,255 | 360 | 214 | 59.4 |
| TX-El Paso County | 53 | 20,393 | 348 | 211 | 60.6 |
| TX-Hidalgo County | 107 | 23,985 | 370 | 255 | 68.9 |
| TX-Rest of State | 51 | 413,647 | 1,080 | 596 | 55.2 |
| Utah | 64 | 73,266 | 379 | 247 | 65.2 |
| Vermont | 7 | 8,540 | 409 | 262 | 64.1 |
| Virginia | 18 | 149,556 | 655 | 373 | 56.9 |
| Washington | 77 | 127,236 | 526 | 309 | 58.7 |
| West Virginia | 19 | 29,917 | 453 | 241 | 53.2 |
| Wisconsin | 44 | 98,535 | 389 | 245 | 63.0 |
| Wyoming | 65 | 10,205 | 421 | 219 | 52.0 |
| U.S. Virgin Islands | 95 | 1,956 | 580 | 327 | 56.4 |
| Guam | 105 | 5,058 | 467 | 282 | 60.4 |
| Puerto Rico | 106 | 50,148 | 617 | 224 | 36.3 |

*Excludes territories.

Table F.2: Estimated Population Totals and Sample Sizes for Age Group by Maternal Education, National Immunization Survey, 2015

| Age Group in Months | Maternal Education | Children with Completed Household Interviews* | Children with Completed Household Interviews* | Children with Adequate Provider Data* | Children with Adequate Provider Data* |
|---------------------|---------------------------|---|---|---------------------------------------|---------------------------------------|
| | | Unweighted Completes | Weighted Completes [†] | Unweighted Completes | Weighted Completes [§] |
| 19-23 | <12 Years | 756 | 273,432 | 469 | 263,247 |
| 19-23 | 12 Years | 1,424 | 448,449 | 807 | 457,944 |
| | >12, Non College Graduate | | | | |
| 19-23 | Graduate | 1,947 | 389,501 | 1,082 | 399,633 |
| 19-23 | College Grad | 3,527 | 607,050 | 1,980 | 597,607 |
| 24-29 | <12 Years | 814 | 298,700 | 482 | 303,578 |
| 24-29 | 12 Years | 1,505 | 484,913 | 833 | 473,346 |
| | >12, Non College Graduate | | | | |
| 24-29 | Graduate | 2,112 | 465,164 | 1,175 | 477,178 |
| 24-29 | College Grad | 3,688 | 668,547 | 2,132 | 663,221 |
| 30-35 | <12 Years | 1,113 | 325,879 | 692 | 334,888 |
| 30-35 | 12 Years | 2,038 | 552,154 | 1,129 | 550,668 |
| | >12, Non College Graduate | | | | |
| 30-35 | Graduate | 2,907 | 466,162 | 1,517 | 441,571 |
| 30-35 | College Grad | 5,144 | 721,590 | 2,869 | 738,657 |
| Total | | 26,975 | 5,701,540 | 15,167 | 5,701,540 |

* Excludes territories.

[†] Weighted by RDDWT_D.

[§] Weighted by PROVWT_D.

Table F.3: Estimated Population Totals and Sample Sizes for Age Group by Poverty Status, National Immunization Survey, 2015

| Age Group in Months | Poverty Status | Children with Completed Household Interviews* | Children with Completed Household Interviews* | Children with Adequate Provider Data* | Children with Adequate Provider Data* |
|------------------------|-------------------------|---|---|---|---|
| | | Unweighted Completes | Weighted Completes [†] | Unweighted Completes | Weighted Completes [§] |
| 19-23 Months | Above poverty, > \$75K | 2,795 | 474,404 | 1,591 | 482,155 |
| 19-23 Months | Above poverty, <= \$75K | 2,621 | 551,486 | 1,470 | 563,074 |
| 19-23 Months | Below poverty | 1,797 | 574,338 | 1,103 | 550,792 |
| 19-23 Months | Unknown | 441 | 118,203 | 174 | 122,411 |
| 24-29 Months | Above poverty, > \$75K | 3,025 | 573,078 | 1,737 | 562,122 |
| 24-29 Months | Above poverty, <= \$75K | 2,751 | 594,221 | 1,534 | 599,758 |
| 24-29 Months | Below poverty | 1,954 | 633,207 | 1,211 | 662,296 |
| 24-29 Months | Unknown | 389 | 116,817 | 140 | 93,148 |
| 30-35 Months | Above poverty, > \$75K | 4,248 | 611,297 | 2,400 | 620,955 |
| 30-35 Months | Above poverty, <= \$75K | 3,754 | 660,064 | 2,003 | 667,292 |
| 30-35 Months | Below poverty | 2,602 | 668,741 | 1,589 | 661,362 |
| 30-35 Months | Unknown | 598 | 125,682 | 215 | 116,176 |
| Total | | 26,975 | 5,701,540 | 15,167 | 5,701,540 |

* Excludes territories.

[†] Weighted by RDDWT_D.

[§] Weighted by PROVWT_D.

Table F.4: Estimated Population Totals and Sample Sizes for Race/Ethnicity by Poverty Status, National Immunization Survey, 2015

| Race/Ethnicity [†] | Poverty Status | Children with Completed Household Interviews* | Children with Completed Household Interviews* | Children with Adequate Provider Data* | Children with Adequate Provider Data* |
|-----------------------------------|-------------------------|---|---|---------------------------------------|---------------------------------------|
| | | Unweighted Completes | Weighted Completes [§] | Unweighted Completes | Weighted Completes [¶] |
| Hispanic | Above poverty, > \$75K | 1,027 | 196,447 | 538 | 187,381 |
| Hispanic | Above poverty, <= \$75K | 1,723 | 399,792 | 951 | 415,299 |
| Hispanic | Below poverty | 2,495 | 803,584 | 1,639 | 812,944 |
| Hispanic | Unknown | 396 | 118,051 | 193 | 105,048 |
| Non-Hispanic White Only | Above poverty, > \$75K | 7,119 | 1,140,847 | 4,135 | 1,160,642 |
| Non-Hispanic White Only | Above poverty, <= \$75K | 5,341 | 968,219 | 3,015 | 989,306 |
| Non-Hispanic White Only | Below poverty | 1,873 | 475,946 | 1,133 | 491,212 |
| Non-Hispanic White Only | Unknown | 642 | 140,748 | 190 | 116,712 |
| Non-Hispanic Black Only | Above poverty, > \$75K | 522 | 95,321 | 239 | 84,766 |
| Non-Hispanic Black Only | Above poverty, <= \$75K | 903 | 224,090 | 417 | 206,887 |
| Non-Hispanic Black Only | Below poverty | 1,128 | 380,213 | 616 | 358,192 |
| Non-Hispanic Black Only | Unknown | 201 | 55,529 | 80 | 66,075 |
| Non-Hispanic Other & Multi-Racial | Above poverty, > \$75K | 1,400 | 226,163 | 816 | 232,443 |
| Non-Hispanic Other & Multi-Racial | Above poverty, <= \$75K | 1,159 | 213,670 | 624 | 218,630 |
| Non-Hispanic Other & Multi-Racial | Below poverty | 857 | 216,544 | 515 | 212,101 |
| Non-Hispanic Other & Multi-Racial | Unknown | 189 | 46,374 | 66 | 43,900 |
| Total | | 26,975 | 5,701,540 | 15,167 | 5,701,540 |

* Excludes territories.

† Race/Ethnicity is self-reported and mutually exclusive.

§ Weighted by RDDWT_D.

¶ Weighted by PROVWT_D.

Table F.5: Estimated Population Totals and Sample Sizes for Age Group by Race/Ethnicity, National Immunization Survey, 2015

| Age Group in Months | Race/Ethnicity of Child [†] | Children with Completed Household Interviews* | Children with Completed Household Interviews* | Children with Adequate Provider Data* | Children with Adequate Provider Data* |
|---------------------|--------------------------------------|---|---|---------------------------------------|---------------------------------------|
| | | Unweighted Completes | Weighted Completes [§] | Unweighted Completes | Weighted Completes [¶] |
| 19-23 Months | Hispanic | 1,579 | 470,006 | 943 | 452,111 |
| 19-23 Months | Non-Hispanic White Only | 4,265 | 815,891 | 2,425 | 831,833 |
| 19-23 Months | Non-Hispanic Black Only | 797 | 227,087 | 396 | 215,264 |
| 19-23 Months | Non-Hispanic Other & Multi-Racial | 1,013 | 205,446 | 574 | 219,224 |
| 24-29 Months | Hispanic | 1,764 | 516,184 | 1,054 | 525,206 |
| 24-29 Months | Non-Hispanic White Only | 4,490 | 933,686 | 2,580 | 947,684 |
| 24-29 Months | Non-Hispanic Black Only | 806 | 234,763 | 386 | 214,139 |
| 24-29 Months | Non-Hispanic Other & Multi-Racial | 1,059 | 232,690 | 602 | 230,294 |
| 30-35 Months | Hispanic | 2,298 | 531,684 | 1,324 | 543,355 |
| 30-35 Months | Non-Hispanic White Only | 6,220 | 976,183 | 3,468 | 978,356 |
| 30-35 Months | Non-Hispanic Black Only | 1,151 | 293,304 | 570 | 286,517 |
| 30-35 Months | Non-Hispanic Other & Multi-Racial | 1,533 | 264,614 | 845 | 257,556 |
| Total | | 26,975 | 5,701,540 | 15,167 | 5,701,540 |

* Excludes territories.

[†] Race/Ethnicity is self-reported and mutually exclusive.

[§] Weighted by RDDWT_D.

[¶] Weighted by PROVWT_D.

Table F.6: Estimated Population Totals and Sample Sizes for Age Group by Gender, National Immunization Survey, 2015

| Age Group in Months | Gender | Children with Completed Household Interviews* | Children with Completed Household Interviews* | Children with Adequate Provider Data* | Children with Adequate Provider Data* |
|---------------------|--------|---|---|---------------------------------------|---------------------------------------|
| | | Unweighted Completes | Weighted Completes [†] | Unweighted Completes | Weighted Completes [§] |
| 19-23 Months | Male | 4,032 | 907,683 | 2,280 | 896,588 |
| 19-23 Months | Female | 3,622 | 810,748 | 2,058 | 821,844 |
| 24-29 Months | Male | 4,199 | 960,425 | 2,411 | 975,049 |
| 24-29 Months | Female | 3,920 | 956,899 | 2,211 | 942,274 |
| 30-35 Months | Male | 5,696 | 1,048,592 | 3,210 | 1,045,063 |
| 30-35 Months | Female | 5,506 | 1,017,193 | 2,997 | 1,020,722 |
| Total | | 26,975 | 5,701,540 | 15,167 | 5,701,540 |

* Excludes territories.

[†] Weighted by RDDWT_D.

[§] Weighted by PROVWT_D.

Table F.7: Estimated Vaccination Coverage* with Individual Vaccines and Selected Vaccination Series Among Children 19-35 Months of Age by State and Estimation Area, National Immunization Survey Q1/2015-Q4/2015†

| | ≥4 DTaP [§] | ≥3 Polio [¶] | ≥1 MMR ^{**} | Hib-FS ^{††} | ≥3 HepB ^{¶¶} | HepB Birth Dose | ≥1 HepA | ≥1 Var ^{***} | ≥4 PCV ^{†††} | Rotavirus | 4:3:1:3*:3:1:4 ^{§§§} |
|------------------------------------|----------------------|-----------------------|----------------------|----------------------|-----------------------|-----------------|------------|-----------------------|-----------------------|------------|-------------------------------|
| U.S. National^{¶¶¶} | 84.6 ± 1.1 | 93.7 ± 0.6 | 91.9 ± 0.8 | 82.7 ± 1.1 | 92.6 ± 0.7 | 72.4 ± 1.4 | 85.8 ± 1.1 | 91.8 ± 0.8 | 84.1 ± 1.1 | 73.2 ± 1.4 | 72.2 ± 1.4 |
| Alabama | 82.2 ± 6.4 | 95.1 ± 2.9 | 95.2 ± 3.5 | 83.3 ± 5.9 | 92.4 ± 3.9 | 83.2 ± 5.4 | 88.9 ± 5.2 | 92.9 ± 4.2 | 85.7 ± 5.5 | 76.2 ± 6.9 | 70.6 ± 7.1 |
| Alaska | 79.7 ± 5.2 | 91.9 ± 3.4 | 89.7 ± 4.0 | 81.6 ± 5.0 | 90.5 ± 3.7 | 65.7 ± 6.1 | 83.5 ± 4.9 | 81.8 ± 5.3 | 80.3 ± 5.2 | 72.0 ± 5.9 | 66.3 ± 6.2 |
| Arizona | 83.7 ± 5.8 | 92.4 ± 3.9 | 90.6 ± 4.1 | 85.5 ± 5.3 | 90.1 ± 4.2 | 78.0 ± 6.0 | 87.2 ± 5.1 | 92.7 ± 3.7 | 81.7 ± 5.8 | 75.1 ± 6.6 | 72.3 ± 6.7 |
| Arkansas | 76.4 ± 6.9 | 91.9 ± 4.4 | 90.2 ± 5.0 | 75.5 ± 6.9 | 94.4 ± 3.1 | 80.6 ± 6.2 | 85.1 ± 5.9 | 89.1 ± 5.2 | 76.1 ± 6.8 | 68.2 ± 7.4 | 66.6 ± 7.5 |
| California | 85.2 ± 5.7 | 96.3 ± 2.4 | 92.8 ± 3.9 | 85.2 ± 5.5 | 94.3 ± 3.4 | 69.1 ± 7.4 | 87.9 ± 5.4 | 93.4 ± 3.8 | 86.2 ± 5.4 | 74.6 ± 7.1 | 75.0 ± 6.9 |
| Colorado | 86.1 ± 5.0 | 92.7 ± 4.1 | 93.6 ± 3.6 | 83.6 ± 5.4 | 92.1 ± 4.1 | 73.1 ± 6.4 | 86.9 ± 5.0 | 92.5 ± 3.9 | 87.4 ± 4.9 | 75.9 ± 6.5 | 75.4 ± 6.2 |
| Connecticut | 90.8 ± 4.5 | 97.9 ± 1.8 | 97.5 ± 2.4 | 89.0 ± 5.0 | 97.3 ± 2.3 | 81.8 ± 6.2 | 97.3 ± 2.2 | 96.8 ± 2.3 | 91.5 ± 4.6 | 77.9 ± 6.7 | 80.6 ± 6.0 |
| Delaware | 89.9 ± 4.5 | 97.0 ± 2.7 | 97.2 ± 2.6 | 85.5 ± 5.5 | 95.1 ± 3.4 | 76.0 ± 6.7 | 96.1 ± 2.8 | 95.8 ± 3.1 | 90.8 ± 4.4 | 81.5 ± 6.1 | 79.3 ± 6.1 |
| Dist. of Columbia | 86.1 ± 5.0 | 90.9 ± 4.3 | 92.4 ± 3.8 | 85.4 ± 5.2 | 89.4 ± 4.6 | 72.7 ± 5.8 | 93.3 ± 3.6 | 94.4 ± 3.3 | 89.9 ± 4.7 | 73.0 ± 6.2 | 76.3 ± 6.0 |
| Florida | 86.0 ± 5.2 | 93.9 ± 3.4 | 90.4 ± 5.0 | 86.0 ± 5.2 | 90.4 ± 4.4 | 56.9 ± 7.6 | 77.7 ± 6.1 | 93.0 ± 3.4 | 85.9 ± 5.3 | 63.8 ± 7.5 | 66.6 ± 7.2 |
| Georgia | 82.3 ± 6.0 | 91.9 ± 4.3 | 90.5 ± 4.7 | 83.5 ± 6.0 | 94.2 ± 3.7 | 80.5 ± 5.7 | 88.2 ± 5.1 | 89.5 ± 5.0 | 84.8 ± 5.6 | 73.8 ± 7.0 | 75.6 ± 6.7 |
| Hawaii | 83.2 ± 5.4 | 93.7 ± 3.5 | 94.7 ± 3.3 | 82.8 ± 5.4 | 91.8 ± 3.6 | 75.5 ± 6.3 | 87.5 ± 4.6 | 91.6 ± 3.9 | 80.6 ± 5.8 | 72.5 ± 6.4 | 73.8 ± 6.3 |
| Idaho | 81.0 ± 5.9 | 88.9 ± 4.9 | 91.2 ± 4.0 | 81.2 ± 5.9 | 89.8 ± 4.6 | 69.8 ± 6.9 | 87.2 ± 4.9 | 88.5 ± 4.6 | 84.4 ± 5.5 | 74.5 ± 6.6 | 71.6 ± 6.8 |
| Illinois | 85.0 ± 3.8 | 93.3 ± 2.5 | 91.6 ± 2.8 | 82.7 ± 3.8 | 93.2 ± 2.3 | 71.2 ± 4.6 | 83.9 ± 3.7 | 91.7 ± 2.7 | 83.9 ± 3.9 | 75.4 ± 4.5 | 70.8 ± 4.7 |
| IL-City of Chicago | 86.2 ± 5.9 | 92.0 ± 4.3 | 90.5 ± 4.7 | 84.1 ± 6.1 | 87.7 ± 5.1 | 82.9 ± 5.9 | 86.4 ± 5.4 | 92.2 ± 4.4 | 84.5 ± 5.9 | 78.3 ± 6.8 | 72.8 ± 7.3 |
| IL-Rest of State | 84.6 ± 4.6 | 93.7 ± 3.0 | 91.9 ± 3.4 | 82.3 ± 4.6 | 95.1 ± 2.5 | 67.2 ± 5.8 | 83.1 ± 4.6 | 91.6 ± 3.3 | 83.8 ± 4.8 | 74.4 ± 5.5 | 70.1 ± 5.8 |
| Indiana | 85.3 ± 5.7 | 94.9 ± 3.4 | 92.0 ± 4.4 | 83.3 ± 5.9 | 91.7 ± 4.6 | 80.0 ± 6.3 | 89.2 ± 5.1 | 94.1 ± 3.8 | 83.4 ± 6.0 | 72.6 ± 7.2 | 74.7 ± 7.0 |
| Iowa | 88.9 ± 4.0 | 95.9 ± 2.4 | 95.5 ± 2.4 | 90.0 ± 3.5 | 96.3 ± 2.1 | 78.2 ± 5.3 | 86.7 ± 4.5 | 92.1 ± 3.6 | 88.1 ± 4.3 | 75.1 ± 5.9 | 77.9 ± 5.5 |
| Kansas | 86.8 ± 4.7 | 94.9 ± 3.3 | 92.3 ± 4.1 | 84.6 ± 5.1 | 95.7 ± 2.8 | 83.3 ± 5.4 | 90.3 ± 4.1 | 92.8 ± 3.5 | 86.2 ± 5.0 | 77.0 ± 6.3 | 75.2 ± 6.3 |
| Kentucky | 87.0 ± 4.8 | 94.9 ± 2.7 | 91.6 ± 4.1 | 83.5 ± 5.2 | 92.9 ± 3.9 | 75.3 ± 6.7 | 74.0 ± 6.5 | 95.7 ± 2.6 | 86.1 ± 4.8 | 65.3 ± 7.1 | 73.0 ± 6.5 |
| Louisiana | 84.4 ± 5.7 | 94.6 ± 3.0 | 92.6 ± 4.3 | 77.2 ± 7.0 | 94.8 ± 2.9 | 75.3 ± 7.1 | 84.8 ± 6.0 | 93.4 ± 3.9 | 84.2 ± 5.8 | 67.7 ± 7.6 | 70.8 ± 7.6 |
| Maine | 92.0 ± 5.0 | 95.0 ± 4.1 | 96.0 ± 3.1 | 84.5 ± 6.6 | 88.9 ± 5.4 | 68.7 ± 7.7 | 81.0 ± 6.8 | 93.8 ± 3.6 | 84.9 ± 6.5 | 71.1 ± 7.7 | 71.8 ± 7.9 |
| Maryland | 87.6 ± 4.5 | 94.4 ± 3.4 | 95.4 ± 2.5 | 87.0 ± 4.8 | 94.0 ± 3.4 | 79.0 ± 6.6 | 93.1 ± 3.2 | 91.5 ± 3.8 | 89.5 ± 4.2 | 76.8 ± 6.4 | 76.8 ± 5.9 |
| Massachusetts | 87.2 ± 5.1 | 96.9 ± 2.2 | 91.8 ± 4.0 | 86.6 ± 5.3 | 96.0 ± 2.3 | 78.4 ± 5.8 | 84.9 ± 5.2 | 92.0 ± 4.0 | 87.2 ± 5.0 | 83.5 ± 5.4 | 78.5 ± 6.0 |
| Michigan | 84.9 ± 5.5 | 92.9 ± 4.0 | 90.6 ± 4.5 | 79.5 ± 6.2 | 93.5 ± 3.8 | 80.0 ± 5.9 | 86.5 ± 5.1 | 89.5 ± 4.7 | 80.5 ± 6.3 | 65.5 ± 7.7 | 67.6 ± 7.3 |
| Minnesota | 85.4 ± 5.0 | 94.5 ± 3.1 | 92.6 ± 3.6 | 83.2 ± 5.3 | 94.7 ± 2.9 | 67.8 ± 6.5 | 90.4 ± 4.2 | 93.4 ± 3.2 | 86.6 ± 5.1 | 82.6 ± 5.4 | 73.2 ± 6.4 |
| Mississippi | 79.6 ± 7.0 | 92.0 ± 5.3 | 89.8 ± 5.3 | 74.3 ± 7.5 | 92.9 ± 4.7 | 77.0 ± 7.3 | 71.6 ± 7.5 | 89.2 ± 5.7 | 78.0 ± 7.2 | 65.9 ± 8.0 | 70.6 ± 7.5 |
| Missouri | 82.6 ± 6.3 | 90.8 ± 5.0 | 91.6 ± 4.3 | 82.7 ± 6.3 | 91.4 ± 4.6 | 75.0 ± 7.4 | 84.1 ± 5.7 | 92.4 ± 4.1 | 83.4 ± 6.2 | 76.1 ± 7.0 | 71.0 ± 7.4 |
| Montana | 78.6 ± 6.5 | 91.1 ± 3.9 | 91.7 ± 3.7 | 81.2 ± 5.7 | 91.3 ± 4.0 | 73.8 ± 6.5 | 76.4 ± 6.6 | 89.1 ± 4.2 | 75.6 ± 6.7 | 67.3 ± 7.0 | 68.1 ± 7.1 |
| Nebraska | 86.9 ± 5.0 | 97.4 ± 2.0 | 95.6 ± 3.0 | 85.5 ± 4.8 | 93.7 ± 3.2 | 72.5 ± 6.8 | 93.4 ± 3.0 | 94.4 ± 2.9 | 87.1 ± 4.5 | 73.0 ± 6.4 | 73.8 ± 6.3 |
| Nevada | 84.7 ± 5.7 | 92.3 ± 4.1 | 93.7 ± 3.7 | 81.7 ± 5.9 | 91.4 ± 4.5 | 77.7 ± 6.8 | 90.2 ± 4.6 | 92.9 ± 4.0 | 82.4 ± 6.1 | 65.5 ± 7.6 | 71.3 ± 7.0 |
| New Hampshire | 88.4 ± 5.4 | 96.9 ± 2.6 | 93.4 ± 3.9 | 88.7 ± 5.2 | 95.8 ± 3.4 | 72.0 ± 7.0 | 85.6 ± 5.7 | 89.6 ± 5.0 | 86.8 ± 5.9 | 80.9 ± 6.2 | 74.1 ± 7.1 |
| New Jersey | 89.8 ± 4.8 | 93.1 ± 4.0 | 92.8 ± 4.4 | 87.6 ± 5.3 | 93.9 ± 3.7 | 63.9 ± 7.2 | 84.2 ± 5.5 | 93.0 ± 4.2 | 86.8 ± 4.9 | 75.2 ± 6.8 | 76.5 ± 6.5 |
| New Mexico | 84.8 ± 6.0 | 93.7 ± 3.7 | 89.7 ± 5.3 | 79.0 ± 6.8 | 91.9 ± 4.4 | 67.8 ± 7.5 | 85.9 ± 6.3 | 86.4 ± 6.1 | 82.9 ± 5.9 | 73.8 ± 6.9 | 70.1 ± 7.9 |
| New York | 87.4 ± 3.3 | 94.9 ± 2.0 | 92.5 ± 2.6 | 81.8 ± 3.9 | 93.8 ± 2.2 | 59.0 ± 4.7 | 80.9 ± 4.0 | 93.4 ± 2.4 | 82.7 ± 3.8 | 73.0 ± 4.5 | 71.9 ± 4.4 |
| NY-City of New York | 85.5 ± 5.0 | 93.7 ± 3.1 | 94.1 ± 2.9 | 80.1 ± 5.7 | 94.1 ± 2.9 | 53.4 ± 6.8 | 81.2 ± 5.6 | 94.1 ± 2.9 | 76.6 ± 6.0 | 71.1 ± 6.4 | 68.2 ± 6.5 |
| NY-Rest of State | 89.2 ± 4.3 | 96.0 ± 2.6 | 90.9 ± 4.3 | 83.5 ± 5.4 | 93.6 ± 3.3 | 64.6 ± 6.6 | 80.5 ± 5.6 | 92.8 ± 3.7 | 88.9 ± 4.4 | 75.0 ± 6.4 | 75.7 ± 6.1 |

| | ≥4 DTaP [§] | ≥3 Polio [†] | ≥1 MMR ^{**} | Hib-FS ^{††} | ≥3 HepB ^{†††} | HepB Birth Dose | ≥1 HepA | ≥1 Var ^{***} | ≥4 PCV ^{†††} | Rotavirus | 4:3:1:3*:3:1:4 ^{§§§} |
|-------------------------------|----------------------|-----------------------|----------------------|----------------------|------------------------|-----------------|------------|-----------------------|-----------------------|------------|-------------------------------|
| North Carolina | 83.9 ± 5.8 | 94.1 ± 3.8 | 94.3 ± 3.3 | 83.6 ± 5.6 | 93.4 ± 4.0 | 81.3 ± 6.0 | 83.7 ± 5.8 | 92.7 ± 3.8 | 82.2 ± 5.9 | 75.9 ± 6.7 | 76.4 ± 6.5 |
| North Dakota | 85.7 ± 5.1 | 96.7 ± 2.2 | 92.8 ± 3.8 | 85.6 ± 5.1 | 96.0 ± 2.5 | 87.5 ± 4.7 | 91.9 ± 3.6 | 95.3 ± 2.7 | 91.2 ± 4.1 | 79.8 ± 6.2 | 80.2 ± 5.7 |
| Ohio | 80.9 ± 6.8 | 91.8 ± 4.5 | 88.1 ± 5.6 | 78.6 ± 7.0 | 92.3 ± 4.3 | 77.9 ± 6.5 | 79.9 ± 7.4 | 86.2 ± 6.0 | 79.1 ± 6.9 | 71.8 ± 7.6 | 68.3 ± 7.9 |
| Oklahoma | 85.7 ± 6.3 | 92.4 ± 5.0 | 92.6 ± 5.1 | 83.7 ± 6.4 | 94.0 ± 4.2 | 80.4 ± 5.9 | 93.2 ± 4.9 | 93.6 ± 4.9 | 82.7 ± 6.4 | 67.2 ± 8.3 | 75.4 ± 7.3 |
| Oregon | 85.8 ± 5.7 | 93.4 ± 3.8 | 94.1 ± 3.7 | 79.1 ± 6.9 | 93.7 ± 3.7 | 72.5 ± 7.0 | 91.8 ± 4.0 | 93.0 ± 3.7 | 81.8 ± 6.4 | 72.9 ± 7.3 | 67.4 ± 7.8 |
| Pennsylvania | 88.7 ± 3.8 | 93.1 ± 3.2 | 90.9 ± 3.7 | 84.1 ± 4.7 | 91.9 ± 3.2 | 73.2 ± 6.0 | 86.1 ± 4.7 | 90.4 ± 4.0 | 88.6 ± 4.0 | 74.3 ± 5.9 | 72.8 ± 5.8 |
| PA-Philadelphia County | 87.2 ± 4.5 | 95.0 ± 2.5 | 93.2 ± 3.5 | 84.8 ± 4.8 | 93.4 ± 3.1 | 77.3 ± 6.1 | 93.9 ± 3.0 | 96.1 ± 2.4 | 82.3 ± 5.5 | 71.5 ± 6.4 | 76.1 ± 6.0 |
| PA-Rest of State | 88.9 ± 4.4 | 92.7 ± 3.8 | 90.5 ± 4.4 | 84.0 ± 5.6 | 91.6 ± 3.7 | 72.5 ± 7.0 | 84.6 ± 5.6 | 89.3 ± 4.7 | 89.8 ± 4.6 | 74.9 ± 6.9 | 72.2 ± 6.8 |
| Rhode Island | 90.5 ± 4.1 | 97.2 ± 2.2 | 94.5 ± 3.2 | 85.2 ± 5.4 | 96.2 ± 2.4 | 73.2 ± 6.4 | 93.1 ± 3.7 | 95.1 ± 3.0 | 90.4 ± 4.3 | 87.6 ± 4.9 | 77.2 ± 6.0 |
| South Carolina | 77.5 ± 7.0 | 91.3 ± 4.8 | 88.5 ± 5.1 | 76.1 ± 6.6 | 90.2 ± 4.9 | 68.9 ± 7.0 | 81.8 ± 5.9 | 91.0 ± 4.6 | 82.0 ± 6.1 | 69.8 ± 7.1 | 68.2 ± 7.3 |
| South Dakota | 83.1 ± 6.0 | 93.6 ± 4.2 | 91.4 ± 4.8 | 82.0 ± 6.2 | 91.8 ± 4.8 | 71.4 ± 7.3 | 77.6 ± 6.6 | 90.3 ± 4.9 | 84.2 ± 5.9 | 73.4 ± 7.0 | 75.6 ± 6.8 |
| Tennessee | 81.2 ± 6.5 | 91.8 ± 4.2 | 90.2 ± 4.8 | 81.1 ± 6.5 | 91.2 ± 4.6 | 64.7 ± 7.9 | 89.0 ± 5.2 | 89.7 ± 4.9 | 83.0 ± 6.3 | 70.9 ± 7.6 | 70.1 ± 7.5 |
| Texas | 82.1 ± 3.6 | 92.5 ± 2.5 | 92.5 ± 2.4 | 80.4 ± 3.7 | 90.7 ± 2.9 | 76.4 ± 3.9 | 90.7 ± 2.9 | 92.3 ± 2.4 | 83.3 ± 3.3 | 75.7 ± 3.9 | 71.2 ± 4.2 |
| TX-Bexar County | 80.1 ± 6.1 | 92.0 ± 4.3 | 89.2 ± 5.0 | 78.3 ± 6.3 | 89.3 ± 4.8 | 72.5 ± 6.7 | 90.0 ± 4.6 | 89.4 ± 4.8 | 78.9 ± 6.3 | 70.7 ± 7.0 | 67.5 ± 7.1 |
| TX-City of Houston | 80.5 ± 6.5 | 93.9 ± 4.0 | 92.8 ± 3.9 | 77.7 ± 6.8 | 89.2 ± 5.2 | 82.6 ± 5.7 | 89.6 ± 4.9 | 91.9 ± 4.3 | 84.1 ± 5.9 | 74.8 ± 6.8 | 70.5 ± 7.3 |
| TX-El Paso County | 82.8 ± 6.1 | 91.2 ± 4.2 | 90.6 ± 4.5 | 83.8 ± 5.7 | 90.5 ± 4.3 | 76.5 ± 7.0 | 92.9 ± 3.8 | 91.9 ± 4.3 | 84.8 ± 5.8 | 73.4 ± 7.1 | 71.6 ± 7.1 |
| TX-Hidalgo County | 82.4 ± 5.8 | 91.9 ± 4.5 | 86.9 ± 5.2 | 81.3 ± 5.9 | 90.5 ± 4.7 | 89.5 ± 4.8 | 90.7 ± 4.3 | 90.6 ± 4.8 | 85.3 ± 5.5 | 73.1 ± 6.4 | 71.6 ± 6.5 |
| TX-Rest of State | 82.5 ± 4.8 | 92.4 ± 3.3 | 93.3 ± 3.1 | 80.9 ± 4.8 | 91.1 ± 3.8 | 74.9 ± 5.2 | 90.8 ± 3.8 | 92.7 ± 3.2 | 83.4 ± 4.4 | 76.6 ± 5.1 | 71.6 ± 5.6 |
| Utah | 82.0 ± 5.8 | 92.2 ± 4.3 | 89.2 ± 5.2 | 79.3 ± 6.2 | 91.7 ± 4.3 | 79.6 ± 6.0 | 86.2 ± 5.5 | 88.8 ± 5.0 | 78.3 ± 6.6 | 74.5 ± 6.7 | 68.1 ± 7.3 |
| Vermont | 89.2 ± 4.2 | 94.6 ± 2.9 | 95.5 ± 2.7 | 86.8 ± 4.8 | 93.6 ± 3.3 | 49.4 ± 6.7 | 82.7 ± 5.0 | 91.5 ± 3.5 | 89.7 ± 4.3 | 72.7 ± 6.2 | 75.6 ± 5.9 |
| Virginia | 80.6 ± 7.0 | 90.2 ± 5.2 | 83.4 ± 6.7 | 73.9 ± 7.7 | 87.8 ± 5.7 | 67.3 ± 8.1 | 76.9 ± 7.5 | 84.2 ± 6.6 | 75.3 ± 7.6 | 67.1 ± 8.3 | 64.4 ± 8.3 |
| Washington | 87.3 ± 4.7 | 92.9 ± 3.5 | 95.3 ± 2.6 | 87.2 ± 4.7 | 91.4 ± 3.5 | 71.2 ± 6.2 | 89.2 ± 3.9 | 92.9 ± 3.1 | 86.5 ± 4.9 | 73.6 ± 6.3 | 77.1 ± 5.7 |
| West Virginia | 78.6 ± 6.8 | 90.9 ± 4.9 | 86.7 ± 5.7 | 75.4 ± 7.1 | 92.6 ± 4.5 | 68.3 ± 7.4 | 85.7 ± 6.0 | 85.9 ± 5.7 | 82.9 ± 6.2 | 69.6 ± 7.5 | 64.9 ± 7.8 |
| Wisconsin | 85.2 ± 5.7 | 93.1 ± 3.9 | 92.4 ± 4.3 | 77.3 ± 6.4 | 90.1 ± 4.6 | 73.1 ± 6.6 | 81.4 ± 6.1 | 90.3 ± 4.5 | 83.2 ± 5.7 | 75.3 ± 6.7 | 68.8 ± 7.1 |
| Wyoming | 83.2 ± 6.1 | 96.2 ± 2.4 | 93.5 ± 3.5 | 82.1 ± 6.0 | 95.9 ± 2.5 | 67.2 ± 7.3 | 73.0 ± 6.8 | 91.2 ± 4.1 | 83.5 ± 5.8 | 76.4 ± 6.6 | 73.3 ± 7.1 |
| U.S. Virgin Islands | 65.8 ± 6.3 | 81.5 ± 5.1 | 77.2 ± 5.5 | 62.3 ± 6.5 | 85.5 ± 4.5 | 79.4 ± 5.3 | 73.3 ± 5.6 | 79.6 ± 5.1 | 61.6 ± 6.4 | 43.3 ± 6.9 | 50.7 ± 6.7 |
| Guam | 73.3 ± 6.1 | 85.5 ± 4.7 | 91.7 ± 3.6 | 64.0 ± 6.7 | 85.8 ± 4.8 | 82.3 ± 5.0 | 87.0 ± 4.4 | 78.7 ± 5.8 | 66.4 ± 6.9 | 40.6 ± 6.5 | 52.9 ± 7.0 |
| Puerto Rico | 82.0 ± 5.8 | 92.6 ± 3.5 | 91.4 ± 3.9 | 71.9 ± 7.5 | 86.6 ± 5.8 | 72.7 ± 7.6 | 85.7 ± 5.6 | 92.1 ± 3.8 | 73.4 ± 6.7 | 64.9 ± 7.6 | 61.0 ± 8.1 |

* Estimates presented as point estimate (%) ± 95% Confidence Interval. Estimate=NA (Not Available) if the unweighted sample size for the denominator was < 30, or (CI half width)/Estimate > 0.588, or (CI half width) > 10.

† Children in the Q1/2015-Q4/2015 National Immunization Survey were born from January 2012 through May 2014.

§ 4 or more doses of DTaP.

† 3 or more doses of any poliovirus vaccine.

** 1 or more doses of measles-mumps-rubella vaccine

†† 4 or more doses of Haemophilus influenzae type b (Hib) vaccine of any type or 2 doses of Hib of Merck types followed by 1+ dose of Hib of any type.

††† 3 or more doses of hepatitis B vaccine.

*** 1 or more doses of varicella at or after child's first birthday, unadjusted for history of varicella illness.

†††† 4 or more doses of PCV.

§§§ 4 or more doses of DTaP, 3 or more doses of any poliovirus vaccine, 1 or more doses of measles-containing vaccine, full series Hib vaccine, 3 or more doses of HepB, 1 or more doses of varicella vaccine, and 4 or more doses of PCV.

††††† U.S. national estimates exclude territories.

Appendix G: Trends in NIS Response Rates and Vaccination Coverage Rates, 1995-2015


Table G.1: Key Indicators* from Landline Sample Household and Provider Data Collection by Survey Year, National Immunization Survey, 1995-2015[†]

| Survey Year | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|-------------|---------------------|------------------------------|-------------------------------|-------------------------|--|
| 1995 | 96.5 | 96.4 | 93.5 | 87.1 | 50.6 |
| 1996 | 94.3 | 96.8 | 94.0 | 85.8 | 63.4 |
| 1997 | 92.1 | 97.9 | 93.8 | 84.6 | 69.7 |
| 1998 | 90.4 | 97.8 | 93.6 | 82.7 | 67.1 |
| 1999 | 88.6 | 97.0 | 93.4 | 80.2 | 65.4 |
| 2000 | 88.1 | 96.0 | 93.1 | 78.7 | 67.4 |
| 2001 | 86.8 | 96.2 | 91.1 | 76.1 | 70.4 |
| 2002 | 84.8 | 96.6 | 90.6 | 74.2 | 67.6 |
| 2003 | 83.6 | 94.0 | 88.7 | 69.8 | 68.9 |
| 2004 | 83.8 | 94.8 | 92.0 | 73.1 | 71.0 |
| 2005 | 83.3 | 92.8 | 84.2 | 65.1 | 63.6 |
| 2006 | 83.3 | 90.5 | 85.6 | 64.5 | 70.4 |
| 2007 | 82.9 | 90.2 | 86.8 | 64.9 | 68.6 |
| 2008 | 82.3 | 90.3 | 85.1 | 63.2 | 71.0 |
| 2009 | 82.9 | 92.4 | 83.2 | 63.8 | 68.7 |
| 2010 | 83.3 | 91.5 | 83.6 | 63.8 | 71.2 |
| 2011 | 83.0 | 90.7 | 81.7 | 61.6 | 72.3 |
| 2012 | 84.1 | 90.7 | 84.6 | 64.5 | 67.9 |
| 2013 | 83.2 | 91.0 | 82.3 | 62.3 | 63.5 |
| 2014 | 82.7 | 92.2 | 82.1 | 62.6 | 63.3 |
| 2015 | 81.9 | 89.9 | 80.3 | 59.1 | 59.7 |

* For the definition of the key indicators see Table 1 of NIS Data User's Guide for the survey year of interest.

[†] Excludes territories.

Figure G.1: Trends in Landline Sample Key Indicators from Household and Provider Data Collection by Survey Year, National Immunization Survey, 1995-2015*



* Excludes territories.

Figure G.1 provides a graphical representation of the data contained in Table G.1. It shows how selected landline sample key indicators from the household and provider data collection performed throughout the years, from 1995 to present. We observe that the trend in the data collection rates is going downward, with the exception of the percentage of children with adequate provider data, which had been essentially flat since 1997 until a drop in 2013. Note that these data reflect the landline sample only.


Table G.2: Key Indicators* from Cell-Phone Sample Household and Provider Data Collection by Survey Year, National Immunization Survey, 2011-2015†

| Survey Year | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|-------------|---------------------|------------------------------|-------------------------------|-------------------------|--|
| 2011 | 47.0 | 76.2 | 70.4 | 25.2 | 66.7 |
| 2012 | 52.4 | 77.5 | 75.5 | 30.6 | 63.9 |
| 2013 | 53.8 | 79.3 | 71.6 | 30.5 | 59.8 |
| 2014 | 58.7 | 78.5 | 72.6 | 33.5 | 58.9 |
| 2015 | 56.3 | 79.3 | 72.2 | 32.2 | 55.5 |

*For the definition of the key indicators see Table 1 of NIS Data User’s Guide for the survey year of interest.

† Excludes territories.

Figure G.2: Trends in Cell-Phone Sample Key Indicators from Household and Provider Data Collection by Survey Year, National Immunization Survey, 2011-2015*



* Excludes territories.

Figure G.2 provides a graphical representation of the data contained in Table G.2. It shows how selected cell-phone sample key indicators from the household and provider data collection performed from 2011 to present. We observe that the rates since the inception of the cell-phone sample have been essentially flat, aside from a declining percentage of children with adequate provider data and a moderate increase in the resolution rate in 2014 that is likely due to the process introduced in 2014 of removing and classifying as non-working cell-phone numbers flagged as having no recent activity.

The response rate is the number of households with a completed household interview divided by the estimated number of eligible households in the sample. Within each sample type (landline or cell phone), the number of eligible households was estimated using the CASRO assumptions; these assumptions are that the rate of households among the unresolved telephone numbers is the same as the observed rate of households among the resolved telephone numbers, and the rate of eligible households among unscreened households is the same as the observed rate of eligible households among screened households. Under these assumptions, within each sample type the CASRO response rate is equal to the product of the resolution rate, the screener completion rate, and the interview completion rate. For the combined samples, we have defined the CASRO response rate as the total number of households with a completed interview divided by the estimated total number of eligible households across both sample types, where the estimated total number of eligible households is equal to the sum of the estimated number of eligible households in the landline sample (using CASRO assumptions) and the estimated number of eligible households in the cell-phone sample (using CASRO assumptions). Table G.3 presents the CASRO response rate calculated in this way for the combined landline and cell-phone samples, by survey year, and Figure G.3 presents a graphical representation. Because the CASRO response rate is lower for the cell-phone sample than for the landline sample, the CASRO response rate for the combined landline and cell-phone samples was lower in years with a larger cell-phone sample and higher in years with a smaller cell-phone sample.


Table G.3: CASRO Response Rate for the Combined Landline and Cell-Phone Samples by Survey Year, National Immunization Survey, 2011-2015*

| Survey Year [†] | CASRO Response Rate (%) |
|--------------------------|-------------------------|
| 2011 | 52.2 |
| 2012 | 41.2 |
| 2013 | 34.4 |
| 2014 | 42.5 |
| 2015 | 34.9 |

* Excludes territories.

[†] Cell-phone sample was added to the NIS in 2011.

Figure G.3: Trend in CASRO Response Rate for the Combined Landline and Cell-Phone Samples by Survey Year, National Immunization Survey, 2011-2015*



* Excludes territories.

Table G.4: Vaccine-Specific Coverage Levels Among Children Age 19-35 Months in the United States by Survey Year, National Immunization Survey, 1995-2015*

| Survey Year [†] | 4+ DTaP | 3+ Polio | 1+ MMR | 3+ Hib [§] | 3+ Hep B | 1+ Varicella [¶] | 4+ PCV | 4:3:1 ^{**} | 4:3:1:3 ^{††} |
|--------------------------|---------|----------|--------|---------------------|----------|---------------------------|--------|---------------------|-----------------------|
| 1995 | 78.4 | 87.8 | 89.8 | 91.2 | 67.9 | N.A. | N.A. | 76.0 | 73.7 |
| 1996 | 81.1 | 91.0 | 90.6 | 91.4 | 81.8 | 12.0 | N.A. | 78.4 | 76.4 |
| 1997 | 81.5 | 90.7 | 90.4 | 92.5 | 83.6 | 25.8 | N.A. | 77.9 | 76.2 |
| 1998 | 83.9 | 90.8 | 92.0 | 93.4 | 87.0 | 43.2 | N.A. | 80.6 | 79.2 |
| 1999 | 83.3 | 89.6 | 91.5 | 93.5 | 88.1 | 57.5 | N.A. | 79.9 | 78.4 |
| 2000 | 81.7 | 89.5 | 90.5 | 93.4 | 90.3 | 67.8 | N.A. | 77.6 | 76.2 |
| 2001 | 82.1 | 89.4 | 91.4 | 93.0 | 88.9 | 76.3 | N.A. | 78.6 | 77.2 |
| 2002 | 81.6 | 90.2 | 91.6 | 93.1 | 89.9 | 80.6 | N.A. | 78.5 | 77.5 |
| 2003 | 84.8 | 91.6 | 93.0 | 93.9 | 92.4 | 84.8 | N.A. | 82.2 | 81.3 |
| 2004 | 85.5 | 91.6 | 93.0 | 93.5 | 92.4 | 87.5 | N.A. | 83.5 | 82.5 |
| 2005 | 85.7 | 91.7 | 91.5 | 93.9 | 92.9 | 87.9 | 53.7 | 83.1 | 82.4 |
| 2006 | 85.2 | 92.8 | 92.3 | 93.4 | 93.3 | 89.2 | 68.4 | 83.1 | 82.2 |
| 2007 | 84.5 | 92.6 | 93.2 | 92.6 | 92.7 | 90.0 | 75.3 | 82.8 | 80.1 |
| 2008 | 84.6 | 93.6 | 92.1 | 90.9 | 93.5 | 90.7 | 80.1 | 82.5 | 79.6 |
| 2009 | 83.9 | 92.8 | 90.0 | 83.6 | 92.4 | 89.6 | 80.4 | 81.5 | 73.4 |
| 2010 | 84.4 | 93.3 | 91.5 | 90.4 | 91.8 | 90.4 | 83.3 | 82.0 | 78.8 |
| 2011 | 84.6 | 93.9 | 91.6 | 94.0 | 91.1 | 90.8 | 84.4 | 82.6 | 81.9 |
| 2012 ^{§§} | 82.5 | 92.8 | 90.8 | 93.0 | 89.7 | 90.2 | 81.9 | 80.5 | 80.0 |
| 2013 | 83.1 | 92.7 | 91.9 | 92.8 | 90.8 | 91.2 | 82.0 | 81.5 | 81.1 |
| 2014 | 84.2 | 93.3 | 91.5 | 92.6 | 91.6 | 91.0 | 82.9 | 82.6 | 82.0 |
| 2015 | 84.6 | 93.7 | 91.9 | 93.2 | 92.6 | 91.8 | 84.1 | 83.2 | 82.6 |

* Excludes territories.

[†] Prior to 2011, estimates are single-frame, landline-sample estimates. From 2011 onward, estimates are dual-frame (landline plus cell-phone) estimates.

[§] Beginning in 2009, the number of doses required to be up-to-date on Hib depends on the manufacturer of the vaccine. However, the figures shown here refer to 3 or more doses of Hib vaccine regardless of manufacturer.


[¶] Varicella was added to the NIS in 1996.

^{**} Four or more doses of DTaP, three or more doses of poliovirus vaccine, and one or more doses of MCV.

^{††} Four or more doses of DTaP, three or more doses of poliovirus vaccine, one or more doses of MCV, and three or more doses of Hib.

^{§§} Revised definition of adequate provider data (APD) implemented.

Figure G.4: Trends in Vaccine-Specific Coverage Levels among Children 19-35 Months of Age in the United States by Survey Year, National Immunization Survey, 1995-2015*†¶



* Excludes territories.

† Prior to 2011, estimates are single-frame, landline-sample estimates. From 2011 onward, estimates are dual-frame (landline plus cell-phone) estimates.

¶ Revised definition of adequate provider data (APD) implemented in 2012.

Figure G.4 provides a graphical representation of the data contained in Table G.4. It displays the trend in vaccine-specific coverage levels among children aged 19 through 35 months from 1995 to present. We observe that the trend in the vaccination coverage levels is slightly upward for the longer-established vaccines, while the early trends for new vaccines are strongly upward. Note that these data reflect the landline sample prior to 2011 and the dual-frame sample thereafter.

Appendix H: Vaccine Type Codes

Table H.1: 2015 NIS Vaccine Type Codes

| Vaccine Code | Description |
|--------------|---------------------------------------|
| 03 | DTaP/DTP-containing, unknown type |
| 04 | DTaP |
| 07 | DTaP-Hib |
| 08 | DTaP-HepB-IPV |
| 20 | OPV |
| 21 | IPV |
| 22 | Polio-containing, unknown type |
| 30 | Measles-mumps-rubella |
| 31 | Measles only |
| 32 | Measles-mumps |
| 33 | Measles-rubella |
| 43 | HepB-Hib |
| 44 | Hib-only, unknown type |
| 60 | HepB-only |
| 70 | Pneumococcal conjugate, unknown type |
| 71 | Pneumococcal polysaccharide |
| 72 | Pneumococcal-containing, unknown type |
| 73 | Pneumococcal conjugate-7 |
| 74 | Pneumococcal conjugate-13 |
| D3 | DTaP-IPV-Hib |
| FL | Seasonal influenza, unknown type |
| FM | Seasonal influenza spray |
| FN | Injected seasonal influenza |
| HB | HepB-containing, unknown type |
| HG | Hib-only (GSK) |
| HI | Hib-containing, unknown type |
| HM | Hib-only (Merck) |
| HS | Hib-only (Sanofi) |
| HY | Hib-MenCY |
| MM | Measles-containing, unknown type |
| RG | Rotarix (GSK) |
| RM | Rotateq (Merck) |
| RO | Rotavirus-containing, unknown type |
| VA | Varicella-containing, unknown type |
| VM | MMR-varicella |
| VO | Varicella-only |

Appendix I: Key NIS Response Rates by Area

Table I.1: Key Indicators* for the Landline Sample by Estimation Area, National Immunization Survey, 2015

| Area | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|----------------------|---------------------|------------------------------|-------------------------------|-------------------------|--|
| U.S. National† | 81.9 | 89.9 | 80.3 | 59.1 | 59.7 |
| Alabama | 83.5 | 89.2 | 81.1 | 60.4 | 62.2 |
| Alaska | 87.8 | 90.9 | 83.6 | 66.7 | 64.2 |
| Arizona | 83.5 | 90.8 | 81.8 | 62.0 | 58.1 |
| Arkansas | 85.8 | 92.1 | 81.7 | 64.5 | 49.2 |
| California | 81.1 | 87.4 | 81.6 | 57.9 | 57.9 |
| Colorado | 83.9 | 89.3 | 81.5 | 61.0 | 58.9 |
| Connecticut | 77.1 | 88.2 | 79.0 | 53.8 | 60.3 |
| Delaware | 71.6 | 88.9 | 74.2 | 47.2 | 50.5 |
| District of Columbia | 82.2 | 89.8 | 80.9 | 59.7 | 61.7 |
| Florida | 81.9 | 90.1 | 80.7 | 59.5 | 52.6 |
| Georgia | 83.0 | 89.3 | 78.4 | 58.1 | 54.7 |
| Hawaii | 85.7 | 87.2 | 68.0 | 50.9 | 56.9 |
| Idaho | 86.7 | 91.1 | 90.5 | 71.5 | 63.2 |
| Illinois | 85.6 | 89.7 | 76.6 | 58.9 | 57.6 |
| IL-City of Chicago | 86.7 | 89.2 | 73.4 | 56.8 | 58.9 |
| IL-Rest of State | 84.1 | 90.3 | 80.8 | 61.4 | 56.1 |
| Indiana | 86.0 | 92.9 | 83.8 | 66.9 | 51.6 |
| Iowa | 87.3 | 92.6 | 89.4 | 72.3 | 67.2 |
| Kansas | 85.3 | 91.5 | 88.1 | 68.8 | 64.6 |
| Kentucky | 83.3 | 91.9 | 81.3 | 62.3 | 73.4 |
| Louisiana | 84.9 | 89.8 | 66.7 | 50.8 | 56.3 |
| Maine | 81.9 | 93.0 | 83.6 | 63.7 | 66.1 |
| Maryland | 77.6 | 88.3 | 77.8 | 53.3 | 52.5 |
| Massachusetts | 73.7 | 88.3 | 86.6 | 56.4 | 60.5 |
| Michigan | 84.5 | 91.3 | 84.6 | 65.3 | 58.1 |
| Minnesota | 85.3 | 91.7 | 81.5 | 63.8 | 59.0 |
| Mississippi | 85.9 | 89.9 | 72.3 | 55.9 | 50.0 |
| Missouri | 84.2 | 91.8 | 86.8 | 67.1 | 51.5 |
| Montana | 86.6 | 91.9 | 87.7 | 69.8 | 67.2 |
| Nebraska | 84.1 | 90.7 | 85.7 | 65.4 | 72.9 |
| Nevada | 78.7 | 90.3 | 84.4 | 60.0 | 58.9 |
| New Hampshire | 78.1 | 91.0 | 87.3 | 62.0 | 53.1 |

| Area | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|------------------------|----------------------------|--------------------------------------|---------------------------------------|---------------------------------|--|
| New Jersey | 76.2 | 87.2 | 80.1 | 53.3 | 55.9 |
| New Mexico | 86.6 | 90.9 | 90.5 | 71.2 | 76.4 |
| New York | 77.3 | 87.3 | 75.4 | 50.8 | 52.7 |
| NY-City of New York | 76.6 | 85.5 | 73.5 | 48.2 | 53.5 |
| NY-Rest of State | 78.0 | 88.8 | 77.2 | 53.5 | 51.9 |
| North Carolina | 81.2 | 90.1 | 83.5 | 61.1 | 66.7 |
| North Dakota | 87.9 | 93.1 | 75.4 | 61.7 | 54.9 |
| Ohio | 84.0 | 91.0 | 75.8 | 58.0 | 48.9 |
| Oklahoma | 84.4 | 91.0 | 90.4 | 69.4 | 61.7 |
| Oregon | 86.0 | 92.4 | 77.8 | 61.8 | 64.7 |
| Pennsylvania | 74.8 | 90.3 | 82.9 | 56.0 | 60.2 |
| PA-Philadelphia County | 73.9 | 88.9 | 81.6 | 53.6 | 49.4 |
| PA-Rest of State | 76.0 | 91.5 | 84.2 | 58.5 | 68.8 |
| Rhode Island | 69.2 | 89.2 | 76.8 | 47.4 | 63.6 |
| South Carolina | 82.8 | 90.3 | 74.1 | 55.4 | 45.9 |
| South Dakota | 88.7 | 91.8 | 80.4 | 65.5 | 65.9 |
| Tennessee | 84.1 | 91.2 | 76.1 | 58.3 | 60.7 |
| Texas | 84.4 | 87.7 | 72.8 | 53.9 | 61.4 |
| TX-Bexar County | 82.0 | 88.3 | 72.0 | 52.2 | 57.6 |
| TX-City of Houston | 85.9 | 86.6 | 70.2 | 52.2 | 54.7 |
| TX-El Paso County | 84.9 | 88.8 | 75.3 | 56.8 | 69.1 |
| TX-Hidalgo County | 85.0 | 86.9 | 77.3 | 57.1 | 74.1 |
| TX-Rest of State | 83.5 | 88.4 | 70.9 | 52.4 | 56.0 |
| Utah | 86.1 | 88.6 | 92.7 | 70.8 | 73.2 |
| Vermont | 81.6 | 92.0 | 90.6 | 68.0 | 72.1 |
| Virginia | 78.8 | 90.2 | 87.3 | 62.1 | 65.5 |
| Washington | 83.5 | 91.2 | 83.7 | 63.7 | 58.1 |
| West Virginia | 76.6 | 91.5 | 79.4 | 55.7 | 59.7 |
| Wisconsin | 83.0 | 92.0 | 76.9 | 58.7 | 54.9 |
| Wyoming | 85.9 | 90.3 | 87.1 | 67.6 | 62.1 |
| U.S. Virgin Islands | 92.1 | 93.4 | 83.8 | 72.1 | 60.9 |
| Guam | 91.5 | 88.1 | 76.6 | 61.7 | 62.8 |
| Puerto Rico | 90.1 | 94.2 | 65.9 | 56.0 | 40.9 |

* For the definition of the key indicators see Table 1 of NIS Data User's Guide.

† Excludes territories.

Table I.2: Key Indicators* for the Cell-Phone Sample by Estimation Area, National Immunization Survey, 2015

| Area | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|----------------------|---------------------|------------------------------|-------------------------------|-------------------------|--|
| U.S. National† | 56.3 | 79.3 | 72.2 | 32.2 | 55.5 |
| Alabama | 59.9 | 79.1 | 68.2 | 32.4 | 53.8 |
| Alaska | 77.8 | 83.0 | 78.9 | 50.9 | 57.7 |
| Arizona | 51.3 | 80.0 | 72.5 | 29.8 | 54.9 |
| Arkansas | 64.5 | 81.4 | 71.0 | 37.3 | 55.6 |
| California | 51.4 | 77.9 | 69.8 | 27.9 | 52.3 |
| Colorado | 53.8 | 78.2 | 75.4 | 31.7 | 55.5 |
| Connecticut | 42.0 | 77.2 | 69.7 | 22.6 | 51.0 |
| Delaware | 48.4 | 78.7 | 73.0 | 27.8 | 58.3 |
| District of Columbia | 53.5 | 79.6 | 75.2 | 32.0 | 53.8 |
| Florida | 51.3 | 78.9 | 71.2 | 28.9 | 49.6 |
| Georgia | 55.2 | 79.8 | 68.8 | 30.3 | 52.7 |
| Hawaii | 51.4 | 76.5 | 70.8 | 27.8 | 55.5 |
| Idaho | 52.2 | 80.6 | 78.8 | 33.1 | 60.9 |
| Illinois | 61.4 | 80.7 | 72.3 | 35.8 | 56.6 |
| IL-City of Chicago | 67.3 | 80.9 | 71.6 | 39.0 | 58.7 |
| IL-Rest of State | 58.7 | 80.6 | 72.8 | 34.4 | 55.8 |
| Indiana | 54.5 | 81.8 | 73.2 | 32.6 | 54.7 |
| Iowa | 58.4 | 83.4 | 78.7 | 38.3 | 62.1 |
| Kansas | 69.5 | 81.1 | 74.3 | 41.8 | 61.8 |
| Kentucky | 57.6 | 79.8 | 67.6 | 31.1 | 52.0 |
| Louisiana | 64.6 | 79.1 | 67.5 | 34.5 | 51.5 |
| Maine | 52.1 | 81.9 | 77.6 | 33.1 | 55.7 |
| Maryland | 53.5 | 77.0 | 75.5 | 31.1 | 55.4 |
| Massachusetts | 50.7 | 77.9 | 70.7 | 27.9 | 58.4 |
| Michigan | 58.3 | 80.5 | 75.4 | 35.4 | 59.8 |
| Minnesota | 54.1 | 80.2 | 74.8 | 32.5 | 54.9 |
| Mississippi | 63.3 | 80.1 | 65.1 | 33.0 | 50.2 |
| Missouri | 59.2 | 80.9 | 75.1 | 35.9 | 57.4 |
| Montana | 61.9 | 83.1 | 77.4 | 39.8 | 63.8 |
| Nebraska | 64.0 | 81.5 | 76.1 | 39.6 | 59.7 |
| Nevada | 53.3 | 78.3 | 70.7 | 29.5 | 56.3 |
| New Hampshire | 48.7 | 79.2 | 76.5 | 29.5 | 52.3 |
| New Jersey | 48.6 | 77.3 | 69.1 | 25.9 | 46.6 |

| Area | Resolution Rate (%) | Screener Completion Rate (%) | Interview Completion Rate (%) | CASRO Response Rate (%) | Children with Adequate Provider Data (%) |
|------------------------|----------------------------|--------------------------------------|---------------------------------------|---------------------------------|--|
| New Mexico | 59.6 | 80.4 | 74.3 | 35.6 | 58.6 |
| New York | 50.5 | 78.2 | 68.4 | 27.0 | 49.9 |
| NY-City of New York | 53.0 | 77.6 | 65.6 | 27.0 | 51.0 |
| NY-Rest of State | 47.9 | 79.0 | 72.2 | 27.3 | 48.8 |
| North Carolina | 54.0 | 78.2 | 72.2 | 30.5 | 52.0 |
| North Dakota | 66.6 | 82.6 | 77.8 | 42.8 | 59.4 |
| Ohio | 52.2 | 79.5 | 74.8 | 31.0 | 52.8 |
| Oklahoma | 64.0 | 79.2 | 75.4 | 38.2 | 58.1 |
| Oregon | 51.6 | 82.1 | 79.7 | 33.8 | 54.8 |
| Pennsylvania | 51.0 | 77.5 | 68.8 | 27.2 | 54.2 |
| PA-Philadelphia County | 51.4 | 77.3 | 69.1 | 27.5 | 57.3 |
| PA-Rest of State | 47.2 | 79.6 | 66.2 | 24.9 | 51.5 |
| Rhode Island | 46.8 | 77.3 | 69.6 | 25.2 | 57.2 |
| South Carolina | 54.3 | 79.2 | 66.1 | 28.4 | 54.2 |
| South Dakota | 63.0 | 82.0 | 76.7 | 39.6 | 51.6 |
| Tennessee | 54.6 | 80.5 | 74.4 | 32.7 | 55.7 |
| Texas | 58.4 | 76.1 | 66.5 | 29.5 | 57.5 |
| TX-Bexar County | 57.0 | 77.2 | 71.1 | 31.3 | 51.4 |
| TX-City of Houston | 52.9 | 75.8 | 67.2 | 26.9 | 60.7 |
| TX-El Paso County | 56.2 | 75.7 | 66.0 | 28.0 | 59.0 |
| TX-Hidalgo County | 65.3 | 75.1 | 62.1 | 30.5 | 68.0 |
| TX-Rest of State | 57.6 | 79.4 | 72.6 | 33.2 | 55.1 |
| Utah | 58.0 | 77.9 | 80.6 | 36.4 | 63.8 |
| Vermont | 49.3 | 81.1 | 77.7 | 31.0 | 61.1 |
| Virginia | 52.3 | 80.2 | 69.2 | 29.1 | 55.0 |
| Washington | 49.0 | 80.1 | 76.8 | 30.1 | 58.9 |
| West Virginia | 57.5 | 79.6 | 72.6 | 33.2 | 51.9 |
| Wisconsin | 58.7 | 80.5 | 75.5 | 35.6 | 64.2 |
| Wyoming | 78.0 | 82.8 | 80.1 | 51.8 | 50.1 |
| U.S. Virgin Islands | 78.3 | 84.5 | 65.3 | 43.2 | 55.3 |
| Guam | 61.8 | 78.3 | 55.2 | 26.7 | 59.3 |
| Puerto Rico | 68.4 | 86.5 | 62.5 | 37.0 | 35.5 |

* For the definition of the key indicators see Table 1 of NIS Data User's Guide.

† Excludes territories.

Table I.3: CASRO Response Rate for the Combined Landline and Cell-Phone Samples* by Estimation Area, National Immunization Survey, 2015

| Area | CASRO Response Rate (%) |
|----------------------|-------------------------|
| U.S. National† | 34.9 |
| Alabama | 34.7 |
| Alaska | 53.8 |
| Arizona | 33.4 |
| Arkansas | 39.5 |
| California | 33.6 |
| Colorado | 33.9 |
| Connecticut | 26.2 |
| Delaware | 30.0 |
| District of Columbia | 33.7 |
| Florida | 31.8 |
| Georgia | 32.6 |
| Hawaii | 29.3 |
| Idaho | 34.9 |
| Illinois | 37.9 |
| IL-City of Chicago | 41.3 |
| IL-Rest of State | 36.3 |
| Indiana | 35.6 |
| Iowa | 40.9 |
| Kansas | 44.3 |
| Kentucky | 33.5 |
| Louisiana | 36.1 |
| Maine | 35.7 |
| Maryland | 40.7 |
| Massachusetts | 32.3 |
| Michigan | 38.5 |
| Minnesota | 35.4 |
| Mississippi | 34.3 |
| Missouri | 39.3 |
| Montana | 42.3 |
| Nebraska | 42.2 |
| Nevada | 32.5 |
| New Hampshire | 32.0 |
| New Jersey | 30.1 |
| New Mexico | 38.3 |

| Area | CASRO Response Rate (%) |
|------------------------|--------------------------------|
| New York | 30.1 |
| NY-City of New York | 29.6 |
| NY-Rest of State | 31.0 |
| North Carolina | 33.3 |
| North Dakota | 44.3 |
| Ohio | 32.9 |
| Oklahoma | 40.7 |
| Oregon | 36.2 |
| Pennsylvania | 30.0 |
| PA-Philadelphia County | 29.0 |
| PA-Rest of State | 35.7 |
| Rhode Island | 27.6 |
| South Carolina | 30.3 |
| South Dakota | 43.3 |
| Tennessee | 35.2 |
| Texas | 31.3 |
| TX-Bexar County | 32.7 |
| TX-City of Houston | 28.7 |
| TX-El Paso County | 30.0 |
| TX-Hidalgo County | 31.7 |
| TX-Rest of State | 40.9 |
| Utah | 39.2 |
| Vermont | 35.4 |
| Virginia | 37.4 |
| Washington | 33.8 |
| West Virginia | 35.4 |
| Wisconsin | 37.8 |
| Wyoming | 53.4 |
| U.S. Virgin Islands | 46.8 |
| Guam | 32.2 |
| Puerto Rico | 38.9 |

* For the definition of the CASRO response rate for the combined landline and cell-phone samples, see footnote 6 of Table 1.

† Excludes territories.