MEDICATION TRACKER

The 12-Dose Regimen for Latent Tuberculosis (TB) Infection

Your Medication Schedule

(Providers: Indicate the appropriate number of pills and day)

Medicine	Number of pills per week	Frequency	Day
Isoniazid: mg Rifapentine: mg	TOTAL:(Isoniazid:, Rifapentine:)	Once a week for 12 weeks (3 months)	M T W Th F S Sun

Your doctor may also add Vitamin B6 to your treatment plan.

Keeping Track of Your Treatment

On the table below, check the box and write the date to show when you took your medicine.

WEEK	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
EXAMPLE 5/7 - 5/13		X 5/8					
Week 1							
Week 2		□					
Week 3		□			□	□	□
Week 4		□					
Week 5	□	□			□	□	□
Week 6		□	□	□	□	□	□
Week 7	□	□				□	□
Week 8	□	□					
Week 9		□				□	
Week 10		□	□	□	□	□	□
Week 11		□	□	□	□	□	□
Week 12		□					



SYMPTOM CHECKLIST

The 12-Dose Regimen for Latent Tuberculosis (TB) Infection

Patient Name:					
Normal Side Effects Most people can take their TB medicines without any problems. The rifapentine medicine may cause your urine (pee), saliva, tears, or sweat to appear an orange-red color. This is normal and the color may fade over time.					
	and CALL your TB doctor or any of the problems below: Skin or whites of your eyes appear yellow Skin rash or itching Bruises, or red or purple spots on your skin that you cannot explain Nosebleeds, or bleeding from your gums or around your teeth Shortness of breath Pain or tingling in your hands, arms, or legs Feelings of sadness or depression				
Please talk to your doctor or nurse if you have treatment for latent TB infection. Doctor/Clinic Contact Information Name of the staff caring for you: Phone number: Address: Hours:	ormation				

