

Prenatal Care and Routinely Recommended Vaccinations

Vaccine	Jan	Feb	Mar	April	May	June	July	Aug	Sept	Oct	Nov	Dec
Tdap												
COVID-19												
Influenza												
RSV												

LEGEND

Administer now	Administer based on schedule – see below
----------------	------------------------------------------

Tdap Vaccine

Vaccine product: Any Tdap vaccine product may be administered.

Schedule: Administer a dose of Tdap, preferably during the early part of gestational weeks 27 through 36, during each pregnancy irrespective of the patient's prior Tdap vaccination history.

2024-25 COVID-19 Vaccine

Vaccine product: Any COVID-19 vaccine product may be administered.

Schedule: Administer the current COVID-19 vaccine as indicated based on the patient's health and vaccination history. Administer vaccine during any trimester, including the 1st trimester.

Influenza Vaccine

Vaccine product: Only administer IIV or RIV vaccine products.

Schedule: Administer a dose of inactivated influenza vaccine (IIV) or recombinant influenza vaccine (RIV) to women who are pregnant during any trimester or will be pregnant during influenza season.

Consider vaccination during July and August for pregnant women in the 3rd trimester. For those in the 1st or 2nd trimester during July and August, waiting to vaccinate until September or October is preferable, unless there is concern that later vaccination might not be possible. Vaccination should continue throughout the season as long as influenza viruses are circulating.

RSV Vaccine

Vaccine product: Only administer Abrysvo (Pfizer) RSV vaccine.*

Schedule: Administer a dose of RSV vaccine (Abrysvo, Pfizer) only between 32 through 36 weeks of gestation during September through January (in most of the continental United States) if the patient was not previously vaccinated.

In jurisdictions where seasonality differs from most of the continental United States, such as Alaska, southern Florida, Guam, Hawaii, Puerto Rico, the U.S.-affiliated Pacific Islands, and the U.S. Virgin Islands, healthcare providers should follow state, local, or territorial guidance on timing of administering the RSV vaccine (Pfizer Abrysvo) for pregnant women.

Additional vaccines may be needed based on:

- Patient's age, health status, risk of exposure, vaccination history such as hepatitis B, meningococcal vaccines, pneumococcal vaccines. See [Immunization Schedules for Healthcare Professionals | CDC](#) for detailed guidance.
- Travel outside the United States. See [Travelers' Health | CDC](#) for detailed guidance.

*Nirsevimab may be administered to the infant as an alternative to vaccinating during pregnancy.

Additional Resources

[Maternal Vaccine Information for Health Care Providers](#)